FORMULAS FOR THE UNRAMIFIED BRAUER GROUP OF A PRINCIPAL HOMOGENEOUS SPACE OF A LINEAR ALGEBRAIC GROUP

Mikhail Borovoi and Boris Kunyavskiĭ

Abstract. For a smooth compactification V of a principal homogeneous space E under a connected linear algebraic group G defined over a field k of characteristic zero, we present two formulas expressing $\operatorname{Br} V/\operatorname{Br} k$ in terms of G.

Introduction

Let G be a connected linear algebraic group over a field k of characteristic zero, E a principal homogeneous space (torsor) under G, and V a smooth complete variety over k containing E as an open subset. The Brauer group $\operatorname{Br} V = H^2_{\text{ét}}(V, \mathbb{G}_m)$ does not depend on the choice of V; this birational invariant of E is denoted $\operatorname{Br}_{\text{nr}}(k(E)/k)$ and is called the unramified Brauer group of E. In this paper we give two formulas for $\operatorname{Br} V/\operatorname{Br} k$ in terms of G.

Formulas for $\operatorname{Br} V/\operatorname{Br} k$ were first given by Voskresenskiĭ [V] and Sansuc [S] in the case when k is a number field. A generalization to an arbitrary ground field was presented in [CT/K]. In all these three papers, G is of some special type: either a torus, or a semisimple group, or a group admitting a finite cover of the type $G_0 \times S$ where G_0 is a semisimple simply connected group and S is a quasi-trivial torus.

In this paper we compute $\operatorname{Br} V/\operatorname{Br} k$ for any connected k-group G. The paper is based on results of [CT/K]. We use the method of z-extensions developed by Kottwitz [K2], [K3].

We now briefly describe our results. Let G^{u} denote the unipotent radical of G. Set $G^{\mathrm{red}} = G/G^{\mathrm{u}}$; it is a reductive group. Let G^{ss} denote the derived group of G^{red}; it is semisimple. Let G^{sc} denote the universal covering of G^{ss}; it is simply connected. Consider the composite map

$$\rho \colon G^{\mathrm{sc}} \to G^{\mathrm{ss}} \to G^{\mathrm{red}}.$$

Let $T \subset G^{\mathrm{red}}$ be a maximal torus. Set $T^{\mathrm{sc}} = \rho^{-1}(T) \subset G^{\mathrm{sc}}$. Denote $L^{-1} = \mathbf{X}^*(T)$, $L^0 = \mathbf{X}^*(T^{\mathrm{sc}})$, where $\mathbf{X}^*(\cdot) = \operatorname{Hom}_{\bar{k}}(\cdot, \mathbb{G}_m)$ stands for the character group, and $\bar{k}$ denotes an algebraic closure of k. Consider the complex

$$L^\bullet = (0 \to L^{-1} \to L^0 \to 0) = (0 \to \mathbf{X}^*(T^{\mathrm{sc}}) \to \mathbf{X}^*(T) \to 0).$$

The first named author partially supported by the Hermann Minkowski Center for Geometry.

Research of the second named author partially supported by the Ministry of Absorption (Israel) and the Israel Science Foundation founded by the Israel Academy of Sciences and Humanities.

The Galois group $\mathfrak{g} = \mathrm{Gal}(\bar{k}/k)$ acts on $L^\bullet$. Let $\mathbb{H}^i(\mathfrak{g}, L^\bullet)$ denote the hypercohomology group of $\mathfrak{g}$ with coefficients in the complex $L^\bullet$. Set

$$Ш^i_\omega(\mathfrak{g}, L^\bullet) = \ker[\mathbb{H}^i(\mathfrak{g}, L^\bullet) \to \prod_\gamma \mathbb{H}^i(\gamma, L^\bullet)]$$

where γ runs over all closed procyclic subgroups of $\mathfrak{g}$. Let $\overline{V} = V \times_k \bar{k}$, and denote by $\mathrm{Pic}\,\overline{V}$ the Picard group; it is a $\mathfrak{g}$-module. Our main results are the following theorem and corollary.

Theorem A. *With the above assumptions and notation,*

$$H^1(k, \mathrm{Pic}\,\overline{V}) = Ш^1_\omega(\mathfrak{g}, L^\bullet).$$

Corollary B. *There is an injection*

$$\mathrm{Br}\,V/\mathrm{Br}\,k \hookrightarrow Ш^1_\omega(\mathfrak{g}, L^\bullet)$$

which is an isomorphism provided either $V(k) \neq \varnothing$*, or* $H^3(k, \bar{k}^\times) = 0$.

Let now $Z(\hat{G})$ denote the center of a connected Langlands dual group for a connected reductive group G, cf. [K2], 1.5. It is a $\mathbf{C}$-group of multiplicative type. It turns out that $Z(\hat{G}) = \ker[L^{-1} \otimes \mathbf{C}^\times \to L^0 \otimes \mathbf{C}^\times]$.

Proposition C.

$$Ш^1_\omega(\mathfrak{g}, L^\bullet) = Ш^1_\omega(\mathfrak{g}, Z(\hat{G})).$$

Thus

$$H^1(k, \mathrm{Pic}\,\overline{V}) = Ш^1_\omega(\mathfrak{g}, Z(\hat{G}^{\mathrm{red}})).$$

We obtain a new case of the following Kottwitz principle [K2]: an invariant of reductive groups which is trivial for semisimple simply connected groups can be computed from the Galois module $Z(\hat{G})$.

The structure of the paper is as follows. In Section 1 we collect required information on linear algebraic groups, Brauer groups, Galois cohomology and hypercohomology and prove Proposition C (Proposition 1.3.2). In Section 2 we state and prove our main results (Theorem A = Theorems 2.1 and 2.4, and Corollary B = Corollary 2.2). In Section 3 we present some comments and remarks relating our results to previously known ones.

Acknowledgements. We thank J.-L. Colliot-Thélène for very valuable remarks. The second named author thanks the Max-Planck-Institut für Mathematik (Bonn) for hospitality and support.

Notation and conventions

Throughout the paper, k denotes a field of characteristic zero, $\bar{k}$ is a fixed algebraic closure of k, $\mathfrak{g} = \mathrm{Gal}(\bar{k}/k)$ is the absolute Galois group of k, $k^\times$ denotes the multiplicative group of k. An algebraic k-torus T is called quasi-trivial if it is a direct product of tori of the form $R_{K/k}\mathbb{G}_m$ where K/k is a finite extension and $R_{K/k}$ stands for Weil's restriction of scalars. We denote by $\mathbf{X}^*(G)$ the group of characters of a linear algebraic group G and by $\mathbf{X}_*(T) = \mathrm{Hom}_{\bar{k}}(\mathbb{G}_m, T)$ the group of cocharacters of a torus T; one can view $\mathbf{X}^*(G)$ and $\mathbf{X}_*(T)$ as $\mathfrak{g}$-modules. For a torus T, $\mathbf{X}^*(T)$ is a $\mathbf{Z}$-free $\mathfrak{g}$-module of finite rank; if T is quasi-trivial, $\mathbf{X}^*(T)$ is a permutation module (i.e. it has a $\mathbf{Z}$-basis permuted by $\mathfrak{g}$). If M is a Galois module, we denote by $H^i(k,M)$ (or by $H^i(\mathfrak{g},M)$) the i-th Galois cohomology group. For a smooth projective k-variety X we denote $\overline{X} = X \times_k \bar{k}$. Set $\operatorname{Pic} X = H^1_{\text{ét}}(X, \mathbb{G}_m)$, $\operatorname{Br} X = H^2_{\text{ét}}(X, \mathbb{G}_m)$, these are the Picard group and the Brauer group of X, respectively. Other notation is explained in the introduction.

1. Preliminaries

1.1. Linear algebraic groups.

1.1.1. Let G be a connected linear algebraic group over a field k of characteristic zero. By Chevalley's theorem [C] the k-variety G is rational, i.e. $\bar{k}$-birationally equivalent to an affine space.

Every unipotent k-group is k-rational, i.e. k-birationally equivalent to an affine space.

Every quasi-trivial k-torus Z is k-rational, and by Hilbert 90 $H^1(K,Z) = 1$ for every extension K of k.

1.1.2. Levi decomposition. For any connected linear algebraic group G over a field k of characteristic zero, there is an isomorphism $G^{\mathrm{u}} \rtimes G^{\mathrm{red}} \xrightarrow{\sim} G$ (Levi decomposition) which gives rise to a k-biregular morphism of varieties

$$G^{\mathrm{u}} \times G^{\mathrm{red}} \xrightarrow{\sim} G.$$

1.1.3. z-extensions. A z-extension of a connected reductive k-group G is an epimorphism of reductive groups $\alpha\colon H \to G$ with kernel Z, such that H^{ss} is simply connected and Z is central and is a quasi-trivial k-torus. The notion of z-extension was introduced by Langlands [L]. We say that a z-extension $\alpha_1\colon H_1 \to G$ dominates a z-extension $\alpha_2\colon H_2 \to G$ if there exists a homomorphism $\phi\colon H_1 \to H_2$ such that $\alpha_2 = \phi \circ \alpha_1$.

Lemma 1.1.4. (Kottwitz) *(1) For every connected reductive k-group G and a cohomology class $\xi \in H^1(k,G)$ there exists a z-extension $\alpha\colon H \to G$ such that $\xi \in \operatorname{im}[\alpha_*\colon H^1(k,H) \to H^1(k,G)]$.*

(2) For every two z-extensions $\alpha_1\colon H_1 \to G$ and $\alpha_2\colon H_2 \to G$ of G there exists a z-extension $\alpha_3\colon H_3 \to G$ that dominates both α_1 and α_2.

(3) Let $G_1 \to G_2$ be a homomorphism, and let $H_i \to G_i$ $(i = 1,2)$ be z-extensions. Then there exists a commutative diagram

$$\begin{array}{ccccc} H_1 & \longleftarrow & H_3 & \longrightarrow & H_2 \\ \downarrow & & \downarrow & & \downarrow \\ G_1 & \xleftarrow{\mathrm{id}} & G_1 & \longrightarrow & G_2 \end{array}$$

in which $H_3 \to G_1$ is a z-extension.

Proof. (1) For a proof of existence of some z-extension of G see [M/S], Prop. 3.1. The existence of a z-extension such that ξ lifts to $H^1(k,H)$ is proved in [K3] in the proof of Theorem 1.2, p. 369.

(2) See [K1], Lemma 1.1(2).

(3) See [K2], Lemma 2.4.4. □

1.2. Birational invariants.

1.2.1. Permutation modules. A permutation $\mathfrak{g}$-module P can be written as a direct sum of induced modules $\mathbf{Z}[\mathfrak{g}/\mathfrak{h}]$, where $\mathfrak{h}$ is a closed subgroup of finite index in $\mathfrak{g}$. By Shapiro's lemma, $H^1(\mathfrak{g}, \mathbf{Z}[\mathfrak{g}/\mathfrak{h}]) = H^1(\mathfrak{h}, \mathbf{Z}) = 0$, hence $H^1(\mathfrak{g}, P) = 0$. Moreover, $H^1(\gamma, P) = 0$ for any closed subgroup $\gamma \subset \mathfrak{g}$.

We also have $Ш^2_\omega(\mathfrak{g}, P) = 0$ (cf. [S], (1.9.1), for the case where k is a number field). Indeed, it suffices to prove this for an induced module $M = \mathbf{Z}[\mathfrak{g}/\mathfrak{h}]$. We have $H^2(\mathfrak{g}, M) = H^2(\mathfrak{h}, \mathbf{Z}) = \mathrm{Hom}(\mathfrak{h}, \mathbf{Q}/\mathbf{Z})$. Since any continuous homomorphism $\mathfrak{h} \to \mathbf{Q}/\mathbf{Z}$ factors through a finite quotient of $\mathfrak{g}$, we may assume that $\mathfrak{h}$ and $\mathfrak{g}$ are finite. Since any non-trivial homomorphism $\mathfrak{h} \to \mathbf{Q}/\mathbf{Z}$ is non-trivial on some cyclic subgroup of $\mathfrak{h}$, we conclude that $Ш^2_\omega(\mathfrak{g}, M) = 0$.

1.2.2. Smooth compactifications. By Hironaka [H], any smooth affine k-variety X can be embedded into a smooth complete k-variety $V(X)$ containing X as an open subset. Indeed, one has to map X biregularly onto a closed subscheme of an affine space, embed it into the projective space, take the projective closure, and resolve its singularities. We call $V(X)$ a smooth k-compactification of X. If V_1 and V_2 are two smooth k-compactifications of X, then there exists an isomorphism of $\mathfrak{g}$-modules $\mathrm{Pic}\,\overline{V}_1 \oplus P_1 \cong \mathrm{Pic}\,\overline{V}_2 \oplus P_2$ where P_1 and P_2 are permutation $\mathfrak{g}$-modules (cf. [V], Th. 1). By 1.2.1, this gives an isomorphism $H^1(k, \mathrm{Pic}\,\overline{V}_1) \to H^1(k, \mathrm{Pic}\,\overline{V}_2)$, and the construction in [V] shows that this isomorphism is canonical. This also shows that $H^1(k, \mathrm{Pic}\,\overline{V}(X))$ is a birational invariant of X.

Moreover, the group $H^1(k, \mathrm{Pic}\,\overline{V}(X))$ is functorial in X. Indeed, let $f\colon X_1 \to X_2$ be a k-morphism of smooth integral k-varieties. We wish to extend f to a k-morphism $f'\colon V_1 \to V_2$ where V_i is a suitable smooth compactification of X_i, $i = 1, 2$. We are very grateful to J.-L. Colliot-Thélène for communicating us the following construction. Let U denote the graph of f in $X_1 \times_k X_2$. Choose smooth compactifications W_i of X_i, $i = 1, 2$. Let W be the closure of U in $W_1 \times_k W_2$. Then U is a smooth open subvariety of W. By Hironaka [H] there exists a proper morphism (desingularization) $\pi\colon V_1 \to W$ such that V_1 is smooth and the restriction $\pi^{-1}(U) \to U$ is an isomorphism. Clearly V_1 is a smooth compactification of X_1. Set $V_2 = W_2$ and define $f'\colon V_1 \to V_2$ to be the composite map $V_1 \to W \to V_2$ where the second arrow is the restriction of the canonical projection $W_1 \times W_2 \to W_2 = V_2$. The map f' induces a homomorphism $H^1(k, \mathrm{Pic}\,\overline{V}_1) \to H^1(k, \mathrm{Pic}\,\overline{V}_2)$, as required.

We prove the following property of the functor $H^1(k, \mathrm{Pic}\,\overline{V}(X))$: if Z is a k-rational variety, then $H^1(k, \mathrm{Pic}\,\overline{V}(X \times_k Z)) \cong H^1(k, \mathrm{Pic}\,\overline{V}(X))$. Indeed, let V_X, V_Z be smooth compactifications of X, Z, respectively. One can then take $V_X \times V_Z$ as a smooth compactification of $X \times Z$. The variety $\overline{V}_Z$ is rational, hence by [CT/S1], Lemme 11, p. 188, the canonical homomorphism $\mathrm{Pic}\,\overline{V}_X \oplus \mathrm{Pic}\,\overline{V}_Z \to \mathrm{Pic}\,(\overline{V}_X \times \overline{V}_Z)$ is an isomorphism. Since Z is k-rational, $H^1(k, \mathrm{Pic}\,\overline{V}_Z) = 0$. Thus $H^1(k, \mathrm{Pic}\,\overline{V}(X \times Z)) \cong H^1(k, \mathrm{Pic}\,\overline{V}(X))$, as required. This isomorphism is induced by the canonical projection $\mathrm{pr}_X\colon X \times Z \to X$.

1.2.3. Brauer group. For a geometrically integral smooth projective k-variety X we have an exact sequence

$$\mathrm{Br}\,k \to \ker[\mathrm{Br}\,X \to \mathrm{Br}\,\overline{X}] \to H^1(k, \mathrm{Pic}\,\overline{X}) \to H^3(k, \bar{k}^\times);$$

if X has a k-point, we have an exact sequence

$$0 \to \mathrm{Br}\,k \to \ker[\mathrm{Br}\,X \to \mathrm{Br}\,\overline{X}] \to H^1(k, \mathrm{Pic}\,\overline{X}) \to 0$$

(cf. [CT/S2], 1.5.0). If X is k-rational, this gives an isomorphism $\operatorname{Br} k \xrightarrow{\sim} \operatorname{Br} X$; if X is a smooth k-compactification of a G-torsor E with $X(k) \neq \varnothing$, this gives an isomorphism

$$\operatorname{Br} X/\operatorname{Br} k \cong H^1(k, \operatorname{Pic} \overline{X}),$$

because by 1.1.1 $\overline{X}$ is rational, and since $\overline{X}$ is projective and rational, $\operatorname{Br} \overline{X} = 0$.

1.3. Hypercohomology.

1.3.1. Let $M^\bullet = (0 \to M^{-1} \to M^0 \to 0)$ be a short complex of $\mathfrak{g}$-modules. We often shorten notation to $(M^{-1} \to M^0)$. We define the hypercohomology $\mathbb{H}^i(\mathfrak{g}, M^\bullet)$ as the cohomology H^i of the ordinary chain complex corresponding to the double complex

$$\begin{array}{ccccccccc}
0 & \longrightarrow & M^0 & \longrightarrow & C^1(\mathfrak{g}, M^0) & \longrightarrow & C^2(\mathfrak{g}, M^0) & \longrightarrow & \dots \\
 & & \uparrow & & \uparrow & & \uparrow & & \\
0 & \longrightarrow & M^{-1} & \longrightarrow & C^1(\mathfrak{g}, M^{-1}) & \longrightarrow & C^2(\mathfrak{g}, M^{-1}) & \longrightarrow & \dots
\end{array}$$

where C^i is the usual group of non-homogeneous continuous i-cochains and the bidegree of M^{-1} is $(-1, 0)$.

For a subgroup $\gamma \subset \mathfrak{g}$ one can define the restriction map $\mathbb{H}^i(\mathfrak{g}, M^\bullet) \to \mathbb{H}^i(\gamma, M^\bullet)$ and define

$$Ш^i_\omega(\mathfrak{g}, M^\bullet) = \ker \left[\mathbb{H}^i(\mathfrak{g}, M^\bullet) \to \prod_\gamma \mathbb{H}^i(\gamma, M^\bullet) \right]$$

where γ runs over all closed procyclic subgroups of $\mathfrak{g}$.

1.3.2. Let T be a maximal k-torus in a connected reductive k-group G, and let $Z(\hat{G})$ denote the center of a connected Langlands dual group for G (see Introduction). Denote $L^{-1} = \mathbf{X}^*(T)$, $L^0 = \mathbf{X}^*(T^{\mathrm{sc}})$, $L^\bullet = (L^{-1} \to L^0)$.

Proposition. *With the above notation*

$$Ш^1_\omega(\mathfrak{g}, L^\bullet) = Ш^1_\omega(\mathfrak{g}, Z(\hat{G})).$$

1.3.3. We first prove the following lemma.

Lemma. *There is a short exact sequence of* $\mathbf{C}$*-groups*

$$1 \to Z(\hat{G}) \to L^{-1} \otimes \mathbf{C}^\times \to L^0 \otimes \mathbf{C}^\times \to 1.$$

Proof. Set $\pi_1(G) = \mathbf{X}_*(T)/\rho(\mathbf{X}_*(T^{\mathrm{sc}}))$, where $\mathbf{X}_*(\cdot) = \operatorname{Hom}_{\bar{k}}(\mathbb{G}_m, \cdot)$ is the cocharacter group. By [B], 1.2, the Galois module $\pi_1(G)$ does not depend on the choice of $T \subset G$. We call $\pi_1(G)$ the algebraic fundamental group of G. By [B], 1.10, we have $\pi_1(G) = \operatorname{Hom}(Z(\hat{G}), \mathbf{C}^\times)$. Hence $Z(\hat{G}) = \operatorname{Hom}(\pi_1(G), \mathbf{C}^\times)$.

By the definition of $\pi_1(G)$, there is an exact sequence

$$0 \to \mathbf{X}_*(T^{\mathrm{sc}}) \to \mathbf{X}_*(T) \to \pi_1(G) \to 0.$$

We thus obtain an exact sequence

$$1 \to \operatorname{Hom}(\pi_1(G), \mathbf{C}^\times) \to \operatorname{Hom}(\mathbf{X}_*(T), \mathbf{C}^\times) \to \operatorname{Hom}(\mathbf{X}_*(T^{\mathrm{sc}}), \mathbf{C}^\times) \to 1,$$

or

$$1 \to Z(\hat{G}) \to L^{-1} \otimes \mathbf{C}^\times \to L^0 \otimes \mathbf{C}^\times \to 1. \quad \square$$

1.3.4. Proof of Proposition 1.3.2. From Lemma 1.3.3 we obtain a quasi-isomorphism of complexes $(Z(\hat{G}) \to 1) \to L^\bullet \otimes \mathbf{C}^\times$. Hence

$$\mathbb{H}^0(\mathfrak{g}, L^\bullet \otimes \mathbf{C}^\times) = \mathbb{H}^0(\mathfrak{g}, Z(\hat{G}) \to 1) = H^1(\mathfrak{g}, Z(\hat{G})). \tag{1.3.4.1}$$

On the other hand, the short exact sequence

$$0 \to \mathbf{Z} \to \mathbf{C} \to \mathbf{C}^\times \to 1$$

induces a short exact sequence of complexes

$$0 \to L^\bullet \to L^\bullet \otimes \mathbf{C} \to L^\bullet \otimes \mathbf{C}^\times \to 1$$

and a hypercohomology exact sequence

$$\mathbb{H}^0(\mathfrak{g}, L^\bullet \otimes \mathbf{C}) \to \mathbb{H}^0(\mathfrak{g}, L^\bullet \otimes \mathbf{C}^\times) \to \mathbb{H}^1(\mathfrak{g}, L^\bullet) \to \mathbb{H}^1(\mathfrak{g}, L^\bullet \otimes \mathbf{C}). \tag{1.3.4.2}$$

Content-Length: 21892 X-Lines: 687 Status: O

We prove that $\mathbb{H}^0(\mathfrak{g}, L^\bullet \otimes \mathbf{C}) = 0$ and $\mathbb{H}^1(\mathfrak{g}, L^\bullet \otimes \mathbf{C}) = 0$. Let $T^{\mathrm{ss}} = T \cap G^{\mathrm{ss}}$. Then $\mathbf{X}^*(T^{\mathrm{ss}})$ is a subgroup of finite index of $\mathbf{X}^*(T^{\mathrm{sc}})$, and so $\mathbf{X}^*(T^{\mathrm{sc}}) \otimes \mathbf{C} = \mathbf{X}^*(T^{\mathrm{ss}}) \otimes \mathbf{C}$. We see that

$$L^\bullet \otimes \mathbf{C} = (\mathbf{X}^*(T) \otimes \mathbf{C} \to \mathbf{X}^*(T^{\mathrm{ss}}) \otimes \mathbf{C}).$$

It follows that $L^\bullet \otimes \mathbf{C}$ is quasi-isomorphic to $(\mathbf{X}^*(G^{\mathrm{tor}}) \otimes \mathbf{C} \to 0)$. Hence

$$\mathbb{H}^0(\mathfrak{g}, L^\bullet \otimes \mathbf{C}) = \mathbb{H}^0(\mathfrak{g}, (\mathbf{X}^*(G^{\mathrm{tor}}) \otimes \mathbf{C} \to 0)) = H^1(\mathfrak{g}, \mathbf{X}^*(G^{\mathrm{tor}}) \otimes \mathbf{C}).$$

But $H^1(\mathfrak{g}, \mathbf{X}^*(G^{\mathrm{tor}}) \otimes \mathbf{C}) = 0$ because $\mathbf{X}^*(G^{\mathrm{tor}}) \otimes \mathbf{C}$ is a uniquely divisible group. Thus $\mathbb{H}^0(\mathfrak{g}, L^\bullet \otimes \mathbf{C}) = 0$. Similarly $\mathbb{H}^1(\mathfrak{g}, L^\bullet \otimes \mathbf{C}) = 0$. From exact sequence (1.3.4.2) we then obtain

$$\mathbb{H}^0(\mathfrak{g}, L^\bullet \otimes \mathbf{C}^\times) = \mathbb{H}^1(\mathfrak{g}, L^\bullet).$$

We see from (1.3.4.1) that

$$H^1(\mathfrak{g}, Z(\hat{G})) = \mathbb{H}^1(\mathfrak{g}, L^\bullet).$$

Similarly, $H^1(\gamma, Z(\hat{G})) = \mathbb{H}^1(\gamma, L^\bullet)$ for every closed subgroup $\gamma \subset \mathfrak{g}$. We conclude that

$$Ш^1_\omega(\mathfrak{g}, L^\bullet) = Ш^1_\omega(\mathfrak{g}, Z(\hat{G})). \quad \square$$

1.3.5. Remark. Proposition 1.3.2 shows that $Ш^1_\omega(\mathfrak{g}, L^\bullet)$ does not depend on the choice of T. Indeed, it only depends on the algebraic fundamental group $\pi_1(G)$ which does not depend on T ([B], 1.2).

2. Main results

Theorem 2.1. *Let k be a field of characteristic zero, $\mathfrak{g} = \mathrm{Gal}(\bar{k}/k)$, G a connected linear algebraic k-group, V a smooth k-compactification of G, $T \subset G$ a maximal k-torus, $L^\bullet = (\mathbf{X}^*(T) \to \mathbf{X}^*(T^{\mathrm{sc}}))$. Then the group $H^1(k, \mathrm{Pic}\,\overline{V})$ is canonically isomorphic to $Ш^1_\omega(\mathfrak{g}, L^\bullet)$, and this isomorphism is functorial in G.*

Corollary 2.2. *With the notation of Theorem* 2.1*, there is an injection*

$$\mathrm{Br}\, V/\mathrm{Br}\, k \hookrightarrow Ш^1_\omega(\mathfrak{g}, L^\bullet)$$

which is an isomorphism provided either $V(k) \neq \varnothing$*, or* $H^3(k, \bar{k}^\times) = 0$.

Proof. By 1.1.1 G is rational, hence V is projective and rational, and $\mathrm{Br}\, \overline{V} = 0$, see 1.2.3. The corollary now follows from 1.2.3 and Theorem 2.1. □

2.3. Proof of Theorem 2.1. We first assume that G is reductive and G^{ss} is simply connected. We use Voskresenskiĭ's exact sequence

$$0 \to \mathbf{X}^*(G) \to P \to \mathrm{Pic}\, \overline{V} \to \mathrm{Pic}\, \overline{G} \to 0 \tag{2.3.1}$$

which is valid for any connected k-group G [V] (see also [S], (9.0.0)). Here P is a permutation $\mathfrak{g}$-module. The exact sequence of $\bar{k}$-groups

$$1 \to \overline{G}^{\mathrm{ss}} \to \overline{G} \to \overline{G}^{\mathrm{tor}} \to 1$$

induces the exact sequence ([S], (6.11.4))

$$0 \to \mathbf{X}^*(G^{\mathrm{tor}}) \to \mathbf{X}^*(G) \to \mathbf{X}^*(G^{\mathrm{ss}}) \to \mathrm{Pic}\, \overline{G}^{\mathrm{tor}} \to \mathrm{Pic}\, \overline{G} \to \mathrm{Pic}\, \overline{G}^{\mathrm{ss}} \to 0.$$

Since $\overline{G}^{\mathrm{ss}}$ is simply connected, we have $\mathrm{Pic}\, \overline{G}^{\mathrm{ss}} = 0$; since G^{ss} is semisimple, we have $\mathbf{X}^*(G^{\mathrm{ss}}) = 0$; since G^{tor} is a torus, we have $\mathrm{Pic}\, \overline{G}^{\mathrm{tor}} = 0$, cf. [S], 6.9. We conclude that $\mathrm{Pic}\, \overline{G} = 0$ and $\mathbf{X}^*(G) = \mathbf{X}^*(G^{\mathrm{tor}})$. Exact sequence (2.3.1) is thus reduced to

$$0 \to \mathbf{X}^*(G^{\mathrm{tor}}) \to P \to \mathrm{Pic}\, \overline{V} \to 0. \tag{2.3.2}$$

We can now use the following fundamental property of the Picard group $\mathrm{Pic}\, \overline{V}$ proved in [CT/K], Prop. 3.2: $H^1(\gamma, \mathrm{Pic}\, \overline{V}) = 0$ for all closed procyclic subgroups $\gamma \subset \mathfrak{g}$. Since P is a permutation $\mathfrak{g}$-module, we obtain

$$H^1(\mathfrak{g}, \mathrm{Pic}\, \overline{V}) = Ш^2_\omega(\mathfrak{g}, \mathbf{X}^*(G^{\mathrm{tor}})) \tag{2.3.3}$$

by [CT/S2], Prop. 9.5(ii).

We now prove that $Ш^2_\omega(\mathfrak{g}, \mathbf{X}^*(G^{\mathrm{tor}})) \cong Ш^1_\omega(\mathfrak{g}, L^\bullet)$. Since G^{ss} is simply connected, one has an exact sequence of tori

$$1 \to T^{\mathrm{sc}} \to T \to G^{\mathrm{tor}} \to 1$$

and the dual exact sequence of character groups

$$0 \to \mathbf{X}^*(G^{\mathrm{tor}}) \to \mathbf{X}^*(T) \to \mathbf{X}^*(T^{\mathrm{sc}}) \to 0.$$

It induces a morphism of complexes

$$(\mathbf{X}^*(G^{\mathrm{tor}}) \to 0) \to L^\bullet$$

which is a quasi-isomorphism. Thus

$$H^{i+1}(\gamma, \mathbf{X}^*(G^{\mathrm{tor}})) = \mathbb{H}^i(\gamma, L^\bullet)$$

for every natural i and every closed subgroup $\gamma \subseteq \mathfrak{g}$. We conclude that

$$Ш^2_\omega(\mathfrak{g}, \mathbf{X}^*(G^{\mathrm{tor}})) \cong Ш^1_\omega(\mathfrak{g}, L^\bullet).$$

Thus $H^1(\mathfrak{g}, \operatorname{Pic} \overline{V}) \cong Ш^1_\omega(\mathfrak{g}, L^\bullet)$, and this isomorphism is functorial in G. The theorem is proved for reductive G with G^{ss} simply connected.

Let now G be an arbitrary connected reductive k-group. Let $H \overset{\alpha}{\to} G$ be a z-extension with kernel Z. Let V_G be a a smooth compactification of G and let V_H be a smooth compactification of H. We have a homomorphism

$$\alpha_*\colon H^1(k, \operatorname{Pic} \overline{V}_H) \to H^1(k, \operatorname{Pic} \overline{V}_G).$$

We prove that α_* is an isomorphism, hence

$$H^1(k, \operatorname{Pic} \overline{V}_H) \cong H^1(k, \operatorname{Pic} \overline{V}_G). \tag{2.3.4}$$

Since Z is a quasi-trivial torus, the map α admits a rational k-section $s\colon G \to H$. Indeed, the obstruction to the existence of such a section lies in $H^1(k(G), Z) = 0$. The rational section s gives rise to a biregular k-isomorphism $i\colon U_H \to U_G \times Z$, where $U_H \subset H$ and $U_G \subset G$ are open k-subvarieties, U_G is an open subvariety on which s is defined, and $U_H = s(U_G) \cdot Z$. The projections are defined as follows: $\mathrm{pr}_{U_G}(h) = \alpha(h)$, $\mathrm{pr}_Z(h) = h \cdot s(\alpha(h))^{-1}$, where $h \in U_H$. Since Z is a quasi-trivial torus, it is k-rational, and by 1.2.2 we obtain a canonical isomorphism $H^1(k, \operatorname{Pic} \overline{V}(U_H)) \xrightarrow{\sim} H^1(k, \operatorname{Pic} \overline{V}(U_G))$ induced by the projection $\mathrm{pr}_{U_G}\colon U_H \to U_G$. This gives us (2.3.4), and we see that (2.3.4) is the canonical isomorphism induced by $\alpha\colon H \to G$. It does not depend on s.

Let $T_G \subset G$ be a maximal torus and set $L^\bullet_G = (\mathbf{X}^*(T_G) \to \mathbf{X}^*(T^{\mathrm{sc}}_G))$. Set $T_H = \alpha^{-1}(T) \subset H$, $L^\bullet_H = (\mathbf{X}^*(T_H) \to \mathbf{X}^*(T^{\mathrm{sc}}_H))$. We prove that

$$Ш^1_\omega(k, L^\bullet_G) \cong Ш^1_\omega(k, L^\bullet_H). \tag{2.3.5}$$

We note that the z-extension $H \to G$ induces an exact sequence of complexes

$$1 \to L^\bullet_G \to L^\bullet_H \to (\mathbf{X}^*(Z) \to 1) \to 1$$

which leads to the following commutative diagram:

$$\begin{array}{ccccccccc}
 & & & & 0 & & 0 & & 0 \\
 & & & & \downarrow & & \downarrow & & \| \\
0 & = & Ш^1_\omega(\mathfrak{g}, \mathbf{X}^*(Z)) & \longrightarrow & Ш^1_\omega(\mathfrak{g}, L^\bullet_G) & \longrightarrow & Ш^1_\omega(\mathfrak{g}, L^\bullet_H) & \longrightarrow & Ш^2_\omega(\mathfrak{g}, \mathbf{X}^*(Z)) \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
0 & = & H^1(\mathfrak{g}, \mathbf{X}^*(Z)) & \longrightarrow & \mathbb{H}^1(\mathfrak{g}, L^\bullet_G) & \longrightarrow & \mathbb{H}^1(\mathfrak{g}, L^\bullet_H) & \longrightarrow & H^2(\mathfrak{g}, \mathbf{X}^*(Z)) \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
0 & = & \prod_\gamma H^1(\gamma, \mathbf{X}^*(Z)) & \longrightarrow & \prod_\gamma \mathbb{H}^1(\gamma, L^\bullet_G) & \longrightarrow & \prod_\gamma \mathbb{H}^1(\gamma, L^\bullet_H) & \longrightarrow & \prod_\gamma H^2(\gamma, \mathbf{X}^*(Z)).
\end{array}$$

In this diagram, the middle and the bottom rows are exact, and the terms in the left column are all zero since $\mathbf{X}^*(Z)$ is a permutation $\mathfrak{g}$-module. The group $Ш^2_\omega(\mathfrak{g}, \mathbf{X}^*(Z))$ is zero by the same reason (see 1.2.1). By diagram chasing one can easily prove that the top row is exact. We have proved isomorphism (2.3.5).

We have already proved that $H^1(k, \operatorname{Pic} \overline{V}_H) \cong Ш^1_\omega(k, L^\bullet_H)$, because H^{ss} is simply connected. Together with isomorphisms (2.3.4) and (2.3.5) we obtain an isomorphism

$$H^1(k, \operatorname{Pic} \overline{V}_G) \cong Ш^1_\omega(k, L^\bullet_G). \tag{2.3.6}$$

Using Lemma 1.1.4(2), one can easily prove that isomorphism (2.3.6) does not depend on the choice of a z-extension $H \to G$. Using Lemma 1.1.4(3), one can easily check that isomorphism (2.3.6) is functorial in G. This establishes the theorem for reductive k-groups.

Let now G be an arbitrary connected k-group. Let V_G be a smooth compactification of G and let $V_{G^{\mathrm{red}}}$ be a smooth compactification of G^{red}. By 1.1.2, there is an isomorphism of k-varieties $G \cong G^{\mathrm{red}} \times G^{\mathrm{u}}$ where G^{u} is k-rational; by 1.2.2, we then obtain

$$H^1(\mathfrak{g}, \operatorname{Pic} \overline{V}_G) \cong H^1(\mathfrak{g}, \operatorname{Pic} \overline{V}_{G^{\mathrm{red}}}). \tag{2.3.7}$$

Let the complex $L^\bullet$ be defined as in the introduction, i.e. in terms of G^{red}. Since G^{red} is reductive, we have already proved that $H^1(\mathfrak{g}, \operatorname{Pic} \overline{V}_{G^{\mathrm{red}}}) \cong Ш^1_\omega(\mathfrak{g}, L^\bullet)$. Together with (2.3.7) we obtain

$$H^1(\mathfrak{g}, \operatorname{Pic} \overline{V}_G) \cong Ш^1_\omega(\mathfrak{g}, L^\bullet).$$

This isomorphism is functorial in G. $\square$

Theorem 2.4. *Let k be a field of characteristic zero, $\mathfrak{g} = \operatorname{Gal}(\bar{k}/k)$, G a connected linear algebraic k-group, E a G-torsor, V a smooth compactification of E, $T \subset G$ a maximal k-torus, $L^\bullet = (\mathbf{X}^*(T) \to \mathbf{X}^*(T^{\mathrm{sc}}))$. Then the group $H^1(k, \operatorname{Pic} \overline{V})$ is isomorphic to $Ш^1_\omega(\mathfrak{g}, L^\bullet)$.*

2.5. Proof of Theorem 2.4. First assume that G is reductive and G^{ss} is simply connected. If $E = G$, Theorem 2.4 coincides with Theorem 2.1. If E has no rational points, one just has to make use of the device of passage to the generic point (see [CT/S3], App. 2B, p. 462–463; cf. [CT/K], proof of Th. 4.1). We reproduce here this argument adapted to our setting. Denote $K = k(E)$, $L = \bar{k}(E)$, and let M be an algebraic closure of K containing L. We have $\operatorname{Gal}(L/K) = \operatorname{Gal}(\bar{k}/k) = \mathfrak{g}$. Let $\mathfrak{g}_1 = \operatorname{Gal}(M/K)$, $\mathfrak{h} = \operatorname{Gal}(M/L)$: $\mathfrak{h}$ is a normal subgroup of $\mathfrak{g}_1$, and $\mathfrak{g} = \mathfrak{g}_1/\mathfrak{h}$. Since $\overline{V}$ is a proper smooth rational variety, the natural inclusions of free abelian groups of finite rank

$$\operatorname{Pic}(V \times_k \bar{k}) \hookrightarrow \operatorname{Pic}(V \times_k L) \hookrightarrow \operatorname{Pic}(V \times_k M)$$

are in fact equalities. Denote this abelian group by $\operatorname{Pic} \overline{V}$. The group $\mathfrak{h}$ acts trivially on $\operatorname{Pic}(V \times_k M)$. We write down the restriction-inflation exact sequence for the extensions $M/L/K$:

$$0 \to H^1(\mathfrak{g}, \operatorname{Pic} \overline{V}) \overset{\inf}{\to} H^1(\mathfrak{g}_1, \operatorname{Pic} \overline{V}) \overset{\mathrm{res}}{\to} H^1(\mathfrak{h}, \operatorname{Pic} \overline{V}) = 0.$$

This gives an isomorphism

$$H^1(\operatorname{Gal}(\bar{k}/k), \operatorname{Pic}(V \times_k \bar{k})) \cong H^1(\operatorname{Gal}(M/K), \operatorname{Pic}(V \times_k M)). \tag{2.5.1}$$

Recall that G is reductive and G^{ss} is simply connected. The K-variety $V\times_k K$ is a smooth compactification of the torsor $E\times_k K$ which has a K-point and is hence isomorphic to $G\times_k K$. Formula (2.3.3) and isomorphism (2.5.1) then show that there is an isomorphism

$$H^1(\mathrm{Gal}(\bar{k}/k),\mathrm{Pic}\,(V\times_k\bar{k}))\cong Ш^2_\omega(\mathrm{Gal}(M/K),\mathbf{X}^*(G^{\mathrm{tor}})). \tag{2.5.2}$$

Since $\mathfrak{h}=\mathrm{Gal}(M/L)$ acts trivially on $\mathbf{X}^*(G^{\mathrm{tor}})$, and $\mathbf{X}^*(G^{\mathrm{tor}})$ is a torsion-free module, we can use the equality $H^1(\mathfrak{h},\mathbf{X}^*(G^{\mathrm{tor}}))=0$ and write down the restriction-inflation exact sequence for H^2:

$$0\to H^2(\mathfrak{g},\mathbf{X}^*(G^{\mathrm{tor}}))\overset{\mathrm{inf}}{\to}H^2(\mathfrak{g}_1,\mathbf{X}^*(G^{\mathrm{tor}}))\overset{\mathrm{res}}{\to}H^2(\mathfrak{h},\mathbf{X}^*(G^{\mathrm{tor}})).$$

Since $Ш^2_\omega(\mathfrak{h},\mathbf{X}^*(G^{\mathrm{tor}}))=0$, we obtain

$$Ш^2_\omega(\mathrm{Gal}(\bar{k}/k),\mathbf{X}^*(G^{\mathrm{tor}}))\cong Ш^2_\omega(\mathrm{Gal}(M/K),\mathbf{X}^*(G^{\mathrm{tor}})). \tag{2.5.3}$$

Putting together isomorphisms (2.5.2) and (2.5.3), we obtain $H^1(\mathfrak{g},\mathrm{Pic}\,\overline{V})\cong Ш^2_\omega(\mathfrak{g},\mathbf{X}^*(G^{\mathrm{tor}}))$. Since G^{ss} is simply connected, we have $Ш^2_\omega(\mathfrak{g},\mathbf{X}^*(G^{\mathrm{tor}}))\cong Ш^1_\omega(\mathfrak{g},L^\bullet)$. Thus $H^1(\mathfrak{g},\mathrm{Pic}\,\overline{V})\cong Ш^1_\omega(\mathfrak{g},L^\bullet)$, and we obtain the theorem for G reductive with G^{ss} simply connected.

Let now G be an arbitrary reductive group and E_G a principal homogeneous space of G. By Lemma 1.1.4(1), there exists a z-extension $\alpha\colon H\to G$ with kernel Z such that the class $\mathrm{Cl}(E_G)\in H^1(k,G)$ is the image of some $\mathrm{Cl}(E_H)\in H^1(k,H)$ where E_H is a principal homogeneous space of H. The cohomology map $\alpha_*\colon H^1(k,H)\to H^1(k,G)$ is represented by the map $E_H\mapsto E_H/Z$. We may therefore assume that $E_G=E_H/Z$. The canonical projection $E_H\to E_G=E_H/Z$ admits a k-rational section, because Z is a quasi-trivial torus. Indeed, the obstruction to the existence of such a section lies in $H^1(k(E_G),Z)$, and this cohomology group is zero by Hilbert 90. Therefore we have a birational isomorphism $f\colon E_G\times Z\to E_H$ given by $f(x,z)=s(x)\cdot z$ where $x\in E_G$ and $z\in Z$. The quasi-trivial torus Z is k-rational. By 1.2.2 the birational isomorphism f gives an isomorphism

$$H^1(\mathrm{Gal}(\bar{k}/k),\mathrm{Pic}\,(V_G\times_k\bar{k}))\cong H^1(\mathrm{Gal}(\bar{k}/k),\mathrm{Pic}\,(V_H\times_k\bar{k}))$$

where V_G (resp. V_H) stands for a smooth compactification of E_G (resp. E_H). Since H^{ss} is simply connected, by the preceding part of the proof we have

$$H^1(\mathrm{Gal}(\bar{k}/k),\mathrm{Pic}\,(V_H\times_k\bar{k}))\cong Ш^2_\omega(\mathrm{Gal}(\bar{k}/k),\mathbf{X}^*(H^{\mathrm{tor}})).$$

As shown in the proof of Theorem 2.1, $Ш^2_\omega(\mathrm{Gal}(\bar{k}/k),\mathbf{X}^*(H^{\mathrm{tor}}))$ is isomorphic to $Ш^1_\omega(\mathfrak{g},L^\bullet)$. Thus $H^1(\mathfrak{g},\mathrm{Pic}\,V_G)\cong Ш^1_\omega(\mathfrak{g},L^\bullet)$. This proves the theorem for any reductive group.

Let now G be an arbitrary (not necessarily reductive) connected k-group. The canonical homomorphism $r\colon G\to G^{\mathrm{red}}$ induces a bijection of Galois cohomology pointed sets $r_*\colon H^1(k,G)\to H^1(k,G^{\mathrm{red}})$, cf. [S], 1.13. The map r_* is represented by the map of torsors $E\mapsto E/G^{\mathrm{u}}$, where E is a torsor under G and E/G^{u} is a torsor under $G/G^{\mathrm{u}}=G^{\mathrm{red}}$. We wish to prove that

$$H^1(k,\mathrm{Pic}\,\overline{V}_E)\cong H^1(k,\mathrm{Pic}\,\overline{V}_{E/G^{\mathrm{u}}}),$$

where V_E and $V_{E/G^{\mathrm{u}}}$ are smooth compactifications of E and E/G^{u}, respectively. Since our functor $Ш^1_\omega(\mathfrak{g}, L^\bullet)$ is, by definition, the same for G and for G^{red}, this will prove the theorem.

We fix a Levi decomposition $G^{\mathrm{u}} \rtimes G^{\mathrm{red}} \xrightarrow{\sim} G$. It defines a natural homomorphism $\varphi\colon G^{\mathrm{red}} \to G$ and a map $\varphi_*\colon H^1(k, G^{\mathrm{red}}) \to H^1(k, G)$, inverse to r_*. We want to describe φ_* in terms of torsors.

Let X be a torsor under G^{red}. Set $Y = X \times G^{\mathrm{u}}$. We define a right action of $G = G^{\mathrm{red}} \ltimes G^{\mathrm{u}}$ on Y by

$$(x, v) \cdot (g, u) = (x \cdot g, v^g \cdot u),$$

where $x \in X$, $v, u \in G^{\mathrm{u}}$, $g \in G^{\mathrm{red}}$, and v^g refers to the right action of G^{red} on G^{u} defined by the Levi decomposition. One can easily check that this is a well-defined action and that the map $X \mapsto Y$ represents φ_*.

Since $Y = X \times G^{\mathrm{u}}$ and G^{u} is k-rational, by 1.2.2 we have $H^1(k, \operatorname{Pic} \overline{V}(Y)) \cong H^1(k, \operatorname{Pic} \overline{V}(X))$. Since φ_* is inverse to r_*, any torsor E of G is isomorphic to Y for $X = E/G^{\mathrm{u}}$. We obtain $H^1(k, \operatorname{Pic} \overline{V}_E) \cong H^1(k, \operatorname{Pic} \overline{V}_{E/G^{\mathrm{u}}})$. This proves the theorem. $\square$

3. Comments and remarks

3.1. If $G = T$ is a torus, we have $Ш^1_\omega(\mathfrak{g}, L^\bullet) = Ш^2_\omega(\mathfrak{g}, \mathbf{X}^*(T))$, and the formula of Theorem 2.1 reduces to

$$H^1(k, \operatorname{Pic} \overline{V}) \cong Ш^2_\omega(k, \mathbf{X}^*(T));$$

cf. [S], Prop. 9.8, in the number field case, [CT/S2] for $E = T$ over an arbitrary field, and [CT/K] in general.

3.2. If G is a semisimple group, we have $Ш^1_\omega(\mathfrak{g}, L^\bullet) = Ш^1_\omega(\mathfrak{g}, \mathbf{X}^*(B))$ where $B = \ker[G^{\mathrm{sc}} \to G]$ is the fundamental group of G, and the formula of Theorem 2.1 reduces to

$$H^1(k, \operatorname{Pic} \overline{V}) \cong Ш^1_\omega(k, \mathbf{X}^*(B)); \tag{3.2.1}$$

cf. [S], 9.6, in the number field case and [CT/K] in general.

3.3. Let now G be a reductive group admitting a special covering $\mu\colon G_0 \times S \to G$ with kernel B, where G_0 is a simply connected group, S is a quasi-trivial torus, and B is a finite group. We show that in this case Theorem 2.1 reduces to formula (3.2.1), cf. [S], Prop. 9.8, in the number field case and [CT/K] in general. Recall that T is a maximal torus of G, $L^\bullet = (\mathbf{X}^*(T) \to \mathbf{X}^*(T^{\mathrm{sc}}))$. We have to prove that

$$Ш^1_\omega(k, \mathbf{X}^*(B)) = Ш^1_\omega(\mathfrak{g}, L^\bullet).$$

Write $\mu^{-1}(T) = T_0 \times S$, where T_0 is a maximal torus of G_0. We have $G^{\mathrm{sc}} = G_0$, $T^{\mathrm{sc}} = T_0$. Set $L^\bullet_1 = (\mathbf{X}^*(T) \to \mathbf{X}^*(T_0) \times \mathbf{X}^*(S))$. Consider an exact sequence

$$1 \to B \to T_0 \times S \to T \to 1.$$

We see that the complexes $(B \to 1)$ and $(T_0 \times S \to T)$ are quasi-isomorphic. Hence the complexes $(0 \to \mathbf{X}^*(B))$ and $L^\bullet_1$ are also quasi-isomorphic, and therefore

$$Ш^1_\omega(k, \mathbf{X}^*(B)) = Ш^1_\omega(\mathfrak{g}, L^\bullet_1).$$

We now consider a short exact sequence of complexes

$$0 \to (0 \to \mathbf{X}^*(S)) \to L_1^\bullet \to L^\bullet \to 0.$$

It induces the following commutative diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & Ш^1_\omega(\mathfrak{g}, L_1^\bullet) & \longrightarrow & Ш^1_\omega(\mathfrak{g}, L^\bullet) & \longrightarrow & Ш^2_\omega(\mathfrak{g}, \mathbf{X}^*(S)) \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
0 = H^1(\mathfrak{g}, \mathbf{X}^*(S)) & \longrightarrow & \mathbb{H}^1(\mathfrak{g}, L_1^\bullet) & \longrightarrow & \mathbb{H}^1(\mathfrak{g}, L^\bullet) & \longrightarrow & H^2(\mathfrak{g}, \mathbf{X}^*(S)) \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
0 = \prod_\gamma H^1(\gamma, \mathbf{X}^*(S)) & \longrightarrow & \prod_\gamma \mathbb{H}^1(\gamma, L_1^\bullet) & \longrightarrow & \prod_\gamma \mathbb{H}^1(\gamma, L^\bullet) & \longrightarrow & \prod_\gamma H^2(\gamma, \mathbf{X}^*(S))
\end{array}$$

The middle and the bottom rows are exact. Since S is a quasi-trivial torus, we have $Ш^2_\omega(\mathfrak{g}, \mathbf{X}^*(S)) = 0$. By diagram chasing, one can show that the top row of the diagram is also exact. Thus

$$Ш^1_\omega(k, \mathbf{X}^*(B)) = Ш^1_\omega(\mathfrak{g}, L_1^\bullet) = Ш^1_\omega(\mathfrak{g}, L^\bullet),$$

as required. □

References

[**B**] M. Borovoi, *Abelian Galois cohomology of reductive groups*, Mem. Amer. Math. Soc. **132** (1998), no. 626, 1–50.

[**C**] C. Chevalley, *On algebraic group varieties*, J. Math. Soc. Japan **6** (1954), 303–324.

[**CT/K**] J.-L. Colliot-Thélène et B. È. Kunyavskiĭ, *Groupe de Brauer non ramifié des espaces principaux homogènes de groupes linéaires*, J. Ramanujan Math. Soc. **13** (1998), 37–49.

[**CT/S1**] J.-L. Colliot-Thélène et J.-J. Sansuc, *La R-équivalence sur les tores*, Ann. Sci. École Norm. Sup. (4) **10** (1977), 175–229.

[**CT/S2**] J.-L. Colliot-Thélène and J.-J. Sansuc, *Principal homogeneous spaces under flasque tori: applications*, J. Algebra **106** (1987), 148–205.

[**CT/S3**] J.-L. Colliot-Thélène et J.-J. Sansuc, *La descente sur les variétés rationnelles,* II, Duke Math. J. **54** (1987), 375–492.

[**H**] H. Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero,* I, II, Ann. of Math. (2) **79** (1964), 109-326.

[**K1**] R. E. Kottwitz, *Rational conjugacy classes in reductive groups*, Duke Math. J. **49** (1982), 785–806.

[**K2**] R. E. Kottwitz, *Stable trace formula: Cuspidal tempered terms*, Duke Math. J. **51** (1984), 611–650.

[**K3**] R. E. Kottwitz, *Stable trace formula: Elliptic singular terms*, Math. Ann. **275** (1986), 365–399.

[**L**] R. P. Langlands, *Stable conjugacy: definitions and lemmas*, Canad. J. Math. **31** (1979), 700–725.

[**M/S**] J. S. Milne and K.-y. Shih, *Conjugates and Shimura varieties*, Hodge Cycles, Motives and Shimura Varieties, Lecture Notes in Math. **900**, Springer, Berlin, 1982, pp. 280–356.

[**S**] J.-J. Sansuc, *Groupe de Brauer et arithmétique des groupes algébriques linéaires sur un corps de nombres*, J. Reine Angew. Math. **327** (1981), 12–80.

[**V**] V. E. Voskresenskiĭ, *Birational properties of linear algebraic groups*, Math. USSR Izv. **4** (1970), 1–17.

Borovoi: Raymond and Beverly Sackler School of Mathematical Sciences, Tel Aviv University, 69978 Tel Aviv, Israel
E-mail address: borovoi@math.tau.ac.il

Kunyavskiĭ: Department of Mathematics and Computer Science, Bar-Ilan University, 52900 Ramat Gan, Israel
E-mail address: kunyav@macs.biu.ac.il