

**ASSOCIATE FORMS, JOINS, MULTIPLICITIES AND  
AN INTRINSIC ELIMINATION THEORY**

by

Federico Gaeta

Preprint – to appear in definitive form in

**TOPICS IN ALGEBRA**

Banach Center Publications, volume 26

PWN – Polish Scientific Publishers, Warsaw.

Max–Planck–Institut  
für Mathematik  
Gottfried–Claren–Str. 26  
5300 Bonn 3  
Federal Republic of Germany

Universidad Complutense  
de Madrid  
Spain



## NOTATIONS

	Occurrence page
$V, W, \dots, V_d = V^c, W_d = W^c \dots, c = n - d$ algebraic subvarieties $\subset \mathbb{P}_n(\mathbb{C})$ of dimension $d$ and codimension $c$ in $\mathbb{P}_n(\mathbb{C})$	5
$\mathbb{P}_n = \mathbb{P}(E_{n+1}), E_{n+1}(n+1)$ dimensional complex vector space	7
$\Pi : V^{(1)} \times V^{(2)} \times \dots \times W^{(h)}$ Cartesian product	
$\delta : V \longrightarrow \Pi = V^h$ diagonal map, (0.8)	8
$\Sigma$ diagonal manifold $\subset \mathbb{P}_n^{(1)} \times \dots \times \mathbb{P}_n^{(h)}$ ,	
$\Sigma = \mathbb{P}\{(x \otimes n) \mid x \in E_{n+1} - 0, n \in \mathbb{C}^h - 0\}$	
$\Delta$ diagonal space $\mathbb{P}(E \otimes (1,1, \dots, 1)) \subset \Sigma \subset \mathbb{P}_n^{(1)} \times \dots \times \mathbb{P}_n^{(h)}$	
$\mathcal{J}, J, J(V \times W), J(V^{(1)} \times \dots \times V^{(h)})$ join of shown varieties Abstract DEF. 1.1	
$\mathcal{J}$ Prejoin	
$J = \overline{J}_p = \bigcup_{\mathbb{P}_{n-1} \in \mathcal{J}} \mathbb{P}_{n-1}$ Full join (2.3)	25
$\mathcal{C}(V^c) = \{\mathbb{P}_{c-1} \subset \mathbb{P}_n \mid \mathbb{P}_{c-1} \cdot V^c \neq \emptyset\}$ , DEF. 0.1, (0.13)	9
$S, Y, N$ the SEVERI CHOW or WEIL-SIEGEL associated form to a $V_d = V^c \subset \mathbb{P}_n$	10
$S = S(x_1, x_2, \dots, x_c), Y = Y(u_1, u_2, \dots, u_{d+1})$	
$N = N(u_1, \dots, u_{d+2}; x) x_j \in E, u_j \in \check{E}$ formulas (0.10), (0.11), (0.12)	
$S(x_1, x_2, \dots, x_c) = \mathbf{S}(x_1 \wedge \dots \wedge x_c)$	40
$x_1 \wedge \dots \wedge x_c \in \overset{\circ}{\wedge} E$ (6.1)	
$Y(u_1, u_2, \dots, u_{d+1}) = Y(u_1 \wedge \dots \wedge u_{d+1})$	
$u_1 \wedge \dots \wedge u_{d+2} \in \overset{d+1}{\wedge} \check{E}$ (6.2)	
$N(u_1, u_2, \dots, u_{d+2}; x) = \mathbb{N}(\langle u_1, x \rangle, \dots, \langle u_{d+2}, x \rangle) = 0$ (6.6)	42

ASSOCIATE FORMS, JOINS, MULTIPLICITIES AND AN INTRINSIC  
ELIMINATION THEORY

by

FEDERICO GAETA

Universidad Complutense de Madrid, Spain

TABLE OF CONTENTS

	Page
NOTATIONS	1
ABSTRACT	4
0. INTRODUCTION	6
I. GENERALITIES ON JOINS	16
1. THE "REDUCTION TO THE DIAGONAL". A PROJECTIVE VERSION	16
2. RECALL OF THE JOIN OF $h$ VARIETIES. RELATION WITH SEGRE'S MODEL OF THE PRODUCT $V^{(1)} \times V^{(2)} \times \dots \times V^{(h)}$	21
3. CASE $n_1 = n_2 = \dots = n_h = n$ . THE DIAGONALS $\Sigma$ , $\Delta$ .	28
4. JOINS AND $h$ -COLLINEATIONS	31

	Page
II. GENERALITIES ON THE COMPLEX $\mathfrak{C}(V)$ ATTACHED TO A $V_c \subset \mathbb{P}_n$ .	
5. THE COMPLEX $\mathfrak{C}(V^c) = \{\mathbb{P}_{c-1} \subset \mathbb{P}_n \mid \mathbb{P}_{c-1} \cap V^c \neq \emptyset\}$	35
6. REVIEW ON ASSOCIATE FORMS	36
III. APPLICATIONS	
FIRST PART: $c \leq n$	
THE EXPONENT INTERSECTION MULTIPLICITY	
7. THE RESTRICTION TO THE DIAGONAL OF $\mathfrak{C}(J(V^{(1)} \times \dots \times V^{(h)}))$ , $c \leq n$	41
8. COMPUTATION OF $F_j$ . BEZOUT'S THEOREM	46
9. ON THE PROOF OF THE THEOREMS	47
10. EQUIVALENCE OF THE EXPONENT MULTIPLICITY WITH VAN DER WAERDEN'S THEORY	47
11. BEZOUT'S THEOREM WITH A NEW DEGENERATION METHOD	50
SECOND PART: $c > n$	
12. A GEOMETRICAL THEORY FOR RESULTANT SYSTEMS	53

	Page
AN INTRINSIC ELIMINATION THEORY	62
13. HISTORICAL APPROACH	62
14. INTRINSIC ELIMINATION THEORY USING WEIL-SIEGEL FORMS	67
BIBLIOGRAPHY	71

A B S T R A C T

*The first time a mathematician hears about "multiplicity"  $m_r(\in \mathbb{Z}^+)$  refers to an  $m_r$  - ple root  $r$  of a polynomial  $f(x)$  or binary form  $\phi(x_0, x_1)$  :*

$$(0.0) \quad f(x) = a_0 \prod_{r \in \mathbb{C}} (x - r)^{m_r} \quad m_r \in \mathbb{Z} \mid m_r \geq 0$$

$$\phi(x_0, x_1) = \prod_{r \in \mathbb{P}_1(\mathbb{C})} \begin{vmatrix} x_0 & x_1 \\ r_0 & r_1 \end{vmatrix}^{m_r}$$

$x - r$  (or  $\begin{vmatrix} x_0 & x_1 \\ r_0 & r_1 \end{vmatrix}$ ) is the trivial associate form (a.f) cf. DEF. 4.6 of the point  $r$  (with affine (or projective) coordinates in the complex affine (or projective) line). It is natural to ask whether or not this exponent is also the natural intersection multiplicity of an irreducible component  $I$  in the proper intersection  $V \cap W$  ( $V, W$  irreducible a.v in  $\mathbb{P}_n(\mathbb{C}) = \mathbb{P}(E)$ ). An affirmative answer is found in [vdW1] only for two irreducible plane

curves. This idea of the exponent intersection multiplicity is developed in this paper in the general case by showing that the Form

$$F_{V \cdot W} = \prod F_I^{m_I} \quad (I \text{ proper irreducible component of } V \cap W)$$

can be computed by restriction of the  $F_J$  (associated to the Join  $J(V \times W) \subset \mathbb{P}(E \oplus E)$ , cf. GAETA, [G.1]) to the diagonal subspace  $\Delta \subset \mathbb{P}(E \oplus E)$ . The method extends naturally to  $h(\geq 2) V^{(j)} \subset \mathbb{P}_n$  provided  $c = \sum c_j \leq n$  ( $c_j = \text{codim } V^{(j)}$  in  $\mathbb{P}_n$ ). The geometric interpretation of  $F_V$  in terms of the complex

$$\mathcal{C}(V^c) = \{P_{c-1} \subset \mathbb{P}_n \mid P_{c-1} \cap V^c \neq \emptyset\}$$

leads naturally to an equivalence of the exponent multiplicity with VAN DER WAERDEN's theory (cf. § 10), [v.d.W 1], [v.d.W, ZAG].

Since  $c = \text{cod } J$  in  $\mathbb{P}(E \oplus \dots \oplus E)$  a natural discussion arises also in the case  $c > n$ . Then the old elimination theory (too much discredited because of its heavy dependence on coordinates) can be replaced by intrinsic constructions, cf. § 13, 14 pages 67-76. Natural applications are made to BEZOUT's theorem § 8, 11 as well to possible future relations with the "length multiplicity" (cf. VOGEL's report here), [vdW-ZAG], [Grö 1, 2].

0. INTRODUCTION

Most of the algebraic varieties needed in this paper will be embedded in a fixed complex projective space  $\mathbb{P}_n(\mathbb{C}) = \mathbb{P}(E_{n+1}) = E - (0)/\mathbb{C}^x$  (with  $E = E_{n+1}$   $(n + 1)$ -dimensional  $\mathbb{C}$ -vector space. The projection  $P : E - (0) \longrightarrow \mathbb{P}_n(\mathbb{C})$  will be denoted also by  $P$  although for a given  $v \in E - 0$  we write simply  $P(v) = (v)$ .

Let  $V, W$  be two irreducible algebraic varieties of  $\mathbb{P}_n(\mathbb{C})$  meeting properly.

Let

$$(0.1) \quad F_{V \cdot W} = \sum_{C=C_d \subset V \cap W} F_C^{i(V,W;C)}$$

be the *associale form* (a.F.), cf. DEF. 4.6; of the intersection cycle

$$(0.2) \quad V \cdot W = \sum_{C=C_d \subset V \cap W} i(V,W;C) C_d$$

where  $F_C$  is the (irreducible) a.F. to the irreducible component  $C_d$  of  $V \cap W$ . *The intersection multiplicities  $i(V,W;C)$  are uniquely determined as the exponents in the prime factor decomposition of  $F_{V \cdot W}$ ; this remark is useless if there is no way of computing intrinsically  $F_{V \cdot W}$  in terms of  $V$  and  $W$  ( $\Leftrightarrow F_V$  and  $F_W$ ). This paper shows that actually  $F_{V \cdot W}$  is uniquely and intrinsically determined in a natural way by restriction to the diagonal space  $\Delta \subset \mathbb{P}(E \oplus E)$  of the  $F_J$  associated to the join  $J = J(V \times W) \subset \mathbb{P}(E \oplus E)$  of  $V$  and  $W$ , § 1, DEF. 1.1, page 19.*

More precisely we have:



$$(0.3) \quad F_{V \cdot W} = \delta^{-1}(F_J | \Delta) \quad \text{cf. § 5}$$

where  $\delta : \mathbb{P}_n \longrightarrow \mathbb{P}(E \oplus E)$  is defined by

$$(0.4) \quad \delta(x) = ((x, x))$$

for any  $x \in E - \{0\}$  ,  $(x) = \mathbb{P}(x) \in \mathbb{P}(E)$  ,  $((x, x)) \in \mathbb{P}(E \oplus E)$  , is the *diagonal injection* and  $\Delta = \delta(\mathbb{P}_n)$  is the *diagonal space* (cf. § 3). The a.F  $F_J$  of  $J(V \times W)$  is *actually intrinsically determined by standard methods* (cf. § 6).

The construction can be extended in several ways:

a) If  $V \cap W$  is improper (0.3) is meaningless since  $V \cdot W$  is not defined as a cycle ,  $\Rightarrow F_{V \cdot W}$  is not defined. However the right hand side of (0.1) is always defined and we have

$$(0.3)' \quad \delta^{-1}(F_J | \Delta) = 0 \quad J = J(V \times W)$$

*iff*  $V \cap W$  is improper. Notice that

$$V \cap W \text{ improper} \Leftrightarrow J(V \times W) \cap \Delta \text{ improper}$$

b) The construction is valid also for finitely many irreducible varieties  $V^{(j)}$  denoted sometimes also by

$$(0.5) \quad V_{d_j} = V^{c_j} \subset \mathbb{P}_n \quad d_j + c_j = n , \quad j = 1, 2, \dots, h$$

where we use a double notation  $V = V_d = V^c$  for an irreducible  $V \subset \mathbb{P}_n$  if there is no ambiguity where the subscript  $d$  indicates the dimension and the superscript  $c$  the codimension:  $(d + c = n)$ , of  $V$  in  $\mathbb{P}_n$ .

The join

$$(0.6) \quad J = J(V^{c_1} \times V^{c_2} \times \dots \times V^{c_n}) \subset \mathbb{P}(E \oplus \dots \oplus E)$$

cf. § 1, DEF.1.1. is also irreducible of codimension  $c$  in  $\mathbb{P}(E \oplus \dots \oplus E)$ . This ambient space of  $J$  can be identified with  $\mathbb{P}(E \otimes \mathbb{C}^h)$  where the  $j$ -th direct summand  $(0, \dots, \underset{j}{E}, \dots)$  is identified with  $E \otimes e_j, e_j = (0, \dots, \underset{j}{1}, \dots, 0)$ . The set-theoretic intersection

$$\bigcap_{j=1}^h V^{c_j}$$

do always exists provided

$$(0.7) \quad c = c_1 + c_2 + \dots + c_n \leq n.$$

We shall assume (0.7) in the first part (cf. TABLE OF CONTENTS). Then we have

$$(0.8) \quad \text{cod} \bigcap_{j=1}^h V^{c_j} \leq n$$

and this *intersection is proper* ( $\Leftrightarrow \text{cod } \Omega = c$ ) iff  $J \cap \Delta$  is proper in  $\mathbb{P}(E \otimes \mathbb{C}^h)$  because if  $C$  runs through the set of irreducible components of  $\Omega$ ,  $\delta(C)$  runs through the set of all irreducible components of  $J \cap \Delta$  and  $\dim C = \dim \delta(C)$ . Then (0.3), (0.3)' can be extended to an arbitrary  $h \geq 2$  as indicated by the following:

THEOREM I.  $\delta^{-1}(F_J | \Delta) = 0$  iff the set-theoretic intersection  $\bigcap_{j=1}^h V^{c_j}$  is improper. Otherwise the intersection cycle  $I = V^{c_1} \cdot V^{c_2} \cdot \dots \cdot V^{c_h}$  is well defined in  $\mathbb{P}_n$  and we have

$$(0.9) \quad F_I = \delta^{-1}(F_J | \Delta) \quad \Delta = \delta(\mathbb{P}_n).$$

The associate Form  $F_J$  can be determined by  $V^{c_1}, V^{c_2}, \dots, V^{c_h}$  ( $\Leftrightarrow F_{V^{(j)}}(c_j, j = 1, 2, \dots, h)$ ) in the standard way (cf. § 6) for any  $h$ , as well as in the case  $h = 2$ .

There are several versions of the associate forms attached to a given pure cycle  $V^c \subset \mathbb{P}_n$  (and for each one the restriction symbol  $F | \Delta$  appearing in (0.9)) has a natural meaning); on the other hand all of them lead to the same intersection- multiplicities. But we shall use only the following three versions of the a.F :

$$(0.10) \quad (x_1, x_2, \dots, x_c) \longmapsto S(x_1, x_2, \dots, x_c) \quad (\text{CAYLEY-SEVERI}), \text{ cf. [C], [P], [S]}$$

$$(0.11) \quad (u_1, u_2, \dots, u_{d+1}) \longmapsto Y(u_1, u_2, \dots, u_{d+1}) \quad (\text{VAN DER WAERDEN-CHOW})$$

[Ch - vdW]

$$(0.12) \quad (u_1, u_2, \dots, u_{d+2}; x) \longmapsto N(u_1, u_2, \dots, u_{d+2}; x) \quad (\text{WEIL-SIEGEL}), \text{ [W], [Si].}$$

where the  $x$ 's  $\in E$ , the  $u$ 's  $\in \check{E} = \text{Hom}_c(E, \mathbb{C})$ . Cf. § 4 and they are defined up to a proportionality factor  $\lambda \in \mathbb{C}^\times$ . It suffices to define them first for an irreducible  $V^c$  and then to extend to the general  $\Gamma^c$  by "prime factor decomposition". All of them can be defined in terms of the  $\mathcal{C}(V^c)$  introduced by the following:

DEFINITION 0.1

*The Complex  $\mathcal{C}(V^c)$  of  $(c-1)$ -dimensional projective subspaces attached to an irreducible  $V^c$  defined by*

$$(0.13) \quad \mathcal{C}(V^c) = \{\mathbb{P}_{c-1} \subset \mathbb{P}_n \mid \mathbb{P}_{c-1} \cap V^c \neq \emptyset\} \quad (\text{cf. } \S 3)$$

(cf. § 4). In fact  $\mathcal{C}(V^c)$  is represented by an irreducible subvariety of codimension one in the Grassmann manifold  $\mathcal{G}(c-1; n)$ . Furthermore  $V^c$ , is recovered from  $\mathcal{C}(V^c)$  as the locus of singular points of  $\mathcal{C}(V^c)$ . Cf. § 4. The proof of our Theorem is a consequence of the following fact:

*If  $\bigcap_{j=1}^h V^{c_j}$  is improper the restriction of the complex  $\mathcal{C}(J)$  attached to  $J_c$ :*

$$\mathcal{C}(J(V^{(1)} \times \dots \times V^{(h)})) = \{\mathbb{P}_{c-1} \subset \mathbb{P}(E \otimes \mathbb{C}^h) \mid \mathbb{P}_{c-1} \cap J \neq \emptyset\}$$

to the diagonal space  $\Delta$  :

$$(0.14) \quad \mathcal{C}(J) \mid \Delta = \{\mathbb{P}_{c-1} \subset \Delta \mid \mathbb{P}_{c-1} \in \mathcal{C}(J)\}$$

is the full Grassmannian  $\mathcal{G}(c+1, \Delta)$  because if  $C^{c'}$  ( $c' < c$ ) is an excendatory irreducible component of  $\bigcap_{j=1}^h V^{(j)}$  then every subspace  $\mathbb{P}_{c-1}$  of  $\Delta$  meets the diagonal image  $\delta(C^{c'})$ .

If the previous  $\cap$  is proper the restriction  $\mathcal{C}(J)|_{\Delta}$  is a proper complex of  $\mathcal{G}(c-1; \Delta)$  and  $\delta^{-1}(\mathcal{C}(J)|_{\Delta})$  is a positive divisor  $\mathcal{C}(I)$  of the Grassmannian  $\mathcal{G}(c-1; \mathbb{P}_n)$  attached to  $I = V^{c_1} \cdot V^{c_2} \cdot \dots \cdot V^{c_n}$  in a natural way:

$$e_I = \sum i_c \mathcal{C}(C^c) \longleftrightarrow I = \sum_{\substack{C^c \subset \bigcap_{j=1}^h V^{c_j} \\ j=1}} i \left( \bigcap_{j=1}^h V^{c_j}; C^c \right) C^c.$$

The intersection multiplicities  $i_C = i \left( \bigcap_{j=1}^h V^c; C \right)$  equal the exponents of the corresponding  $F_C$ 's. In fact we recall in § 4 that all the  $F_V$  (0,10,11,12) are defined in terms of  $\mathcal{C}(V)$  by means of conjugation conditions (cf. DEF. 4.1). It suffices to assume first  $V$  irreducible. Namely:  $S$ , the CAYLEY-SEVERI form of  $V^c (= V_d)$  is the conjugation condition with respect to  $\mathcal{C}(V^c)$  of  $c$  Points  $(x_j)_{j=1,2,\dots,c}$ .  $Y$  (the original *zugeordnete Form*, (now usually called CHOW form of  $V_d$ ), (cf. § 5 and [S])) is the conjugation condition of  $d+1$  hyperplanes and the WEIL-SIEGEL form  $N$  (cf. [SI]). (SIEGEL's *Normalgleichung* of  $V$ ) is the conjugation condition of  $d+2$  hyperplanes  $(u_j) \in \mathbb{P}(E^V)$   $j=0,1,\dots,d+1$ , and one point  $(x)$  with respect to  $\mathcal{C}(V)$ . In terms of the exterior algebra:

$S, Y, N$  vanish if (cf. [BOU])  $x_1 \wedge x \wedge \dots \wedge x_c = 0$  or  $u_1 \wedge \dots \wedge u_{d+1} = 0$  or  $x \lrcorner u_1 \wedge \dots \wedge u_{d+2}$  vanish. If this is not the case any non zero product

$$\bigwedge_{j=1}^c x_j \quad \bigwedge_{j=1}^{d+1} u_j \quad x \perp \left[ \bigwedge_{j=0}^{d+1} u_j \right]$$

represents (in the well-known way) a projective subspace  $\mathbb{P}_{c-1} \subset \mathbb{P}_n$ . Then  $S = 0$  (resp.  $Y = 0$ ,  $N = 0$ ) iff such  $\mathbb{P}_{c-1} \in \mathcal{E}(V)$ . Cf. § 5, 6 for further details.

If  $V = \sum_{I > 0} m_I I \in \mathbb{P}_V$  is defined by  $F_I = \prod_{I \in \text{irr. dim } I=d} F_I^{m_I}$  ( $F = S, Y, N$ ). In any case the  $F$  is well defined up to a factor  $\lambda \in \mathbb{C}^\times$ .

In any case the restrictions  $S_j | \Delta, Y_j | \Delta, N_j | \Delta$  are well defined taking in (0.9)  $(x_j) \in \Delta | u_j | \Delta$ ,  $j = 1, 2, \dots$ .

The condition  $c \leq n$  of (0.7) – essential to define the previous restrictions to the diagonal space is not necessary in order to define the join  $J = J(V^{(1)} \times \dots \times V^{(h)})$  of  $h$  irreducible varieties  $V^{(j)} \subset \mathbb{P}(E)$ ,  $j = 1, 2, \dots, h$  cf. DEF. 1.1, page 19. In the case  $c > n$  the given varieties – in general position – do not meet but when  $c_1 = c_2 = \dots = c_h = 1$  the existence and discussion of a non empty intersection

$$(0.15) \quad \bigcap_{j=1}^h V \neq \emptyset$$

is precisely the goal of the old elimination theory! Accordingly we devote § 12 to such a problem also with arbitrary  $c_j$ 's – *but*  $c > 0$ . Under this hypothesis the compatibility condition (0.14) – equivalent to  $J \cap \Delta \neq \emptyset$  can be expressed by the following condition:

THEOREM II. *The  $h$  given irreducible varieties  $V^{(j)} \subset \mathbb{P}_n$ ,  $j = 1, 2, \dots, h$  with*

$c = \sum_{j=1}^n c_j > n$  *meet iff the diagonal space  $\Delta$  is singular for the complex  $\mathfrak{C}(J)$  attached to  $J = J(V^{(1)} \times \dots \times V^{(h)})$  ( $\Leftrightarrow$  every  $\mathbb{P}_{c-1}$  satisfying  $\Delta \subset \mathbb{P}_{c-1}$  belongs to  $\mathfrak{C}(J)$ ).*

In particular for  $c = n + 1$  we have:

*The  $h$  varieties  $V^{(j)}$  meet iff the diagonal space  $\Delta$  belongs to the complex  $\mathfrak{C}(J)$  attached to the join  $J = J(V^{(1)} \times \dots \times V^{(h)})$ .*

*This condition implies a single equation in the coefficients of  $F_J$  reducing to  $R = 0$  where  $R = R(f_1, f_2, \dots, f_{n+1})$  is the resultant of the  $n + 1$  hypersurfaces  $H_1, H_2, \dots, H_{n+1}$  if  $c_1 = c_2 = \dots = c_{n+1} = 1$ .*

*In the case  $c > n + 1$  the singularity condition of  $\Delta$  can be expressed by the identical vanishing of a covariant in agreement with GRAM'S theorem of invariant theory, cf. [WE].*

The distribution of matters is sufficiently indicated in the TABLE OF CONTENTS, page 2.

In the last part of the paper I review some results of the Author (cf. [G2.] [G3]) regarding a replacement of the usual KRONECKER elimination procedure by the explicit computation of the CAYLEY-SEVERI forms  $S_1(x_1, x_2, \dots, x_c)$  attached to an irreducible component  $I = I^c$  of codimension  $c$  of the ZARISKI-closed set represented by an arbitrary system

$$(0.16) \quad f_1 = 0 \quad f_2 = 0 \quad \dots \quad f_r = 0$$

of homogeneous polynomial equations in the homogeneous coordinates  $x_0, x_1, \dots, x_n$  in  $\mathbb{P}_n$ . The method rests on the fact that the "elimination of the variables"  $x_0, x_1, \dots, x_i$  represents geometrically the projection of a variety from a certain space of the projective

coordinate frame to the opposite face. If we replace these – indeed very particular projections – (essentially attached to the coordinate frame) by appropriate generic projections we obtain the indicated algorithm. But *the easy transition from the S to the N forms gives back the old KRONECKER elimination theory with respect to a generic frame "Φ" built in" in the formulas*, (instead of mentioning it but never written as before). I believe that this shows that the WEIL–SIEGEL forms are the best ones—although the CHOW forms seem to be the most famous. This inclusion of  $\Phi$  is actually accomplished by means of an arbitrary basis  $u_0, u_1, \dots, u_n$  of  $\check{E}$  acting as coordinate forms for points in

$$\mathbb{P}(E); x \mapsto (\langle u_0, x \rangle, \langle u_1, x \rangle, \dots, \langle u_n, x \rangle) \in \mathbb{P}(\mathbb{C}^{n+1})$$

for a fixed projective frame with current coordinates functions  $(x_0, x_1, \dots, x_n)$ .

In order to see that it suffices to represent the projection center  $\mathbb{P}_{c-2}$  defined in  $S_c$  by  $x_2 \wedge x_3 \wedge \dots \wedge x_c$  (with  $(x_1) = (x)$  acting as a current variable point of the projecting cone of a  $V^c$  from  $\mathbb{P}_{c-2}$ ) with hyperplane coordinates  $u_0, u_1, \dots, u_{d+1}$  in such a way that  $x_2 \wedge \dots \wedge x_c$  and  $x \lrcorner u_0 \wedge \dots \wedge u_{d+1}$  are dual, i.e. they represent the same  $\mathbb{P}_{c-2}$ .

We try to use standard notations as much as possible. Some non–standard ones are listed in the Notation sheet in page 0.

ACKNOWLEDGMENTS. The *ruled join* or *join* was introduced by the Author in [G.1] trying to compare  $F_V, F_W$  with  $F_V \times W$  (or  $F_{V \cdot W}$  when the intersection cycle does exist) with the name *prodotto rigato*, but actually similar ideas were frequent in the Italian School also in symmetric squares; for instance the symmetric square of a smooth curve was represented frequently by the variety of chords containing the



tangential surface as representative of the diagonal. But it was necessary also to recover lost properties of the "*zweifach projektive Raume*"  $\mathbb{P}_{m,n}$  remarking that the "point"  $(v,w) = (\lambda v, \mu w)$  of  $\mathbb{P}_{m,n}$  is essentially the same as the line  $\lambda(v,0) + \mu(0,w)$  but certain natural subspaces, such as  $\Delta$  do not appear in  $\mathbb{P}_{m,n}$ . Cf. § 1.2 for more details. This construction was also used by FULTON [F], [F-L] to illustrate his intersection theory and in the study of the topology of algebraic subvarieties of  $\mathbb{P}_n$ . A few years ago VOGEL [V.1], [V.2], [F-V] tried successfully to recover the "length multiplicity" – rejected previously for well-known reasons with a sort of reduction to the diagonal using the double projective space  $\mathbb{P}_{m,n}$ . KLEIMANN – in a letter to VOGEL [K] *recommended him to do precisely what I did in the join construction*. As a consequence I am coming back to this old technique. I hope to establish a link of the exponent multiplicity (previously used by VAN DER WAERDEN's elementary cases of BEZOUT's theorem by means of resultants) with the length multiplicity. The pleasant atmosphere and the kind invitation of the BANACH Center of the Polish Academy of Sciences is certainly a good encouragement in this direction.

The last part (of page 68) is just sketched – although the methods are very similar to those of [G.2], [G.3]. We shall come back to this with full details in [G.3] with an application to the SCHOTTKY problem (where the SIEGEL form appears in [SI]).

I am indebted to the wonderful facilities of the Max-Planck-Institute in Bonn – in particular to the extreme patience of the typist Frau Wolf-Gazo who made a beautiful job with them.

I. GENERALITIES ON JOINS

The reduction to the diagonal (cf. formula (1.1) below) introduced by C. SEGRE and SEVERI (fixed points of correspondences) and widely used later in Topology was applied by WEIL [W] and others to local intersection multiplicity theories. The global extension to varieties in a projective space has some difficulties due to the fact that the diagonal  $\Sigma$  is not anymore a linear space.  $\Sigma$  is a SEGRE variety:

$$\Sigma = \mathbb{P}\{x \otimes n \mid x \in E - \{0\}, n \in \mathbb{C}^h - \{0\}\},$$

cf. § 1, 3. We show [G.1], § 1, 2, 3 that a naturally chosen generator  $\Delta \subset \Sigma \subset \mathbb{P}(E \otimes \mathbb{C}^h)$ .

$$(1.0) \quad \Delta = \mathbb{P}\{x \otimes (1,1, \dots, 1) \mid x \in E - \{0\}\} = \mathbb{P}\{(x,x, \dots, x) \mid x \in E - \{0\}\}$$

plays the same role as in the affine case, although it is essential to introduce the space  $\mathbb{P}(E \otimes \mathbb{C}^h)$  instead of the "h-fach projektive Raum" of [vdW1], [vdW-ZAG], [H-P]. The affine formulas (1.4) lead naturally to the (1.4)' suggesting the definition of the *join* (cf. DEF. 1.1) and the *projective reduction to the diagonal*, cf. formula (1.10) in page 20.

1. THE "REDUCTION TO THE DIAGONAL" . A PROJECTIVE VERSION.

Let  $A_j \neq \emptyset$ ,  $j = 1, 2, \dots, h$  be  $h$  non empty subsets of an ambient set  $E$ . Let  $\Pi = E \times E \times \dots \times E$  be the  $h^{\text{th}}$  Cartesian power of  $E$ . We have:

$$(1.1) \quad \delta\left(\bigcap_{j=1}^h A_j\right) = A_1 \times A_2 \times \dots \times A_h \cap \Delta$$

where  $\delta: E \hookrightarrow \Pi$  is the *diagonal injection*  $\delta(x) = (x, x, \dots, x)$ ,  $\forall x \in E$  and  $\Delta = \delta(E)$  is the *diagonal* of  $\Pi$ .

This simple remark has many applications in algebraic Geometry and it is regarded as a "reduction" (in spite of the fact that  $\Pi$  seems more complicated than  $E$ ) because of the following reasons:

a) *If  $E$  is an algebraic variety and the  $A_j$  are all subvarieties,  $\Pi$  is also an algebraic variety and  $A_1 \times \dots \times A_h$  and  $\Delta$  are algebraic subvarieties of  $\Pi$  with  $\Delta$  independent of the  $A_j$ .*

b) *The subvarieties of  $\Pi$  are graphs of algebraic  $h$ -correspondences on  $E$ , in particular they might be graphs of maps and  $\Delta$  is the graph of the identity. If we can "move"  $\Pi$  in an algebraic system, it is possible to move the  $A_j$  to generic positions  $\bar{A}_j$ ,  $j = 1, 2, \dots, h$  in such a way that we can predict geometric statements on the original  $A_j$ 's by a subsequent specialization.*

c) *In particular: if  $E$  is an affine space,  $\Pi$  is another one and  $\Delta$  is a linear subspace of  $\Pi$  with  $\dim E = \dim \Delta$ . In this case  $\delta(I)$  is an irreducible component of  $A_1 \times \dots \times A_n \cap \Delta$  iff  $I$  is an irreducible component of  $\bigcap_{j=1}^h A_j$ . Accordingly  $\Pi \cap \Delta$  is proper iff  $\bigcap_{j=1}^h A_j$  is proper. Since the definition of the intersection multiplicities looks easier if one of the intersecting varieties is a linear space the diagonal provides a way to define*

$$(1.2) \quad i(A_1 \cdot \dots \cdot A_h; I) = i(\Pi \cdot \Delta; \delta(I))$$

i.e.: *It suffices to know how to define  $\Pi \cdot \Delta$  ( $h = 2$ ) and  $\Delta$  a linear space.* Cf. [W], [F]. The affine case is sufficient for all the local theories.

If  $E$  is a projective space  $\mathbb{P}_n$ ,  $\Pi$  and  $\Delta$  are not projective spaces, but SEGRE varieties, cf. [SE], [H-P]. However the explicit description of  $\Delta$  in the affine case leads naturally to the "join construction" (cf. Introduction) as follows: Let us assume  $h = 2$ . Then  $\Delta$  is characterized by the system of linear equations

$$(1.3) \quad x_j - y_j = 0 \quad j = 1, 2, \dots, n$$

if  $(x_1, \dots, x_n)$  and  $(y_1, y_2, \dots, y_n)$  are current affine coordinates in the two copies of  $E$ . If  $f_i(x) = 0$ ; and  $g_j(x) = 0$  are two systems of equations defining  $A_1, A_2$  the system

$$(1.4) \quad f_j(x) = 0 \quad g_h(y) = 0$$

defines  $A_1 \times A_2$ . (1.4) and (1.3) together define  $\Pi \cap \Delta$ .

In the projective case the  $x, y$  can be regarded as absolute coordinates in the  $\mathbb{C}$ -vector space  $E = E_{n+1}$  or as homogeneous coordinates in  $\mathbb{P}_n = \mathbb{P}(E)$  and (1.3) is replaced by

$$(1.5) \quad \text{rank} \begin{bmatrix} x_0 & x_1 & \dots & x_n \\ y_0 & y_1 & \dots & y_n \end{bmatrix} = 1 \Rightarrow \begin{vmatrix} x_i & x_j \\ y_i & y_j \end{vmatrix} = 0 \quad 0 \leq i < j \leq n$$

Then the (1.4) can be replaced by

$$(1.4)' \quad f_j(\lambda x) = 0 \quad g_h(\mu y) = 0$$

where all the  $f_j$  and  $g_h$  are homogeneous and  $\lambda, \mu$  are two independent non zero proportionality factors. Moreover the equations (1.5) define the SEGRE variety representing  $\mathbb{P}(E) \times \mathbb{P}(\mathbb{C}^1)$  ( $\Leftrightarrow$  locus of  $x \otimes (\lambda, \mu) \in \mathbb{P}(E \otimes E) = \mathbb{P}(E \otimes \mathbb{C}^2)$ ). Cf. § 3 for further details.

#### REMARKS

- 1) *We do not need the homogeneous equations  $f_i(x) = 0 \quad g_j(y) = 0$  anymore to establish (1.1).*
- 2)  *$A_1, A_2$  can be arbitrary non empty subsets of  $\mathbb{P}(E)$ .*
- 3) *It suffices to define the two injections  $i_{1,i_2} : i_{1,i_2} \hookrightarrow \mathbb{P}(E) \hookrightarrow \mathbb{P}(E \otimes E)$  by*

$$(1.6) \quad i_1(x) = (x, 0) \quad i_2(x) = (0, x)$$

$i_{1,i_2}$  have the following properties:

$$i_1(\mathbb{P}(E)) = \mathbb{P}(E \otimes 0) = \mathbb{P}(E \otimes (1, 0))$$

$$(1.7) \quad i_2(\mathbb{P}(E)) = \mathbb{P}(0 \otimes E) = \mathbb{P}(E \otimes (0, 1))$$

$$(1.8) \quad i_1(\mathbb{P}(E)) \cap i_2(\mathbb{P}(E)) = \phi$$

$$(1.9) \quad \Delta = \mathbb{P}(E \otimes (1, 1))$$

In other words *the two copies of  $\mathbb{P}(E)$  in  $\mathbb{P}(E \oplus E) = \mathbb{P}(E \oplus \mathbb{C}^2)$  do not meet;* accordingly *any ordered pair  $(P, Q) \in \mathbb{P}(E) \times \mathbb{P}(E)$  can be represented by the line joining  $i_1(P)$  with  $i_2(Q)$  and conversely any line joining one point of  $\mathbb{P}(E \oplus 0)$  with another one of  $\mathbb{P}(0 \oplus E)$  represents a uniquely defined ordered pair  $(P, Q)$ .*

More generally we have the following formal definition of the join (used already before).

DEFINITION 1.1

4) The join of  $A_1, A_2$ , denoted by  $J(A_1 \times A_2)$ , *is the locus of all (always well defined!) lines joining points of  $i_1(A_1)$  with points of  $i_2(A_2)$ .* In particular;  $J(\mathbb{P}(E) \times \mathbb{P}(E))$  *is the subvariety of  $\mathbb{P}(E \oplus E)$  consisting of lines joining points of  $\mathbb{P}(E \oplus 0)$  and  $\mathbb{P}(0 \oplus E)$ .*

5) The following natural generalizations are possible

$$(\phi \neq) A_1 \subset \mathbb{P}(E) \mid (\phi \neq) A_2 \subset \mathbb{P}(F) \Rightarrow J(A_1 \times A_2) \subset \mathbb{P}(E \oplus F)$$

because  $i_1 : \mathbb{P}(E) \hookrightarrow \mathbb{P}(E \oplus F)$   $i_2 : \mathbb{P}(F) \hookrightarrow \mathbb{P}(E \oplus F)$  are still valid.

6) We can consider any finite number  $h$  of non-empty subsets

$$A_j \subset \mathbb{P}(E_{(j)}) \quad j = 1, 2, \dots, h.$$

We shall consider this general set up in § 2 in order to clarify the relationship between the diagonal subspace  $\Delta$  and the diagonal variety  $\Sigma$  in § 3.

*The "reduction to the diagonal" in  $\mathbb{P}(E)$  has finally the following expression:*

$$(1.10) \quad \delta(A_1 \times A_2) = J(A_1 \times A_2) \cap \Delta$$

where  $A_1, A_2$  are arbitrary non empty subsets of  $\mathbb{P}(E)$ ,  $\Delta$  is the diagonal space (cf. (1.9), and  $\delta : \mathbb{P}(E) \longleftrightarrow \mathbb{P}(E \oplus E)$  is defined by (0.4), :

$$(1.11) \quad \delta(x) = ((x,x) = (x \otimes (1,1)) \quad \forall x \in E - \{0\} .$$

REMARK

We see that in the formula (1.10) one needs the points of  $\mathbb{P}(E \oplus E)$ , for instance those  $((x,x) \in \Delta$ , not just the lines  $\lambda(x,0) + \mu(0,y)$ . *This justifies our preference for the join construction rather than the use of the two-way projective space  $\mathbb{P}_{m,n}$ ; in  $\mathbb{P}_{m,n}$  the previous line is the "point"  $(x,y) \sim (\lambda x, \mu y)$ ,  $\lambda \neq 0, \mu \neq 0$ .*

2. RECALL OF THE JOIN OF  $h$  VARIETIES. RELATION WITH THE SEGRE MODEL OF THE PRODUCT  $V^{(1)} \times V^{(2)} \times \dots \times V^{(h)}$ .

Let  $\mathbb{P}(E_j) = E_j - \{0\} / \mathbb{C}^\times$ ,  $j = 1, 2, \dots, h$  be  $h (\geq 2)$  complex projective spaces generated by the corresponding vector spaces  $E_j$ . Let  $\mathbb{P}(S)$  be the quotient projective space of the direct sum

$$(2.1) \quad S = E_1 \oplus E_2 \oplus \dots \oplus E_h .$$

Let us call  $S_j = (0, \dots, E_j, \dots, 0)$   $j = 1, 2, \dots, h$ .  $\mathbb{P}(S)$  is the ambient projective space containing copies  $\mathbb{P}(S_j) = i_j(\mathbb{P}(E_j))$ ,  $j = 1, 2, \dots, h$  of the given spaces  $\mathbb{P}(E_j)$  satisfying the following properties (already checked for  $h = 2$ ) :

a) For every ordered  $h$ -tuple  $(x_1, x_2, \dots, x_h) \in \prod_{j=1}^h \mathbb{P}(E_j)$  the corresponding images  $i_j(x_j)$  ( $j = 1, 2, \dots, h$ ) are linearly independent

b) The  $S_{h-1} = S_{h-1}(x_1, x_2, \dots, x_h)$  space spanned by the  $x_j$  meet  $\mathbb{P}(S_j)$  precisely in the point  $x_j$ :

$$S_{h-1} \cap \mathbb{P}(S_j) = x_j \quad j = 1, 2, \dots, h.$$

As a consequence we have:

c) There is a bijection of  $\prod_{j=1}^h \mathbb{P}(E_j)$  with the subset  $\mathcal{J}(\mathbb{P}(E_1) \times \dots \times \mathbb{P}(E_h)) \subset \mathcal{G}(h-1; \mathbb{P}(E_1 \oplus \dots \oplus E_h))$  of the sharon Grassmannian of  $(h-1)$ -spaces defined by:

$$(2.2) \quad \mathcal{J} = \mathcal{J}(\mathbb{P}(E_1) \times \dots \times \mathbb{P}(E_h)) = \{ \mathbb{P}_{h-1} \subset \mathbb{P}(E_1 \oplus \dots \oplus E_h) \mid \mathbb{P}_{h-1} \cap i_j(\mathbb{P}(E_j)) = x_j, \quad j = 1, 2, \dots, h \}$$

for  $j = 1, 2, \dots, h$ .  $\mathcal{J}$  is closely related to  $J$  by

DEFINITION 2.1  $J = J(\mathbb{P}(E_1) \times \dots \times \mathbb{P}(E_h))$  is defined in terms of  $\mathcal{J}$  (cf. (2.2)) by

$$(2.3) \quad J = J(\mathbb{P}(E_1) \times \dots \times \mathbb{P}(E_h)) = \{ \mathbb{P}_{h-1} \subset \mathbb{P}(E_1 \oplus \dots \oplus E_h) \mid \mathbb{P}_{h-1} \in \mathcal{J} \}$$

is called the *ruled join* (or just *join*) of the given spaces  $\mathbb{P}(E_1), \mathbb{P}(E_2), \dots, \mathbb{P}(E_h)$ .

DEF. 2.1 is the extension of DEF. 1.1 page 19 for any  $h \geq 2$ .



A vector of  $S$  is regarded as an ordered  $h$ -tuple  $(v_1, v_2, \dots, v_h)$  with  $v_j \in E_j$ ,  $j = 1, 2, \dots, h$ . Let  $i_j : E_j \hookrightarrow S$  be the natural injection defined by

$$(2.4) \quad i_j(v) = (0, 0, \dots, \overset{j}{\underbrace{v}}, \dots, 0) \quad v \in E_j$$

where  $i_j(\mathbb{P}(E_j)) = \mathbb{P}(S_j) = \mathbb{P}(0, \dots, E_j, \dots, 0)$ . We shall use the same symbol  $i_j$  for the corresponding maps between projective spaces.

$$(2.5) \quad i_j : \mathbb{P}(E_j) \hookrightarrow \mathbb{P}(S) \quad i_j(\mathbb{P}(E_j)) = \mathbb{P}(S_j) \quad j = 1, 2, \dots, h.$$

It is easy to check both conditions a), b) for the  $h$  copies  $\mathbb{P}(S_1), \mathbb{P}(S_2), \dots, \mathbb{P}(S_h)$  of given projective spaces  $\mathbb{P}(E_j)$ . In fact any ordered  $h$ -tuple

$$(x_1, \times x_2 \times \dots \times x_h) \in \prod_{j=1}^h \mathbb{P}(E_j) \quad ((E_j - \{0\}))$$

defines an  $h$ -tuple of linearly independent vectors  $i_j(x_j) \in S_j \quad j = 1, 2, \dots, h \quad (\Leftrightarrow \bigwedge_{j=1}^h i_j(x_j) \neq 0)$ . They define a subspace  $S(x_1, x_2, \dots, x_h)$  of dimension  $h - 1$  in  $\mathbb{P}(S)$  - the projection in  $\mathbb{P}(S)$  of the  $h$ -dimensional vector space locus of points of the type:

$$(2.6) \quad (\lambda_1(v_1, 0, \dots, 0) + \lambda_2(0, v_2, \dots, 0) + \dots + \lambda_h(0, 0, \dots, v_h))$$

in such a way that

$$(2.7) \quad S(x_1, \dots, x_h) \cap \mathbb{P}(S_j) = (x_j) \quad j = 1, 2, \dots, h$$

and conversely.

Another  $(y_1) \times \dots \times (y_h) \in \prod_{j=1}^h \mathbb{P}(E_j) (E_j - \{0\})$  defines the same  $h$ -tuple of points in  $\mathbb{P}(S_1) \times \dots \times \mathbb{P}(S_h)$  and also the same  $S_{h-1}$  iff  $y_j = \lambda_j x_j \quad \lambda_j \in \mathbb{C}^\times$   $j = 1, 2, \dots, h$ , (i.e. iff  $(x_1, x_2, \dots, x_h) \sim (y_1, \dots, y_h)$  as points of the "*h-way projective space of*  $\mathbb{P}_{n,n, \dots, n}$  (cf. Introduction, [H-P], [vdW1], [vdW-ZAG]); in other words:

$$S(x_1, \dots, x_h) = S(y_1, \dots, y_h) \Leftrightarrow y_j = \lambda_j x_j \quad j = 1, 2, \dots, h.$$

This construction leads to two modifications of DEF. 2.1 obtained taking into account rather than the  $\mathbb{P}_{h-1}$  of  $\mathcal{J}$  some set of points in  $\mathbb{P}(E_1 \oplus \dots \oplus E_h)$

DEF.2.1' The *pre-join*

$$J_p = J_p(\mathbb{P}(E_1) \times \dots \times \mathbb{P}(E_h)) = \{((x_1, \dots, x_h)) \in \mathbb{P}(E_1 \oplus \dots \oplus E_h) \mid x_j \neq 0 \quad j = 1, 2, \dots, h\}.$$

DEF.2.2' The *full-join*  $J$  is the ZARISKI closure of  $J_p$  :

$$(2.3)' \quad J = \overline{J_p} = \bigcup_{\mathbb{P}_{h-1} \in \mathcal{J}} \mathbb{P}_{h-1}$$

However, in spite of the differences between  $\mathcal{J}$ ,  $J_p$ ,  $J$  the context will indicate without confusion which one we need, and we prefer the simplest notation  $J$ .

#### REMARK

The name *ruled join* ("*prodotta rigata*") is clear since an  $h$ -tuple of

$\mathbb{P}(E_1) \times \dots \times \mathbb{P}(E_h)$  is not represented by a point of another space but by a  $\mathbb{P}_{h-1}$ , i.e. *by a line for*  $h = 2$  (cf. Introduction).

*The product*  $\mathbb{P}(E_1) \times \dots \times \mathbb{P}(E_h)$  *is represented also by the quotient set*

$$(2.7) \quad J_p / \sim = J_p / \mathbb{C}^x \times \dots \times \mathbb{C}^x = \prod_{j=1}^h (E_j - \{0\} / \mathbb{C}^x \times \dots \times \mathbb{C}^x$$

usually called the *r-way projective space*  $\mathbb{P}_{n_1, n_2, \dots, n_h}$  where  $n_j = \dim \mathbb{P}(E_j)$  by [vdW], [vdW-ZAG]; s. also [H-P].

REMARKS:

1) Since there is a bijection between "points"  $(v_1, \dots, v_h)$  of  $\mathbb{P}_{n_1, n_2, \dots, n_h}$  and  $(h-1)$ -dimensional subspaces of type  $S(v_1, v_2, \dots, v_h)$  the relation between  $J_p$   $J_p / \sim = \mathbb{P}_{n_1, n_2, \dots, n_h}$  and  $J$  is very close (cf. DEF. 2,1). The reason of our preference of  $J$  over  $\mathbb{P}_{n_1, n_2, \dots, n_h}$  is due to the fact that in the interpretation of the *reduction to the diagonal* (cf. § 1) we need  $J$  (rather than  $J_p$  or  $\mathcal{J}$ ) and the subset  $\Delta \subset \mathbb{P}(E \oplus \dots \oplus E)$  (which do not belong to  $\mathbb{P}_{n_1, \dots, n_h}$ ). In other words *the equivalence relation defining*  $\mathbb{P}_{n_1, \dots, n_h}$  *loses the points of*  $\mathbb{P}(E \oplus \dots \oplus E)$  *needed essentially in the reduction to the diagonal.*

EXAMPLE. The product  $\mathbb{P}_1 \times \mathbb{P}_1 = \mathbb{P}(E)$  ( $\dim E = 2$ ) is represented by the set (*line congruence*)  $J$  of lines joining pairs of points of  $\mathbb{P}(S_1) = \mathbb{P}(E \oplus 0)$  and

$\mathbb{P}(S_2) = \mathbb{P}(0 \oplus E_2)$ . The two lines  $\mathbb{P}(S_1)$ ,  $\mathbb{P}(S_2)$  do not meet and conversely *any line of this congruence determines uniquely the pair of points (A,B)*.




Fig. 1

The relation of the ruled model  $J(\mathbb{P}(E_1) \times \dots \times \mathbb{P}(E_h))$  with the usual SEGRE model  $\Sigma_{n_1, \dots, n_h}$  is very simple. It suffices to show it for  $h = 2$ :

Let  $J(m,n) = J(\mathbb{P}_m \times \mathbb{P}_n)$  be the join and let  $\Sigma_{m,n} \subset \mathbb{P}(E_1 \oplus E_2)$  be the SEGRE model; let us recall that  $\Sigma_{m,n}$  is the image of the set of ( $\neq 0$ ) monomial elements  $x \otimes y (x \in E_1, y \in E_2)$  in the tensor product  $E_1 \otimes E_2$  by the canonical projection  $E_1 \otimes E_2 \longrightarrow \mathbb{P}(E_1 \otimes E_2)$  in such a way that the pair  $(x) \times (y) \in \mathbb{P}(E_1) \times \mathbb{P}(E_2)$  is represented by  $(x \otimes y) \in \mathbb{P}(E_1 \otimes E_2)$ . The Grassmann coordinates of the line joining  $(x,0)$  with  $(0,y)$  are the two-minors of the matrix

$$(2.8) \quad \begin{bmatrix} x^0 & x^1 & \dots & x^m & 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & y^{0'} & y^{1'} & \dots & y^{n'} \end{bmatrix}$$

where we choose a couple of basis in  $E_1$ ,  $E_2$  labelling the coordinates with the indices  $0, 1, \dots, m, 0', 1', \dots, n'$ ; we have  $p^{ij} = p^{i'j'} = 0$  but

$$(2.9) \quad p^{ij'} = x^i y^{j'} = \text{coordinates of } x \otimes y.$$

In other words: *The products  $\alpha^i y^j$  representing the coordinates of  $x \otimes y$  in a canonical basis represent also the essential Grassmann coordinates of the line joining  $(i_1(x))$  with  $(i_2(y))$ .*

Intrinsically: *we can identify  $x \otimes y$  with  $i_1(x) \wedge i_2(y)$  inside  $E_1 \oplus E_2$ ; similarly we have for any  $h \geq 2$*

$$(2.10) \quad x_1 \otimes x_2 \otimes \dots \otimes x_h \otimes i_1(x_1) \wedge i_2(x_2) \wedge \dots \wedge i_h(x_h) \text{ in } E_1 \oplus E_2 \oplus \dots \oplus E_h \cdot \text{ Cf. [SG], [B] . [H-P]}$$

*The join of  $h$  irreducible subvarieties  $V^{(j)} \subset \mathbb{P}(E_j)$  is naturally defined by restriction as follows:*

DEF. 2.2 Let  $i_j(V^{(j)}) \subset \mathbb{P}(S_j)$  be the corresponding copies of the  $h$  given subvarieties. *The join  $J(V^{(1)} \times \dots \times V^{(h)})$  of  $V^{(1)}, V^{(2)}, \dots, V^{(h)}$  is the restriction of  $J = J(\mathbb{P}(E_1) \times \dots \times \mathbb{P}(E_{h-1}))$  to the  $\mathbb{P}_{h-1}$  subspaces of  $J$  joining points of the  $i_j(V^{(j)})$ ,  $j = 1, 2, \dots, h$ .*  
 $(V^{(1)} \times \dots \times V^{(h)}) = \{\mathbb{P}_{h-1} \in J(\mathbb{P}(E_1) \times \dots \times \mathbb{P}(E_h)) \mid \mathbb{P}_{h-1} \cap S_j = V^{(j)} \text{ } j = 1, 2, \dots, h\}$ .

We shall use the following properties of  $J(V^{(1)} \times \dots \times V^{(h)})$ :

1)  $J(V^{(1)} \times V^{(2)} \times \dots \times V^{(h)})$  is irreducible if  $V^{(j)}$  is irreducible (for  $j = 1, 2, \dots, h$ ). Moreover:

$$(2.11) \quad \dim J(V^{(1)} \times V^{(2)} \times \dots \times V^{(h)}) = d_1 + d_2 + \dots + d_h + h - 1$$

where  $d_j = \dim V^j$ ,  $j = 1, 2, \dots, h$ .

$$\begin{aligned} 2) \quad J(V^{(1)} \times \dots \times V^{(h)}) &= \\ &= \bigcap_{j=1}^h J(\mathbb{P}(E_1) \times \dots \times \mathbb{P}(E_{j-1}) \times V^{(j)} \times \mathbb{P}(E_{j+1}) \times \dots \times \mathbb{P}(E_h)). \end{aligned}$$

3) The codimension  $c$  of  $J(V^{(1)} \times \dots \times V^{(h)})$  in  $\mathbb{P}(E_1 \oplus \dots \oplus E_h)$  is equal to the sum of the codimensions  $c_j = n - d_j$ ,  $j = 1, 2, \dots, h$

$$(2.12) \quad c = c_1 + c_2 + \dots + c_h.$$

### 3. CASE $n_1 = n_2 = \dots = n_h = n$ . THE DIAGONALS $\Sigma$ , $\Delta$

The case  $E_1 = E_2 = \dots = E_h = E$ ,  $S = E \oplus E \oplus \dots \oplus E$ ,  $\dim E = n + 1$  is particularly important in the intersection problems, because then we need to consider the representation of the abstract diagonal

$D = \{P_1 \times P_2 \times \dots \times P_n \in \mathbb{P}(E) \times \mathbb{P}(E) \times \dots \times \mathbb{P}(E) \mid P_1 = P_2 = \dots = P_n\}$  in the abstract product  $D$  is represented by the SEGRE model  $\Sigma_{n_1 n_2, \dots, n_h}$  is a VERONESE variety  $V(D)$  (cf. [B])

$$(3.1) \quad V(D) = \{\lambda(x_1 \otimes x_2 \otimes \dots \otimes x_h) \in \Sigma_{n, n, \dots, n} \mid x_1 = x_2 = \dots = x_n \neq 0\}$$

In the join the image of  $(\lambda_1 x) \times (\lambda_2 x) \times \dots \times (\lambda_h x)$   $\lambda_j \neq 0$   $j = 1, 2, \dots$  is the subspace  $S(x, x, \dots, x)$ , thus the image of  $D$  is

$$(3.2) \quad \Sigma_D = U\{S(x_1, x_2, \dots, x_h) \mid x_1 = x_2 = \dots = x_h\}$$

$\Sigma = \Sigma_D$  is a SEGRE variety model of  $\mathbb{P}(E) \times \mathbb{P}(\mathbb{C}^h)$ . In order to see that it is convenient to introduce the following identifications:

$$(3.3) \quad S = E \otimes \mathbb{C}^h \quad S_j = E \otimes u_j \quad S = \bigoplus_{j=1}^h S_j$$

where  $u_j = (0, 0, \dots, \underset{j}{1}, \dots, 0)$ ,  $j = 1, 2, \dots, h$ .

$$(3.4) \quad (x_1, x_2, \dots, x_h) \in J \Leftrightarrow (x_1 \otimes u_1, x_2 \otimes u_2, \dots, x_h \otimes u_h)$$

(3.4) implies in the diagonal case  $x_1 = x_2 = \dots = x_h = x \neq 0$ .

$$(3.5) \quad (\lambda_1 x, \lambda_2 x, \dots, \lambda_h x) \Leftrightarrow x \otimes (\lambda_1, \lambda_2, \dots, \lambda_h).$$

The generating spaces  $\mathbb{P}(E) \otimes (\lambda_1, \lambda_2, \dots, \lambda_h)$   $((\lambda_1, \dots, \lambda_h) \in \mathbb{P}(\mathbb{C}^h))$  and  $(x) \otimes \mathbb{P}(\mathbb{C}^h)$  are represented by

$$\mathbb{P}\{x \otimes (\lambda_1, \dots, \lambda_h) \mid x \in E\} \text{ and } \mathbb{P}\{x \otimes (\lambda_1, \dots, \lambda_h) \mid (\lambda_1, \dots, \lambda_h) \in \mathbb{C}^h\}$$

respectively. The latter is the image of the abstract diagonal point

$(x) \times (x) \times \dots \times (x)$ , i.e. by the span of the  $h$  copies of  $(x)$  in  $\mathbb{P}(S_j)$   $j = 1, 2, \dots, h$ .

*The former is a copy of  $\mathbb{P}(E)$ , the copy maps being*

$$(x) \longmapsto (x) \otimes (\lambda_1, \dots, \lambda_h).$$

In particular we have the following distinguished copies

$$\begin{aligned} \mathbb{P}(S_j) &= \mathbb{P}(E \otimes u_j) \subset \Sigma_D \quad j = 1, 2, \dots, h \\ \Delta &= \mathbb{P}(E \otimes (1, 1, \dots, 1)) = \mathbb{P}\{(x, x, \dots, x) \mid x \in E\} \subset \Sigma_D \subset \Gamma(E \otimes \mathbb{C}^h) \end{aligned}$$

$\Delta$  is the diagonal space (cf. Introduction) not to be confused with  $\Sigma_D$ .

*The reduction to the diagonal for  $h$  arbitrary non empty subsets  $A_1, A_2, \dots, A_n$  of  $\mathbb{P}(E)$  has the final form:*

$$(3.6) \quad \delta\left(\bigcap_{j=1}^h A_j\right) = J(A_1 \times A_2 \times \dots \times A_h) \cap \Delta$$

where  $J$  is the full join:  $J(A_1 \times \dots \times A_h) = \{\mathbb{P}_{h-1} \mid \mathbb{P}_{h-1} \in J(A_1 \times A_2 \times \dots \times A_h)\}$ .

Let us come back to our interesting case  $A_j = V^{(j)}$  irreducible algebraic subvariety of  $\mathbb{P}(E)$  of dimension  $d_j$  and codimension  $c_j$ . We know (cf. formula (2.12)) that  $\text{cod } J$  in  $\mathbb{P}(E \otimes \mathbb{C}^h)$  is equal to  $c = c_1 + c_2 + \dots + c_h$ . Then our discussions lead naturally to the two cases  $c \geq n$  and  $c < n$ .

If  $c \leq n$  is always  $\bigcap V^{(j)} \neq \emptyset \Leftrightarrow J(V^{(1)} \times \dots \times V^{(h)}) \cap \Delta \neq \emptyset$ .

If  $c > n$   $\bigcap V^{(j)} = \emptyset$  for the  $V^{(j)}$  in generic position  $\Leftrightarrow$  the diagonal space  $\Delta$  does not meet the join:

$$(3.7) \quad J(V^{(1)} \times \dots \times V^{(h)}) \cap \Delta = \emptyset \Leftrightarrow \bigcap_{j=1}^h V^{(j)} = \emptyset.$$



4. JOINS AND h-COLLINEATIONS.

The h-way projective space  $\mathbb{P}_{n_1, n_2, \dots, n_h} = \prod_{j=1}^h (E_j - \{0\}) / \mathbb{C}^x \times \dots \times \mathbb{C}^x$  where  $\dim E_j = n_j + 1$  was introduced by VAN DER WAERDEN [Ch-vdW] to study the correspondences in  $\mathbb{P}_{n_1} \times \mathbb{P}_{n_2} \times \dots \times \mathbb{P}_{n_h}$  (cf. also [H-P], Vol. I, Chapter V, § 10 and specifically Vol. II, Ch XI). An irreducible correspondence in

$$\prod_{n_1, n_2, \dots, n_h} = \mathbb{P}_{n_1} \times \dots \times \mathbb{P}_{n_h}$$

is an irreducible subvariety of this product. The natural way to study them is to introduce the systems of homogeneous polynomial equations; a polynomial  $f \in \mathbb{C} [x^{(1)}, x^{(2)}; \dots; x^{(h)}]$  (where  $x^{(j)} = (x_0^{(j)}, x_1^{(j)}, \dots, x_{n_j}^{(j)})$ ,  $j = 1, 2, \dots, h$ ) is called homogeneous of degree  $(m_1, m_2, \dots, m_h)$  iff

$$(4.1) \quad f(\lambda_1 x^{(1)}, \lambda_2 x^{(2)}, \dots, \lambda_h x^{(h)}) = \lambda_1^{m_1} \lambda_2^{m_2} \dots \lambda_h^{m_h} f(x^{(1)}, \dots, x^{(h)}).$$

In the interpretation of the points of  $\mathbb{P}_{n_1, n_2, \dots, n_h}$  as  $(h-1)$ -subspaces of  $J(\mathbb{P}_{n_1} \times \dots \times \mathbb{P}_{n_h})$  any subvariety of  $\mathbb{P}_{n_1, \dots, n_h}$  might be regarded as a Grassmannian subvariety:

$$\mathcal{S} \subset \mathcal{J}(\mathbb{P}_{n_1} \times \dots \times \mathbb{P}_{n_h}) \subset \mathcal{G}(h-1; \mathbb{P}(E_1 \oplus \dots \oplus E_h)).$$

The transition of  $\mathcal{J}$  to  $J$  originates a ruled variety

$$(4.2) \quad S = \bigcup_{\mathbb{P}_{h-1} \in \mathcal{S}} \mathbb{P}_{h-1}$$

We shall omit the easy transition of the language developed in [H-P] for  $\mathbb{P}_{n_1, n_2, \dots, n_h}$  to our "join" - interpretation with the exception of the *h-collineations* among  $h$  copies of  $\mathbb{P}(E) = \mathbb{P}_n$ : *they have some special properties closely related to the subspaces of  $\mathbb{P}(E \otimes \mathbb{C}^h)$  which will enable us in § 10 to show the equivalence of the exponent multiplicity with VAN DER WAERDEN's.*

Let us recall the following ones:

4) Let  $P = (v_1, v_2, \dots, v_h) \in \mathbb{P}(E \otimes \mathbb{C}^h)$  be one point of  $\mathbb{P}(J_p)$  ( $\Leftrightarrow v_j \neq 0$ ,  $j = 1, \dots, h$ ). Then there is one and only one  $\mathbb{P}_{h-1} \in \mathcal{J}(\mathbb{P}(E) \times \dots \times \mathbb{P}(E))$  containing  $P$ .

Let  $U \subset \mathbb{P}(J_p)$  be a *J-unisecant variety*  $\Leftrightarrow U$  does not contain two different points belonging to the same  $\mathbb{P}_{h-1} \in \mathcal{J}(\mathbb{P}_n \times \overset{h}{\dots} \times \mathbb{P}_n)$ . Then  $U$  represents in a natural way the same *h-correspondence* that the ruled variety  $R$  locus of  $\mathbb{P}_{h-1} \in \mathcal{J}(\mathbb{P}(E) \times \dots \times \mathbb{P}(E))$  meeting  $U$ :

$$R = \bigcup_{\mathbb{P}_{h-1} \in \mathcal{J}(\mathbb{P}(E) \times \dots \times \mathbb{P}(E))} \mathbb{P}_{h-1} \cap U \neq \emptyset$$

Let  $\mathcal{D}$  be the collineation group of  $\mathbb{P}(E \otimes \mathbb{C}^h)$  in itself represented by homogeneous diagonal matrices:  $\text{diag}(\underbrace{\lambda_1 \dots \lambda_1}_{n+1}; \underbrace{\lambda_2 \dots \lambda_2}_{n+1}, \dots, \underbrace{\lambda_h \dots \lambda_h}_{(n+1)})$  with  $h$  non zero scalars  $\lambda_j$ ,  $j = 1, 2, \dots, h$ .

Then  $U$  and  $\gamma U$  represent the same correspondence for any

$$\gamma = D_{\lambda_1 \lambda_2 \dots \lambda_h} \in \mathcal{D}.$$

EXAMPLE

The diagonal space  $\Delta$  has the two properties we want:  $\Delta \in \mathbb{P}(J_p)$  and  $\Delta$  does not contain two different points of the same  $\mathbb{P}_{h-1}$  of  $J(\mathbb{P}(E) \times \dots \times \mathbb{P}(E))$ . In this case  $R_\Delta = \Sigma$ .  $\Delta$  and  $\Sigma$  represent both the *diagonal* ( $\Leftrightarrow$  "identity") in the abstract product  $\mathbb{P}(E) \times \dots \times \mathbb{P}(E)$ .

However there are other linear spaces  $\mathbb{P}(E \otimes \mathbb{C}^h)$  having this property, for instance those (replacing  $\Delta$ ) obtained "moving" the  $h$  identifications

$i_j : \mathbb{P}(E) \longrightarrow \mathbb{P}(S_j)$ . Let us replace them by  $h$  arbitrary non-degenerate collineations  $\gamma_j : \mathbb{P}(E) \longrightarrow \mathbb{P}(S_j)$ ,  $j = 1, 2, \dots, h$ . Then we have: The correspondence  $\gamma$ , locus of  $(\gamma_1(P), \gamma_2(P), \dots, \gamma_h(P)) \in \mathbb{P}(E)$  will be called a *non-degenerate h-collineation*. It is represented by a SEGRE variety  $\Sigma_\gamma$  (reducing to  $\Sigma$  for  $\gamma = i_j$ ,  $j = 1, 2, \dots, h$ ) whose vertical  $(h-1)$ -spaces belong to  $J(\mathbb{P}(E) \times \dots \times \mathbb{P}(E))$ . Any horizontal one  $H \neq \mathbb{P}(S_1), \mathbb{P}(S_2), \dots, \mathbb{P}(S_h)$  represents  $\gamma$ , i.e.  $H \subset (J_p) : H$  has no two different points in the same  $\mathbb{P}_{h-1}$  of the join and  $R_H = \Sigma_\gamma$ .

In the case  $h = 2$   $\gamma_2 \gamma_1^{-1} (\gamma_1 \gamma_2^{-2})$  represent a collineation  $\mathbb{P}(S_1) \longrightarrow \mathbb{P}(S_2)$  (or its inverse  $\mathbb{P}(S_2) \longrightarrow \mathbb{P}(S_1)$ ).

Let us see this properties more closely using basis:

Let  $B_j(u_0^{(j)}, u_1^{(j)}, \dots, u_n^{(j)})$ ,  $j = 1, 2, \dots, h$  be a basis of  $E (\Leftrightarrow \bigwedge_{i=0}^n u_i^{(j)} \neq 0 \quad j = 1, 2, \dots, h)$ . Then we have:

Such  $h$  bases define a *non-degenerate h-collineation* where  $(x_1, x_2, \dots, x_h)$  correspond if  $x_j$  has the same homogeneous coordinates in  $B_j$  for  $j = 1, 2, \dots, h$ . But the  $h$  vectors

$$(u_j^{(1)}, u_j^{(2)}, \dots, u_j^{(h)}) \in J =$$

are linearly independent and they define a  $S_n \subset \mathbb{P}(E \otimes \mathbb{C}^h)$ . The  $h$  bases  $(\lambda B_1, \lambda B_2, \dots, \lambda B_h)$  define the same  $S$  for any  $\lambda \neq 0$ . Let  $\lambda_1, \lambda_2, \dots, \lambda_h$  be  $h$  different non zero scalars. Then  $(\lambda_1 B_1, \lambda_2 B_2, \dots, \lambda_h B_h)$  define a different  $S'_n = D S_n$  where  $D = D_{\lambda_1, \lambda_2, \dots, \lambda_h}$ . But  $S_n$  and  $S'_n$  define the same  $h$ -collineation.

EXAMPLE. For  $h = 2$  we have:

If  $(\lambda, \mu) \neq (0, 0)$   $(\lambda B_1, \mu B_2)$  define a subspace  $S_{n-1}$  representing the non degenerate collineation  $(B_1, B_2)$ .  $(\lambda', \mu')$  defines the same  $S_{n-1}$  iff  $(\lambda', \mu') = \nu (\lambda, \mu)$ .

$(B_1, B_2)$  and  $(B'_1, B'_2)$  define the same collineation iff  $B'_1 = B_1 T$   $B'_2 = B_2 T$  where  $T$  is a  $(n+1) \times (n+1)$  matrix with  $\det T \neq 0$ .

Then we can see that  $R_S = R_{S'}$  is a SEGRE variety.

Let us introduce back coordinate systems  $(x_0, x_1, \dots, x_n)$  in  $E$  as well as  $(x_0^{(j)} \dots x_n^{(j)})$  in  $S_j$  interpreted as homogeneous coordinates when needed. Then for  $\gamma_1, \gamma_2, \dots, \gamma_n$  non degenerate we can assign to any set of  $h$  non singular matrices  $G_1, G_2, \dots, G_h$  the  $n$ -subspaces  $S_n$  of  $\mathbb{P}(E \otimes \mathbb{C}^h)$  generated by the  $n+1$  rows of  $G_1 G_2 \dots G_h$ . The non singularity condition  $\det G_j \neq 0$  is equivalent to the fact that  $S_n \cap (S_j) = \emptyset$  where

$$S_j = E \dots \underset{j}{0} \dots E.$$

Thus  $S_n \subset \mathbb{P}(J_p) \Leftrightarrow \det G_j \neq 0$  for  $j = 1, 2, \dots, n$ .

$$(G_1 G_2 \dots G_h) \text{ and } (G_1 T G_2 T \dots G_h T)$$

are two different bases of  $S_n$  if  $\det T \neq 0$ , and we can assume either one  $G_j = \mathbb{P}_n$ . If  $\lambda_j \neq 0$  for  $j = 1, 2, \dots, n$  ( $\lambda_1 G_1, \lambda_2 G_2, \dots, \lambda_h G_h$ ) defines  $D_{\lambda_1 \lambda_2 \dots \lambda_h} S_n$  with

$$D_{\lambda_1 \dots \lambda_h} \in \mathcal{D}.$$

Let us forget now the condition  $\det G_j \neq 0$  for some (or all  $j$ ) but keeping the fact that  $\text{rank}(G_1, G_2 \dots G_h) = n + 1$ . Then the condition  $S_n \subset \mathbb{P}(J_p)$  fails  $\Leftrightarrow S_n$  meets some  $\mathbb{P}(S_j)$ . However we can assign to  $S_n$  a correspondence  $\Gamma(S_n)$  where  $(x_1, x_2, \dots, x_h) \in \Gamma$  iff the  $\mathbb{P}_{h-1}$  space  $\{\lambda_1 x_1 + \mu_2 x_2 + \dots + \mu_h x_h\}$  meets  $S_n$  (we cannot insure anymore that it meets in a single point).

EXAMPLE. Let  $S_\alpha, S_\beta$  be two subspaces of  $\mathbb{P}_n$ . Then  $J(S_\alpha \times S_\beta)$  is a subspace of dimension  $\alpha + \beta + 1$  of  $\mathbb{P}(E \oplus E)$ , but  $J(S_\alpha \times S_\beta) \cap \mathbb{P}(S_1) = i_1(S_\alpha)$   
 $J(S_\alpha \times S_\beta) \cap \mathbb{P}(S_2) = i_2(S_\beta)$  if  $(x) \in S_\alpha$   $(y) \in S_\beta$  the whole line  $\lambda(x, 0) + \mu(0, y)$  is contained in  $J(S_\alpha \times S_\beta)$ .

## II. GENERALITIES ON THE COMPLEX $\mathcal{C}(V)$ ATTACHED TO A $V \subset \mathbb{P}_n$ .

We shall complete with appropriate references some of the information already given in the Introduction. It is wellknown that not every complex in  $\mathcal{G}(c-1; n)$  is attached to a  $V$ . Such particular complexes are indeed very special; they will be called *nucleated* with nucleus  $V^c$ . The characteristic nuclearity conditions for a  $\mathcal{C} \subset \mathcal{G}(c-1; n)$  can be expressed by a system of homogeneous polynomial equations – the so called CHOW equations (cf. [Ch–vdW]) they are use; to prove that the set of positive cycles of codimension  $c$  in  $\mathbb{P}_n$  is ZARISKI closed.

5. THE COMPLEX  $\mathcal{C}(V^c)$  OF  $\mathbb{P}_{c-1}$ . RECALL ON ZUGEORDNETE FORMEN.

The word *complex of subspaces*  $\mathbb{P}_d$  in  $\mathbb{P}_n$  ( $0 \leq d \leq n$ ) is used here in the XIX<sup>th</sup> century sense—namely as a synonym of *Grassmann divisor* (in  $\mathcal{G}(d; \mathbb{P}_n)$ ). We identify  $\mathcal{C}$  with its image in the Grassmann embedding

$$(5.1) \quad \mathcal{G} = \mathcal{G}(d; n) \longleftrightarrow \mathbb{P}(\overset{d+1}{\wedge} E_{n+1}) \quad \mathbb{P}_n = \mathbb{P}(E_{n+1}).$$

A  $\mathbb{P}_d(\mathbb{C} \mathbb{P}_n)$  can be determined uniquely by  $d + 1$  linearly independent points in  $\mathbb{P}_n$  or by  $n - d$  l.i. hyperplanes meeting at  $\mathbb{P}_d$ . Accordingly we define the *conjugation conditions* with respect to a complex  $\mathcal{C}$  of  $d$ -spaces in  $\mathbb{P}_n$  as follows:

DEF. 5.1.  $d + 1$  linearly independent points  $P_1, P_2, \dots, P_{d+1}$  of  $\mathbb{P}_n$  are called *conjugate with respect to  $\mathcal{C}$*  iff if the unique  $S_d \ni P_j$  ( $j = 1, 2, \dots, d + 1$ ) belongs to  $\mathcal{C}$ .

DEF. 5.2  $n - d$  linearly independent hyperplanes  $H_1, H_2, \dots, H_{n-d}(\mathbb{C} \mathbb{P}_n)$  are called *conjugate with respect to  $\mathcal{C}$*  iff the unique  $S_d = H_1 \cap H_2 \cap \dots \cap H_{n-d}$  belongs to  $\mathcal{C}$ .

The conjugation condition of  $d + 1$  points with respect to an irreducible  $\mathcal{C}(\mathbb{C} \mathcal{G}(d; n))$  (cf. DEF. 5.1) can be determined by a single irreducible equation

$$(5.2) \quad F(x_1, x_2, \dots, x_{d+1}) = 0$$

where  $F$  is a polynomial homogeneous of the same degree  $g$  with respect to each one of the  $d + 1$  variable vectors  $x_j \in E_{n+1}$  representing the points  $P_j$ ,  $j = 1, 2, \dots, h$ .

Similarly we have another plurihomogeneous form  $G$  (with the same  $g$  for the

$n - d$  variables  $u^j \in \check{E}$  (dual of  $E_{n+1}$ ), such that

$$(5.3) \quad G(u^1, u^2, \dots, u^{n-d}) = 0$$

characterizes the conjugation condition of the  $H_j$  ( $= \mathbb{P}(u^j)$ ),  $j = 1, 2, \dots, n - d$ ; of DEF. 4.2.  $F$  and  $G$  can be written uniquely as  $\mathbb{C}$ -linear combination of standard monomials  $p(S)$ ,  $q(\Sigma)$  of degree  $g$  (cf. [H-P] vol. II, Ch. XIV, page 377) in the Grassmann coordinates of  $\mathbb{P}_d(\mathbb{P}_d^\perp)$   $p^{i_1 i_2 \dots i_{d+1}}$ ,  $(q_{j_1 j_2 \dots j_{n-d}})$

$$(5.4) \quad F = \sum \lambda_S p(S) \quad G = \sum_{\Sigma} \mu_{\Sigma} q(\Sigma)$$

$F$  and  $G$  are uniquely determined by  $\mathcal{C}$  (up to a  $\mathbb{C}^\times$ -factor). Accordingly  $(\lambda_1, \lambda_2, \dots)$  or  $(\mu_1, \mu_2, \dots)$  are well defined homogeneous coordinates representing  $\mathcal{C}$ . The procedure is extended to arbitrary positive Grassmann divisors by prime factor decomposition  $F = \prod F_j^{m_j}$ ,  $G = \prod G_j^{m_j}$ . Both expressions (4.4) are not essentially different because of the well known identities between the  $p$  and  $q$ .

When  $\mathcal{C} = \mathcal{C}(V)$  ( $d = c - 1$ ) (cf. Introduction, page 6 and Abstract, page 4) these conjugation conditions (5.2), (5.4) define the CAYLEY-SEVERI *form* (or the CHOW *form* respectively) of  $V = V^c = V_d$ . We emphasize that the number of vectors ( $\Leftrightarrow$  belonging to  $E$ ) in (5.2) is equal to the codimension  $c$  of  $V^c$ , *thus it gives back the equation of a  $V^1$  (i.e. of a hypersurface), for  $c = 1$* . The CHOW forms of  $V$  contain a number of covectors (belonging to  $\check{E}$ ) equal to  $\dim V + 1$ . Then (5.2), resp. (5.3) *represent the characteristic conditions for a  $S$  to meet  $V$*  (where  $S_{c-1}$  is uniquely determined by  $c$  points, resp. as intersection of  $d + 1$  hyperplanes).

In order to introduce the formal definition for nucleated complexes (DEF. 5.5) we shall need to consider certain exceptional behaviour of points and  $S$ -spaces ( $m > d$ ) with respect to a complex of  $d$ -spaces .

DEF. 5.3. Let  $P$  be a point of  $\mathbb{P}_n$  .  $P$  is called *singular with respect to  $\mathcal{C}$*  iff every  $S_d \ni P$  belongs to  $\mathcal{C}$  .

DEF. 5.4. The subspace  $S_m$  ( $m > d$ ) of  $\mathbb{P}_m$  is called *singular with respect to  $\mathcal{C}$*  iff every  $S_d \subset S_m$  belongs to  $\mathcal{C}$  . Cf. [S] .

We shall introduce now formally *the complex  $\mathcal{C}(V^c)$  attached to an irreducible subvariety  $V^c$  of codimension  $c$  in  $\mathbb{P}_n$*  . It is necessary to check first the following property:

*The set*

$$(5.5) \quad \mathcal{C}(V^c) = \{ \mathbb{P}_{c-1} \subset \mathbb{P}_n \mid \mathbb{P}_{c-1} \cap V^c \neq \emptyset \} \subset \mathcal{G}(c-1; \mathbb{P}_n)$$

*is an irreducible complex in  $\mathcal{G}(c-1; \mathbb{P}_n)$  . The variety  $V^c$  is the locus of singular points of  $\mathcal{C}(V^c)$  (cf. [S], [H-P], vol. I, II); i.e.  $\mathcal{C}(V^c)$  is nucleated with locus  $V^c$  .*

DEF. 5.5. The set  $\mathcal{C}(V^c)$  defined by (5.6) is called *the complex attached to  $V^c$*  .

EXAMPLES.

1) For  $c = 1$   $\mathcal{C}(V^1)$  is just *the set of points of the irreducible hypersurface  $V^1 \subset \mathbb{P}_n$*  .

2) For  $c = 2$   $\mathcal{C}(V^2)$  *is the set of lines meeting  $V^2$*  . For instance if  $V^2 = \Gamma_1$  is an irreducible curve of  $\mathbb{P}_3$  ,  $\mathcal{C}(\Gamma)$  *is the complex of lines of  $\mathbb{P}_3$  meeting  $\Gamma$*  .



DEF. 5.6 *The conjugation conditions of points (or hyperplanes) with respect to  $\mathfrak{C}(V^c)$  are called the CAYLEY–SEVERI form (or CHOW – form) of  $V^c$ .*

$$(5.6) \quad S(x_1, \dots, x_c) = 0$$

$$(5.7) \quad Y(u^1, u^2, \dots, u^{d+1}) = 0$$

Actually  $Y = 0$  is the first systematic "*yugeordnete Form*" (cf. [vdW–ZAG]) or *associated form*. In the case of an irreducible plane curve  $\Gamma$  the left hand side of (5.7) is the resultant  $R(f; u, v)$  where  $f = 0$  represents  $\Gamma$  and  $u, v$  are linear forms.

In the introduction we mentioned also the *characteristic form* (WEIL) (or *Normalgleichung*) (SIEGEL) (valid also for a non nucleated  $\mathfrak{C}$ ) containing  $\dim V + 2$  covectors and a single vector; in the general case we have this "mixed" equation:

$$(5.8) \quad N(u_1, u_2, \dots, u_{d+2}; x) = 0.$$

#### REMARK

SEVERI pointed out in [S] that  $S = 0$  is the real generalization of the equation of an irreducible hypersurface  $V^1$ , since the number of vector variables equals the codimension. But  $S = 0$  was described by CAYLEY (as early as 1860) for conics in  $\mathbb{P}_3$  of  $[C_1], [C_2]$ . If we keep fixed  $c - 1$  linearly independent variables  $a_1 \dots a_{c-1}$  in  $S = 0$  in such a way that  $(a_1 \wedge \dots \wedge a_{c-1})$  does not meet  $V^c$  then

$$(5.9) \quad S(a_1, \dots, a_{c-1}; x) = 0$$

represents the projecting cone of  $V^c$  from  $\mathbb{P}_{c-2}$ ; accordingly  $V^c$  is recovered from  $S = 0$  as the intersection of all the projecting cones of  $V^c$  from the  $\mathbb{P}_{c-2}(\cap V^c = \phi)$ . If we replace  $a_1 \wedge \dots \wedge a_{c-1} \in \overset{c-1}{\Lambda} E$  by the corresponding  $u_1 \wedge \dots \wedge u_{d+2} \in \overset{d+2}{\Lambda} \check{E}$  we have the WEIL–SIEGEL equation (5.8)


$$(5.10) \quad N(u_1, u_2, \dots, u_{d+2}; \mathbf{x}) = 0$$

representing  $V^c = V_d$  as the intersection of all the projecting cones from generic spaces  $(u_1 \wedge \dots \wedge u_{d+2})$  non meeting  $V^c$ .

EXAMPLES (for an irreducible curve  $\Gamma$  in  $\mathbb{P}_3$ )

$$S(x_1, x_2) = 0 \quad Y(u, v) = 0 \quad N(u_0, u_1, u_2; \mathbf{x}) = 0$$

*represent*  $\Gamma$  *via the complex*  $\mathcal{C}(\Gamma)$  (cf. Fig. 2), where a line  $\ell \in \mathcal{C}(\Gamma)$  is defined by a couple of points (or of planes  $(c = 2, d = 1)$ ).  $\Gamma$  appears as intersection of all its projecting cones from outside points  $P = (a) = (u_0 \wedge u_1 \wedge u_2)$  given by a single  $a \in E$  or as intersection of three linearly independent planes  $(u_0), (u_1), (u_2)$



6. REVIEW ON ASSOCIATE FORMS. We shall recall here the main properties of the a.f. needed subsequently referring – for further details to the original papers [vdW–Ch], the Einführung [vdW 1] (with the 2<sup>nd</sup> historical appendix), the ZAG book, H–P II, CH X, § 6, 7, 8 and SEVERI'S comments in his paper on Grassmannians [S]. First of all there are uniquely defined linear combinations of the standard power products of Grassmann coordinates  $p^{i_1 i_2 \dots i_c}$  (or  $p_{j_0 j_1 \dots j_d}$ ) of  $x_1 \wedge x_2 \wedge \dots \wedge x_c$  (or  $u_0 \wedge u_1 \wedge \dots \wedge u_d$ ) in  $\overset{c}{\Lambda} E$  (or  $\overset{d+1}{\Lambda} \check{E}$ ) such that

$$(6.1) \quad S(x_1, x_2, \dots, x_c) = S(x_1 \wedge \dots \wedge x_c)$$

$$(6.2) \quad Y(u_0, u_1, \dots, u_j) = Y(u_0 \wedge u_1 \wedge \dots \wedge u_d)$$

and the transition between the right hand sides of (6.1), (6.2) is given by the well-known formulas of type

$$S_b(x_1, x_2, \dots, x_{c-k}) = S(x_1, x_2, \dots, x_{c-k} b_{c-k+1} \dots b_c)$$

where  $S_b$  is the CAYLEY–SEVERI form of the projecting cone of  $V$  from the  $S_{k-1}$ –subspace represented by  $(b_{c-k+1} \wedge \dots \wedge b_c)$  with  $S_{k-1} \cap V = \phi$ . The identical vanishing takes place iff  $S_{k-1} \cap V \neq \phi$ .

In particular for  $k = c - 1$  we obtain back the original CAYLEY'S idea of representing  $V$  as intersection of all its projecting cones of  $V$  from  $S_{c-2}$  projecting centers not meeting  $V$ .

The fact that  $V$  is the locus of singular points of the complex  $\mathcal{C}(V)$  gives raise to a canonical system of equations of  $V$  expressing the fact that for a point  $(a) \in V$  the equation of the projecting cone from  $(a)$  vanishes identically  $\Leftrightarrow (a)$  is a singular point of  $\mathcal{C}(V)$ .

In order to get the properties of the WEIL–SIEGEL form (*characteristic form = Normalgleichung*) it is convenient to represent the projection center  $S_{c-2} (\cap V = \phi)$  as a complete intersection of  $d + 2$  hyperplanes

$$(u_0) (u_1) \dots (u_{d+1}) \quad (\in \mathbb{P}(\check{E})).$$

This will give us an identity of type:

$$(6.4) \quad S(x_1, x_2, \dots, x_c) = N(u_0, u_1, \dots, u_{d+1}; x).$$

We shall give a more explicit expression of (6.4) using the fact that we can write:

$$(6.5) \quad v_0 \wedge v_1 \wedge \dots \wedge v_d = x \lrcorner u_0 \wedge u_1 \wedge \dots \wedge u_{d+1} .$$

(cf. [BOU]) if we normalize conveniently  $x$ , where the point  $(x)$  belongs to the intersection of the  $d+2$  linearly independent hyperplanes  $\langle u_j, x \rangle = 0$ ,  $j = 0, 1, \dots, d+1$ , namely

$$(6.4)' \quad S(x; x_2, \dots, x_c) = N(u_0, u_1, \dots, u_{d+1}; x) = \mathbb{N}(\xi_0, \xi_1, \dots, \xi_{d+1})$$

$$(6.6) \quad \xi_j = \langle u_j, x \rangle = \sum_{i=0}^n u_{ji} x_i \quad j = 0, 1, \dots, d+1 .$$

The form  $\mathbb{N}$  contains coefficients depending on  $u_0, u_1, \dots, u_{d+1}$  that can be determined explicitly. (6.5) has *the following remarkable geometric interpretation* :

Let  $\mathbb{P}(E/E_{c-1})$  be the quotient projective space of  $E$  with respect to the subspace  $E_{c-1}$  represented by  $x_2 \wedge \dots \wedge x_c \in \bigwedge^{c-2} E$  (which is also represented by  $(v_0 \wedge \dots \wedge v_{d+1}) (\in \bigwedge^{d+2} \check{E})$ ).

$$(6.7) \quad \dim E/E_{c-1} = d+2 \Leftrightarrow \dim \mathbb{P}(E/E_{c-1}) = d+1 .$$

The  $d+2$  forms  $u_j$  linearly independent of  $\check{E}$  can be regarded also as forms in  $E/E_{c-1}$  because  $E_{c-1}$  is defined by  $\langle u_j, x \rangle = 0$  for,  $j = 0, \dots, d+1$  :  
 $\langle u_j, x \rangle = \langle u_j, x + y \rangle \quad \forall y \in E_{c-1} .$

As a consequence: *The  $d+2$  forms  $\xi_j$  ( $j = 0, 1, \dots, d+1$ ) are homogeneous coordinates in the quotient projective space  $\mathbb{P}_{d+1} = \mathbb{P}(E/E_{c-1})$  and .*

*The equation:*

$$(6.8) \quad \mathbb{N}(\xi_0, \xi_2, \dots, \xi_{d+1}) = 0$$

cf. (6.4)' represents a hypersurface model of  $V_d$  lying in  $\mathbb{P}_{d+1} = \mathbb{P}(E/E_{c-1})$  whose points are naturally mapped to the generators of the projecting cone of  $V$  from  $\mathbb{P}(E_{c-1})$ .

Since  $E_{c-1}$  can be any vector subspace of  $E$  such that  $\mathbb{P}(E_{c-1}) \cap V = \phi$  we have a refinement of CAYLEY'S idea in the sense that given *one of those* CAYLEY'S projection centers  $\mathbb{P}(E_{c-1})(\cap V = \phi)$  (6.8) defines an ordinary equation of a hypersurface  $H_{E_{c-1}}$  model of  $V$ , for every choice of forms  $u_j$  ( $j = 0, 1, \dots, d+1$ ) defining  $E_{c-1}$ .

The points of this hypersurface correspond bijectively with the  $\mathbb{P}_{c-1}$  generators of the CAYLEY cone of center  $\mathbb{P}(E_{c-1})$ . For a generic choice of  $\mathbb{P}(E_{c-1})$  the generic generator of this projecting cone contains just one point of  $V$ , the exceptional ones correspond bijectively with the singular points of  $H_{E_{c-1}}$ . In particular if  $d = \dim V = n - 1$  the  $n + 1$  linearly independent forms  $u_0, u_1, \dots, u_n$  in  $\check{E}$  define a coordinate system in  $\check{E} \Rightarrow$  a projective system in  $\mathbb{P}_n$ ; thus in this case

$$(6.9) \quad \mathbb{N}(\langle u_0, x \rangle, \langle u_1, x \rangle, \dots, \langle u_n, x \rangle) = 0$$

with  $\langle u_j, x \rangle = \sum_{k=0}^n u_{jk} x^k$  defines the equation of the hypersurface  $V$  in this

coordinate system, or in the language of invariants:

(6.9) represents all the possible equations of the hypersurface  $V$  for all the choices with  $u_0 \wedge \dots \wedge u_n \neq 0$ .

### EXAMPLES

In the case of Fig. 2, page 41, any triple of linearly independent linear forms  $u_0$ ,  $u_1$ ,  $u_2$  define a projective coordinate system with  $(u_j)$  ( $j = 0,1,2$ ) as coordinate planes and  $(u_0 + u_1 + u_2)$  as the unit "line" in the abstract plane  $\mathbb{P}(E_4/E_1)$ , where  $P = \mathbb{P}(E_1)$  is any point of  $\mathbb{P}_3 = \mathbb{P}(E_4)$  outside  $V$  intersection of the three planes  $\langle u_j, x \rangle = 0$ . The equations

$$\mathbb{N}(\xi_0, \xi_1, \xi_2) = 0 \quad N(\langle u_0, x \rangle, \langle u_1, x \rangle, \langle u_2, x \rangle) = 0$$

represent a model of  $V$  in  $\mathbb{P}(E_4/E_1)$  or the projecting cone of  $V$  with vertex  $P$ .

### REMARK

The explicit computation of  $\mathbb{N}$  in terms of  $\mathbb{S}$  can be achieved expressing (6.5) in coordinates; replacing  $p^{i_0 i_1 \dots i_d}$  by

$$(6.15)' \quad p^{i_0 i_1 \dots i_d} = \sum x^i q^{i i_0 i_1 \dots i_d}$$

where  $q^{j_0 j_1 \dots j_{d+1}}$  are the coordinates of  $x \perp u_0 \wedge u_1 \wedge \dots \wedge u_{d+1}$  leading to

$$(6.10) \quad p^{i_0 i_1 \dots i_d} = \begin{vmatrix} \langle u_0, x \rangle & u_{0 i_0} & u_{0 i_1} & \dots & u_{0 i_d} \\ \langle u_1, x \rangle & u_{1 i_0} & u_{1 i_1} & \dots & u_{1 i_d} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \langle u_{d+1}, x \rangle & u_{d+1, i_0} & u_{d+1, i_1} & \dots & u_{d+1, i_d} \end{vmatrix}.$$

### III. APPLICATIONS

The construction of  $J = J(V^{(1)} \times \dots \times V^{(h)})$ ,  $V^{(j)} \subset \mathbb{P}_n$ ,  $j = 1, 2, \dots, h$ , has two natural applications depending on the codimension of  $J$  in  $\mathbb{P}(E \otimes \mathbb{C}^h)$ . Cf. (2.12), page 27; if  $c \leq n$  the  $h$  given varieties  $V^{(j)}$  always meet in  $\mathbb{P}_n$  ( $\Leftrightarrow \Delta$  always meet  $J$  in  $\mathbb{P}(E \otimes \mathbb{C}^h)$ ). If  $c > n$  the given varieties do not meet ( $\Leftrightarrow \Delta \cap J = \emptyset$ ) if they are in generic position, but the discussion of their meeting gives a new form to the old "elimination theory" which can be made intrinsic. We shall divide the paper in two parts according to both possibilities:

In part I, page 46, § 7, 8, 9, 10 we deal with the case  $c \leq n$ . If  $S \in \mathcal{C}(J)$ ,  $J = J(V^{(1)} \times \dots \times V^{(h)})$ , since  $\dim S = c - 1 < n$ , it makes sense to introduce the restriction to the diagonal space  $\Delta$  in  $\mathbb{P}(E \otimes \mathbb{C}^h)$  (cf. § 3, page 27). Such a restriction is trivial ( $\Leftrightarrow \mathcal{C}(J)|\Delta = \mathcal{G}(c-1; \Delta)$ ) iff the intersection  $\bigcap_{j=1}^h V^{(j)}$  is improper. Otherwise, there is a well defined complex  $\mathcal{C}(J)|\Delta$  whose pull-back to  $\mathbb{P}(E)$  by  $\delta^{-1}$  gives the natural definition of  $\mathcal{C}(I)$ , where  $I = V^{(1)} \cdot V^{(2)} \cdot V^{(h)}$  is the intersection cycle. The prime factor decomposition gives the intersection multiplicity as the exponent of either one of  $S_C$ ,  $Y_C$  or  $N_C$  (any two of them agree) for any irreducible component  $C$  of  $I$ . See our main definition DEF. 7.1, page 51. In particular we can prove BEZOUT'S theorem since  $\deg J = \prod_{j=1}^h \deg V^{(j)}$  can be proved with a rigorous degeneration method using the characteristic transversality condition for multiplicity one.

*The announced equivalence of the exponent intersection multiplicity with the original one of VAN DER WAERDEN follows easily from the interpretation of the  $S_n$  subspaces of  $\mathbb{P}(E \otimes \mathbb{C}^h)$  as representatives of  $h$ -collineations (cf § 4).*



FIRST PART

THE EXPONENT INTERSECTION MULTIPLICITY

7. RESTRICTION TO THE DIAGONAL OF  $\mathcal{C}(J(V^{(1)} \times V^{(2)} \times \dots \times V^{(h)}))$ ,  $c \leq n$ .

Let us come back to the constructions of § 0,1. Let  $J = J(V^{(1)} \times \dots \times V^{(h)})$  be the join of the  $h$  shown irreducible varieties of codimensions  $c_j$  in  $\mathbb{P}_n$  satisfying (0.7). Let us consider the complex  $\mathcal{C}(J)$  of  $(c-1)$ -dimensional subspaces of  $\mathbb{P}(E \otimes \mathbb{C}^h) = \mathbb{P}_{h(n+1)-1}$  attached to  $J$ :

$$(7.1) \quad \mathcal{C}(J) = \{ \mathbb{P}_{c-1} \subset \mathbb{P}(E \otimes \mathbb{C}^h) \mid \mathbb{P}_{c-1} \cap J \neq \emptyset \} \subset \mathcal{G}(c-1; h(n+1)-1)$$

then, since  $c \leq n$  the restriction to the  $n$ -dimensional diagonal space  $\Delta$ :

$$(7.2) \quad \mathcal{C}(J) \mid \mathcal{G}(c-1, \Delta)$$

makes sense. We shall distinguish two cases:

1) *If  $\bigcap_{j=1}^h V^{(j)}$  is improper  $\Leftrightarrow$  if there is at least one excedentary irreducible component  $X \Leftrightarrow \text{cod } X < n$  then*

$$\delta(X) \cap \mathbb{P}_{c-1} \neq \emptyset, \quad \forall \mathbb{P}_{c-1} \subset \Delta.$$

2) On the contrary: *If*  $\bigcap_{j=1}^h V^{(j)}$  *is proper we can construct some subspace*  
 $\mathbb{P}_{c-1} \subset \Delta$  *satisfying*

$$\delta(\bigcap V^{(j)}) \cap \mathbb{P}_{c-1} = \phi .$$

It suffices to take the diagonal image of a  $\mathbb{P}_{c-1}$  of  $\mathbb{P}_n$  not meeting  $\bigcap_{j=1}^h V^{(j)}$ .

In other words, we have proved the following:

LEMMA. *The diagonal space*  $\Delta$  *of*  $\mathbb{P}_{h(n+1)-1}$  *(cf. § 3) is a singular space of*  $\mathcal{C}$   
 (J) *iff*  $\bigcap_{j=1}^h V^{(j)}$  *is improper. Otherwise the restriction of*  $\mathcal{C}(J)$  *to*  $\Delta$  *defines the*  
*complex (7.4) below which will be attached to the intersection cycle*  $I$  *by the formula*

$$(7.4) \quad \mathcal{C}(I) = \delta^{-1}(\mathcal{C}(J)|\Delta)$$

where  $I = V^{(1)} \cdot V^{(2)} \cdot \dots \cdot V^{(h)}$ .

#### MAIN DEFINITION 1:

DEF. 7.1 *The complex of*  $S_{c-1}$  *subspaces of*  $\mathbb{P}_n$  *defined by (7.4) is called the complex*  
*attached to the (well-defined) intersection cycle*  $I = V^{(1)} \cdot V^{(2)} \cdot \dots \cdot V^{(h)}$  *of the*  $h$   
~~*given properly intersecting varieties.*~~

#### REMARK

The effective restriction  $\mathcal{C}(J) = \delta^{-1}(\mathcal{C}(J)|\Delta)$  can be achieved by means of either one of the associated forms discussed in § 6, namely:

We know that the SEVERI form  $S_J$  attached to  $J$  contains  $c$  covariant vector variables  $x_1, x_2, \dots, x_c$ . It suffices to take  $x_j \in \Delta$  for  $j = 1, 2, \dots, c$  to get the desired restriction. For  $Y_j$  there are  $d + 1$  hyperplane variables representing a  $\mathbb{P}_{c-1} \in \mathcal{C}(J)$  where  $d = h(n + 1) - 1 - c$ . The corresponding number for a  $\mathbb{P}_{c-1} \subset \Delta$  is  $n - c + 1$ . The difference  $(n + 1) - (h - 1)$  equals the number of equations of type

$$(7.5) \quad x_j^{(r)} - x_j^{(1)} = 0 \quad r = 2, \dots, h; \quad j = 0, 1, \dots, n$$

defining  $\Delta$ . Thus, we shall define a  $\mathbb{P}_{c-1} \subset \Delta$  with forms containing the (7.4). The rest define the same  $\mathbb{P}_{c-1}$  as a subspace of  $\Delta$ .

Similarly the WEIL-SIEGEL form suffices to restrict the generic projection center of dimension  $c - 2$  - in the ambient space of  $J$  - to the diagonal subspace  $\Delta$ .

In the three cases we have prime factor decompositions of  $S|\Delta$ ,  $Y|\Delta$ ,  $N|\Delta$  with prime factors  $S_C$ ,  $Y_C$ ,  $N_C$  attached bijectively to all the proper irreducible components  $C$  of  $\prod_{j=1}^h V^{(j)}$  and equal exponents  $i_C$ :

$$(7.6) \quad S_I = \prod S_C^{i_C} \quad Y_I = \prod Y_C^{i_C} \quad N_I = \prod N_C^{i_C}.$$

Such equality is indeed a consequence of the transformation formulas between  $S_I$ ,  $Y_I$ ,  $N_I$  studied in § 6.

## MAIN DEFINITION 2

DEF. 7.2. The positive integer  $i_C$  well defined by either one of the (7.5) in an intrinsic way is called the *exponent intersection multiplicity of  $C$  in  $I$*  (cf. (7.4)).

8. COMPUTATION OF  $F_j$ . BEZOUT'S THEOREM. The computation of the CHOW form  $Y_V$  on any irreducible  $V \subset \mathbb{P}_n$  is based on the theory of the u-resultant (cf. [H-P], I). It can be applied to  $J = J(V \times W)$  when we give any two systems of equations in  $(x)$ ,  $(y)$  to represent  $V$  and  $W$ . From  $Y_V$  we can construct  $S_V$  and  $N_V$ . A direct computation of any  $S_V$  with  $\text{cod } V = c$  can be obtained by

$$(8.1) \quad S_V(x_1, x_2, \dots, x_c) = \text{h.c.d} (\dots, R_k, \dots)$$

where the  $R_k$  are resultant forms with respect to  $\lambda_1, \dots, \lambda_c$  in the equations

$$f_k \left( \sum_{j=1}^c \lambda_j x_j \right) = 0$$

obtained by the specialization  $x \longmapsto \sum_{j=1}^c \lambda_j x_j$  in the equations  $\dots f_k(x) = 0 \dots$  representing  $V$ .

REMARK *It is remarkable, very simple and essentially "new" (since the N-form is not widely used in the literature) that the equation*

$$(8.2) \quad N \left( \sum_{k=0}^n u_{jk} x_k \right) = 0 \quad j = 0, 1, \dots, d+1$$

*can be obtained immediately observing that the transcendence degree of the projecting cone  $\Gamma(V)$  of  $V$  is equal to  $d+1$ . Accordingly: The  $d+2$  restrictions*

$$\left( \sum_{k=0}^n u_{jk} x_k \right) \in \Gamma(V)$$

$j = 0, \dots, d + 1$  are algebraically dependent.

For instance, let  $V$  be an irreducible algebraic curve in  $\mathbb{P}_n$ . Then we write immediately an irreducible equation

$$(8.3) \quad F(\langle u_{1,x} \rangle, \langle u_{2,x} \rangle, \langle u_{3,x} \rangle) = 0$$

representing  $V$  as intersection of all the projecting cones from  $\mathbb{P}_{n-3}$  - subspaces, complete intersections of the three hyperplanes.

$$(8.4) \quad \langle u_{j,x} \rangle = 0, \quad j = 1, 2, 3$$

where  $\langle u_{j,x} \rangle = \sum_{k=0}^n u_{jk} x_k = 0$ .

In particular if  $V$  is a canonical curve - non hyperelliptic - of genus  $g$  in  $\mathbb{P}_{g-1}$  we can take three generic holomorphic differentials to define the WEIL-SIEGEL form. We shall apply elsewhere this remark to the SCHOTTKY problem, cf. [G.4].

#### REMARK

The following natural question arises; let  $F_j$  be associate forms (of the same kind  $S, Y, N$ ) corresponding to  $h$  algebraic irreducible  $V^{(j)} \subset \mathbb{P}_n$ . Can we compute  $F_J$  in terms of the  $F_j$ ? (where  $J = J(V^{(1)} \times \dots \times V^{(h)})$ ). If the  $V^{(j)}$  are all hypersurfaces:  $c_j = 1$  and  $c = h \leq n$ , the answer is positive because  $F_J =$  Resultant form with respect to  $\lambda_1, \lambda_2, \dots, \lambda_0$  of the  $c$  equations



$G(\lambda x + \mu y) = \sum_{j=0}^g b_j \lambda^j \mu^{f-j}$  where  $F(x) = 0$ ,  $G(x) = 0$  are the equations of the projecting cones of  $V(W)$  from  $\mathbb{P}_{c-2}$  ( $\mathbb{P}_{c_2-2}$ ) respectively. An immediate consequence of this property is the following:

*The degree of  $J(V \times W)$  is equal to the product of the degrees of  $J(V)$ ,  $J(W)$*

$$(8.7) \quad \deg J(V \times W) = \deg V \cdot \deg W .$$

The intersection  $J \cap \Delta$  has the same degree; accordingly we have:

BEZOUT'S THEOREM. *Let  $V \cdot W$  be the intersection cycle of two irreducible algebraic varieties  $V$ ,  $W_{c_n}$  meeting properly. We have:*

$$(8.8) \quad \deg V \cdot W = \deg V \cdot \deg W$$

*as a consequence of (8.7).*

9. ON THE PROOF OF THE THEOREMS. In the expository part of the introduction and in the exposition of the adaptation of the reduction to the diagonal in the projective case (§ 1,2,3) we gave already all the necessary ingredients to prove Th. I. but – since, there we lacked some technical tools, for instance the relation between the diagonal space  $\Delta$  and the diagonal  $\Sigma$ , (cf. (9.1), below) the more precise recall on associate forms, etc.:

$$(9.1) \quad \Sigma = \text{SEGRE variety} = \mathbb{P}\{x \otimes (\lambda_1 \dots \lambda_h) \mid x \in E, (\lambda_1 \dots \lambda_h) \in \mathbb{C}^h\}$$

with  $\Delta = \mathbb{P}(E \otimes (1,1, \dots, 1))$  - it is convenient for the reader to have now a complete version of the proof. On the other hand with the same procedure we shall prove also Th. II, although we shall come back to it in § 12.

Proof of Th. I: If the set-theoretic intersection  $\bigcap_{j=1}^h V^{c_j}$  with  $c = \sum_{j=1}^h c_j \leq n$  is improper there is at least one excedentary irreducible component  $X$  of codimension less than  $c$ ; as a consequence every  $\mathbb{P}_{c-1} \subset \mathbb{P}_n$  meets  $I$ ; this is equivalent to the fact that every  $\mathbb{P}_{c-1} \subset \Delta$  meets  $\delta(I)$  i.e.  $\Delta$  is a singular space for  $\mathcal{C}(J(C))$ , where

$$\mathcal{I}(C) \leq \mathcal{G}(c-1; \mathbb{P}(E \otimes \mathbb{C}^h)).$$

cf. DEF. 5.4 page 37.

On the other hand iff  $\bigcap_{j=1}^h V^{c_j}$  is proper it is always possible to find a  $\mathbb{P}_{c-1} \subset \mathbb{P}_n$  such that  $\mathbb{P}_{c-1} \cap (\bigcap_{j=1}^h V^{(j)}) = \emptyset$ . This implies there exists some  $\mathbb{P}_{c-1} \subset \Delta$  which do not meet  $J(V^{(1)} \times \dots \times V^{(h)})$ ; in other words we have a proper restriction to  $\Delta$  of the complex  $\mathcal{C}(J)$  attached to  $J$  iff  $\bigcap$  is proper; such restriction can be effectively computed by restriction to  $\Delta$  of either one of the equivalent form  $S_J, Y_J, N_J$ . The prime factor decompositions determine uniquely the exponents of the irreducible factors  $C$  (cf. § 5,6); each irreducible factor has the form  $S_C, Y_C$  on  $N_C$  appearing with the same well defined exponent

$$i_I = i(V^{(1)}, V^{(2)}, \dots, V^{(h)}; I).$$



10. EQUIVALENCE OF THE EXPONENT MULTIPLICITY WITH VAN DER WAERDEN'S THEORY.

The exponent multiplicity theory enables an easy transition (in both directions) between the so-called *static* and the *dynamic* multiplicity theories, (cf. [F]) roughly speaking it is equivalent *to move the intersecting varieties*  $V, W$  *or to move the diagonal space*  $\Delta$ . But  $\Delta$  belong to  $\mathcal{G}(n; 2n + 1)$  and such *motion* is quite well understood. On the other hand a generic  $S_n \subset \mathbb{P}(E \oplus E)$  represents a non singular collineation  $\gamma$  in  $\mathbb{P}_n$  (where  $S_n, S'_n$  represent the same collineations iff they are equivalent under the group  $\mathcal{D}$ ) of collineations of  $\mathbb{P}(E \oplus E)$  in itself (cf. § 4).

The original VAN DER WAERDEN'S multiplicity theory (cf. ZAG-papers, the historical survey [vdW2] and [H-P] (vol. II)) relies precisely in a motion of the pair  $(V, W)$  of irreducible subvarieties in  $\mathbb{P}_n$  to  $(\gamma_1 V, \gamma_2 W)$  by means of generic collineations  $\gamma_1, \gamma_2 \in \mathbb{P} GL(E)$ .  $(\gamma_1 V, \gamma_2 W)$  gives essentially the same as  $(\gamma \gamma_1 V, \gamma \gamma_2 W)$  where  $\gamma \in GL(E)$ , thus we can consider also  $(V, \gamma_1^{-1} \gamma_2 W)$  or  $(\gamma_2^{-1} \gamma_1 V, W)$  (with the inconvenience of a subsequent proof of the symmetry of  $i(I; V.W)$  when we permute  $V$  and  $W$ . Anyway the intersection multiplicity  $i(I; V.W)$  (for  $I$  irreducible component of  $V \cdot W$ ) is defined in vdW s' theory by specialization when  $(\gamma_1 \gamma') \longrightarrow$  Identity.

The equivalence of the exponent multiplicity with van der WAERDEN'S appears naturally when we replace the motion of  $\gamma$  with the (equivalent) "motion of  $\Delta$ ". We shall make explicit this equivalence:

We recall that a non singular collineation  $\gamma : \mathbb{P}_n \longrightarrow \mathbb{P}_n$  ( $\mathbb{P}_n = \mathbb{P}(E)$ ) is uniquely defined by a pair of bases  $B, B'$  where  $\gamma = \mathbb{P}(L)$ ,  $L \in GL(E)$  and  $L$  is uniquely

defined by  $B' = LB$ . Let us construct the  $n + 1$  vectors

$$(10.1) \quad (b_j, b'_j) \in E \oplus E \quad j = 0, 1, \dots, n.$$

They form a basis of the subspace  $E \oplus E$ . The bases  $(\lambda_1 B_1, \lambda_2 B_2)$ ,  $\lambda_1, \lambda_2 \in C^\times$  define another subspace  $D_{\lambda_1, \lambda_2} L$  representing the same collination  $\gamma = \mathbb{P}(L)$  as  $L$ .

$$D_{\lambda_1, \lambda_2} L = L \text{ iff } \lambda_1 = \lambda_2.$$

Two points  $P = (x)$ ,  $P' = (x')$  correspond in  $\gamma$  iff

$$(10.2) \quad x = \sum_0^n \lambda_j b_j \quad x' = \sum_0^n \lambda_j b'_j$$

with  $(\lambda_0, \lambda_1, \dots, \lambda_n) \neq (0, 0, \dots, 0)$ ;  $(x, x') \in L$ .

Conversely any point  $(\neq 0)$ .  $(x, x') \in L$  defines a pair of corresponding points  $(x)$ ,  $(x')$  in  $\mathbb{P}_n$ .

The set-theoretic intersection  $L \cap J(V \times W)$  can be interpreted as the set of pairs  $(x) \times (x') \in V \times W$  with  $(x') = \gamma(x)$ . The specialization  $\gamma \rightarrow \text{id}$ , will give back  $\Delta \cap J(V \times W)$  leading to the natural definition of the intersection cycle  $V \cdot W$ .

The precise nature of the equivalence between the exponent multiplicity and the original VdW's can be retraced quoting the following paragraph of the historical survey [vdW] or the Einführung [vdW] (page 276): "*Applying a projective transformation*":  
 ..... "*therefore I proposed in 1928 to bring  $V$  and  $W$  into a generic relative position by applying to one of them a projective transformation  $T$  with indeterminate coefficients. The transformed variety  $T$  intersects  $W$  in a finite number of points. If  $T$  is specialized to the identity, the points of intersection of  $TV$  and  $W$  specialize to*

*points of intersection of V and W. If V and W meet in a finite number of points, each of these appears with a certain multiplicity, which may be defined to be the intersection multiplicity ... "*

In order to adapt VdW's words to our procedure with the join in  $\mathbb{P}(E \oplus E)$  let us assume now  $\dim V.W = 0 \Leftrightarrow c = n$ . Then to the projective  $T$  let us associate the  $S_n \subset \mathbb{P}(E \oplus E)$  defined by the  $n + 1$  points of the  $(n + 1) \times 2(n + 1)$  matrix:

$$(10.3) \quad (1_{n+1} T)$$

where  $1_{n+1}$  is the  $(n + 1) \times (n + 1)$  unit matrix. The specialization  $T \longrightarrow$  Identity specializes (10.3) to  $(1_{n+1} \ 1_{n+1})$  defining the diagonal space.

In the general case for any  $c < n$ , the intersection of  $J(V \times W) \cdot S_c$  reduces the problem to  $J(V.S_c \times W.S_c) \subset S_c$  where  $V.S_c \subset S_c$ ,  $W.S_c \subset S_c$  and we have again the previous case:  $\dim V.W = 0$ .

In the discussion with the complex  $\mathcal{C}(J)$ , we need to consider a variable  $S_{c-1}$ . The same reduction to  $V.S_{c-1} \times W.S_{c-1} \subset \mathbb{P}(E_c \oplus E_c)$  ( $S_{c-1} = \mathbb{P}(E_c)$ ) leads to a case discussed in PART II of this paper, because now for the varieties in generic position  $V.S_{c-1} \cap W.S_{c-1} = \emptyset$  and  $\text{cod } J(V.S_{c-1} \times W.S_{c-1})$  in  $\mathbb{P}(E_c \oplus E_c)$  is equal to  $c = \dim S_{c-1} + 1$ .

## 11. BEZOUT'S THEOREM WITH A NEW DEGENERATION METHOD

The original discovery of the property.

$$(11.1) \quad \text{deg } F.G = f g \quad f = \text{deg } F, g = \text{deg } G$$

of the intersection cycle  $F.G$  of two irreducible algebraic curves in  $\mathbb{P}_2$  was obtained in a pure heuristic – non rigorous way – by degeneration of  $F, G$  in generic sets of  $f$  (resp.  $g$ ) different lines—intersecting together in  $fg$  simple points. I don't believe that anybody thought of this remark as a proof, but it has been always interesting whether this can be transformed indeed in a proof. We shall show here that by means of a certain degeneration (not of  $F, G$ , but of a secant space of complementary dimension) we can prove that

$$(11.2) \quad \text{deg } J(V \times W) = \text{deg } V \cdot \text{deg } W$$

where  $V, W$  are again two irreducible varieties  $V \subset \mathbb{P}(E_1), W \subset \mathbb{P}(E_2)$ . In fact it is well known that we can choose subspaces  $L \subset \mathbb{P}(E_1), M \subset \mathbb{P}(E_2)$  such that the intersection cycles consists of different simple points:

$$(11.3) \quad V.L = P_1 + P_2 + \dots + P_f \quad W.M = Q_1 + Q_2 + \dots + Q_g$$

$$f = \text{deg } V, g = \text{deg } W \quad P_i \neq P_j, Q_i \neq Q_j \quad i \neq j.$$

On the other hand the join  $J(L \times M)$  is a subspace of  $\mathbb{P}(E_1 \oplus E_2)$  of dimension equal to  $\text{cod } V + \text{cod } W + 1$ . The set theoretic intersection consists of  $fg$  lines  $J(P_i \times Q_j)$ .

$$(11.4) \quad J(V \times W) \cap J(L \times N) = \cup J(P_i \times Q_j)$$

$$i = 1, 2, \dots, f \quad j = 1, 2, \dots, g.$$

The transversality criterion for multiplicity one in each  $P_i$  or  $Q_j$  implies the transversality condition for the line  $J(P_i \times Q_j)$ . As a consequence we have:

$$(11.5) \quad J(V \times W) \cdot J(L \times N) = \sum J(P_i \times Q_j)$$

$$i = 1, 2, \dots, f \quad j = 1, 2, \dots, g.$$

In the same way we can see that we can choose a hyperplane  $\sum u_i x_i + \sum v_j y_j = 0$  in  $\mathbb{P}(E_1 \oplus E_2)$  transversal to each fixed  $J(P_i \times Q_j)$  because the opposite implies

$$(11.6) \quad \lambda(\sum u_k \xi_k^{(i)}) + \mu(\sum v_i \eta_i^{(j)}) = 0$$

where  $i = 1, 2, \dots, \deg V$ ,  $j = 1, 2, \dots, \deg W$ .

## SECOND PART: $c > n$

Let us consider now the case  $c > n$ . Then if the given irreducible  $V^{(j)} \subset \mathbb{P}_n$  are generically located the intersection is empty, i.e. we have

$$(11.7) \quad \bigcap_{j=1}^h V^{(j)} = \phi \Leftrightarrow J(V^{(1)} \times \dots \times V^{(h)}) \cap \Delta = \phi.$$

The complex  $\mathcal{C}(J)$  consists then of spaces of dimension  $c - 1 \geq n$  and our task is just to express the exceptional behaviour:

$$(11.8) \quad \bigcap_{j=1}^h V^{(j)} \neq \phi \Leftrightarrow J \cap \Delta \neq \phi$$

in terms of associate forms.

The extreme case  $c = n + 1$  appears in our treatment because then a  $\mathbb{P}_{c-1}$  is a  $\mathbb{P}_n$  and in particular the alternative (11.7) or (11.8) is equivalent to *the diagonal space  $\Delta$  does not belong to  $\mathcal{C}(J)$  iff the intersection is empty or  $\Delta \in \mathcal{C}(J)$  iff  $\bigcap_{j=1}^h V^{(j)} \neq \phi$ .*

For  $c > n + 1$  the property  $\Delta \cap J \neq \phi$  implies that every  $\mathbb{P}_{c-1}$  containing  $\Delta$  meets  $J$ :

$$\mathbb{P}_{c-1} \supset \Delta \Rightarrow \mathbb{P}_{c-1} \cap J \neq \phi$$

but the converse property is true:

*If every  $\mathbb{P}_{c-1} \supset \Delta$  meets  $J$  then  $\Delta$  meets  $J$  (equivalently if  $\Delta \cap J = \phi$  it is possible to find a  $\mathbb{P}_{c-1} \supset \Delta$  such that  $\mathbb{P}_{c-1} \cap J = \phi$ . This property leads naturally to express the condition  $J \cap \Delta \neq \phi$  by the identical vanishing of a covariant, as indicated in the introduction.*

12. A GEOMETRICAL THEORY FOR RESULTANT SYSTEMS.

In the particular case  $c_1 = c_2 = \dots = c_h = 1$ ,  $h = c > n$  we come back to the compatibility conditions of a system of  $h = c > n$  homogeneous polynomial equations

$$(12.0) \quad F_1 = 0 \quad F_2 = 0 \quad \dots \quad F_h = 0$$

of degrees  $m_1, m_2, \dots, m_h$ .

It is well-known that in the extreme case  $c = n + 1 = h$  the compatibility conditions are characterized by the vanishing on a single equation

$$(12.1) \quad R = 0$$

where  $R$  is a polynomial homogeneous of degree  $m/m_j$  in the  $\binom{n + m_j}{n}$  indeterminate coefficients of a generic form of degree  $m_j$  where

$$(12.2) \quad m = m_1 m_2 \dots m_{n+1}$$

$R = 0$  is equivalent to  $\Delta \cap J \neq \emptyset$  where

$$J = J(H_1 \times \dots \times H_{n+1})$$

as before and  $F_j = 0$  defines the irreducible hypersurface  $H_j$ ,  $j = 1, 2, \dots, n + 1$ .

Thus in the general case  $c = n + 1$ , the characteristic condition  $J \cap \Delta \neq \emptyset \Leftrightarrow \in \mathcal{C}(J)$  is a generalization of the equation  $R = 0$ .

In the case  $c > n + 1$  we checked already in the introduction that  $\Delta \cap J \neq \emptyset$  is equivalent to the identical vanishing of the CAYLEY-SEVERI form  $S_J$  (S.

DEF. 5.6 page 39) for  $U_0, U_1, \dots, U_n, X_1, X_2, \dots, X_{c-n-1}$  where

$U_j = (u_j, u_j, \overset{h}{\dots}, u_j) \in \Delta, j = 0, 1, \dots, n$  and the  $x_i$  are arbitrary vectors of  $E \otimes \mathbb{C}^h$ ;  
 $i = 1, 2, \dots, c-n-1$ , i.e.

$$(12.3) \quad S_J(U_0, U_2, \dots, U_n; X_1, X_2, \dots, X_{c-n-1}) \equiv 0$$

iff  $\bigcap_{j=1}^h V^{(j)} \neq \phi \Leftrightarrow J(V^{(1)} \times \dots \times V^{(h)}) \cap \Delta \neq \phi$ .

In particular in the "elimination case" again  $c_1 = c_2 = \dots = c_h = 1, h = c > n + 1$  the condition (12.2) is a covariant in the coefficients of the forms  $F_j$  of degree  $m_j$  containing  $c - n - 1$  arbitrary series of variables  $X_1, X_2, \dots, X_{c-n-1}$ . The coefficients of the power products in these  $X$ 's gives a system of resultant forms. We hope to study in the near future the relation between this invariant-theoretic approach and the classical ones.

### AN INTRINSIC ELIMINATION THEORY

13. HISTORICAL APPROACH. The elimination theory has been completely "eliminated" from algebraic Geometry! I believe that the main reason is that it was not intrinsic enough; as a matter of fact it was always presented in relation with a coordinate system. For instance the HENZELT-NOETHER sophistication of the KRONECKER elimination method was presented as follows: ([H - N]).

"Let  $m \subset K[x_1, \dots, x_n]$  be a polynomial ideal. We can associate to  $m$  a "resultant form"

$$(13.1) \quad R_m = R^{(1)}(x_1, \dots, x_n) \dots R^{(1)}(x_1, \dots, x_n) \dots R^{(n)}(x_n) \equiv 0 \ (m)$$



in such a way that  $R$  vanishes for all the solutions of  $m$  and only for them. If  $n \supset m$  and  $R_m = R_n$  then  $m = n$ ."

We can appreciate that the  $x_i$  are explicitly used in the statement and in a given order.

The geometrical meaning of the  $R^{(1)}$  is clear.  $R^{(1)}$  represents the irreducible components of  $V = V(m)$  of dimension equal to one precisely if

$$(13.2) \quad R^{(1)} = \prod F_{r_k}^{m_k}$$

the hypersurface  $F_{1k} = 0$  is an irreducible hypersurface contained in the solution variety  $V = V(m)$  and conversely any such hypersurface appears as a prime factor of  $R^{(1)}$ ,  $R^{(2)}(x_2, \dots, x_n) = 0$  represents the projection in the hyperplane  $x_1 = 0$  of the locus of irreducible components of codimension two, ..... and

$$R^{(1)}(x_i, x_{i+2}, \dots, x_n) = 0$$

appears as the projection in the coordinate space  $x_1 = 0, x_2 = 0, \dots, x_{i-1} = 0$  of the locus of irreducible components of  $V = V(m)$  of codimension equal to  $i : i = 1, 2, \dots, n$ . More precisely if we want to deal again with projective varieties in  $\mathbb{P}_n$  we need to introduce the homogeneous coordinates  $x_0, x_1, \dots, x_n$  and to assume that  $m$  is homogeneous. Besides *it is necessary to assume that the projective frame of reference is generically located with respect to  $V$* . If this is not the case *it is necessary to apply previously a generic projective transformation to achieve this goal*. We emphasize that:

*Such generic projective coordinate changes were never written in the notations ; as a consequence they did not appear in the formulas; accordingly the results are wrongly applied when the reference frame is badly located with respect to the variety defined by (12.0) again the results are misleading.*

The fact that the homogeneous  $R^{(1)} = R(x_{i-1}, \dots, x_n)$  appear as a projection from the coordinate space joining the vertices  $P_0, P_1, \dots, P_{i-2}$  (assumed previously as "well-located") suggests naturally the idea of projecting the locus  $\Gamma^{(1)}$  of irreducible components of  $V$  of codimension equal to  $i$  from a generic  $P_{i-2}$ . But this is CAYLEY'S idea. As a consequence the Author in the two papers  $[G_2], [G_3]$  replace the original problem of "elimination" by the following one:

*Let*

$$(13.3) \quad F_j = c \quad j = 1, 2, \dots, r$$

*be an arbitrary basis of the homogeneous ideal  $m$ . We shall compute the CAYLEY-SEVERI forms  $S^{(i)}$  of  $\Gamma^{(i)}$ ;  $i = 1, 2, \dots$ .*

$$(13.4) \quad S^{(1)}(x) = 0 \quad S^2(x_1, x_2) = 0, \dots, \quad S^{(i)}(x_1, \dots, x_i) = 0 \dots$$

*following the same steps as the traditional KRONECKER elimination method.*

The first step is obvious;  $S_1 = h \cdot c \cdot d(F_1, F_2, \dots, F_r)$  i.e. the hypersurface component appears in the same way as in the KRONECKER method. The elimination of one variable (which one?) depends on the choice of a well-located ( $\Leftrightarrow$  not belonging to  $\Gamma^{(2)}$ ) vertex of the projective frame. If we choose a generic projection center  $(y)$

we are reduced to the first step again because such a cone has codimension one. This can be achieved in an elementary way writing  $F_j = S^{(1)} G_j$ , then  $G_j = G_j(\lambda x + \mu y)$  and a resultant system in  $(\lambda, \mu)$ :

$$(13.5) \quad G_{2k}(x, y) = 0 \quad k = 1, 2, \dots, r_2.$$

Then

$$S^{(2)} = \text{hcd}(G_1, G_2, \dots, G_{r_2}).$$

In such a way – by induction we construct associate systems of equations.

$$F_{ck}(x_1, x_2, \dots, x_c) = 0 \quad k = 1, 2, \dots, r_k$$

where  $F_{1k} = F_k$   $r_1 = r$ . Then  $S_c = \text{h.c.d}(F_{ck} \dots)$ .

With this procedure we can attach to any system (13.3) the associate forms to the  $\Gamma^{(i)}$   $i = 1, 2, \dots, n$ . The prime factor decomposition of  $S_c(x_1, \dots, x_c)$  gives all the CAYLEY–SEVERI forms of the irreducible components of codimension  $c$  of  $V$  *with a certain intrinsic exponent depending only on  $m$ .*

We refer to [G2] [G3] for more details. There is a curious paradoxon in this procedure pointed out already in [G2]: instead of decreasing the number of coordinates by successive "elimination" of  $x_0, x_1, \dots, x_n$  we increase by  $n + 1$  homogeneous coordinates of  $(x_1, x_2, \dots, x_c \dots)$  in every step. However, let us recall that there exists an expression

$$S_c = S_c( \dots, p, \overset{i_1 i_2 \dots i_2}{\dots} \dots )$$

unique if we assume that all the power products of the  $p$ 's are standard. Let us specialize the projection points—coming back to the elimination theory, assuming them to be the vertices of the projective reference frame  $P_0, P_1 \dots$  (assuming again that they are well-located to avoid identical vanishing ...). Then we have the coordinate matrix

$$\begin{bmatrix} 1 & 0 & 0 & \dots & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \dots & 1 & \dots & 0 \\ \alpha_0 & \alpha_1 & \dots & \alpha_{c-1} & \alpha_c & \dots & \alpha_n \end{bmatrix}$$

and we remark that *the only not zero Grassmann coordinates are  $x_c, x_{c+1}, \dots, x_n$ . The  $x_0, x_1, \dots, x_{c-1}$  are "eliminated" again.*

Since the three types of associate forms can be transformed among them it is not difficult to compute the CHOW or the WEIL-SIEGEL forms. We are definitely interested in the latter because we shall prove in [G4] (cf. § 14 for a short Introduction) that *the computation of these forms is equivalent to use the KRONECKER elimination method with generic projective coordinate systems explicitly given in the formulas by means of basis  $u_0, u_1, \dots, u_n$  of the dual vector space  $\check{E}$ .*

The generic coordinates

$$\xi_j = \langle u_j, x \rangle = \sum_{k=0}^n u_{jk} x_k \quad j = 0, 1, \dots, n$$

of any vector  $x \in E$  (can be interpreted as the projective coordinates of the point  $(x)$ ) (whenever  $x \neq 0$ ). The elimination of the generic variables  $\xi_0, \xi_1, \dots, \xi_{i-1}$  leads naturally to forms of type.

$$N(\xi_1, \xi_{i+1}, \dots, \xi_n)$$

and we know that the  $\xi_j'$ , represent actually homogeneous coordinates in the more sophisticated projective space  $\mathbb{P}(E/E_{n-i})$  where  $E_{n-i}$  is defined by  $\langle u_j, x \rangle = 0$  for  $j = 0, 1, \dots, i-1$  cf. § 6. Actually the projection on the coordinate space opposite to  $E_{n-i}$  is not needed. The genericity insures that

$$\mathbb{P}(E_{n-i}) \cap V(\mathfrak{m}) - \Gamma^{(1)} - \Gamma^{(2)} - \dots - \Gamma^{(i-1)} = \phi.$$

#### 14. INTRINSIC ELIMINATION THEORY USING WEIL-SIEGEL FORMS.

Let us replace the CAYLEY-SEVERI forms by the corresponding WEIL-SIEGEL ones using formulas of type:

$$(14.1) \quad S_c(x, x_0, \dots, x_{c-2}) = N_c(u_0, u_1, \dots, u_{d+1}; x) = \mathbb{W}(\langle u_0, x \rangle, \dots, \langle u_{d+1}, x \rangle)$$

where  $x \in E - \{0\}$  is regarded as variable in the cone  $S(x, x_0, \dots, x_{c-2}) = 0$  of vertex  $\mathbb{P}_{c-2}$  spanned by  $(x_0), (x_1); \dots; (x_{c-2})$  ( $\Leftrightarrow$  intersection of the  $d+2$  hyperplanes  $(u_j)$ ,  $j = 0, 1, \dots, d+1$ ). The variables  $\xi_j = \langle u_j, x \rangle$  are again the homogeneous coordinates in the abstract  $(d+1)$ -dimensional space  $\mathbb{P}(E/E_{c-1})$  where  $\mathbb{P}(E_{c-1}) = \mathbb{P}_{c-2}$ .

If we take the  $n + 1$  forms  $u_0, u_1, \dots, u_n \in \check{E}$  dual to  $x_0, x_1, \dots, x_n \in E$   $x_j \neq 0$   
 $u_j \neq 0$  we have the following sequence of WEIL-SIEGEL forms:

$$\begin{aligned}
 S_1(x) &= N(x; u_0, u_1, \dots, u_n) = \mathbb{N}(\xi_0, \xi_1, \dots, \xi_n) \\
 S_2(x, x_0) &= N(x; u_1, u_2, \dots, u_n) = \mathbb{N}(\xi_1, \dots, \xi_n) \\
 (14.2) \quad S_3(x, x_0, x_1) &= N(x; u_2, \dots, u_n) = \mathbb{N}(\xi_2, \dots, \xi_n) \\
 &\dots\dots\dots \\
 S_c(x; x_0, x_1, \dots, x_{c-2}) &= N(x; u_{c-1}, u_c, \dots, u_n) = \mathbb{N}(\xi_{c-1}, \dots, \xi_n) \\
 &\dots\dots\dots
 \end{aligned}$$

We remark that formally, when we read the (14.2) from top to bottom we have:

$$(14.3) \quad N(u_0, u_1, \dots, u_n; x) = \mathbb{N}(\xi_0, \xi_1, \dots, \xi_n) = 0$$

as *the equation of the hypersurface* (or better, *of the divisor attached to  $\Gamma^{(i)}$* ) written in the projective coordinate system  $((u_0), (u_1), \dots, (u_n); (u_0 + u_1 + \dots + u_n))$  which can be regarded as "indeterminate": more precisely *if we write*

$$\langle u_j, x_j \rangle = \sum_{k=0}^n u_{jk} x_k$$

*we have the  $(n + 1) \times (n + 1)$  matrix  $(u_{jk})$  representing the projective coordinate "system" of Introduction but written in the formula instead of being ignored.*

If we write the system (13.3) in this "invariant way"

$$F_j = F_j(x; u_0, u_1, \dots, u_n) = F_j(\langle u_0, x \rangle \dots \langle u_n, x \rangle)$$

we can perform equally the first step of KRONECKER'S elimination method:

$$N_1(x; u_0, u_1, \dots, u_n) = M_1(\xi_0, \xi_1, \dots, \xi_n) .$$

The next step is to divide each  $F_j$  by  $M_1$ ,  $F_j = M_1 G_j$ . Then we can "eliminate  $\xi_0$ " (but within the generic projective frame  $(u_0), (u_1), \dots, (u_n); (u_0 + \dots + u_n)$ ). The new system

$$G_j(x; u_0, u_1, \dots, u_n) = G_j(\xi_0, \xi_1, \dots, \xi_n)$$

represents the variety  $V - \Gamma^{(1)}$  of codimension two which is not contained in  $\xi_1 = 0$ . Let us cut  $V - \Gamma^{(1)}$  with this hyperplane; we shall have only the useful "generic" variables  $\xi_1, \xi_2, \dots, \xi_n$ , i.e. we have a system of type:

$$G_j(x; u_1, u_2, \dots, u_n) = \tilde{G}_j(\xi_1, \xi_2, \dots, \xi_n)$$

representing the projecting cone of  $V - \Gamma^{(1)}$  from the intersection point of the  $n$  hyperplanes  $\langle u_j, x \rangle = 0$   $j = 1, 2, \dots, n$  (of vertex  $P_0$  in the corresponding generic projective frame). Then h.c.d. of the  $G_j(\xi_1 \dots, \xi_n)$  will give us back the WEIL-SIEGEL form attached to  $\Gamma^{(2)}$ , i.e. to the cycle of codimension 2 represented by  $m$ .

In other words we can prove the announced result:

*The systematic computation of the WEIL–SIEGEL forms  $\mathbb{N}_1(\xi_{c-1}, \xi_c, \dots, \xi_n)$  for  $c = 1, 2, \dots, n$  is equivalent with the old KRONECKER elimination method but with the projective projective coordinate system built in the formulas.*

#### REMARKS

1) We emphasize the use of the quotient projective spaces  $\mathbb{P}(E/E_{c-1})$  corresponding to coordinate spaces  $\mathbb{P}(E_{c-1})$  instead of the projection or the face opposite to  $\mathbb{P}(E_{c-1})$ .

2) In order to check all the necessary cautions we follow [vdW1] IV Kap. § 31, page 116; as well as the second Edition of vdW's Algebra.

The first steps are possible because we know, that the coefficient of the highest power of each  $x_i$  is  $\neq 0$  (because the corresponding projection space never met the projecting variety. The resultant systems of relative prime forms cannot be identically zero). The coefficient of  $x_2^g$  for a WEIL–SIEGEL form is equal to

$$\pm Y(u_0, \dots, u_i, \dots, u_{j+1}) \neq 0$$

for  $(i = 1, 2, \dots, \text{etc.})$ .



BIBLIOGRAPHY

- [B] BERTINI, *Geometria proiettiva degli Spazii*, Principato, Messina, 1923.
- [B-V] BODA-VOGEL, *On system of parameters, local intersection multiplicity and Bezout's theorem*, Proc. Am Math. Soc. 781 (1890) 1–7.
- [BOU] BOURBAKI, *Algèbre multilinéaire*, 1044 Act. Sci. Ind. Hermann, Paris, 1968.
- [C] CAYLEY, *On a new analytical representation of curves in space*, Collected papers, 4, 446, Quarterly Journal Pure, Appl. math. Vol. III, 1860, and Vol. V (1862), 81–86.
- [Ch-vdW] CHOW-VAN DER WAERDEN, ZAG-IX, *Ueber zugeordnete Formen und algebraische Systeme von algebraischen Mannigfaltigkeiten*, Math. Annalen 113, 5(1937) 692–704 (ZAG book, page 212).
- [dC-E-P] DE CONCINI-EISENBUD-PROCESI, *Hodge Algebras*, Astérisque 91, 1982.
- [F-V] FLENNER-VOGEL, *Connectivity and its applications to improper intersections in  $\mathbb{P}_n$* , Mathemata Göttingensis, Heft 53, Oktober 1988.
- [F] FULTON, *Intersection theory*, Ergebnisse der Math., 3 Folge, Bd. 2.
- [F-L] FULTON-LAZARSELD, *Connectivity and its application in algebraic geometry*, LNM, Springer 862 (1981), 26–92.
- [G1] GAETA, *Sul calcolo effettivo della forma associata  $F(W_{\alpha+\beta}^{gh})$  all'intersezione di due cicli effettivi puri  $U_{\alpha}^g, V_{\beta}^1$  di  $S_n$  in funzione*

delle  $F(U_{\alpha}^g)$ ,  $F(V_{\beta}^h)$  relativi ai cicli secanti, Atti Acc. Naz. Lincei, 8, 24 (1958), 269–276.

- [G2] GAETA, *On a new tensorial algorithm replacing the elimination theory*, Tensor, 1963, Vol. 13, No. 1.
- [G3] " *Sopra un aspetto proiettivamente invariante del metodo d'eliminazione di Kronecker*, Rend. Acc. Naz. Lincei (8), 18, 1955, pág. 148–150.
- [G4] " *Intrinsic elimination theory with WEIL–SIEGEL forms and applications to the SCHOTTKY problem* (to appear).
- [Gr] GREEN–MORRISON, *The equations defining Chow varieties*, Duke math. Journal, September, 1986.
- [Grö1] GRÖBNER, *Moderne algebraische Geometrie*, Springer Wien–Innsbruck, 1949.
- [Grö2] " *Algebraische Geometrie*, I, II, Mannheim; B1 1968, 1970.
- [H–M] HENTZELT–NOETHER, *Zur Theorie der Polynomideale und Resultanten*, Math. Ann. 88 (1923), 553–79 or E. NOETHER Ges. Abh. page 409.
- [H–P] HODGE–PEDOE, *Methods of algebraic Geometry*, vol. 1–2 Cambridge University Press, 1952.
- [Hu] HURWITZ, *Ueber die Fraegheitsformen eines algebraischen Modells*, Annali di Matematica pura ed. applicata, serie III, t. 20, 1913 and, *Gesammelte Abhandlungen*, Band II, LXXXVI, 586.
- [K] KAPFERER, *Ueber Resultanten und Resultanten-Systeme*, Sitzungsberichte der mathematisch–naturwissenschaftlichen Abteilung der Bayrischen Akademie der Wissenschaften, Jahrgang 1929, 179–200.

- [KL] KLEIMAN, *Letter to Vogel*.
- [M] MARTENS, *Ueber die bestimmenden Eigenschaften der Resultante von  $n$  Formen mit  $n$  Voraenderlichen*, Sitzungsberichte k. Ak. der Wiss., XCIII Band, 525–566.  
*Zur Theorie der Elimination*, Ibidem, 1173–1228 and 1345–1386.
- [N] E. NOETHER, *Gesammelte Abhandlungen*, Springer, 1983 in particular papers on Elimination theory 18, 22, 24, 25:  
18: *Ueber eine Arbeit des im Kriege gefallenen H. Hentzelt zur Eliminationstheorie*, J. Ber. d. BMV 30 (1921), S. 101.  
22: Bearbeitung von K. Hentzelt: *Zur Theorie der Polynomideale und Resultanten*, Math. Ann. 88 (1923), S. 53–79.  
24: *Eliminationstheorie und allgemeine Idealttheorie*, Math. Ann. 90 (1923), S. 229–261  
25: *Eliminationstheorie und Idealttheorie*, J. Ber. d. DMV 33, S. 116–120.
- [P] PEDOE, *On a new analytical representation of curves in space*, Proc. Camb. Phil. Society, 43 (1947), 455–458.
- [SG] SEGRE, C. *Sulla varietà che rappresenta le coppie dei punti . . . . .*, Rend. Circ. Mat. Palermo 5 (1891), 192–204.
- [S] SEVERI, *Memorie scelte*, Zuffi, Bologna 195, in particular: 22 *Sulla varietà che rappresenta gli spazi subordinati . . . . .*, page 405; (or Ann. de Mat. s. III, vol. 24, pag. 89–120, 1915) and comments by SEVERI in page 438.
- [SI] SIEGEL, *Moduln abelscher Funktionen*, Nachrichten der Ak. der Wiss. in Göttingen, Math. – phys. Klasse 1960, Nr. 25, 365–427 and *Gesammelte Abhandlungen*, Vol. III, 77, pag. 373.
- [S–V] STÜCKRAD–VOGEL, *Buchbaum-rings and applications* VEB Deutscher Verlag der Wissenschaften, Berlin, 1986.

- [vdW] VAN DER WAERDEN, *Moderne Algebra*, 2<sup>te</sup> Auflage, Grund. Math. Wiss. Springer Verlag.
- [vdW1] " *Einführung in die algebraische Geometrie* , Grund. Math. Wiss. Berlin, Band 51, 2<sup>nd</sup> Ed, Springer Verlag 1973 and the Appendix II, pag. 271,
- [vdW2] *The Foundation of algebraic Geometry from Severi to Andre Weil* , (cf. also Archives for History of exact Sciences) Vol. 7, Number 3, 1971, 171-180.
- [vdW3] Math. Ann. 115 (1938), 619-642.
- [vdW-ZAG] *Zur algebraischen Geometrie* (Selected papers), Springer, 1983.
- [V1] VOGEL, *Lectures on results on Bézout's Theorem* , Lecture Notes of the Tata Institute of Fundamental Research. Bombay, n. 74, Springer Verlag, 1984.
- [V2] Proceedings of this Conference.
- [W] WEIL, *Foundations of algebraic Geometry*, 2<sup>nd</sup> Ed. A. Math. Soc. Coll. Publications, 29, Providence RI, USA, 1962.
- [WE] WEITZENBÖCK, *Invariantentheorie*, Noordhoff, Gröningen, 1923.