

**ON THE SHARPNESS OF NOVIKOV TYPE  
INEQUALITIES FOR MANIFOLDS WITH FREE  
ABELIAN FUNDAMENTAL GROUPS**

A.V. Pazhitnov

Max-Planck-Institut  
für Mathematik  
Gottfried-Claren-Straße 26  
D-5300 Bonn 3

Federal Republic of Germany

Institute of chemical physics  
AN USSR, 117977  
Moscow, Kosygin str., 4

USSR

MPI/90-61



ON THE SHARPNESS OF NOVIKOV TYPE INEQUALITIES  
FOR MANIFOLDS WITH FREE ABELIAN FUNDAMENTAL GROUPS

UDC 515.164.174

Pazhitnov A.V.

In papers [1], [2] S.P. Novikov has constructed an analogue of Morse theory for "multivalued Morse functions", i.e. for closed but, generally, non-exact Morse 1-forms on smooth manifolds. Recall that a closed 1-form  $\omega$  on a smooth manifold  $M^n$  is called a Morse form, if all the zeros of  $\omega$  are nondegenerate, or, equivalently, if locally  $\omega = dh$ , where  $h$  is a Morse function. In this case the index  $\text{ind}_c \omega$  of each zero  $c$  of the form  $\omega$  is defined. Denote  $m_p(\omega)$  the number of zeros of the form  $\omega$  of index  $p$ .

One of the basic problems of this theory in the finite-dimensional case is the following: for a given cohomology class  $\xi \in H^1(M, \mathbb{R})$  find the numbers  $c_p(M, \xi)$ , where  $0 \leq p \leq n$ , providing the lower estimates (sharp if possible) for the Morse numbers  $m_p(\omega)$  of any form  $\omega$ , belonging to the class  $\xi$ :

$$m_p(\omega) \geq c_p(M, [\omega]), \quad 0 \leq p \leq n \quad (0.1)$$

Recall that the estimates (0.1) are said to be sharp for a manifold  $M$  and a class  $\xi$  if there exists a Morse 1-form  $\omega$ , belonging to  $\xi$ , such that inequalities (0.1) appear to be equalities for all  $p$ .

For  $\xi = 0$  the estimates (0.1) are provided by Morse inequalities; in this case  $c_p(M^n, 0) = b_p(M) + q_p(M) + q_{p-1}(M)$ , where  $b_p(M)$  stands for the rank of  $H_p(M)$  and  $q_p(M)$  - for

the torsion number of  $H_p(M)$  (i.e. the minimal number of generators of torsion subgroup  $\text{Tors } H_p(M)$ ).

For  $\pi_1 M^n = 0$ ,  $n \geq 6$  these estimates are sharp (Smale's theorem [3]).

For any cohomology class  $\xi \in H^1(M, \mathbb{R})$  (and also for any form  $\omega : [\omega] = \xi$ ) we define the irrationality degree of  $\xi$  to be the maximal number of  $\mathbb{Q}$ -linearly independent periods of (or, equivalently, the rank of  $\text{Im}(\xi : \pi_1 M \rightarrow \mathbb{R})$ ). The forms  $\omega$  of irrationality degree 1 will be called rational.

For a class  $\xi \neq 0$  having irrationality degree 1 (these are exactly the multiples of the integer classes) the estimates (0.1) were suggested by S.P. Novikov [1], [2] (cf. later). The sharpness of these estimates for the case  $\pi_1 M^n = \mathbb{Z}$ ,  $n \geq 6$  has been proved by M.Sh. Farber [4].

For  $\pi_1 M = \mathbb{Z}$  any Morse 1-form is up to the positive constant the differential of the Morse map  $M^n \rightarrow S^1$ ; thus the sharpness problem is equivalent to the problem of constructing a Morse map  $f : M \rightarrow S^1$  with a minimal number of critical points of all indices. The necessary and sufficient condition of existence of a map  $f : M \rightarrow S^1$  without critical points (i.e. fibration) was supplied by W. Browder and J. Levine [5] (it's easy to check the equivalence of this condition to the condition arising from Novikov inequalities).

The main purpose of the present paper is to obtain the estimates of the type (0.1) for the forms of arbitrary irrationality degree and to prove the sharpness of the estimates obtained for generic cohomology classes  $[\omega]$  and manifolds  $M^n$  satisfying  $\pi_1 M^n = \mathbb{Z}^m$ ,  $n \geq 6$  (and some additional homotopical

restrictions; for precise formulation see theorem 0.1 below). These results were partly announced in [6].

We begin with definitions (cf. [1],[2]).

Let  $G$  be a group,  $\xi: G \rightarrow \mathbb{R}$  - a homomorphism. Denote by  $\Lambda$  the group ring  $\mathbb{Z}[G]$ . Consider the set of all linear combinations (infinite in general)  $\lambda = \sum_{g \in G} n_g g$ , such that the intersection of  $\text{supp } \lambda$  (where  $\text{supp } \lambda = \{g | n_g \neq 0\} \subset G$ ) with any set  $\{g | \xi(g) \geq c\}$  is finite. It is easy to see that the resulting abelian group is a ring. We denote it by  $\Lambda_{\xi}^{-}$ . (Note, that  $\Lambda_{\xi}^{-}$  is the completion of  $\Lambda$  with respect to the system of subrings, but not ideals.)

For  $G = \mathbb{Z}^{\ell}$  the group ring  $\Lambda = \mathbb{Z}[G] = \mathbb{Z}[t_1^{\pm 1}, \dots, t_{\ell}^{\pm 1}]$  is the ring of Laurent polynomials with integer coefficients; the completion  $\Lambda_{\xi}^{-}$  is the ring of all the power series  $\lambda = \sum_I \lambda_I t^I$  (where  $I = (i_1, \dots, i_{\ell}) \in \mathbb{Z}^{\ell}$ ,  $\lambda_I \in \mathbb{Z}$ ), such that for any  $c$  the set  $\text{supp } \lambda$  contains only finite number of indices  $I$ , satisfying  $\xi(I) \geq c$ . The ring  $\mathbb{Z}[\mathbb{Z}]$  will be further denoted by  $L$ . For  $\ell = 1$ ,  $\xi \neq 0$  the ring  $\Lambda_{\xi}^{-}$  (denoted further by  $L$ ) is the ring of all integer Laurent power series with the finite negative part (we suppose that  $\xi(1) < 0$ ).

Next we recall some results from [1],[2].

Let  $\omega$  be a Morse 1-form on a manifold  $M$  ( $M$  having an arbitrary fundamental group); denote  $\xi \in H^1(M, \mathbb{R})$  its cohomology class. Consider the minimal covering  $p: \tilde{M}_{\xi}^n \rightarrow M$ , for which the pullback of  $\omega$  is exact:  $p^* \omega = df$ . This covering corresponds to the subgroup  $\text{Ker}(\xi: \pi_1 M \rightarrow \mathbb{R}) \subset \pi_1 M$  and is regular with the structure group  $\mathbb{Z}^{\ell}$ , where  $\ell$  is irrationality degree of  $\omega$ . The critical points of  $f: \tilde{M}_{\xi}^n \rightarrow \mathbb{R}$  and the paths of

steepest descent of  $f$  give rise to Novikov complex  $C_*(\bar{M}_\xi^n, \omega)$ . The latter is an analogue of Morse complex of a single-valued function  $f$  on a compact manifold  $M$ . Novikov complex is a free finitely generated complex over the ring  $\Lambda_\xi^-$ , where  $\Lambda = \mathbb{Z}[\mathbb{Z}^{\mathbb{Z}}]$  (note that homomorphism  $\xi$  factors through  $\mathbb{Z}^{\mathbb{Z}}$  by definition). The number of free  $\Lambda_\xi^-$ -generators of  $C_p(\bar{M}_\xi^n, \omega)$  equals  $m_p(\omega)$ . Homology  $H_*(C_*(\bar{M}_\xi^n, \omega))$  is isomorphic to homology  $H_*(C_*(\bar{M}_\xi^n) \otimes_{\Lambda} \Lambda_\xi^-)$ .

Consider for instance a cohomology class  $\xi$  of irrationality degree 1. The covering  $M_\xi^n \rightarrow M^n$  is infinite cyclic. The ring  $\Lambda_\xi^-$  is the ring  $\hat{\Lambda} = \mathbb{Z}[[t]][t^{-1}]$ , which is known to be a principal ideal domain. For any finitely generated module  $M$  over a principal ideal domain  $R$  the rank  $b(M)$  and the torsion number  $q(M)$  are defined; for any free finitely generated  $R$ -complex  $C_*$  the number of free generators  $\mu(C_p)$  is not greater than

$$b(H_p(C_*)) + q(H_p(C_*)) + q(H_{p-1}(C_*))$$

Hence, for rational forms

$$m_p(\omega) \geq b(H_p(\bar{M}_\xi^n) \otimes_{\hat{\Lambda}} \hat{\Lambda}) + q(H_p(\bar{M}_\xi^n) \otimes_{\hat{\Lambda}} \hat{\Lambda}) + q(H_{p-1}(\bar{M}_\xi^n) \otimes_{\hat{\Lambda}} \hat{\Lambda}) \quad (0.2)$$

(see [1], [2]; see also [4] for another proof).

J.-C. Sikorav proved (see for a proof §1 of the present paper) that for any  $k$  the completion  $\Lambda_\xi^-$  of the ring  $\Lambda = \mathbb{Z}[\mathbb{Z}^k]$  with respect to a homomorphism  $\xi : \mathbb{Z}^k \rightarrow \mathbb{R}$  of a maximal irrationality degree  $k$  is a principal ideal domain.

Therefore the same argument as above enables us to obtain the

analogues of (0.2) for the forms of arbitrary irrationality degree.

We will need still another variant of these inequalities, dealing with forms of arbitrary irrationality degree, but arising from maximal free abelian covering. To produce it we need one more algebraic lemma.

Namely, let  $m = \text{rk} H_1(M)$  and consider a class  $\xi \in H^1(M, \mathbb{R})$  of the maximal irrationality degree  $m$ . Denote by  $b_p(M, \xi)$  the rank and by  $q_p(M, \xi)$  the torsion number of the module  $H_p(C_*(\bar{M}) \otimes_{\Lambda} \Lambda_{\xi}^{-1})$ , where  $\bar{M} \rightarrow M$  stands for a  $\mathbb{Z}^m$ -covering, corresponding to the homomorphism  $\pi_1 M \rightarrow H_1 M / \text{Tors } H_1 M, \Lambda = \mathbb{Z}[\mathbb{Z}^m]$ . For  $[\omega] = \xi$  we have

$$m_p(\omega) \geq b_p(M, \xi) + q_p(M, \xi) + q_{p-1}(M, \xi) \quad (0.3)$$

One easily proves that  $b_p(M, \xi)$  does not depend on  $\xi$ . J.-C. Sikorav has also proven that  $q_p(M, \xi)$  does not depend on  $\xi$  in any connected component of the complement in  $H^1(M, \mathbb{R}) = \mathbb{R}^m$  to the finite union  $\bigcup \Gamma_i$  of hyperplanes  $\Gamma_i$ , each of which is determined by a linear equation with integer coefficients (see §1 of the present paper). Note, by the way, that the definition implies

$$q_p(M, [\omega]) = q_p(M, c[\omega]), \quad c > 0.$$

Now let  $\xi$  be any element of  $H^1(M, \mathbb{R}) \setminus \bigcup \Gamma_i$ . We set by definition  $q_p(M, \xi) = q_p(M, \xi')$ , where  $\xi'$  is an arbitrary maximally irrational class, sufficiently close to  $\xi$ . Suppose that  $\omega$  is a Morse 1-form, such that  $[\omega] \in H^1(M, \mathbb{R}) \setminus \bigcup \Gamma_i$ . Approximating  $\omega$  with the maximally irrational forms we obtain the

inequalities (0.3) also for the ~~( )~~ arbitrary cohomology class  $[\omega] \in H^1(M, \mathbb{R}) \setminus \bigcup_i \Gamma_i$ .

Our notations differ here from that of [1], [2]. In these papers  $b_p(M, \ )$ ,  $q_p(M, \ )$  stand for numbers which we denoted by  $b(H_p(\bar{M}_\xi) \otimes_L \hat{L})$ ,  $q(H_p(\bar{M}_\xi) \otimes_L \hat{L})$ . Still these notations correspond rather well to one another. Namely, lemma 2.6 from §2 asserts that for rational cohomology classes  $\xi$ , belonging to some dense conical open set  $U^1$ , we have

$$b_p(M, \xi) = b(H_p(\bar{M}_\xi) \otimes_L \hat{L}), \quad q_p(M, \xi) = q(H_p(\bar{M}_\xi) \otimes_L \hat{L}).$$

Now we can state the main theorem of the present paper.

Theorem 0.1. Let  $M^n$  be a smooth compact connected manifold without boundary,  $\pi_1 M^n = \mathbb{Z}^m$ ,  $n \geq 6$ . Suppose that for some  $r$ :  $2 \leq r \leq n-4$  the homology  $H_p(\tilde{M}^n)$  of the universal cover vanishes for  $r-1 \leq p \leq r+2$ .

Then there exists an open dense conical subset  $U \subset H^1(M, \mathbb{R}) = \mathbb{R}^m$ , such that any  $\gamma \in U$  can be realized by a Morse 1-form  $\omega : [\omega] = \gamma$ , which has the minimal possible number of zeros of any index  $p$  in the class  $\gamma$ , this number being equal to the righthand side of (0.3).

Applying lemma 2.6 from §2 we immediately deduce from here the result concerning the sharpness of classical Novikov inequalities (0.1).

Corollary 0.2. Under the assumptions of theorem 0.1 there

---

1) A subset  $U \subset \mathbb{R}^m$  is called conical if  $x \in U \Rightarrow tx \in U$  for every  $t > 0$ .



exists the conical dense open set  $V \subset H^1(M, \mathbb{R}) = \mathbb{R}^m$ , such that for any rational cohomology class  $\gamma \in V$  the Novikov inequalities (0.1) are sharp (i.e. any rational  $\gamma \in V$  can be realized by a Morse form  $\omega$  with  $m_p(\omega)$  equal to the righthand side of (0.1)).

Corollary 0.3. For a smooth manifold  $M^n$  with  $\pi_1 M^n = \mathbb{Z}^m$ ,  $n \geq 6$  and  $\tilde{M}^n$  four-connected, there exists a conical dense open set  $U \subset H^1(M, \mathbb{R})$ , such that any integer class  $\xi \in U$  can be realized by a Morse map  $f : M \rightarrow S^1$ , which has the minimal possible number of zeros of any index  $p$  in the class  $\xi$ .

Denote by  $m_p([\omega])$  the righthand side of (0.3). In the proof of theorem 0.1 we use actually not the vanishing of the universal cover homology, but the weaker condition  $m_r(\gamma) = m_{r+1}(\gamma) = m_{r+2}(\gamma) = 0$ . Hence we get

Corollary 0.4. For a smooth manifold  $M^n$  with  $\pi_1 M^n = \mathbb{Z}^m$ ,  $n \geq 6$ , the set of cohomology classes  $\gamma \in H^1(M, \mathbb{R})$ , realizable (up to multiplicative constant) by a fibration  $M \rightarrow S^1$ , is contained in the open cone  $V \subset H^1(M, \mathbb{R})$ , which is determined by the condition  $m_*(\gamma) = 0$ . This cone contains an open dense subset  $V_0$ , any rational class of which is realized by a fibration.

This corollary gives a partial generalization (for dimensions  $\geq 6$ ) of Thurston's result [7], concerning the fibrations of 3-manifolds over a circle.

It is natural to ask if one can weaken the assumptions of theorem 0.1, keeping the conclusion. The Morse 1-form  $\omega$  is called minimal if it has the minimal possible number of zeros of all indices in its cohomology class  $[\omega]$ . If  $\pi_1 M^n = \mathbb{Z}^m$ ,  $n \geq 6$

and  $\gamma \in H^1(M, \mathbb{R})$  is an arbitrary cohomology class the problem of existence of a minimal form in  $\gamma$  seems to be rather difficult. Indeed, if  $\gamma = 0$  this problem is just the problem of existence of minimal Morse functions on the manifolds with free abelian fundamental groups. The latter problem is not yet completely solved; a detailed treatment can be found in [8],[9]. Taking into account the theorem 0.1 we can describe the situation as following. We treat Morse 1-forms as multivalued Morse functions; the monodromy of each function of this type is given by  $m$  real numbers. The condition of zero monodromy (corresponding to usual Morse functions) proves to be too rigid for the known methods to deform the function into minimal one. If the class  $\gamma$  is of a general position (actually it is sufficient that  $\gamma$  is close enough to some class  $\gamma_0$ , having the  $\mathbb{Q}$ -linearly independent periods) the Morse form belonging to  $\gamma$  can be deformed into minimal one.

It's now natural to state the conjecture: the set of cohomology classes  $\gamma \in H^1(M, \mathbb{R}) = \mathbb{R}^m$ , realizable by minimal Morse forms, contains the complement to finite union of hyperplanes, determined by linear equations with integer coefficients. This hyperplanes correspond to "uncomfortable" monodromy conditions. The set of these hyperplanes must contain the hyperplanes  $\Gamma_1$  mentioned above and maybe smth. else.

The restriction

$$m_{r_c}(\gamma) = m_{r_c+1}(\gamma) = m_{r_c+2}(\gamma) = 0$$

is imposed for technical reasons. Still the author does not know if it is removable.

Now we'll sketch the main idea of the proof of theorem 0.1 and the contents of the paper.

§§ 1, 2 include the proof (not published before) of J.-C. Sikorav's theorems on the euclidean property of Novikov ring and on the numbers  $q_*(M, \xi)$ . We also prove here that in many cases one can replace the Novikov ring  $\Lambda_{\xi}^-$  by a suitable localization  $\Lambda_{(\xi)}$  of the ring  $\Lambda$ . (This is essential for the proof of theorem 0.1.) We also prove here that for a generic rational class  $\xi$  the number  $b_p(M, \xi)$  coincide with the rank of the module  $H_p(\bar{M}_{\xi}) \otimes_{\hat{L}} \hat{L}$ , and the number  $q_p(M, \xi)$  - with its torsion number.

In § 3 we prove the Poincare duality formula for Novikov homology.

§§ 4, 5 contain some auxiliary material. We recall here the results on Morse functions on the cobordisms, due to V.V. Sharko (§ 4) and prove some algebraic lemmas (§ 5).

In § 6 we produce two another proofs of inequalities (0.3). They use only Morse theory for the functions on the compact manifolds with boundary. These proofs are formally independent of the properties of Novikov complex, which was discussed above to clarify the roots of the present work. Thus the present work is self-contained. The first proof makes use of the algebraic lemmas of § 2 and reduces the problem to the case of cyclic covering; then we refer to [4]. The second proof is independent of [4]. In § 6 we also state some algebraic conjecture; if it holds, the second proof provides the general Morse type estimates for rational forms, similar to [8].

§§ 7, 8, 9 are devoted directly to the proof of theorem 0.1.

It is sufficient to prove this theorem for the rational

cohomology classes of general position. We'll prove it for any rational class  $\gamma$ , which is sufficiently close to the maximally irrational class  $\gamma'$ . The point is that in this case the modules  $H_*(\tilde{M}^n, \Lambda_{(\gamma)})$  have the resolutions of length two (although the ring  $\Lambda_{(\gamma)}$  is not the principal ideal domain). This enables us to apply here the scheme of proof of sharpness due to Browder-Levine-Farrell-Farber (§§ 7, 8). Instead of Smale's theorem on the minimal functions on simply-connected manifolds we use the corresponding Sharko's result [8].

The Browder-Levine-Farrell-Farber scheme does not work through directly. We use here the non-simply-connected surgery and while constructing the Morse form sought we come across the obstruction of Farrell type [10]. (To define this obstruction correctly we need the vanishing of  $H_*(M^n)$  in four successive dimensions.) The obstruction lies in a zero group  $C(\mathbb{Z}[\mathbb{Z}^{m-1}])$  (see [10]) but we need to realize this vanishing geometrically which requires some extra arguments (§ 9).

The author is grateful to S.P. Novikov for attention to the work and valuable discussions. The author is also grateful to J.-C. Sikorav and V.V. Sharko for the very timely information of their results and also to P.M. Akhmetiev, O.Ya. Viro, V.L. Kobel'skii, A.S. Mischenko, Yu.P. Soloviev, V.G. Turaev and V.V. Sharko for valuable discussions.

§ 1. The Novikov ring is euclidean

Denote  $\mathbb{Z}[\mathbb{Z}^{\ell}] = \mathbb{Z}[t_1^{\pm 1}, \dots, t_{\ell}^{\pm 1}]$  by  $\Lambda$  and let  $\xi: \mathbb{Z}^{\ell} \rightarrow \mathbb{R}$  be a homomorphism. Recall from the introduction that Novikov ring  $\Lambda_{\xi}^{-}$  consists of power series  $\lambda = \sum a_I t^I$  (where  $I = (i_1, \dots,$

$i_0) \in \mathbb{Z}^l$ , such that for every  $c$  the domain  $\xi \gg c$  contains only finite number of indices  $I$ , belonging to  $\text{supp } \lambda$  (recall that  $\text{supp } \lambda = \{I \mid a_I \neq 0\}$ ).

Theorem 1.1. (J.-C. Sikorav). If  $\xi : \mathbb{Z}^l \rightarrow \mathbb{R}$  is a monomorphism, then  $\Lambda_{\xi}^{-}$  is euclidean.

Proof. First of all note that injectivity of  $\xi$  implies that for any  $c \in \mathbb{R}$  the set  $\{\xi(x) = c\}$  contains at most one element. Therefore any element  $\lambda$  of the ring  $\Lambda_{\xi}^{-}$  contains a principal term  $a_{I_0} t^{I_0}$ , which is uniquely determined. For  $\lambda \in \Lambda_{\xi}^{-}$  we define the height  $h(\lambda) \in \mathbb{R}$  and the norm  $\|\lambda\| \in \mathbb{Z}$ , setting  $h(\lambda) = \xi(I_0)$ ,  $\|\lambda\| = |a_{I_0}|$ . We'll prove that  $\Lambda_{\xi}^{-}$  is euclidean with respect to this norm.

Suppose that  $A, B \in \Lambda_{\xi}^{-}$ . We are to divide  $A$  by  $B$  with a remainder. Without losing generality we may assume that the principal terms of these power series are  $a_0 \cdot \mathbb{1}$ ,  $b_0 \cdot \mathbb{1}$  ( $\mathbb{1}$  stands for the unit of the group  $\mathbb{Z}$ ).

Divide  $a_0$  by  $b_0$  with a remainder:  $a_0 = m_0 b_0 + q_0$ .

If  $q_0 \neq 0$ , then  $A = m_0 B + Q$  where  $\|Q\| = |a_0 - m_0 b_0| = |q_0| < |b_0|$  and the division is over. If not, apply the same procedure to the power series  $Q$ . Going on in the same fashion, we construct the sequence of polynomials  $M_i$  and the sequence of power series  $Q_i$ , such that  $A = M_i B + Q_i$ ,  $M_{i+1} = M_i + \mu_{i+1}$ , where  $\mu_{i+1}$  is a monomial, located lower (with respect to  $\xi$ ), than any monomial of  $M_i$ ;  $h(\mu_{i+1}) = h(Q_i)$ . If at some step we obtain  $\|Q_i\| < |b_0| = \|B\|$ , then our sequence stops and the division is over. If this never happens, consider the power series

$$M = \sum_{i=0}^{\infty} \mu_i. \quad \text{I claim that } M \in \Lambda_{\xi}^{-}.$$

Indeed, let  $\varepsilon = |h(B - b_0 \cdot 1)|$ . Suppose that we've already proved that only finite number of monomials  $\mu_i$  is located above the level ( $\xi = -N\varepsilon$ ), and let  $\mu_{n+1}$  be the first monomial lying below this level. Denote  $K$  the number of monomials of the power series  $Q_n$ , lying in the stratum  $-(N+1)\varepsilon < \xi \leq -N\varepsilon$ . It is clear that  $\mu_{n+1+K}$  lies below the level  $\xi = -(N+1)\varepsilon$ . Thus  $M \in \Lambda_{\xi}^-$  and it's clear that  $A = MB$ .

Remark 1.2. Consider  $\mathbb{Z}^{\ell}$  as a lattice in  $\mathbb{R}^{\ell}$  and extend  $\xi$  to a linear functional on  $\mathbb{R}^{\ell}$ . Consider the cone  $C \subset \mathbb{R}^{\ell}$ , formed by intersection of a finite number of half-spaces and lying in the domain ( $\xi \leq 0$ ). One deduces easily from the proof of theorem 1.1 that if all the monomials of  $a, b$  are contained in  $C$ , then the monomials of  $Q, M$  also are contained in  $C$ .

Definition 1.3. Let  $\gamma : \mathbb{Z}^{\ell} \rightarrow \mathbb{R}$  be any homomorphism (not necessary injective). Define the multiplicative subset  $S_{\gamma} \subset \Lambda$  as following:  $S_{\gamma} = \{1 + P\}$ , where all the monomials of  $P$  belong to the domain  $\gamma < 0$ . (If  $\gamma$  is injective,  $S$  is just the set of polynomials with the principal term equal to 1.) Set  $\Lambda_{(\gamma)} = S_{\gamma}^{-1}\Lambda$ .

Theorem 1.4. Let  $\xi : \mathbb{Z}^{\ell} \rightarrow \mathbb{R}$  be a monomorphism. Then  $\Lambda_{(\xi)}$  is euclidean.

Proof. We introduce some notations. Suppose that  $e = \{e_1, \dots, e_{\ell}\}$  is a collection of independent integer vectors in  $\mathbb{R}^{\ell}$ , such that  $\xi(e_i) < 0$ . Denote by  $M(e)$  the set of all linear combinations of  $e_i$  with integer nonnegative coefficients, by  $C(e)$  - the cone in  $\mathbb{R}^{\ell}$ , generated by the vectors  $e_i$ ; by  $M(e)$  - intersection of  $\mathbb{Z}^{\ell}$  and  $C(e)$ ; by  $\mathbb{Z}[M(e)]$ ,  $\mathbb{Z}[\tilde{M}(e)]$  - the subrings of  $\Lambda$ , generated by monomials whose exponents belong

to  $M(e)$  (correspondingly to  $\tilde{M}(e)$ ). The ring  $\mathbb{Z}[M(e)]$  is isomorphic to the polynomial ring  $\mathbb{Z}[u_1, \dots, u_\ell]$ . The ring  $\mathbb{Z}[\tilde{M}(e)]$  is finitely generated as a module over its sub ring  $\mathbb{Z}[M(e)]$ , therefore it is noetherian.

Consider the ideal  $I \subset \mathbb{Z}[\tilde{M}(e)]$ , consisting of all Laurent polynomials having the zero coefficient at  $\mathbf{1}$ . The ring of all the power series in monomials, belonging to  $\tilde{M}(e)$ , coincides with the completion of  $\mathbb{Z}[\tilde{M}(e)]$  with respect to  $I$  (and is contained in  $\Lambda_{\xi}^-$ ). Consider the localization

$$S(e)^{-1} \mathbb{Z}[\tilde{M}(e)], \text{ where } S(e) = \mathbf{1} + I \subset \mathbb{Z}[\tilde{M}(e)].$$

The ring  $\mathbb{Z}[\tilde{M}(e)]^{\wedge}$  is a faithfully flat module over the ring  $S(e)^{-1} \mathbb{Z}[\tilde{M}(e)]$  (since the latter is Zarisky ring; see [11, ch.III, §3]). This implies that the equation  $ax = b$ , where  $a, b \in S(e)^{-1} \mathbb{Z}[\tilde{M}(e)]$ , has a solution in  $\mathbb{Z}[\tilde{M}(e)]^{\wedge}$  if and only if it has a solution in  $S(e)^{-1} \mathbb{Z}[\tilde{M}(e)]$ .

Now we can prove the theorem 1.4. The ring  $\Lambda_{(\xi)}$  is contained in  $\Lambda_{\xi}^-$  and inherits from there the norm  $\| \cdot \|$ .

We'll show that  $\Lambda_{(\xi)}$  is euclidean with respect to this norm. It suffices to divide  $a$  by  $b$ , where  $a, b \in \Lambda$ ,  $h(a) = h(b) = 0$ . Apply now the division procedure described above. If it finishes after finite number of steps, we have  $a = mb + q$ , where  $m, q \in \Lambda$  and the division is over. If not, we have  $a = bx$ , where  $x \in \Lambda_{\xi}^-$ . Choose any collection  $e = \{e_1, \dots, e_\ell\}$  of vectors for which  $\text{supp } a, \text{supp } b \subset C(e)$ . Then  $a, b \in \mathbb{Z}[\tilde{M}(e)]$  and remark 1.2 implies that  $x \in \mathbb{Z}[\tilde{M}(e)]^{\wedge}$ , hence

$$x \in S(e)^{-1} \mathbb{Z}[\tilde{M}(e)] \subset \Lambda_{(\xi)}.$$

Q.E.D.

Remark 1.5. The ring  $\Lambda_{\xi}^{-}$  is faithfully flat module over its subring  $\Lambda_{(\xi)}$ , since they both are principal ideal domains and  $a \in \Lambda_{(\xi)}$  is invertible in  $\Lambda_{(\xi)}$  if and only if it is invertible in  $\Lambda_{\xi}^{-}$  (see [11, § 1]).

Thus  $\Lambda_{\xi}^{-}$  and  $\Lambda_{(\xi)}$  have the same homological properties. The ring  $\Lambda_{(\xi)}$  has some technical advantages over the ring  $\Lambda_{\xi}^{-}$ ; the replacement of  $\Lambda_{\xi}^{-}$  by  $\Lambda_{(\xi)}$  is used essentially in the proof of the exactness theorem.

Remark 1.6. For  $\ell = 1$  the faithful flatness of  $\Lambda_{\xi}^{-} = \mathbb{Z}[[t]][t^{-1}]$  over  $\Lambda_{(\xi)} = S^{-1}\mathbb{Z}[t, t^{-1}]$  is well known (see [11, ch. 3]). This property implies that the equation  $Px = Q$  where  $P, Q \in \mathbb{Z}[t, t^{-1}]$  is solvable in  $\mathbb{Z}[[t]][t^{-1}]$  if and only if it is solvable in  $S^{-1}\mathbb{Z}[t, t^{-1}]$  (where  $S$  is the multiplicative subset  $\{1 + tP(t)\}$ ). This fact was known already to Hurwitz (see [12, problem 156]).

## §2. The numbers $b_*(\xi), q_*(\xi)$

Suppose that  $C_*$  is a free finitely generated complex over  $\Lambda = \mathbb{Z}[z^{\ell}]$  and  $\xi : \mathbb{Z}^{\ell} \rightarrow \mathbb{R}$  is a monomorphism. Consider the complex  $S^{-1}C_* = C_* \otimes_{\Lambda} \Lambda_{(\xi)}$ . The homology modules

$$H_*(C_* \otimes_{\Lambda} \Lambda_{(\xi)}) = H_*(C_*) \otimes_{\Lambda} \Lambda_{(\xi)}$$

are finitely generated over principal ideal domain  $\Lambda_{(\xi)}$ ; therefore the rank  $b_p(C_*, \xi)$  and the torsion number  $q_p(C_*, \xi)$  of the module  $H_p(C_*) \otimes_{\Lambda} \Lambda_{(\xi)}$  are defined (the torsion number of a module is by definition the minimal number of generators in the submodule of torsion elements). For a manifold  $M^n$  and a maximally irrational  $\xi \in H^1(M, \mathbb{R})$  we set  $b_p(M, \xi) = b_p(C_*(\bar{M}), \xi)$ ,



$q_p(M, \xi) = q_p(C_*(\bar{M}), \xi)$ , where  $\bar{M} \rightarrow M$  is the maximal free abelian cover. Note that in the introduction we defined these numbers using the modules  $H_*(C_*(\bar{M}) \otimes_{\Lambda} \Lambda_{\xi}^-)$ ; these two definitions are the same. Indeed, the faithful flatness of  $\Lambda_{\xi}^-$  over  $\Lambda_{(\xi)}$  implies

$$\begin{aligned} H_*(C_* \otimes_{\Lambda} \Lambda_{\xi}^-) &= H_*(C_* \otimes_{\Lambda_{(\xi)}} \Lambda_{(\xi)} \otimes_{\Lambda_{(\xi)}} \Lambda_{\xi}^-) = \\ &= H_*(C_* \otimes_{\Lambda_{(\xi)}} \Lambda_{(\xi)}) \otimes_{\Lambda_{(\xi)}} \Lambda_{\xi}^- = (H_*(C_*) \otimes_{\Lambda_{(\xi)}} \Lambda_{(\xi)}) \otimes_{\Lambda_{(\xi)}} \Lambda_{\xi}^- \end{aligned}$$

Next we'll study the behaviour of  $b_p(C_*, \xi)$ ,  $q_p(C_*, \xi)$ , where  $C_*$  is fixed and  $\xi$  varies. For this purpose we need lemma 2.1. Denote by  $M$  the set of all monomorphisms  $\mathbb{Z}^{\ell} \rightarrow \mathbb{R}$ ,  $M \subset \mathbb{R}^{\ell}$ .

Lemma 2.1. Let  $a_1, \dots, a_n \in \Lambda$ . Then the set of  $\xi$ , for which the greatest common divisor of elements  $a_1, \dots, a_n$  (abbreviated further as g.c.d.  $(a_1, \dots, a_n)$ ) in  $\Lambda_{(\xi)}$  is equal to 1, is the intersection of  $M$  and several components (maybe none at all) of the complement  $U = \mathbb{R}^{\ell} \setminus \bigcup_i \Gamma_i$  in  $\mathbb{R}^{\ell}$  to the finite union of some integer hyperplanes (i.e. hyperplanes, determined by linear equations with integer coefficients)  $\Gamma_i \subset \mathbb{R}^{\ell}$ .

Proof. Denote by  $A$  the g.c.d. of elements  $a_1, \dots, a_n$  in the unique factorization domain  $\Lambda$ . Then the set sought consists of those  $\xi$ 's for which the principal coefficient of polynomial  $A$  with respect to  $\xi$  is equal to 1. Denote by  $\langle A \rangle$  the convex hull in  $\mathbb{R}^{\ell}$  of the subset  $\text{supp } A \subset \mathbb{Z}^{\ell}$ . For an edge  $\gamma$  of the polyhedron  $\langle A \rangle$  denote by  $\Gamma_{\gamma}$  the hyperplane in  $\text{Hom}(\mathbb{Z}^{\ell}, \mathbb{R}) = \mathbb{R}^{\ell}$ , consisting of all the homomorphisms  $\xi$ , vanishing on  $\gamma$ . The lemma is now obvious: the set  $U$  is  $\mathbb{R}^{\ell} \setminus \bigcup_i \Gamma_i$ .

Theorem 2.2. (J.-C. Sikorav). 1. The number  $b_p(C_*, \xi)$  does not depend on  $\xi$ ; it is equal to the rank of the module  $H_p(C_*) \otimes_{\Lambda} \{\Lambda\}$  over the fraction field  $\{\Lambda\}$  of  $\Lambda$ .

2. There exists a finite collection of integer hyperplanes  $\Gamma_i \subset \text{Hom}(\mathbb{Z}^l, \mathbb{R}) = \mathbb{R}$ , such that  $q_p(C_*, \xi)$  does not depend on  $\xi$  in any connected component of  $\mathbb{R}^l \setminus \bigcup_i \Gamma_i$ .

Proof. 1. The module

$$H_p(C_* \otimes_{\Lambda} \Lambda_{(\xi)}) = H_p(C_*) \otimes_{\Lambda} \Lambda_{(\xi)}$$

can be presented as a sum of a free module of rank  $b_p(C_*, \xi)$  and a torsion module. When we pass to  $\{\Lambda\}$ , which means additional localization, the torsion module disappears and the free module of rank  $b_p(C_*, \xi)$  survives.

2. Recall that the  $p$ th torsion number  $q_p$  for a complex

$$C_{p-1} \xleftarrow{\partial_{p-1}} C_p \xleftarrow{\partial_p} C_{p+1}$$

over a principal ideal domain can be calculated as follows. Consider the matrix  $D$  of the homomorphism  $\partial_p$ . Denote by  $d_r$  the g.c.d. of the  $r$ -minors of  $D$  and by  $\delta$  - the greatest  $r$  for which  $d_r = 1$ . Then  $q_p = \text{rk } D - \delta$ .

Now our assertion is easily deduced from lemma 2.1.

Remark 2.3. One can prove (similar to [13], [14]) that the number  $b_p(C_*, \xi)$  is equal to the dimension of  $p$ th homology of  $C_*$  with coefficients in 1-dimensional local system, determined by a generic representation  $\rho : t_i \rightarrow \rho(t_i) \in \mathbb{C}$ . The numbers  $b_p(C_*, \xi)$  can be computed in terms of usual homology with real coefficients and Massey operations, see [13].

Now let  $\xi$  be a homomorphism  $\mathbb{Z}^l \rightarrow \mathbb{R}$ , which is not con-

tained in  $\bigcup_i \Gamma_i$ . Define the numbers  $b_p(C_*, \xi) = b_p(C_*)$ ,  $q_p(C_*, \xi)$  by setting  $b_p(C_*, \xi) = b_p(C_*, \xi')$ ,  $q_p(C_*, \xi) = q_p(C_*, \xi')$ , where  $\xi'$  is any maximally irrational homomorphism sufficiently close to  $\xi$ .

The numbers  $b_p(M)$ ,  $q_p(M, \xi)$ <sup>2)</sup> are defined and calculated in terms of maximal free abelian covering  $\bar{M} \rightarrow M$ . It appears however that for a generic class  $\xi$  they can be computed in terms of cyclic covering  $M \rightarrow M$  and coincide with corresponding classical Novikov numbers of [1], [2].

To prove this we need a simple lemma (which we'll use also many times in the sequel).

Lemma 2.4. Let  $M, N$  be finitely generated modules over a noetherian commutative ring  $W$ , and  $S$  be a multiplicative subset of  $W$ . For  $\sigma \in W$  we denote  $S_{(\sigma)}$  the multiplicative subset, generated by  $\sigma$ .

Then 1) if  $f : S^{-1}M \rightarrow S^{-1}N$  is a homomorphism of  $S^{-1}W$ -modules then there exist  $\sigma \in W$  and  $f_{(\sigma)} : S_{(\sigma)}^{-1}M \rightarrow S_{(\sigma)}^{-1}N$ , such that  $S^{-1}f_{(\sigma)} = f$ ;

2) if  $\theta \in W$  and  $f_{(\theta)}, f'_{(\theta)} : S_{(\theta)}^{-1}M \rightarrow S_{(\theta)}^{-1}N$  are homomorphisms of  $S_{(\theta)}^{-1}W$ -modules, such that  $S_{(\theta)}^{-1}f_{(\theta)} = f = S_{(\theta)}^{-1}f'_{(\theta)}$ , then there exists  $\sigma' \in W$ , such that  $S_{(\sigma', \theta)}^{-1}f_{(\theta)} = S_{(\sigma', \theta)}^{-1}f'_{(\theta)}$ ;

3) if  $f : S^{-1}M \rightarrow S^{-1}N$  is an isomorphism then there exists  $\sigma \in W$  and  $f_{(\sigma)} : S_{(\sigma)}^{-1}M \rightarrow S_{(\sigma)}^{-1}N$ , such that  $f_{(\sigma)}$  is an isomorphism and  $S^{-1}f_{(\sigma)} = f$ .

Proof. 1) Denote by  $M_0, N_0$  the kernels of localization

---

2) Our definitions here differ from the standard ones: usually  $b_p(M)$  denotes the  $p$ th Betti number of  $M$ .

maps  $M \rightarrow S^{-1}M$ ,  $N \rightarrow S^{-1}N$ ; they are finitely generated, since  $W$  is noetherian.

Consider any  $\gamma \in S$ , annihilating both  $M_0$  and  $N_0$ . The localization maps  $S_{(\gamma)}^{-1}M \rightarrow S^{-1}M$ ,  $S_{(\gamma)}^{-1}N \rightarrow S^{-1}N$  are injective. Pick any finite system of generators  $m_i$  of  $M$ . There exists  $\gamma \in S$ , such that  $f(m_i) \in S_{(\gamma)}^{-1}N$ . Thus the modules  $S_{(\gamma\lambda)}^{-1}M$ ,  $S_{(\gamma\lambda)}^{-1}N$  are the submodules of  $S^{-1}M$  and correspondingly of  $S^{-1}N$ , and  $f$  sends one of them into the other. Now we set  $\sigma = \gamma\lambda$ ,  $f_{(\sigma)} = f|_{S_{(\sigma)}^{-1}M}$ .

The first claim of the lemma is proved.

2) The homomorphisms  $S_{(\gamma)}^{-1}f_{(\theta)}$  and  $S_{(\gamma)}^{-1}f'_{(\theta)}$  both are restrictions of the homomorphism  $f : S^{-1}M \rightarrow S^{-1}N$  to submodule  $S_{(\theta\gamma)}^{-1}M \subset S^{-1}M$ .

3) follows from the first two points.

Corollary 2.5. Let  $C_*$  and  $D_*$  be finitely generated free complexes over a commutative noetherian ring  $W$ . Suppose that  $S^{-1}C_* \sim S^{-1}D_*$ . Then for some  $\sigma \in W$  we have

$$S_{(\sigma)}^{-1}C_* \sim S_{(\sigma)}^{-1}D_*$$

Return now to numbers  $b_*$ ,  $q_*$ . Let  $C_*$  be a free finitely generated complex over the ring  $\Lambda = \mathbb{Z}[\mathbb{Z}^k]$ . Let  $\xi : \mathbb{Z}^k \rightarrow \mathbb{R}$  be a homomorphism of irrationality degree 1. The image of  $\xi$  is isomorphic to  $\mathbb{Z}$ . Denote by  $(\xi)$  the homomorphism of the group rings  $\Lambda \rightarrow \mathbb{Z}[\mathbb{Z}] = L$  which is obtained from the composition of  $\xi : \mathbb{Z}^k \rightarrow \mathbb{Z}$  with  $(-1) : \mathbb{Z} \rightarrow \mathbb{Z}$  by passing to group rings ( $(-1)$  is due to our sign conventions). Denote by  $S \subset L$  the multiplicative subset, consisting of Laurent polynomials with prin-

cipal coefficient 1. Set

$$\beta_p(C_*, \xi) = \tau^k (H_p(C_* \otimes_{\Lambda} S^{-1}L)),$$

$$\alpha_p(C_*, \xi) = q (H_p(C_* \otimes_{\Lambda} S^{-1}L)).$$

Here the structure of  $\Lambda$ -module on  $S^{-1}L$  is defined via homomorphism  $(\xi) : \Lambda \rightarrow L \subset S^{-1}L$  (so the  $S^{-1}L$ -module  $H_p(C_* \otimes_{\Lambda} S^{-1}L)$  depends on  $\xi$ ).

Lemma 2.6. Any maximally irrational homomorphism  $\gamma : \mathbb{Z}^k \rightarrow \mathbb{R}$  possesses an open conical neighbourhood  $U(\gamma)$  in the set  $\text{Hom}(\mathbb{Z}^k, \mathbb{R})$ , such that the following equalities hold for  $\xi \in U(\gamma)$ :

$$\beta_p(C_*, \gamma) = \beta_p(C_*, \xi), \quad q_p(C_*, \gamma) = \alpha_p(C_*, \xi)$$

Proof. Consider for all  $p$  isomorphisms

$$H_p(C_*) \otimes_{\Lambda(\gamma)} \approx \left( \bigoplus_{i=1}^{b_p(C_*, \gamma)} \Lambda(\gamma) \right) \oplus \left( \bigoplus_{j=1}^{q_p(C_*, \gamma)} \Lambda(\gamma) / a_j^{(p)} \Lambda(\gamma) \right)$$

where  $a_j^{(p)} \in \Lambda = \mathbb{Z}[\mathbb{Z}^k]$  is divisible in  $\Lambda$  by  $a_{j-1}^{(p)}$  and  $\gamma$ -principal terms of Laurent polynomials  $a_j^{(p)}$  are equal to  $\alpha_j^{(p)} \cdot 1$ , where  $\alpha_j^{(p)} \in \mathbb{Z}$ ,  $\alpha_j^{(p)} \neq \pm 1$ . Consider the free finitely generated  $\Lambda$ -complex  $D_*$ , defined as follows. The module  $D_p$  is the sum of free modules  $F_p, E_p, B_p$  of the ranks correspondingly  $b_p(C_*, \gamma), q_p(C_*, \gamma), q_{p-1}(C_*, \gamma)$ . The differential  $d_p : D_p \rightarrow D_{p-1}$  vanishes on  $F_p \oplus E_p$  and sends the  $k$ th free generator  $b_k$  of the module  $B_p$  to the element  $a_k^{(p)} \cdot e_k$  where  $e_k$  is the  $k$ th free generator of the module  $E_p$ . It is known that any complex over principal ideal domain is homotopy equivalent to a standard one like this; from this we easily deduce

that  $C_* \otimes \Lambda(\gamma)$  is homotopy equivalent to  $D_* \otimes \Lambda(\gamma)$ . Since  $\Lambda$  is noetherian,  $C_*$  and  $D_*$  become equivalent when localized with respect to the multiplicative subset generated by a single element  $\sigma \in S_\gamma$  (see lemma 2.6). Consider now any rational homomorphism  $\xi: \mathbb{Z}^k \rightarrow \mathbb{Q}$  which is close enough to  $\gamma$ , so that

- 1) each polynomial  $a_j^{(p)}$  has only one  $\xi$ -principal term, namely  $\alpha_j^{(p)} \cdot 1$ ;
- 2)  $\sigma \in S_\xi$ .

The complexes  $C_* \otimes_{\Lambda} \Lambda(\xi)$ ,  $D_* \otimes_{\Lambda} \Lambda(\xi)$  are homotopy equivalent over  $\Lambda(\xi)$ . The homomorphism  $(\xi): \Lambda \rightarrow S^{-1}\Lambda$  can be factored through  $S_\xi^{-1}\Lambda = \Lambda(\xi)$  and to calculate the homology of  $C_* \otimes S^{-1}\Lambda$  we can use the complex

$$D_* \otimes \Lambda(\xi) \otimes S^{-1}\Lambda = D_* \otimes_{\Lambda} S^{-1}\Lambda$$

The complex  $D_*$  is described above; using 1) we easily get

$$H_p(D_* \otimes S^{-1}\Lambda) \approx \left( \bigoplus_{i=1}^{b_p(C_*, \gamma)} S^{-1}\Lambda \right) \oplus \left( \bigoplus_{j=1}^{q_p(C_*, \gamma)} S^{-1}\Lambda / \tilde{a}_j^{(p)} S^{-1}\Lambda \right),$$

where  $\tilde{a}_j^{(p)} \in S^{-1}\Lambda$ ,  $\tilde{a}_j^{(p)} : \tilde{a}_{j-1}^{(p)}$ ; the  $\gamma$ -principal coefficient of  $\tilde{a}_j^{(p)}$  is equal to  $\alpha_j^{(p)} \neq \pm 1$ . Now the lemma follows easily.

The argument applied here will often be used in the sequel.

Remark 2.7. It is easy to see that for any connected manifold  $M$  and any maximally irrational  $\xi \in H^1(M, \mathbb{R})$  the numbers  $b_0(\xi)$  and  $q_0(\xi)$  vanish. Indeed, the module  $H_0(\bar{M}) \approx \mathbb{Z}$  is annihilated by any element of  $\mathbb{Z}[\mathbb{Z}^\ell]$  of the type  $1 - t$ , where  $t \in \mathbb{Z}^\ell$ . Choosing  $t \in \mathbb{Z}^\ell$  with  $\xi(t) < 0$  we get  $S^{-1}H_0(\bar{M}) = 0$ .

§ 3. Duality properties

Recall first the formulation of Poincare duality for non-simply-connected manifolds.

Let  $M^n$  be a smooth manifold (not necessary orientable). The universal covering  $\tilde{M}^n$  is equipped with the fundamental  $n$ -cycle  $U$  (infinite in general). The intersection  $\cap U$  defines an isomorphism  $D$  between cohomology  $H_c^*(\tilde{M}^n, \mathbb{Z})$  with compact support and homology  $H_{n-*}(M^n, \mathbb{Z})$ . Define the automorphism  $\chi$  of the group ring  $\mathbb{Z}[\pi_1 M]$  by setting  $\chi(g) = \varepsilon(g)g^{-1}$ , where  $g \in \pi_1 M$ ,  $\varepsilon(g) = -1$  if the orientation of  $M$  is changed along  $\gamma$  and  $\varepsilon(g) = 1$  if not.

To simplify the statements we'll suppose that  $\pi_1 M$  is abelian (we'll need only this case in the present paper).

Homology and cohomology groups of  $M$  are the  $\mathbb{Z}[\pi_1 M]$ -modules, and the isomorphism  $D$  is subjected to the following commutativity relation:  $D(gx) = \chi(g)D(x)$ . Isomorphisms of that kind are called  $\chi$ -isomorphisms. Thus we have the  $\chi$ -isomorphism of  $\mathbb{Z}[\pi_1 M]$ -modules

$$D: H_c^*(\tilde{M}^n, \mathbb{Z}) \longrightarrow H_{n-*}(\tilde{M}^n, \mathbb{Z}).$$

Note that there exists the natural isomorphism of  $\mathbb{Z}[\pi_1 M]$ -modules

$$H_c^*(\tilde{M}^n, \mathbb{Z}) \rightarrow H_* \left( \text{Hom}_{\mathbb{Z}[\pi_1 M]} (C_*(\tilde{M}^n), \mathbb{Z}[\pi_1 M]) \right).$$

For a  $\mathbb{Z}[\pi_1 M]$ -module  $G$  we denote the modules

$$H_* \left( \text{Hom}_{\mathbb{Z}[\pi_1 M]} (C_*(\tilde{M}^n), G) \right),$$

$$H_* \left( C_*(\tilde{M}^n) \otimes_{\mathbb{Z}[\pi_1 M]} G \right)$$

by  $H^*(M^n, G), H_*(M^n, G)$ .

Thus we have the  $\chi$ -isomorphism of  $\mathbb{Z}[\pi_1 M]$ -modules

$$D: H^*(M, \mathbb{Z}[\pi_1 M]) \rightarrow H_{n-*}(\tilde{M}) \quad (3.1)$$

Now we turn to the case  $\pi_1 M = \mathbb{Z}^l$ . Set  $\Lambda = \mathbb{Z}[\pi_1 M]$ . Let  $\gamma$  be a homomorphism  $\mathbb{Z}^l \rightarrow \mathbb{R}$  (not necessarily injective). It is clear that  $\chi$  sends  $S_\gamma$  to  $S_{-\gamma}$  and thus defines an isomorphism  $\chi: \Lambda_{(\gamma)} \rightarrow \Lambda_{(-\gamma)}$ . Localizing  $D$ , we get the following lemma.

Lemma 3.1. There exists a  $\chi$ -isomorphism

$$D_\gamma: H^p(M^n, \Lambda_{(\gamma)}) \rightarrow H_{n-p}(M^n, \Lambda_{(-\gamma)}) \quad (3.2)$$

of  $\Lambda_{(\gamma)}$ -module (left) and  $\Lambda_{(-\gamma)}$ -module (right).

Module  $G$  over a ring  $W$  is called principal, if it is isomorphic to a direct sum of modules of the type  $W/aW$  (where  $a \in W$ ). Suppose now that all the modules  $H_i(M^n, \Lambda_{(\gamma)})$  are principal for  $r \leq p$  and fix the decompositions

$$H_r(M^n, \Lambda_{(\gamma)}) \cong \left( \bigoplus_{i=1}^{\beta_r} \Lambda_{(\gamma)} \right) \oplus \left( \bigoplus_{i=1}^{\alpha_r} \Lambda_{(\gamma)} / a_i^{(r)} \Lambda_{(\gamma)} \right)$$

$$0 \leq r \leq p.$$

Applying lemma 5.1 from §5 (which provides the standard presentation for a complex with principal homology) to the complex  $C_*(\tilde{M}^n) \otimes \Lambda_{(\gamma)}$ , we get

$$H^r(M^n, \Lambda_{(\gamma)}) \cong \left( \bigoplus_{i=1}^{\beta_r} \Lambda_{(\gamma)} \right) \oplus \left( \bigoplus_{j=1}^{\alpha_{r-1}} \Lambda_{(\gamma)} / a_j^{(r-1)} \Lambda_{(\gamma)} \right), \quad 0 \leq r \leq p. \quad (3.3)$$

Now the Poincare duality (3.2) implies



$$H_q(M^n, \Lambda(-\gamma)) \cong \left( \bigoplus_{i=1}^{\beta_{n-q}} \Lambda(-\gamma) \right) \oplus \bigoplus_{i=1}^{\alpha_{n-q-1}} \left( \bigoplus_{i=1}^{\alpha_{n-q-1}} \Lambda(-\gamma) / \chi(a_i^{\alpha_{n-q-1}}) \Lambda(-\gamma) \right), n-p \leq q \leq n. \quad (3.4)$$

Corollary 3.2. For a monomorphism  $\gamma : \mathbb{Z}^l \rightarrow \mathbb{R}$ ,

$$b_k = b_{n-k}, \quad q_k(\gamma) = q_{n-k-1}(-\gamma)$$

Proof. First equality follows immediately from the above. To prove the second we recall that for any module  $N$  over a principal ideal domain  $R$  the number  $q_k(N)$  equals the number of nonzero ideals  $J$  in any decomposition  $N = R/J_1 \oplus \dots \oplus R/J_n$  where  $J_i \subset J_{i+1}$ . We can choose decomposition (3.3) to be of this kind, then the decomposition (3.4) is also of this kind, Q.E.D.

#### § 4. V.V. Sharko's results concerning Morse function on non-simply-connected cobordisms

In this paragraph we recall from [8] some results, which we'll use later. Let  $(W; V_0, V_1)$  be a manifold with boundary  $\partial W$ , consisting of two components  $V_0, V_1$ . Any regular Morse function  $f : W \rightarrow \mathbb{R}$ , constant on  $V_0$  and on  $V_1$ , together with the suitable gradient-like vector field gives rise to a Morse complex, defined over  $\mathbb{Z}[\pi_1 W]$ . This complex is simply homotopy equivalent to  $C_*(\tilde{W}^n, \tilde{V}_0)$ , where  $\tilde{W}^n$  stands for the universal covering of  $W$ . In some cases the inverse also holds, i.e. the given free finitely generated complex  $C_*$  over  $\mathbb{Z}[\pi_1 W]$ , simply homotopy equivalent to  $C_*(\tilde{W}^n, \tilde{V}_0)$  can be realized as a Morse complex for some Morse function on  $(W; V_0, V_1)$ .

Theorem 4.1. (see [8, proposition 6.1]). Suppose that

$\pi_1(V_0) \rightarrow \pi_1(W) \leftarrow \pi_1(V_1)$  are isomorphisms. Then any free finitely generated  $\mathbb{Z}[\pi_1 W]$ -complex of the type

$$C_* = \{0 \leftarrow C_2 \leftarrow C_3 \leftarrow \dots \leftarrow C_{n-2} \leftarrow 0\} \quad (4.1)$$

simply homotopy equivalent to  $C_*(\tilde{W}_n, \tilde{V}_0^{n-1})$ , can be realized as a Morse complex of some regular Morse function  $f : W \rightarrow \mathbb{R}$ , which is constant on  $V_1$  and on  $V_0$ .

The idea of the proof is the following (see [8]). First we choose a regular Morse function  $f$  on the cobordism  $(W; V_0, V_1)$  without critical points of indices  $0, 1, n-1, n$  (this is possible since  $\pi_1(V_0) \rightarrow \pi_1(W) \leftarrow \pi_1(V_1)$  are isomorphisms). Denote the corresponding Morse complex by  $C_*(f_0)$ . The complexes  $C_*, C_*(f_0)$  are simply homotopy equivalent. By Cockroft-Swan [15]  $C_*(f_0) \oplus D_1 \approx C_* \oplus D_2$ , where complexes  $D_i$  ( $i = 0, 1$ ) are direct sums of complexes of the type  $0 \leftarrow F \xleftarrow{\text{id}} F \leftarrow 0$ . The complexes  $D_1, D_2$  can be chosen so as to concentrate in dimensions  $2 \leq * \leq n-2$  (as  $C_*, C_*(f_0)$  do). The procedure of adding (or subtracting) the complex  $0 \leftarrow F \xleftarrow{\text{id}} F \leftarrow 0$  can be realized by corresponding changes of the function  $f_0$ , so that we obtain at an end a function  $f$  with the Morse complex  $C_*$ . The details can be found in [8].

Remark. We'll need this theorem for  $\pi_1 W = \mathbb{Z}$ . In this case the notions of the homotopy equivalence and the simple homotopy equivalence coincide. Therefore, any complex of the type (4.1), homotopy equivalent to  $C_*(\tilde{W}_n, \tilde{V}_0^{n-1})$  can be realized as a Morse complex of some function.

The paper [8] contains also the results concerning minimal

Morse functions. We reproduce them here partially. We won't need them in the proof of theorem 1 and we'll make use of them only in §6 and in remark 2 of §7.

For some classes of fundamental groups any cobordism  $(W; V_0, V_1)$ , where  $\pi_1(V_0) \rightarrow \pi_1(W) \leftarrow \pi_1(V_1)$  are isomorphisms, possesses a Morse function, having a minimal possible (among all the Morse functions) number of critical points of all indices. Namely, let  $\text{Wh}(\pi_1 W) = 0$ . Theorem 4.1 implies that existence of such function is guaranteed if we find among the free finitely generated  $\mathbb{Z}[\pi_1 W]$ -complexes of the type (4.1) the complex  $C_*^0$ , which has the minimal possible number of generators in each dimension. The following theorem can be proved purely algebraically.

Theorem 4.2 (see [8, proposition 4.8]). Let  $Q$  be an IBN-ring (which means that free modules  $Q^n$  and  $Q^m$  are not isomorphic if  $m \neq n$ ) and also an s-ring (which means that for any finitely generated  $Q$ -module  $N$  the condition  $N \oplus Q^n \approx Q^{\ell}$  implies  $N \approx Q^{\ell-n}$ ).

Then for any free finitely generated  $Q$ -complex  $C_*$  there exists a free finitely generated  $Q$ -complex  $C_*'$ , homotopy equivalent to  $C_*$ , which has the minimal possible number of generators in each dimension among free finitely generated complexes, homotopy equivalent to  $C_*$ . The complex  $C_*'$  is called minimal.

There is also a simple criterion to decide whether a given complex is minimal (in its homotopy type) or not.

Definition. The pair  $(N, M)$  of modules over a ring  $Q$ , where  $N \subset M$ , is called irreducible if there exists no module  $F \subset N$ , which is a direct summand of  $M$ . The pair  $(N, M)$  is

called strongly irreducible, if for any  $n \geq 0$  the greatest possible rank of a free submodule  $F$  of  $N \oplus Q^n$ , which is a direct summand of  $M \oplus Q^n$ , equals  $n$ . Our terminology differs here from that of [8].

Theorem 4.3 (see [8, theorem 4.7]). A  $Q$ -complex

$$C_* = \{0 \leftarrow C_0 \xleftarrow{\partial_1} \dots \xleftarrow{\partial_n} C_n \leftarrow 0\}$$

is minimal if and only if for each  $i$  the pair  $(\partial_i(C_i), C_{i-1})$  is strongly irreducible. (We imply that  $Q$  is an IBN, s-ring.)

If the ring  $Q$  possesses a non-trivial homomorphism to a field, then the rank of a free module is correctly defined. All the group rings  $\mathbb{Z}[G]$  enjoy this property. The ring  $\mathbb{Z}[\mathbb{Z}^l]$  is also an s-ring, see [16], [17].

### §5. Algebraic lemmas

First of all we prove the lemma (already used in §3) on the standard presentation for complexes.

Lemma 5.1. Let  $W$  be a commutative ring and

$$C_* = \{0 \leftarrow C_1 \xleftarrow{\dots} \xleftarrow{\partial_n} C_n \leftarrow 0\}$$

be a free finitely generated  $W$ -complex. Suppose that for  $p \leq k$  the  $W$ -modules  $H_p(C_*)$  have the free resolutions of length 2 :

$$0 \leftarrow H_p(C_*) \xleftarrow{\pi_p} F_p \xleftarrow{\varphi_p} G_p \leftarrow 0$$

Then  $C_*$  is homotopy equivalent to a free finitely generated complex

$$C'_* = \{0 \leftarrow C'_1 \xleftarrow{\dots} \xleftarrow{\partial'_n} C'_n \leftarrow 0\}$$

such that

1)  $C'_* = C_*$  if  $* \geq k+2$  ;

2) for  $p \leq k$  we have  $C'_p = G_{p-1} \oplus F_p$  ; furthermore,

$$\partial'_p|_{F_p} = 0, (\partial'_p|_{G_{p-1}})' = \varphi_{p-1}: G_{p-1} \rightarrow F_{p-1} \subset C'_{p-1}$$

and  $\text{Im}(\partial'_{p+1}: C'_{p+1} \rightarrow C'_p) \subset C'_p$  coincides with  $\text{Im} \varphi_p \subset F_p$ .

Proof is by induction on  $p$ . For  $p = 0$  the assertion is obvious. To produce the induction step we suppose that the assertion is proved for all  $k < m$  and prove it for  $k = m$ .

Suppose that  $C_*$  satisfies the assumptions of our lemma for  $k = m$ . Find a complex  $C'_*$  which satisfies the conclusion of lemma for  $k = m - 1$ . The image

$$\text{Im}(\partial'_m: C'_m \rightarrow C'_{m-1})$$

is a free module  $\text{Im} \varphi_{m-1} \approx G_{m-1}$ , hence  $C'_m$  can be decomposed as  $G_{m-1} \oplus K_m$ , where  $\partial'_m|_{G_{m-1}} = \text{id}$ . We may assume that  $K_m$  is a free finitely generated  $W$ -module (having added if necessary a complex  $0 \leftarrow G_{m-1} \xleftarrow{\text{id}} G_{m-1} \leftarrow 0$ , located in dimensions  $m, m+1$ , to  $C'_*$ ). The complex  $C'_*$  splits into the sum of two complexes

$$E_* = \{0 \leftarrow C'_1 \leftarrow \dots \leftarrow C'_{m-2} \leftarrow C'_{m-1} \leftarrow 0\}$$

$$D_* = \{0 \leftarrow \dots \leftarrow 0 \leftarrow K_m \leftarrow \dots\}$$

( $D_*$  is located in dimensions  $\geq m$ ). Here  $H_m(D_*) \approx H_m(C'_*) \approx H_m(C_*)$ .

Now add the complex  $0 \leftarrow F_m \xleftarrow{\text{id}} F_m \leftarrow 0$ , located in dimensions  $m, m+1$ , to  $D_*$  and consider the resulting complex  $D'_*$ .

There is an epimorphism

$$D'_m = K_m \oplus F_m \xrightarrow{P} H_m(D'_*) \approx H_m(C_*),$$

where  $P|_{F_m} = 0$ ,  $P|_{K_m}$  is a natural projection.

The map  $P' : K_m \oplus F_m \rightarrow H_m(C_*)$ , defined by  $P'|_{K_m} = 0$ ,  $P'|_{F_m} = \pi_m$ , is another epimorphism onto the same module.

By [8], lemma 1.10 there exists an isomorphism

$$\varphi : K_m \oplus F_m \longrightarrow K_m \oplus F_m$$

such that  $P' \circ \varphi = P$ . That means that we can find another decomposition  $D'_m = K_m \oplus F_m$  (i.e. choose another free basis), such that the projection  $P : D'_m \rightarrow H_m(D'_*) = H_m(C_*)$  is given by  $P|_{K_m} = 0$ ,  $P|_{F_m} = \pi_m$ . It is clear now that we can split the complex  $0 \leftarrow K_m \leftarrow K_m \leftarrow 0$  (located in dimensions  $m, m+1$ ) off the  $D'_*$ . In the resulting complex

$$D''_m = \{ 0 \leftarrow \dots \leftarrow F_m \xleftarrow{\partial''_{m+1}} \dots \}$$

the image  $\text{Im } \partial''_{m+1}$  coincides with  $\text{Im } \varphi_m \subset F_m$ , since the projection  $D''_m \rightarrow H_m(D''_m) \approx H_m(C_*)$  coincides with  $\pi_m$ . It is clear now that the complex  $E_* \oplus D''_*$  satisfies the points 1), 2) of the conclusion for  $k = m$ .

Remark. We do not prove (and do not use) assertions concerning simple homotopy equivalence.

In order to state our next lemma, we introduce some notations.

Set  $\Lambda = \mathbb{Z}[\mathbb{Z}^l]$  and let  $\gamma : \mathbb{Z} \rightarrow \mathbb{R}$  be a rational homomorphism (i.e.  $r \mathbb{k} \text{ Im } \gamma = 1$ ). We'll study the localizations  $S^{-1}A$  of  $\Lambda$ -modules  $A$ . We can assume that there is chosen a system of generators  $(t, t_1, \dots, t_{l-1})$  for a group  $\mathbb{Z}^l$ , such that  $\gamma$  is the projection of  $\mathbb{Z}^l$  onto the direct summand  $\mathbb{Z}$ , generat-

ed by  $t$ , and that  $\gamma(t) = -1$ .

Set

$$R = \mathbb{Z}[t_1^{\pm 1}, \dots, t_{\ell-1}^{\pm 1}] = \mathbb{Z}[\mathbb{Z}^{\ell-1}],$$

$$P = R[t], \quad S = \{1 + tQ(t) \mid Q(t) \in R[t]\}, \quad (5.1)$$

$$K = S^{-1}P, \quad S' = \{t^n \mid n \in \mathbb{N}\}, \quad \Gamma = S'^{-1}S^{-1}P = S'^{-1}\Lambda.$$

Lemma 5.2. Let  $A$  be a finitely generated  $P$ -module, such that  $S'^{-1}A$  is a principal  $\Gamma$ -module. Fix a decomposition

$$S'^{-1}A = \left( \bigoplus_{i=1}^{\ell} \Gamma(e_i) \right) \oplus \left( \bigoplus_{j=1}^{\rho} (\Gamma/a_j\Gamma)(f_j) \right),$$

where  $a_j = a_{j,0} + a_{j,1}t + \dots$ ,  $a_{j,k} \in P$ ,  $a_{j,0} \neq 0$  is non-invertible element of  $R$ .

Then there exists a finitely generated  $P$ -module  $B \subset A$ , such that

$$1) \quad S^{-1}B \approx \left( \bigoplus_{i=1}^{\ell} K(e'_i) \right) \oplus \left( \bigoplus_{j=1}^{\rho} (K/a_jK)(f'_j) \right),$$

$$2) \quad t^N A \subset B \text{ for some } N.$$

Proof. Note first that we may assume that  $e_i$  and  $f_i$  belong to  $A$ . Furthermore, having multiplied  $e_i$  and  $f_i$  by  $t^N$ , where  $N$  is large enough, we may assume that the module  $P(e_i, f_j) \subset A$  is free of  $t$ -torsion.

We'll show that  $P(e_i, f_j)$  satisfies 1).

Consider the  $K$ -submodule  $K(e_i, f_j)$  of the module  $S^{-1}A$ .

It is easy to see that

$$K(e_i, f_j) \approx \left( \bigoplus_{i=1}^{\ell} (S^{-1}P)(e_i) \right) \oplus \left( \bigoplus_{j=1}^{\rho} (S^{-1}P/a_j S^{-1}P)(f_j) \right)$$

We show that  $K(e_i, f_j)$  is the  $S$ -localization of  $P$ -module  $P(e_i, f_j)$ . Indeed,  $K(e_i, f_j)$  is an  $S^{-1}P$ -module, hence there exists an epimorphism

$$\varphi: S^{-1}(P(e_i, f_j)) \rightarrow K(e_i, f_j) \subset S_{\mathcal{Y}}^{-1}A$$

Now we show that  $\varphi$  is an isomorphism.

Denote by  $e_i, f_j$  the images of  $e_i, f_j \in P(e_i, f_j)$  under the map of this module to its localization; they generate over  $K$  the entire module  $S^{-1}(P(e_i, f_j))$ . Consider any  $x = \sum \alpha_i e_i + \sum \beta_j f_j$ , belonging to  $\text{Ker } \varphi$ , where  $\alpha_i, \beta_j \in P$ . Observe that  $\alpha_i e_i = 0 = \beta_j f_j$  in the module  $S^{-1}A$ , which means that

$$t^{N_i}(1+tQ_i(t))\alpha_i e_i = 0 = t^{M_j}(1+tW_j(t))\beta_j f_j$$

in module  $A$  for some natural  $N_i, M_j$  and  $Q_i(t), W_j(t) \in P$ . The module  $P(e_i, f_j)$  is free of  $t$ -torsion by construction, therefore

$$(1+tW_j(t))\beta_j f_j = 0 = (1+tQ_i(t))\alpha_i e_i$$

in  $A$ . It is clear now that  $x = 0$  and thus,  $\text{Ker } \varphi = 0$  and  $P(e_i, f_j)$  satisfies the requirement 1) of the conclusion.

Observe next that the localization  $S_{\mathcal{Y}}^{-1}P(e_i, f_j)$  coincides with the entire module  $S_{\mathcal{Y}}^{-1}A$ , and  $A$  is finitely generated. Therefore there exists an element  $t^N(1+tQ(t))$  of  $P$  such that  $t^N(1+tQ(t))a \in P(e_i, f_j)$  for all  $a \in A$ . Consider now the module

$$B = \{a \in A \mid \exists Q(t) \in P: (1+tQ(t))a \in P(e_i, f_j)\}$$



The noetherian property implies that  $B$  is finitely generated; it's obvious that  $S^{-1}B \approx S^{-1}P(e_i, f_j)$ . By construction  ${}^t N_A \subset B$ . Lemma is proved.

For a  $P$ -module  $M$  we denote by  $\text{Tor}_t M$  the submodule of elements, annihilated by some degree of  $t$ , by  ${}^t M$  - the submodule of elements, annihilated by  $t$ , by  $M_t$  - the factormodule  $M/tM$ .

Lemma 5.3. For a  $P$ -module  $M$  the localization map  $M \rightarrow S^{-1}M$  induces

1) an isomorphism  $M/tM \xrightarrow{\approx} S^{-1}M/tS^{-1}M$ ,

1a) an isomorphism  $M/t^q M \xrightarrow{\approx} S^{-1}M/t^q S^{-1}M$  for any natural  $q$ ,

2) a monomorphism  ${}^t M \rightarrow {}^t(S^{-1}M)$ ,

2a) a monomorphism  $\text{Tor}_t M \rightarrow \text{Tor}_t(S^{-1}M)$ .

Proof. 1) Injectivity: suppose that  $m \in M$  and  $m = \frac{t}{1+tQ(t)}n$  in  $S^{-1}M$ . Then

$$(1+tQ(t))\tilde{m} = tn + x$$

where  $(1+tQ_1(t))x = 0$ . Therefore  $(1+tQ_2(t))m = tn'$ , hence  $m \in tM$ .

$$\text{Surjectivity: } \frac{m}{1+tQ(t)} = m - t \frac{Q(t)m}{1+tQ(t)}$$

1a) Note that if  $n \in M$ ,  $Q(t) \in P$  the following equality holds in the module  $S^{-1}M/t^q S^{-1}M$ :

$$\frac{n}{1+tQ(t)} = (1-tQ(t) + \dots \pm (tQ(t))^{q-1})n$$

Now the surjectivity is straightforward. Further if  $m = \frac{t^q}{1+tQ(t)}n$ ,

where  $n, m \in M$ , the equality

$$m = t^q (1 - tQ(t) + \dots \pm (tQ(t))^{q-1}) n$$

holds in  $S^{-1}M/t^q S^{-1}M$  and injectivity is also proved.

2) if  $tm = 0$  and  $m \mapsto 0$  in  $S^{-1}M$ , then

$$(1 + tQ(t))m = 0 \Rightarrow m = 0.$$

2a) if  $t^N m = 0$  and  $(1 + tQ(t))m = 0$ , then

$$(t^{N-1} + t^N Q(t))m = 0 \Rightarrow t^{N-1} m = 0;$$

proceeding further in a similar way, we find  $m = 0$ .

Lemma 5.4. Let  $a = a_0 + a_1 t + \dots + a_N t^N \in P$ , where  $a_i \in R$ ,  $a_0 \neq 0$  is non-invertible element of  $R$ ,  $q$  is a natural number. Then

1) the factors  $K_q = K/(a, t^q)$ ,  $P_q = P/(a, t^q)$  of the rings  $K, P$  by the ideals, generated by  $a, t^q$ , are isomorphic as rings and as  $P$ -modules;

2) the ring  $P_q$  is isomorphic as an  $R$ -module to the module  $R^q/F(R^q)$ , where  $F: R^q \rightarrow R^q$  is injective and is given by the matrix

$$\begin{pmatrix} a_0 & a_1 & a_2 & \dots & a_q \\ & a_0 & a & \dots & a_{q-1} \\ \circ & & & \dots & \\ & & & & a_0 \end{pmatrix} \quad (5.2)$$

Thus the  $R$ -module  $P_q$  has a free resolution of length 2 ;

3) the multiplication map  $t^\ell: P_q \rightarrow P_{\ell+q}$  (where  $\ell > 0$ ) is injective and induces an epimorphism

$$\text{Ext}_R^1(P_{\ell+q}, R) \longrightarrow \text{Ext}_R^1(P_q, R).$$

The module  $\text{Hom}_R(P_q, R)$  is trivial.

Proof. 1) The embedding  $P \subset K$  induces a map  $P_q \rightarrow K_q$  which is a ring homomorphism and a  $P$ -module homomorphism. The map

$$P/(t^q) \longrightarrow K/(t^q)$$

is an isomorphism (see lemma 5.3), hence the map

$$(P/(t^q))/(a) \longrightarrow (K/(t^q))/(a)$$

also is an isomorphism.

2) The ring  $P/(t^q) = R[t]/(t^q)$  is a free  $R$ -module with basis  $1, t, \dots, t^{q-1}$ .

The multiplication by  $a \in P$  is given in this basis by the matrix (5.2). Since  $a_0 \neq 0$  and  $R$  has no zero divisors,  $F$  is a monomorphism.

3) Suppose that  $x \in \text{Ker}(t^l : P_q \rightarrow P_{l+q})$ , i.e.  $t^l x = Na + Mt^{l+q}$  where  $x, N, M \in P$ . The ring  $P$  is a unique factorization domain and  $t^l$  does not divide  $a$ , therefore  $N = t^l N'$ . Furthermore, since  $P$  has no zero divisors,  $x = N'a + Mt^q$ , hence  $x = 0$  in  $P_q$ .

Next observe that the map  $t^l : P_q \rightarrow P_{l+q}$  lifts to resolutions in a following way:

$$\begin{array}{ccccccc} 0 & \leftarrow & P_q & \leftarrow & R[t]/(t^q) & \xleftarrow{a} & R[t]/(t^q) \leftarrow 0 \\ & & \downarrow t^l & & \downarrow t^l & & \downarrow t^l \\ 0 & \leftarrow & P_{l+q} & \leftarrow & R[t]/(t^{q+l}) & \xleftarrow{a} & R[t]/(t^{q+l}) \leftarrow 0 \end{array}$$

The right arrow is a monomorphism onto the direct summand. This implies the surjectivity of the map of Ext's.

The equality  $\text{Hom}_R(P_q, R) = 0$  is obvious.

Lemma 5.5. 1) The ring  $K = S^{-1}P$  is an IBN-ring and an s-ring.

2) A free finitely generated K-complex

$$C_* = \{0 \leftarrow C_1 \leftarrow \dots \leftarrow C_e \leftarrow 0\}$$

is minimal if and only if the free finitely generated R-complex

$$C_* / tC_* = \{0 \leftarrow C_1 / tC_1 \leftarrow \dots \leftarrow C_e / tC_e \leftarrow 0\}$$

is minimal.

Proof. 1) We can embed  $K$  into its field of fractions, hence  $K$  is an IBN-ring. The ring  $R = K/tK$  is an s-ring, therefore to establish the s-property for  $K$  it is enough to prove the following: for a finitely generated  $K$ -module  $N$ , which is free of  $t$ -torsion any collection  $(n_1, \dots, n_s)$  of the elements of  $N$ , which forms a basis of  $N/tN$  over  $R$ , is a basis of  $N$  over  $K$ .

To prove this we consider the free module  $F(e_1, \dots, e_s)$  and the homomorphism  $\varphi : F(e_1, \dots, e_s) \rightarrow N$ , sending  $e_i$  to  $n_i$ . Denote  $\pi$  the projection  $N \rightarrow N/tN$ . Since  $\pi \circ \varphi$  is surjective,  $N/\text{Im} \varphi = t(N/\text{Im} \varphi)$ .

The element  $t$  belongs to the radical of  $K$ , and, using Nakayama's lemma we get  $N/\text{Im} \varphi = 0$ , i.e.  $\varphi$  is surjective. Next we show that  $\varphi$  is a monomorphism. Indeed, suppose that  $\sum a_i \cdot n_i = 0$  where  $a_i \in K$ , and some  $a_i$  is nonzero. We can assume that  $a_i \in P$  and that there exists the number  $i$  for which the free term of the polynomial  $a_i$  is not equal to zero (here we use that  $N$  is free of  $t$ -torsion). Reducing this equality modulo  $t$  we obtain the contradiction.

2) By the minimality criterion (see the theorem 4.3) it

suffices to prove that for a homomorphism  $\varphi : F_1 \rightarrow F_2$  where  $F_1, F_2$  are finitely generated free  $K$ -modules the pair  $(\text{Im } \varphi, F_2)$  is strongly irreducible if and only if the pair  $(\text{Im } \varphi_t, F_2/tF_2)$  is strongly irreducible (where  $\varphi_t : F_1/tF_1 \rightarrow F_2/tF_2$  denotes the homomorphism  $\varphi$  reduced modulo  $t$ ).

Suppose that the pair  $(\text{Im } \varphi, F_2)$  is irreducible. We'll show that the pair  $(\text{Im } \varphi_t, F_2/tF_2)$  is irreducible. (The inverse implication is obvious.) Indeed suppose that opposite is true, i.e. there exist elements  $(e_1, \dots, e_k, f_1, \dots, f_n)$ , such that  $f_j \in \text{Im } \varphi$  and the images  $\bar{e}_1, \bar{f}_j$  in  $F_2/tF_2$  form the basis of this module. By p. 1) the elements  $e_1, \dots, e_k, f_1, \dots, f_n$  form the basis of  $F_2$ ; we get the contradiction.

The assertion concerning the strong irreducibility is proved on the same lines.

#### §6. Novikov type inequalities

The simplest way of proving (0.3) is the following. Let  $\omega$  be a maximally irrational form. We can find a rational form  $\omega'$ , belonging to a very small neighbourhood of  $\omega$ , such that  $m_p(\omega') = m_p(\omega)$ . Now we apply lemma 2.6 and reduce the problem to proving Novikov inequalities (0.1); the latter are treated in [1], [2], see also [4].

We produce here one more proof of the Novikov type inequalities, which in our opinion clarifies the general homological reasons for these inequalities to arise.

Recall first from Morse theory some well-known facts. Let  $(W; V_0, V_1)$  be a compact manifold with boundary  $\partial W = V_0 \sqcup V_1$ ,  $f$  - a Morse function on  $W$ , constant on  $V_0$  and on  $V_1$ . The func-

tion  $f$  gives rise to a relative cell complex  $(K, V_0)$ , homotopy equivalent to  $(W, V_0)$ . The number of relative  $p$ -cells of  $(K, V_0)$  is equal to  $m_p(f)$ . Consider any regular covering  $\bar{K}$  over  $K$  with a structure group  $G$  and denote by  $\bar{V}_0$  the preimage of  $V_0$  in  $K$ . The chain complex of the relative cell complex  $(K, V_0)$  is a free  $\mathbb{Z}[G]$ -complex with  $m_p(f)$  free  $\mathbb{Z}[G]$ -generators in dimension  $p$ . Any homotopy equivalence  $\varphi: (K, V_0) \rightarrow (W, V_0)$  (which we assume to be cellular) induces an isomorphism of  $\mathbb{Z}[G]$ -modules  $H_*(\bar{K}, \bar{V}_0) \rightarrow H_*(\bar{W}, \bar{V}_0)$ . This implies that there exists a free finitely generated  $\mathbb{Z}[G]$ -complex  $C'_* \sim C_*(\bar{W}, \bar{V}_0)$ , such that  $\mu(C'_p) = m_p(f)$ .

Consider now an arbitrary rational Morse 1-form  $\omega$  on  $M$  and set  $[\omega] = \lambda$ .

We may assume (having multiplied  $\omega$  by a constant) that  $\omega = df$ , where  $f$  is a map of  $M$  to  $S^1$ .

Consider the cyclic covering  $\bar{M} \rightarrow M$ , on which  $\omega$  becomes exact:  $\omega = d\bar{f}$ ,  $\bar{f}: \bar{M} \rightarrow \mathbb{R}$ . Denote  $V$  the preimage in  $M$  of any regular value  $c$  of the map  $f$  (or, equivalently, the preimage in  $M$  of the corresponding regular value of  $\bar{f}: \bar{M} \rightarrow \mathbb{R}$ ).

Denote  $t$  the generator of the deck transformation group of  $M$ ; set  $V^- = \{x \in \bar{M} \mid \bar{f}(x) \leq c\}$ ,  $V^+ = \{x \in \bar{M} \mid \bar{f}(x) \geq c\}$ . We can assume that  $tV^- \subset V^-$ ; set  $W = tV^+ \cap V^-$ . The manifold  $M$  is the countable union of "bricks"  $t^k W$ ,  $k \in \mathbb{Z}$ . There is a Morse function  $f$  on the cobordism  $(W; V, tV)$ .

Consider now the regular covering  $\hat{M} \xrightarrow{p} M$  with a structure group

$$H_1(M, \mathbb{Z}) / \text{Tors } H_1(M, \mathbb{Z}) = \mathbb{Z}^m$$

It can be factored through the cyclic covering:  $\hat{M} \rightarrow \bar{M} \rightarrow M$ .  
 The composition  $\hat{M} \xrightarrow{p} \bar{M} \xrightarrow{f} R$  will be denoted by  $\hat{f}$ , the  $p$ -preimages of  $w, v, v^+, v^-$  - by  $\hat{w}, \hat{v}, \hat{v}^+, \hat{v}^-$  correspondingly. Choose now a triangulation of  $W$  and extend it to a triangulation of  $M$ . Then  $v^+, v^-, \hat{v}^+, \hat{v}^-$  also get the triangulations. The chain complex  $C_*(\hat{V}^-)$  is a free finitely generated  $P$ -complex (recall from (5.1) that  $R = \mathbb{Z}[\mathbb{Z}^{m-1}]$ ,  $P = R[t]$ ); the  $R$ -complex  $C_*(\hat{V}^-)/tC_*(\hat{V}^-)$  is the same as  $C_*(\hat{W}, t\hat{V})$ . The manifold  $W$  is a covering of  $W$  and by the above we get the following:

There exists a free finitely generated  $P$ -complex  $C_* = C_*(\hat{V}^-)$ , such that the  $R$ -complex  $C_*/tC_*$  is homotopy equivalent to an  $R$ -complex  $D_*$ , having exactly  $m_p(f)$  generators of dimension  $p$ .

Consider now the complex  $S^{-1}C_*$  (we use the notations (5.1)). It is free finitely generated  $K$ -complex and since  $K$  is an IBN,  $s$ -ring (by lemma 5.5), we can find a minimal  $K$ -complex  $C_*^0$  in the homotopy type of  $S^{-1}C_*$  (see § 4). By the same lemma the complex  $C_*^0/tC_*^0$  is minimal in the homotopy type of

$$\begin{aligned} S^{-1}C_*(\hat{V}^-)/tS^{-1}C_*(\hat{V}^-) &\cong C_*(\hat{V}^-)/tC_*(\hat{V}^-) \cong \\ &\cong C_*(\hat{V}^-, t\hat{V}^-) \cong C_*(\hat{W}, t\hat{V}) . \end{aligned}$$

Note that  $m_p(\omega) = \mu(D_p)$ . The  $R$ -complex  $D_*$  is homotopy equivalent to  $C_*(\hat{W}, t\hat{V})$ ; this homotopy type contains a minimal complex  $C_*^0/tC_*^0$ , having exactly  $\mu(C_p^0)$  generators in each dimension  $p$ .

Note further that the localized complex  $S'^{-1}C_*^0$  (recall that  $S'^{-1}$  denote the localization with respect to  $t$ ) is homotopy equivalent to

$$S'^{-1} S^{-1} C_* = S^{-1} S'^{-1} C_* = S^{-1} C_*(\hat{M})$$

Thus we have constructed a free finitely generated K-complex  $C_*^0$ , such that

$$S'^{-1} C_*^0 \sim S^{-1} C_*(\hat{M}) \tag{6.1}$$

$$m_p(\omega) \geq \mu_K(C_*^0).$$

This implies of course

$$m_p(\omega) \geq \mu_\Gamma(C_*^1) \tag{6.2}$$

for some free finitely generated  $\Gamma$ -complex  $C_*^1$ , homotopy equivalent to  $S^{-1} C_*(\hat{M})$ .

Conjecture. The ring  $\Gamma = S'^{-1} S^{-1} P$  is an s-ring, i.e. any finitely generated stably free module is free<sup>3)</sup>.

If this conjecture holds, then the homotopy type of any finitely generated  $\Gamma$ -complex  $N_*$  contains a minimal complex, numbers of generators of which are the invariants of homotopy type of  $N_*$  (see § 4). Denote these numbers by  $m_p(N_*)$ . Then we get

$$m_p(\omega) \geq m_p(S_{[\omega]}^{-1} C_*(\hat{M})).$$

Next we deduce (0.3) from (6.2). Let  $\lambda$  be a rational cohomology class,  $\lambda \in H^1(M, \mathbb{Q})$ ,  $C_*^1$  be a free finitely generated

3) The validity of s-property was analyzed for the rings of the similar type, see [18]. However for  $\Gamma$  itself the conjecture seems not settled.

to be



$S^{-1}S^{-1}P$ -complex, homotopy equivalent to  $S_{\lambda}^{-1}C_{*}(\hat{M})$ .

Note that  $C_{*}^1$  is an  $S_{\lambda}$ -localization of some free finitely generated  $\Lambda$ -complex  $C_{*}^2$ . The  $\Lambda$ -complexes  $C_{*}(\hat{M})$  and  $C_{*}^2$  become homotopy equivalent already after localization with respect to some element  $\sigma \in S_{\lambda}$  (see corollary 2.5). For any maximally irrational class  $\lambda' \in H^1(M, \mathbb{R})$ , sufficiently close to  $\lambda$ , the element  $\sigma$  belongs also to  $S_{\lambda'}$ , hence

$$S_{\lambda'}^{-1}C_{*}(\hat{M}) \sim S_{\lambda'}^{-1}C_{*}^2.$$

The ring  $S_{\lambda'}^{-1}\Lambda$  is a principal ideal domain (see theorem 1.4), therefore the number of generators of  $C_{*}^2$  in dimension  $p$  is not greater than

$$b_{\cdot}(S_{\lambda'}^{-1}H_p(\hat{M})) + q(S_{\lambda'}^{-1}H_p(\hat{M})) + q(S_{\lambda'}^{-1}H_{p-1}(\hat{M})),$$

and, having recollected (6.2), we obtain (0.3).

### §7. Reduction to a surgery problem

Let  $\gamma$  be a maximally irrational cohomology class. Then  $\Lambda_{(\gamma)}$  is a principal ideal domain (theorem 1.1), hence

$$S_{\gamma}^{-1}H_p(\tilde{M}, \mathbb{Z}) \cong H_p(M, \Lambda_{(\gamma)}) = \left( \bigoplus_{i=1}^{b_p(\gamma)} \Lambda_{(\gamma)} \right) \oplus \left( \bigoplus_{j=1}^{q_p(\gamma)} \Lambda_{(\gamma)} / a_j^{(p)} \Lambda_{(\gamma)} \right).$$

We may assume that  $a_j^{(p)} \in \Lambda$ , thus we get

$$S_{\gamma}^{-1}H_p(\tilde{M}, \mathbb{Z}) \cong S_{\gamma}^{-1} \left[ \left( \bigoplus_{i=1}^{b_p(\gamma)} \Lambda \right) \oplus \left( \bigoplus_{j=1}^{q_p(\gamma)} \Lambda / a_j^{(p)} \Lambda \right) \right].$$

Together with lemma 2.4 this implies that modules

$H_p(\tilde{M}, \mathbb{Z})$  and

$$\left( \bigoplus_{i=1}^{b_p(\gamma)} \Lambda \right) \oplus \left( \bigoplus_{j=1}^{q_p(\gamma)} \Lambda / a_j^{(p)} \Lambda \right)$$

become isomorphic after localization with respect to multiplicative set, generated by some element  $\sigma \in S_\lambda$ . The  $\sigma$  belongs also to the set  $S_\lambda$  for every  $\lambda$  (in particular, rational), sufficiently close to  $\gamma$ .

Thus for the elements  $\lambda$  of some open dense set  $U \subset H^1(M, \mathbb{R}) = \mathbb{R}^m$  we get

$$S_\lambda^{-1} H_p(\tilde{M}, \mathbb{Z}) \cong \left( \bigoplus_{i=1}^{b_p} S_\lambda^{-1} \Lambda \right) + \left( \bigoplus_{j=1}^{q_p} S_\lambda^{-1} \Lambda / a_j^{(p)} S_\lambda^{-1} \Lambda \right) \quad (7.1)$$

The number  $q_p$  depends of course on  $\lambda$ ; in the complement in  $H^1(M, \mathbb{R})$  to  $\Gamma_i$  the number  $q_p$  equals  $q_p(\gamma)$  (see introduction). To obtain the proof of theorem 0.1 it suffices to prove the following

**Theorem 7.1.** Let  $\pi_1 M^n = \mathbb{Z}^m$ ,  $n \geq 6$  and suppose that  $\lambda$  is a rational cohomology class,  $\lambda \in H^1(M, \mathbb{Q})$ , such that (7.1) holds and  $m_r = m_{r+1} = m_{r+2} = 0$  for some  $r : 2 \leq r \leq n-4$ . Then there exists a Morse map  $f : M^n \rightarrow S^1$ , inducing from the fundamental class  $i$  of circle the element, which is a multiple of  $\lambda$ , such that  $m_p(f) = m_p = b_p + q_p + q_{p-1}$  (for  $0 \leq p \leq n$ ).

It is sufficient to prove this theorem under the additional assumption that  $\lambda$  is an integer cohomology class which is defined by the projection of  $H_1(M, \mathbb{Z}) = \mathbb{Z}^m$  onto the first direct summand  $\mathbb{Z}$ . In the course of this and two subsequent sections we suppose that this condition holds, without stating this

any more.

Let  $V^{n-1} \subset M^n$  be a connected smooth submanifold of codimension 1,  $\nu$  be a normal vector field on  $V$ . The pair  $(V^{n-1}, \nu)$  is called an admissible splitting <sup>4)</sup> if

1)  $\pi_1(V^{n-1}) \rightarrow \pi_1(M^n)$  is a monomorphism onto the subgroup  $\text{Ker } \lambda \approx \mathbb{Z}^{m-1}$ ;


2) the Pontryagin-Thom construction with respect to  $\nu$  determines a map  $M^n \rightarrow S^1$ , representing the class  $\lambda \in H^1(M, \mathbb{Z})$ .

The existence of admissible splittings is proved in [10] under the assumption that the homotopy fiber of  $\lambda : M \rightarrow S^1$  has a finite type (one can show that for  $\pi_1 M = \mathbb{Z}^m$  this is equivalent to the following: the class  $\lambda$  satisfies (7.1) with  $b_p = q_p = 0$  for all  $p$ ).

The same proof is valid for arbitrary cohomology classes. (Note also that the results of [19] imply that for any Morse map  $f : M^n \rightarrow S^1$ , which represents  $\lambda$  and has no critical points of indices 0,  $n$  the level surfaces  $f^{-1}(c)$  are connected and  $\pi_1(f^{-1}(c)) \rightarrow \text{Ker } \lambda$  is an epimorphism.)

Consider the infinite cyclic covering  $p : \hat{M}^n \rightarrow M^n$ , corresponding to  $\lambda$ . For an admissible splitting  $V^{n-1}$  the pre-image  $p^{-1}(V^{n-1})$  consists of countably infinite number of copies of  $V^{n-1}$ , which divide  $\hat{M}^n$  into countably infinite number of "bricks"  $W^n$ ;  $\partial W^n \approx V^{n-1} \cup tV^{n-1}$  (where  $t$  is a generator of the structure group of the covering), see the picture below. Now fix any copy of  $V^{n-1} \subset \hat{M}^n$ , it divides  $\hat{M}^n$  into two parts:

<sup>4)</sup> We will omit  $\nu$  in the notations if no confusion is possible.



$V^+$  and  $V^-$ . (We assume here and elsewhere that for every admissible splitting  $V$  the lifting of  $V$  into  $\hat{M}^n$  is chosen and fixed. In this case the notations  $V^+$ ,  $V^-$  etc. make sense.) The intersection  $V^+ \cap V^-$  equals  $V^{n-1}$ , the vector  $\nu$  points into  $V^+$  and  $tV^- \subset V^-$ .

Furthermore,

$$\pi_1(W^n) \approx \pi_1(V^{n-1}) \approx \pi_1(\hat{M}^n) \approx \mathbb{Z}^{m-1}$$

the universal covering  $\tilde{M}^n$  is divided by  $\tilde{V}^{n-1}$  into two parts:  $\tilde{V}^+$  and  $\tilde{V}^-$  (all this can be found in [10]).

One easily sees that any triangulation of  $M$ , such that  $V^{n-1}$  is a subcomplex, determines a free finitely generated  $P$ -complex  $C_*(\tilde{V}^-)$  (in the notations (5.1)); the complexes  $S^{-1}C_*(\tilde{V}^-)$  and  $C_*(\tilde{M}^n)$  coincide, and, therefore

$$H_*(\tilde{M}^n) \approx S^{-1}H_*(\tilde{V}^-)$$

The factorcomplex  $C_*(\tilde{V}^-)/tC_*(\tilde{V}^-)$  is the chain complex of the triangulation of the pair  $(\tilde{W}^n, t\tilde{V}^{n-1})$ .

Suppose now that  $\lambda$  satisfies (7.1) for  $p \leq k$ , and that

$$a_j^{(p)} = a_{j,0}^{(p)} + a_{j,1}^{(p)} t + \dots$$

where  $a_{j,0}^{(p)}$  are nonzero, noninvertible elements of  $R$ . An admissible splitting  $V$  will be called  $k$ -regular if for  $p \leq k$

$$S^{-1}H_p(\tilde{V}^-, \mathbb{Z}) \approx \left( \bigoplus_{i=1}^{b_p} S^{-1}P \right) \oplus \left( \bigoplus_{j=1}^{q_p} S^{-1}P / a_j^{(p)} S^{-1}P \right) \quad (7.2)$$

Lemma 7.2. Suppose that  $\pi_1 M^n = \mathbb{Z}^m$ ,  $n \geq 6$ ,  $\lambda \in H^1(M, \mathbb{Z})$  satisfies (7.1) for  $p \leq k$ , where  $k \leq n-4$ . Let  $V$  be an admis-

sible splitting. Then there exists a  $k$ -regular splitting  $V_0$ , obtained from  $V$  by surgical modifications of indices  $\leq k+1$ , made inside  $V^+ \subset \widehat{M}^n$ .

The proof of the lemma will be given in §8 (the main ideas were outlined in [6]).

Now we'll deduce from this lemma the existence of a Morse form in a class  $\gamma$  with the required number of zeros of indices  $k$ .

First of all note that given an admissible splitting  $V$  and a Morse function  $f$  on the cobordism  $(W; V, tV)$ , we can produce from this data a Morse map  $\tilde{f} : M \rightarrow S^1$ , belonging to  $\lambda$  (for this purpose we change  $f$  in the small neighbourhood of the boundary  $V^{n-1} \cup tV^{n-1}$  so that it becomes a projection on the second factor of the collar:  $V^{n-1} \times [0, \varepsilon] \rightarrow [0, \varepsilon]$ , afterwards we glue together  $V$  and  $tV$ ). The map  $\tilde{f}$  has the same critical points as  $f$ .

So we proceed as to construct a Morse function with the required Morse numbers on the cobordism  $(W_0; V_0, tV_0)$ , where  $W_0$  is the part of  $\widehat{M}$ , lying between  $V_0$  and  $tV_0$ .

First calculate the homology  $H_p(\tilde{W}_0, t\tilde{V}_0; \mathbb{Z})$ :

$$H_p(\tilde{W}_0, t\tilde{V}_0) \approx H_p(\tilde{V}_0^-, t\tilde{V}_0^-) \approx H_p(\tilde{V}_0^-) / tH_p(\tilde{V}_0^-)$$

(First identification is due to excision axiom. To get the second we observe that the map  $H_p(t\tilde{V}_0^-) \rightarrow H_p(\tilde{V}_0^-)$  of the exact sequence of the pair  $(V_0, tV_0)$  coincides with multiplication by  $t$  and the latter is injective since the module  $B_p = H_p(\tilde{V}_0^-)$  is free of  $t$ -torsion.)

Furthermore

$$H_p(\tilde{V}_0^-) / tH_p(\tilde{V}_0^-) \approx B_p / tB_p \approx \left( \bigoplus_{i=1}^{b_p} R \right) \oplus \left( \bigoplus_{j=1}^{q_p} R/a_{j,0}^{(p)} R \right), \quad p \leq k$$

(the last identification follows from lemma 5.3). Besides that,  $b_0 = q_0 = 0$  (remark 2.7) and  $b_1 = q_1 = 0$  since  $M$  is simply-connected; hence from corollary 3.2 we conclude that  $b_n = q_n = b_{n-1} = q_{n-1} = 0$ . Thus the homology modules  $H_*(\tilde{W}_0, t\tilde{V}_0)$  vanish outside the dimensions  $2 \leq * \leq n-2$  and are principal  $\mathbb{Z}[\pi_1(W_0)]$ -modules for  $p \leq k$ . According to lemma 5.1 we can find a complex  $C_* = \{0 \leftarrow C_2 \leftarrow \dots \leftarrow C_{n-2} \leftarrow 0\}$  which is homotopy equivalent to  $C_*(\tilde{W}_0, t\tilde{V}_0)$ , and has  $b_i + q_i + q_{i-1}$  free generators in dimension  $i$  (where  $2 \leq i \leq k \leq n-4$ ).

According to the theorem 4.1 and the remark following it we can realize  $C_*$  as the Morse complex of some Morse function  $f$  on the cobordism  $(W_0, tV_0)$ . This function gives rise to a 1-form, which satisfy the requirements. Thus we have proved the following assertion.

Theorem 7.3 (see [6])<sup>5)</sup>. Suppose that  $\pi_1 M^n = \mathbb{Z}^m$ ,  $n \geq 6$ . Then any element  $\gamma$  of some dense open conical set  $U \subset H^1(M, \mathbb{R})$  can be realized by a Morse form  $\omega \in \Omega^1(M)$ , which has the least possible number of zeros of indices  $p$  where  $0 \leq p \leq n-4$  among all the forms of the class  $\gamma$ . This number equals  $b_p + q_p(\bar{\gamma}) + q_{p-1}(\bar{\gamma})$ , where  $\bar{\gamma}$  is any maximally irrational cohomology class sufficiently close to  $\gamma$ .

Now we'll use duality and do surgery also "from another end".

---

5) The assumptions here are weaker than in theorem 0.1, but the result is concerned only with the indices  $p$ :  $0 \leq p \leq n-4$ .

In addition to (5.1) we introduce some new notations:

$$\begin{aligned} \bar{P} &= R[t^{-1}], \quad \bar{S} = \{1 + t^{-1} Q(t^{-1}) \mid Q(t^{-1}) \in R[t^{-1}]\}, \\ \bar{K} &= \bar{S}^{-1} \bar{P}, \quad \bar{S}' = \{t^{-n} \mid n \in \mathbb{N}\}, \quad \bar{\Gamma} = \bar{S}'^{-1} \bar{S}^{-1} \bar{P} = S_{(-\lambda)}^{-1} \Lambda. \end{aligned}$$

Note that if  $(V, \nu)$  is an admissible splitting with respect to  $\lambda$ , then  $(V, -\nu)$  is an admissible splitting with respect to  $(-\lambda)$  and all the results above hold for  $(-\lambda)$  (with  $t$  replaced by  $t^{-1}$ ). Note that  $(V, \nu)^- = (V, -\nu)^+$ .

If the class  $\lambda$  satisfies (7.1) for  $p \leq k$  then according to (3.4) the following holds:

$$\begin{aligned} S_{(-\lambda)}^{-1} H_s(\tilde{M}^n, \mathbb{Z}) &\cong \left( \bigoplus_{i=1}^{b_s} S_{(-\lambda)}^{-1} \Lambda \right) \oplus \\ &\oplus \left( \bigoplus_{j=1}^{q_{n-s-1}} (S_{(-\lambda)}^{-1} \Lambda) / (\chi(a_j^{(n-s-1)}) S_{(-\lambda)}^{-1} \Lambda) \right) \end{aligned}$$

for  $s \geq n-k$  (recall that  $b_s = b_{n-s}$ ).

Suppose now that  $1 \leq r \leq n-4$  and  $(V, \nu)$  is an admissible splitting. Applying lemma 7.2 (where  $k = r$ ) we get an  $r$ -regular splitting  $(V_0, \nu_0)$ . Applying lemma 7.2 to the splitting  $(V_0, -\nu_0)$ , cohomology class  $(-\lambda)$  and  $k = n - r - 3$  we get an  $(n - r - 3)$ -regular splitting, say  $(V_1, -\nu_1)$ , corresponding to  $(-\lambda)$ . Note that  $V_1$  is obtained from  $V_0$  by a sequence of surgical modifications of indices  $\leq n-r-2$ , consequently, the homology of  $(V_1, \nu_1)^-$  coincide with that of  $(V_0, \nu_0)^-$  in dimensions  $r$ . Hence  $(V_1, \nu_1)$  is also an  $r$ -regular splitting.

An admissible splitting  $(V, \nu)$  is called  $r$ -biregular if  $(V, \nu)$  is  $r$ -regular with respect to  $\lambda$  and  $(V, -\nu)$  is  $(n-r-3)$ -regular with respect to  $(-\lambda)$ .



We have proved that  $r$ -biregular splittings exist for  $1 \leq r \leq n-4$ .

Remarks. 1. In §9 we'll deduce from the above that under the assumptions of theorem 7.3 every element  $\gamma$  of some dense open conical  $U \subset H^1(M, \mathbb{R})$  can be realized by a Morse form  $\omega$  having a minimal Morse numbers of all indices except two adjacent ones, say  $r, r+1$  where  $1 \leq r \leq n-4$ .

2. Here we'll show that any  $(n-3)$ -regular splitting  $V$  is also  $n$ -regular. Consider the free finitely generated complex  $S^{-1}C_*(\tilde{V}^-)$ . The ring  $K = S^{-1}P$  is an IBN,  $s$ -ring (lemma 5.5), hence the homotopy type of this complex contains a minimal free finitely generated  $K$ -complex  $C_*^0$ . Note that  $C_0^0 = C_1^0 = C_{n-1}^0 = C_n^0 = 0$ . Indeed, the complex  $C_*^0/tC_*^0$  is  $R$ -minimal (see lemma 5.5) and belongs to the homotopy type of

$$S^{-1}C_*(\tilde{V}^-)/t S^{-1}C_*(\tilde{V}^-) = C_*(\tilde{V}^-, t\tilde{V}^-)$$

There exists a Morse function  $f$  on the cobordism  $(W; V, tV)$ , which has no critical points of indices  $0, 1, n-1, n$ . The corresponding complex  $C_*(f)$  has no generators in dimensions  $0, 1, n-1, n$  and since  $C_*^0/tC_*^0$  is minimal and  $C_*^0/tC_*^0 \sim C_*(f) \sim C_*(\tilde{W}, t\tilde{V})$  we get  $C_i^0/tC_i^0 = 0$  for  $i = 0, 1, n-1, n$ . Therefore,  $C_i^0 = 0$  for  $i = 0, 1, n-1, n$ .

The homology modules  $H_s(C_*^0)$  have the resolutions of length two for  $s \leq n-3$ . Therefore there exists a complex  $C_*^1$ , in the homotopy type of  $C_*^0$ , which is standard in dimensions  $* \leq n-3$  and moreover  $C_n^1 = 0, C_{n-1}^1 = B \oplus Z$ , where  $\partial|B$  is injective  $\partial|Z = 0$ . Therefore

$$Z \approx H_{n-2}(C_*^1) \approx H_{n-2}(S^{-1}C_*(\tilde{V}^-)), \mu(Z) = b_{n-2}$$

Now we realize  $C_*^1/tC_*^1$  by a Morse function and get our assertion.

For  $\pi_1 M^n = \mathbb{Z}$  the  $(n-3)$ -regular splittings always exist (see [4]). The author does not know if the same holds for  $\pi_1 M = \mathbb{Z}^m$ ,  $m > 1$  and any  $\lambda$ .

To cope with the dimensions left we are to impose the restriction  $m_r = m_{r+1} = m_{r+2} = 0$ , where  $m_s = b_s + q_s + q_{s-1}$ . This restriction is an analogue (for this three particular dimensions) of the Farrell condition for existence of a map  $f : M^n \rightarrow S^1$ , which realizes a given cohomology class and has no critical points at all.

Lemma 7.4. Suppose  $\pi_1 M^n = \mathbb{Z}^m$ ,  $n \geq 6$ , the integer cohomology class  $\lambda$  satisfies (7.1) and  $(V, \nu)$  is an  $(r-1)$ -biregular splitting (here  $1 \leq r \leq n-2$ ).

Suppose further that  $m_r = m_{r+1} = 0$ .

Then 1)  $H_r(\tilde{M}^n, \tilde{V}^-) = 0$

2)  $H_{r+1}(\tilde{M}^n, \tilde{V}^-)$  is a free finitely generated  $R$ -module and  $t$  is its nilpotent endomorphism.

Thus the module  $Q = H_{r+1}(\tilde{M}^n, \tilde{V}^-)$  together with the endomorphism  $t$  is an object of the category  $\mathcal{G}(\mathbb{Z}[\mathbb{Z}^{m-1}], \text{id})$  which consists of free finitely generated  $\mathbb{Z}[\mathbb{Z}^{m-1}]$ -modules and their nilpotent endomorphisms (see [10]). The corresponding Grothendieck group vanishes (see [10]), therefore our pair  $(Q, t)$  is equal to zero modulo relations in this group. To realize this equivalence in geometric setting we need the third "critical point free" dimension.

Lemma 7.5. Suppose that the assumptions of lemma 7.4 hold, and, moreover,  $m_{r+2} = 0$ ,  $2 \leq r \leq n-4$ .

Then there exists an  $(r-1)$ -biregular splitting  $(V, \nu)$ , such that

$$H_r(\tilde{M}, \tilde{V}^-) = 0, \quad H_{r+1}(\tilde{M}, \tilde{V}^-) = 0.$$

The proofs of lemmas 7.4 and 7.5 will be presented in §9. Now we'll deduce from them the theorem 7.1. Let  $(V, \nu)$  be an admissible splitting constructed in lemma 7.5,  $(W; V, tV)$  be a corresponding brick in  $\hat{M}$ . Compute the homology  $H_*(\tilde{W}, t\tilde{V})$ . For  $p \leq r-1$

$$H_p(\tilde{W}, t\tilde{V}) \approx \left( \bigoplus_{i=1}^{b_p} R \right) \oplus \left( \bigoplus_{j=1}^{q_p} R / a_{j,0}^{(p)} R \right)$$

(see 7.2). Since  $(V, -\nu)$  is  $(n-r-2)$ -regular,

$$H_s(\tilde{W}, \tilde{V}) \approx \left( \bigoplus_{i=1}^{b_s} R \right) \oplus \left( \bigoplus_{j=1}^{q_{n-s-1}} R / \chi(a_{j,0}^{(n-s-1)}) R \right),$$

for  $s \leq n-r-2$ .

Apply now the Poincare duality to the manifold  $W$  with two components  $V, tV$  of the boundary.

The  $R$ -modules  $H_s(\tilde{W}, \tilde{V})$  are principal for  $s \leq n-r-2$  and it is easy to calculate cohomology  $H^s(\tilde{W}, \tilde{V})$  for  $s \leq n-r-2$ .

Applying the Poincare duality arguments from §3 we get

$$H_p(\tilde{W}, t\tilde{V}) \approx \left( \bigoplus_{i=1}^{b_p} R \right) \oplus \left( \bigoplus_{j=1}^{q_p} R / a_{j,0}^{(p)} R \right), \quad p \geq r+2$$

Since  $m_{r+2} = 0$ ,

$$H_{r+2}(\tilde{W}, t\tilde{V}) = H_{r+2}(\tilde{V}^-, t\tilde{V}^-) = 0$$

From the exact sequence of the triple  $(t^{-1}\tilde{V}^-, \tilde{V}^-, t\tilde{V}^-)$  we obtain  $H_{r+2}(t^{-1}\tilde{V}^-, t\tilde{V}^-) = 0$  and (by induction)  $H_{r+2}(t^{-n}\tilde{V}^-, t\tilde{V}^-) = 0$ .

Passing to direct limit in  $n \rightarrow \infty$  we get  $H_{r+2}(\tilde{M}, t\tilde{V}^-) = 0$ .

From the exact sequence of the triple  $(\tilde{M}, \tilde{V}^-, t\tilde{V}^-)$  (using  $H_*(\tilde{M}, \tilde{V}^-) = H_*(\tilde{M}, t\tilde{V}^-) = 0$  for  $* = r, r+1, r+2$ ) we get

$$H_{r+1}(\tilde{V}^-, t\tilde{V}^-) = H_r(\tilde{V}^-, t\tilde{V}^-) = 0$$

Now we collect together results of our computations and see that  $H_p(\tilde{W}, t\tilde{V})$  is isomorphic to  $(\bigoplus_{i=1}^{b_p} R) \oplus (\bigoplus_{j=1}^{q_p} R/a_{j,0}^{(p)} R)$  for all  $p$ .

Now we apply the same argument as in proof of theorem 7.3 and the theorem 7.1 is proven.

### 8. Construction of p-regular splitting

This section is devoted to the proof of lemma 7.2 We prove it by induction in  $k$ .

Note first that every admissible splitting  $V$  is 1-regular. Indeed, since  $\tilde{V}^-$  is connected, the map  $t : \tilde{V}^- \rightarrow \tilde{V}^-$  induces the identity homomorphism in the group  $H_0(\tilde{V}^-)$ , hence  $S^{-1}H_0(\tilde{V}^-) = 0$ . The same argument proves  $b_0 = q_0 = 0$ . Furthermore, the commutativity of  $\pi_1 M^n$  implies that  $t$  induces the identity homomorphism also in the group  $H_1(\tilde{V}^-)$  (which is isomorphic to  $\pi_1(\tilde{V}^-)$ ), therefore  $S^{-1}H_1(\tilde{V}^-) = 0$ .

The induction step will be produced by means of lemma 8.1 and lemma 8.2. In both lemmas we assume that  $\pi_1 M^n = \mathbb{Z}^m$ ,  $n \geq 6$  and that  $\lambda$  satisfies (7.1) for  $p \leq k$ .

Lemma 8.1. Let  $k \leq n-4$  and suppose that an admissible

splitting  $V$  is  $(k-1)$ -regular (i.e.  $V$  satisfies (7.2) for  $p \leq k-1$ ). Then there exists an admissible splitting  $V_0$ , obtained from  $V$  by a sequence of surgical modifications inside  $V^+$  of indices  $\leq k+1$ , such that

- 1) the  $P$ -modules  $H_p(\tilde{V}_0^-)$  and  $H_p(\tilde{V}^-)$  are isomorphic for  $p \leq k-1$ ,
- 2) the  $P$ -module  $H_k(\tilde{V}_0^-)$  is isomorphic to  $H_k(\tilde{V}^-)/\text{Tor}_t H_k(\tilde{V}^-)$  (recall that  $\text{Tor}_t M$  denotes the submodule of all elements of  $M$ , annihilated by some power of  $t$  and  ${}^t M$  denotes the submodule of elements, annihilated by  $t$ ).

Lemma 8.2. Let  $k \leq n-3$  and suppose that  $V$  is a  $(k-1)$ -regular splitting, such that  $A = H_k(\tilde{V}^-)$  is free of  $t$ -torsion. Suppose that  $B$  is a  $P$ -submodule of  $A$ , such that  $tA \subset B \subset A$  and  $B/tA$  is a cyclic  $R$ -module.

Then there exists an admissible splitting  $V_0$ , obtained from  $V$  by a sequence of surgical modifications inside  $V^+$  of indices  $\leq k+1$ , such that

- 1) the  $P$ -modules  $H_p(\tilde{V}_0^-)$  and  $H_p(\tilde{V}^-)$  are isomorphic for  $p \leq k-1$ ,
- 2) there is an epimorphism of  $P$ -modules  $H_k(\tilde{V}_0^-) \rightarrow B$  with the kernel  ${}^t H_k(\tilde{V}_0^-) = \text{Tor}_t H_k(\tilde{V}_0^-)$ .

Both lemma proved, the induction step is made as follows. Let  $V$  be a  $(k-1)$ -regular splitting,  $k \leq n-4$ . Having applied lemma 8.1 we may assume that  $P$ -module  $H_k(\tilde{V}^-)$  have no  $t$ -torsion. Consider the  $P$ -module  $A_k = H_k(\tilde{V}^-)$ . The condition (7.1) holds for  $p \leq k$ , hence by lemma 5.3 there exists a submodule  $B_k \subset A_k$  such that

$$\begin{cases} t^r A_k \subset B_k \subset A_k & \text{for some } r \\ S^{-1} B_k \approx \left( \bigoplus_{i=1}^r S^{-1} P \right) \oplus \left( \bigoplus_{j=1}^{q_k} S^{-1} P / a_j^{(k)} S^{-1} P \right) \end{cases} \quad (8.1)$$

Choose the filtration of  $B_k$

$$t^r A_k = B_k^{(0)} \subset B_k^{(1)} \subset \dots \subset B_k^{(m)} = B_k$$

such that  $B_k^{(i)}$  are  $P$ -submodules in  $B_k$ , and factormodules  $B_k^{(i)}/B_k^{(i-1)}$  are cyclic  $R$ -modules and  $tB_k^{(i)} \subset B_k^{(i-1)}$ .

Since  $A_k$  has no  $t$ -torsion, the module  $B_k^{(1)}$  is contained in  $t^{r-1} A_k$ . Applying lemma 8.2 to the admissible splitting  $t^{r-1} V$  and the modules  $t^r A_k \subset B_k^{(1)} \subset t^{r-1} A_k$  and killing afterwards the  $t$ -torsion in the resulting homology  $H_k(\tilde{V}_0^-)$  (by means of lemma 8.1) we get an admissible splitting  $V_1$ , such that  $H_s(\tilde{V}_1^-) \approx H_s(\tilde{V}^-)$  for  $s \leq k-1$  and  $H_k(\tilde{V}_1^-) \approx B_k^{(1)}$ . Applying, further, the lemma 8.2 to the manifold  $V_1$  and the modules  $tB_k^{(1)} \subset tB_k^{(2)} \subset B_k^{(1)}$  (and killing afterwards the  $t$ -torsion by means of lemma 8.1) we get an admissible splitting  $V_2$ , such that  $H_s(\tilde{V}_2^-) \approx H_s(\tilde{V}_1^-)$  for  $s \leq k-1$  and  $H_k(\tilde{V}_2^-) \approx tB_k^{(2)}$ . Since the  $P$ -module  $A_k$  has no  $t$ -torsion,  $tB_k^{(2)} \approx B_k^{(2)}$ . We proceed further in a similar way and in  $m$  steps get an admissible splitting  $V_k$ , satisfying the conclusion of lemma 7.2.

The proof of lemma 8.1.

0. First of all we note that since  $P$  is noetherian and  $\text{Tor}_t H_k(\tilde{V}^-)$  is a finitely generated module, it suffices to construct for any  $\alpha \in {}^t H_k(\tilde{V}^-)$  a new admissible splitting  $V_0$ , satisfying the requirement 1) and the requirement

$$H_k(\tilde{V}_0^-) \approx H_k(\tilde{V}^-) / (\alpha)$$

Indeed, suppose that  $t^N \text{Tor}_{tH_k}(\tilde{V}^-) = 0$ . Performing several times the mentioned construction we get a manifold  $V'$ , satisfying  $t^{N-1} \text{Tor}_{tH_k}(\tilde{V}'^-) = 0$ , and we end by induction in  $N$ .

1. By repeating the argument, exhibited during the proof of theorem 7.3, we find first of all

$$H_p(\tilde{W}, t\tilde{V}) \approx H_p(\tilde{V}^-, t\tilde{V}'^-) \approx \left( \bigoplus_{i=1}^{b_p} R \right) \oplus \bigoplus_{j=1}^{q_p} R / a_{j,0}^{(p)} R, \quad p \leq k-1. \quad (8.2)$$

Furthermore, the complex  $C_*(\tilde{W}, t\tilde{V}) = C_*(\tilde{V}^-, t\tilde{V}'^-)$  is homotopy equivalent to a free finitely generated  $R$ -complex

$$C_* = \{ 0 \xleftarrow{\partial_2} C_2 \xleftarrow{\dots} \xleftarrow{\partial_{n-2}} C_{n-2} \xleftarrow{0} \}$$

which through dimensions  $\leq k-1$  is of the standard type, corresponding to the representation (8.2). This means that for  $r \leq k-1$  we have  $C_r = (R)^{b_p} \oplus (R)^{q_p} \oplus (R)^{q_{p-1}}$ ,  $\partial|(R)^{b_p} = 0$ ,  $\partial|(R)^{q_p} = 0$ , and the differential  $\partial|(R)^{q_{p-1}} : (R)^{q_{p-1}} \rightarrow (R)^{q_{p-1}} \subset C_{r-1}$  is given by a diagonal matrix with diagonal entries  $a_{j,0}^{(r-1)}$ . Besides, the image

$$\partial_k : C_k \rightarrow C_{k-1} \approx R^{b_{k-1}} \oplus R^{q_{k-1}} \oplus R^{q_{k-2}}$$

coincides with the submodule  $(\bigoplus_{j=1}^{q_{k-1}} a_{j,0}^{(k-1)} R) \subset (R)^{q_{k-1}}$  (see lemma 5.1). This implies that  $\text{Ker } \partial_k$  splits off as a direct summand:  $C_k = \text{Ker } \partial_k \oplus (R)^{q_{k-1}}$  that and  $\partial_k|(R)^{q_{k-1}}$  is a diagonal injective operator. Having added to  $C_*$  if necessary the complex  $0 \leftarrow (R)^{q_{k-1}} \leftarrow (R)^{q_{k-1}} \leftarrow 0$ , located in dimensions  $(k, k+1)$ , we may assume that  $\text{Ker } \partial_k$  is a free  $R$ -module.

According to theorem 4.1 and the remark following it we can realize  $C_*$  by a regular Morse function on the cobordism  $(W; V, tV)$ . This function gives rise to a handle decomposition of the pair  $(W, tV)$ . Denote by  $Y$  the result of attaching to  $V$  of all the handles of indices  $\leq k-1$  and those  $q_{k-1}$  handles of indice  $k$ , which correspond to the direct summand  $(R)^{q_{k-1}} \subset C_k$ .

The upper boundary of  $Y$  (which forms the result of corresponding surgical modification of  $tV$ ) will be denoted by  $V_1$  (see the picture). We have attached only the handles of indices  $i$ , where  $2 \leq i \leq n-4$ , hence  $\pi_1(Y) \approx \pi_1(V_1) \approx \mathbb{Z}^{m-1}$  and  $V_1$  is again an admissible splitting.

Lemma 8.1.1. 1) The embedding  $\tilde{V}_1^- \subset \tilde{V}^-$  induces an isomorphism  $H_s(\tilde{V}_1^-) \rightarrow H_s(\tilde{V}^-)$  for  $s \leq k-1$ .

2) The embedding  $(t\tilde{V})^- \subset \tilde{V}_1^-$  induces an isomorphism  $H_s(t\tilde{V}^-) \rightarrow H_s(\tilde{V}_1^-)$  for  $s \geq k$ .

The proof of lemma 8.1.1. 1) Consider the segment of the exact sequence of the couple  $(\tilde{V}^-, V_1^-)$  :

$$H_{s+1}(\tilde{V}^-, \tilde{V}_1^-) \rightarrow H_s(\tilde{V}_1^-) \rightarrow H_s(\tilde{V}^-) \rightarrow H_s(\tilde{V}^-, \tilde{V}_1^-).$$

The manifold  $W$  is obtained from  $Y$  by attaching handles of indices  $\geq k$ , hence for  $s \leq k-1$  we have  $H_s(\tilde{W}, \tilde{Y}) \approx H_s(\tilde{V}^-, \tilde{V}_1^-) = 0$  and 1) is proved for  $s \leq k-2$ .

Further, note that the boundary operator

$$\partial : H_k(\tilde{W}, \tilde{Y}) \rightarrow H_{k-1}(\tilde{Y}, t\tilde{V})$$

vanishes by construction (recall that the cellular decomposition of  $(\tilde{W}, \tilde{Y})$  starts with  $k$ -dimensional cells, having zero boundary in the cell complex of  $(W, tV)$ ; the  $k$ -dimensional cells with



nonzero boundary are included into  $Y$ ). Therefore the image of differential  $H_k(\tilde{V}^-, \tilde{V}_1^-) \rightarrow H_{k-1}(\tilde{V}_1^-)$  is contained in  $\text{Im}(H_{k-1}(t\tilde{V}^-) \rightarrow H_{k-1}(\tilde{V}_1^-))$  and also (by obvious reasons) in  $\text{Ker}(H_k(\tilde{V}_1^-) \rightarrow H_{k-1}(\tilde{V}^-))$ . But  $V$  is  $(k-1)$ -regular, hence the homology  $H_{k-1}(\tilde{V}^-)$  has no  $t$ -torsion and the map  $H_{k-1}(t\tilde{V}^-) \rightarrow H_{k-1}(\tilde{V}^-)$  has no kernel. Consequently

$$\text{Im}(\partial: H_k(\tilde{V}^-, \tilde{V}_1^-) \rightarrow H_{k-1}(\tilde{V}_1^-)) = 0$$

and this proves p.1).

2) Consider the segment of the exact sequence of the pair  $(\tilde{V}_1^-, t\tilde{V}^-)$ :

$$H_{s+1}(\tilde{V}_1^-, t\tilde{V}^-) \rightarrow H_s(t\tilde{V}^-) \rightarrow H_s(\tilde{V}_1^-) \rightarrow H_s(\tilde{V}_1^-, t\tilde{V}^-).$$

By construction the cell decomposition of the pair  $(\tilde{V}_1^-, t\tilde{V}^-) = (\tilde{Y}, t\tilde{V})$  contains only the cells of dimensions  $\geq k$ . Furthermore, the boundary operator in dimension  $k$  is injective, thus  $H_s(\tilde{V}_1^-, t\tilde{V}^-) = 0$  for  $s \geq k$ . This implies our assertion.

2. Now we turn directly to proof of lemma 8.1.

We'll consider the admissible splitting  $tV$  instead of  $V$  and kill the element  $\alpha \in {}^tH_k(t\tilde{V}^-)$ ,  $\alpha \in \text{Ker}(H_k(t\tilde{V}^-) \rightarrow H_k(\tilde{V}^-))$ .

Consider the embedding of manifolds  $V_1 \rightarrow X$ , where  $X = W \setminus \text{Int } Y = V^- \cap V_1^+$  (see the picture). By excision  $H_*(\tilde{X}, \tilde{V}_1) \approx H_*(\tilde{V}^-, \tilde{V}_1^-)$ . The first nonzero homology  $H_*(\tilde{V}^-, \tilde{V}_1^-)$  appears in dimension  $k$ , therefore the strong Hurewicz theorem for simply-connected pairs implies that the Hurewicz map

$$H: \pi_{k+1}(\tilde{X}, \tilde{V}_1) \rightarrow H_{k+1}(\tilde{X}, \tilde{V}_1)$$

is surjective. Pick an element  $a$  of  $H_{k+1}(\tilde{X}, \tilde{V}_1)$  such that

$\partial a \in H_k(\tilde{V}_1^-)$  is homologous in  $\tilde{V}_1^-$  to the element  $\mathcal{A}$  and an element  $A$  of  $\pi_{k+1}(X, V_1) \approx \pi_{k+1}(\tilde{X}, \tilde{V}_1^-)$ , such that  $H(A) = a$ .

According to the corollary 1.1 of [20] (Siebenmann's theorem) any element of  $\pi_i(Q^q, P)$ , where  $Q$  is a manifold and  $P$  is a component of  $\partial Q$ , can be realized by a smooth embedding of the disc  $(D^i, S^{i-1}) \rightarrow (Q^q, P)$ , provided  $i \leq q-3$ ,  $\pi_j(Q^q, P) = 0$  for  $j \leq 2i-q+1$ .

Set for our purposes  $Q^q = X$ ,  $P = V_1$ ,  $q = n$ ,  $i = k+1$ . The groups  $\pi_*(X, V_1) = \pi_*(\tilde{X}, \tilde{V}_1^-)$  vanish for  $* \leq k-1$ . Observe now that  $2i-q+1 = 2(k+1)-n+1 = k + (k-n) + 3 \leq k-1$ , and thus the assumptions of Siebenmann theorem hold. Realize now the element  $A \in \pi_{k+1}(X, V_1)$  by a smoothly embedded disc, consider a small tubular neighbourhood of this disc and attach it to  $V_1$  (see the picture). Denote by  $V_0$  the boundary of manifold thus obtained;  $V_0$  is the result of a surgical modification of  $V_1$  with respect to the sphere  $\partial A$ .

The embedding  $V^- \subset V_0^-$  induces an isomorphism  $H_s(\tilde{V}^-) \xrightarrow{\cong} H_s(\tilde{V}_0^-)$  for  $s \leq k-1$  and for  $s = k$  we have  $H_k(\tilde{V}_0^-) = H_k(\tilde{V}^-)/(\mathcal{A})$  (here  $(\mathcal{A})$  stands for the  $\mathbb{R}$ -submodule in  $H_k(\tilde{V}^-)$ , generated by  $\mathcal{A}$ ). Recall now that for  $s \leq k-1$  the homology modules  $H_s(\tilde{V}_1^-)$  are isomorphic to  $H_s(\tilde{V}^-) \approx H_s(t\tilde{V}^-)$  and if  $s = k$  then  $H_k(\tilde{V}_1^-) \approx H_k(t\tilde{V}^-)$ ; consequently  $H_k(\tilde{V}_0^-) \approx H_k(t\tilde{V}^-)/(\mathcal{A})$  and using p. 0 we get the manifold sought. Lemma 8.1 is proved.

Proof of lemma 8.2. Consider the admissible splitting  $V_1 \subset W$ , constructed in p.1 of the proof of lemma 8.1.

We have the embeddings

$$H_k(t\tilde{V}^-) = tA \subset B \subset A = H_k(\tilde{V}^-)$$

here  $A/tA \approx H_k(\tilde{V}^-, t\tilde{V}^-)$ . Consider a generator  $m$  of the  $R$ -module  $B/tA \subset A/tA$  and the image  $m'$  of  $m$  in the module  $H_k(\tilde{V}^-, \tilde{V}_1^-) \approx H_k(\tilde{X}, \tilde{V}_1)$ . We've shown in the course of proof of lemma 8.1, that  $H_k(\tilde{X}, \tilde{V}_1)$  is a first nontrivial homology module of the pair  $(\tilde{X}, \tilde{V}_1)$  and that it is isomorphic to  $\pi_k(\tilde{X}, \tilde{V}_1) \approx \pi_k(X, V_1)$ . By the virtue of argument similar to that of the proof of lemma 8.1 we can realize  $m' \in \pi_k(X, V_1)$  by a smoothly embedded disc  $(D^k, S^{k-1})$ . Attach now the corresponding handle  $\mu$  (see the picture). Denote by  $V_0$  the upper boundary of the manifold obtained;  $V_0$  is the result of the surgical modification of  $V_1$  with respect to the sphere  $\partial m'$ . Now we'll show that  $V_0$  satisfies the conclusion of lemma 8.2.

Observe that  $\partial m' \in H_{k+1}(\tilde{V}_1^-)$  vanishes (otherwise the map  $H_{k-1}(\tilde{V}_1^-) \rightarrow H_{k-1}(\tilde{V}^-)$  would have a non trivial kernel, and this is impossible since the embedding  $\tilde{V}_1^- \rightarrow \tilde{V}^-$  induces an isomorphism in  $(k-1)$ -homology and  $H_{k-1}(\tilde{V}^-)$  has no  $t$ -torsion). Thus we get from the exact sequence of the pair  $(\tilde{V}_0^-, \tilde{V}_1^-)$

$$1) H_s(\tilde{V}_0^-) \approx H_s(\tilde{V}^-) \quad \text{for } s \leq k-1$$

2)  $H_k(\tilde{V}_0^-) \approx H_k(\tilde{V}_1^-) \oplus R(\bar{m})$ , where  $R(\bar{m})$  is a free module generated by an element  $\bar{m}$ , which is sent by the composition

$$H_k(\tilde{V}_0^-) \longrightarrow H_k(\tilde{V}^-) \longrightarrow H_k(\tilde{V}^-, \tilde{V}_1^-)$$

to the element  $m'$ .

Compute now the image of  $H_k(\tilde{V}_0^-)$  under the map  $\tilde{V}_0^- \rightarrow \tilde{V}^-$ . Since  $H_k(t\tilde{V}^-) \rightarrow H_k(\tilde{V}_1^-)$  is an isomorphism, the image of  $H_k(\tilde{V}_1^-) \rightarrow H_k(\tilde{V}_0^-)$  equals the image of  $H_k(t\tilde{V}^-)$  in  $H_k(\tilde{V}^-) = A$ , i.e. the submodule  $tA \subset A$ .

The homology  $H_k(\tilde{V}_1^-, t\tilde{V}^-)$  vanishes by construction (see

the proof of lemma 8.1), thus the injectivity of  $H_k(\tilde{V}^-, t\tilde{V}^-) \rightarrow H_k(\tilde{V}^-, \tilde{V}_1^-)$  follows immediately from the exact sequence of the triple  $(\tilde{V}^-, \tilde{V}_1^-, t\tilde{V}^-)$ . Therefore the projection of  $H_k(\tilde{V}^-) \approx A$  onto  $H_k(\tilde{V}^-, t\tilde{V}^-) \approx A/tA$  sends the image of  $\bar{m}$  in  $H_k(\tilde{V}^-)$  to the element  $m \in A/tA$ .

Therefore the image of  $H_k(\tilde{V}_0^-)$  in  $A = H_k(\tilde{V}^-)$  equals  $tA + (m) = B$ .

Now we'll compute the kernel of  $H_k(\tilde{V}_0^-) \rightarrow A$ . If  $x \in H_k(\tilde{V}_0^-)$  goes to zero via the map  $\tilde{V}_0^- \rightarrow \tilde{V}^-$  then  $tx \in H_k(\tilde{V}_0^-)$  goes to zero via the map  $t\tilde{V}_0^- \rightarrow t\tilde{V}^- \subset \tilde{V}_0^-$ ; that means  $tx = 0$ . On the other hand,  $A$  lacks  $t$ -torsion, hence

$$\text{Tor}_t H_k(\tilde{V}_0^-) \subset \text{Ker}(H_k(\tilde{V}_0^-) \rightarrow H_k(\tilde{V}^-)) \subset {}^t H_k(\tilde{V}_0^-)$$

Lemma 8.2 is proved.

### 9. The surgery in the dimensions left

The proof of lemma 7.4 will be split into several lemmas.

Lemma 9.1. Let  $M^n$  be a manifold,  $\pi_1 M^n = \mathbb{Z}^m$ . Suppose that a class  $\gamma \in H^1(M, \mathbb{Z})$  is represented by an epimorphism  $\mathbb{Z}^m \rightarrow \mathbb{Z}$  and that  $S^{-1}H_q(M^n, \mathbb{Z}) = 0$ .

Then for any admissible splitting  $V$  the module  $H_q(\tilde{M}^n, \tilde{V}^-)$  is a finitely generated  $R$ -module (we use the notations (5.1)) isomorphic to  $\text{Ker}(H_{q-1}(\tilde{V}^-) \rightarrow H_{q-1}(\tilde{M}))$ . If  $V$  is  $(q-1)$ -regular then  $H_q(\tilde{M}^n, \tilde{V}^-) = 0$ .

Proof of lemma 9.1. The condition  $S^{-1}H_q(\tilde{M}^n) = 0$  means that every cohomology class  $x \in H_q(\tilde{M}^n)$  is annihilated by some polynomial  $1+tQ(t)$ ,  $Q(t) \in R[t] = \mathbb{Z}[\mathbb{Z}^{m-1}][t]$ , hence  $x = -tQ(t)x$ .

Hence  $H_q(\tilde{V}^-) \rightarrow H_q(\tilde{M}^n)$  is an epimorphism. Furthermore,

$$\text{Ker}(H_{q-1}(\tilde{V}^-) \rightarrow H_{q-1}(\tilde{M})) \approx \text{Tor}_t H_{q-1}(\tilde{V}^-)$$

and since  $H_{q-1}(\tilde{V}^-)$  is finitely generated over  $P$ ,  $\text{Tor}_t H_q(\tilde{V}^-)$  is finitely generated over  $R$ . Now the lemma follows directly from the exact sequence of the pair  $(\tilde{M}^n, \tilde{V}^-)$ .

For an  $(r-1)$ -regular splitting  $V$  the map  $H_{r-1}(\tilde{V}^-) \rightarrow H_{r-1}(\tilde{M})$  is injective, hence  $H_r(\tilde{M}, \tilde{V}^-) = 0$  and we get the point 1) of lemma 7.4.

Lemma 9.2. Suppose that  $M^n$  is a manifold,  $\pi_1 M^n = \mathbb{Z}^m$ ,  $\gamma \in H^1(M, \mathbb{Z})$  is an epimorphism  $\mathbb{Z}^m \rightarrow \mathbb{Z}$ , the modules  $S^{-1}H_p(\tilde{M}^n)$  satisfy (7.1) for  $p \leq k$  and  $(V, \nu)$  is  $k$ -regular.

Then for all natural  $q$ ,

1) the  $R$ -modules  $H_p(\tilde{V}^-, t^q \tilde{V}^-)$  have resolutions of length 2 for  $p \leq k$ ,

2) the homomorphism

$$H^p(\tilde{V}^-, t^{q+l} \tilde{V}^-) \rightarrow H^p(t^q \tilde{V}^-, t^{q+l} \tilde{V}^-)$$

induced by embedding of the pairs is an epimorphism for  $p \leq k$ <sup>6)</sup>.

Proof. 1) Compute first the homology  $H_p(\tilde{V}^-, t^q \tilde{V}^-)$ . Since  $V$  is  $k$ -regular, the embedding  $t^q \tilde{V}^- \subset \tilde{V}^-$  induces in  $p$ -homology a monomorphism, which equals  $t^q : H_p(\tilde{V}^-) \rightarrow H_p(\tilde{V}^-)$ . Hence

$$H_p(\tilde{V}^-, t^q \tilde{V}^-) \approx H_p(\tilde{V}^-) / t^q H_p(\tilde{V}^-).$$

Since (7.2) holds for  $H_p(\tilde{V}^-)$  we can apply lemma (5.4) to com-

6) We mean the cohomology of corresponding universal coverings with coefficients in the  $R = \mathbb{Z}[\pi_1 V^-]$ -module  $R$ , or equivalently the cohomology with compact supports.

pute the above factor.

We get

$$\begin{aligned}
 H_p(\tilde{V}^-, t^q \tilde{V}^-) &\approx \left( \bigoplus_{i=1}^{b_p} P / t^q P \right) \oplus \left( \bigoplus_{j=1}^{q_p} P / (a_j^{(p)}, t^q) P \right) \approx \\
 &\approx \left( \bigoplus_{i=1}^{b_p} R^q \right) \oplus \left( \bigoplus_{j=1}^{q_p} R^q / F_j^{(p)}(R^q) \right) \quad (9.1)
 \end{aligned}$$

where  $F_j^{(p)} : R^q \rightarrow R^q$  is a monomorphism, given by the matrix

$$\begin{pmatrix}
 a_{j,0}^{(p)} & a_{j,1}^{(p)} & \dots & a_{j,q}^{(p)} \\
 & a_{j,0}^{(p)} & \dots & a_{j,q-1}^{(p)} \\
 & & \dots & \\
 & & & a_{j,0}^{(p)}
 \end{pmatrix}$$

The assertion 1) is proved.

Next we note that for a free finitely generated complex  $C_*$ , such that for  $p \leq k$  the module  $H_p(C_*)$  has a free resolution of length 2, the spectral sequence

$$\begin{aligned}
 E_2^{p,s} &= \text{Ext}^p(H_s(C_*), R) \Rightarrow \\
 &\Rightarrow H_{p+s}(\text{Hom}_R(C_*, R)) = H^{p+s}(C_*, R)
 \end{aligned}$$

degenerates in  $E_2$  for  $p \leq k$  and there exists a functorial exact sequence

$$0 \rightarrow \text{Ext}^1(H_{p-1}(C_*), R) \rightarrow H^p(C_*, R) \rightarrow \text{Hom}(H_p(C_*), R) \rightarrow 0$$

The embedding  $(t^q \tilde{V}^-, t^{q+l} \tilde{V}^-) \subset (\tilde{V}^-, t^{q+l} \tilde{V}^-)$  induces the homomorphism of these exact sequences which is an epimorphism

on the left (by virtue of lemma 5.4.3) and an epimorphism on the right (obviously), hence the middle map

$$H^p(\tilde{V}^-, t^{q+l}\tilde{V}^-) \rightarrow H^p(t^q\tilde{V}^-, t^{q+l}\tilde{V}^-)$$

is an epimorphism. Lemma 9.2 is proved.

Lemma 9.3. Let  $\pi_1 M = \mathbb{Z}^m$ ,  $n \geq 6$ ,  $\lambda \in H^1(M, \mathbb{Z})$  be an epimorphism  $\mathbb{Z}^m \rightarrow \mathbb{Z}$ ,  $1 \leq r \leq n-2$ ,  $m_r = m_{r+1} = 0$ ,  $(V, \nu)$  be an  $(r-1)$ -biregular splitting.

Then there exists a natural number  $q_0$  such that for  $q > q_0$  the embedding  $(\tilde{V}^-, t^q\tilde{V}^-) \rightarrow (\tilde{M}, t^q\tilde{V}^-)$  induces an isomorphism

$$H_{r+1}(\tilde{V}^-, t^q\tilde{V}^-) \xrightarrow{\cong} H_{r+1}(\tilde{M}, t^q\tilde{V}^-)$$

and the differential of the exact sequence of the triple  $(\tilde{M}, \tilde{V}^-, t^q\tilde{V}^-)$  induces an isomorphism

$$H_{r+1}(\tilde{M}, \tilde{V}^-) \cong H_r(\tilde{V}^-, t^q\tilde{V}^-)$$

Proof. Consider the exact sequence of the triple  $(\tilde{M}, \tilde{V}^-, t^q\tilde{V}^-)$ :

$$\begin{aligned} &\rightarrow H_{r+2}(\tilde{M}, t^q\tilde{V}^-) \rightarrow H_{r+2}(\tilde{M}, \tilde{V}^-) \xrightarrow{\partial_{r+1}} H_{r+1}(\tilde{V}^-, t^q\tilde{V}^-) \\ &\rightarrow H_{r+1}(\tilde{M}, t^q\tilde{V}^-) \rightarrow H_{r+1}(\tilde{M}, \tilde{V}^-) \xrightarrow{\partial_r} \\ &\rightarrow H_r(\tilde{V}^-, t^q\tilde{V}^-) \rightarrow H_r(\tilde{M}, t^q\tilde{V}^-). \end{aligned}$$

According to lemma 9.1  $H_r(\tilde{M}, t^q\tilde{V}^-) = 0$  and  $H_{r+1}(\tilde{M}, \tilde{V}^-) \cong H_{r+1}(\tilde{M}, t^q\tilde{V}^-)$  is finitely generated over  $R$ , hence for  $q$  sufficiently large the embedding  $(\tilde{M}, t^q\tilde{V}^-) \subset (\tilde{M}, \tilde{V}^-)$  induces the

zero map in  $(r+1)$ -homology. That implies the second statement of the lemma.

To prove the first it suffices to verify the surjectivity of the map  $H_{r+2}(\tilde{M}, t^{q\tilde{V}^-}) \rightarrow H_{r+2}(\tilde{M}, \tilde{V}^-)$ . For this it suffices in turn to verify the surjectivity of the map  $H_{r+2}(t^{-l}\tilde{V}^-; t^{q\tilde{V}^-}) \rightarrow H_{r+2}(t^{-l}\tilde{V}^-, \tilde{V}^-)$  for all natural  $l, q$ .

Consider now the manifold  $Z = t^{qV^+} \cap t^{-l}V^-$  with the boundary  $t^{qV} \cup t^{-l}V$ ; the manifold  $Z_0 = V^+ \cap t^{-l}V^-$  with the boundary  $V \cup t^{-l}V$  and the embedding of pairs  $(Z, t^{qV}) \subset (Z, V^- \cap t^{qV^+})$ . This embedding is a map of degree 1, i.e. it sends the infinite fundamental cycle  $U_Z \in H_n^{\text{inf}}(\tilde{Z}, t^{q\tilde{V}} \cup t^{-l}\tilde{V})$  to the infinite fundamental <sup>(cycle)</sup>  $U_{Z_0} \in H_n^{\text{inf}}(\tilde{Z}_0, \tilde{V} \cup t^{-l}\tilde{V})$ . Thus we obtain a commutative diagram

$$\begin{array}{ccc} H_{r+2}(t^{-l}\tilde{V}^-, t^{q\tilde{V}^-}) & \xleftarrow{\cap U_Z} & H_c^{n-r-2}(t^{q\tilde{V}^+}, t^{-l}\tilde{V}^+) \\ \downarrow & & \downarrow \\ H_{r+2}(t^{-l}\tilde{V}^-, \tilde{V}^-) & \xleftarrow{\cap U_{Z_0}} & H_c^{n-r-2}(\tilde{V}^+, t^{-l}\tilde{V}^+) \end{array}$$

The horizontal arrows are isomorphisms by Poincaré duality, and it suffices to show that the righthand arrow is an epimorphism. Since  $(V, \nu)$  is  $(r-1)$ -biregular,  $(V, -\nu)$  is  $(n-r-2)$ -regular and we deduce the required assertion from lemma 9.2 p.2).

Lemma 9.4. Under the assumptions of lemma 9.3 for any  $q$  there exists a Morse function  $\varphi$  on the cobordism  $(V^- \cap t^{qV^+}, t^{qV})$ , such that the differentials  $\partial_r : C_r \rightarrow C_{r-1}$ ,  $\partial_{r+2} : C_{r+2} \rightarrow C_{r+1}$  of the corresponding complex  $C_*(\varphi) = \{0 \leftarrow C_2 \leftarrow \dots \leftarrow C_{n-2} \leftarrow 0\}$  vanish.

Proof. We proceed similarly to the proof of lemma 8.1. Denote by  $W_q$  the manifold  $\tilde{V}^- \cap t^{qV^+}$  with the boundary



$V \cup t^q V$ . By lemma 9.2 the  $R$ -modules  $H_p(\tilde{W}_q, t^q \tilde{V})$  have the resolutions of length 2 for  $p \leq r-1$ . Represent the complex  $C_*(\tilde{W}_q, t^q \tilde{V})$  up to homotopy by a standard one  $C_*^0$  (lemma 5.1) and realize  $C_*^0$  as a Morse complex of a regular function  $f$  on the cobordism  $W$ . Denote by  $V_1$  a level surface  $f^{-1}(c)$ , separating the critical points of indices  $\leq r-1$  from critical points of indices  $\geq r$ .  $V_q$  is the result of a surgical modifications of indices  $\leq r-1$  of the manifold  $t^q V$ . Set  $Y_q = V_q^- \wedge t^q V_q^+$ ,  $X_q = V_q^+ \wedge V_q^-$  (these notations are similar to the notations of lemma 8.1; the picture illustrates after necessary corrections the present situation as well). The Morse function  $(-f)$  gives rise to a handle decomposition of the pair  $(W_q, V)$ , and  $X_q$  is precisely the result of attaching all the handles of indices  $\leq n-r$  to  $V$ . The Morse complex of  $(-f) | (X_q, V)$  looks like  $\{0 \leftarrow D_2 \leftarrow \dots \leftarrow D_{n-r} \leftarrow 0\}$ . Furthermore, the embedding  $(X_q, V) \rightarrow (W_q, V)$  induces an isomorphism in the homology  $H_s(\tilde{X}_q, \tilde{V}) \rightarrow H_s(\tilde{W}_q, \tilde{V})$  for  $s \leq n-r$ . Apply now lemma 7.2 to  $(n-r-2)$ -regular splitting  $(V, -V)$  to get a Morse function  $f$  on the cobordism  $(X_q, V)$ , such that its Morse complex  $\{0 \leftarrow \tilde{D}_2 \leftarrow \dots \leftarrow \tilde{D}_{n-r} \leftarrow 0\}$  is standard in dimensions  $\leq n-r-2$ . Since  $q_{n-r-2} = q_{r+1} = 0$  the differential  $\partial_{n-r-1}: \tilde{D}_{n-r-1} \rightarrow \tilde{D}_{n-r-2}$  vanishes.

Define now a Morse function  $\varphi$  by setting  $\varphi | Y_q = f$ ,  $\varphi | X_q = -f$  ( $V_q$  is a level surface for  $f$  and for  $\tilde{f}$  and we may assume that  $f | V_q = -\tilde{f} | V_q$  and that in a neighbourhood of  $V_q$  both  $f$  and  $(-\tilde{f})$  coincide with the coordinate, normal to  $V_q$ , so that the definition makes sense).

The Morse complex of this function coincides with  $C_*(f)$

for  $* \leq r-1$  and with  $C_*(-\tilde{f})$  for  $* \geq r+1$ . Hence the differential  $\partial_{n-r-1}$  vanishes. By construction

$$C_{r-1}(f) = R^{b_{r-1} \cdot q} \oplus B,$$

where  $\partial | R^{b_{r-1} \cdot q} = 0$ ,  $\partial | B$  is injective,  $R^{b_{r-1} \cdot q} \rightarrow \rightarrow H_{r-1}(C_*(f)) \approx H_{r-1}(\tilde{W}_q, t^{q\tilde{V}})$  is an isomorphism. (Here we use  $q_{r-1} = 0$ .) Therefore the differential  $\partial_r$  must vanish. Lemma 9.4 is proved.

Proof of lemma 7.4. We are to prove p.2 of the statement (p.1 was proved in lemma 9.1). For this we'll construct a free (maybe infinitely generated)  $\mathbb{Z}[\mathbb{Z}^{m-1}]$ -complex  $D_*$ , such that homology of  $D_*$  is isomorphic to  $H_*(\tilde{M}, \tilde{V}^-)$  and the differentials  $\partial_r, \partial_{r+2}$  of this complex vanish. From this the lemma follows easily (similarly to [10, lemma 3.5]). Indeed,  $H_r(D_*) = H_r(\tilde{M}, \tilde{V}^-) = 0$ , implies  $H_{r+1}(\tilde{M}, \tilde{V}^-) \oplus D_r = D_{r+1}$ , i.e.  $H_{r+1}(\tilde{M}, \tilde{V}^-)$  is projective. By lemma 9.3 this module is finitely generated, hence (by Suslin-Quillen theorem) free.

To construct  $D_*$  we pick a number  $N = q_0$  and consider the embeddings

$$(t^{-N}V^-, V^-) \subset (t^{-2N}V^-, V^-) \subset \dots \tag{9.2}$$

The union of this sequence is  $(\tilde{M}, \tilde{V}^-)$ ; the direct limit commutes with homology and we get

$$\lim_{\rightarrow} H_* (t^{-N} \tilde{V}^-, \tilde{V}^-) = H_* (\tilde{M}, \tilde{V}^-).$$

Choose a triangulation of this pairs in such a way that every pair is a subcomplex of the next one. The factorcomplex of the

kth pair by (k-1)th is  $(t^{-2kN} V^-, t^{-2(k-1)N} V^-)$ . We have proved in lemma 9.3 that the differential maps  $H_{r+1}(\tilde{M}, \tilde{V}^-)$  isomorphically onto  $H_r(\tilde{V}^-, t^{N\tilde{V}^-})$ , hence the differential  $\partial_r : H_r(\tilde{V}^-, t^{N\tilde{V}^-}) \rightarrow H_{r-1}(t^{N\tilde{V}^-})$  vanishes. In the same lemma we have proved that  $H_{r+2}(t^{-2kN\tilde{V}^-}, \tilde{V}^-)$  is mapped surjectively onto  $H_{r+2}(t^{-2kN\tilde{V}^-}, t^{-2(k-1)N\tilde{V}^-})$ . This implies that the differential  $\partial_r, \partial_{r+2}$  of the exact sequence of the triple  $(t^{-2kN\tilde{V}^-}, t^{-2(k-1)N\tilde{V}^-}, \tilde{V}^-)$  vanish.

We'll need a purely algebraic

Lemma 9.5. Let  $X_* \subset Y_*$  be free finitely generated complexes, such that  $X_n$  are direct summands of  $Y_n$ . Suppose that  $X_*$  and  $Z_* = Y_*/X_*$  are homotopy equivalent to the free finitely generated complexes  $X'_*$  and correspondingly  $Z'_*$ , such that rth and (r+2)th differentials of  $X'_*$  and  $Z'_*$  vanish. Suppose further that the differentials

$$\delta_r : H_r(Z'_*) \rightarrow H_{r-1}(X'_*), \quad \delta_{r+2} : H_{r+2}(Z'_*) \rightarrow H_{r+1}(X'_*)$$

in the exact sequence of the pair  $(Y_*, X_*)$  vanish.

Then there exists a pair of free finitely generated complexes  $\tilde{X}_* \subset \tilde{Y}_*$ , homotopy equivalent to the pair  $X_* \subset Y_*$ , and such that the modules  $X_n$  are direct summands in  $Y_n$ , and the differentials  $\partial_r$  and  $\partial_{r+2}$  of the complex  $\tilde{Y}_*$  vanish.

Proof of lemma 9.5. According to Cockroft-Swan [15] the complexes  $X_*$  and  $X'_*$  (as well as  $Z_*$  and  $Z'_*$ ) can be made isomorphic by adding to them several complexes of the type  $0 \leftarrow F \xleftarrow{\text{id}} F \leftarrow 0$ , where  $F$  is some free module. From this we easily deduce that there exists a pair of complexes  $X'_* \subset Y'_*$  (the modules  $X'_n$  being the direct summands of  $Y'_n$ ), such that

$Y'_*/X'_* \approx Z'_*$ . Consider now the free generators  $z_i$  of the module  $Z'_*$ . Our assumptions imply  $\partial z_i \in X'_{r-1}$ ; furthermore, these elements are homologous to zero in this complex, hence they are zero itself (since  $\partial_r | X'_r = 0$ ). The same argument proves that  $\partial_{r+2}$  vanishes.

Return now to the sequence (9.2). According to lemma 9.5 there exists a sequence  $X_*^{(1)} \subset X_*^{(2)} \subset \dots$ , where  $X_*^{(i)}$  are free finitely generated complexes,  $X_n^{(i)}$  being the direct summands in  $X_n^{(i+1)}$ , and the homotopy equivalences

$$C_*(t^{-N}V^-, V^-) \subset C_*(t^{-2N}V^-, V^-) \subset \dots$$

$$X_*^{(1)} \subset X_*^{(2)} \subset \dots$$

Now we set  $D_* = X_*^{(i)} = \varinjlim X_*^{(i)}$  and lemma 7.4 is proved.

Proof of lemma 7.5. It will occupy the rest of § 9. We obtain the proof by reproducing the argument of [10] in our setting. We exhibit the argument here, elaborating on these parts which need modification.

Denote by  $\mathcal{G}(R)$  the category formed by pairs  $(F, f)$  where  $F$  is a free finitely generated  $R$ -module and  $f$  is a nilpotent endomorphism of  $F$ . (As always  $R = \mathbb{Z}[\mathbb{Z}^{m-1}]$ . The definition is somewhat simplified in comparison with [10]: we omit the automorphism  $\alpha$  since fundamental groups are abelian and we use the free modules since all  $R$ -projectives are  $R$ -free.) Denote by  $C(R)$  the set of equivalence classes of isomorphism classes of objects from  $\mathcal{G}(R)$  with respect to equivalence relation, generated by two relations:

1)  $(F, f) \sim (F \oplus F', f \oplus 0)$

2) if the sequence  $0 \rightarrow (F_2, f_2) \rightarrow (F_1, f_1) \rightarrow (F_0, f_0) \rightarrow 0$  is exact, then  $(F_1, f_1) \sim (F_2, f_2) \oplus (F_0, f_0)$ .

It is proved in [10] that  $c(R) = 0$ .

Suppose now that the assumptions of lemma 7.4 hold. Then the module  $H_{r+1}(\tilde{M}, \tilde{V}^-)$  together with the endomorphism  $t$  gives rise to an object of  $\mathcal{G}(R)$ , which we denote by  $c(V, \nu)$ . It vanishes when we pass to  $C(R)$ . We will now prove (following [10]), that this equivalence to zero can be realized geometrically, so that we can construct an  $(r-1)$ -biregular splitting  $(V', \nu')$  with  $c(V', \nu') = 0$  (i.e. satisfying the conclusion of lemma 7.5).

The ring  $R$  is a subring of  $\mathbb{Z}[\pi_1 M]$ , stable with respect to automorphism  $\chi$  (see § 3). Denote by  $\overline{\text{Hom}}(M, N)$  the set of all  $\chi$ -homomorphisms of  $R$ -module  $N$  into  $R$ -module  $M$ . For any  $\mathcal{G}(R)$ -object  $X = (F, f)$  we define the dual  $X^* = (\text{Hom}(F, R), \pm f^*)$ , where we choose  $(+)$  if  $t \in \pi_1 M$  preserves the orientation and  $(-)$  if not.

Lemma 9.6. Under the assumptions of lemma 7.4

$$c(V, \nu)^* = c(V, -\nu)$$

Proof. The manifold  $(V, -\nu)$  is an  $(n-r-2)$ -regular splitting and

$$c(V, -\nu) = (H_{n-r}(\tilde{M}, \tilde{V}^+), t^{-1}).$$

We know from lemma 9.3 that  $H_{r+1}(\tilde{M}, \tilde{V}^-) \approx H_r(\tilde{V}^-, t^q \tilde{V}^-)$  for  $q$  sufficiently large, hence

$$c(V, \nu) \approx (H_r(\tilde{V}^-, t^q \tilde{V}^-), t).$$

The same lemma implies that

$$c(V, -\nu) \approx (H_{n-r}(t^q \tilde{V}^+, \tilde{V}^+), t^{-1})$$

for  $q$  sufficiently large. The Poincare duality implies that there is a  $\chi$ -isomorphism.

$$(H_c^r(\tilde{V}^-, t^{q\tilde{V}^-}), \pm t) \approx (H_{n-r}(t^{q\tilde{V}^+}, \tilde{V}^+), t)$$

(the sign (+) appears if  $t$  is orientation preserving, otherwise (-) appears). The lemma 9.4 implies that we can choose a cell decomposition of a pair  $(\tilde{V}^-, t^{q\tilde{V}^-})$  such that in the resulting chain complex  $C_*(\tilde{V}^-, t^{q\tilde{V}^-})$  the differential  $\partial_{r+2}$  vanishes. Moreover, lemma 9.3 and p.2 of lemma 7.4 imply that  $H_r(\tilde{V}^-, t^{q\tilde{V}^-})$  is a free module. Hence we obtain

$$(H_c^r(\tilde{V}^-, t^{q\tilde{V}^-}), \pm t) \approx (\text{Hom}(H_r(\tilde{V}^-, t^{q\tilde{V}^-}), R), \pm t)$$

and since  $\text{Hom}(M, R)$  is  $\chi$ -isomorphic to  $\text{Hom}(M, R)$ , the lemma follows.

A triangular object of  $\mathcal{C}(R)$  is by definition an object  $(F, f)$  together with a filtration  $0 = F_0 \subset F_1 \subset \dots \subset F_n = F$ , such that all the factors  $F_{i+1}/F_i$  are free modules of rank 1 and  $f(F_{i+1}) \subset F_i$ .

The basic role in realizing geometrically the relations is played by the following lemma.

Lemma 9.7. Suppose that all the assumptions of lemma 7.5 hold and  $c(V, \nu) = (P, f)$ . Suppose given an exact sequence

$$0 \rightarrow (Q, \psi) \rightarrow (F, \varphi) \xrightarrow{p} (P, f) \rightarrow 0$$

of the objects of  $\mathcal{C}(R)$ , where  $(F, \varphi)$  is a triangular object.

Then there exists an  $r$ -biregular splitting  $(V_0, \nu_0)$ , such that

$$c(V_0, \nu_0) = (Q, \psi)$$

Proof. Denote  $P_i$  the image of  $F_i$  in  $P$  (where  $\{F_i\}$  is the filtration of  $F$ , mentioned above).

Lemma 9.8. Let  $i \leq n$ . There exists an admissible splitting  $(V_i, \mathcal{V}_i)$ , obtained from  $(V, \mathcal{V})$  by a sequence of surgical modifications of indices  $\leq r-1$  and of indice  $(r+1)$  (all the modifications take place inside  $V^+$ ), such that  $H_s(\tilde{V}_i^-) \approx H_s(\tilde{V}^-)$  for  $s \leq r-1$  and the map  $H_{r+1}(\tilde{V}_i^-, \tilde{V}^-) \rightarrow H_{r+1}(\tilde{M}, \tilde{V}^-)$  is isomorphic to the map  $p_i = p|_{F_i} : F_i \rightarrow P$ .

First we deduce lemma 9.7 from lemma 9.8.

Set  $i = n$ . We'll show that  $(V_n, \mathcal{V}_n)$  is an  $r$ -biregular splitting. Consider the exact sequence of the triple  $(\tilde{M}, \tilde{V}_n^-, \tilde{V}^-)$ :

$$\begin{aligned} H_{r+1}(\tilde{V}_n^-, \tilde{V}^-) &\rightarrow H_{r+1}(\tilde{M}, \tilde{V}^-) \rightarrow \\ &\rightarrow H_{r+1}(\tilde{M}, \tilde{V}_n^-) \rightarrow H_r(\tilde{V}_n^-, \tilde{V}^-). \end{aligned}$$

The left arrow is the epimorphism  $F_i \rightarrow P_i$ , the right group vanishes. Therefore  $H_{r+1}(\tilde{M}, \tilde{V}_n^-) = 0$ . Since  $V_n$  is  $(r-1)$ -regular and  $m_r = 0$  we deduce from lemma 9.1 that  $H_r(\tilde{M}, \tilde{V}_n^-) = 0$ . Hence  $H_r(\tilde{V}_n^-, \tilde{V}_n^-) = 0$ , and, consequently,  $V_n$  is  $r$ -regular. Further,  $\tilde{V}_n^-$  is obtained from  $\tilde{V}^-$  by attaching handles of indices  $r+1$ , hence the homology of  $\tilde{V}^+$  did not change through the dimensions  $n-r-3$ , and consequently,  $V_n$  is  $r$ -biregular. Furthermore,  $m_{r+2} = 0$  and lemma 9.1 implies that the  $R$ -module  $H_{r+2}(\tilde{M}, \tilde{V}^-)$  is finitely generated, which implies that for  $N$  sufficiently large there exists an epimorphism  $H_{r+2}(t^{-N}\tilde{V}^-, \tilde{V}^-) \rightarrow H_{r+2}(\tilde{M}, \tilde{V}^-)$ . But  $(V, \mathcal{V})$  is  $(n-r-2)$ -regular and  $m_{n-r-2}(-\lambda) = 0$ , hence (by lemma 9.2)  $H_{r+2}(t^{-N}\tilde{V}^-, \tilde{V}^-) = 0$ . Now our conclusion follows from the exact sequence

$$H_{r+2}(\tilde{M}, \tilde{V}^-) \rightarrow H_{r+2}(\tilde{M}, \tilde{V}_n^-) \rightarrow H_{r+1}(\tilde{V}_n^-, \tilde{V}^-) \rightarrow H_{r+1}(\tilde{M}, \tilde{V}^-).$$

Proof of lemma 9.8. Induction in  $i$ . Suppose that we've constructed an admissible splitting  $V_i$ , satisfying the required properties. We'll identify  $F_i$  with  $H_{r+1}(\tilde{V}_i^-, \tilde{V}^-)$ , the homomorphism  $H_{r+1}(\tilde{V}_i^-, \tilde{V}^-) \rightarrow H_{r+1}(\tilde{M}, \tilde{V}^-)$ , induced by embedding of the pairs, - with  $p|_{F_i}$ , the image  $p(H_{r+1}(\tilde{V}_i^-, \tilde{V}^-)) \subset H_{r+1}(\tilde{M}, \tilde{V}^-)$  - with  $P_i$ . Choose an element  $e_{i+1} \in F_{i+1}$  such that  $F_{i+1} = F_i \oplus R(e_{i+1})$ . Denote  $te_{i+1}$  by  $x \in H_{r+1}(\tilde{V}_i^-, \tilde{V}^-)$ ,  $p(e_{i+1})$  - by  $y \in H_{r+1}(\tilde{M}, \tilde{V}^-)$ . Note that  $p(x) = ty$ .

We prove first that there exists an element  $\sigma \in H_{r+1}(t^{-1}\tilde{V}_i^-, \tilde{V}^-)$ , such that  $t\sigma \in H_{r+1}(\tilde{V}_i^-, \tilde{V}^-)$  equals  $x$  and image  $j_*\sigma$  in  $H_{r+1}(\tilde{M}, \tilde{V}^-)$  equals  $y$ .

Indeed, choose some chains  $\bar{y} \in C_{r+1}(\tilde{M})$ ,  $\bar{x} \in C_{r+1}(\tilde{V}_i^-)$ , which represent  $y$  and  $x$ . The chain  $t\bar{y}$  is homologous to  $p(x)$  modulo  $\tilde{V}^-$ , i.e.  $t\bar{y} = p(\bar{x}) + v + \partial u$ , where  $v \in C_{r+1}(\tilde{V}^-)$ ,  $u \in C_{r+1}(\tilde{M})$ .

The chain  $\bar{\sigma} = t^{-1}(\bar{x} + v)$  is a relative cycle in  $C_{r+1}(t^{-1}\tilde{V}_i^-, \tilde{V}^-)$  and the homology class  $\sigma$  of  $\bar{\sigma}$  satisfy the requirement.

Consider now the admissible splitting  $t^{-1}V_i$  and apply to it the procedure described in p.1 of the proof of lemma 8.1 (where  $k = r$ ). We get the manifold  $V_1 \subset t^{-1}V_i^-$ .

Note that  $V_1^-$  is obtained from  $V^-$  by attaching handles of indices  $\leq r-1$ , since  $m_r = 0$ . The  $\tilde{V}_1^-$ ,  $t^{-1}\tilde{V}_1^-$  are simply connected and the first nonzero homology group of  $(t^{-1}\tilde{V}_1^-, \tilde{V}_1^-)$  sits in dimension  $r$ . Therefore the Hurewicz homomorphism



is surjective. Consider the element  $\sigma' \in \pi_{r+1}(t^{-1}V_i^-, V_1^-)$ , such that  $H(\sigma')$  equals to the  $\sigma$ , reduced modulo  $V_1^-$ . By the same argument as in the proof of lemma 8.1 we realize  $\sigma'$  by the embedded disc  $D = (D^{r+1}, S^r)$  and attach to  $V_1^-$  a small tubular neighbourhood at this disc. The upper boundary of the manifold thus obtained will be denoted  $V_{i+1}$ . We claim that this manifold satisfies the conclusion of lemma 9.8 for the number  $(i+1)$ .

Indeed, if  $s \leq r-1$ , then

$$H_s(\tilde{V}_{i+1}^-) \approx H_s(\tilde{V}_1^-) \approx H_s(t^{-1}\tilde{V}_i^-) \approx H_s(\tilde{V}_i^-)$$

Note further that the R-module  $H_{r+1}(\tilde{V}_{i+1}^-, \tilde{V}^-)$  contains an element  $S'$ , such that the image of  $S'$  in  $H_{r+1}(\tilde{V}_{i+1}^-, \tilde{V}_1^-)$  equals  $H(\sigma')$  and the image of  $S'$  in  $H_{r+1}(t^{-1}\tilde{V}_i^-, \tilde{V}^-)$  equals  $\sigma$ . Indeed, the chains  $D \in Z_{r+1}(t^{-1}\tilde{V}^-, \tilde{V}_1^-)$  and  $\bar{\sigma} \in Z_{r+1}(t^{-1}\tilde{V}_1^-, \tilde{V}^-)$  are homologous modulo  $\tilde{V}_1^-$ , i.e.  $D = \bar{\sigma} + R + \partial u$ , where  $R \in C_{r+1}(\tilde{V}_1^-)$ . Now set  $S' = D - R$ .

Consider the exact sequence of the triple  $(\tilde{V}_{i+1}^-, \tilde{V}_1^-, \tilde{V}^-)$ :

$$\begin{aligned} &\dots \rightarrow H_{r+2}(\tilde{V}_{i+1}^-, \tilde{V}_1^-) \rightarrow H_{r+1}(\tilde{V}_1^-, \tilde{V}^-) \rightarrow \\ &\rightarrow H_{r+1}(\tilde{V}_{i+1}^-, \tilde{V}^-) \xrightarrow{P} H_{r+1}(\tilde{V}_{i+1}^-, \tilde{V}_1^-) \rightarrow H_r(\tilde{V}_1^-, \tilde{V}^-) \rightarrow \dots \\ &\hspace{15em} \approx \text{?} \\ &\hspace{15em} R(\sigma) \end{aligned}$$

The first module from the right and the first module from the left vanish. Thus

$$\begin{aligned} H_{r+1}(\tilde{V}_{i+1}^-, \tilde{V}^-) &\approx H_{r+1}(\tilde{V}_1^-, \tilde{V}^-) \oplus R(\sigma) \approx \\ &\approx H_{r+1}(\tilde{V}_i^-, \tilde{V}^-) \oplus R(\sigma) \end{aligned}$$

Now we extend the identification  $F_i = H_{r+1}(\tilde{V}_1^-, \tilde{V}^-)$  to an isomorphism of the R-modules  $\varphi : F_{i+1} \rightarrow H_{r+1}(\tilde{V}_{i+1}^-, \tilde{V}^-)$  by setting  $\varphi(e_{i+1}) = S$ . Since  $tS = x$  in  $H_{r+1}(\tilde{V}_1^-, \tilde{V}^-) = F_i$ , the map  $\varphi$  commutes with the action of  $t$ ; since the cycle  $S$  is homologous to  $y$  modulo  $\tilde{V}^-$ , the composition of  $\varphi$  and the map  $H_{r+1}(\tilde{V}_{i+1}^-, \tilde{V}^-) \rightarrow H_{r+1}(\tilde{M}, \tilde{V}^-)$  equals  $P_{i+1} : F_{i+1} \rightarrow P_{i+1}$ .

The induction step is over.

Now we show how to realize geometrically the relation 2) of the definition of  $\mathcal{C}(R)$ . More precisely, suppose that the assumptions of lemma 7.5 hold. Let  $(V, \nu)$  be an  $(r-1)$ -biregular splitting, such that  $c(V, \nu) = (F_1, f_1)$ . Let

$$0 \rightarrow (F_2, f_2) \rightarrow (F_1, f_1) \rightarrow (F_0, f_0) \rightarrow 0$$

be an exact sequence of objects from  $\mathcal{C}(R)$ . We'll show that there exists an  $(r-1)$ -biregular splitting  $(V', \nu')$ , such that  $c(V', \nu') = (F_2 \oplus F_0, f_2 \oplus f_0)$ .

Let

$$0 \rightarrow (P_0, \varphi_0) \rightarrow (Q_0, \psi_0) \rightarrow (F_0, f_0) \rightarrow 0,$$

$$0 \rightarrow (P_2, \varphi_2) \rightarrow (Q_2, \psi_2) \rightarrow (F_2, f_2) \rightarrow 0$$

be the exact sequences of objects from  $\mathcal{C}(R)$ , where  $(Q_0, \psi_0)$ ,  $(Q_2, \psi_2)$  are triangular (such sequences exist, see [10, lemma 1.2]). Since  $F_0, F_2$  are free we have  $F_1 = F_0 \oplus F_2$  and  $f_1$  is given in this representation by the matrix  $\begin{pmatrix} f_0 & g \\ 0 & f_2 \end{pmatrix}$ , where  $g$  is a homomorphism  $F_2 \rightarrow F_0$ . This enables us to construct

the following exact sequences of objects of  $\mathcal{C}(R)$ :

$$0 \rightarrow (P_2 \oplus P_0, \varphi_2 \oplus \varphi_0) \rightarrow (Q_2 \oplus Q_0, \psi_2 \oplus \psi_0) \rightarrow (9.3) \\ \rightarrow (F_2 \oplus F_0, f_2 \oplus f_0) \rightarrow 0,$$

$$0 \rightarrow (P_2 \oplus P_0, \varphi_2 \oplus \varphi_0) \rightarrow (Q_2 \oplus Q_0, \gamma) \rightarrow (F_2 \oplus F_0, f_1) \rightarrow 0$$

where the middle objects are both triangular. The details (in slightly different notations) can be found in [10, p. 338].

Next we apply lemma 9.7 and find an  $r$ -biregular manifold

$(V_0, \nu_0)$ , such that  $c(V_0, \nu_0) = (P_2 \oplus P_0, \varphi_2 \oplus \varphi_0)$ . The manifold  $(V_0, -\nu_0)$  is an  $(n-r-3)$ -biregular splitting and

$$c(V_0, -\nu_0) = (P_2^* \oplus P_0^*, \pm(\varphi_2 \oplus \varphi_0)^*) \quad (\text{by lemma 9.6}).$$

Since  $r \geq 2$  and  $(n-r-3) + 1 \leq n-4$  we can apply lemma 9.7

again, this time - to the exact sequence, dual to the first

sequence from (9.3). Now we get a manifold  $(V_1, \nu_1)$ , such

that  $c(V_1, \nu_1) = (F_2 \oplus F_0, f_2 \oplus f_0)^*$ , and one more applica-

tion of lemma 9.6 completes the proof.

References

1. S.P. Novikov, Multivalued functions and functionals. An analogue of the Morse theory, Dokl. Akad. Nauk SSSR 260 (1981), 31-35; English transl. in Soviet Math. Dokl. 24 (1981).
2. S.P. Novikov. The Hamiltonian formalism and a multivalued analogue of Morse theory, Uspekhi Mat. Nauk 37 (1982), No.5 (227), 3-49; English transl. in Russian Math. Surveys 37 (1982).
3. S. Smale, On the structure of manifolds, Amer. J. Math. 84 (1962), 387-399.
4. M.Sh. Farber, The exactness of Novikov inequalities, Funktsional. Anal. i Prilozhen. 19 (1985) No.1, 49-59; English transl. in Functional Anal. Appl. 19 (1985).
5. W. Browder and J.P. Levine, Fiberings manifolds over  $S^1$ , Comment. Math. Helv. 40 (1966), 153-160.
6. A.V. Pazhitnov, On the exactness of Novikov type inequalities for the case  $\pi_1 M^n = \mathbb{Z}^m$  and Morse forms within a generic cohomology class, Dokl. Akad. Nauk SSSR 306 (1989), 544-548; English transl. in Soviet Math. Dokl.
7. W.P. Thurston, A norm on the homology of 3-manifolds, Memoirs of the A.M.S., 339 (1986), 99-130.
8. V.V. Sharko, K-theory and Morse theory I, Preprint N° 86.39, Institute of Math. AN USSR, 1986 (Russian).
9. V.V. Sharko, K-theory and Morse theory II, Preprint N° 86.40, Institute of Math. AN USSR, 1986 (Russian).
10. F.T. Farrell, The Obstruction to Fiberings a Manifold over a Circle, Indiana Univ. Math. Jour., 21 (1971), 315-346.

11. N. Bourbaki, *Algebre commutative*, Hermann, 1961.
12. G. Polya and G. Szegö, *Aufgaben und lehrsätze aus der analysis, Zweiterband*, Springer-Verlag, 1964.
13. S.P. Novikov, Bloch homology, critical points of functions and closed 1-forms, *Dokl. Akad. Nauk SSSR*, 287 (1986), 1321-1324; English transl. in *Soviet Math. Dokl.* 33 (1986).
14. A.V. Pazhitnov, An analytic proof of the real part of Novikov's inequalities, *Dokl. Akad. Nauk SSSR*, 293 (1987), 1305-1307; English transl. in *Soviet Math. Dokl.* 35 (1987).
15. W. Cockroft and R. Swan, On the homotopy type of certain two-dimensional complexes, *Proc. London Math. Soc.*, 11 (1961), 306-311.
16. A.A. Suslin, Projective modules over polynomial rings are free, *Dokl. Akad. Nauk*, 229 (1976), 1063-1066; (Russian).
17. D. Quillen, Projective modules over polynomial rings, *Invent. Math.*, 36 (1976), 167-171.
18. M. Kumar, Stably free modules, *Amer. J. Math.*, 107 (1985), 1439-1443.
19. G. Levitt, 1-formes fermées singulieres et groupe fondamental, *Invent. Math.*, 88 (1987), 635-667.
20. J.F.P. Hudson, Embedding of bounded manifolds, *Proc. Camb. Phil. Soc.*, 72 (1972), 11-20.

Institute of chemical physics  
AN USSR, 117977,  
Moscow, Kosygin str., 4.

Received 23/Nov./88.