COMPLEXIFICATION OF VIRO THEOREM AND TOPOLOGY OF REAL COMBINATORIAL HYPERSURFACES

Ilia Itenberg
IRMAR, Université de Rennes 1
Campus de Beaulieu
F-35042 Rennes cedex
France

Eugenii Shustin*
Tel Aviv University
School of Mathematical Sciences
Ramat Aviv, 69978 Tel Aviv
Israel

Abstract

We generalize the Viro construction of real algebraic hypersurfaces, based on the use of convex subdivisions of Newton polyhedra, to arbitrary subdivisions and show that the combinatorial hypersurfaces appearing from arbitrary subdivisions satisfy the same topological restrictions (congruences, inequalities etc.) as algebraic varieties. Our construction can be seen as a complex version of the Viro gluing theorem.

Introduction

The Viro method of gluing of polynomials appeared to be the most powerful construction of real algebraic varieties with prescribed topology [32, 33, 36] (see also [13], 11.5, [20, 23]). It provides a nice interaction of real algebraic geometry, toric geometry and combinatorics, and gives rise to various generalizations and applications [4, 6, 19, 28, 29, 30, 31].

*A part of the present work was done during the stay of the second author at the Fields Institute, Toronto, and at the NSF Science and Technology Research Center for the Computation and Visualization of Geometric Structures, funded by NSF/DMS89-20161. The work was completed during the stay of both authors at Max-Planck Institut für Mathematik. The authors thank these funds and institutions for hospitality and financial support.

Consider the simplest example of Viro's construction. Let $T_d \subset \mathbb{R}^2$, $d \in \mathbb{N}$, be the triangle with vertices $(0,0)$, $(0,d)$, $(d,0)$,

$$\tau: \quad T_d = \Delta_1 \cup ... \cup \Delta_N$$

be a triangulation with the set of vertices $V \subset \mathbb{Z}^2$, and $\sigma : V \to \{\pm 1\}$ be any function. Out of this combinatorial data we construct piecewise-linear plane curves. Denote by $T_d^{(1)}$, $T_d^{(2)}$ and $T_d^{(3)}$ the copies of T_d under reflections with respect to the coordinate axes and the origin. Take in $T_d^{(1)}$, $T_d^{(2)}$ and $T_d^{(3)}$ triangulations symmetric to τ, and define σ at the vertices of new triangulations by

$$\sigma(\varepsilon_1 i, \varepsilon_2 j) = \varepsilon_1^i \varepsilon_2^j \sigma(i,j), \quad (i,j) \in V, \quad \varepsilon_1, \varepsilon_2 = \pm 1.$$

Now in any triangle of the triangulation of $T_d \cup T_d^{(1)} \cup T_d^{(2)} \cup T_d^{(3)}$, having vertices with different values of σ, we draw the mean line separating the vertices with different signs. The union $C(\tau, \sigma)$ of all these mean lines is a broken line homeomorphic to a disjoint union of circles and segments. Introduce natural maps:

$$\Phi : T_d \cup T_d^{(1)} \cup T_d^{(2)} \cup T_d^{(3)} \to \mathbb{R}P^2, \quad \Psi : \mathrm{Int}(T_d \cup T_d^{(1)} \cup T_d^{(2)} \cup T_d^{(3)}) \to \mathbb{R}^2,$$

where Φ is continuous onto, identifying antipodal points on $\partial(T_d \cup T_d^{(1)} \cup T_d^{(2)} \cup T_d^{(3)})$, and Ψ is a homeomorphism. We call the curves $\Phi(C(\tau,\sigma)) \subset \mathbb{R}P^2$, $\Psi(C(\tau,\sigma)) \subset \mathbb{R}^2$ *projective* and *affine T-curves of degree d.*

The Viro theorem states that a projective (resp., affine) T-curve of degree d is isotopic in $\mathbb{R}P^2$ (resp., in $\mathbb{R}^2$) to a nonsingular algebraic projective (resp., affine) curve of degree d, providing that the triangulation τ is *convex*, *i.e.*, there exists a convex piecewise-linear function $\nu : T_d \to \mathbb{R}$, whose linearity domains are $\Delta_1, \dots, \Delta_N$ (sometimes such triangulations are called *coherent*; see [13]). The Viro theorem, in fact, endows the combinatorial broken line $C(\tau,\sigma)$ with a rich structure, which implies a number of restrictions on the topology of $C(\tau,\sigma)$ (see an account of known results in [25, 34, 35, 38]).

On the other hand, there are *non-convex* triangulations (see the simplest classical example in Figure 1 [7]). There are examples of T-curves beyond the range of known algebraic curves [26], and there is some similarity between T-curves and algebraic curves: up to degree 6 all projective T-curves are isotopic to algebraic ones of the same degree [9], T-curves satisfy some consequences of the Bezout theorem [9] and the Harnack inequality [18, 15]. The natural question arising is if any T-curve is isotopic to an algebraic curve of the same degree, and if not, how far T-curves may differ from algebraic curves.

In the present paper, we propose a complex version of the general Viro construction. This version uses not necessarily convex subdivisions of Newton polyhedra. The construction produces PL-hypersurfaces. We call them *C-hypersurfaces.* They share a lot of topological properties with algebraic hypersurfaces. In particular, for any T-curve, we construct its *complexification*, *i.e.*, an equivariant surface in $\mathbb{C}P^2$ or $\mathbb{C}^2$, whose real part coincides with the given T-curve, and show that an equivariant smoothing of this complexification is a *flexible curve* in the sense of [35]. As a



Figure 1: A non-convex triangulation

corollary, we obtain that arbitrary T-curves satisfy all *topological* restrictions known for real algebraic curves.

In section 1 we describe the construction of complex and real C-hypersurfaces. In section 2 we study the topological properties of complex C-hypersurfaces of degree d in $\mathbb{C}P^n$ (*i.e.*, constructed using a subdivision of the simplex in $\mathbb{R}^n$ with vertices $(0,0,...,0)$, $(d,0,...,0)$, ..., $(0,...,0,d)$) and of double coverings of $\mathbb{C}P^n$ ramified along complex C-hypersurfaces. We prove the following statements.

- A C-hypersurface of degree d in $\mathbb{C}P^n$ is an orientable manifold, homologous to an algebraic hypersurface of degree d in $\mathbb{C}P^n$.
- Any C-hypersurface M in $\mathbb{C}P^n$ is isotopic to a close smooth manifold M_{sm} of codimension 2 in $\mathbb{C}P^n$. If M is real, *i.e.* invariant under the complex conjugation in $\mathbb{C}P^n$, the isotopy can be made equivariant.
- Given a (real) C-hypersurface M, the tangent bundle to its smoothing M_{sm} is (equivariantly) isotopic to a (equivariant) bundle of complex hyperplanes. In particular, M_{sm} possesses an (equivariant) almost complex structure.
- A C-hypersurface in $\mathbb{C}P^n$ is simply connected if $n > 2$.
- For a C-hypersurface of degree d in $\mathbb{C}P^n$ one has $\pi_1(\mathbb{C}P^n \backslash M) = \mathbb{Z}/d\mathbb{Z}$ if $n \geq 2$.
- Let $M \subset \mathbb{C}P^n$ be a C-hypersurface and $M' \subset \mathbb{C}P^n$ a nonsingular algebraic hypersurface, both of degree d. Then $H_*(M)$ and $H_*(M')$ are isomorphic as graded groups.
- Let n be a positive odd number, M a C-hypersurface of degree d in $\mathbb{C}P^n$ and M' a nonsingular algebraic hypersurface of degree d in $\mathbb{C}P^n$. Then the lattices

$(H_{n-1}(M), B_M)$ and $(H_{n-1}(M'), B_{M'})$ (where $B_M : H_{n-1}(M) \times H_{n-1}(M) \to \mathbb{Z}$ and $B_{M'} : H_{n-1}(M') \times H_{n-1}(M') \to \mathbb{Z}$ are the intersection forms of M and M', respectively) are isomorphic.

- A C-surface of degree d in $\mathbb{C}P^3$ is homeomorphic to a nonsingular algebraic surface of degree d in $\mathbb{C}P^3$.
- Any C-curve in $\mathbb{C}P^2$ is isotopic to an algebraic curve of the same degree.

Section 3 is devoted to topology of real C-hypersurfaces in $\mathbb{C}P^n$. We prove for them the generalized Harnack inequality and the Gudkov-Rokhlin and Gudkov-Krahnov-Kharlamov congruences. For real C-surfaces in $\mathbb{C}P^3$ we also prove inequalities similar to the Comessatti ones.

1 Construction of complex and real C-hypersurfaces

1.1 Notations and definitions

Further on the term *polyhedron* (*polygon*) means a convex polyhedron (polygon) in $\mathbb{R}^n$, $n \geq 2$, whose vertices have nonnegative integral coordinates.

Given a polynomial

$$F = \sum_{i_1,...,i_n} A_{i_1...i_n} z_1^{i_1} \cdot ... \cdot z_n^{i_n} \ ,$$

by $\Delta(F)$ we denote its Newton polyhedron, *i.e.*, the convex hall of the set

$$\{(i_1, ..., i_n) \in \mathbb{R}^n \ : \ A_{i_1...i_n} \neq 0\} \ .$$

The truncation of F on a face δ of $\Delta(F)$ is the polynomial

$$F^\delta = \sum_{(i_1,...,i_n)\in\delta} A_{i_1...i_n} z_1^{i_1} \cdot ... \cdot z_n^{i_n} \ .$$

A polynomial $F \in \mathbb{C}[z_1, ..., z_n]$ is called *nondegenerate*, if F and any truncation F^δ on a proper face δ of $\Delta(F)$ have a nonsingular zero set in $(\mathbb{C}^*)^n$ (cf. [32]).

1.2 Complexification of the momentum map

Let $\Delta \subset \mathbb{R}^n$ be a polyhedron, and let $\mu_\Delta : (\mathbb{R}^*)^n \to I(\Delta)$ be the momentum map (see [1, 2, 12], [13], 6.1), where $I(\Delta)$ is the complement in Δ of the union of all its proper faces,

$$\mu_\Delta(x_1, ..., x_n) = \frac{\sum_{(i_1,...,i_n)\in\Delta} x_1^{i_1}...x_n^{i_n} \cdot (i_1, ..., i_n)}{\sum_{(i_1,...,i_n)\in\Delta} x_1^{i_1}...x_n^{i_n}} \ . \tag{1}$$

Split the complex torus in the product $(\mathbb{C}^*)^n = (\mathbb{R}_+^*)^n \times (S^1)^n$:

$$(z_1, ..., z_n) \in (\mathbb{C}^*)^n \ \mapsto \ (|z_1|, ..., |z_n|) \in (\mathbb{R}_+^*)^n, \ \left(\frac{z_1}{|z_1|}, ..., \frac{z_n}{|z_n|}\right) \in (S^1)^n \ .$$

Put $\mathbb{C}(I(\Delta)) = I(\Delta) \times (S^1)^n \subset (\mathbb{C}^*)^n$, and define the *complexification* $\mathbb{C}\Delta$ *of* Δ to be the closure of $\mathbb{C}I(\Delta)$ in $\mathbb{C}^n$.

Proposition 1.1 *The complexification $\mathbb{C}\Delta$ of Δ is a singular PL-manifold with boundary. The singular set of $\mathbb{C}\Delta$ is the union of $\mathbb{C}\delta$ over all faces $\delta \subset \Delta$, which are intersections of Δ with coordinate planes of dimension $> n - \dim\Delta + \dim\delta$. The real part $\mathbb{R}\Delta$ of $\mathbb{C}\Delta$ is the union of Δ with all its symmetric copies with respect to the coordinate hyperplanes.*

Proof. Straightforward. $\Box$

Define *the complex momentum map* $\mathbb{C}\mu_\Delta : (\mathbb{C}^*)^n \to \mathbb{C}(I(\Delta))$ by

$$\mathbb{C}\mu_\Delta(x_1v_1, ..., x_nv_n) = (\mu_\Delta(x_1, ..., x_n), (v_1, ..., v_n)) \in I(\Delta) \times (S^1)^n = \mathbb{C}(I(\Delta)), \quad (2)$$
$$(x_1, ..., x_n) \in (\mathbb{R}_+^*)^n, \ (v_1, ..., v_n) \in (S^1)^n \ .$$

As an easy consequence of classical results we obtain the following statement.

Proposition 1.2 *The map $\mathbb{C}\mu_\Delta$ is surjective and commutes with the complex conjugation* Conj. *It is a diffeomorphism when* $\dim\Delta = n$. *The real part of* $\mathbb{C}(I(\Delta))$ *is the image of* $(\mathbb{R}^*)^n$.

Below we use the following interaction of the complex momentum map with the action of $\mathbb{Z}^n$ in $\mathbb{R}^n$

$$\overline{a} = (a_1, ..., a_n) \in \mathbb{Z}^n, \ \overline{t} = (t_1, ..., t_n) \in \mathbb{R}^n \quad \mapsto \quad \overline{a} + \overline{t} = (t_1 + a_1, ..., t_n + a_n) \in \mathbb{R}^n \ ,$$

and the actions of $SL(\mathbb{Z}, n)$ in $\mathbb{R}^n$ and $(\mathbb{R}^*)^n$

$$A \in SL(\mathbb{Z}, n), \ \overline{x} \in \mathbb{R}^n \quad \mapsto \quad A\overline{x} \in \mathbb{R}^n \ ,$$

$$A \in SL(\mathbb{Z}, n), \ \overline{x} = (x_1, ..., x_n) \in (\mathbb{R}^*)^n \ \mapsto \ \overline{x}^A = \left(\prod_{i=1}^n x_i^{a_{i1}}, ..., \prod_{i=1}^n x_i^{a_{in}}\right) \ .$$

Proposition 1.3 *Let $\Delta \subset \mathbb{R}^n$ be a polygon. If $\overline{a} = (a_1, ..., a_n) \in \mathbb{Z}^n$, $A \in SL(\mathbb{Z}, n)$, and $\overline{a} + A\Delta$ lies in the nonnegative orthant $\mathbb{R}_+^n$, then*

$$\mathbb{C}\mu_{\overline{a}+A\Delta}(\overline{z}) = (\overline{a} + A\mu_\Delta(\overline{x}^A), \overline{v}) \ , \quad (3)$$

where $\overline{z} = (\overline{x}, \overline{v}) \in (\mathbb{R}_+^*)^n \times (S^1)^n$.

Proof. The result follows from the definition

$$\mathbb{C}\mu_{\overline{a}+A\Delta}(\overline{z}) = (\mu_{\overline{a}+A\Delta}(\overline{x}), \overline{v}) \ ,$$

and the classically known relation

$$\mu_{\overline{a}+A\Delta}(\overline{x}) = \overline{a} + A\mu_\Delta(\overline{x}^A) \quad \Box$$

1.3 Real and complex chart of a polynomial

Let $F \in \mathbb{C}[z_1, ..., z_n]$ be a nondegenerate polynomial with Newton polyhedron $\Delta(F) = \Delta$. The closure $\mathbb{C}Ch(F) \subset \mathbb{C}\Delta$ of the set $\mathbb{C}\mu_\Delta(\{F = 0\} \cap (\mathbb{C}^*)^n)$ is called the *(complex) chart* of the polynomial F. If F is real then $\mathbb{R}Ch(F) = \mathbb{C}Ch(F) \cap \mathbb{R}\Delta$ is called the *real chart* of F.

This definition is a modification of the classical one [23, 32] (cf. [27]).

Proposition 1.4 *Suppose that Δ belongs to the open positive orthant $(\mathbb{R}^*_+)^n = \{(x_1, ..., x_n) \in \mathbb{R}^n : x_1 > 0, \ldots, x_n > 0\}$. Then the set $\mathbb{C}Ch(F)$ is a PL-submanifold in $\mathbb{C}\Delta$ of codimension 2 with boundary $\partial\mathbb{C}Ch(F) = \mathbb{C}Ch(F) \cap \partial\mathbb{C}\Delta$. It is smooth in $\mathbb{C}I(\Delta)$, and, for any proper face δ of Δ of positive dimension,*

$$\mathbb{C}Ch(F) \cap \mathbb{C}\delta = \mathbb{C}Ch(F^\delta) \ . \tag{4}$$

If F is real then $\mathbb{C}Ch(F)$ is invariant with respect to Conj*, and $\mathbb{R}Ch(F)$ is a PL-submanifold in $\mathbb{R}\Delta$ of codimension 1 with boundary $\partial\mathbb{R}Ch(F) = \mathbb{R}Ch(F) \cap \partial\mathbb{R}\Delta$.*

Remark 1.5 *If Δ intersects with coordinate hyperplanes, then the statement of Proposition 1.4 holds true when substituting $\mathbb{C}\Delta \backslash \mathrm{Sing}(\mathbb{C}\Delta)$ and $\mathbb{R}\Delta \backslash \mathrm{Sing}(\mathbb{C}\Delta)$ for $\mathbb{C}\Delta$ and $\mathbb{R}\Delta$, respectively, but we will not use this below.*

Proof of Proposition 1.4.

(i) If $\dim \Delta = n$ then by Proposition 1.2, $\mathbb{C}Ch(F) \cap \mathbb{C}(I(\Delta))$ is a smooth manifold of codimension 2.

So, we have to consider only the behavior of $\mathbb{C}Ch(F)$ on $\partial\mathbb{C}\Delta$.

(ii) Assume that $\dim \Delta < n$ and $A \in SL(\mathbb{Z}, n)$ is such that $A\Delta \subset (\mathbb{R}^*_+)^n$ is parallel to the coordinate hyperplane $i_n = 0$. By (3) there is a diffeomorphism which takes the pair $(\mathbb{C}\Delta, \mathbb{C}Ch(F(\overline{z})))$ onto the pair $(\mathbb{C}(A\Delta), \mathbb{C}Ch(F(\overline{z}^A)))$. Now, if $i_n\big|_\Delta = a$, then $\Delta' = \Delta - (0, ..., 0, a) \subset \{i_n = 0\}$, $F(\overline{z}) = z_n^a F'(\overline{z})$, and by formula (3) the pair $(\mathbb{C}\Delta, \mathbb{C}Ch(F))$ naturally splits into the product $(C\Delta', \mathbb{C}Ch(F')) \times S^1$, thereby one reduces the dimension of the ambient space.

(iii) To prove (4) (for $\dim \Delta = n$), we note that any point $\overline{z}^0 \in \mathbb{C}Ch(F) \cap \partial\mathbb{C}\Delta$ is a limit of $\mathbb{C}\mu_\Delta(\gamma(t))$ as $t > 0$, $t \to 0$, where

$$\gamma(t) = (\lambda_1 t^{k_1} + O(t^{k_1+1}), ..., \lambda_n t^{k_n} + O(t^{k_n+1}))$$

is a curve lying in $\{F = 0\} \cap (\mathbb{C}^*)^n$ with $\lambda_1 ... \lambda_n \neq 0$, $(k_1, ..., k_n) \in \mathbb{Z}^n \backslash \{0\}$. Let δ be the maximal (proper) face of Δ, where the function $(i_1, ..., i_n) \in \Delta \mapsto i_1k_1 + ... + i_nk_n$ achieves its minimum (equal to l). Then

$$0 = F(\gamma(t)) = F^\delta(\lambda_1, ..., \lambda_n)t^l + O(t^{l+1}) \quad \Longrightarrow \quad F^\delta(\lambda_1, ..., \lambda_n) = 0 \ ,$$

$\gamma(t) = (\overline{x}, \overline{v}) \in (\mathbb{R}^*_+)^n \times (S^1)^n$, where

$$\overline{x} = (|\lambda_1|t^{k_1} + O(t^{k_1+1}), ..., |\lambda_n|t^{k_n} + O(t^{k_n+1})), \ \overline{v} = \left(\frac{\lambda_1}{|\lambda_1|} + O(t), ..., \frac{\lambda_n}{|\lambda_n|} + O(t)\right) .$$

So, one derives

$$\mathbb{C}\mu_\Delta(\gamma(t)) = (\mu_\Delta(\overline{x}), \overline{v}) = \left(\frac{\sum_{(i_1,...,i_n)\in\Delta}(i_1, ..., i_n)x_1^{i_1}...x_n^{i_n}}{\sum_{(i_1,...,i_n)\in\Delta} x_1^{i_1}...x_n^{i_n}},\ \overline{v} \right)$$

$$= \left(\frac{t^l \cdot \sum_{(i_1,...,i_n)\in\delta}(i_1, ..., i_n)|\lambda_1|^{i_1}...|\lambda_n|^{i_n} + O(t^{l+1})}{t^l \cdot \sum_{(i_1,...,i_n)\in\delta} |\lambda_1|^{i_1}...|\lambda_n|^{i_n} + O(t^{l+1})},\ \overline{v} \right) \quad \overset{t\to 0}{\longrightarrow} \quad \mathbb{C}\mu_\delta(\lambda_1, ..., \lambda_n).$$

Hence $\mathbb{C}Ch(F) \cap \mathbb{C}\delta \subset \mathbb{C}Ch(F^\delta)$.

Now let $(\lambda_1, ..., \lambda_n) \in (\mathbb{C}^*)^n$, $F^\delta(\lambda_1, ..., \lambda_n) = 0$, where δ is a proper face of Δ, and let $(k_1, ..., k_n) \in K(\delta)$, where $K(\delta) \subset \mathbb{Z}^n \backslash \{0\}$ consists of vectors $(k_1, ..., k_n)$ such that δ is the maximal face of Δ on which the functional $(i_1, ..., i_n) \in \Delta \mapsto i_1k_1 + ... + i_nk_n$ achieves its minimum. Then $F^\delta(\lambda_1 t^{k_1}, ..., \lambda_n t^{k_n}) = t^l F^\delta(\lambda_1, ..., \lambda_n) = 0$, $l = i_1k_1 + ... + i_nk_n$, $(i_1, ..., i_n) \in \delta$, and there exist smooth functions $\varphi_1(t) = \lambda_1 + O(t)$, ..., $\varphi_n = \lambda_n + O(t)$ such that the curve $\gamma(t) = (t^{k_1}\varphi_1(t), ..., t^{k_n}\varphi_n(t))$ lies on $\{F = 0\}$. Indeed,

$$F(\gamma(t)) = t^\lambda F^\delta(\varphi_1, ..., \varphi_n) + t^{\lambda+1}G(t, \varphi_1, ..., \varphi_2) = 0$$

$$\iff \quad F^\delta(\varphi_1, ..., \varphi_n) + tG(t, \varphi_1, ..., \varphi_2) = 0\ ,$$

hence the existence of $\varphi_1, ..., \varphi_n$ defined for small $t > 0$ follows from the implicit function theorem and the fact that $F^\delta = 0$ is nonsingular in $(\mathbb{C}^*)^n$. The above argument shows that

$$\mathbb{C}\mu_\delta(\lambda_1, ..., \lambda_n) = \lim_{t\to 0} \mathbb{C}\mu_\Delta(\gamma(t))\ .$$

(iv) Assume that $\dim \Delta = n$. Let $\delta \subset \Delta$ be a proper face, $\dim \delta = s > 0$, and $\overline{w} = (y_1v_1^0, ..., y_nv_n^0) \in \mathbb{C}I(\delta) \cap \mathbb{C}Ch(F)$, where $(y_1, ..., y_n) \in I(\delta)$, $(v_1^0, ..., v_n^0) \in (S^1)^n$. We will describe $\mathbb{C}Ch(F)$ in a neighborhood $U_{\overline{w}}$ of the point $\overline{w}$ in $\mathbb{C}\Delta$.

Represent $\mathbb{R}^n$ as $\mathbb{R}^s \times \mathbb{R}^{n-s}$ and denote by $\mathrm{pr}_s : \mathbb{R}^n \to \mathbb{R}^s$, $\mathrm{pr}_{n-s} : \mathbb{R}^n \to \mathbb{R}^{n-s}$ the natural projections. According to Proposition 1.3, we can assume that $\mathrm{pr}_{n-s}(\delta)$ is a point $(p_{s+1}, ..., p_n)$ with positive integers $p_{s+1}, ..., p_n$, so $\widetilde{\delta} = \mathrm{pr}_s(\delta) = \delta - (0, ..., 0, p_{s+1}, ..., p_n)$, and $\widetilde{\Delta}\backslash\widetilde{\delta}$ lies in the positive orthant. Then $F(z_1, ..., z_n) = z_{s+1}^{p_{s+1}}...z_n^{p_n}\widetilde{F}(z_1, ..., z_n)$, where

$$\widetilde{F}(z_1, ..., z_n) = \widetilde{F}^{\widetilde{\delta}}(z_1, ..., z_s) + \sum_{(i_1,...,i_n)\in\Lambda} A_{i_1...i_n} z_1^{i_1}...z_s^{i_s} \cdot M_{i_{s+1}...i_n}\ , \tag{5}$$

$$\Lambda = \{(i_1, ...i_n) \in \widetilde{\Delta}\ :\ i_{s+1}, ..., i_n > 0\} \cap \mathbb{Z}^n, \quad M_{i_{s+1}...i_n} = z_{s+1}^{i_{s+1}}...z_n^{i_n}\ .$$

Note that there are uniquely defined $x_1^0, ..., x_s^0 > 0$ such that $(y_1, ..., y_s) = \mu_{\widetilde{\delta}}(x_1^0, ..., x_s^0)$. The condition on $\overline{z} = (\overline{x}, \overline{v}) \in (\mathbb{R}_+^*)^n \times (S^1)^n = (\mathbb{C}^*)^n$ such that $\mathbb{C}\mu_\Delta(\overline{x}, \overline{v}) \in U_{\overline{w}}$ can be expressed as follows. For $\overline{v} = (v_1, ..., v_n)$ it, clearly, is

$$|v_1 - v_1^0| < \varepsilon,\ ...,\ |v_n - v_n^0| < \varepsilon, \tag{6}$$

where $\varepsilon > 0$ is sufficiently small. For $\overline{x} = (x_1, ..., x_n)$ we have that $\mu_{\widetilde{\Delta}}(\overline{x})$, equal to

$$\frac{\sum_{(i_1,...,i_s)\in\widetilde{\delta}} x_1^{i_1}...x_s^{i_s} \cdot (i_1, ..., i_s, 0, ..., 0) + \sum_{(i_1,...,i_n\in\Lambda} x_1^{i_1}...x_s^{i_s} \cdot |M_{i_{s+1}...i_n}| \cdot (i_1, ..., i_n)}{\sum_{(i_1,...,i_s)\in\widetilde{\delta}} x_1^{i_1}...x_s^{i_s} + \sum_{(i_1,...,i_n\in\Lambda} x_1^{i_1}...x_s^{i_s} \cdot |M_{i_{s+1}...i_n}|} ,$$

is close to $(y_1, ..., y_s, 0, ..., 0)$. This means that in the latest expression the summands over Λ are small with respect to the sums over $\widetilde{\delta}$; hence $\mathrm{pr}_s(\mu_{\widetilde{\Delta}}(\overline{x}))$ is close to $\mu_{\widetilde{\delta}}(x_1, ..., x_s)$, which means that $(x_1, ..., x_s)$ is close to $(x_1^0, ..., x_s^0)$. Coming back to the sums over Λ, we obtain that $M_{i_{s+1}...i_n}$ must be close to zero, $(i_{s+1}, ..., i_n) \in \mathrm{pr}_{n-s}(\Lambda)$. So, without loss of generality the conditions on $\overline{x}$ can be expressed as

$$|x_l - x_l^0| < \varepsilon,\ l = 1, ..., s, \quad x_{s+1}^{i_{s+1}}...x_n^{i_n} < \varepsilon,\ (i_{s+1}, ..., i_n) \in \mathrm{pr}_{n-s}(\Lambda) . \tag{7}$$

Let $U' \subset (\mathbb{R}_+^*)^n$ and $U'' \subset (S^1)^n$ be given by (7) and (6), respectively, and note that $U' = \mathrm{pr}_s(U') \times \mathrm{pr}_{n-s}(U')$. Put $\sigma = \mathrm{pr}_{n-s}(\widetilde{\Delta})$. Then μ_σ takes $\mathrm{pr}_{n-s}(U')$ onto a set $U_0\backslash\partial\sigma$, where U_0 is a neighborhood of the origin (a vertex of σ !) in σ.

Since $\widetilde{F}$ is nondegenerate, we can assume that

$$\frac{\partial\widetilde{F}}{\partial x_1}(x_1^0, ..., x_s^0, 0, ..., 0) = \frac{\partial\widetilde{F}^{\widetilde{\delta}}}{\partial x_1}(x_1^0, ..., x_s^0) \neq 0 .$$

Hence, by (5) and the implicit function theorem, the hypersurface $\{F = 0\}$ can be described in $U' \times U''$ (with, possibly, smaller ε) by equations

$$x_1 = x_1^0 + G, \quad v_1 = v_1^0 + H , \tag{8}$$

where G and H are vanishing at zero smooth functions of $x_2 - x_2^0$, ..., $x_s - x_s^0$, $v_2 - v_2^0$, ..., $v_n - v_n^0$, and $M_{i_{s+1}...i_n}$, $(i_{s+1}, ..., i_n) \in \mathrm{pr}_{n-s}(\Lambda)$, where $x_2, ..., x_n$, $v_2, ..., v_n$ are arbitrary satisfying (6) and (7).

Now we show that $U_{\overline{w}}$ is homeomorphic to $\mathrm{pr}_s(U') \times U_0 \times U''$ via an extension of $\mathbb{C}\mu_\Delta$. Indeed, $U_{\overline{w}}\backslash\partial\mathbb{C}\Delta$ is homeomorphic via $\mathbb{C}\mu_\Delta$ to $\mathrm{pr}_s(U') \times \mathrm{pr}_{n-s}(U') \times U'' \simeq \mathrm{pr}_s(U'') \times (U_0\backslash\partial\sigma) \times U''$. As we saw, the points in $U_{\overline{w}} \cap \partial\Delta$ are $\lim_{t\to 0} \mathbb{C}\mu_\Delta(\overline{x}(t), \overline{v}(t))$ for all curves $\overline{x}(t) \in U'$, $\overline{v}(t) \in U''$, $t > 0$, such that $\lim_{t\to 0} \overline{v}(t) \in U''$, $\lim_{t\to 0}(x_1(t), ..., x_s(t)) \in \mathrm{pr}_s(U')$ and

$$x_{s+1}(t) = \lambda_1 t^{k_1} + O(t^{k_1+1}),\ ...,\ x_n(t) = \lambda_{n-s} t^{k_{n-s}} + O(t^{k_{n-s}+1}) .$$

Restriction (7) means, first, that the vector $(k_1, ..., k_{n-s})$ belongs to the cone generated by the interior normal vectors to the $(s-1)$-dimensional faces of σ, which contain the origin, and, second, that $\lambda_1^{i_{s+1}}...\lambda_{n-s}^{i_n} < \varepsilon$ for all $(i_{s+1}, ..., i_n) \in \mathrm{pr}_{n-s}(\Lambda)$ orthogonal to $(k_1, ..., k_{n-s})$. On the other hand, the same data determine a point $\lim_{t\to 0} \mu_\sigma(x_{s+1}(t), ..., x_n(t)) \in U_0 \cap \partial\sigma$. This gives a one-to-one correspondence between $U_{\overline{w}} \cap \partial\Delta$ and $\mathrm{pr}_s(U') \times (U_0 \cap \partial\sigma) \times U''$, which respects the combinatorial structure: the interior of a face $\psi \subset U_{\overline{w}} \cap \partial\Delta$ corresponds to $\mathrm{pr}_s(U') \times \mathrm{pr}_{n-s}(I\widetilde{\psi}) \times U''$, where $\widetilde{\psi} = \psi - (0, ..., 0, p_{s+1}, ..., p_n)$. The above correspondence provides a homeomorphism of $U_{\overline{w}}$ and $\mathrm{pr}_s(U') \times U_0 \times U''$, since close points in $U_{\overline{w}} \cap \partial\Delta$ come from close curves $\gamma(t)$ which in turn bear close points in $\mathrm{pr}_s(U') \times (U_0 \cap \partial\sigma) \times U''$. Consequently, we can introduce in U_0 coordinates $\theta_1, ..., \theta_{n-s}$ such that

$$U_0 = \{0 \le \theta_1 < \varepsilon, \ -\varepsilon < \theta_j < \varepsilon, \ j = 2, ..., n-s\}, \quad U_0 \cap \partial\sigma = \{\theta_1 = 0\} \ ,$$

and $M_{i_{s+1}...i_n}$, $(i_{s+1}, ..., i_n) \in \mathrm{pr}_{n-s}(\Lambda)$, are continuous functions of $\theta_1, ..., \theta_{n-s}$, vanishing at zero. Thus, formulae (8) allow us to parameterize $\mathbb{C}Ch(F) \cap U_{\overline{w}}$ by the following subset in $\mathrm{pr}_s(U') \times U_0 \times U''$:

$$x_1 = x_1^0 + G', \quad v_1 = v_1^0 + H' \ ,$$

where G' and H' are vanishing at zero smooth functions of $x_2, ..., x_s, v_2, ..., v_n, \theta_1, ..., \theta_{n-s}$. This subset is a subvariety of codimension 2 with a boundary on $\mathrm{pr}_s(U') \times (U_0 \cap \partial\sigma) \times U''$. Hence $\mathbb{C}Ch(F) \cap U_{\overline{w}}$ is a subvariety of codimension 2 in $U_{\overline{w}}$ with boundary on $U_{\overline{w}} \cap \partial\Delta$.

(v) The statement of Lemma for $\mathbb{R}Ch(F)$ can be proven as in the complex case. $\square$

Corollary 1.6 *If* $\dim \Delta = 2$, $\Delta \subset \mathbb{R}^2$, *then* $\mathbb{C}Ch(F)$ *is a smooth surface with boundary.*

1.4 Digression: real resolution of toric varieties

Let $\Delta \subset (\mathbb{R}_+^*)^n$ be an n-dimensional polyhedron, F be a nondegenerate polynomial with Newton polyhedron Δ. Denote by $\mathrm{Tor}_{\mathbb{C}}(\Delta)$ the toric variety over $\mathbb{C}$ defined by Δ, and by $Z(F)$ the (closed) hypersurface in $\mathrm{Tor}_{\mathbb{C}}(\Delta)$ defined by F.

Proposition 1.7 *There exists an equivariant surjective map* $\nu : \mathbb{C}\Delta \to \mathrm{Tor}_{\mathbb{C}}(\Delta)$ *such that*

- $\nu(\mathbb{C}Ch(F)) = Z(F)$,
- $\nu\big|_{\mathbb{C}I(\Delta)}$ *takes* $\mathbb{C}I(\Delta)$ *diffeomorphically to* $(\mathbb{C}^*)^n \subset \mathrm{Tor}_{\mathbb{C}}(\Delta)$,
- *for any face* δ, $\dim \delta = s < n$, *the map* $\nu\big|_{\mathbb{C}I(\delta)}$ *is a submersion of* $\mathbb{C}I(\delta)$ *onto* $(\mathbb{C}^*)^s \subset \mathrm{Tor}_{\mathbb{C}}(\delta) \subset \mathrm{Tor}_{\mathbb{C}}(\Delta)$.

Proof. Let us represent $\mathrm{Tor}_{\mathbb{C}}(\Delta)$ as the closure of the variety

$$\{(z_1^{i_1}...z_n^{i_n})_{(i_1,...,i_n)\in\Delta\cap\mathbb{Z}^n} \ : \ (z_1, ..., z_n) \in (\mathbb{C}^*)^n\} \subset \mathbb{C}P^{N-1} \ ,$$

where $N = \#(\Delta \cap \mathbb{Z}^n)$ (see, for example [12]). We define $\nu\big|_{\mathbb{C}I(\Delta)} = (\mathbb{C}\mu_\Delta)^{-1}$, which is a required diffeomorphism of $\mathbb{C}I(\Delta)$ and $(\mathbb{C}^*)^n \subset \mathrm{Tor}_{\mathbb{C}}(\Delta)$, and extend it to $\partial\mathbb{C}\Delta$. Namely, given a face $\delta \subset \Delta$ and a point $\overline{w} = \mathbb{C}\mu_\delta(\overline{z}) \in \mathbb{C}I(\delta)$, $\overline{z} = (z_1, ..., z_n) \in (\mathbb{C}^*)^n$, we put $\nu(\overline{w}) = (a_{i_1...i_n})_{(i_1,...,i_n)\in\Delta\cap\mathbb{Z}^n}$, where

$$a_{i_1...i_n} = z_1^{i_1}...z_n^{i_n}, \ (i_1, ..., i_n) \in \delta, \quad a_{i_1...i_n} = 0, \ (i_1, ..., i_n) \in \Delta\backslash\delta \ .$$

Clearly, $\nu\big|_{\mathbb{C}I(\delta)}$ satisfies the required property. It remains only to explain that ν is continuous. Indeed, in the previous notation $\overline{w} = \lim_{t\to\infty} \mathbb{C}\mu_\Delta(\gamma(t))$, where

$$\gamma(t) = (z_1 t^{k_1} + O(t^{k_1-1}), ..., z_n t^{k_n} + O(t^{k_n-1})), \quad t > 0 \ ,$$

is a curve in $(\mathbb{C}^*)^n$ with $(k_1, ..., k_n) \in \mathbb{Z}^n\backslash\{0\}$ belonging to the dual cone of δ with respect to Δ. Then one can easily check that $\lim_{t\to\infty} \gamma(t) \in \mathbb{C}P^{N-1}$ is just $\nu(\overline{w})$ defined as above. $\Box$

1.5 Gluing of charts

Let $\mathcal{S}$ be a subdivision of a polyhedron $\Delta \subset \mathbb{R}^n$ into equidimensional polyhedra: $\Delta = \Delta_1 \cup ... \cup \Delta_N$ (*i.e.*, $\Delta_i \cap \Delta_j$ is empty or a common proper face), and let $\mathcal{A} = \{A_{\overline{i}},\ \overline{i} \in \Delta \cap \mathbb{Z}^n\}$ be a collection of complex numbers such that $A_{\overline{i}} \ne 0$ if $\overline{i}$ is a vertex of some Δ_i, $1 \le i \le N$. Further on, speaking on subdivisions of polyhedra and corresponding collections of numbers, we always assume the above properties.

Assume that the polynomials

$$F_k(z_1, ..., z_n) = \sum_{(i_1,...,i_n)\in\Delta_k} A_{i_1...i_n} z_1^{i_1} \cdot ... \cdot z_n^{i_n} \ , \quad k = 1, ..., N,$$

are nondegenerate. We define the *gluing* of the complex charts of $F_1, \dots, F_N$ (below referred to as a *C-hypersurface in* $\mathbb{C}\Delta$) by

$$\mathbb{C}Ch(\mathcal{S}, \mathcal{A}) = \bigcup_{k=1}^N \mathbb{C}Ch(F_k) \ .$$

If all the polynomials $F_1, \dots, F_N$ are real, we define the *gluing* of the real charts of $F_1, \dots, F_N$ (which is called a *C-hypersurface in* $\mathbb{R}\Delta$) by

$$\mathbb{R}Ch(\mathcal{S}, \mathcal{A}) = \bigcup_{k=1}^N \mathbb{R}Ch(F_k) \ .$$

Proposition 1.8 *Let $\Delta \subset (\mathbb{R}^*_+)^n$. Then the set $\mathbb{C}Ch(\mathcal{S}, \mathcal{A})$ is a PL-manifold of codimension 2 in $\mathbb{C}\Delta$ with boundary $\partial\mathbb{C}Ch(\mathcal{S}, \mathcal{A}) = \partial\mathbb{C}\Delta \cap \mathbb{C}Ch(\mathcal{S}, \mathcal{A})$. If all the numbers in $\mathcal{A}$ are real, then $\mathbb{C}Ch(\mathcal{S}, \mathcal{A})$ is invariant with respect to* Conj, *the set $\mathbb{R}Ch(\mathcal{S}, \mathcal{A})$ is a PL-manifold of codimension 1 in $\mathbb{R}\Delta$ with boundary $\partial\mathbb{R}Ch(\mathcal{S}, \mathcal{A}) = \partial\mathbb{R}\Delta \cap \mathbb{R}Ch(\mathcal{S}, \mathcal{A})$.*

Remark 1.9 *(a) Neither $\mathbb{C}Ch(\mathcal{S}, \mathcal{A})$, nor $\mathbb{R}Ch(\mathcal{S}, \mathcal{A})$ are smooth in general. Indeed, for the polynomials*

$$F_1(x, y) = x(y - x - 1), \quad F_2(x, y) = xy - x + 1$$

with the Newton triangles

$$\Delta_1 = [(1, 0), (1, 1), (2, 0)], \quad \Delta_2 = [(0, 0), (1, 0), (1, 1)]$$

the closures of curves

$$\mu_{\Delta_1}(\{F_1=0\})=\left\{\left(\frac{2+3x}{2+2x}\ ,\ \frac{x+1}{2+2x}\right)\ :\ x>0\right\},$$

$$\mu_{\Delta_2}(\{F_2=0\})=\left\{\left(\frac{2x-1}{2x}\ ,\ \frac{x-1}{2x}\right)\ :\ x>0\right\}$$

have the common limit point $(1,1/2)$ *on the edge* $[(1,0),(1,1)]$, *and different tangent lines*

$$y=\frac{1}{2},\quad y=x-\frac{1}{2}$$

at this point.

(b) Similarly to Proposition 1.4, the statement of Proposition 1.8 holds for Δ *intersecting coordinate hyperplanes, when substituting* $\mathbb{C}\Delta\backslash\mathrm{Sing}(\mathbb{C}\Delta)$ *and* $\mathbb{R}\Delta\backslash\mathrm{Sing}(\mathbb{C}\Delta)$ *for* $\mathbb{C}\Delta$ *and* $\mathbb{R}\Delta$*, respectively.*

Proof of Proposition 1.8. In view of Proposition 1.4, we have to study only the behavior of $\mathbb{C}Ch(\mathcal{S},\mathcal{A})$ of faces $\delta\subset\Delta_k\cap\Delta_l$, $k\ne l$, $\dim\delta=s>0$.

Let $I(\delta)\subset I(\Delta)$, $\delta=\Delta_1\cap...\cap\Delta_k$, $\delta\not\subset\Delta_{k+1}\cup...\cup\Delta_N$. Pick a point $w\in\mathbb{C}I(\delta)\cap\mathbb{C}Ch(\mathcal{S},\mathcal{A})$. Following the proof of Proposition 1.4, we impose assumptions of step (iv) and obtain that a neighborhood U_w of w in $\mathbb{C}\Delta$ is the union of neighborhoods $U_{w,1},...,U_{w,k}$ of w in $\mathbb{C}\Delta_1,...,\mathbb{C}\Delta_k$, respectively, so that $U_{w,m}$ is parameterized by $\mathrm{pr}_s(U'_m)\times U_{0,m}\times U''$, where $\mathrm{pr}_s(U'_1)=...=\mathrm{pr}_s(U'_k)$, U'' is common for all $m=1,...,k$, and $U_{0,m}$ is a neighborhood of the point $\mathrm{pr}_{n-s}(\delta)=(p_{s+1},...,p_n)$ in $\mathrm{pr}_{n-s}(\Delta_m)$, and, moreover, these parameterizations are compatible on the common faces of $\Delta_1,...,\Delta_k$. In turn, $\mathbb{C}Ch(\mathcal{S},\mathcal{A})\cap U_{w,m}$ is parameterized by

$$\prod_{j=2}^{s}\{|x_j-x_j^0|<\varepsilon\}\times U_{0,m}\times\prod_{j=2}^{n}\{|v_j-v_j^0|<\varepsilon\}\ ,$$

and by (4) these parameterizations are compatible along common faces of $\Delta_1,...,\Delta_k$. Hence $\mathbb{C}Ch(\mathcal{S},\mathcal{A})\cap U_w=\bigcup_m\mathbb{C}Ch(\mathcal{S},\mathcal{A})\cap U_{0,m}$ is parameterized by

$$\prod_{j=2}^{s}\{|x_j-x_j^0|<\varepsilon\}\times\bigcup_{m=1}^{k}U_{0,m}\times\prod_{j=2}^{n}\{|v_j-v_j^0|<\varepsilon\}\ ,$$

which is homeomorphic to a $(2n-2)$-ball.

Similarly one treats the case $\delta\subset\partial\Delta$ and the real case. $\Box$

Definition 1.10 *A homeomorphism or an isotopy in* $\mathbb{C}\Delta$ *are called tame if they respect the filtration of* $\mathbb{C}\Delta$ *by* $\mathbb{C}\delta$*, where* δ *runs over all faces of* Δ*, i.e., the restrictions of a tame homeomorphism or a tame isotopy to any* $\mathbb{C}\delta$ *are a homeomorphism, or an isotopy in* $\mathbb{C}\delta$*. In addition, we call such objects equivariant if they commute with* Conj.

Remark 1.11 *For given Δ and $\mathcal{S}$, and different $\mathcal{A}, \mathcal{A}' : \Delta \cap \mathbb{Z}^n \to \mathbb{C}$ the C-hypersurfaces $\mathbb{C}Ch(\mathcal{S}, \mathcal{A})$ and $\mathbb{C}Ch(\mathcal{S}, \mathcal{A}')$ are tame isotopic in $\mathbb{C}\Delta$ (not equivariantly, in general). Indeed, one can connect $\mathcal{A}$ and $\mathcal{A}'$ by a family $\mathcal{A}_t$, $t \in [0,1]$ such that the polynomials $F_{k,t}$ are nondegenerate for all $k = 1, ..., N$, $t \in [0,1]$.*

1.6 Projectivization of charts

Let d and n be positive integers. Denote by T_d^n the simplex in $\mathbb{R}^n$ with vertices $(0,0,...,0)$, $(d,0,...,0)$, ..., $(0,...,0,d)$. Observe that $\mathbb{C}T_d^n$ is homeomorphic to a closed ball in $\mathbb{C}^n$. The group $S^1 = \{|v| = 1\}$ acts freely on $\partial\mathbb{C}T_d^n$ by

$$v \in S^1,\ (z_1, ..., z_n) \in \partial\mathbb{C}T_d^n \ \mapsto\ (z_1 v, ..., z_n v) \in \partial\mathbb{C}T_d^n\ .$$

Proposition 1.12 *The quotient $\mathbb{C}T_d^n / S^1$ is equivariantly homeomorphic to the projective space $\mathbb{C}P^n$ so that the faces of T_d^n naturally correspond to the coordinate planes in $\mathbb{C}P^n$.*

Proof. The momentum map $\mathbb{C}\mu_{T_d^n}$ takes $\mathbb{C}^n$ diffeomorphically onto $\mathrm{Int}(\mathbb{C}T_d^n)$. Its inverse extends up to a surjective map $\nu_d^n : \mathbb{C}T_d^n \to \mathbb{C}P^n$ in the following way. Consider a point $(0, z_1, ..., z_n) \in \mathbb{C}P^n \backslash \mathbb{C}^n$, the line $\gamma(t) = (z_1 t, ..., z_n t) \in \mathbb{C}^n$, $t \in (0, \infty)$, and $\lim_{t\to\infty} \mathbb{C}\mu_{T_d^n}(\gamma(t))$. Then $\lim_{t\to\infty} \mathbb{C}\mu_{T_d^n}(\gamma(t)) = \mathbb{C}\mu_\delta(z_1, ..., z_n)$, where

$$\delta = \left\{ \sum_{z_k \neq 0} i_k = d \right\} \cap \bigcap_{z_k = 0} \{i_k = 0\}$$

is a face of T_d^n. Then we put $\nu_d^n(\mathbb{C}\mu_\delta(z_1, ..., z_n) = (0, z_1, ..., z_n) \in \mathbb{C}P^n$. One can easily verify that ν_d^n is well-defined, surjective, continuous, commutes with Conj, sends orbits of the S^1-action on $\partial\mathbb{C}T_d^n$ into the points of the hyperplane $\{z_0 = 0\}$ in $\mathbb{C}P^n$, and establishes a one-to-one correspondence between the faces of T_d^n and the coordinate planes in $\mathbb{C}P^n$. $\Box$

Remark 1.13 *Further on we always assume $\mathbb{C}T_d^n / S^1$ to be identified with $\mathbb{C}P^n$ via the map ν_d^n.*

Definition 1.14 *In the notation of section 1.5, given a subdivision $T_d^n = \Delta_1 \cup ... \cup \Delta_N$ and a collection of complex numbers $\mathcal{A} = \{A_{\bar{i}} : \bar{i} \in T_d^n \cap \mathbb{Z}^n\}$, define the projective gluing of charts as*

$$P\mathbb{C}Ch(\mathcal{S}, \mathcal{A}) = \nu_d^n(\mathbb{C}Ch(\mathcal{S}, \mathcal{A})) \subset \mathbb{C}P^n\ .$$

This object is called a complex combinatorial hypersurface (C-hypersurface, in short) of degree d in $\mathbb{C}P^n$. If the numbers $\mathcal{A} = \{A_{\bar{i}} : \bar{i} \in T_d^n \cap \mathbb{Z}^n\}$ are real, the C-hypersurface is called real and its real part $\nu_d^n(\mathbb{R}Ch(\mathcal{S}, \mathcal{A}))$ is denoted by $P\mathbb{R}Ch(\mathcal{S}, \mathcal{A})$.

Proposition 1.15 *If F is a nondegenerate polynomial of degree d in n variables, then $P\mathbb{C}Ch(F) \subset \mathbb{C}P^n$ coincides with the projective closure $P\{F=0\}$ of the affine hypersurface $\{F=0\}$.*

Proof. Straightforward from $\nu_d^n(\mathbb{C}Ch(F)) = P\{F=0\}$ and the fact that $\mathbb{C}Ch(F) \cap \partial(\mathbb{C}T_d^n)$ is invariant with respect to the S^1 action. □

Definition 1.16 *A homeomorphism, or an isotopy in $\mathbb{C}P^n$ are called tame if they respect the filtration of $\mathbb{C}P^n$ by the coordinate planes, i.e., their restrictions to any coordinate plane are a homeomorphism, or an isotopy in that plane. In addition, we call such objects equivariant if they commute with* Conj.

Proposition 1.17 *In the previous notation, $P\mathbb{C}Ch(\mathcal{S},\mathcal{A})$ is a PL-submanifold in $\mathbb{C}P^n$, of codimension 2. It is invariant with respect to* Conj *if all the numbers in $\mathcal{A}$ are real.*

Proof. First, we show that the C-hypersurface $\mathbb{C}Ch(\mathcal{S},\mathcal{A})$ is a PL-manifold in $\mathbb{C}T_d^n$ of codimension 2 with boundary

$$\partial\mathbb{C}Ch(\mathcal{S},\mathcal{A}) = \mathbb{C}Ch(\mathcal{S},\mathcal{A}) \cap \partial\mathbb{C}T_d^n \ ,$$

and if all the numbers in $\mathcal{A}$ are real $\mathbb{C}Ch(\mathcal{S},\mathcal{A}$ is invariant with respect to Conj, and $\text{Fix(Conj)} = \mathbb{R}Ch(\mathcal{S},\mathcal{A})$ is a PL-manifold in $\mathbb{R}T_d^n$ of codimension 1 with boundary

$$\partial\mathbb{R}Ch(\mathcal{S},\mathcal{A}) = \mathbb{R}Ch(\mathcal{S},\mathcal{A}) \cap \partial\mathbb{R}T_d^n \ .$$

By Proposition 1.8 it is enough to consider only the gluing of charts on the coordinate hyperplanes. For any polyhedron $\Delta \subset \mathbb{R}^n$, put $\Delta_t = (t,...,t)+\Delta$, $t \ge 0$.

The family $\mathbb{C}\mu_{\Delta_{k,t}}$, $0 \le t \le 1$, connects $\mathbb{C}\mu_{\Delta_k}$ with $\mathbb{C}\mu_{\Delta_{k,1}}$, $k=1,...,N$, and defines an isotopy of $\text{Cl}(\mathbb{C}\mu_{\Delta_{k,t}}(F_k=0))$, the closure of $\mathbb{C}\mu_{\Delta_{k,t}}(F_k=0)$, $0<t\le 1$, which degenerates into $\mathbb{C}Ch(F_k)$, $k=1,...,N$. The same holds for any proper face δ of $\Delta_1,...,\Delta_N$ and the corresponding truncation of $F_1,...,F_N$.

Since $\text{Sing}(\mathbb{C}\Delta_{k,1}) = \emptyset$, $k=1,...,N$, the set $\mathcal{K}_1 = \bigcup_{k=1}^N \mathbb{C}Ch(z_1...z_nF_k)$ is a PL-manifold with boundary on $\partial\mathbb{C}\widetilde{T}_d^n$, according to Proposition 1.4 (where $\widetilde{T}_d^n = (1,...,1)+T_d^n$). Denote some faces of $\widetilde{T}_d^n$ as follows:

$$\widetilde{T}_d^n(k) = \widetilde{T}_d^n \cap \{i_k = 1\}, \quad \widetilde{T}_d^n(k_1,...,k_s) = \bigcap_{j=1}^s \widetilde{T}_d^n(k_j) \ .$$

Put $\delta = \widetilde{T}_d^n(k)$. By (3)

$$\mathbb{C}\delta \cong \mathbb{C}T_d^{n-1} \times S^1. \tag{9}$$

Similarly, by (3) $\mathcal{K}_1 \cap \mathbb{C}\delta$ is a PL-manifold of dimension $2n-3$ with boundary on $\partial\mathbb{C}\delta$, satisfying

$$\mathcal{K}_1 \cap \mathbb{C}\delta \cong \left(\bigcup_{l=1}^N \mathbb{C}Ch(z_1...z_nF_l^\delta)\right) \times S^1, \tag{10}$$

where the product structure is compatible with that in (9). Using the product representations (9) and (10), we define

$$D^n(k) = \{(w_1, ..., w_n) \ : \ (w_1, ..., w_{k-1}, w_{k+1}, ..., w_n) \in \mathbb{C}T_d^{n-1}, \ |w_k| \le 1\}$$
$$\cong \mathbb{C}T_d^{n-1} \times D^2 \ ,$$

where D^2 is the closed 2-dimensional unit ball, and inside $D^n(k)$ the $(2n-2)$-dimensional manifold $M(k)$:

$$\{(w_1, ..., w_n) \ : \ (w_1, ..., w_{k-1}, 1, w_{k+1}, ..., w_n) \in \bigcup_{l=1}^{N} \mathbb{C}Ch(z_1 ... z_n F_l^\delta), \ |w_k| \le 1\}$$
$$\cong \left(\bigcup_{l=1}^{N} \mathbb{C}Ch(\widetilde{F}_l^\delta) \right) \times D^2$$

with boundary on $\partial D^n(k)$ and such that $M(k) \cap \mathbb{C}\delta = \bigcup_{l=1}^{N} \mathbb{C}Ch(z_1 ... z_n F_l^\delta)$.

Similarly, given $\delta = \widetilde{T}_d^n(k_1, ..., k_s)$, we have by (3),

$$\mathbb{C}\delta \cong \mathbb{C}T_d^{n-s} \times (S^1)^s, \quad \mathcal{K}_1 \cap \mathbb{C}\delta \cong \left(\bigcup_{l=1}^{N} \mathbb{C}Ch(z_1 ... z_n F_l^\delta) \right) \times (S^1)^s,$$

and one defines

$$D^n(k_1, ..., k_s) = \{(w_1, ..., w_n) \ : \ (w_j)_{j \ne k_1, ..., k_s} \in \mathbb{C}T_d^{n-s}, \ |w_{k_1}|, ..., |w_{k_s}| \le 1\}$$
$$\cong \mathbb{C}T_d^{n-s} \times (D^2)^s \ ,$$

and the $(2n-2)$-dimensional manifold $M(k_1, ..., k_s)$:

$$\{(w_1, ..., w_n) \ : \ (w_j)_{j \ne k_1, ..., k_s} \in \bigcup_{l=1}^{N} \mathbb{C}Ch(z_1 ... z_n F_l^\delta), \ |w_{k_1}|, ..., |w_{k_s}| \le 1\}$$
$$\cong \left(\bigcup_{l=1}^{N} \mathbb{C}Ch(z_1 ... z_n F_l^\delta) \right) \times (D^2)^s$$

with boundary on $\partial D^n(k_1, ..., k_s)$ and such that

$$M(k_1, ..., k_s) \cap \mathbb{C}\delta = \bigcup_{l=1}^{N} \mathbb{C}Ch(z_1 ... z_n F_l^\delta) \ .$$

It can easily be shown that $\mathcal{T}_1 = \mathbb{C}\widetilde{T}_d^n \cup \bigcup_{s=1}^{n} \bigcup_{1 \le k_1 < ... < k_s} D^n(k_1, ..., k_s)$ is the convex hull of $\mathbb{C}\widetilde{T}_d^n$ in $\mathbb{C}^n$, and $\mathcal{K}_1 \cup \bigcup_{s=1}^{n} \bigcup_{1 \le k_1 < ... < k_s} M(k_1, ..., k_s)$ is a PL-manifold in $\mathcal{T}_1$ of codimension 2 with boundary on $\partial \mathcal{T}_1$.

Now, for any $t \in (0, 1)$, in the same way one constructs similar objects:

- complexifications of polyhedra

$$\mathbb{C}\Delta_{k,t} = \mathrm{Cl}(\mathbb{C}\mu_{\Delta_k,t}((\mathbb{C}^*)^n)), \quad \mathbb{C}T^n_d(t) = \bigcup_{k=1}^{N} \mathbb{C}\Delta_{k,t},$$

- gluing of charts $\mathcal{K}_t = \bigcup_{k=1}^{N} \mathrm{Cl}(\mathbb{C}\mu_{\Delta_k,t}(\{F_k = 0\}))$, which is a PL-manifold of codimension 2 in $\mathbb{C}T^n_d(t)$ with boundary on $\partial\mathbb{C}T^n_d(t)$,
- completion of $\mathbb{C}T^n_d(t)$ up to its convex hull

$$\mathcal{T}_t = \mathbb{C}T^n_d(t) \cup \bigcup_{s=1}^{n} \bigcup_{1\le k_1<\ldots<k_s} D^n_t(k_1, ..., k_s),$$

  where

$$\begin{aligned} D^n_t(k_1, ..., k_s) &= \{(w_1, ..., w_n) \ : \ (w_j)_{j\neq k_1,...,k_s} \in \mathbb{C}T^{n-s}_d, \ |w_{k_1}|, ..., |w_{k_s}| \le t\} \\ &\cong \mathbb{C}T^{n-s}_d \times (D^2(t))^s \ , \end{aligned}$$

  $D^2(t)$ is a disc of radius t,
- a $(2n-2)$-dimensional manifold

$$\begin{aligned} M_t(k_1, ..., k_s) = \Big\{ &(w_1, ..., w_n) \ : \\ &(w_j)_{j\neq k_1,...,k_s} \in \bigcup_{l=1}^{N} \mathrm{Cl}(\mathbb{C}\mu_{\delta_t}(\{F^\delta_l = 0\})), \ |w_{k_1}|, ..., |w_{k_s}| \le t \Big\} \\ &\cong \left(\bigcup_{l=1}^{N} \mathrm{Cl}(\mathbb{C}\mu_{\delta_t}(\{F^\delta_l = 0\})) \right) \times (D^2(t))^s \end{aligned}$$

- a PL-manifold

$$\mathcal{K}_t \cup \bigcup_{s=1}^{n} \bigcup_{1\le k_1<\ldots<k_s} M_t(k_1, ..., k_s) \tag{11}$$

  of codimension 2 in $\mathcal{T}_t$ with boundary on $\partial\mathcal{T}_t$.

If t varies from 1 to 0, $\mathcal{T}_t$ contracts from $\mathcal{T}_1$ to $\mathbb{C}T^n_d$ and the manifold (11) naturally contracts into the C-hypersurface $\mathbb{C}Ch(\mathcal{S}, \mathcal{A})$ with boundary on $\partial\mathbb{C}T^n$.

To complete the proof of Proposition 1.17 we have to verify only that $\partial\mathbb{C}Ch(\mathcal{S}, \mathcal{A})$ is invariant with respect to the S^1 action on $\partial\mathbb{C}T^n_d$, and the quotient $\mathbb{C}Ch(\mathcal{S}, \mathcal{A})/S^1$ is a closed manifold.

First, note that $\partial\mathbb{C}Ch(\mathcal{S}, \mathcal{A})$ is invariant with respect to the S^1 action. Indeed, $\partial\mathbb{C}Ch(\mathcal{S}, \mathcal{A})$ consists of charts of homogeneous polynomials which are invariant with respect to the S^1 action on $\partial\mathbb{C}T^n_d$. Second, for any edge of T^n_d there exists a combination of an automorphism of $\mathbb{Z}^n$ and shifts which puts this edge on a coordinate axis and the adjacent faces of T_d^n on the corresponding coordinate planes. Since such a transformation is compatible with ν_d^n defined via the momentum map, the question on the behavior of $\mathbb{C}Ch(\mathcal{S},\mathcal{A})$ on $\partial\mathbb{C}T_d^n$ is reduced to that on coordinate planes, which has been treated above. $\square$

Remark 1.18 *Similarly to Remark 1.11, two C-hypersurfaces in $\mathbb{C}P^n$ of the same degree and with the same subdivision $\mathcal{S}$ of T_d^n are tame isotopic in $\mathbb{C}P^n$ (not equivariantly, in general).*

2 Topology of complex C-hypersurfaces

2.1 Complexification of the Viro theorem

Theorem 2.1 (Affine complex Viro theorem). *In the notation of section 1.5, if $\Delta \subset (\mathbb{R}_+^*)^n$ and a subdivision $\mathcal{S} \ : \ \Delta = \Delta_1 \cup ... \cup \Delta_N$ is defined by a convex piecewise-linear function $\nu : T_d^n \to \mathbb{R}$, then $\mathbb{C}Ch(\mathcal{S},\mathcal{A})$ is tame isotopic in $\mathbb{C}\Delta$ (equivariantly, if all the numbers in $\mathcal{A}$ are real) to $\mathbb{C}Ch(F)$, where*

$$F(z_1,...,z_n,t) = \sum_{(i_1,...,i_n)\in\Delta} A_{i_1...i_n} t^{\nu(i_1,...,i_n)} z_1^{i_1}\cdot ... \cdot z_n^{i_n}$$

and $t = \text{const} > 0$ *is sufficiently small.*

Proof. Our argument is similar to that from the proof of the Viro gluing theorem [32] (see also [13], Chapter 11, and [23]).

Without loss of generality one can suppose that $\dim\Delta = n$, and ν is positive integral-valued at integral points. Consider the polynomial $\hat{F}(z_1,...,z_n,t) = (1+t)F(z_1,...,z_n,t)$ as a polynomial in $t, x_1,...,x_n$, and denote by $\hat{\Delta}$ its Newton polyhedron in $\mathbb{R}^{n+1}$. The graph of ν is the "lower" part of $\partial\hat{\Delta}$, naturally projected onto $\Delta \subset \mathbb{R}^n = \{i_{n+1} = 0\} \subset \mathbb{R}^{n+1}$. Denote by $\hat{\Delta}_k$ the part of the graph of ν projected onto Δ_k, $k = 1,...,N$. Denote by B the part of $\hat{\Delta}$ projected onto $\partial\Delta$. Clearly, B can be viewed as $\partial\Delta\times[0,1]$.

Step 1. Introduce the set

$$H_0 = \{(w_1,...,w_n,\nu(|w_1|,...,|w_n|)) \ : \ (|w_1|,...,|w_n|)\in\Delta\} = \bigcup_{k=1}^N \hat{\Delta}_k\times(S^1)^n \subset \partial\mathbb{C}\hat{\Delta} \ ,$$

and the family of hypersurfaces $H_a = \text{Cl}(\mathbb{C}\mu_{\hat{\Delta}}(\{t=a\}) \subset \mathbb{C}\hat{\Delta}$, $a\in(0,\varepsilon)$. We shall show that the family H_a, $a\in[0,\varepsilon)$, is an isotopy in $\mathbb{C}\hat{\Delta}$.

The map $\mathbb{C}\mu_{\hat{\Delta}}$ takes $\{t=a\}\subset(\mathbb{C}^*)^{n+1}$ diffeomorphically into $\mathbb{C}I(\hat{\Delta})$. Considering $\lim_{\tau\to 0}\mu_{\hat{\Delta}}(\gamma(\tau))$ for all curves

$$\gamma(\tau) = \left(\lambda_1 x_1^{k_1},...,\lambda_n x_n^{k_n}, a\right) \subset (\mathbb{R}_+^*)^{n+1}, \quad \tau > 0,$$

with $\lambda_1,...,\lambda_n>0$, $(k_1,...,k_n)\ne 0$, one can show (as in the proof of Proposition 1.4) that H_a being the closure of $\mathbb{C}\mu_{\hat\Delta}(\{t=a\})$ is a PL-manifold with boundary $\partial H_a=H_a\cap B$, which projects onto $\partial\Delta$. Moreover, $H_a\cap H_b=\emptyset$, $a\ne b$. The projection on the first n coordinates provides a homeomorphism of H_a, $a>0$, and $\mathbb{C}\Delta$. Indeed, in $\mathbb{C}I(\hat\Delta)$ this fact is reduced to the claim that

$$(x_1,...,x_n)\in(\mathbb{R}^*_+)^n\mapsto\frac{\sum_{(i_1,...,i_n)\in\Delta}x_1^{i_1}...x_n^{i_n}a^{\nu(i_1,...,i_n)}\cdot(i_1,...,i_n)}{\sum_{(i_1,...,i_n)\in\Delta}x_1^{i_1}...x_n^{i_n}a^{\nu(i_1,...,i_n)}}\in I(\Delta)$$

is a homeomorphism, which holds since the above map is a kind of the momentum map [1, 2]. Similarly one shows that the projection is one-to-one on $\partial H_a\subset B$, taking into account that the limit of $\mu_{\hat\Delta}$ on a face of B is the momentum map of this face. To finish the proof one should note that $\bigcup_{a>0}H_a$ fill the intersection of $\mathbb{C}I(\hat\Delta)$ with the space $\mathrm{Im}(w_{n+1})=0$, where $w_1,...,w_{n+1}$ are coordinates in $(\mathbb{C}^*)^{n+1}\supset\mathbb{C}\hat\Delta$.

Step 2. Now we show that $\mathbb{C}Ch(\hat F)\cap H_a$, $a\in[0,\varepsilon)$, is an isotopy in $\mathbb{C}\hat\Delta$. For $\varepsilon>0$ small enough, $\{\hat F=0\}\cap\{t=a\}=\{F=0\}\cap\{t=a\}$ and the intersection is transverse in $(\mathbb{C}^*)^{n+1}$. This implies that the family $\mathbb{C}Ch(\hat F)\cap H_a$, $a\in(0,\varepsilon)$, is an isotopy in $\mathbb{C}\hat\Delta$.

Now we pick a point $\overline{w}=(y_1v_1^0,...,y_nv_n^0,y_{n+1})\in H_0\cap\mathbb{C}Ch(\hat F)$, where $(y_1,...,y_n)\in I(\delta)$ for some s-dimensional face of $\Delta_1,...,\Delta_N$, $y_{n+1}=\nu(y_1,...,y_n)$, and $(v_1^0,...,v_n^0)\in(S^1)^n$. Following the proof of Proposition 1.4 and using transformations

$$(x_1,...,x_n,t)\mapsto\left(t^{\beta_1}\prod_{k=1}^n x_k^{\alpha_{k1}},...,t^{\beta_n}\prod_{k=1}^n x_k^{\alpha_{kn}},t\right)$$

with $A=(\alpha_{kj})\in SL(\mathbb{Z},n)$, we can get δ to be parallel to the coordinate s-plane $\{i_1=...=0\}$, $\nu\big|_\delta=\nu_0$, and $\nu\big|_{\Delta\backslash\delta}>\nu_0$. Then we introduce a parameterization of a neighborhood $U_{\overline{w}}$ of the point $\overline{w}$ in $\mathbb{C}\hat\Delta\cap\{\mathrm{Im}(w_{n+1})=0\}$ by $\mathrm{pr}_s(U')\times U_0\times U''$ via an extension of the map $\mathbb{C}\mu_{\hat\Delta}$ such that, in the notation of the proof of Proposition 1.4, $\mathrm{pr}_s(U')$ is given by the first s inequalities in (7), U'' is given by (6), and U_0 is a neighborhood of the point $\mathrm{pr}_{n-s+1}(\delta)$ in $\mathrm{pr}_{n-s+1}(\hat\Delta)$. Here $\mathrm{Int}(U_0)$ is identified with the domain $\mathrm{pr}_{n-s+1}(U')\subset(\mathbb{R}^*_+)^{n-s+1}$, given in the coordinates $x_{s+1},...,x_n,t$ by the relations

$$M_{i_{s+1},...,i_n,i_{n+1}}=x_{s+1}^{i_{s+1}}...x_n^{i_n}t^{i_{n+1}}<\varepsilon,\quad(i_{s+1},...,i_{n+1})\in\mathrm{pr}_{n-s+1}(\hat\Delta\backslash\Delta)\cap\mathbb{Z}^{n-s+1}\ ,$$

and $\mathbb{C}Ch(\hat F)\cap\mathrm{Int}(U_{\overline{w}})$ is parameterized in $\mathrm{pr}_s(U')\times\mathrm{Int}(U_0)\times U''$ by equations (8), where G and H are vanishing at zero smooth functions of $x_2,...,x_s$, $v_1,...,v_n$ and $M_{i_{s+1},...,i_n,i_{n+1}}$, $(i_{s+1},...,i_{n+1})\in\mathrm{pr}_{n-s+1}(\hat\Delta\backslash\Delta)\cap\mathbb{Z}^{n-s+1}$.

Denote $\widetilde\Delta=\delta\times\mathrm{pr}_{n-s+1}(\hat\Delta)$. Clearly, $\hat\Delta$ and $\widetilde\Delta$ coincide in a neighborhood of $\overline{w}$. Moreover, The maps $\mathbb{C}\mu_{\hat\Delta}$ and $\mathbb{C}\mu_{\widetilde\Delta}$ are connected by an isotopy on the domain $\mathrm{pr}_s(U')\times\mathrm{Int}(U_0)\times U''$: such an isotopy can be written explicitly by supplying the non-common summands in the formulae for $\mathbb{C}\mu_{\hat\Delta}$ and $\mathbb{C}\mu_{\widetilde\Delta}$ by a parameter running over $[0,1]$. This isotopy extends up to equivariant tame isotopy on $\mathrm{pr}_s(U')\times U_0\times U''$, so that replacing $\mathbb{C}\mu_{\hat\Delta}$ by $\mathbb{C}\mu_{\widetilde\Delta}$, one replace $U_{\overline{w}}$ by another neighborhood $\widetilde U_{\overline{w}}$ of $\overline{w}$

in $\mathbb{C}\widetilde{\Delta} \cap \{\mathrm{Im}(w_{n+1})\}$. Note that the map $\mathbb{C}\mu_{\widetilde{\Delta}}$ splits on $\mathrm{pr}_s(U') \times U_0 \times U''$ into the product of

$$\mu_\delta : \mathrm{pr}_s(U') \to \delta, \quad \mu_{\mathrm{pr}_{n-s+1}(\hat{\Delta})} \to \mathrm{pr}_{n-s+1}(\hat{\Delta}), \quad \mathrm{Id} : U'' \to U''.$$

Together with the result of Step 1 this allows us to introduce in U_0 coordinates $t, \theta_1, ..., \theta_{n-s}$ so that

$$U_0 = \{0 \le t < \varepsilon, \ -\varepsilon < \theta_j < \varepsilon\}, \quad U_0 \cap \partial(\mathrm{pr}_{n-s+1}(\hat{\Delta})) = \{t = 0\} \ ,$$

and $\mathbb{C}\mu_{\widetilde{\Delta}}(H_a) = \mathrm{pr}_s(U') \times \{t = a\} \times U''$. In addition, $\mu_{\mathrm{pr}_{n-s+1}(\hat{\Delta})}$ expresses $M_{i_{s+1},...,i_n,i_{n+1}}$, $(i_{s+1}, ..., i_{n+1}) \in \mathrm{pr}_{n-s+1}(\hat{\Delta}\backslash\Delta) \cap \mathbb{Z}^{n-s+1}$, as continuous functions of $t, \theta_1, ..., \theta_{n-s}$, vanishing at zero. Hence the closure of $\mathbb{C}\mu_{\widetilde{\Delta}}(\{F = 0\})$ in $\widetilde{U_{\overline{w}}}$ is given by equations

$$x_1 = x_1^0 + G', \quad v_1 = v_1^0 + H', \tag{12}$$

where G' and H' are vanishing at zero smooth functions of $x_2, ..., x_s$, $v_2, ..., v_n$, $t, \theta_1, ..., \theta_{n-s}$. The variety (12) intersects any hypersurface $\mathrm{pr}_s(U') \times \{t = a\} \times U''$, $a \in [0, \varepsilon)$, transversally in $\mathrm{pr}_s(U') \times U_0 \times U'' \simeq \widetilde{U_{\overline{w}}} \simeq U_{\overline{w}}$, thereby proving that $\mathbb{C}Ch(\hat{F}) \cap H_a$, $a \in [0, \varepsilon)$, is an isotopy in $\mathbb{C}\hat{\Delta}$.

Step 3. For $a \ne 0$, the projection of $\mathbb{C}Ch(\hat{F}) \cap H_a$ into $\mathbb{C}\Delta$ is the closure of the image of $\{F = 0\} \cap \{t = a\} \subset (\mathbb{C}^*)^{n+1}$ by the map

$$\mathbb{C}\mu_{\Delta,a}(\overline{x}, \overline{v}) = \frac{\sum_{(i_1,...,i_n)\in\Delta}(i_1v_1, ..., i_nv_n) \cdot \left(x_1^{i_1}...x_n^{i_n} \sum_{(i_1,...,i_n,k)\in\hat{\Delta}} a^k\right)}{\sum_{(i_1,...,i_n)\in\Delta}\left(x_1^{i_1}...x_n^{i_n} \sum_{(i_1,...,i_n,k)\in\hat{\Delta}} a^k\right)} \ . \tag{13}$$

The map $\mathbb{C}\mu_{\Delta,a}$ is connected with $\mathbb{C}\mu_\Delta$ on $\{t = a\} \simeq (\mathbb{C}^*)^n$ by the isotopy

$$\frac{\sum_{(i_1,...,i_n)\in\Delta}(i_1v_1, ..., i_nv_n) \cdot \left(x_1^{i_1}...x_n^{i_n}\left(\sum_{(i_1,...,i_n,k)\in\hat{\Delta}} a^k(1-\tau) + \tau\right)\right)}{\sum_{(i_1,...,i_n)\in\Delta}\left(x_1^{i_1}...x_n^{i_n}\left(\sum_{(i_1,...,i_n,k)\in\hat{\Delta}} a^k(1-\tau) + \tau\right)\right)} \ , \tag{14}$$

providing a tame isotopy of the projection of $\mathbb{C}Ch(\hat{F}) \cap H_a$ with $\mathbb{C}Ch(F\big|_{t=a}) \subset \mathbb{C}\Delta$. Similarly, the projection of $\mathbb{C}Ch(\hat{F}) \cap \mathbb{C}\hat{\Delta}_k \cap \{\mathrm{Im}(w_{n+1}) = 0\}$ into $\mathbb{C}\Delta_k$, $1 \le k \le N$, is the closure of the image of $\{F^{\hat{\Delta}_k} = 0\} \cap \{t = 1\}$ by the map $\mathbb{C}\mu_{\hat{\Delta}_k}$ which coincides with the closure of the image of $\{F_k = 0\}$ by the map $\mathbb{C}\mu_{\Delta_k}$, *i.e.*, $\mathbb{C}Ch(F_k)$.

So, Theorem is proven in the complex case. If all the numbers in $\mathcal{A}$ are real, then the isotopy constructed is equivariant. □

Theorem 2.2 (Projective complex Viro theorem). *Let, in the notation of Theorem 2.1, $\Delta = T_d^n$ and the subdivision $\mathcal{S}$ be defined by a convex piecewise-linear function $\nu : T_d^n \to \mathbb{R}$. Then $P\mathbb{C}Ch(\mathcal{S}, \mathcal{A})$ is tame isotopic in $\mathbb{C}P^n$ (equivariantly, if all the numbers in $\mathcal{A}$ are real) to $P\mathbb{C}Ch(F)$, where*

$$F(z_1, ..., z_n, t) = \sum_{i_1+...+i_n \le d} A_{i_1...i_n} t^{\nu(i_1,...,i_n)} z_1^{i_1} \cdot ... \cdot z_n^{i_n} = 0,$$

and $t = \mathrm{const} > 0$ is sufficiently small.

Proof. We multiply all the polynomials by $z_1...z_n$ and apply Theorem 2.1 to the simplex $\widetilde{T}^n_d = T^n_d+(1,...,1)$ and correspondingly shifted $\mathcal{S},\mathcal{A}$. Then we note that the isotopy of $\mathbb{C}Ch(z_1...z_nF)$ and $\bigcup_{k=1}^N \mathbb{C}Ch(z_1...z_nF_k)$ in $\mathbb{C}\widetilde{T}^n_d$ is tame and compatible with the action of $(S^1)^n$ on $\partial\mathbb{C}\widetilde{T}^n_d$. This allows us to take quotient by this action as was done in the proof of Proposition 1.17 and obtain the required isotopy. $\square$

Proposition 2.3 *Let two subdivisions $\mathcal{S}=\{\Delta_k,\ k=1,...,N\}$ and $\mathcal{S}'=\{\Delta_{kl},\ l=1,...,r_k,\ k=1,...,N\}$ of a polyhedron $\Delta\subset(\mathbb{R}^*_+)^n$ satisfy*

$$\Delta_k=\bigcup_{l=1}^{r_k}\Delta_{kl},\quad k=1,...,N,$$

so that the subdivision $\mathcal{S}'$ is given by piecewise-linear function $\nu:\Delta\to\mathbb{R}$, whose restrictions $\nu_k=\nu\big|_{\Delta_k}$, $k=1,...,N$, are convex. Then the varieties $\mathbb{C}Ch(\mathcal{S},\mathcal{A})$ and $\mathbb{C}Ch(\mathcal{S}',\mathcal{A}')$ are tame isotopic, provided $\mathcal{A},\mathcal{A}':\Delta\to\mathbb{C}$ define nondegenerate polynomials F_k, F_{kl}, $l=1,...,r_k$, $k=1,...,N$. Similarly, given two subdivisions $\mathcal{S}$, $\mathcal{S}'$ of T^n_d satisfying the previous assumptions, the C-hypersurfaces of degree d constructed out of $\mathcal{S},\mathcal{A}$ and $\mathcal{S}',\mathcal{A}'$ are tame isotopic in $\mathbb{C}P^n$.

Proof. Consider the case $\Delta\subset(\mathbb{R}^*_+)^n$. Fix $a>0$. Let the maps $\mathbb{C}\mu_{\Delta_k,a}:(\mathbb{C}^*)^n\to\mathbb{C}\Delta_k$, $k=1,...,N$, be defined by (13), where summations run over Δ_k and $\hat{\Delta}_k$, constructed as in the proof of Theorem 2.1 by means of ν_k. Put

$$F_{k,a}=\sum_{(i_1,...,i_n)\in\Delta_k}A'_{i_1...i_n}z_1^{i_1}...z_n^{i_n}a^{\nu(i_1,...,i_n)},\quad k=1,...,N.$$

As shown in the proof of Theorem 2.1, the closure $\mathbb{C}Ch_a(F_{k,a})$ of $\mathbb{C}\mu_{\Delta_k,a}(\{F_{k,a}=0\})$ in $\mathbb{C}\Delta_k$ is tame isotopic to $\cup_l\mathbb{C}Ch(F_{kl})$. Moreover, for any face $\delta=\Delta_k\cap\Delta_j$, the isotopies in $\mathbb{C}\Delta_k$ and $\mathbb{C}\Delta_l$ coincide on $\mathbb{C}\delta$. Then, for each $k=1,...,N$, we connect $\mathbb{C}\mu_{\Delta_k,a}$ with $\mathbb{C}\mu_{\Delta_k}$ by an isotopy (14), thereby obtaining the required isotopy of $\mathbb{C}Ch(\mathcal{S},\mathcal{A})$ and $\mathbb{C}Ch(\mathcal{S}',\mathcal{A}')$.

The same argument proves the statement for $\Delta=T^n_d$. $\square$

A subdivision $\Delta=\Delta_1\cup...\cup\Delta_N$ is called *maximal* if it cannot be refined. In this case all the integral points in Δ are vertices of $\Delta_1,...,\Delta_N$, and these polyhedra are simplices.

Corollary 2.4 *Given polyhedron $\Delta\subset(\mathbb{R}^*_+)^n$, any C-hypersurface in $\mathbb{C}\Delta$ is tame isotopic to a C-hypersurface constructed out of a maximal subdivision of Δ. The same is true for C-hypersurfaces in $\mathbb{C}P^n$.*

Proof. Let $\Delta=\Delta_1\cup...\cup\Delta_N$, and $f:\Delta\to\mathbb{R}$ be a smooth convex function. Define a piecewise-linear function $\nu:\Delta\to\mathbb{R}$ as follows: put $\nu(\bar{i})=f(\bar{i})$, $\bar{i}\in\Delta\cap\mathbb{Z}^n$, then define the graph of $\nu\big|_{\Delta_k}$ to be the "lower" part of the convex hull of $\{(\bar{i},\nu(\bar{i})\ :\ \bar{i}\in\Delta_k\}$. The function ν defines a maximal subdivision of Δ inscribed into the initial subdivision and satisfying the conditions of Proposition 2.3, which completes the proof. $\square$

Corollary 2.5 *Any C-curve in $\mathbb{C}P^2$ is tame isotopic (not equivariantly, in general) to an algebraic curve of the same degree.*

Proof. By Corollary 2.4 we can assume that a C-curve in $\mathbb{C}P^2$ is constructed out of a maximal triangulation of T_d^2. We shall transform any given triangulation into a convex triangulation, so that in each transformation step the conditions of Proposition 2.3 hold true.

Let $\mathcal{S}$ be a triangulation of T_d^2, $O(\mathcal{S})$ denote the star of the origin O with respect to this triangulation, and outside $O(\mathcal{S})$ the triangulation $\mathcal{S}$ is maximal. We construct a triangulation $\mathcal{S}'$ of T_d^2 such that $O(\mathcal{S}') \supset O(\mathcal{S})$ and $O(\mathcal{S}') \neq O(\mathcal{S})$. Then, in finitely many steps we come to $O(\mathcal{S}) = T_d^2$, which corresponds to a convex triangulation, and the required statement will follow from Theorem 2.2 and Proposition 2.3.

Let $O, P_1, ..., P_r$ be all the vertices of $O(\mathcal{S})$ numbered successively clockwise along $\partial O(\mathcal{S})$. Any segment $[P_i, P_{i+1}]$ either lies on ∂T_d^2, or is an edge of a unique triangle $T_i \in \mathcal{S}$, $T_i \not\subset O(\mathcal{S})$.

(i) Assume that, for some $i = 1, ..., r$, the vertex $Q \neq P_i, P_{i+1}$ of T_i lies between the straight lines(OP_i) and (OP_{i+1}), and $Q \notin (OP_i) \cup (OP_{i+1})$ (see Figure 2a). Then we change the subdivision of T_d^2 as shown in Figure 2b,c. These changes satisfy the conditions of Proposition 2.3 and lead to a triangulation with a strongly greater star of O.

(ii) Assume that, for some $i = 1, ..., r-1$, $T_i = T_{i+1}$, *i.e.*, the vertices of the latest triangle are P_i, P_{i+1}, P_{i+2} (see Figure 2d). Then we perform the transformation shown in Figure 2e, once again increasing the star of O.

(iii) Assume that there are no triangles T_i as in (i), (ii). Then any triangle T_i is either "left", *i.e.*, the vertex Q lies on (OP_i) or above (OP_i), or "right", *i.e.*, Q lies on (OP_{i+1}) or below (OP_{i+1}). If there exist "left" triangles, consider the "left" triangle T_i with the minimal i. If $i = 1$, we have the situation shown in Figure 2f. Then we change triangulation as shown in Figure 2g, increasing the star of O. If $i > 1$, then the triangle T_{i-1} must be "right", which means that we have a situation shown in Figure 2h. Then we change triangulation as shown in Figure 2i, increasing the star of O. $\square$

2.2 First properties of C-hypersurfaces

The real part of a real C-hypersurface M in $\mathbb{C}P^n$ is denoted by $\mathbb{R}M$.

Proposition 2.6 *A complex C-hypersurface M of degree d in $\mathbb{C}P^n$ is an orientable manifold, homologous to an algebraic hypersurface of degree d in $\mathbb{C}P^n$. If M is real its real part $\mathbb{R}M$ is a closed manifold, mod 2 homologous to the real point set of a real algebraic hypersurface of degree d in $\mathbb{R}P^n$.*

Proof. The Jacobian of the real momentum map $\mu_\Delta : (\mathbb{R}_+^*)^n \to \mathrm{Int}(\Delta)$, $\mu_\Delta(\overline{x}) = (\mu_\Delta^{(1)}(\overline{x}), ..., \mu_\Delta^{(n)}(\overline{x}))$, is positive. Hence the Jacobian of the complex momentum map



Figure 2: Transformation of a triangulation

$\mathbb{C}\mu_\Delta : (\mathbb{C}^*)^n \to \mathbb{C}I(\Delta)$ is positive in the coordinates

$$x_1 = |z_1|,\ v_1 = \frac{z_1}{|z_1|},\ \dots\ ,\ x_n = |z_n|,\ v_n \frac{z_n}{|z_n|}\ ,$$

because this is a diffeomorphism, and at a point with $v_1 = \dots = v_n = 1$ one can easily compute

$$\det\left(\frac{D(\mathbb{C}\mu_\Delta)}{D(x_1, v_1, \dots, x_n, v_n)}\right) = \det\left(\frac{D(\mu_\Delta)}{D(x_1, \dots, x_n)}\right) \cdot \prod_{j=1}^n \mu_\Delta^{(j)}(x_1, \dots, x_n) > 0\ .$$

This means, in particular, that $\mathbb{C}\mu_\Delta$ canonically defines an orientation of images of complex subvarieties of $(\mathbb{C}^*)^n$. Therefore, the open subset

$$\bigcup_{k=1}^N (\mathbb{C}Ch(F_k) \cap \mathbb{C}I(\Delta_k))$$

of $\mathbb{C}Ch(\mathcal{S}, \mathcal{A})$ is canonically orientable. So, to complete the proof of orientability, one should verify that these orientations are compatible when gluing the charts $\mathbb{C}Ch(F_k)$, $\mathbb{C}Ch(F_j)$ with $\Delta_k \cap \Delta_j = \delta$ being a common facet (the gluing along faces of lower dimensions does not affect the orientation). In other words, the orientations of $\mathbb{C}Ch(F_k^\delta) = \mathbb{C}Ch(F_j^\delta)$ induced by $\mathbb{C}Ch(F_k)$ and $\mathbb{C}Ch(F_j)$ are opposite.

Without loss of generality suppose that δ is contained in a hyperplane $i_1 = a > 0$, $\Delta_k \subset \{i_1 \le a\}$, and $\Delta_j \subset \{i_1 \ge a\}$. Then by construction, $\mathbb{C}\Delta_k$ induces on $\mathbb{C}\delta$ an orientation defined by the form

$$-dv_1 \wedge dx_2 \wedge dv_2 \wedge \dots \wedge dx_n \wedge dv_n,$$

and $\mathbb{C}\Delta_j$ induces on $\mathbb{C}\delta$ the opposite orientation defined by

$$dv_1 \wedge dx_2 \wedge dv_2 \wedge \dots \wedge dx_n \wedge dv_n.$$

On the other hand, a coorienting 2-vector bundle on $\mathbb{C}Ch(F_k) \cap \mathbb{C}I(\Delta_k)$ can continuously be extended to a coorienting 2-bundle on $\mathbb{C}Ch(F_k^\delta) \subset \mathbb{C}I(\delta)$, and the same for Δ_j. Indeed, let the complex straight line Λ

$$z_2 = \dots = z_n = xv, \quad z_1 = x_0 = \text{const}, \quad x \in (0, \infty),\ |v| = 1,$$

with small $z_0 > 0$ meet the hypersurfaces $\{F_k = 0\}$, $\{F_k^\delta = 0\}$ transversally. Then the family of surfaces Σ_λ, $\lambda \in [0, 1]$,

$$\left\{ \frac{\sum_{(i_1,\dots,i_n)\in\Delta_k} x^{i_2+\dots+i_n} x_0^{i_1} \lambda^{a-i_1} \cdot (i_1, i_2 v, \dots, i_n v)}{\sum_{(i_1,\dots,i_n)\in\Delta_k} x^{i_2+\dots+i_n} x_0^{i_1} \lambda^{a-i_1}}\ :\ x \in (0, \infty),\ |v| = 1 \right\}$$

is a diffeotopy, connecting $\Sigma_0 = \mathbb{C}\mu_\delta(\Lambda)$, which coorients $\mathbb{C}Ch(F_k^\delta)$ in $\mathbb{C}I(\delta)$, and $\Sigma_1 = \mathbb{C}\mu_{\Delta_k}(\Lambda)$, which coorients $\mathbb{C}Ch(F_k)$ in $\mathbb{C}I(\Delta_k)$.

Comparing this with the previous remark on orientations of $\mathbb{C}\delta$, one completes the proof of the orientability of a complex C-hypersurface.

At last, $[\mathbb{C}Ch(\mathcal{S},\mathcal{A})] \in H_{2n-2}(\mathbb{C}P^n)$ and $[\mathbb{R}Ch(\mathcal{S},\mathcal{A})] \in H_{n-1}(\mathbb{R}P^n, \mathbb{Z}/2)$ can easily be computed by induction considering the intersection of the complex and real charts with the coordinate planes. $\Box$

Proposition 2.7 *Any (equivariant) C-hypersurface M is (equivariantly) tame isotopic to a close smooth manifold M_{sm} of codimension 2 in $\mathbb{C}P^n$.*

Proof. Let $M = \mathbb{C}Ch(\mathcal{S},\mathcal{A})/S^1$, where $\mathcal{S} = \{\Delta_1, ..., \Delta_N\}$, $\mathcal{A} = \{A_{\overline{i}} \ : \ \overline{i} \in T^n_d \cap \mathbb{Z}^n\}$. Put $F_m = \sum_{\overline{i}\in\Delta_m} A_{\overline{i}}\overline{z}^{\overline{i}}$, $m = 1, ..., N$. We construct two nonvanishing $\mathbb{R}$-linearly independent sections $\overline{s}$ and $\overline{s}'$ of the bundle $T\mathbb{C}P^n\big|_{U(M)}$, where $U(M)$ is a neighborhood of M in $\mathbb{C}P^n$, such that the 2-subbundle $\text{Span}_\mathbb{R}\{\overline{s},\overline{s}'\} \subset T\mathbb{C}P^n\big|_M$ is transverse to $T\mathbb{C}Ch(F_m)$ in $T\mathbb{C}P^n\big|_{\mathbb{C}Ch(F_m)}$ for any $m = 1, ..., N$. This will imply the existence of a smooth $2(n-1)$-manifold M_{sm} isotopic (equivariantly, if $\text{Conj}(M) = M$) to M, close to M and transverse to the 2-bundle $\text{Span}_\mathbb{R}\{\overline{s},\overline{s}'\}$.

(i) First, we shift T^n_d into $\widetilde{T}^n_d = (1, ..., 1) + T^n_d$ (shifting respectively $\mathcal{S},\mathcal{A}$ as well), and construct sections $\overline{s}$ and $\overline{s}'$ of $T\mathbb{C}^n$ defined in a neighborhood of $\mathbb{C}Ch(\mathcal{S},\mathcal{A})$, with the above properties and an additional one: invariance with respect to the S^1 action on $\partial\mathbb{C}\widetilde{T}^n_d$. The latter allows us to obtain the required sections of $T\mathbb{C}P^n$ along the procedure described in the proof of Proposition 1.17.

(ii) Fix $m = 1, ..., N$ and consider the hypersurface $\widetilde{F}_m(\overline{z}) = F_m(e^{z_1}, ..., e^{z_n}) = 0$ in $\mathbb{C}^n$. Define vector fields $\overline{s}_1$ and $\overline{s}'_1$ on $\mathbb{C}^n$ by

$$\overline{s}_1(\overline{z}) = \frac{\text{Conj}(\text{grad}\widetilde{F}_m)}{|\text{Conj}(\text{grad}\widetilde{F}_m)|}, \quad \overline{s}'_1(\overline{z}) = \frac{\text{Conj}(\text{grad}\widetilde{F}_m)\sqrt{-1}}{|\text{Conj}(\text{grad}\widetilde{F}_m)\sqrt{-1}|} \ .$$

They do not vanish along $\{\widetilde{F}_m = 0\}$ and span a 2-bundle orthogonal to $T\{\widetilde{F}_m = 0\}$ in $T\mathbb{C}^n\big|_{\{\widetilde{F}_m=0\}}$. These vector fields are $2\pi\sqrt{-1}$-periodic in each coordinate $z_1, ..., z_n$, and their normalized images $\overline{s}$ and $\overline{s}'$ by the differential $D(\mathbb{C}\mu_{\Delta_m} \circ \exp)$ of the map $\mathbb{C}\mu_{\Delta_m} \circ \exp \, : \, \mathbb{C}^n \to \mathbb{C}\Delta_m$ give a 2-bundle $\text{Span}_\mathbb{R}\{\overline{s},\overline{s}'\}$ on $\mathbb{C}I(\Delta_m) \cap \mathbb{C}Ch(F_m)$ transverse to the tangent bundle $T(\mathbb{C}I(\Delta_m) \cap \mathbb{C}Ch(F_m))$. We will show that these vector fields continuously extend to $\bigcup_{m=1}^N (\partial\mathbb{C}\Delta_m)$.

(iii) Let δ be a face of Δ_m, $\overline{w}$ be a point in $\mathbb{C}Ch(F_m) \cap \mathbb{C}I(\delta)$, and $U_{\overline{w}} \subset \mathbb{C}\Delta_m$ be a neighborhood of $\overline{w}$ as introduced in the step (iv) of the proof of Proposition 1.4. We claim that, for any point $\overline{z} \in \mathbb{C}^n$ such that $\mathbb{C}\mu_\Delta(e^{\overline{z}}) \in U_{\overline{w}} \cap \mathbb{C}Ch(F_m) \cap \mathbb{C}I(\Delta_m)$, one has

$$\left|\left| \frac{D(\mathbb{C}\mu_{\Delta_m} \circ \exp)_{\overline{z}}}{||D(\mathbb{C}\mu_{\Delta_m} \circ \exp)_{\overline{z}}||} - \frac{D(\mathbb{C}\mu_\delta \circ \exp)_{\overline{\alpha}}}{||D(\mathbb{C}\mu_\delta \circ \exp)_{\overline{\alpha}}||} \right|\right| < K\varepsilon \ , \tag{15}$$

and

$$\left| \frac{\text{grad}(F_m \circ \exp)}{|\text{grad}(F_m \circ \exp)|}(\overline{z}) - \frac{\text{grad}(F^\delta_m \circ \exp)}{|\text{grad}(F^\delta_m \circ \exp)|}(\overline{\alpha}) \right| < K\varepsilon \ , \tag{16}$$

where $\overline{\alpha} \in \mathbb{C}^n$ is some point, satisfying $\mathbb{C}\mu_\Delta(\exp(\overline{\alpha})) = \overline{w}$, and $K > 0$ depends only on the coefficients of F and on the point $\overline{w}$.

(iv) To show (15), note that $\mathbb{C}\mu_{\Delta_m}\circ\exp$ splits into $\mu_{\Delta_m}\circ\exp:\mathbb{R}^n\to I(\Delta_m)$ and $\exp:\mathbb{R}^n\sqrt{-1}\to(S^1)^n$. Inequality (15) for the differential of $\exp:\mathbb{R}^n\sqrt{-1}\to(S^1)^n$ instead of $\mathbb{C}\mu_{\Delta_m}\circ\exp$ immediately follows from (6). Then, acting by a transformation from $SL(n,\mathbb{Z})$ and a shift, we move Δ_m,δ into $\widetilde{\Delta}_m,\widetilde{\delta}$ such that $\widetilde{\delta}$ lies in a coordinate s-plane. According to (7),

$$|e^{(\overline{i},\mathrm{Re}\overline{z})}-e^{(\overline{i},\mathrm{Re}\overline{\alpha})}|<K_1\varepsilon,\ \overline{i}\in\widetilde{\delta}\cap\mathbb{Z}^n,\quad e^{(\overline{i},\mathrm{Re}\overline{z})}<K_1\varepsilon,\ \overline{i}\in(\widetilde{\Delta}_m\backslash\widetilde{\delta})\cap\mathbb{Z}^n,$$

where $K_1>0$ depends only on Δ_m, thereby this implies (15), because $D(\mu_{\widetilde{\Delta}_m}\circ\exp)$ is represented by the matrix

$$\left(\frac{\sum_{\overline{i}\in\widetilde{\Delta}_m}i_pi_qe^{(\overline{i},\mathrm{Re}\overline{z})}\cdot\sum_{\overline{i}\in\widetilde{\Delta}_m}e^{(\overline{i},\mathrm{Re}\overline{z})}-\sum_{\overline{i}\in\widetilde{\Delta}_m}i_pe^{(\overline{i},\mathrm{Re}\overline{z})}\cdot\sum_{\overline{i}\in\widetilde{\Delta}_m}i_qe^{(\overline{i},\mathrm{Re}\overline{z})}}{(\sum_{\overline{i}\in\widetilde{\Delta}_m}e^{(\overline{i},\mathrm{Re}\overline{z})})^2}\right)_{p,q=1,\ldots,n},$$

and $D(\mu_{\widetilde{\delta}}\circ\exp)$ is not trivial.

(v) Similarly, $F_m(e^{\overline{z}})=e^{(\overline{i}_0,\overline{z})}\sum_{\overline{i}\in\Delta}A_{\overline{i}}e^{(\overline{i},\overline{z})}$, where by (6) and (7)

$$|e^{(\overline{i},\overline{z})}-e^{(\overline{i},\overline{\alpha})}|<K_2\varepsilon,\ \overline{i}\in\delta\cap\mathbb{Z}^n,\quad |e^{(\overline{i},\overline{z})}|<K_2\varepsilon,\ \overline{i}\in(\Delta\backslash\delta)\cap\mathbb{Z}^n,$$

where $K_2>0$ depends on Δ_m. So, we obtain

$$\mathrm{grad}(F_m\circ\exp)(\overline{z})=e^{(\overline{i}_0,\overline{z})}\left(\sum_{\overline{i}\in\delta}A_{\overline{i}}\overline{i}e^{(\overline{i},\overline{z})}+\sum_{\overline{i}\in\Delta\backslash\delta}A_{\overline{i}}\overline{i}e^{(\overline{i},\overline{z})}\right),$$

which immediately implies (16), because $\sum_{\overline{i}\in\delta}A_{\overline{i}}\overline{i}e^{(\overline{i},\overline{\alpha})}=\mathrm{grad}(F_m^\delta\circ\exp)(\overline{\alpha})\ne0$.

(vi) Relations (15) and (16) provide continuous extension of the vector fields $\overline{s}$ and $\overline{s}'$ on $\bigcup_{m=1}^N(\partial\mathbb{C}\Delta_m)$ so that they remain $\mathbb{R}$-linearly independent and belong to $T\mathbb{C}I(\delta)$ along $\mathbb{C}I(\delta)$ for any proper face δ of Δ_m, $m=1,...,N$. Moreover, the restrictions of $\overline{s}$ and $\overline{s}'$ on $\mathbb{C}I(\delta)$ depend only on δ and F_m^δ, common for all $\Delta_m\supset\delta$, hence they are compatible with S^1 action on $\partial\mathbb{C}\widetilde{T}_d^n$, and we are done. $\Box$

Proposition 2.8 *Given a (equivariant) C-hypersurface M, the tangent bundle to its smoothing M_{sm} is (equivariantly) isotopic to a (equivariant) bundle of complex hyperplanes. In particular, M_{sm} possesses an (equivariant) almost complex structure. If M is real, the above isotopy has fixed intersection with $T\mathbb{R}P^n\big|_{M_{sm}}$ (equal to $T\mathbb{R}M_{sm}$).*

Proof. For any $\overline{w}\in M_{sm}$ denote by $\overline{s}_2(\overline{w})$ the projection of $\overline{s}(\overline{w})$ to $(T_{\overline{w}}M_{sm})^\perp\subset T_{\overline{w}}\mathbb{C}P^n$. This is a nonvanishing section of $T_{\overline{w}}\mathbb{C}P^n\big|_{M_{sm}}$, which in the equivariant case is real along $\mathbb{R}M_{sm}=\mathbb{R}P^n\cap M_{sm}$ and orthogonal to $T_{\overline{w}}\mathbb{R}M_{sm}$. Assuming that $\overline{w}=\mathbb{C}\mu_{\Delta_m}(\overline{z})\in\mathbb{C}\Delta_m$, one obtains $\overline{s}_2(\overline{w})=D(\mathbb{C}\mu_{\Delta_m})_{\overline{z}}(\overline{b}(\overline{z}))$, where $\overline{b}(\overline{z})\in T_{\overline{z}}(\mathbb{C}^*)^n\backslash T_{\overline{z}}(\{F_m=\mathrm{const}\})$. Clearly, the 2-plane $\mathrm{Span}_{\mathbb{R}}\{\overline{b}(\overline{z}),\overline{b}(\overline{z})\sqrt{-1}\}$ is transversal to $T_{\overline{z}}(\{F_m=\mathrm{const}\})$ in $T_{\overline{z}}(\mathbb{C}^*)^n$. Hence the two plane $L^2_{\overline{w}}=\mathrm{Span}_{\mathbb{R}}\{\overline{s}_2(\overline{w}),\overline{s}_2'(\overline{w}\}$, where $\overline{s}_2'(\overline{w})=D(\mathbb{C}\mu_{\Delta_m})_{\overline{z}}(\overline{b}(\overline{z})\sqrt{-1})$, is transversal to $T_{\overline{w}}M_{sm}$ in $T_{\overline{w}}\mathbb{C}P^n$.

We construct a required isotopy in two steps.

First, we (equivariantly) move $T_{\overline{w}}M_{sm}$ to $L^{2n-2}_{\overline{w}} = (L^2_{\overline{w}})^\perp \subset T_{\overline{w}}\mathbb{C}P^n$, for any $\overline{w} \in M_{sm}$. This, evidently, is possible, since by construction, $T_{\overline{w}}M_{sm}$ is transverse to $L^2_{\overline{w}}$.

Now we show that it is possible to move the distribution L^{2n-2} on M_{sm} to the distribution of complex hyperplanes $(\mathrm{Span}_{\mathbb{R}}\{\overline{s}_2, \overline{s}_2\sqrt{-1}\})^\perp$ in $T\mathbb{C}P^n\big|_{M_{sm}}$.

Lemma 2.9 *In the above notation, for any $\overline{w} \in M_{sm}$, no vector $(1-\lambda)\overline{s}_2'(\overline{w}) + \lambda\overline{s}_2\sqrt{-1}$, $0 \le \lambda \le 1$, is $\mathbb{R}$-proportional to $\overline{s}_2(\overline{w})$.*

Proof. Since $\exp : \mathbb{C}^n \to (\mathbb{C}^*)^n$ preserves the complex structure on $T\mathbb{C}^n$ and $T(\mathbb{C}^*)^n$, we can replace $D(\mathbb{C}\mu_{\Delta_m})$ in the above formulae by $D(\widetilde{\mathbb{C}\mu}_{\Delta_m})$, where $\widetilde{\mathbb{C}\mu}_{\Delta_m} : \mathbb{C}^n \to \mathbb{C}^n$ is defined by

$$\mathbb{C}^n = \mathbb{R}^n \oplus \mathbb{R}^n\sqrt{-1} \overset{\log\widetilde{\mu}_{\Delta_m}\oplus\mathrm{Id}}{\longrightarrow} \mathbb{R}^n \oplus \mathbb{R}^n\sqrt{-1} = \mathbb{C}^n \ , \tag{17}$$

$$\log\widetilde{\mu}_{\Delta_m} = (\log\mu_{1,\Delta_m}, ..., \log\mu_{n,\Delta_m}), \quad \mu_{\Delta_m} = (\mu_{1,\Delta_m}, ..., \mu_{n,\Delta_m}) \ .$$

Clearly $\exp\circ\widetilde{\mathbb{C}\mu}_{\Delta_m} = \mathbb{C}\mu_{\Delta_m}\circ\exp$.

Assume that $\overline{s}_2(\overline{w}) = (1-\lambda)\overline{s}_2'(\overline{w}) + \lambda\overline{s}_2(\overline{w})\sqrt{-1}$ for some $\lambda \in (0,1)$ and $\overline{w} \in \bigcup_{m=1}^N \mathbb{C}I(\Delta_m)$. Splitting $\overline{b}(\overline{w})$ into $\overline{b}_r + \overline{b}_i\sqrt{-1}$ with $\overline{b}_r, \overline{b}_i \in \mathbb{R}^n$, we obtain in view of (17)

$$\begin{aligned} \overline{s}_2(\overline{w}) &= D(\log\mu_{\Delta_m})(\overline{b}_r) + \overline{b}_i\sqrt{-1}, \\ \overline{s}_2(\overline{w})\sqrt{-1} &= -\overline{b}_i + D(\log\mu_{\Delta_m})(\overline{b}_r)\sqrt{-1}, \\ \overline{s}_2'(\overline{w}) &= -D(\log\mu_{\Delta_m})(\overline{b}_i) + \overline{b}_r\sqrt{-1} \ . \end{aligned}$$

The latest equality splits in two equations for $\overline{b}_r, \overline{b}_i$ which imply that

$$((\lambda-\lambda^2)\mathrm{Id} + 2(1+\lambda+\lambda^2)D(\log\mu_{\Delta_m}) + (\lambda-\lambda^2)D(\log\mu_{\Delta_m})^2)(\overline{b}_r) = 0, \quad \overline{b}_r \ne 0 \ ;$$

hence the operator $D(\log\mu_{\Delta_m})$ has a negative eigenvalue

$$\frac{-(1+\lambda+\lambda^2) \pm \sqrt{(1+\lambda+\lambda^2)^2 - (\lambda-\lambda^2)^2}}{\lambda-\lambda^2} \ ,$$

but this is impossible. Indeed, the matrix of $D(\log\mu_{\Delta_m})$ is a product of

$$A = \left(\frac{\sum_{\overline{i}\in\Delta} i_p i_q e^{(\overline{i},\overline{x})} \cdot \sum_{\overline{i}\in\Delta} e^{(\overline{i},\overline{x})} - (\sum_{\overline{i}\in\Delta} i_p e^{(\overline{i},\overline{x})})(\sum_{\overline{i}\in\Delta} i_q e^{(\overline{i},\overline{x})})}{(\sum_{\overline{i}\in\Delta} e^{(\overline{i},\overline{x})})^2}\right)_{p,q=1,...,n} \ ,$$

$$B = \mathrm{diag}\left(\frac{\sum_{\overline{i}\in\Delta} i_p e^{(\overline{i},\overline{x})}}{\sum_{\overline{i}\in\Delta} e^{(\overline{i},\overline{x})}}\right)_{p=1,...,n} \ .$$

Here both A and B are positive definite, for instance,

$$(A\overline{b}, \overline{b}) = \frac{\sum_{\overline{i},\overline{j}\in\Delta_m} e^{(\overline{i},\overline{x})}e^{(\overline{j},\overline{x})}(\sum_{p=1}^n (i_p - j_p)b_p)^2}{(\sum_{\overline{i}\in\Delta_m} e^{(\overline{i},\overline{x})})^2} > 0 \ ,$$

hence AB rotates any vector by an angle $< \pi$, and we are done.

The same argument prove the required statement when $\overline{w} \in \bigcup_{m=1}^N(\partial\mathbb{C}\Delta_m)$ in view of (15) and (16). $\Box$

2.3 Algebraic covering

Introduce the map $\Pi_m : \mathbb{C}P^n \to \mathbb{C}P^n$, $[z_0 : \ldots : z_n] \mapsto [z_0^m : \ldots : z_n^m]$.

Proposition 2.10 *For any C-hypersurface M of degree d in $\mathbb{C}P^n$ there exists a number $m_0 > 0$ such that for any integer $m > m_0$ the preimage $\Pi_m^{-1}(M)$ of M is a PL-variety tame isotopic to a smooth algebraic hypersurface of degree md in $\mathbb{C}P^n$.*

Proof. Let M be defined by polynomials $F_i(z_1, ..., z_n)$, $i = 1, ..., N$, with Newton polyhedra $\Delta_1, ..., \Delta_N$, where $\Delta_1 \cup ... \cup \Delta_N = T_d^n$. The polynomials

$$\widetilde{F}_i(z_1, ..., z_n) = F_i(z_1^m, ..., z_n^m), \quad i = 1, ..., N,$$

have the Newton polyhedra $m\Delta_i$, $i = 1, ..., N$, and define a C-hypersurface $\widetilde{M}$ of degree md.

Lemma 2.11 *The variety $\widetilde{M}$ is tame isotopic in $\mathbb{C}P^n$ to $\Pi_m^{-1}(M)$.*

Proof. Let $\Delta \subset \mathbb{R}^n$ be a polyhedron of dimension n. Let us show that the diffeomorphism $\mu_\Delta \Pi_m \mu_{m\Delta}^{-1} : I(m\Delta) \to I(\Delta)$, where $\Pi_m : (\mathbb{R}_+)^n \to (\mathbb{R}_+)^n$ acts as $(x_1, ..., x_n) \mapsto (x_1^m, ..., x_n^m)$, extends to a homeomorphism $\varphi : m\Delta \to \Delta$ such that, for any proper face $\sigma \subset \Delta$, the map $\varphi|_{m\sigma} : m\sigma \to \sigma$ is a homeomorphism depending only on σ (and not on Δ). Indeed, since Π_m commutes with $SL_n(\mathbb{Z})$ acting on $(\mathbb{R}_+)^n$, we can replace Δ by its image under some $g \in SL_n(\mathbb{Z})$ such that the given face σ will lie on the coordinate hyperplane $\{i_1 = 0,\ i_2 \cdot ... \cdot i_n \neq 0\}$. Then $\mu_{t(\Delta)}$ and $\mu_{mt(\Delta)}$ extend by the formulas similar to (1) to homeomorphisms of $(\mathbb{R}_+)^n \cup \{x_1 = 0,\ x_2 \cdot ... \cdot x_n \neq 0\}$ and $I(t(\Delta)) \cup I(t(\sigma))$ and $I(mt(\Delta)) \cup I(mt(\sigma))$, respectively. This gives us a homeomorphism $I(m\Delta) \cup I(m\sigma) \to I(\Delta) \cup I(\sigma)$, which does not depend on the choice of t, and we are done.

Consider now the following commutative diagram:

$$\begin{array}{ccccc}
\coprod_{1\le k\le N}(\mathbb{C}^*)^n & \xrightarrow{\coprod_{1\le k\le N}\mathbb{C}\mu_{m\Delta_k}} & \mathbb{C}T^n_{md} & \xrightarrow{\mathbb{C}\mu} & \mathbb{C}P^n \\
\downarrow{\scriptstyle \coprod_{1\le k\le N}\Pi_m} & & \downarrow{\scriptstyle \varphi_m} & & \downarrow{\scriptstyle \widetilde{\Pi}_m} \\
\coprod_{1\le k\le N}(\mathbb{C}^*)^n & \xrightarrow{\coprod_{1\le k\le N}\mathbb{C}\mu_{\Delta_k}} & \mathbb{C}T^n_{d} & \xrightarrow{\mathbb{C}\mu} & \mathbb{C}P^n
\end{array}$$

where $\coprod_{1\le k\le N}(\mathbb{C}^*)^n$ means the disjoint union of N copies of $(\mathbb{C}^*)^n$. The map φ_m is defined on any $\mathbb{C}\Delta_k$ as follows:

$$(x_1, ..., x_n) \in \Delta_k \mapsto \mu_{\Delta_k}\Pi_m\mu_{m\Delta_k}^{-1}(x_1, ..., x_n) \in \Delta_k,$$

$$(v_1, ..., v_n) \in (S^1)^n \mapsto (v_1^m, ..., v_n^m) \in (S^1)^n\ ,$$

with $\mu_{\Delta_k}\Pi_m\mu_{m\Delta_k}^{-1}$ extended on the whole Δ_k. This definition is correct since the extensions coming from Δ_k and Δ_j with a common face are the same on the common face, as shown above. The map $\widetilde{\Pi}_m$ is defined by this diagram. Let us show that $\widetilde{\Pi}_m$

is tame isotopic to Π_m. First, $\Pi_m(x_0 v_0, ..., x_n v_n) = (x_0^m v_0^m, ..., x_n^m v_n^m)$ is tame isotopic to $\pi'_m(x_0 v_0, ..., x_n v_n) = (x_0 v_0^m, ..., x_n v_n^m)$. On the other hand, the homeomorphism

$$\coprod_{1 \le k \le N} \mu_{\Delta_k} \Pi_m \mu_{m\Delta_k}^{-1} : T_{md}^n \to T_d^n$$

in the definition of φ_m is tame isotopic to the homothety, turning φ_m into

$$\varphi'_m(x_1 v_1, ..., x_n v_n) = \frac{1}{m}(x_1 v_1^m, ..., x_n v_n^m) ,$$

and completing the proof of Lemma, since $\pi'_m = \mathbb{C}\mu \circ \varphi'_m \circ (\mathbb{C}\mu)^{-1}$, and $\varphi_m^{-1}(\mathbb{C}Ch(F_k)) = \mathbb{C}Ch(F_k(z_1^m, ..., z_n^m))$. $\square$

Lemma 2.12 *There exists m_0 such that, for any $m \ge m_0$, the subdivision $T_{md}^n = m\Delta_1 \cup ... \cup m\Delta_N$ admits a convex refinement.*

Proof. Let Γ be the graph of a smooth convex function of n variables, say $f(i_1, ..., i_n) = i_1^2 + ... + i_n^2$, and let $\mathrm{pr} : \Gamma \to \mathbb{R}^n$ be the projection. Denote by $\mathrm{sk}^1(\Delta)$ the 1-skeleton of the subdivision $\Delta_1 \cup ... \cup \Delta_N$. Clearly, $\mathrm{pr}^{-1}\left(\bigcup_k \mathrm{sk}^1(\Delta_k)\right)$ lies on the boundary of its convex hull. The same is true for

$$\mathrm{pr}^{-1}\left(\bigcup_{k=1}^N \mathrm{sk}^1(\Delta_k) \cap \frac{1}{m}\mathbb{Z}^n\right) . \qquad (18)$$

If m is big enough, one can define a required refinement by the piecewise-linear convex function whose graph is the lower part of the boundary of the convex hull of the set (18). $\square$

Now to finish the proof of Proposition 2.10 it remains to apply Theorem 2.2 and Proposition 2.3. $\square$

Denote by $\chi(X)$ the Euler characteristic of X, by $\mathrm{sign}(X)$ the signature of X if X is a manifold whose dimension is divisible by 4, and by χ_d^n (resp., sign_d^n, if n is odd) the Euler characteristic (resp., the signature) of a smooth algebraic hypersurface of degree d in $\mathbb{C}P^n$.

Corollary 2.13 *Any C-hypersurface M of degree d in $\mathbb{C}P^n$ satisfies*

$$\chi(M) = \chi_d^n, \quad \mathrm{sign}(M) = \mathrm{sign}_d^n .$$

Proof.

(i) The equality $\chi(M) = \chi_d^n$ follows immediately by induction from Proposition 2.10 and the behavior of the Euler characteristic under ramified coverings.

(ii) Suppose that n is odd and consider a smooth algebraic hypersurface M' of degree d in $\mathbb{C}P^n$. Let m_0 be as in Proposition 2.10. Take any prime number

$p > m_0$. Note that $\widetilde{M'} = \Pi_p^{-1}(M') \subset \mathbb{C}P^n$ is an algebraic hypersurface of degree pd tame isotopic to $\widetilde{M} = \Pi_p^{-1}(M)$. The deck transformation group of the coverings $\Pi_p : \widetilde{M} \to M$ and $\Pi_p : \widetilde{M'} \to M'$ is $G \simeq (\mathbb{Z}/p)^n$. By formula (19) [17]

$$\mathrm{sign}(M) = \frac{1}{|G|}\sum_{g\in G}\mathrm{sign}(g, \widetilde{M}), \quad \mathrm{sign}(M') = \frac{1}{|G|}\sum_{g\in G}\mathrm{sign}(g, \widetilde{M'}) \ . \tag{19}$$

For $g = \mathrm{Id} \in G$, we have $\mathrm{sign}(g, \widetilde{M}) = \mathrm{sign}(\widetilde{M}) = \mathrm{sign}(\widetilde{M'}) = \mathrm{sign}(g, \widetilde{M'})$. Pick $g \neq \mathrm{Id}$. The twisted signatures $\mathrm{sign}(g, \widetilde{M})$ and $\mathrm{sign}(g, \widetilde{M'})$ depend on the embedding of $\mathrm{Fix}(g)$ in $\widetilde{M}$ and $\widetilde{M'}$, respectively, and the action of g in the tangent and normal bundles of $Y = \mathrm{Fix}(g) \subset \widetilde{M}$ and $Y' = \mathrm{Fix}(g) \subset \widetilde{M'}$ (see Theorem 6.12 [3]). We do not compute the twisted signatures along Theorem 6.12 [3], but use Hirzebruch's formula for the signature of ramified coverings. Namely, let $\langle g\rangle \subset G$ be the cyclic group of order p generated by g. According to [16],

$$p \cdot \mathrm{sign}(\widetilde{M}/\langle g\rangle) = \sum_{i=0}^{p-1}\mathrm{sign}(g^i, \widetilde{M})$$

is a universal function of p and the signatures of self-intersections $Y \circ Y$, $Y \circ Y \circ Y$, ... Note that $Y = \mathrm{Fix}(g) \subset \widetilde{M}$ and $Y' = \mathrm{Fix}(g) \subset \widetilde{M'}$ are the intersections of $\widetilde{M}$ and $\widetilde{M'}$, respectively, with the same collection of coordinate planes. Hence, due to the tame isotopy of $\widetilde{M}$ and $\widetilde{M'}$, one has $\mathrm{sign}(Y \circ Y) = \mathrm{sign}(Y' \circ Y')$, $\mathrm{sign}(Y \circ Y \circ Y) = \mathrm{sign}(Y' \circ Y' \circ Y')$, and so on. Therefore for any $g \in G$ we have

$$\sum_{i=0}^{p-1}\mathrm{sign}(g^i, \widetilde{M}) = \sum_{i=0}^{p-1}\mathrm{sign}(g^i, \widetilde{M'}) \ ,$$

which immediately implies the required equality of the right hand sides in (19), since $G \simeq (\mathbb{Z}/p)^n$ can be decomposed in the union of cyclic subgroups such that the only intersection of any two of these subgroups is $\mathrm{Id} \in G$. $\square$

Corollary 2.14 *Let M be a C-hypersurface of degree d in $\mathbb{C}P^n$. Then*

- *M is simply connected if $n > 2$,*
- *$\pi_1(\mathbb{C}P^n \backslash M) = \mathbb{Z}/d\mathbb{Z}$ if $n \geq 2$.*

Proof.

(i) To show that M is simply connected for $n > 2$, note that an algebraic hypersurface of dimension greater than 1 is simply connected. Let $m > m_0$ be as in Proposition 2.10. Consider a loop γ in M. We can move it slightly so that it does not meet the coordinate hyperplanes in $\mathbb{C}P^n$. Since $\Pi_m^{-1}(M)$ is simply connected, $m^n \cdot [\gamma]$ is contractible in M. Similarly, $(m+1)^n \cdot [\gamma]$ is contractible in M, and we are done, because m^n and $(m+1)^n$ are coprime.

(ii) Since an affine nonsingular algebraic hypersurface in $\mathbb{C}^{n+1}$, $n \ge 2$, is simply connected, the previous argument shows that $\hat{M} \backslash \mathbb{C}P^n \subset \mathbb{C}^{n+1}$ is simply connected, where $\hat{M}$ is a C-hypersurface in $\mathbb{C}P^{n+1}$ and $\mathbb{C}P^n$ is a coordinate hyperplane in $\mathbb{C}P^{n+1}$.

Let a C-hypersurface M of degree d in $\mathbb{C}P^n$ be defined by a subdivision $\mathcal{S}:\ T_d^n = \Delta_1 \cup ... \cup \Delta_N$ and collection of numbers $\mathcal{A} : T_d^n \cap \mathbb{Z}^n \to \mathbb{C}$. Embed T_d^n into T_d^{n+1} as the face $T_d^{n+1} \cap \{i_{n+1} = 0\}$, take the subdivision $\widetilde{\mathcal{S}}:\ T_d^{n+1} = \widetilde{\Delta}_1 \cup ... \cup \widetilde{\Delta}_N$ with $\widetilde{\Delta}_k$ being the cone over Δ_k with the vertex $(0, ..., 0, d)$, $k = 1, ..., N$, and define $\widetilde{\mathcal{A}}$: $T_d^{n+1} \cap \mathbb{Z}^{n+1} \to \mathbb{C}$ by $\widetilde{A}_{i_1...i_n 0} = A_{i_1...i_n}$, $\widetilde{A}_{0...0d} = -1$, $\widetilde{A}_{i_1...i_n i_{n+1}} = 0$ if $0 < i_{n+1} < d$. These data define nondegenerate polynomials $\widetilde{F}_k(z_1, ..., z_{n+1}) = F_k(z_1, ..., z_n) - z_{n+1}^d$ with Newton polyhedra $\widetilde{\Delta}_k$, $k = 1, ..., N$, whose charts can be glued into a C-hypersurface $\widetilde{M}$ of degree d in $\mathbb{C}P^{n+1}$. The required statement $\pi_1(\mathbb{C}P^n \backslash M) = \mathbb{Z}/d\mathbb{Z}$ is a corollary of the following statement.

Lemma 2.15 *There is a $\mathbb{Z}/d\mathbb{Z}$-covering $\widetilde{M} \backslash \{z_{n+1} = 0\} \to \mathbb{C}P^n \backslash M$.*

Proof. We construct a $\mathbb{Z}/d\mathbb{Z}$-covering $\Phi : \mathbb{C}Ch(\widetilde{\mathcal{S}}, \widetilde{\mathcal{A}}) \backslash \{w_{n+1} = 0\} \to \mathbb{C}T_d^n \backslash \mathbb{C}Ch(\mathcal{S}, \mathcal{A})$ and then show that it commutes with the S^1-action on $\partial\mathbb{C}Ch(\widetilde{\mathcal{S}}, \widetilde{\mathcal{A}}) \backslash \{z_{n+1} = 0\}$ and $\partial\mathbb{C}T_d^n \backslash \mathbb{C}Ch(\mathcal{S}, \mathcal{A})$, thereby defining the required covering.

Let $(w_1, ..., w_n, w_{n+1}) \in \mathbb{C}Ch(\widetilde{\mathcal{S}}, \widetilde{\mathcal{A}})$, $w_{n+1} \ne 0$. Then $(w_1, ..., w_{+1}) \in \mathbb{C}I(\widetilde{\delta})$, where $\widetilde{\delta}$ is a cone over a face δ of Δ_k, $1 \le k \le N$, with the vertex at $(0, ..., 0, d)$. Hence $(w_1, ..., w_{n+1}) = \mathbb{C}\mu_{\widetilde{\delta}}(z_1, ..., z_{n+1})$ for some $(z_1, ..., z_{n+1}) \in (\mathbb{C}^*)^{n+1}$. So, we define

$$\Phi(w_1, ..., w_{n+1}) = \mathbb{C}\mu_\delta(z_1, ..., z_n) \ .$$

The map Φ is well defined. If $\delta = \Delta_1, \ldots, \Delta_N$, then $\mathbb{C}\mu_{\widetilde{\delta}}$ and $\mathbb{C}\mu_\delta$ are diffeomorphisms. If δ lies in an $(n-1)$-plane $\alpha_1 i_1 + ... + \alpha_n i_n = \beta$, $i_{n+1} = 0$, then $\widetilde{\delta}$ lies in an n plane $\alpha_1 i_1 + ... + \alpha_n i_n + \beta i_n/d = \beta$. This means that $(\mathbb{C}\mu_{\widetilde{\delta}})^{-1}(w_1, ..., w_{n+1})$ contains the family $(z_1 t^{\alpha_1}, ..., z_n t^{\alpha_n}, z_{n+1}^{\beta/d})$, $t \in \mathbb{R}_+^*$, but $\mathbb{C}\mu_\delta$ takes the family $(z_1 t^{\alpha_1}, ..., z_n t^{\alpha_n})$, $t \in \mathbb{R}_+^*$, to one point.

The map Φ is continuous. If, for some $k = 1, ..., N$ and a curve $\widetilde{\gamma}(t) \in (\mathbb{C}^*)^{n+1}$:

$$(\lambda_1 t^{k_1} + O(t^{k_1+1}), ..., \lambda_n t^{k_n} + O(t^{k_n+1}), \lambda_{n+1} t^{k_{n+1}} + O(t^{k_{n+1}+1})), \quad t > 0,$$

one has $\lim_{t\to 0} \mathbb{C}\mu_{\widetilde{\Delta}_k}(\widetilde{\gamma}(t)) = \mathbb{C}\mu_{\widetilde{\delta}}(\lambda_1, ..., \lambda_n, \lambda_{n+1})$, then $\lim_{t\to 0} \mathbb{C}\mu_{\Delta_k}(\gamma(t)) = \mathbb{C}\mu_\delta(\lambda_1, ..., \lambda_n)$, where

$$\gamma(t) = (\lambda_1 t^{k_1} + O(t^{k_1+1}), ..., \lambda_n t^{k_n} + O(t^{k_n+1})) \in (\mathbb{C}^*)^n, \quad t > 0 \ .$$

The map Φ is surjective. Indeed, if $(w_1, ..., w_n) \in \mathbb{C}T_d^n \backslash \mathbb{C}Ch(\mathcal{S}, \mathcal{A})$, then $(w_1, ..., w_n) = \mathbb{C}\mu_\delta(z_1, ..., z_n)$ for a face δ of Δ_k $(1 \le k \le N)$, where $(z_1, ..., z_n) \in (\mathbb{C}^*)^n$, $F_k^\delta(z_1, ..., z_n) \ne 0$. Then there exists $z_{n+1} \ne 0$ such that $F_k^\delta(z_1, ..., z_n) = z_{n+1}^d$; hence $\mathbb{C}\mu_{\widetilde{\delta}}(z_1, ..., z_n, z_{n+1})$ belongs to $\mathbb{C}Ch(F_k^{\widetilde{\delta}}) \backslash \{w_{n+1} = 0\}$ and $\Phi(\mathbb{C}\mu_{\widetilde{\delta}}(z_1, ..., z_n, z_{n+1})) = (w_1, ..., w_n)$.

The map Φ is a $\mathbb{Z}/d\mathbb{Z}$-covering. Indeed, for any point $(w_1, ..., w_n) \in \mathbb{C}T_d^n \backslash \mathbb{C}Ch(\mathcal{S}, \mathcal{A})$ its preimage $\Phi^{-1}(w_1, ..., w_n)$ consists of d distinct points $(w'_1, ..., w'_n, w'_{n+1}\omega)$ with some fixed $w'_1, ..., w'_{n+1}$ and any d-th root of unity ω.

At last, note that if $\Phi(w'_1, ..., w'_n, w'_{n+1}) = (w_1, ..., w_n)$, then $\Phi(w'_1 v, ..., w'_n v, w'_{n+1} v) = (w_1 v, ..., w_n v)$ for arbitrary $v \in S^1$, which completes the proof. $\Box$

Proposition 2.16 *Let $M \subset \mathbb{C}P^n$ be a C-hypersurface and $M' \subset \mathbb{C}P^n$ a nonsingular algebraic hypersurface, both of degree d. Then $H_*(M)$ and $H_*(M')$ are isomorphic as graded groups.*

Proof. Take m_0 as in Proposition 2.10. Choose $m > m_0$ in such a way that m is coprime with all the orders of elements in $\text{Tors}H_*(M)$, and consider $M_m = \Pi_m^{-1}(M)$. Pick a homology class $\alpha \in H_i(M)$. Note that $m^n \alpha$ belongs to the image of $(\Pi_m)_* : H_i(M_m) \to H_i(M)$. For an odd i different from $n-1$ we have $H_i(M_m) = 0$ and, hence, $H_i(M) = 0$. In particular, $H_*(M)$ has no torsion. Any group $H_i(M)$ with even i $(0 \le i \le 2n-2)$ contains a nontrivial element: the fundamental class of the intersection of M with coordinate hyperplanes taken in appropriate number. Thus, $H_i(M)$ is isomorphic to $H_i(M_m) \simeq \mathbb{Z}$ if i is even and different from $n-1$. The fact that the groups $H_{n-1}(M)$ and $H_{n-1}(M')$ are isomorphic follows now from the equality $\chi(M) = \chi(M')$ proven in Corollary 2.13. $\Box$

Proposition 2.17 *Let n be a positive odd number, M a C-hypersurface of degree d in $\mathbb{C}P^n$ and M' a nonsingular algebraic hypersurface of degree d in $\mathbb{C}P^n$. Then the lattices $(H_{n-1}(M), B_M)$ and $(H_{n-1}(M'), B_{M'})$ (where $B_M : H_{n-1}(M) \times H_{n-1}(M) \to \mathbb{Z}$ and $B_{M'} : H_{n-1}(M') \times H_{n-1}(M') \to \mathbb{Z}$ are the intersection forms on M and M', respectively) are isomorphic.*

Proof. The lattices $H_{n-1}(M)$ and $H_{n-1}(M')$ are unimodular and have the same rank and signature (Corollary 2.13). It remains to show that these two lattices have the same parity, since for $d = 1$ we have $H_{n-1}(M) \simeq H_{n-1}(M') \simeq \mathbb{Z}$, and for $d \ge 2$ the lattice $H_{n-1}(M')$ is indefinite. Let us show, first, that the lattices $H_{n-1}(M)$ and $H_{n-1}(\widetilde{M})$ have the same parity, where $\widetilde{M} = \Pi_m^{-1}(M)$, m_0 is as in Proposition 2.10 and $m > m_0$ is odd.

(i) If α^2 is odd, $\alpha \in H_{n-1}(M)$, then $((\Pi_m)^! \alpha)^2 = m^{2n} \alpha^2$ is odd as well.

(ii) If β^2 is odd, $\beta \in H_{n-1}(\widetilde{M})$, then γ^2 is odd, where $\gamma = \sum_{g \in G} g_* \beta$, and $G = (\mathbb{Z}/m)^n$ is the deck transformation group of Π_m. Furthermore, $\gamma = (\Pi_m)^! \alpha$ for certain $\alpha \in H_{n-1}(M)$, and α^2 is odd.

Similarly, the lattices $H_{n-1}(M')$ and $H_{n-1}(\widetilde{M'})$, where $\widetilde{M'} = \Pi_m^{-1}(M')$, have the same parity. Since $\widetilde{M}$ and $\widetilde{M'}$ are isotopic in $\mathbb{C}P^n$, we obtain that the lattices $H_{n-1}(M)$ and $H_{n-1}(M')$ have the same parity. $\Box$

Corollary 2.18 *A C-surface of degree d in $\mathbb{C}P^3$ is homeomorphic to a nonsingular algebraic surface of degree d in $\mathbb{C}P^3$.*

Proof. Note that a C-surface of degree d in $\mathbb{C}P^3$ is simply connected (see 2.13), smoothable (see Proposition 2.7) and have the same intersection form as an algebraic surface of degree d in $\mathbb{C}P^3$ (see 2.17). It remains to apply Freedman's theorem (see, for example, [11]). $\square$

2.4 Ramified double coverings

Proposition 2.19 *Let M be a C-hypersurface of even degree d in $\mathbb{C}P^n$. Then there exists a closed simply connected $2n$-dimensional PL-manifold Y and a map $\Pi : Y \to \mathbb{C}P^n$ which is a double covering ramified along M.*

Proof. The existence of a ramified double covering $\Pi : Y \to \mathbb{C}P^n$ with Y being a PL-manifold follows generically from the fact that $[M] \in H_{2n-2}(\mathbb{C}P^n)$ is an even class. However, we prefer to give an explicit construction of Y.

Namely, we repeat the construction of step (ii) in the proof of Corollary 2.14, taking the simplex $T^{n+1}_{d,2} \subset \mathbb{R}^{n+1}$ with vertices $(0,...,0)$, $(d,0,...,0)$, ..., $(0,...,0,d,0)$, $(0,...,0,2)$ instead of T^{n+1}_d. So, T^n_d embeds into $T^{n+1}_{d,2}$ as the face $T^{n+1}_{d,2} \cap \{i_{n+1} = 0\}$, the subdivision $T^{n+1}_{d,2} = \widetilde{\Delta}_1 \cup ... \cup \widetilde{\Delta}_N$ is defined as the cone over the subdivision $T^n_d = \Delta_1 \cup ... \cup \Delta_N$ with the vertex at $(0,...,0,2)$, and $\widetilde{\mathcal{A}} : T^{n+1}_{d,2} \cap \mathbb{Z}^{n+1} \to \mathbb{C}$ is defined by $\mathcal{A}_{i_1...i_n0} = A_{i_1...i_n}$, $\widetilde{\mathcal{A}}_{i_1...i_n1} = 0$, $\widetilde{\mathcal{A}}_{0...02} = -1$. These data define a PL-manifold $\mathbb{C}Ch(\widetilde{\mathcal{S}}, \widetilde{\mathcal{A}})$ with boundary $\partial\mathbb{C}Ch(\widetilde{\mathcal{S}}, \widetilde{\mathcal{A}}) = \mathbb{C}Ch(\widetilde{\mathcal{S}}, \widetilde{\mathcal{A}}) \cap \partial\mathbb{C}T^{n+1}_{d,2}$, which is the gluing of the charts of the polynomials $\widetilde{F}_k(z_1,...,z_n,z_{n+1}) = F_k(z_1,...,z_n) - z^2_{n+1}$, $k = 1,...,N$, and, as in the proof of Lemma 2.15, there exists a double covering $\Phi : \mathbb{C}Ch(\widetilde{\mathcal{S}}, \widetilde{\mathcal{A}}) \to \mathbb{C}T^n_d$ ramified along $\mathbb{C}Ch(\mathcal{S}, \mathcal{A})$.

Now note that S^1 acts on $\partial\mathbb{C}T^{n+1}_{d,2}$ by

$$(v,\ (w_1,...,w_n,w_{n+1}) \mapsto (w_1v,...,w_nv,w_{n+1}v^{d/2}), \quad v \in S^1 \ .$$

The manifold $\partial\mathbb{C}Ch(\widetilde{\mathcal{S}}, \widetilde{\mathcal{A}})$ is invariant with respect to this action, because it is defined by quasihomogeneous polynomials $f(z_1,...,z_{n+1})$ satisfying

$$f(z_1\tau,...,z_n\tau,z_{n+1}\tau^{d/2}) = \tau^d f(z_1,...,z_{n+1}),$$

which is compatible with the S^1 action. By the same reason

$$\Phi(w_1v,...,w_nv,w_{n+1}v^{d/2}) = (w'_1v,...,w'_nv), \quad v \in S^1 \ ,$$

as far as $\Phi(w_1,...,w_n,w_{n+1}) = (w'_1,...,w'_m)$, $(w_1,...,w_{n+1}) \in \partial\mathbb{C}Ch(\widetilde{\mathcal{S}}, \widetilde{\mathcal{A}})$. Hence Φ reduces to a double covering $\Pi : Y \to \mathbb{C}P^n$, where $Y = \mathbb{C}Ch(\widetilde{\mathcal{S}}, \widetilde{\mathcal{A}})/S^1$, ramified along M. It remains to show that Y is a closed PL-manifold. Indeed, for any edge of $T^{n+1}_{d,2}$ there exists a combination of an automorphism of $\mathbb{Z}^{n+1}$, leaving the

$(n+1)$-st axis fixed, and shifts which puts this edge on a coordinate axis and the adjacent faces of $T^n_{d,s}$ on the corresponding coordinate planes. One can easily verify that the orbits of the S^1 action on that edge and adjacent faces contract into points, whose neighborhoods in Y are homeomorphic to $\mathbb{R}^{2n}$ as shown in the proof of Proposition 1.17.

At last, we show that $\pi_1(Y)=0$. Any loop in $\mathbb{C}P^n$ through a point $\overline{w}\in M$ lifts to a loop on Y through $\overline{w}$, because $\overline{w}$ is covered in Y by itself only. Since any loop in $\mathbb{C}P^n$ is contractible, so does its lifting in Y. $\Box$

Proposition 2.20 *Let n and d be positive even numbers, M be a C-hypersurface of degree d in $\mathbb{C}P^n$, and Y be a double covering of $\mathbb{C}P^n$ ramified along M. Then*

$$\chi(Y)=\chi^n_{d,2},\quad \mathrm{sign}(Y)=\mathrm{sign}^n_{d,2},\quad H_*(Y)\simeq H_*(Y'),$$

and the lattices $H_n(Y)$ and $H_n(Y')$, equipped with the intersection forms, are isomorphic, where Y' is the double covering of $\mathbb{C}P^n$ ramified along a nonsingular algebraic hypersurface of degree d, and $\chi^n_{d,2}$ and $\mathrm{sign}^n_{d,2}$ are, respectively, the Euler characteristic and the signature of Y'.

Proof. The equalities $\chi(Y)=\chi(Y')$ and $\mathrm{sign}(Y)=\mathrm{sign}(Y')$ follow immediately from the additivity of the Euler characteristic, the Hirzebruch formula for the signature of ramified coverings and Corollary 2.13. The isomorphism $H_*(Y)\simeq H_*(Y')$, by [10], reduces to $H_{2i-1}(Y)=0$, $i=1,...,n$, $H_{2i}(Y)=\mathbb{Z}$, $i=0,...,n$, $i\neq n/2$, $H_n(Y)\simeq\mathbb{Z}^r$, $r=\mathrm{rk}H_n(Y')$. It can be proven as Proposition 2.16, using a ramified covering of Y similar to that constructed in Proposition 2.10. The same ramified covering applied as in the proof of Proposition 2.17, gives a lattice isomorphism $H_n(Y)\simeq H_n(Y')$. $\Box$

Proposition 2.21 *Let n and d be positive even numbers and M be a real C-hypersurface of degree d in $\mathbb{C}P^n$. Then $\mathbb{R}P^n=\mathbb{R}M_+\cup\mathbb{R}M_-$, where $\mathbb{R}M_+$ and $\mathbb{R}M_-$ are compact manifolds with boundary such that*

$$\partial\mathbb{R}M_+=\partial\mathbb{R}M_-=\mathbb{R}M_+\cap\mathbb{R}M_-=\mathbb{R}M\ .$$

The ramified double covering $\Pi:Y\to\mathbb{C}P^n$ admits an action of the group $\mathbb{Z}/2\mathbb{Z}\oplus\mathbb{Z}/2\mathbb{Z}=\{\mathrm{Id},\tau,\mathrm{Conj}_+,\mathrm{Conj}_-\}$, where τ is the deck transformation of Π and $\Pi\circ\mathrm{Conj}_+=\Pi\circ\mathrm{Conj}_-=\mathrm{Conj}\circ\Pi$. In addition,

$$\mathrm{Fix}(\tau)=M,\quad \mathbb{R}Y_\pm\overset{\mathrm{def}}{=}\mathrm{Fix}(\mathrm{Conj}_\pm)=\Pi^{-1}(\mathbb{R}M_\pm)\ .$$

Proof. In the framework of the construction in the proof of Proposition 2.19, we define $\mathbb{R}Ch(\mathcal{S},\mathcal{A})_+$ (resp., $\mathbb{R}Ch(\mathcal{S},\mathcal{A})_-$) as the union of the closures of $\mathbb{C}\mu_{\Delta_k}(\{F_k\geq 0\}\cap(\mathbb{R}^*)^n)$ (resp., $\mathbb{C}\mu_{\Delta_k}(\{F_k\leq 0\}\cap(\mathbb{R}^*)^n)$), $k=1,...,N$. The action of S^1 on $\partial\mathbb{C}T^n_d$ reduces to the action of $S^0=\{\pm 1\}$ on $\partial\mathbb{R}T^n_d$ which preserves

$\mathbb{R}Ch(\mathcal{S},\mathcal{A})_+$ and $\mathbb{R}Ch(\mathcal{S},\mathcal{A})_-$, thus defining $\mathbb{R}M_\pm = \mathbb{R}Ch(\mathcal{S},\mathcal{A})_\pm/S^0$. The involutions $\tau, \mathrm{Conj}_+, \mathrm{Conj}_-$ on Y one gets from the involutions on $\mathbb{C}Ch(\widetilde{\mathcal{S}},\widetilde{\mathcal{A}})$:

$$(w_1,...,w_n,w_{n+1}) \mapsto (w_1,...,w_n,-w_{n+1}) \ ,$$
$$(w_1,...,w_n,w_{n+1}) \mapsto \mathrm{Conj}(w_1,...,w_n,w_{n+1}) \ ,$$
$$(w_1,...,w_n,w_{n+1}) \mapsto \mathrm{Conj}(w_1,...,w_n,-w_{n+1}) \quad \Box$$

Now we have to make a digression on the topology of smooth (not necessarily algebraic) hypersurfaces in $\mathbb{R}P^n$. Following Viro [37], we define a *rank* of a connected smooth hypersurface in $\mathbb{R}P^n$ to be the maximal integer r such that the homomorphism induced in r-dimensional homology with $\mathbb{Z}/2\mathbb{Z}$ coefficients by the inclusion of the hypersurface into the projective space is nontrivial. It is easy to see that the intersection of a hypersurface of rank r in $\mathbb{R}P^n$ with a transversal projective subspace P of dimension $k \geq n-r$ is a hypersurface of rank $r-n+k$ in P. Similarly, we define a *rank* of a connected component of the complement of a hypersurface in $\mathbb{R}P^n$. Clearly, the rank of a connected component C of the complement of a two-sided hypersurface (*i.e.*, a hypersurface dividing its tubular neighborhood) is greater or equal to the rank of each component of the boundary ∂C.

A component C of the complement of a two-sided hypersurface is called *principal* if the rank of C is greater then the rank of each component of ∂C.

Proposition 2.22 *Let S be a two-sided hypersurface in $\mathbb{R}P^n$. Then there exists at most one principal component of $\mathbb{R}P^n \setminus S$.*

Proof. Suppose that $\mathbb{R}P^n \setminus S$ has two principal components C_1 and C_2. Let r_1 (resp., r_2) be the maximal rank of a connected component of ∂C_1 (resp., ∂C_2). Assume that $r_1 \leq r_2$. Consider a projective subspace P of $\mathbb{R}P^n$ transversal to S and of dimension $n-r_1$. All the connected components of $P \cap \partial C_1$ are of rank 0 in P. Note that the ranks of $P \cap C_1$ and $P \cap C_2$ in P are positive. Each connected component of $P \cap \partial C_1$ divides P into two parts. One of them has rank 0 in P, and is called the *interior* of the component. The other one contains $P \cap C_1$ (and has a positive rank). Therefore, $P \cap C_2$ is contained in the interior of some component of $P \cap \partial C_1$, and thus should have rank 0. $\Box$

Clearly, the real parts of a real algebraic hypersurface of even degree and of a real C-hypersurface of even degree are two-sided in $\mathbb{R}P^n$. Switching if necessary $\mathbb{R}M_+$ and $\mathbb{R}M_-$, we suppose from now on that $\mathbb{R}M_-$ contains the principal component of $\mathbb{R}P^n \setminus \mathbb{R}M$, if this principal component does exist. For any manifold X denote by $b_*(X)$ the total Betti number $\dim_{\mathbb{Z}/2\mathbb{Z}} H_*(X;\mathbb{Z}/2\mathbb{Z})$ of X.

Theorem 2.23 *Let n be a positive even number, and M a real C-hypersurface of degree d in $\mathbb{C}P^n$. Then $b_*(\mathbb{R}Y_+) = b_*(\mathbb{R}M)$ and $b_*(\mathbb{R}Y_-) = b_*(\mathbb{R}M)+2$.*

Proof. From the Smith exact sequence (see, for example, [5, 38]) applied to the deck transformation of $\mathbb{R}Y_\pm$, it easily follows that $b_*(\mathbb{R}Y_\pm) = b_*(\mathbb{R}M) + 2a$, where a is the number of principal components in $\mathbb{R}M_\pm$. It remains to use Proposition 2.22

and note that in our case $\mathbb{R}P^n \setminus \mathbb{R}M$ should have a principal component. Indeed, if $\mathbb{R}P^n \setminus \mathbb{R}M$ does not have a principal component, one has $b_*(\mathbb{R}Y_+) = b_*(\mathbb{R}M)$ and $b_*(\mathbb{R}Y_-) = b_*(\mathbb{R}M)$, and thus, $\chi(\mathbb{R}Y_+) \equiv \chi(\mathbb{R}Y_-) \mod 4$. The last congruence is impossible, because $\chi(\mathbb{R}Y_\pm) = 2\chi(\mathbb{R}M_\pm)$ and $\chi(\mathbb{R}M_+) + \chi(\mathbb{R}M_-) = \chi(\mathbb{R}P^n) = 1$. □

Remark 2.24 *In particular, the statement of Theorem 2.23 holds for real algebraic hypersurfaces of even degree in even dimensional projective space.*

3 Topology of real C-hypersurfaces

3.1 Generalized Harnack inequalities

Let M' be a nonsingular algebraic hypersurface of degree d in $\mathbb{C}P^n$. Denote by b_d^n the total Betti number $b_*(M') = \dim_{\mathbb{Z}/2\mathbb{Z}} H_*(M'; \mathbb{Z}/2\mathbb{Z})$ of M'. One has (see, for example, [8])

$$b_d^n = \frac{(d-1)^{n+1} - (-1)^{n+1}}{d} + d + (-1)^{n+1}.$$

Theorem 3.1 *For a real C-hypersurface M of degree d in $\mathbb{C}P^n$ one has*

$$b_*(\mathbb{R}M) = b_d^n - 2a(\mathbb{R}M), \tag{20}$$

where $a(\mathbb{R}M)$ is a nonnegative integer. Furthermore, if n and d are both even, then

$$b_*(\mathbb{R}Y_-) = b_{d,2}^n - 2a(\mathbb{R}M), \tag{21}$$

$$b_*(\mathbb{R}Y_+) = b_{d,2}^n - 2a(\mathbb{R}M) - 2, \tag{22}$$

where Y is the double covering of $\mathbb{C}P^n$ ramified along M, $\mathbb{R}Y_\pm$ are the fixed point sets of two liftings $\mathrm{Conj}_\pm$ to Y of the complex conjugation Conj in $\mathbb{C}P^n$, and $b_{d,2}^n$ is the total Betti number of the double covering of $\mathbb{C}P^n$ ramified along a nonsingular hypersurface of degree d.

Proof. The statement follows from the Smith-Floyd inequality

$$\dim_{\mathbb{Z}/2\mathbb{Z}} H_*(\mathrm{Fix}(\tau); \mathbb{Z}/2\mathbb{Z}) \le \dim_{\mathbb{Z}/2\mathbb{Z}} H_*(X; \mathbb{Z}/2\mathbb{Z}) \tag{23}$$

(where X is a compact CW-complex and $\tau : X \to X$ is an involution) and the congruence

$$\dim_{\mathbb{Z}/2\mathbb{Z}} H_*(X; \mathbb{Z}/2\mathbb{Z}) \equiv \dim_{\mathbb{Z}/2\mathbb{Z}} H_*(\mathrm{Fix}(\tau); \mathbb{Z}/2\mathbb{Z}) \mod 2; \tag{24}$$

see for details [24, 38]. We apply the Smith inequality and the above congruence to the involutions Conj on M and $\mathrm{Conj}_\pm$ on Y. In order to get (20) we use Proposition 2.16. To prove (21) and (22) we notice that according to Theorem 2.23 one has $b_*(\mathbb{R}Y_+) = b_*(\mathbb{R}M)$ and $b_*(\mathbb{R}Y_-) = b_*(\mathbb{R}M) + 2$. In addition,

$$b_*(Y) = \chi(Y) = 2\chi(\mathbb{C}P^n) - \chi(M) = 2(n+1) - (2n - b_*(M)) = 2 + b_*(M) \ ,$$

and it remains to apply Proposition 2.20. □

3.2 Congruences

Theorem 3.2 *Let M be a real C-hypersurface of degree d in $\mathbb{C}P^n$. If n is odd then*

$$\chi(\mathbb{R}M) \equiv \mathrm{sign}_d^n + \begin{cases} 0, & \text{if } a(\mathbb{R}M) = 0, \\ \pm 2, & \text{if } a(\mathbb{R}M) = 1 \end{cases} \mod 16 . \tag{25}$$

If n and d are even, then

$$\chi(\mathbb{R}M_-) \equiv \frac{\mathrm{sign}_{d,2}^n}{2} + \begin{cases} 0, & \text{if } a(\mathbb{R}M) = 0, \\ \pm 1, & \text{if } a(\mathbb{R}M) = 1 \end{cases} \mod 8 , \tag{26}$$

$$\chi(\mathbb{R}M_+) \equiv \frac{\mathrm{sign}_{d,2}^n}{2} \pm 1 \mod 8, \text{ if } a(\mathbb{R}M) = 0. \tag{27}$$

Proof. The statement can be proven as the Rokhlin and Gudkov-Krahnov-Kharlamov congruences in the algebraic case [14, 21, 24, 38].

(i) Let $n = 2k+1$ and $a(\mathbb{R}M) = 0$. As in [24] (see also [14, 38]), the latter implies the splitting of $H_{n-1}(M)$ into the orthogonal sum $H_+ \oplus H_-$ of unimodular eigenlattices corresponding to the eigenvalues ± 1 of $\mathrm{Conj}_* : H_{n-1}(M) \to H_{n-1}(M)$.

Lemma 3.3 *The signature of involution* $\mathrm{sign}(\mathrm{Conj}_*) = \mathrm{sign}(H_+) - \mathrm{sign}(H_-)$ *is equal to* $(-1)^k \chi(\mathbb{R}M)$.

Proof. By the Atiyah-Singer formula (see [3, 24, 38]) $\mathrm{sign}(\mathrm{Conj}_*)$ is equal to $\mathbb{R}M \circ \mathbb{R}M$, the selfintersection of $\mathbb{R}M$ in M. Let us show that $\mathbb{R}M \circ \mathbb{R}M = (-1)^k \chi(\mathbb{R}M)$. First, we smooth M as in Proposition 2.7. Then we take a tangent vector field V on $\mathbb{R}M_{sm}$ having only finitely many singular points which are all nondegenerate and lie outside $\mathbb{R}\delta$ for any proper face δ of the polyhedra $\Delta_1, ..., \Delta_N$ in the subdivision of T_d^n. Extend the vector field $J(V)$ (where J is the almost complex structure on M_{sm} defined in Proposition 2.8) to a neighborhood of $\mathbb{R}M_{sm}$ in M_{sm} and slightly move $\mathbb{R}M_{sm}$ along geodesics in M_{sm} tangent to the field obtained. The result has transversal intersection points with $\mathbb{R}M_{sm}$ at the singular points of V. The intersection indices are equal to the multiplied by $(-1)^k$ indices of the singular points of V (see [24, 38]). $\square$

Denote by H_e the lattice H_+ (resp., H_-) if k is odd (resp., k is even). From the existence of an almost complex structure on a smoothing of M (Proposition 2.8), it follows that H_e is even (see, for example, [38]). Let σ_e be the signature of H_e. Since H_e is unimodular and even, the signature σ_e is divisible by 8. To finish the proof in the case $a(\mathbb{R}M) = 0$, it remains to note that according to Lemma 3.3 we have $\mathrm{sign}(M) - \chi(\mathbb{R}M) = 2\sigma_e$.

The prove (27) and (26) in the case $a(\mathbb{R}M) = 0$, we apply the same arguments to $(Y, \mathrm{Conj}\_)$ and (Y, Conj_+) and use the relations $\chi(\mathbb{R}Y_\pm) = 2\chi(\mathbb{R}M_\pm)$, $b_*(\mathbb{R}Y_+) = b_*(\mathbb{R}M)$ and $b_*(\mathbb{R}Y_-) = b_*(\mathbb{R}M) + 2$.

(ii) Let now $n = 2k+1$ and $a(\mathbb{R}M) = 1$. In this case (see, for example, [21, 38]), the discriminants of H_+ and H_- are ± 2 (from the Smith theory we get the inequality $\mid \operatorname{discr}(H_\pm) \mid \leq 2$ and then use the congruence $\chi(M) \equiv (-1)^k \operatorname{sign}(M) \mod 4$ to show that $\mid \operatorname{discr}(H_\pm) \mid \neq 1$). Using Lemma 3.3 and the fact that the signature of an even lattice with discriminant ± 2 is congruent to $\pm 1 \mod 8$ we immediately obtain the statement required.

To prove (26) in the case $a(\mathbb{R}Y_-) = 1$, we again apply the previous arguments to (Y, Conj_-). $\square$

3.3 Comessatti inequality for real C-surfaces

Theorem 3.4 *Let M be a real C-hypersurface of degree d in $\mathbb{C}P^n$ and M' be a nonsingular algebraic hypersurface of degree d in $\mathbb{C}P^n$. Then*

$$2 - h^{1,1}(M') \leq \chi(\mathbb{R}M) \leq h^{1,1}(M').$$

Proof. The arguments are completely similar to the proof of the Comessatti inequality in the case of real algebraic surfaces.

Let again H_+ and H_- be eigensublattices of $H_2(M)$ corresponding to the eigenvalues ± 1 of $\mathrm{Conj}_* : H_2(M) \to H_2(M)$. Denote by $a_\pm^+$ (resp., $a_\pm^-$) the number of positive (resp., negative) squares in the diagonal form over $\mathbb{Q}$ of the restriction of $B : H_2(M) \times H_2(M) \to \mathbb{Z}$ to $H_\pm$. We have

$$\begin{aligned} a_+^+ + a_+^- + a_-^+ + a_-^- &= \dim H_2(M), \\ a_+^+ - a_+^- + a_-^+ - a_-^- &= \operatorname{sign}(M), \\ a_+^+ + a_+^- - a_-^+ - a_-^- &= \chi(\mathbb{R}M) - 2, \\ a_+^+ - a_+^- - a_-^+ + a_-^- &= -\chi(\mathbb{R}M). \end{aligned}$$

The first two equations are clear. The third and fourth ones follow from the Lefschetz fixed point theorem and Atiyah-Singer theorem, respectively. We obtain that

$$a_+^- = (\dim H_2(M) - \operatorname{sign}(M) + 2\chi(\mathbb{R}M) - 2)/4 \geq 0,$$

$$a_-^- = (\dim H_2(M) - \operatorname{sign}(M) - 2\chi(\mathbb{R}M) + 2)/4 \geq 0.$$

These inequalities together with the Hodge index relations give the required statement. $\square$

3.4 Topology of real C-curves

Let M be an oriented smooth connected closed surface in $\mathbb{C}P^2$. Then M is called a *flexible curve of degree d* (see [35]) if

- it realizes $d[\mathbb{C}P^1] \in H_2(\mathbb{C}P^2)$,

- the genus of M is equal to $(d-1)(d-2)/2$,
- M is invariant under the complex conjugation,
- the field of tangent planes to M on $M \cap \mathbb{R}P^2$ can be equivariantly deformed to the field of lines in $\mathbb{C}P^2$ tangent to $M \cap \mathbb{R}P^2$.

According to Propositions 2.6, 2.8 and Corollary 2.13 (a smoothing of) a real C-curve of degree d in $\mathbb{C}P^2$ is a flexible curve of degree d. Thus, all the restrictions on the topology of flexible curves are applicable to real C-curves. We formulate here in the framework of real C-curves the principal known restrictions on flexible curves. A big list of restrictions on the topology of flexible curves can be found in [35].

Let us start from definitions. The standard definitions applicable to real algebraic curves can be naturally extended to real C-curves. A real C-curve A of degree d in $\mathbb{C}P^2$ is called an *M-curve* or *maximal* if $\mathbb{R}A$ has $(d-1)(d-2)/2+1$ connected components. A real C-curve A of degree d in $\mathbb{C}P^2$ is called an *$(M-i)$-curve* if $\mathbb{R}A$ has $(d-1)(d-2)/2+1-i$ connected components. A connected component of the real part of a real C-curve of degree d in $\mathbb{C}P^2$ is called an *oval* if it divides $\mathbb{R}P^2$ into two parts. The part homeomorphic to a disk is called the *interior* of the oval. All the connected components of the real part of a real C-curve of an even degree in $\mathbb{C}P^2$ are ovals. Exactly one connected component of the real part of a real C-curve of an odd degree in $\mathbb{C}P^2$ is not an oval. This component is called *nontrivial*. An oval is *even* (resp., *odd*) if it lies inside of an even (resp., odd) number of other ovals of the curve. The numbers of even and odd ovals of a curve are denoted by p and n, respectively. The Euler characteristic of a connected component of the complement in $\mathbb{R}P^2$ of the real part of a real C-curve is called the *characteristic* of an oval bounding the component from outside.

Theorem 3.5
- Harnack inequality. *The number of connected components of the real part of a real C-curve of degree d in $\mathbb{C}P^2$ is at most $(d-1)(d-2)/2+1$.*
- Gudkov-Rokhlin congruence. *For a maximal real C-curve of degree $d=2k$ in $\mathbb{C}P^2$ one has*
$$p-n \equiv k^2 \mod 8.$$
- Gudkov-Krahnov-Kharlamov congruence. *Let A be a real C-curve of degree $d=2k$ in $\mathbb{C}P^2$. If A is an $(M-1)$-curve, then*
$$p-n \equiv k^2 \pm 1 \mod 8.$$
- Strengthened Petrovsky inequalities. *For a real C-curve A of degree $d=2k$ in $\mathbb{C}P^2$ one has*
$$p-n^- \leq 3k(k-1)/2+1, \quad n-p^- \leq 3k(k-1),$$
where p^- (resp., n^-) is the number of even (resp., odd) ovals of $\mathbb{R}A$ with negative characteristic.

- Strengthened Arnold inequalities. *For a real C-curve A of degree $d = 2k$ in $\mathbb{C}P^2$ one has*

$$p^- + p^0 \leq (k^2 - 3k + 3 + (-1)^k)/2, \quad n^- + n^0 \leq (k^2 - 3k + 2)/2,$$

where p^0 (resp., n^0) is the number of even (resp., odd) ovals of $\mathbb{R}A$ with characteristic 0.

A real C-curve A in $\mathbb{C}P^2$ is said to be of *type I* if its real part $\mathbb{R}A$ divides A into two parts; otherwise, the curve is of *type II.* For a curve A of type I, the orientations of two halves of $A \setminus \mathbb{R}A$ induce on $\mathbb{R}A$ two opposite orientations which are called *complex orientations*. Note that a real C-curve A is of type I if and only if all the algebraic curves used in the gluing of A are of type I and complex orientations on the real parts of these curves can be chosen in such a way that they glue into an orientation of $\mathbb{R}A$.

A pair of ovals of the real part of a real C-curve in $\mathbb{C}P^2$ is *injective* if one of them is inside of the other one. A collection of ovals is called a *nest* if any two of them form an injective pair. An injective pair of ovals of a real C-curve is *positive* (resp., *negative*) if the complex orientations of the ovals are induced (resp., are not induced) from some orientation of the annulus bounded by the ovals. Take an oval of a real C-curve of type I and of an odd degree in $\mathbb{C}P^2$, and consider the Möbius band which is the complement in $\mathbb{R}P^2$ of the interior of the oval. The oval is called *positive* (resp., *negative*) if the integer homology class realized in the Möbius band by the oval equipped with a complex orientation differs in sign (resp., coincides) with the class of the doubled nontrivial component equipped with the complex orientation.

Theorem 3.6

- Klein congruence. *Let A be a real C-curve of type I in $\mathbb{C}P^2$. If A is an $(M - i)$-curve, then $i \equiv 0 \mod 2$.*

- Arnold congruence. *A real C-curve of even degree $d = 2k$ satisfies*

$$p - n \equiv k^2 \mod 4 .$$

- Rokhlin-Mishachev formulae. *For a real C-curve A of type I and of degree $d = 2k$ in $\mathbb{C}P^2$ one has*

$$2(\Pi^+ - \Pi^-) = l - k^2,$$

where l is the number of ovals of $\mathbb{R}A$, and Π^+ and Π^- are the numbers of positive and negative injective pairs, respectively. For a real C-curve A of type I and of degree $d = 2k + 1$ in $\mathbb{R}P^2$ one has

$$2(\Pi^+ - \Pi^-) + \Lambda^+ - \Lambda^- = l - k(k + 1),$$

where Λ^+ and Λ^- are the numbers of positive and negative ovals, respectively.

- Kharlamov- Marin congruence. *Let A be a real C-curve of degree $d = 2k$ in $\mathbb{C}P^2$. If A is an $(M - 2)$-curve and $p - n \equiv k^2 + 4 \mod 8$, then A is of type I.*

Harnack inequality in the case of real C-curves constructed using a primitive (*i.e.*, such that all its triangles are of area 1/2) triangulation was, first, proved in [18] and then in a different way in [15]. Rokhlin-Mishachev formulae in the case of real C-curves constructed using a primitive triangulation was proved in [22].

It is interesting that one restriction on real C-curves is not proved in the case of flexible curves. This restriction was proved in [9]:

consider a real C-curve A of degree d in $\mathbb{C}P^2$; if A is glued from threenomials, then the sum of the depths of any two nests of $\mathbb{R}A$ is at most $d/2$.

In the case of real algebraic curves this statement is known as Hilbert's theorem and is an immediate corollary of the Bezout theorem.

References

[1] Atiyah, M. F.: Convexity and commuting Hamiltonians. *Bull. Lond. Math. Soc.* **14** (1982), 1-15.

[2] Atiyah, M. F.: Angular momentum, convex polyhedra and algebraic geometry. *Proc. Edinburgh Math. Soc.* **26** (1983), 121-138.

[3] Atiyah, M. F., and Singer, I. M.: The index of elliptic operators: III. *Ann. of Math.* **87** (1968), 546-604.

[4] Biran, P.: Constructing new ample divisors out of old ones. *Duke Math. J.* **98** (1999), no. 1, 113-135.

[5] Bredon, G. E.: Introduction to compact transformation groups. *Pure and Applied Mathematics* **46**. Academic Press. New York. 1972.

[6] Chevallier, B.: Secteurs et déformations locales des courbes réelles. *Math. Ann.* **307** (1997), no. 1, 1-28.

[7] Connelly, R. and Henderson, D. W.: A convex 3-complex not simplicially isomorphic to a strictly convex complex. *Proc. Camb. Phil. Soc.* **88** (1980), no. 2, 299-306.

[8] Danilov, V. I. and Khovanskii, A. G.: Newton polyhedra and an algorithm for computing Hodge-Deligne numbers, *Izv. Acad. Nauk SSSR* **50** (1986), (Russian), English transl., Math. USSR Izvestiya **29:2** (1987), 279–298.

[9] de Loera, J. A. and Wicklin, F. J.: On the need of convexity in patchworking. *Adv. in Appl. Math.* **20** (1998), 188-219.

[10] Fary, I.: Cohomologie des variétés algébriques. *Ann. of Math.* **65** (1957), no. 1, 21-73.

[11] Freedman, M. H., and Quinn, F.: *Topology of 4-manifolds.* Princeton Mathematical Series, 39. Princeton Univ. Press, Princeton, NJ, 1990.

[12] Fulton, W.: *Introduction to toric varieties* (Ann. Math. Studies 131), Princeton Univ. Press, Princeton N.J., 1993.

[13] Gelfand, I. M., Kapranov, M. M., Zelevinski, A. V.: *Discriminants, resultants and multidimensional determinants.* Birkhäuser, Boston, 1994.

[14] Gudkov, D. A.: Topology of real projective algebraic varieties. *Rus. Math. Surveys* **29** (1974), no. 4, 3-79.

[15] Haas, B:. *The Ragsdale conjecture for maximal T-curves.* Preprint, 1997, Universität Basel.

[16] Hirzebruch, F.: The signature of ramified coverings. In: *Global Analysis (Papers in honor of K. Kodaira).* Univ. Tokyo Press, Tokyo, 1969, pp. 253-265.

[17] Hirzebruch, F.: The signature theorem: reminiscences and recreation. In: *Prospects in Mathematics (Ann. Math. Stud. 70)*, Princeton Univ. Press, Princeton, 1971, pp. 3-31.

[18] Itenberg, I.: Counterexamples to Ragsdale conjecture and T-curves. In: *Contemporary Math.* **182** *(Proc. Conf. Real Alg. Geom., December 17-21, 1993, Michigan, ed. S.Akbulut)*, AMS, Providence, R.I., 1995, pp. 55-72.

[19] Itenberg, I. and Shustin, E.: Singular points and limit cycles of planar polynomial vector fields. *Duke Math. J.*, to appear.

[20] Itenberg, I., and Viro, O.: Patchworking algebraic curves disproves the Ragsdale conjecture. *Math. Intelligencer* **18** (1996), no. 4, 19-28.

[21] Kharlamov, V. M.: New relations for the Euler characteristic of real algebraic manifolds. *Func. Anal. Appl.* **7** (1973), 147-150.

[22] Parenti, P.: *Combinatorics of dividing T-curves.* Ph. D. Thesis, 1999, Università di Pisa.

[23] Risler, J.-J.: Construction d'hypersurfaces réelles [d'après Viro]. *Séminaire N.Bourbaki*, no. 763, vol. 1992-93, Novembre 1992.

[24] Rokhlin, V. A.: Congruences modulo 16 in Hilbert's 16th problem *Funct. Anal. Appl.* **6** (1972), 301-306.

[25] Rokhlin, V. A.: Complex topological characteristics of real algebraic curves. *Russ. Math. Surveys 33* (1978), no. 5, 85-98.

[26] Santos, F.: *Improved counterexamples to the Ragsdale conjecture.* Preprint, Universidad de Cantabria, 1994.

[27] Shustin, E.: Topology of real plane algebraic curves. In: *Proc. Intern. Conf. Real Algebraic Geometry, Rennes, June 24-29 1991* (Lect. Notes Math. **1524**, Springer, 1992), pp. 97-109.

[28] Shustin, E.: Real plane algebraic curves with prescribed singularities. *Topology* **32** (1993), no. 4, 845-856.

[29] Shustin, E.: Critical points of real polynomials, subdivisions of Newton polyhedra and topology of real algebraic hypersurfaces. *Amer. Math. Soc. Transl. (2)* **173** (1996), 203-223.

[30] Shustin, E.: Gluing of singular and critical points. *Topology* **37** (1998), no. 1, 195-217.

[31] Sturmfels, B.: Viro's theorem for complete intersections. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **21** (1994), no. 3, 377-386.

[32] Viro, O. Ya.: Gluing of algebraic hypersurfaces, smoothing of singularities and construction of curves. In: *Proc. Leningrad Int. Topological Conf., Leningrad, Aug. 1983*, Nauka, Leningrad, 1983, p. 149-197 (Russian).

[33] Viro, O. Ya.: Gluing of plane real algebraic curves and construction of curves of degrees 6 and 7. *Lect. Notes Math.* **1060**, Springer, Berlin etc., 1984, p. 187-200.

[34] Viro, O. Ya.: Real plane curves of degrees 7 and 8: new prohibitions. *Math. USSR Izvestia* **23** (1984), 409-422.

[35] Viro, O. Ya.: Progress in the topology of real algebraic varieties over the last six years. *Rus. Math. Surv.* **41** (1986), no. 3, 55-82.

[36] Viro, O. Ya.: Real algebraic plane curves: constructions with controlled topology. *Leningrad Math. J.* **1** (1990), 1059-1134.

[37] Viro, O. Ya.: Mutual position of hypersurfaces in projective space (to appear in *Arnold's Festschrift*).

[38] Wilson, G.: Hilbert's sixteenth problem. *Topology* **17** (1978), no. 1, 53-73.