

HOMOGENEOUS FOLIATIONS: TOPOLOGY OF LEAVES AND APPLICATIONS TO POLYGONAL BILLIARDS

FERRÁN VALDEZ

ABSTRACT. The aim of this paper is to describe the topology of the leaves of a homogeneous foliation on \mathbf{C}^2 with isolated singularity and reduced tangent cone. This description leads to a complete topological classification of translation surfaces associated to a polygonal billiard.

Keywords: Homogeneous foliation, polygonal billiard.

Mathematical subject classification: 37F75, 37D50.

1. INTRODUCTION

Let $\overline{P} \subset \mathbf{R}^2$ be a polygon and $Q(\overline{P})$ the set of its vertices. Through this article we consider polygons without vertices $P := \overline{P} \setminus Q(\overline{P})$. Let $S(P)$ denote the translation real surface arising by unfolding billiard trajectories inside P . In general, there exist a "non dicritical" holomorphic homogeneous foliation $\mathcal{F} = \mathcal{F}(P)$ on \mathbf{C}^2 whose generic leaves are isomorphic, as translation surfaces, to the surface $S(P)$ [7]. Before precisising what genericity and generality mean, (see §2 and §4 respectively) we recall what is known about the topology of the orientable surface $S(P)$:

Lemma 1. [4] *Let the angles of P be $\pi m_i/n_i$, $i = 1, \dots, k$, where m_i and n_i are coprime, and N be the last common multiple of n_i 's. Then*

$$\text{genus } S(P) = 1 + \frac{N}{2} \left(k - 2 - \sum \frac{1}{n_i} \right)$$

In this article we give complete *topological* description of the orientable surfaces $S(P)$ when the angles of P are not necessarily rational. We achieve this by proving the following

Theorem 1. *Let \mathcal{F} be an homogeneous holomorphic foliation on \mathbf{C}^2 with isolated singularity and reduced tangent cone. Then, every non singular leaf which is not a separatrix is homeomorphic to either:*

- (1) $\mathbf{C} \setminus \mathbf{Z}[i]$, endowed with the standart topology.
- (2) $S \setminus E$, where S is a compact RIEMANN surface and $E \subset S$ is a finite set of points.

- (3) $M \setminus E'$, where M is the LOCH NESS monster, i.e. the plane to which we stick a countable set of handles, and $E' \subseteq M$ is a countable and discrete set of points.
- (4) M , the LOCH NESS monster.

The foregoing theorem completes a result presented in [7]. From the same reference, we deduce that any translation surface $S(P)$ is homeomorphic to either (2), (3) or (4). It is relevant to remark that, up to punctures, all the topological surfaces in our list are contained in the milestone

Theorem 2. [3] *Let \mathcal{G} be an orientable dimension 2 lamination on an arbitrary compact space. Then, if \mathcal{G} has no compact leaf, a non countable set of its leaves are diffeomorphic to either:*

- (1) *The plane \mathbf{R}^2 ,*
- (2) *The LOCH NESS monster,*
- (3) *The cylinder $\mathbf{R} \times \mathbf{S}^1$,*
- (4) *JACOB'S STAIR, i.e. the cylinder to which we stick a countable set of handles in both directions,*
- (5) *CANTOR'S TREE, i.e. the sphere \mathbf{S}^2 to which we remove a CANTOR set,*
- (6) *CANTOR'S FLORID TREE, i.e., a CANTOR'S TREE to which we stick a countable number of handles in all directions.*

2. GENERALITIES

In this section we precise our language and notation. A holomorphic foliation on \mathbf{C}^2 is said to be *homogeneous* if it is invariant under the natural action of the homothetic transformation group $\{T_k(z_1, z_2) := k(z_1, z_2) \mid k \in \mathbf{C}^*\}$. In this article we consider homogeneous holomorphic foliations on \mathbf{C}^2 with isolated singularity and having $\nu + 1$, $\nu \geq 2$, complex lines through the singularity as leaves. We suppose that this set of invariant lines, usually called the *tangent cone* of the foliation, is given by the zeros of the irreducible polynomial $P_{\nu+1} = \prod_{j=1}^{\nu+1} (z_2 - a_j z_1)$, $a_j \in \mathbf{C}$, $a_i \neq a_j, \forall i, j$, and set $a = (a_1, \dots, a_{\nu+1})$. From [2], this foliation is given by the holomorphic 1-form

$$(1) \quad \frac{\omega_{a,\lambda}}{P_{\nu+1}} = \sum_{j=1}^{\nu+1} \lambda_j \frac{d(z_2 - a_j z_1)}{z_2 - a_j z_1},$$

where $\lambda = (\lambda_1, \dots, \lambda_{\nu+1})$ is a point of the affine hyperplane $\{(\lambda_1, \dots, \lambda_{\nu+1}) \in \mathbf{C}^n \mid \sum_{j=1}^{\nu+1} \lambda_j = K, \lambda_j \neq 0 \forall j\}$ and $K \in \mathbf{Z} \setminus 0$ is an arbitrary normalization constant whose choice does not affect

the foliation defined by (1). We denote this foliation $\mathcal{F}_{a,\lambda}$. We remark that

$$(2) \quad F_{a,\lambda}(z_1, z_2) = \prod_{j=1}^{\nu+1} (z_2 - a_j z_1)^{\lambda_j}$$

is a first integral for $\mathcal{F}_{a,\lambda}$. This expression is usually multivaluated. Let \sim_λ be the equivalence relation induced by the \mathbf{Z}^3 action on \mathbf{C}^* defined by $((n_1, n_2, n_3), z) \longrightarrow e^{2\pi i \sum_j n_j \lambda_j} z$. The connected components of the fibers $F_{a,\lambda} : \mathbf{C}^2 \longrightarrow \mathbf{C}^* / \sim_\lambda$ are the leaves of $\mathcal{F}_{a,\lambda}$ in the complement of the tangent cone. Abusing notation we will write $F_{a,\lambda}(z_1, z_2) = k, k \in \mathbf{C}^* / \sim_\lambda$, for such a fiber. A straightforward computation shows that if $k, k' \in \mathbf{C}^*$, the homothety $T_{k/k'}$ defines a diffeomorphism between $F_{a,\lambda}(z_1, z_2) = k$ and $F_{a,\lambda}(z_1, z_2) = k'$. In particular, each leaf in $\mathcal{F}_{a,\lambda}$ is invariant by any homothety $T_{e^{2\pi i \sum_j n_j \lambda_j}}$, $n_j \in \mathbf{Z}, \forall j$.

We will call *generic leaf* any leave in the complement of the tangent cone of $\mathcal{F}_{a,\lambda}$. We denote it by $\mathcal{L} \in \mathcal{F}_{a,\lambda}$. The foliation $\mathcal{F}_{a,\lambda}$ presents, up to diffeomorphism, three kinds of leaves: a (*singular*) *point*, \mathbf{C}^* (*separatrices*) and a *generic leaf* \mathcal{L} .

3. PROOF, THEOREM 1

The topological type of the generic leaf of $\mathcal{F}_{a,\lambda}$ depends only in the parameters $\lambda_1, \dots, \lambda_{\nu+1}$. Indeed, let's consider $\widetilde{\mathcal{F}}_{a,\lambda}$, the extension of the foliation $\mathcal{F}_{a,\lambda}$ to HIRZEBRUCH'S first surface. This complex ruled surface, which we denote $\widetilde{\mathbf{CP}}(2)$, is obtained by blowing up the origin of the affine chart $\mathbf{C} = \{(z_1, z_2)\}$ of $\mathbf{CP}(2)$. A leaf in $\widetilde{\mathcal{F}}_{a,\lambda}$ is called generic if it comes from a generic leaf in $\mathcal{F}_{a,\lambda}$. Let $\pi : \widetilde{\mathbf{CP}}(2) \longrightarrow \mathbf{P}^1(\mathbf{C})$ be the natural projection onto the exceptional divisor. For each $j = 1, \dots, \nu + 1$, we denote as well a_j the intersection of the complex line $z_2 = a_j z_1$ with the exceptional divisor. Let $B_a := \mathbf{P}^1(\mathbf{C}) \setminus \{a_1, \dots, a_{\nu+1}\}$. We remark that $\widetilde{\mathcal{F}}_{a,\lambda}$ is transversal to the fibration defined by π on B_a . Furthermore, for every generic leaf $\mathcal{L} \in \widetilde{\mathcal{F}}_{a,\lambda}$, the restriction $\pi|_{\mathcal{L}} : \mathcal{L} \longrightarrow B_a$ is a covering. The monodromy of this covering is given by the image in $Diff(\mathbf{C}, 0)$ of the holonomy representation of $\widetilde{\mathcal{F}}_{a,\lambda}$ with respect to the fibration defined by π on B_a . This holonomy representation can be explicitly calculated: let $\{\gamma_j\}_{j=1}^{\nu+1}$ be generators for $\pi_1(B_a, *)$, having index δ_i^j with respect to

each $a_j, j = 1, \dots, \nu+1$, and such that $\gamma_i \cap \gamma_j = *$ whenever $i \neq j$. The holonomy representation morphism $hol : \pi_1(B_a, *) \longrightarrow Diff(\mathbf{C}, 0)$ of $\widetilde{\mathcal{F}_{a,\lambda}}$ is given by:

$$(3) \quad hol(\gamma_j) = z \rightsquigarrow e^{2\pi i \lambda_j / K} z.$$

This is clear from (2). In particular, the monodromy action of the covering π_1 is abelian.

Definition 1. We will say that the set $\lambda = (\lambda_1, \dots, \lambda_{\nu+1})$ is:

- Not rational, when $\lambda_j \in \mathbf{C}^* \setminus \mathbf{Q}$ for all $j = 1, \dots, \nu+1$.
- Semi-rational, when there exists a pair $i \neq j$ such that $\lambda_i \in \mathbf{C}^* \setminus \mathbf{Q}$ and $\lambda_j \in \mathbf{Q}$.
- Rational, when $\lambda_j \in \mathbf{Q}$ for all $j = 1, \dots, \nu+1$.

Definition 2. [6] A real surface S is of infinite genus if there is no bounded subset S' such that $S \setminus S'$ is of genus zero.

Following [Ibid], the ideal boundary of a real surface S (not necessarily compact) is a nested triple of topological spaces $B''(S) \subset B'(S) \subset B(S)$. This triple codes the topological type of the surface, that is:

Theorem 3 (Kerékjártó). [Ibid] Let S and \widehat{S} be two separable real surfaces of the same genus and orientability class. Then S and \widehat{S} are homeomorphic if and only if their ideal boundaries (considered as triples of spaces) are topologically equivalent.

The topological spaces $B'(S)$ and $B''(S)$ are the not planar and not orientable parts of the whole ideal boundary $B(S)$. In the context of this article, $B''(\mathcal{L}) = \emptyset$, for the generic leaf of the foliation $\mathcal{F}_{a,\lambda}$ is oriented by its natural complex structure. Up to homeomorphism, the LOCH NESS monster is the only infinite genus real surface whose ideal boundary is just a point.

Lemma 2. If λ is not rational, then the generic leaf $\mathcal{L} \in \mathcal{F}_{a,\lambda}$ is homeomorphic to a LOCH NESS monster.

Proof. First we claim that, for every compact $K \subset \mathcal{L}$ there exists a compact subset $K \subset K' \subset \mathcal{L}$ such that $\mathcal{L} \setminus K'$ is connected by arcs. From the definition of *boundary component*, this implies that the ideal boundary of \mathcal{L} is just a point. Indeed, let $t_0 \in \mathbf{P}^1(\mathbf{C}) \setminus \{a_1, \dots, a_{\nu+1}, *\}$. Let $C_1, \dots, C_{\nu+1}$ be line segments joining a_j to t_0 for every $j = 1, \dots, \nu+1$ and such that $C_i \cap C_j = t_0$, whenever $i \neq j$. Without loss of generality, we suppose as well that the cardinality of $C_i \cap \gamma_j$ is given by KRONECKER'S delta δ_{ij} . The open set

$$(4) \quad U := \mathbf{P}^1(\mathbf{C}) \setminus \bigcup_{j=1}^{\nu+1} C_j$$

is simply connected. Let $\mathbf{P}^1(\mathbf{C}) \simeq \mathbf{S}^2$ be the standard identification of the complex projective line with RIEMANN'S sphere and $\mathbf{S}^2 \hookrightarrow \mathbf{R}^3$ a fixed smooth embedding. Then, the restriction of the standard metric to the image of this embedding induces a complete metric on the exceptional divisor that we denote δ . For every positive real number $\tau < \inf_{i \neq j, i, j=1, \dots, \nu+1} \delta(a_i, a_j)$, the complement in U of the set $\bigcup_{j=1}^{\nu+1} \{t \in \mathbf{P}^1(\mathbf{C}) \mid \delta(t, a_j) < \tau\}$ is closed and simply connected. We denote this set by U_τ .

For every point ξ in a fiber $\pi_1^{-1}(t)$, $t \neq a_1, \dots, a_{\nu+1}, t_0$, let $U_{\tau, \xi}$ be the lifting of U_τ to the leaf \mathcal{L} satisfying $U_{\tau, \xi} \cap \xi = \xi$. We define $K_{\tau, \xi}$ to be the closure of $U_{\tau, \xi}$ in \mathcal{L} . This set is compact in \mathcal{L} .

Let

$$(5) \quad \tau(K) := \frac{1}{2} \inf_{j=1, \dots, \nu+1} \delta(\pi(K), a_j).$$

Since K is compact, there exists a finite set of points $P = \{p_1, \dots, p_s\} \subset B_a$ such that

$$(6) \quad K \subset K' := \bigcup_{\xi \in K \cap \pi_1^{-1}(P)} K_{\tau(K), \xi}.$$

The set $K \cap \pi_1^{-1}(P)$ is finite and therefore K' is compact. We claim that, for every positive real number $\tau < \inf_{i \neq j, i, j=1, \dots, \nu+1} \delta(a_i, a_j)$ and finite subset $Z \subset \pi_1^{-1}(t)$, $t \neq a_1, \dots, a_{\nu+1}, t_0$, the set $\mathcal{L} \setminus \bigcup_{\xi \in Z} K_{\tau, \xi}$ is arcwise connected. This implies that $\mathcal{L} \setminus K'$ is arcwise connected as well.

Indeed, for every point η in a fiber $\pi_1^{-1}(t)$, $t \neq a_1, \dots, a_{\nu+1}, t_0$, let U_η be the lifting of U (4) satisfying $U_\eta \cap \eta = \eta$. We denote $\overline{U_\eta}$ the closure in \mathcal{L} of U_η . Clearly, $\mathcal{L} = \bigcup_{\eta \in \pi_1^{-1}(t)} \overline{U_\eta}$. Consider two distinct points $q \in \overline{U_\eta}$ and $q' \in \overline{U_{\eta'}}$. If $\eta = \eta'$ and $\bigcup_{\xi \in Z} K_{\tau, \xi} \cap \overline{U_\eta}$ has empty interior, then the points q and q' can be joined by an arc in \mathcal{L} . When $\eta \neq \eta'$, the point q can be joined by an arc to a point \tilde{q} such that $\pi(\tilde{q}) \cap \pi(K_{\tau, \xi}) = \emptyset$. Without loss of generality, we assume that $\pi(\tilde{q})$ is contained in a small neighborhood in exceptional divisor of a point a_i , for some $i = 1, \dots, \nu + 1$. Let be $\gamma(\tilde{q})$ a simple loop passing through $\pi(\tilde{q})$ and contained in this small neighborhood. There exist a lift of $\gamma(\tilde{q})$ to the generic leaf \mathcal{L} joining \tilde{q} to a point $\hat{q} \in \overline{U_{\eta''}}$ such that $\bigcup_{\xi \in Z} K_{\tau, \xi} \cap \overline{U_{\eta''}}$ has empty interior. This is true since $\lambda_i \in \mathbf{C}^* \setminus \mathbf{Q}$, for all i , and therefore the cyclic subgroup $\{hol(\gamma_i)^n\}_{n \in \mathbf{Z}}$ of $Diff(\mathbf{C}, 0)$ is torsion-free. Proceeding analogously, the point q' can be joined by an arc to a point $\hat{q}' \in \overline{U_{\eta''}}$. This leads to the previous case in which $\eta = \eta'$. We conclude that a path between the points $q \neq q'$ always exist.

Lemma 3. *If there exist a pair $i \neq j$, $1 \leq i, j \leq \nu + 1$, such that γ_i, γ_j and $\gamma_i\gamma_j$ are not contained in $\text{hol}^{-1}(\text{Id}_{\mathbf{C}})$ (3), then \mathcal{L} has genus different from zero.*

Proof, lemma 3. Let $m \in \pi_1(B_a, *)$ and \tilde{m} be a lifting of this loop to the generic leaf \mathcal{L} via de projection π_{\perp}^{-1} . We denote by $U_m \subset \mathcal{L}$ the analytic continuation of this lifting to the simply connected domain U (4). Let $[\gamma_i, \gamma_j] := \gamma_i\gamma_j\gamma_i^{-1}\gamma_j^{-1}$ and consider the following cases:

- (1) $\gamma_i\gamma_j^{-1} \in \text{hol}^{-1}(\text{Id}_{\mathbf{C}})$. Then $U_{[\gamma_i, \gamma_j]}$ is homeomorphic to a torus minus a disc.
- (2) $\gamma_i\gamma_j^{-1} \notin \text{hol}^{-1}(\text{Id}_{\mathbf{C}})$. Then $U_{[\gamma_i, \gamma_j]}$ is homeomorphic to the annular domain $\{z \in \mathbf{C} \mid 1 < |z| < 2\}$. The domain $U_{[\gamma_i^{-1}, \gamma_j][\gamma_i, \gamma_j]}$ is homeomorphic to torus minus a disc, for the conditions $\gamma_i^2 \in \text{hol}^{-1}(\text{Id}_{\mathbf{C}})$ and $\gamma_j^2 \in \text{hol}^{-1}(\text{Id}_{\mathbf{C}})$ are mutually exclusive. \square

Such a pair of generators γ_i, γ_j always exists whenever $\lambda_j \in \mathbf{C}^* \setminus \mathbf{Q}$, for all j . Suppose the opposite, that is, for every $i \neq j$, $1 \leq i, j \leq \nu + 1$, we have $\gamma_i\gamma_j \in \text{hol}^{-1}(\text{Id}_{\mathbf{C}})$. Then, for every such a pair, $\lambda_i + \lambda_j = q_{ij} \in \mathbf{Z}$. Since $\sum_{j=1}^{\nu+1} \lambda_j \in \mathbf{Z}$, such integral relations imply that $\lambda_{j_0} \in \mathbf{Z}$, for some $1 \leq j_0 \leq \nu + 1$, which is a contradiction.

Whenever a pair $\{\lambda_i, \lambda_j\}$ satisfying the hypothesis of lemma 3 co-exists with $\lambda_k \in \mathbf{C}^* \setminus \mathbf{Q}$, k not necessarily different from i or j , the action of the free-torsion group $\{\text{hol}(\gamma_k)^n\}_{n \in \mathbf{Z}}$ on the fibers of the covering $\pi_{\perp} : \mathcal{L} \rightarrow B_a$ generates a countable family of handles. In particular, every generic leaf $\mathcal{L} \in \mathcal{F}_{a, \lambda}$, for which λ is not rational, has infinite genus. This concludes the proof of lemma 2. \square

Semi-rational case, the set E' . Without loss of generality we assume that $\lambda = (\lambda_1, \dots, \lambda_s, \lambda_{s+1}, \dots, \lambda_{\nu+1})$, where $\lambda_j \in \mathbf{Q}$ for $1 \leq j \leq s$ and $\lambda_j \in \mathbf{C}^* \setminus \mathbf{Q}$ for $s + 1 \leq j$. In the chart $z_2 = tz_1$ of $\mathbf{CP}(2)$, the first integral (2) has the form $F_{a, \lambda}(z_1, t) := z_1^K \prod_{j=1}^{\nu+1} (t - a_j)^{\lambda_j}$. Let $i \leq s$ and $\lambda_i = \frac{p_i}{q_i} \in \mathbf{Q}$. Then, any generic leaf $\mathcal{L} \in \widetilde{\mathcal{F}}_{a, \lambda}$, is given by a connected component of

$$(7) \quad z_1^K (t - a_i)^{p_i/q_i} \prod_{j \neq i} (t - a_j)^{\lambda_j} = z_0,$$

where $z_0 \in \mathbf{C}^*$ is constant. For $t \sim a_i$, $\prod_{j \neq i} (t - a_j)^{\lambda_j}$ is a unity. Then, up to a coordinate change, the generic leaf \mathcal{L} defined by (7) is given, in a neighborhood $t \sim a_i$, by a countable set of algebraic branches of the form

$$(8) \quad z_1^K (t - a_i)^{p_i/q_i} = \text{constant}.$$

For every such a branch, we compactify the abstract surface \mathcal{L} by adding the points corresponding to the limit $t \rightarrow a_i$ in (8). We denote $\overline{\mathcal{L}}$ topological surface obtained from this local compactification process when the index i varies in $\{1, \dots, s\}$. We define $E' := \overline{\mathcal{L}} \setminus \mathcal{L}$.

For example, if λ is rational, then $\overline{\mathcal{L}}$ is a compact orientable surface whose genus can be calculated from (3) using the RIEMANN-HURWITZ formula. For instance, if $\lambda \in \{(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}), (\frac{1}{2}, \frac{1}{4}, \frac{1}{4}), (\frac{1}{2}, \frac{1}{3}, \frac{1}{6})\}$, then $\overline{\mathcal{L}}$ is homeomorphic to the standard torus $\mathbf{S}^1 \times \mathbf{S}^1$.

Lemma 4. *If λ is semi-rational, then the ideal boundary of $\overline{\mathcal{L}}$ is just a point.*

Proof. As before, we claim that every compact subset $K \subset \overline{\mathcal{L}}$ can be covered by a compact subset $K' \subset \overline{\mathcal{L}}$ such that $\overline{\mathcal{L}} \setminus K'$ is connected by arcs. In deed, for positive real number $\tau < \inf_{i \neq j, i, j = s+1, \dots, \nu+1} \delta(a_i, a_j)$, we define V_τ to be the complement in U of the set $\bigcup_{j=s+1}^{\nu+1} \{t \in \mathbf{P}^1(\mathbf{C}) \mid \delta(t, a_j) < \tau\}$. As in the proof of lemma 2, we define $V_{\tau, \xi}$ to be the lifting of V_τ to a point ξ in a generic fiber of $\pi|$.

We consider $\tau(K)$ as in (5), but taking the infimum over the set $j = s+1, \dots, \nu+1$. To define K' it is sufficient to consider a the closure in $\overline{\mathcal{L}}$ of a set

$$(9) \quad \bigcup_{\xi \in Z} V_{\tau, \xi},$$

where $Z \subset \pi|^{-1}(t)$, $t \neq a_1, \dots, a_{\nu+1}, t_0$, is a "sufficiently large" finite subset. The rest of the proof is analog to the proof of lemma 2. Indeed, considering different cases, any two points $q \neq q'$ in $\overline{\mathcal{L}}$ can be joined by an arc, for the action of any free-torsion group $\{\text{hol}(\gamma_j)^n\}_{n \in \mathbf{Z}}$, $j = s+1, \dots, \nu+1$, on the covering $\pi| : \mathcal{L} \rightarrow B_a$ permits to scape to any of compact subsets K' in $\overline{\mathcal{L}}$ previously constructed. \square

Corollary 1. *Let λ be a semi-rational. Then, the ideal boundary of the generic leaf $\mathcal{L} \in \mathcal{F}_{a, \lambda}$ is homeomorphic to $0 \cup \{1/n\}_{n \in \mathbf{N}} \subset \mathbf{C}$ endowed with the subspace topology.*

If two coordinates λ_i, λ_j of a semi-rational point λ meet the hypothesis of lemma 3, then the ideal boundary of the generic leaf of $\mathcal{F}_{a, \lambda}$ presents a distinguished point $B'(\mathcal{L})$, corresponding to 0 in terms of the preceding corollary, representing the not planar end of \mathcal{L} . In such a case, \mathcal{L} is homeomorphic to the LOCH NESS monster punctured in a countable set of points. In a semi-rational point λ , such two coordinates always exist in the complement of the set:

$$(10) \quad \lambda_j \in \mathbf{Z}, \quad \forall j \leq s \quad \text{and} \quad \lambda_j + \lambda_k \in K\mathbf{Z}, \quad \forall s+1 \leq j, k.$$

Lemma 5. *If a semirational point $\lambda = (\lambda_1, \dots, \lambda_s, \lambda_{s+1}, \dots, \lambda_{\nu+1})$ satisfies (10), then the generic leaf of $\mathcal{F}_{a,\lambda}$ is homeomorphic to a LOCH NESS monster punctured in a countable set of points.*

Proof. First we prove the lemma in the illustrative case $\nu = 2$. We assume that $(a_1, a_2, a_3) = (0, 1, \infty)$. Let $W := \mathbf{P}^1(\mathbf{C}) \setminus ([\infty, 0] \cup [1, \infty])$. From condition $\lambda_1 + \lambda_2 + \lambda_3 = K$, we deduce that it is sufficient to consider the action of the holonomy generators $z \rightsquigarrow e^{2\pi i \lambda_j / K} z$, $j = 2, 3$ to reconstruct the generic leaf \mathcal{L} from W . We identify W with figure 1




Figure 1.

Then, the generic leaf \mathcal{L} is homeomorphic to




Figure 2.

which clearly shows that \mathcal{L} is homeomorphic to $\mathbf{C} \setminus \mathbf{Z}[i]$. When ν takes values greater than 2 but $\nu - s$ remains equal to 2, a new discrete set of punctures appears in the leaf \mathcal{L} , for the image on every generator γ_j , $j \leq s$, in $Diff(\mathbf{C}, 0)$ is trivial.

We remark that, if $\nu - s > 1$, there is no semirational point λ satisfying (10). Indeed, in such a case set $\lambda_j = p_j - \lambda_{s+1}$, $p_j \in \mathbf{Z}$, for all $s + 1 < j$. Then $\sum_{j=1}^{\nu+1} \lambda_j = K$ becomes

$$(1 - (\nu - s))\lambda_{s+1} = K',$$

for a certain $K' \in \mathbf{Z}$. This is a contradiction, for λ is semirational. \square

4. APPLICATIONS TO POLYGONAL BILLIARDS

Let $\lambda_1\pi, \dots, \lambda_{\nu+1}\pi$ be the angles of the non degenerated $\nu + 1$ -side polygon P , $\nu \geq 2$. That is, $0 < \lambda_j < 2$, for all $j = 1, \dots, \nu + 1$ and $\sum_{j=1}^{\nu+1} \lambda_j = \nu - 1$. Let $\hat{n} := (n_1, \dots, n_{\nu+1}) \in \mathbf{Z}^{\nu+1}$ and $\psi(\hat{n}) := \sum_{j=1}^{\nu+1} n_j \lambda_j$. We define $G := \psi^{-1}(\mathbf{Z})$ and $\tilde{\psi}(g) := e^{\frac{2\pi i}{\nu-1} \phi(g)}$, for every

$g \in G$. Then $\tilde{G} := \tilde{\psi}(G)$ is a subgroup of $G_{\nu-1} := \{z \in \mathbf{C} \mid z^{\nu-1} = 1\}$. Polygons for which $\tilde{G} = \{1\}$ are called *reasonable*. For example, all triangles are reasonable. Rectangles are *not reasonable*. This nomenclature comes from the same reference as the following

Theorem 4. [7] *Let P be a reasonable polygon. Then, there exist $a \in \mathbf{C}^{\nu+1}$ such that the translation surface $S(P)$ and the generic leaf of $\mathcal{F}_{a,\lambda}$ are isomorphic as translation surfaces.*

The determination of the parameter $a \in \mathbf{C}^{\nu+1}$ is a SCHWARZ-CHRISTOFFEL parameter problem [3]. Above, the translation surface structure of a leaf $\mathcal{L} \in \mathcal{F}_{a,\lambda}$ is defined by the restriction of a holomorphic vector field in \mathbf{C}^2 generating $\text{Ker}(\omega_{a,\lambda}(p))$, for every $p \in \mathbf{C}^2$. The action:

$$(11) \quad \mathbf{C}^2 \times \tilde{G} \ni ((z_1, z_2), \tilde{g}) \rightsquigarrow \tilde{g}(z_1, z_2)$$

has no fixed points except for the origin. It leaves invariant each connected component in the "fibers" of the first integral $F_{a,\lambda}$. We denote \mathbf{C}_λ^2 the quotient of $\mathbf{C}^2 \setminus \{z_1 = z_2 = 0\}$ by the preceding action. Let $\pi_\lambda : \mathbf{C}^2 \rightarrow \mathbf{C}_\lambda^2$ be the natural projection. Then, $\{\pi_\lambda(\mathcal{L}) \mid \mathcal{L} \in \mathcal{F}_{a,\lambda}\}$ is a non singular foliation on \mathbf{C}_λ^2 . We denote it $\widehat{\mathcal{F}}_{a,\lambda}$. The image by π_λ of a generic leaf in $\mathcal{F}_{a,\lambda}$ is called generic as well.

Theorem 5. [7] *Let P be a not reasonable. Then, there exist $a \in \mathbf{C}^{\nu+1}$ such that the translation surface $S(P)$ and the generic leaf of the foliation $\widehat{\mathcal{F}}_{a,\lambda}$ are isomorphic as translation surfaces.*

Here the translation surface structure any generic leaf in $\widehat{\mathcal{F}}_{a,\lambda}$ is inherited from any generic leaf in $\mathcal{F}_{a,\lambda}$ contained in its preimage via the projection π_λ . By definition, if $\mathcal{L} \in \mathcal{F}_{a,\lambda}$ is generic, then $\pi_\lambda|_{\mathcal{L}}$ is a finite covering. The monodromy group of this covering is isomorphic to \tilde{G} .

Corollary 2. *Let P be a non degenerated polygon of angles $\lambda_1\pi, \dots, \lambda_{\nu+1}\pi$ without vertices.*

- (1) *If λ is algebraic, then $S(P)$ is homeomorphic to a compact RIEMANN surface punctured in a finite set of points.*
- (2) *If λ is semi-rational, then $S(P)$ is homeomorphic to a LOCH NESS monster punctured in a countable set of points.*
- (3) *If λ is not rational, then $S(P)$ is homeomorphic to a LOCH NESS monster.*

Certainly, since we are dealing with non-degenerated polygons, there is no λ in this case satisfying the conditions (10). For (2)-(3)

above, it is sufficient to remark that, if M is a LOCH NESS monster and $M \rightarrow N$ is a covering map whose fibers are finite, then N is also a LOCH NESS monster.

REFERENCES

- [1] Cerveau, D. *Densité des feuilles de certaines équations de Pfaff à 2 variables*. Ann. Inst. Fourier (Grenoble), 33 (1983), no. 1, :185-194.
- [2] Cerveau, D. and Mattei, J.-F. *Formes intégrables holomorphes singulières*. Astérisque, 97. Société Mathématique de France, Paris, 1982. 193pp.
- [3] Driscoll, T. A.; Trefethen, L. N. *Schwarz-Christoffel mapping*. Cambridge monographs on Applied and Computational Mathematics, 8. Cambridge University Press, Cambridge, 2002.
- [3] Ghys, E. *Topologie des feuilles génériques*. Ann. of Math. (2) 141 (1995), no. 2, 387–422.
- [4] Masur, H.; Tabachnikov S. *Rational billiards and flat structures*. Handbook of dynamical systems. Vol. 1A. 1015–1089, North-Holland, Amsterdam, 2002.
- [5] Paul, E. *Étude topologique des formes logarithmiques fermées*. Invent. Math. 95 (1989), no. 2, 395–420.
- [6] Richards, I. *On the classification of noncompact surfaces*. Trans. Amer. Math. Soc., 106, 1963, 259–269.
- [7] Valdez, F. *Feuilletages homogènes et billiards polygonaux*. PhD. Thesis, Université de Rennes 1, 2007.
- [8] Zemljakov, A.N. and Katok, A.B. *Topological transitivity of billiards in polygons*. Mat. Zametki. 18 (1975), no. 2, 291–300.

MAX PLANCK INSTITUT FÜR MATHEMATIK VIVATSGASSE 7. 53111, BONN, GERMANY.

E-mail address: ferran@mpim-bonn.mpg.de