


# Max-Planck-Institut für Mathematik Bonn

A remark on connective  $K$ -theory

by

Nikita A. Karpenko





# A remark on connective $K$ -theory

Nikita A. Karpenko

Max-Planck-Institut für Mathematik  
Vivatsgasse 7  
53111 Bonn  
Germany

Mathematical & Statistical Sciences  
University of Alberta  
Edmonton  
Canada



# A REMARK ON CONNECTIVE $K$ -THEORY

NIKITA A. KARPENKO

ABSTRACT. Let  $X$  be a smooth algebraic variety over an arbitrary field. Let  $\varphi_X$  be the canonical surjective homomorphism of the Chow ring of  $X$  onto the ring associated with the Chow filtration on the Grothendieck ring  $K(X)$ . We remark that  $\varphi_X$  is injective if and only if the connective  $K$ -theory  $CK(X)$  coincides with the terms of the Chow filtration on  $K(X)$ . As a consequence,  $CK(X)$  turns out to be computed for numerous flag varieties (under semisimple algebraic groups) for which the injectivity of  $\varphi_X$  had already been established. This especially applies to the so-called *generic* flag varieties  $X$  of many different types, identifying for them  $CK(X)$  with the terms of the explicit Chern filtration on  $K(X)$ .

## CONTENTS

1. Introduction	1
2. The remark	2
3. Applications to flag varieties	4
References	4

## 1. INTRODUCTION

Let  $F$  be an arbitrary field, let  $G$  be a split semisimple algebraic group over  $F$ , and let  $P$  be one of its parabolic subgroups. For any  $G$ -torsor  $E$  over any extension field of  $F$ , the quotient  $X := E/P$  is a variety of parabolic subgroups (a *flag variety* for short) in the (possibly non-split) semisimple group  $\mathrm{Aut}_G E$ , a twisted form of  $G$  over the extension. We call the flag variety  $X$  *generic*, provided that  $E$  is a (standard) generic  $G$ -torsor, i.e., the generic fiber of the quotient map  $\mathrm{GL}(n) \rightarrow \mathrm{GL}(n)/G$  for an embedding of  $G$  into  $\mathrm{GL}(n)$ .

Assume that  $P$  is *special*, i.e., all  $P$ -torsors over all extension fields of  $F$  are trivial. (For instance,  $P$  can be a Borel subgroup of  $G$ .) The following conjecture appears first in [6, §1] in form of a question. It deals with the canonical (surjective) homomorphism of graded rings

$$\varphi_X: \mathrm{CH}(X) \rightarrow \mathrm{Chow}K(X),$$

---

*Date:* 11 June 2019.

*Key words and phrases.* (Connective)  $K$ -theory; Chow groups; algebraic groups; generic torsors; projective homogeneous varieties. *Mathematical Subject Classification (2010):* 19L41; 14C25; 20G15.

This work has been accomplished during author's stay at the Max-Planck Institute for Mathematics in Bonn.

where  $\text{CH}(X)$  is the Chow ring,  $K(X)$  is the Grothendieck ring of  $X$ , and  $\text{Chow}K(X)$  is the ring associated with the Chow filtration (i.e., the filtration by codimension of supports of coherent sheaves) on  $K(X)$ .

**Conjecture 1.1** ([5, Conjecture 1.1]). *The homomorphism  $\varphi_X$  is an isomorphism.*

Being recently disproved for  $G = \text{Spin}(17)$  by Yagita in [10] (see also [4]), Conjecture 1.1 has been confirmed for many other  $G$ . (For instance, the 2-local version for  $G$  of type  $E_7$  is proved in the very [4].) An overview of some positive cases is given in [5]. (On the other hand, for many  $G$  it is still unknown if the above conjecture holds or fails.)

For an arbitrary smooth variety  $X$ , the homomorphism  $\varphi_X$  provides a sort of connection between the Chow theory of  $X$  and its  $K$ -theory. Another standard way to connect those two theories goes through the *connective  $K$ -theory*  $CK(X)$  (see §2). In this note we remark that Conjecture 1.1 can be expressed in terms of  $CK(X)$ . Namely, we prove (see Theorem 2.2) that the injectivity of  $\varphi_X$  actually means  $CK(X)$  coincides with the terms of the Chow filtration on  $K(X)$ .

Note that  $K(X)$  is computed for arbitrary flag variety  $X$ , but not the Chow filtration, which is a finer invariant and remains quite mysterious. However, for a generic flag variety  $X$  given by a special parabolic  $P$ , as in Conjecture 1.1, the Chow filtration coincides with the explicitly computable Chern filtration (more known under the name of gamma filtration), introduced by Grothendieck (see §3). So, Conjecture 1.1 for a given  $X$  turns out to be equivalent to the complete computation of  $CK(X)$ .

## 2. THE REMARK

For any smooth algebraic variety  $X$  over an arbitrary field  $F$  (of arbitrary characteristic), we write  $CK(X) = \bigoplus_{i \in \mathbb{Z}} CK^i(X)$  for the connective  $K$ -theory ring of  $X$ , graded by codimension. Our main reference for the connective  $K$ -theory is [2] (see also [1]) and our  $CK^i(X)$  is the group  $CK^{i,-i}(X)$  of [2, §6.4]. (We only work with small cohomology theories and, in particular, do not use the higher connective  $K$ -theory groups here.) To recall the definition of  $CK^i(X)$ , let  $M^i(X)$  be the Grothendieck group of the category of coherent sheaves on  $X$  with codimension of support  $\geq i$ . Then  $CK^i(X)$  is defined as the image of the homomorphism  $M^i(X) \rightarrow M^{i-1}(X)$  mapping the class of a sheaf to the class of itself.

Since  $M^i(X)$  is the Grothendieck group  $K(X)$  for  $i \leq 0$ ,  $CK^i(X)$  is identified with  $K(X)$  for such  $i$ . Also note that  $CK^i(X) = 0$  for  $i > \dim X$ .

The Grothendieck group  $K(X)$  is actually also a ring (with multiplication given by tensor product of locally-free sheaves) and it is endowed with the Chow filtration (see [8]), i.e., the filtration by codimension of supports of coherent sheaves:

$$K(X) = \dots = K^{(-1)}(X) = K(X)^{(0)} \supset K^{(1)}(X) \supset \dots$$

Since  $K^{(i)}(X) \cdot K^{(j)}(X) \subset K^{(i+j)}(X)$  for any  $i, j \in \mathbb{Z}$ , we may consider a graded ring

$$K^0(X) := \bigoplus_{i \in \mathbb{Z}} K^{(i)}(X),$$

where  $K^{(i)}(X) = 0$  for  $i > \dim X$ . Note that, unlike  $CK$ , the localization sequence

$$K^{(-\dim Y)}(Y) \rightarrow K^0(X) \rightarrow K^0(U) \rightarrow 0$$

for the theory  $K^0$ , relating the theory of  $X$  with the theory of a smooth closed subvariety  $Y \subset X$  and its open complement  $U$ , is not always exact at the term  $K^0(X)$ . (Exactness of the localization sequence for the connective  $K$ -theory is a part of [2, Theorem 5.1].)

Finally, we are considering the Chow ring  $\text{CH}(X) = \bigoplus_{i \in \mathbb{Z}} \text{CH}^i(X)$  of rational equivalence classes of algebraic cycles on  $X$ , graded by codimension of cycles. Here we also have  $\text{CH}^i(X) = 0$  for  $i > \dim X$ . Besides,  $\text{CH}^i(X) = 0$  for  $i < 0$ .

The connective  $K$ -theory “connects”  $\text{CH}(X)$  with  $K(X)$ , or, more precisely, with  $K^0(X)$  by means of canonical surjective homomorphisms of graded rings

$$CK(X) \rightarrow \text{CH}(X) \quad \text{and} \quad \psi_X: CK(X) \rightarrow K^0(X).$$

By [2, Theorem 7.1], the kernel of the first one is generated by the *Bott element*  $\beta \in CK^{-1}(X)$  defined as the unit of the ring  $K(X)$ , considered as an element of  $K^{(-1)}(X) = CK^{-1}(X)$ .

Abusing notation, let us consider the Laurent polynomial ring  $K(X)[\beta^{\pm 1}]$  in one variable  $\beta$  (which we continue to call Bott element). The ring  $K^0(X)$  can be defined as the subring of  $K(X)[\beta^{\pm 1}]$  consisting of the polynomials  $\sum_{i \in \mathbb{Z}} a_i \beta^i$  with  $a_i \in K^{(-i)}(X)$  for all  $i$ . Since  $\beta$  is invertible in  $K(X)[\beta^{\pm 1}]$ , it is not a zero divisor in  $K^0(X)$ .

Again by [2, Theorem 7.1], the composition

$$CK(X) \xrightarrow{\psi_X} K^0(X) \hookrightarrow K(X)[\beta^{\pm 1}]$$

is the localization of the ring  $CK(X)$  with respect to the element  $\beta \in CK(X)$ . In particular,  $\psi_X$  is an isomorphism if and only if  $\beta$  is not a zero divisor in  $CK(X)$ .

The quotient  $K^0(X)/\beta K^0(X)$  is the graded ring  $\text{Chow}K(X)$  associated with the Chow filtration on  $K(X)$ . The canonical surjective homomorphism of graded rings

$$\varphi_X: \text{CH}(X) \rightarrow \text{Chow}K(X),$$

mapping the class of a closed subvariety to the class of its structure sheaf, fits into the commutative square

$$(2.1) \quad \begin{array}{ccc} CK(X) & \xrightarrow{\psi_X} & K^0(X) \\ \downarrow & & \downarrow \\ \text{CH}(X) & \xrightarrow{\varphi_X} & \text{Chow}K(X). \end{array}$$

We recall that the kernel of  $\varphi_X$  consists of elements of finite order. More precisely, the kernel on  $\text{CH}^i(X)$  is trivial for  $i \leq 2$  and is killed by  $(i-1)!$  for  $i \geq 1$ , [3, Example 15.3.6].

**Theorem 2.2.** *For any given smooth algebraic variety  $X$  (over an arbitrary field), the homomorphism  $\psi_X$  is an isomorphism if and only if  $\varphi_X$  is.*

*Proof.* The homomorphism  $\psi_X$  induces  $\varphi_X$  by modding out the ideals in  $CK(X)$  and in  $K^0(X)$  generated by the Bott element. So,  $\varphi_X$  is an isomorphism provided that  $\psi_X$  is.

Conversely, let us assume that  $\text{Ker}(\varphi_X) = 0$  and let us take an element  $x_0 \in CK(X)$  vanishing in  $K^0(X)$  under  $\psi_X$ . Note that  $x_0$  is concentrated in positive degrees:

$$x_0 \in CK^{>0}(X) := \bigoplus_{i>0} CK^i(X).$$

(We do not need to assume it to be homogeneous.) From the commutative square (2.1), we conclude that  $x$  vanishes also in  $\mathrm{CH}(X)$ , so that  $x_0 = \beta x_1$  for some  $x_1 \in \mathrm{CK}^{>1}(X)$ . Since  $\beta \in K^0(X)$  is not a zero divisor,  $x_1$  also vanishes in  $K^0(X)$  under  $\psi_X$  so that  $x_1 = \beta x_2$  and  $x_0 = \beta^2 x_2$  for some  $x_2 \in \mathrm{CK}^{>2}(X)$ . Continuing this way, we manage to write  $x_0$  as  $x_0 = \beta^i x_i$  with some  $x_i \in \mathrm{CK}^{>i}(X)$  for any  $i \geq 0$ . But  $\mathrm{CK}^{>i}(X)$  is trivial for  $i = \dim X$ . It follows that  $x_0$  and  $\mathrm{Ker}(\psi_X)$  are trivial.  $\square$

**Remark 2.3.** Replacing the integer coefficients by rational coefficients for the cohomology theories in the above considerations, we come to the situation, where  $\varphi_X$  is an isomorphism for any  $X$ . It follows that  $\psi_X$  with rational coefficients is always an isomorphism as well. Turning back to the integer coefficients, we see that every element in the kernel of  $\psi_X$  is of finite order.

### 3. APPLICATIONS TO FLAG VARIETIES

Now we fix a semisimple algebraic group  $G$  over  $F$  and consider a projective homogeneous variety (*flag variety* for short)  $X$  under  $G$ . In other terms,  $X$  is a variety of parabolic subgroups in  $G$ . We fix an algebraic closure  $\bar{F}$  of  $F$  and write  $\bar{X}$  for  $X_{\bar{F}}$ . Let us write down an extended version of Theorem 2.2 which holds for such  $X$ :

**Theorem 3.1.** *The following conditions on  $X$  are equivalent.*

- (1) *The homomorphism  $\varphi_X$  is an isomorphism.*
- (2) *The homomorphism  $\psi_X$  is an isomorphism.*
- (3) *The group  $\mathrm{CK}(X)$  is torsion-free.*
- (4) *The change of field homomorphism  $\mathrm{CK}(X) \rightarrow \mathrm{CK}(\bar{X})$  is injective.*

*Proof.* We already know by Theorem 2.2 that (1) and (2) are equivalent. By Remark 2.3, (3) implies (2). Since the group  $K^0(X)$  is torsion-free (by [9]), (2) implies (3) as well. By transfer argument, (3) implies (4). Finally, the group  $\mathrm{CK}(\bar{X})$  is torsion-free (e.g., because  $\mathrm{CH}(\bar{X})$  is torsion-free), implying that  $\varphi_{\bar{X}}$  and  $\psi_{\bar{X}}$  are isomorphisms; consequently (4) implies (3) as well.  $\square$

To get the most from Theorem 3.1, let us put more restrictions on  $X$ : assume that  $X$  is a *generic* flag variety (as defined in the introduction) given by a split semisimple group  $G$  and a *special* parabolic subgroup  $P \subset G$ . By [7, Corollary 7.4], the Chow filtration on  $K(X)$  coincides in this case with the Chern filtration. Therefore  $\mathrm{CK}(X)$  is given by the terms of the Chern filtration as long as Conjecture 1.1 holds for  $G$ .

On the other hand, the counter-example of [10] (see also [4]) provides by Theorem 3.1 a generic flag variety  $X$  (given by the spinor group  $\mathrm{Spin}(17)$ ) with non-trivial torsion in  $\mathrm{CK}(X)$ .

**ACKNOWLEDGEMENTS.** Theorem 3.1 has been inspired by [10, Lemma 7.5]. I thank Alexander Merkurjev for useful comments.

### REFERENCES

- [1] CAI, S. A simple description of algebraic connective  $K$ -theory. PhD Thesis, Los Angeles, USA, 2006, 108 pages.



- [2] CAI, S. Algebraic connective  $K$ -theory and the niveau filtration. *J. Pure Appl. Algebra* 212, 7 (2008), 1695–1715.
- [3] FULTON, W. *Intersection theory*, second ed., vol. 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1998.
- [4] KARPENKO, N. A. A counter-example by Yagita. Preprint (22 May 2019, 9 pages), available on author's web page.
- [5] KARPENKO, N. A. Chow ring of generic flag varieties. *Math. Nachr.* 290, 16 (2017), 2641–2647.
- [6] KARPENKO, N. A. Chow ring of generically twisted varieties of complete flags. *Adv. Math.* 306 (2017), 789–806.
- [7] KARPENKO, N. A. On generically split generic flag varieties. *Bull. Lond. Math. Soc.* 50 (2018), 496–508.
- [8] KARPENKO, N. A., AND MERKURJEV, A. S. Chow filtration on representation rings of algebraic groups. *International Mathematics Research Notices*, to appear.
- [9] PANIN, I. A. On the algebraic  $K$ -theory of twisted flag varieties. *K-Theory* 8, 6 (1994), 541–585.
- [10] YAGITA, N. The Gamma filtration for the Spin groups. arXiv:1811.08288 [math.KT], 17 Nov 2018, 27 pages.

MATHEMATICAL & STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, CANADA  
E-mail address: karpenko at ualberta.ca, web page: [www.ualberta.ca/~karpenko](http://www.ualberta.ca/~karpenko)