


Max-Planck-Institut für Mathematik Bonn

Three embeddings of the Klein simple group into the
Cremona group of rank three

by

Ivan Cheltsov
Constantin Shramov



Three embeddings of the Klein simple group into the Cremona group of rank three

Ivan Cheltsov
Constantin Shramov

Max-Planck-Institut für Mathematik
Vivatsgasse 7
53111 Bonn
Germany

University of Edinburgh
Edinburgh EH9 3JZ
United Kingdom

Steklov Institute of Mathematics
Moscow 119991
Russia

THREE EMBEDDINGS OF THE KLEIN SIMPLE GROUP INTO THE CREMONA GROUP OF RANK THREE

IVAN CHELTSOV AND CONSTANTIN SHRAMOV

ABSTRACT. We show that the Cremona group of rank three has at least three non-conjugate subgroups that are isomorphic to the Klein simple group consisting of 168 elements.

Unless explicitly stated otherwise, varieties are assumed to be projective, normal and complex.

1. INTRODUCTION

The Cremona group of rank n , usually denoted by $\text{Cr}_n(\mathbb{C})$, is the group of birational automorphisms of the complex projective space \mathbb{P}^n . One easily checks that

$$\text{Cr}_n(\mathbb{C}) = \text{Aut}(\mathbb{P}^n) \iff n = 1.$$

The group $\text{Cr}_2(\mathbb{C})$ is well studied (see [9]), but we know little about $\text{Cr}_n(\mathbb{C})$ for $n \geq 3$ (cf. [31]).

Problem 1.1. Describe finite subgroups in the group $\text{Cr}_n(\mathbb{C})$ up to isomorphism.

Problem 1.2. Describe finite subgroups in the group $\text{Cr}_n(\mathbb{C})$ up to conjugation.

For $\text{Cr}_1(\mathbb{C})$, the solutions to both Problems 1.1 and 1.2 is given by Platonic solids.

Example 1.3 ([33]). Suppose that G is a finite group. Then

$$\text{Cr}_1(\mathbb{C}) \text{ has a subgroup isomorphic to } G \iff G \in \left\{ \mathbb{Z}_m, \mathbb{Z}_2 \times \mathbb{Z}_2, D_n, A_4, S_4, A_5 \right\},$$

where $m \geq 2$ and $n \geq 4$. Two finite subgroups in $\text{Cr}_1(\mathbb{C})$ are conjugate if they are isomorphic.

For the group $\text{Cr}_2(\mathbb{C})$, the solution to Problem 1.1 is very close to being complete (see [9]), but the solution to Problem 1.2 is known only for some classes of subgroups.

Example 1.4 ([2, Theorem B.2]). Let G be a finite simple non-abelian subgroup. Then

$$\text{Cr}_2(\mathbb{C}) \text{ has a subgroup isomorphic to } G \iff G \in \left\{ A_5, \text{PSL}_2(\mathbb{F}_7), A_6 \right\},$$

and $\text{Cr}_2(\mathbb{C})$ has 3, 2 and 1 non-conjugate subgroups isomorphic to $A_5, \text{PSL}_2(\mathbb{F}_7), A_6$, respectively.

For $\text{Cr}_n(\mathbb{C})$ and $n \geq 3$, it seems hopeless to solve Problems 1.1 and 1.2 in full generality.

Theorem 1.5 ([31, Theorem 1.3]). Suppose that G is a finite simple non-abelian subgroup. Then

$$\text{Cr}_3(\mathbb{C}) \text{ has a subgroup isomorphic to } G \iff G \in \left\{ A_5, \text{PSL}_2(\mathbb{F}_7), A_6, A_7, \text{PSL}_2(\mathbb{F}_8), \text{PSU}_4(\mathbb{F}_2) \right\}.$$

Arguing as in the proof of [31, Theorem 1.3], one can easily obtain the following result.

Theorem 1.6 ([5, Corollary 1.11]). The group $\text{Cr}_3(\mathbb{C})$ contains 1, 2 and at most 2 non-conjugate subgroups that are isomorphic to the groups $\text{PSL}_2(\mathbb{F}_8), \text{PSU}_4(\mathbb{F}_2)$ and A_7 , respectively.

So far, nothing is known about the number of non-conjugate subgroups in $\text{Cr}_3(\mathbb{C})$ that are isomorphic to the group A_5 . As for the group A_6 , we have the following statement.

The authors thank I. Dolgachev, Th. Köppe, A. Kuznetsov and Yu. Prokhorov for their help.
The first author thanks the Max-Planck-Institute für Mathematik at Bonn for hospitality.

Theorem 1.7 ([5]). The group $\mathrm{Cr}_3(\mathbb{C})$ has at least 5 non-conjugate subgroups isomorphic to A_6 .

The main purpose of this paper is to prove the following result.

Theorem 1.8. The group $\mathrm{Cr}_3(\mathbb{C})$ has at least 3 non-conjugate subgroups isomorphic to $\mathrm{PSL}_2(\mathbb{F}_7)$.

Let G be a finite group such that $G \cong \mathrm{PSL}_2(\mathbb{F}_7)$. Then any embedding $G \hookrightarrow \mathrm{Cr}_3(\mathbb{C})$ is arising from some rational threefold admitting a faithful action of the group G (for details see [9], [31]), and the first examples of such embeddings are coming from representation theory.

Example 1.9. Up to conjugation, the group $\mathrm{Aut}(\mathbb{P}^3)$ has two subgroups isomorphic to $\mathrm{PSL}_2(\mathbb{F}_7)$:

- (I) a subgroup that fixes a point in \mathbb{P}^3 , which arises from a faithful reducible four-dimensional representation of the group $\mathrm{PSL}_2(\mathbb{F}_7)$,
- (II) a subgroup that does not fix any point in \mathbb{P}^3 , which arises from a faithful irreducible four-dimensional representation of the group $\mathrm{SL}_2(\mathbb{F}_7)$ (see [6]).

The next example of an embedding $G \hookrightarrow \mathrm{Cr}_3(\mathbb{C})$ comes from the Klein quartic curve (see [21]), which is the unique curve of genus 3 that admits a non-trivial action of the group $\mathrm{PSL}_2(\mathbb{F}_7)$.

Example 1.10 ([23], [15], [25], [24]). Let \mathfrak{C} be the quartic curve in \mathbb{P}^2 that is given by

$$xy^3 + yz^3 + zx^3 = 0 \subset \mathbb{P}^2 \cong \mathrm{Proj}(\mathbb{C}[x, y, z]),$$

put $X = \mathrm{VSP}(\mathfrak{C}, 6)$ (see Definition A.3). Then X is a rational smooth Fano threefold such that

$$\mathrm{Aut}(X) \cong \mathrm{Aut}(\mathfrak{C}) \cong \mathrm{PSL}_2(\mathbb{F}_7),$$

the group $\mathrm{Pic}(X)$ is generated by $-K_X$, and $-K_X^3 = 22$ (see Section 4, Appendix A, [16]).

Remark 1.11. The Fano variety constructed in Example 1.10 is a compactification of the moduli space of $(1, 7)$ -polarized abelian surfaces (see [23], [15] and [25] for details).

In Sections 5 and 6, we prove the following two results, which imply Theorem 1.8.

Theorem 1.12. Let G be a subgroup in $\mathrm{Aut}(\mathbb{P}^3)$ such that $G \cong \mathrm{PSL}_2(\mathbb{F}_7)$ is of type (II). Then

- the variety \mathbb{P}^3 is G -birationally rigid (see [2, Definitions A.3]),
- there is a G -equivariant birational non-biregular involution $\tau \in \mathrm{Cr}_3(\mathbb{C})$ such that

$$\mathrm{Bir}^G(\mathbb{P}^3) = \langle G, \tau \rangle \cong \mathrm{PSL}_2(\mathbb{F}_7) \times \mathbb{Z}_2.$$

Theorem 1.13. Let X be the threefold constructed in Example 1.10. Put $G = \mathrm{Aut}(X)$. Then

$$\mathrm{Bir}^G(X) = \mathrm{Aut}^G(X) = G \cong \mathrm{PSL}_2(\mathbb{F}_7),$$

and the threefold X is G -birationally rigid (see [2, Definitions A.3]).

Corollary 1.14. Let G and G' be subgroups in $\mathrm{Aut}(\mathbb{P}^3)$ such that $G \cong G' \cong \mathrm{PSL}_2(\mathbb{F}_7)$. Then

$$G \text{ and } G' \text{ are conjugate in } \mathrm{Cr}_3(\mathbb{C}) \iff G \text{ and } G' \text{ are conjugate in } \mathrm{Aut}(\mathbb{P}^3).$$

As a by-product of the proofs of Theorems 1.12 and 1.13, we obtain the following result.

Theorem 1.15. Let V be either \mathbb{P}^3 or a threefold constructed in Example 1.10. Then

$$\sup \left\{ \lambda \in \mathbb{Q} \left| \begin{array}{l} \text{the log pair } (V, \lambda D) \text{ has log canonical singularities} \\ \text{for every } G\text{-invariant effective } \mathbb{Q}\text{-divisor } D \sim_{\mathbb{Q}} -K_V \end{array} \right. \right\} = 1,$$

where G is a subgroup in $\mathrm{Aut}(V)$ such that $G \cong \mathrm{PSL}_2(\mathbb{F}_7)$, and G is of type (II) if $V \cong \mathbb{P}^3$.

Proof. Note that $|-K_V|$ contains a G -invariant surface (see Theorem 3.3 and Lemma 4.10). Then

$$\sup \left\{ \lambda \in \mathbb{Q} \left| \begin{array}{l} \text{the log pair } (V, \lambda D) \text{ has log canonical singularities} \\ \text{for every } G\text{-invariant effective } \mathbb{Q}\text{-divisor } D \sim_{\mathbb{Q}} -K_V \end{array} \right. \right\} \leq 1.$$

Suppose that the equality we want to prove is wrong. Then there is an effective \mathbb{Q} -divisor R on the variety V such that R is G -invariant, the log pair (V, R) is not log canonical and $R \sim_{\mathbb{Q}} -K_V$.

Let us show that the existence of the divisor R leads to a contradiction. Put

$$\mu = \sup \left\{ \epsilon \in \mathbb{Q} \left| \text{the log pair } (V, \epsilon R) \text{ is log canonical} \right. \right\} < 1,$$

let S be a minimal center of log canonical singularities of the log pair $(V, \mu R)$ (see [18], [4]), and let Z be the G -orbit of the subvariety S . Then $\dim(S) \leq 1$. This is obvious if V is a threefold constructed in Example 1.10, and this follows from Theorem 3.5 in the case $V = \mathbb{P}^3$.

Take $\epsilon \in \mathbb{Q}$ such that $1 > \epsilon \gg 0$. By Lemma 2.13, there is a G -invariant \mathbb{Q} -divisor D such that

$$D \sim_{\mathbb{Q}} \epsilon R \sim_{\mathbb{Q}} -\epsilon K_V,$$

the singularities of the log pair (V, D) are log canonical, and every minimal center of log canonical singularities of the log pair (V, D) is an irreducible component of the subvariety Z .

Let $\mathcal{I}(V, D)$ be the multiplier ideal sheaf of the log pair (V, D) . Then the sequence

$$0 \rightarrow H^0(\mathcal{O}_V \otimes \mathcal{I}(V, D)) \rightarrow H^0(\mathcal{O}_V) \rightarrow H^0(\mathcal{O}_Z) \rightarrow 0$$

is exact by Theorem 2.2. In particular, we see that $Z = S$.

The variety V does not contain G -invariant points. This is obvious in the case when $V = \mathbb{P}^3$, and this follows from Theorem 4.8 if V is a threefold constructed in Example 1.10.

Thus, we see that S must be a curve. Then S is a smooth and rational curve Theorem 2.10, which is impossible, because the group G cannot act non-trivially on \mathbb{P}^1 . \square

Applying [34, Theorem 2.1], [3, Theorem A.3] and Theorem 1.15, we immediately obtain the following result (cf. [35, Corollary 1.3], [11, §5.3], [11, Theorem 5.4.3]).

Corollary 1.16. Let X be a threefold constructed in Example 1.10. Then X is Kähler–Einstein.

Applying [4, Theorem 3.15] and Theorem 1.15, we obtain the following result.

Corollary 1.17. Let \hat{G} be a subgroup in $\mathrm{SL}_4(\mathbb{C})$ such that $\hat{G} \cong \mathrm{SL}_2(\mathbb{F}_7)$. Then the quotient singularity \mathbb{C}^4/\hat{G} is weakly-exceptional (see [4, Definition 3.8]).

As a by-product of the proof of Theorem 1.13, we give an example of a smooth $K3$ surface admitting a faithful action of the group $\mathrm{PSL}_2(\mathbb{F}_7)$ (see Lemma 4.10), that must be a new example, since we were not able to find this example in the literature.

Remark 1.18. We do not know whether or not the surface provided by Lemma 4.10 is isomorphic as a non-polarized smooth $K3$ surface to one of three smooth $K3$ surfaces that admit faithful actions of the group $\mathrm{PSL}_2(\mathbb{F}_7)$ that have been mentioned in [26] and [29].

We organize this paper in the following way:

- in Section 2, we recall several well-known preliminary results,
- in Section 3, we collect results about the action of the group $\mathrm{PSL}_2(\mathbb{F}_7)$ on \mathbb{P}^3 ,
- in Section 4, we collect results about the threefold constructed in Example 1.10,
- in Section 5, we prove Theorem 1.12 using results obtained in Section 3,
- in Section 6, we prove Theorem 1.13 using results obtained in Section 4,
- in Appendix A, we describe Mukai's construction of Fano threefolds of degree 22,
- in Appendix B, we collect elementary results about the groups $\mathrm{PSL}_2(\mathbb{F}_7)$ and $\mathrm{SL}_2(\mathbb{F}_7)$.

2. PRELIMINARIES

Let X be a variety with at most Kawamata log terminal singularities (see [19, Definition 3.5]), let B_X be an formal \mathbb{Q} -linear combination of prime divisors and mobile linear systems

$$B_X = \sum_{i=1}^r a_i B_i + \sum_{j=1}^s c_j \mathcal{M}_j,$$

where B_i and \mathcal{M}_j are a prime Weil divisor and a linear system on the variety X that has no fixed components, respectively, and $a_i \in \mathbb{Q}_{\geq 0} \ni c_j$. Suppose that B_X is a \mathbb{Q} -Cartier divisor.

Definition 2.1. We say that B_X and (X, B_X) are mobile if $a_1 = a_2 = \dots = a_r = 0$.

Let $\pi: \bar{X} \rightarrow X$ be a log resolution for the log pair (X, B_X) , let \bar{B}_i and $\bar{\mathcal{M}}_j$ be the proper transforms of the divisor B_i and the linear system \mathcal{M}_j on the variety \bar{X} , respectively. Then

$$K_{\bar{X}} + \sum_{i=1}^r a_i \bar{B}_i + \sum_{j=1}^s c_j \bar{\mathcal{M}}_j \sim_{\mathbb{Q}} \pi^*(K_X + B_X) + \sum_{i=1}^m d_i E_i,$$

where E_i is an exceptional divisor of the morphism π , and d_i is a rational number. Put

$$\mathcal{I}(X, B_X) = \pi_* \left(\sum_{i=1}^m [d_i] E_i - \sum_{i=1}^r [a_i] B_i \right),$$

and recall that $\mathcal{I}(X, B_X)$ is known as the multiplier ideal sheaf (see [20, Section 9.2]).

Theorem 2.2 ([20, Theorem 9.4.8]). Let H be a nef and big \mathbb{Q} -divisor on X such that

$$K_X + B_X + H \sim_{\mathbb{Q}} D$$

for some Cartier divisor D on the variety X . Then $H^i(\mathcal{I}(X, B_X) \otimes D) = 0$ for every $i \geq 1$.

Let $\mathcal{L}(X, B_X)$ be a subscheme given by $\mathcal{I}(X, B_X)$. Put $\text{LCS}(X, B_X) = \text{Supp}(\mathcal{L}(X, B_X))$.

Remark 2.3. If the log pair (X, B_X) is log canonical, than the subscheme $\mathcal{L}(X, B_X)$ is reduced.

Let Z be an irreducible subvariety of the variety X .

Definition 2.4 ([17, Definition 1.3]). The subvariety Z is said to be a center of log canonical singularities (non-log canonical singularities, respectively) of the log pair (X, B_X) if

- either $a_i \geq 1$ ($a_i > 1$, respectively) and $Z = B_i$ for some $i \in \{1, \dots, r\}$,
- or $d_i \leq -1$ ($d_i < -1$, respectively) and $Z = \pi(E_i)$ for some $i \in \{1, \dots, m\}$ and some π .

let $\text{LCS}(X, B_X)$ and $\text{NLCS}(X, B_X)$ be the sets of centers of log canonical and non-log canonical singularities of the log pair (X, B_X) , respectively. Then $\text{NLCS}(X, B_X) \subseteq \text{LCS}(X, B_X)$.

Theorem 2.5 ([7, Theorem 3.1]). Suppose that $\dim(X) = 2$, the set $\text{NLCS}(X, B_X)$ contains a point $P \in X \setminus \text{Sing}(X)$, the boundary B_X is mobile and $s = 1$. Then

$$\text{mult}_P(M_1 \cdot M'_1) > 4/c_1^2,$$

where M_1 and M'_1 are general curves in the linear system \mathcal{M}_1 .

Let us denote by $\text{NLCS}(X, B_X)$ the proper subset of the variety X such that

$$P \in \text{NLCS}(X, B_X) \iff \exists S \in \text{NLCS}(X, B_X) \text{ such that } P \in S.$$

Definition 2.6 ([3, Definition 2.2]). The subvariety Z is said to be a center of canonical singularities (non-canonical singularities, respectively) of the log pair (X, B_X) if

$$Z = \pi(E_i)$$

and $d_i \leq 0$ ($d_i < 0$, respectively) for some $i \in \{1, \dots, m\}$ and some choice of the morphism π .

Let $\mathbb{C}\mathbb{S}(X, B_X)$ and $\mathbb{N}\mathbb{C}\mathbb{S}(X, B_X)$ be the sets of centers of canonical and non-canonical singularities of the log pair (X, B_X) , respectively. Then $\mathbb{N}\mathbb{C}\mathbb{S}(X, B_X) \subseteq \mathbb{C}\mathbb{S}(X, B_X)$.

Theorem 2.7 ([7, Corollary 3.4]). Suppose that $\dim(X) = 3$, the set $\mathbb{N}\mathbb{C}\mathbb{S}(X, B_X)$ contains a point $P \in X \setminus \text{Sing}(X)$, the boundary B_X is mobile and $s = 1$. Then

$$\text{mult}_P(M_1 \cdot M'_1) > 4/c_1^2,$$

where M_1 and M'_1 are general surfaces in the linear system \mathcal{M}_1 .

Let us denote by $\mathbb{N}\mathbb{C}\mathbb{S}(X, B_X)$ the proper subset of the variety X such that

$$P \in \mathbb{N}\mathbb{C}\mathbb{S}(X, B_X) \iff \exists S \in \mathbb{N}\mathbb{C}\mathbb{S}(X, B_X) \text{ such that } P \in S.$$

Lemma 2.8. Suppose that X is smooth at a general point of the subvariety Z . Then

- if $Z \in \mathbb{C}\mathbb{S}(X, B_X)$, then $Z \in \mathbb{L}\mathbb{C}\mathbb{S}(X, 2B_X)$,
- if $Z \in \mathbb{N}\mathbb{C}\mathbb{S}(X, B_X)$, then $Z \in \mathbb{N}\mathbb{L}\mathbb{C}\mathbb{S}(X, 2B_X)$.

Proof. This is obvious, because X is smooth at a general point of the subvariety Z . □

Suppose that $Z \in \mathbb{L}\mathbb{C}\mathbb{S}(X, B_X)$ and (X, B_X) is log canonical along the subvariety Z .

Lemma 2.9 ([17, Proposition 1.5]). Let Z' be a center in $\mathbb{L}\mathbb{C}\mathbb{S}(X, B_X)$ such that $Z' \neq Z$ and

$$\emptyset \neq Z \cap Z' = \sum_{i=1}^k Z_i \subsetneq Z,$$

where Z_i is an irreducible subvariety. Then $Z_i \in \mathbb{L}\mathbb{C}\mathbb{S}(X, B_X)$ for every $i \in \{1, \dots, k\}$.

Suppose that Z is a minimal center in $\mathbb{L}\mathbb{C}\mathbb{S}(X, B_X)$ (see [17], [18], [4, Definition 2.8]).

Theorem 2.10 ([18, Theorem 1]). Let Δ be an ample \mathbb{Q} -Cartier \mathbb{Q} -divisor on X . Then

- the variety Z is normal and has at most rational singularities,
- there exists an effective \mathbb{Q} -divisor B_Z on the variety Z such that

$$(K_X + B_X + \Delta)|_Z \sim_{\mathbb{Q}} K_Z + B_Z,$$

and (Z, B_Z) has Kawamata log terminal singularities.

Let G be a finite subgroup of the group $\text{Aut}(X)$.

Lemma 2.11. Let P , C and S be a point, curve and surface in X , respectively. Suppose that

$$\text{Sing}(X) \not\ni P \in C \subset S$$

and $\dim(X) = 3$. Suppose that P and C are G -invariant, and either $G \cong A_4$ or $G \cong \mathbb{Z}_7 \rtimes \mathbb{Z}_3$. Then

$$\text{mult}_P(C) \geq 3$$

and the surface S is singular at the point $P \in X$.

Proof. Let $\gamma: U \rightarrow X$ be a blow up of the threefold X at the point P , let E be the γ -exceptional divisor, and let \bar{C} be the proper transforms of the curve C on the threefold U . Then

$$\text{mult}_P(C) \geq |\bar{C} \cap E|.$$

The group G naturally acts on $E \cong \mathbb{P}^2$. This action comes from a faithful three-dimensional representation of the group G , which must be irreducible, because the group G does not have two-dimensional irreducible representations and the group G is not abelian. Thus

$$|\bar{C} \cap E| \geq 3,$$

and the points of the set $\bar{C} \cap E$ are not contained in a single line in $E \cong \mathbb{P}^2$, which immediately implies that the surface S must be singular at the point $P \in X$. □

Suppose that B_X is G -invariant. Recall that (X, B_X) is log canonical.

Remark 2.12. Let g be an elements in G . Then $g(Z) \in \text{LCS}(X, B_X)$. By Lemma 2.9, we have

$$Z \cap g(Z) \neq \emptyset \iff Z = g(Z).$$

Suppose that B_X is ample. Take an arbitrary rational number $\epsilon > 1$.

Lemma 2.13. There is a G -invariant linear system \mathcal{B} on the variety X that has no fixed components, and there are rational numbers ϵ_1 and ϵ_2 such that $1 \geq \epsilon_1 \gg 0$ and $1 \gg \epsilon_2 \geq 0$ and

$$\text{LCS}(X, \epsilon_1 B_X + \epsilon_2 \mathcal{B}) = \left(\bigsqcup_{g \in G} \{g(Z)\} \right) \bigsqcup \text{NLCS}(X, \epsilon_1 B_X + \epsilon_2 \mathcal{B}),$$

the log pair $(X, \epsilon_1 B_X + \epsilon_2 \mathcal{B})$ is log canonical along $g(Z)$ for every $g \in G$, the equivalence

$$\epsilon_1 B_X + \epsilon_2 \mathcal{B} \sim_{\mathbb{Q}} \epsilon B_X$$

holds, and $\text{NLCS}(X, \epsilon_1 B_X + \epsilon_2 \mathcal{B}) = \text{NLCS}(X, B_X)$.

Proof. See the proofs of [17, Theorem 1.10] and [18, Theorem 1]. □

Suppose that X is a smooth irreducible curve in \mathbb{P}^3 of genus g and degree d .

Theorem 2.14 ([14, Theorem 6.4]). If X is not contained in a hyperplane in \mathbb{P}^3 , then

$$g \leq \begin{cases} \frac{(d-2)^2}{4} & \text{if } d \text{ is even,} \\ \frac{(d-1)(d-3)}{4} & \text{if } d \text{ is odd.} \end{cases}$$

Suppose, in addition, that $G \cong \text{PSL}_2(\mathbb{F}_7)$.

Lemma 2.15. Let Σ be a G -orbit of a point in X . Then $|\Sigma| \in \{24, 42, 56, 84, 168\}$.

Proof. This follows from Lemma B.1, since stabilizer subgroups of all points in X are cyclic. □

Lemma 2.16. Suppose that $g \leq 30$. Then

$$g \in \{3, 8, 10, 15, 17, 19, 22, 24, 29\},$$

and the number of G -orbits in X consisting of 24, 42, 56, 84 points can be described as follows:

genus g	24 points	42 points	56 points	84 points
3	1	0	1	1
8	0	1	1	0
10	1	1	0	1
15	0	2	1	0
15	0	0	1	3
17	1	0	2	0
19	2	0	0	1
22	0	3	0	0
22	0	1	0	3
24	1	1	1	0
29	0	0	2	2

Proof. It follows from the classification of finite subgroups of the group $\mathrm{PGL}_2(\mathbb{C})$ that $g \neq 0$, and it follows from the non-solvability of the group G that $g \neq 1$.

Let $\Gamma \subset G$ be a stabilizer of a point in X . Then $\Gamma \cong \mathbb{Z}_k$ for $k \in \{1, 2, 3, 4, 7\}$ by Lemma B.1. Put $\bar{X} = X/G$. Then \bar{X} is a smooth curve of genus \bar{g} . The Riemann–Hurwitz formula gives

$$2g - 2 = 168(2\bar{g} - 2) + 84a_2 + 112a_3 + 126a_4 + 144a_7,$$

where a_k is the number of G -orbits in X with a stabilizer of a point isomorphic to \mathbb{Z}_k .

Since $a_k \geq 0$, one has $\bar{g} = 0$, and

$$2g - 2 = -336 + 84a_2 + 112a_3 + 126a_4 + 144a_7,$$

which easily implies the required assertions. \square

Let L be a G -invariant line bundle on the curve X (see [8, §1]).

Lemma 2.17. If $\deg(L) \leq 23$ and L is G -linearizable (see [8, §1]), then $h^0(\mathcal{O}_X(L)) \notin \{1, 2, 4, 5\}$.

Proof. If L is G -linearized, then there is a natural linear action of the group G on $H^0(\mathcal{O}_X(L))$, and the required assertion follows from Lemma 2.15 and Appendix B. \square

Theorem 2.18 ([8, Theorem 2.4]). If $g = 3$, then there is $\theta \in \mathrm{Pic}(\mathfrak{C})$ such that $2\theta \sim K_X$ and

$$\mathrm{Pic}^G(X) = \langle \theta \rangle.$$

Thus, if $g = 3$, then $\deg(L)$ is even by Theorem 2.18, because we assume that L is G -invariant.

Lemma 2.19. Suppose that $g = 8$ and L is G -linearizable. Then $7 \mid \deg(L)$.

Proof. Suppose that $7 \nmid \deg(L)$. Then there are integers a and b such that the equality

$$14a + b\deg(L) = 8$$

holds. Put $D = aK_X + bL$. Then $\deg(D) = 8$ and D is a G -linearizable line bundle.

By the Riemann–Roch theorem, the Clifford theorem (see [14, Theorem 5.4]) and Lemma 2.17, we have $h^0(\mathcal{O}_X(D)) = 3$. Then $h^0(\mathcal{O}_X(K_X - D)) = 2$, which contradicts Lemma 2.17. \square

Lemma 2.20. Suppose that $g = 10$ and L is G -linearizable. Then $3 \mid \deg(L)$.

Proof. Suppose that $3 \nmid \deg(L)$. Then there are integers a and b such that the equality

$$18a + b\deg(L) = 10$$

holds. Put $D = aK_X + bL$. Then $\deg(D) = 10$ and D is a G -linearizable line bundle.

It follows from Lemma 2.17 that $h^0(\mathcal{O}_X(K_X - D)) \neq 5$. Then

$$h^0(\mathcal{O}_X(D)) = 3$$

by the Riemann–Roch theorem, the Clifford theorem and Lemma 2.17.

By Lemma 2.15, the linear system $|D|$ is base point free.

Let $\phi: X \rightarrow \mathbb{P}^2$ be a morphism that is given by the linear system $|D|$. Then

$$\deg(\phi(X)) \in \{2, 5, 10\},$$

which is impossible by Lemma B.5 since the action of the group G on \mathbb{P}^2 comes from an irreducible three-dimensional representation of the group G , and the curve $\phi(X)$ is G -invariant. \square

3. PROJECTIVE SPACE

Let ζ be a primitive seventh root of unity, let \hat{G} be a subgroup in $\mathrm{SL}_4(\mathbb{C})$ such that

$$\hat{G} = \left\langle \left(\begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & \zeta & 0 & 0 \\ 0 & 0 & \zeta^4 & 0 \\ 0 & 0 & 0 & \zeta^2 \end{array} \right), \frac{1}{\sqrt{-1}} \left(\begin{array}{cccc} 1 & 2 & 2 & 2 \\ 1 & \zeta + \zeta^6 & \zeta^2 + \zeta^5 & \zeta^3 + \zeta^4 \\ 1 & \zeta^2 + \zeta^5 & \zeta^3 + \zeta^4 & \zeta + \zeta^6 \\ 1 & \zeta^3 + \zeta^4 & \zeta + \zeta^6 & \zeta^2 + \zeta^5 \end{array} \right) \right\rangle,$$

and let us denote by the symbol U_4 the corresponding faithful four-dimensional representation of the group \hat{G} (cf. Appendix B). Then $\hat{G} \cong \mathrm{SL}_2(\mathbb{F}_7)$, and U_4 is irreducible (see [22], [6]).

Let $\phi: \mathrm{SL}_4(\mathbb{C}) \rightarrow \mathrm{Aut}(\mathbb{P}^3)$ be a natural projection. Put $G = \phi(\hat{G})$. Then $G \cong \mathrm{PSL}_2(\mathbb{F}_7)$.

Remark 3.1. It follows from [1, Chapter VII] that $\mathrm{Aut}^G(\mathbb{P}^3) = G$.

Lemma 3.2. Let P be a point in \mathbb{P}^3 , and let Σ be its G -orbit. Suppose that $|\Sigma| \leq 41$. Then

- either $|\Sigma| = 8$ and the orbit Σ is unique,
- or $|\Sigma| = 24$ and the orbit Σ is unique,
- or $|\Sigma| = 28$ and there are exactly two possibilities for the orbit Σ .

Proof. It follows from Corollary B.2 that

$$|\Sigma| \in \{7, 8, 14, 21, 24, 28\},$$

because the representation U_4 is irreducible (so that $|\Sigma| \neq 1$). Let G_P be a stabilizer subgroup in G of the point P , and let \hat{G}_P be the preimage of the subgroup G_P under ϕ . If $|\Sigma| = 21$, then

$$\hat{G}_P \cong 2.D_4,$$

which is impossible by Lemma B.9. If $|\Sigma| \in \{7, 14\}$, then \hat{G}_P has a subgroup isomorphic to $2.A_4$, which is also impossible by Lemma B.9. Thus, we see that $|\Sigma| \in \{8, 24, 28\}$.

Suppose that $|\Sigma| = 8$. Then it follows from Lemmas B.1 and B.7 that

$$G_P \cong \mathbb{Z}_7 \rtimes \mathbb{Z}_3,$$

the orbit Σ does exist, the point P is the unique G_P -invariant point in \mathbb{P}^3 , and Σ is unique, since all subgroups of the group G that are isomorphic to $\mathbb{Z}_7 \rtimes \mathbb{Z}_3$ are conjugate by Lemma B.1.

Suppose that $|\Sigma| = 24$. Then $G_P \cong \mathbb{Z}_7$ and $\hat{G}_P \cong \mathbb{Z}_{14}$. Take any $g \in \hat{G}_P$ such that $\hat{G}_P = \langle g \rangle$, and let R_n be a one-dimensional representation of the group \hat{G}_P such that g acts on R_n by multiplication by $-\zeta^n$. For a suitable choice of g , we have isomorphism of \hat{G}_P -representations

$$U_4 \cong R_0 \oplus R_1 \oplus R_2 \oplus R_4,$$

which implies that \mathbb{P}^3 contains exactly 3 different points besides $P \in \mathbb{P}^3$, say P_1, P_2 and P_3 , that are fixed by the group G_P . There is a unique subgroup $H \subset G$ such that

$$\mathbb{Z}_7 \rtimes \mathbb{Z}_3 \cong H \supset G_P,$$

and we may assume that the point P_1 is H -invariant (that is, corresponds to the subrepresentation R_0). Then its G -orbit consists of eight points. Thus, we see that $\{P, P_2, P_3\}$ is a H -orbit, which implies that the orbit Σ exists and it is unique.

Suppose that $|\Sigma| = 28$. Then $G_P \cong \mathrm{S}_3$. There is a unique point $P' \in \mathbb{P}^3$ such that

$$P \neq P'$$

and P' is fixed by G_P by Lemma B.9. But G contains exactly 28 subgroups isomorphic to S_3 , and it easily follows from Lemma B.9 that no two of these subgroups can fix the same point, which implies that there are exactly two G -orbits in \mathbb{P}^3 consisting of 28 points. \square

Let Σ_8, Σ_{24} and $\Sigma_{28} \neq \Sigma'_{28}$ be G -orbits in \mathbb{P}^3 consisting of 8, 24 and 28 points, respectively. The group \hat{G} naturally acts on $\mathbb{C}[x_1, x_2, x_3, x_4]$. Put

$a = x_2x_3x_4, b = x_2^3x_3 + x_3^3x_4 + x_4^3x_2, c = x_2^2x_3^3 + x_3^2x_4^3 + x_4^2x_2^3, d = a^2 + x_2x_3^5 + x_3x_4^5 + x_4x_2^5,$
and $e = 7ab + x_2^7 + x_3^7 + x_4^7$. Furthermore, put $\Phi_4 = 2x_1^4 + 6ax_1 + b$ and

$$\Phi_6 = 8x_1^6 - 20ax_1^3 - 10bx_1^2 - 10cx_1 - 14a^2 - d,$$

$$\Phi_8 = x_1^8 - 2ax_1^5 + bx_1^4 + 2cx_1^3 + (6a^2 + d)x_1^2 + 2abx_1 + ac,$$

$$\Phi'_8 = x_1^8 + 14ax_1^5 - 7bx_1^4 + 14cx_1^3 - 7dx_1^2 + ex_1,$$

$$\begin{aligned} \Phi_{14} = & 48x_1^{14} + 168ax_1^{11} + 308bx_1^{10} - 1596cx_1^9 + 126(42a^2 + 11d)x_1^8 - 8(37e + 490ab)x_1^7 + \\ & + 196(12ac + 5b^2)x_1^6 + 196(15ad - 13bc)x_1^5 + 14(182c^2 - 86ae - 7bd)x_1^4 + \\ & + 28(11be - 42cd)x_1^3 + 14(21d^2 - 16ce)x_1^2 + 14dex_1 - e^2. \end{aligned}$$

Theorem 3.3 ([22, Theorem 1]). The forms $\Phi_4, \Phi_6, \Phi_8, \Phi'_8$ and Φ_{14} are \hat{G} -invariant.

Remark 3.4. There are no \hat{G} -invariant two-dimensional vector subspaces in $\mathbb{C}[x_1, x_2, x_3, x_4]$ that consist of linear, quadratic, cubic, quartic, quintic or sextic forms (see [8, Appendix 1]).

Let F_i be a surface in \mathbb{P}^3 that is given by the equation

$$\Phi_i(x_1, x_2, x_3, x_4) = 0 \subset \mathbb{P}^3 \cong \text{Proj}\left(\mathbb{C}[x_1, x_2, x_3, x_4]\right),$$

and let F'_8 be a surface in \mathbb{P}^3 that is given by $\Phi'_8(x_1, x_2, x_3, x_4) = 0$.

Theorem 3.5 ([22, Theorem 1]). The following assertions hold:

- there are no G -invariant odd degree surfaces in \mathbb{P}^3 ,
- there are no G -invariant quadric surfaces in \mathbb{P}^3 ,
- the only G -invariant quartic surface in \mathbb{P}^3 is the surface F_4 .

One can check that the surface F_4 is smooth.

Lemma 3.6 (cf. [22, Theorem 1]). The sets $F_4 \cap F_6 \cap F'_8, F_4 \cap F_6 \cap F_{14}, F_4 \cap F'_8 \cap F_{14}$ are finite.

Proof. This follows from explicit computations. \square

There is a G -invariant irreducible smooth curve $C_6 \subset \mathbb{P}^3$ of genus 3 and degree 6 such that

$$\Sigma_{24} = C_6 \cap F_4,$$

and C_6 is an intersection of cubic surfaces in \mathbb{P}^3 (see [12, page 154], [8, Example 2.8]).

Lemma 3.7. Let C be a G -invariant curve in \mathbb{P}^3 such that $\deg(C) \leq 6$. Then $C = C_6$.

Proof. The required assertion must be well-known (cf. [8, §6]). However, we were unable to find an appropriate reference. So, we decided to give the proof here, which is very easy.

By Corollary B.2, we may assume that the curve C is irreducible.

If the curve C is singular, then $|\text{Sing}(C)| \geq 8$ by Lemma 3.2, which easily leads to a contradiction by applying Lemma 2.15 to the normalization of the curve C . Then C is smooth.

Since U_4 is an irreducible representation of the group \hat{G} , the curve C is not contained in a plane in \mathbb{P}^3 . Then C is a curve of genus 3 and degree 6 by Theorem 2.14 and Lemma 2.16.

By Theorem 2.18, there is a unique G -invariant line bundle of degree 6 on the curve C , which implies that the embedding $C \hookrightarrow \mathbb{P}^3$ is unique up to the action of the group $\text{Aut}^G(\mathbb{P}^3)$. But

$$\text{Aut}^G(\mathbb{P}^3) = G$$

by Remark 3.1, which implies that $C = C_6$. \square

Note that $C_6 \cap \Sigma_8 = \emptyset$ by Lemma 2.15.

Lemma 3.8. There is a non-biregular involution $\tau \in \text{Bir}^G(\mathbb{P}^3)$ such that the diagram

$$(3.9) \quad \begin{array}{ccc} & V & \\ \alpha \swarrow & & \searrow \beta \\ \mathbb{P}^3 & \text{---} \tau \text{---} & \mathbb{P}^3 \end{array}$$

commutes and $\langle G, \tau \rangle \cong G \times \mathbb{Z}_2$, where α and β are blow ups of the curve C_6 .

Proof. The existence of (3.9) is well-known (see [8, Remark 6.8]). The isomorphism

$$\langle G, \tau \rangle \cong \text{PSL}_2(\mathbb{F}_7) \times \mathbb{Z}_2$$

follows from the last three lines of the proof of [8, Lemma 6.4]. \square

Let us introduce a G -invariant curve in \mathbb{P}^3 , which has never been mentioned in the literature.

Lemma 3.10. There is a G -invariant irreducible curve $C_{14} \subset \mathbb{P}^3$ of degree 14 such that $\Sigma_8 \subset C_{14}$.

Proof. Let \mathfrak{C} be the genus 3 curve introduced in Example 1.10. Then

$$\text{Aut}(\mathfrak{C}) \cong G,$$

which implies that \mathfrak{C} admits a natural action of the group G . It follows from Theorem 2.18 that

$$\text{Pic}^G(\mathfrak{C}) = \langle \theta \rangle,$$

where θ is a G -invariant line bundle of degree 2. By [8, Lemma 6.4], there exists an isomorphism

$$H^0(\mathcal{O}_{\mathfrak{C}}(7\theta)) \cong U_4 \oplus U_8,$$

where U_8 is an irreducible eight-dimensional representation of the group \hat{G} (see Appendix B).

The linear system $|7\theta|$ gives a G -equivariant embedding $\rho: \mathfrak{C} \hookrightarrow \mathbb{P}^{11}$ such that there exist unique G -invariant linear subspaces Π_3 and Π_7 in \mathbb{P}^{11} of dimensions 3 and 7, respectively. Then

$$\rho(\mathfrak{C}) \cap \Pi_7 = \emptyset$$

by Lemma 2.15. Let $\iota: \mathbb{P}^{11} \dashrightarrow \Pi_3$ be a G -equivariant projection from Π_7 , put $C_{14} = \iota \circ \rho(\mathfrak{C})$, and identify Π_3 with our \mathbb{P}^3 . Then $C_{14} \subset \mathbb{P}^3$ is an irreducible G -invariant curve of degree 14.

Let H be subgroup in G such that $H \cong \mathbb{Z}_7 \rtimes \mathbb{Z}_3$. Then there is a H -invariant subset $\Sigma_3 \subset \mathfrak{C}$ such that $|\Sigma_3| = 3$. Note that Σ_3 is a subset of the G -orbit of length 24, which implies that

$$\Sigma_8 \subset C_{14} \iff |\iota \circ \rho(\Sigma_3)| = 1$$

by Lemma 3.2. Let us show that $\iota \circ \rho(\Sigma_3)$ consists of a single point.

Let T be a vector subspace in $H^0(\mathcal{O}_{\mathfrak{C}}(7\theta))$ that consists of sections vanishing at the subset Σ_3 , and let L_i be a \hat{G} -subrepresentation in $H^0(\mathcal{O}_{\mathfrak{C}}(7\theta))$ such that $L_i \cong U_i$ for $i \in \{4, 8\}$. Then

$$|\iota \circ \rho(\Sigma_3)| = 1 \iff \dim(L_4 \cap T) = 3$$

by the construction of the map $\iota \circ \rho$. Let us show that $\dim(L_4 \cap T) = 3$.

Take a subgroup $\hat{H} \subset \hat{G}$ such that $\phi(\hat{H}) = H$. Then it follows from Lemma B.7 that

$$L_8 \cong V_3 \oplus V_3' \oplus V_1' \oplus V_1''$$

and $L_4 \cong V_3 \oplus V_1$ as representations of the group \hat{H} , where $V_3, V_3', V_1, V_1', V_1''$ are different irreducible representations of dimensions 3, 3, 1, 1, 1, respectively. Thus, there is an isomorphism

$$T \cong V_3 \oplus V_3 \oplus V_3',$$

because T does not contain one-dimensional \hat{H} -subrepresentations, since the curve \mathfrak{C} does not contain H -invariant subsets consisting of $\deg(7\theta) - 3 = 11$ points. Hence $L_4 \cap T \cong V_3$. \square

Let \mathcal{Q} be a linear system of quadric surfaces in \mathbb{P}^3 that pass through Σ_8 . Put

$$\Sigma_8 = \{O_1, O_2, O_3, O_4, O_5, O_6, O_7, O_8\} \subset \mathbb{P}^3.$$

Example 3.11. The log pair $(\mathbb{P}^3, 2\mathcal{Q})$ is canonical. But $\text{NLCS}(\mathbb{P}^3, 4\mathcal{Q}) = \Sigma_8$.

Let $\pi: U \rightarrow \mathbb{P}^3$ be the blow up of the subset Σ_8 , let E_i be the exceptional divisor of the birational morphism π such that $\pi(E_i) = O_i$ for every i . Then there is a commutative diagram

$$(3.12) \quad \begin{array}{ccc} & U & \\ \pi \swarrow & & \searrow \eta \\ \mathbb{P}^3 & \dashrightarrow & \mathbb{P}^2, \\ & \psi & \end{array}$$

where ψ is a rational map that is given by \mathcal{Q} , and η is an elliptic fibration.

Lemma 3.13. The following assertions hold:

- the curve C_{14} has an ordinary triple point at every point of the set Σ_8 ,
- the proper transform of the curve C_{14} on the threefold U is smooth,
- the curve C_{14} is smooth outside of the points of the set Σ_8 ,
- the map $\psi: \mathbb{P}^3 \dashrightarrow \mathbb{P}^2$ induces a birational map $C_{14} \dashrightarrow \psi(C_{14})$,
- the curve $\psi(C_{14})$ is a smooth curve of genus 3 and degree 4,
- the intersection $C_6 \cap C_{14}$ is empty.

Proof. Let \bar{C}_{14} and \bar{Q} be the proper transforms of the curve C_{14} and a general surface in \mathcal{Q} on the threefold U , respectively. Then $\text{mult}_{O_i}(C_{14}) \geq |\bar{C}_{14} \cap E_i| \geq 3$ by Lemma 2.11. Thus

$$4 \geq 28 - 8|\bar{C}_{14} \cap E_i| \geq 2\text{deg}(C_{14}) - \sum_{i=1}^8 E_i \cdot \bar{C}_{14} = \left(\pi^*(H) - \sum_{i=1}^8 E_i\right) \bar{C}_{14} = \bar{Q} \cdot \bar{C}_{14} \geq 0,$$

which implies that \bar{C}_{14} has an ordinary triple point at every point of the set Σ_8 . Moreover,

$$4 = 2\text{deg}(C_{14}) - 24 = \left(\pi^*(H) - \sum_{i=1}^8 E_i\right) \bar{C}_{14} = \bar{Q} \cdot \bar{C}_{14},$$

which implies that $\eta(\bar{C}_{14})$ is a smooth curve of genus 3 and degree 4 by Lemma B.5. Therefore

$$\bar{C}_{14} \cong \eta(\bar{C}_{14}),$$

which implies that C_{14} is smooth outside of the points of the set Σ_8 .

Let us show that $C_6 \cap C_{14} = \emptyset$. Suppose that $C_6 \cap C_{14} \neq \emptyset$. Then

$$|C_6 \cap C_{14}| \geq 56$$

by Lemma 2.16, since $\Sigma_8 \not\subset C_6$. Let S be a general cubic surface in \mathbb{P}^3 such that $C_6 \subset S$. Then

$$42 = S \cdot C_{14} \geq \sum_{O \in C_6 \cap C_{14}} \text{mult}_O(S) \text{mult}_O(C_{14}) \geq 56,$$

because $C_{14} \not\subset S$. Thus, the intersection $C_6 \cap C_{14}$ is empty. \square

Lemma 3.14. Let C be a G -invariant curve such that $\text{deg}(C) \leq 15$ and $\Sigma_8 \subset C$. Then $C = C_{14}$.

Proof. Let \bar{C} and \bar{Q} be the proper transforms of the curve C and a general surface in \mathcal{Q} on the threefold U , respectively. Then $\text{mult}_{O_i}(C) \geq |\bar{C} \cap E_i| \geq 3$ by Lemma 2.11. Then

$$6 \geq 30 - 8|\bar{C} \cap E_i| \geq 2\text{deg}(C) - \sum_{i=1}^8 E_i \cdot \bar{C} = \left(\pi^*(H) - \sum_{i=1}^8 E_i\right) \bar{C} = \bar{Q} \cdot \bar{C} \geq 0,$$

which implies that $E_i \cdot \bar{C} = |\bar{C} \cap E_i| = 3$.

We may assume that C is a G -orbit of an irreducible curve $\Gamma \subset \mathbb{P}^3$. Then

$$6 \geq 2\deg(C) - 24 = \left(\pi^*(H) - \sum_{i=1}^8 E_i\right)\bar{C} = \bar{Q} \cdot \bar{C} = \delta \deg(\eta(\bar{C}))$$

for some positive integer δ . We may assume that $\delta = 1$ if \bar{C} is contracted by η . Thus, we have

$$\deg(C) \in \{12, 13, 14, 15\},$$

which implies that either $\deg(\Gamma) \leq 2$ or $\Gamma = C$ (see Corollary B.2).

By Lemma B.5, we have $\deg(\psi(C)) \notin \{1, 2, 3\}$. Then $\deg(C) \in \{12, 14, 15\}$ and $\delta = 1$. But

$$(3.15) \quad 6 \geq 2\deg(C) - 24 = \deg(\eta(\bar{C})),$$

which implies that $\deg(\eta(\bar{C})) \in \{0, 4, 6\}$ by Lemma B.5.

Recall that G -invariant quartic curve and sextic curve in \mathbb{P}^2 are irreducible (see Lemma B.5), which easily implies using (3.15) that either $\deg(\Gamma) = 1$ and $\deg(C) = 12$, or $\Gamma = C$.

If $\deg(\Gamma) = 1$ and $\deg(C) = 12$, then it follows from (3.15) that

$$|\eta(\bar{C})| \leq 12,$$

which is impossible, since G -orbit of every point in \mathbb{P}^2 consists of at least 21 points by Lemma B.6.

We see that $\Gamma = C$. Then \bar{C} is not contracted by η , since there is no G -invariant point in \mathbb{P}^2 .

Suppose that $\deg(C) = 15$. Then $\bar{C} \cong \psi(C)$ and $\deg(\psi(C)) = 6$, which immediately implies that $\psi(C)$ is a smooth curve of genus 10 by Lemma B.5. Then there is a natural monomorphism

$$U_4 \cong H^0\left(\mathcal{O}_U(\pi^*(H))\right) \hookrightarrow H^0\left(\mathcal{O}_{\bar{C}} \otimes \mathcal{O}_U(\pi^*(H))\right) \cong \mathbb{C}^6,$$

which contradicts Lemma 2.15, since $\mathrm{SL}_2(\mathbb{F}_7)$ has no irreducible two-dimensional representations.

We see that $\deg(C) = 14$. Thus $\bar{C} \cong \psi(C)$, which implies that \bar{C} is a smooth curve of genus 3.

Arguing as in the proof of Lemma 3.10, we see that $C = \sigma(C_{14})$ for some $\sigma \in \mathrm{Aut}^G(\mathbb{P}^3)$. But

$$\mathrm{Aut}^G(\mathbb{P}^3) = G$$

by Remark 3.1. Hence, we must have $C = C_{14}$, since the curve C_{14} is G -invariant. \square

Lemma 3.16. Let \mathcal{D} be the linear system of quintic surfaces containing C_{14} . Then

- the linear system \mathcal{D} is not empty and does not have fixed components,
- a general surface in \mathcal{D} has a double point in every point of the set Σ_8 ,
- all curves contained in the base locus of the linear system \mathcal{D} are disjoint from C_{14} ,
- general surface in \mathcal{D} is smooth in a general point of the curve C_{14} ,
- any two general surfaces in \mathcal{D} are not tangent to each other along the curve C_{14} .

Proof. Let \bar{C}_{14} be the proper transform of the curve C_{14} on the threefold U , and let \mathcal{I} be the ideal sheaf of the curve \bar{C}_{14} . Put $R = \pi^*(5H) - \sum_{i=1}^8 E_i$. Then

$$h^0\left(\mathcal{O}_U(R) \otimes \mathcal{I}\right) \geq h^0\left(\mathcal{O}_U(R)\right) - h^0\left(\mathcal{O}_{\bar{C}_{14}} \otimes \mathcal{O}_U(R)\right) = 48 - h^0\left(\mathcal{O}_{\bar{C}_{14}} \otimes \mathcal{O}_U(R)\right) = 4,$$

which implies that $\dim(\mathcal{D}) \geq 3$.

Let D be a general surface in \mathcal{D} . Then D is irreducible, because the linear system \mathcal{D} does not have fixed components by Theorem 3.5 and \mathcal{D} is not composed from a pencil by Remark 3.4.

Let \bar{D} be its proper transform of the surface D on the threefold U , and let Γ be a general fiber of the elliptic fibration η . Then

$$20 - 8\mathrm{mult}_{O_1}(\mathcal{D}) = \left(\pi^*(5H) - \sum_{i=1}^8 \mathrm{mult}_{O_i}(\mathcal{D})E_i\right) \cdot \Gamma = \bar{D} \cdot \Gamma \geq 0$$

which implies that $\text{mult}_{O_1}(\mathcal{D}) \leq 2$. Thus $\text{mult}_{O_1}(\mathcal{D}) = 2$ by Lemma 2.11.

Let D' be a general surface in \mathcal{D} such that $D \neq D'$. Hence there is $\mu \in \mathbb{Z}_{>0}$ such that

$$D \cdot D' = \mu C_{14} + B + Z,$$

where Z is a curve not contained in the base locus of the linear system \mathcal{D} , and B is a curve contained in the base locus of the linear system \mathcal{D} such that $C_{14} \not\subseteq \text{Supp}(B)$. Then

$$25 = \deg(D \cdot D') = \mu \deg(C_{14}) + \deg(B) + \deg(Z) = 14\mu + \deg(B) + \deg(Z) \geq 20 + \deg(Z)$$

by Lemma 3.7. Thus, we see that $\mu = 1$ and $\deg(B) + \deg(Z) \leq 11$, which implies, in particular, that the surfaces D and D' are not tangent to each other along C_{14} , since $\mu = 1$.

Note that B is G -invariant, because \mathcal{D} is G -invariant. Therefore $\deg(B) \geq 6$ by Lemma 3.7.

If the base locus of the linear system \mathcal{D} consists of the curve C_{14} , then C_{14} is a scheme-theoretic intersection of surfaces in \mathcal{D} outside some finite subset of the curve C_{14} , because $\mu = 1$.

Let \bar{Z} and \bar{B} be the proper transforms of the curves Z and B on the threefold U , respectively, and let \bar{Q} be a proper transform of a general quadric surface in \mathcal{Q} on the threefold U . Then

$$0 \leq \bar{Q} \cdot \bar{B} = 2\deg(B) - \sum_{i=1}^8 E_i \cdot \bar{B} \leq 22 - 8 \left| \text{Supp}(\bar{B}) \cap E_i \right| \leq 22 - 8 \times \begin{cases} 3 & \text{if } O_i \in \text{Supp}(B), \\ 0 & \text{if } O_i \notin \text{Supp}(B), \end{cases}$$

which implies that $\Sigma_8 \cap \text{Supp}(B) = \emptyset$. But

$$3 + \text{mult}_{O_i}(Z) = \text{mult}_{O_i}(C_{14}) + \text{mult}_{O_i}(B) + \text{mult}_{O_i}(Z) = \text{mult}_{O_i}(D_1 \cdot D_2) \geq 4,$$

which implies that $\Sigma_8 \subset \text{Supp}(Z)$. Hence

$$2\deg(Z) - 8 \geq 2\deg(Z) - \sum_{i=1}^8 E_i \cdot \bar{Z} = \bar{Q} \cdot \bar{Z} \geq 0,$$

which implies that $\deg(Z) \geq 4$, and $\deg(Z) = 4$ if and only if \bar{Z} is contracted by η . But

$$10 \leq 6 + \deg(Z) \leq \deg(B) + \deg(Z) \leq 11,$$

which implies that either $\deg(B) = 7$ and $\deg(Z) = 4$, or $\deg(B) = 6$ and $\deg(Z) = 5$.

If $\deg(Z) = 4$, then \bar{D} is contracted by the morphism η to a curve of degree d , then

$$\bar{D} \sim d \left(\pi^*(2H) - \sum_{i=1}^8 E_i \right),$$

which is a contradiction. Thus, we see that $\deg(B) = 6$ and $\deg(Z) = 5$.

By Lemma 3.7, we have $B = C_6$. Therefore $C_6 \cap C_{14} = \emptyset$ by Lemma 3.13. \square

Let us study some properties of the subset $\Sigma_{28} \subset \mathbb{P}^3$, which also hold for Σ'_{28} .

Lemma 3.17. Let Z be a G -invariant curve in \mathbb{P}^3 such that $\Sigma_{28} \subset Z$. Then

$$\deg(Z) \geq 16$$

if the set Σ_{28} imposes independent linear conditions on quartic surfaces in \mathbb{P}^3 .

Proof. Suppose that the set Σ_{28} imposes independent linear conditions on quartic surfaces in \mathbb{P}^3 , and suppose that $\deg(Z) \leq 15$. Let us derive a contradiction.

Without loss of generality, we may assume that Z is a G -orbit of an irreducible curve Z_1 .

Put $Z = \sum_{i=1}^r Z_i$, where Z_i is an irreducible curve in \mathbb{P}^3 and $r \in \mathbb{Z}_{>0}$. Then

$$\deg(Z) = \sum_{i=1}^r \deg(Z_i) = r \deg(Z_1) \leq 15,$$

which implies that $r \in \{1, 7, 8, 14\}$ by Corollary B.2. Thus, if $r \neq 1$, then $\deg(Z_i) \in \{1, 2\}$.

If $\deg(Z_1) = 2$ and $r \neq 1$, then $r = 7$, which contradicts Lemma 3.2. Hence Z_1 is a line if $r \neq 1$. Let Γ be the stabilizer subgroup in G of the curve Z_1 . If $r = 7$, then $\Gamma \cong S_4$, which implies that

$$|Z_1 \cap \Sigma_{28}| \geq 6,$$

which is impossible, since Σ_{28} impose independent linear conditions on quartic surfaces.

Let $\hat{\Gamma}$ be the smallest subgroup of the group \hat{G} such that $\phi(\hat{\Gamma}) = \Gamma$. Then

- if $r = 8$, then $\hat{\Gamma} \cong 2 \cdot (\mathbb{Z}_7 \rtimes \mathbb{Z}_3)$, which contradicts Lemma B.7.
- if $r = 14$, then $\hat{\Gamma} \cong 2 \cdot A_4$, which contradicts Lemma B.9.

Thus, we see that the curve Z is irreducible.

By Lemma 2.15, the curve Z must be singular at every point of the set Σ_{28} , which implies that $Z \neq C_{14}$ by Lemma 3.13. Hence $\Sigma_8 \not\subset Z$ by Lemma 3.14.

Let A be a point in Σ_{28} . Then there exists a quartic surface $S \subset \mathbb{P}^3$ such that S contain all points of the set $\Sigma_{28} \setminus \{A\}$ and the surface S does not contain the point A . Therefore

$$4\deg(Z) = S \cdot Z \geq \sum_{O \in \Sigma_{28} \setminus \{A\}} \text{mult}_O(Z) \geq \sum_{O \in \Sigma_{28} \setminus \{A\}} 2 \geq 54,$$

which implies that $\deg(Z) \geq 14$. Thus, either $\deg(Z) = 14$ or $\deg(Z) = 15$.

If $\deg(Z) = 15$, then $Z \subset F_4 \cap F_6 \cap F'_8$ by Lemma 2.15, since 60, 90 and 120 are not equal to

$$24n_1 + 42n_2 + 56n_3 + 84n_4 + 168n_5$$

for any non-negative integers n_1, n_2, n_3, n_4 and n_5 . Then $\deg(Z) = 14$ by Lemma 3.6.

Let \bar{Z} be the normalization of the curve Z , and let g be the genus of the curve \bar{Z} . Then

$$(3.18) \quad g \leq 66 - |\text{Sing}(Z)| \leq 38,$$

because the projection from a general points of the curve Z gives us a birational isomorphism between Z and a plane curve of degree 13 with at least $|\text{Sing}(Z)|$ singular points.

Note that G naturally acts on both curves Z and \bar{Z} . Let us show that $Z \setminus \Sigma_{28}$ is smooth.

Suppose that $\text{Sing}(Z) \neq \Sigma_{28}$. Let Λ be a G -orbit of a point in $\text{Sing}(Z) \setminus \Sigma_{28}$. Then

$$|\Lambda| \leq 66 - |\Sigma_{28}| = 38,$$

by (3.18), and $|\Lambda| \in \{24, 28\}$ by Lemma 3.2. If $\Lambda = C_6 \cap F_4$, then

$$42 = S \cdot Z \geq \sum_{O \in \Lambda} \text{mult}_O(S) \text{mult}_O(Z) \geq 48,$$

where S is a general cubic surface such that $C_6 \subset S$. Thus, we see that $|\Lambda| \neq 24$ by Lemma 3.2, which implies that $|\Lambda| = 28$. Therefore, it follows from (3.18) and Lemma 2.16 that

$$g \in \{3, 8, 10\},$$

and the points of the set $\Lambda \cup \Sigma_{28}$ must be singular points of the curve Z of multiplicity two, which implies that \bar{Z} has at least two G -orbits consisting of 56 points, which contradicts Lemma 2.16.

Thus, we see that Z is smooth outside of the set Σ_{28} .

Let B be a sufficiently general point of the curve Z , let \mathcal{M} be a linear system consisting of all quartic surfaces in \mathbb{P}^3 that contain the set $\Sigma_{28} \cup B$. Then $\dim(\mathcal{M}) \geq 35 - 29 = 6$.

Let M be a general surface in \mathcal{M} . If $Z \not\subset M$, then

$$56 = M \cdot Z \geq \text{mult}_B(M) \text{mult}_B(Z) + \sum_{O \in \Sigma_{28}} \text{mult}_O(M) \text{mult}_O(Z) \geq 57,$$

which is a contradiction. Thus, the curve Z is contained in the base locus of the linear system \mathcal{M} , which implies that the linear system \mathcal{M} is G -invariant, because Z is G -invariant.

The linear system \mathcal{M} does not have fixed components by Theorem 3.5 and \mathcal{M} is not composed from a pencil by Remark 3.4. This implies, in particular, that the surface M is irreducible.

Let M' be another general surface in the linear system \mathcal{M} . Put

$$M \cdot M' = \mu Z + \Upsilon,$$

where $\mu \in \mathbb{Z}_{>0}$, and Υ is an effective one-cycle such that $Z \not\subset \text{Supp}(\Upsilon)$. Then

$$16 = \deg(M \cdot M') = 14\mu + \deg(\Upsilon),$$

which implies that $\mu = 1$, and the base locus of the linear system \mathcal{M} contain no curves except Z , because there are no G -invariant lines or conics in \mathbb{P}^3 . Thus $\text{mult}_Z(\mathcal{M}) = 1$ and $\deg(\Upsilon) = 2$, which implies, in particular, that the surface M is uniruled.

Let M'' be another general surface in the linear system \mathcal{M} . Then

$$M \cap M' \cap M'' = C + \Upsilon \cap M'',$$

which immediately implies that the base locus of the linear system \mathcal{M} consists of at most 8 points outside of the curve Z . Thus, it follows from Lemma 3.2 that

- either the base locus of the linear system \mathcal{M} consists of the curve Z ,
- or the base locus of the linear system \mathcal{M} consists of the curve Z and the set Σ_8 .

Let Q be a general surface in the linear system \mathcal{Q} . If $\Sigma_8 \subset M \cap M'$, then

$$32 = Q \cdot M \cdot M' = Q \cdot Z + \sum_{O \in \Sigma_8} \text{mult}_O(Q) \text{mult}_O(\Upsilon) = 28 + \sum_{O \in \Sigma_8} \text{mult}_O(Q) \text{mult}_O(\Upsilon) \geq 36,$$

which implies that $\Sigma_8 \not\subset M \cap M'$. Hence Z is the base locus of the linear system \mathcal{M} .

By adjunction formula, we see that the singularities of the surface M are not canonical, which implies that $(\mathbb{P}^3, \mathcal{M})$ is not canonical by [19, Theorems 4.8.1 and 7.9] (cf. [19, Theorem 7.5]). But

$$Z \notin \text{NCS}(\mathbb{P}^3, \mathcal{M}),$$

because $\text{mult}_Z(\mathcal{M}) = 1$. Thus, the set $\text{NCS}(\mathbb{P}^3, \mathcal{M})$ contains a point $P \in Z$. Thus

$$\text{mult}_P(\Upsilon) \geq 4 - \text{mult}_P(Z) = \begin{cases} 2 & \text{if } P \in \Sigma_{28}, \\ 3 & \text{if } P \notin \Sigma_{28}, \end{cases}$$

since $\text{mult}_P(\mathcal{M}) \geq 2$. Let Θ be the G -orbit of the point P . Then $|\Theta| \geq 28$ by Lemma 3.2, and

$$8 = M \cdot M' \cdot M'' - M'' \cdot Z = M'' \cdot \Upsilon \geq \sum_{O \in \Theta} \text{mult}_O(M'') \text{mult}_O(\Upsilon) \geq 56,$$

which is a contradiction. □

Let \mathcal{R} be a linear system consisting of all quartic surfaces in \mathbb{P}^3 that pass through Σ_{28} .

Lemma 3.19. If the set Σ_{28} imposes independent linear conditions on quartic surfaces in \mathbb{P}^3 , then every curve in the base locus of the linear system \mathcal{R} contains no points of the set Σ_{28} .

Proof. Let C be an irreducible curve in \mathbb{P}^3 such that C is contained in the base locus of the linear system \mathcal{R} , let Z be the G -orbit of the curve C , let R_1 and R_2 be general surfaces in \mathcal{R} . Put

$$R_1 \cdot R_2 = \mu Z + \Upsilon,$$

where $\mu \in \mathbb{Z}_{>0}$, and Υ is an effective one-cycle such that $C \not\subset \text{Supp}(\Upsilon)$. Then

$$16 = \deg(R_1 \cdot R_2) = \mu \deg(Z) + \deg(\Upsilon),$$

which implies that $\deg(Z) \leq 15$ by Remark 3.4.

It follows from Lemma 3.17 that $\Sigma_{28} \not\subset Z$ and $C \cap \Sigma_{28} = \emptyset$ if the points of the set Σ_{28} impose independent linear conditions on quartic surfaces in \mathbb{P}^3 . □

We do not know whether or not Σ_{28} imposes independent linear conditions on quartic surfaces.

4. COMPACTIFIED MODULI SPACE OF $(1, 7)$ -POLARIZED ABELIAN SURFACES

Put $\epsilon = -1/2 + \sqrt{-7}/2$. Let \mathfrak{C} be a curve in \mathbb{P}^2 that is given by the equation

$$(4.1) \quad x^4 + y^4 + z^4 + 3\epsilon(x^2y^2 + x^2z^2 + y^2z^2) = 0 \subset \mathbb{P}^2 \cong \text{Proj}(\mathbb{C}[x, y, z]),$$

which implies that \mathfrak{C} is isomorphic to the curve described in Example 1.10 (see [21, page 55]).

Remark 4.2. The Hessian curve of the curve \mathfrak{C} is a smooth sextic curve (see [10, Example 6.1.1]).

Remark 4.3. The quartic \mathfrak{C} is not degenerate (see [10, Definition 2.8]).

Put $\bar{\epsilon} = -1/2 - \sqrt{-7}/2$. Let $\psi: \text{SL}_3(\mathbb{C}) \rightarrow \text{Aut}(\mathbb{P}^2) \cong \text{PGL}_3(\mathbb{C})$ be a natural projection. Put

$$A = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad D = \frac{1}{2} \begin{pmatrix} -1 & 1 & \bar{\epsilon} \\ \epsilon & \epsilon & 0 \\ -1 & 1 & -\bar{\epsilon} \end{pmatrix},$$

put $G = \psi(\langle A, B, C, D \rangle)$, put $X = \text{VSP}(\mathfrak{C}, 6)$ (see Definition A.3). By [21, page 55], we have

$$\text{Aut}(\mathfrak{C}) \cong G \cong \langle A, B, C, D \rangle \cong \text{PSL}_2(\mathbb{F}_7).$$

Theorem 4.4 ([23, Theorem 4.4]). The threefold X is a smooth Fano threefold such that

$$\text{Pic}(X) = \mathbb{Z}[-K_X]$$

and $(-K_X)^3 = 22$.

The action of the group G on the plane \mathbb{P}^2 induces its natural action on the threefold X that turns the vector space $H^0(\mathcal{O}_X(-K_X))$ into a 14-dimensional representation of the group G .

Theorem 4.5. In the notation of Appendix B, one has $H^0(\mathcal{O}_X(-K_X)) \cong I \oplus W_6 \oplus W_7$.

Proof. We may assume that $W \cong W_3$. By Theorem A.6, one has

$$U_7 \cong \text{Sym}^3(W_3^\vee) / W_3 \cong W_7 \cong U_7^\vee,$$

where all isomorphisms are isomorphisms of G -representations. Hence, we have

$$U_{14} \cong \Lambda^3(U_7) / (W_3^\vee \otimes U_7^\vee) \cong \Lambda^3(W_7) / (W_3^\vee \otimes W_7) \cong U_{14}^\vee,$$

which implies the required assertion by Corollary B.4. \square

Corollary 4.6. There is a unique G -invariant surface in $| -K_X |$.

Let us identify X with its anticanonical image in \mathbb{P}^{13} (see Appendix A).

Lemma 4.7. Let $Q \subset X$ be a surface swept out by the lines contained in X . Then

$$Q \sim -2K_X,$$

the surface Q is irreducible, and the surface Q is birational to $\mathfrak{C} \times \mathbb{P}^1$.

Proof. By [30], the surface Q is reduced and $Q \sim -2K_X$. By [32, Theorem 6.1], it is birational to

$$\mathfrak{C}' \times \mathbb{P}^1,$$

where \mathfrak{C}' is the image of the curve \mathfrak{C} under the Scorza map. But $\mathfrak{C}' \cong \mathfrak{C}$, because \mathfrak{C} is the only curve of genus 3 that admits a faithful action of the group $\text{PSL}_2(\mathbb{F}_7)$. \square

In the remaining part of this section we are going to prove the following result.

Theorem 4.8. Let Σ be a G -orbit of a point in X such that $|\Sigma| \leq 20$. Then

$$|\Sigma| \in \{8, 14\},$$

and if $|\Sigma| = 8$, then Σ is unique.

Remark 4.9. There exist finitely many points in X whose G -orbits consists of 14 points.

Before proving Theorem 4.8, let us use Theorem 4.8 to prove the following result.

Lemma 4.10. Let F be the unique G -invariant surface in $|-K_X|$. Then F is smooth.

Proof. The minimal resolution of the surface F admits a faithful action of the group $\mathrm{PSL}_2(\mathbb{F}_7)$, which implies that F is smooth by [29, Claim 2.1] if it has at most canonical singularities.

Let us show that F has at most canonical singularities (Du Val singularities).

It is well-known that F has canonical singularities if and only if the surface F has rational singularities (see [19, Theorem 1.11]). By Theorem 1.15, the log pair (X, F) is log canonical.

Suppose that F has worse than canonical singularities. Let us derive a contradiction.

Let S be a minimal center in $\mathrm{LCS}(X, F)$. Then

$$\dim(S) \in \{0, 1\}$$

by Theorem 2.10. Let Z be the G -orbit of the subvariety S . Then $Z \subset F$.

Choose any $\epsilon \in \mathbb{Q}$ such that $2 \gg \epsilon > 1$. Then arguing as in the proof of [17, Theorem 1.10], we can find a G -invariant \mathbb{Q} -divisor D such that

$$D \sim_{\mathbb{Q}} \epsilon F \sim_{\mathbb{Q}} -\epsilon K_X,$$

the log pair (X, D) is log canonical, and every minimal center in $\mathrm{LCS}(X, D)$ is an irreducible component of the subvariety Z (see Lemma 2.13).

Let H be a sufficiently general hyperplane section of the threefold X , and let \mathcal{I}_Z be the ideal sheaf of the subvariety Z . Then there is an exact sequence of G -representations

$$0 \rightarrow H^0(\mathcal{O}_X(H) \otimes \mathcal{I}_Z) \rightarrow H^0(\mathcal{O}_X(H)) \rightarrow H^0(\mathcal{O}_Z \otimes \mathcal{O}_X(H)) \rightarrow 0$$

by Theorem 2.2. Put $q = h^0(\mathcal{O}_X(H) \otimes \mathcal{I}_Z)$. One has

$$(4.11) \quad h^0(\mathcal{O}_Z \otimes \mathcal{O}_X(H)) = 14 - q \in \{6, 7, 13\},$$

because $q \in \{1, 7, 8\}$ by Theorem 4.5, since $Z \subset F$. Hence S is not a point by Theorem 4.8.

We see that S is a curve. Moreover, S is a smooth curve of genus g such that

$$-K_X \cdot S = \deg(S) \geq 2g - 1$$

by Theorem 2.10. By Remark 2.12, the curve Z is a disjoint union of smooth irreducible curves.

Let r be the number of connected components of the curve Z . Put $d = \deg(S)$. Then

$$(4.12) \quad g \leq d - g + 1 \leq r(d - g + 1) = 14 - q \in \{6, 7, 13\}$$

by (4.11), which implies that $d \leq 13 + g \leq 27$.

Let us show that $r = 1$.

Suppose that $r \geq 2$. Then $r \geq 7$ and $r \neq 13$ by Corollary B.2 Thus $r = 7$ and

$$d - g + 1 = 1$$

by (4.12), which implies that $d = g$. But $d \geq 2g - 1$, so that $d = g = 1$, which is absurd.

We see that $r = 1$. There is a natural faithful action of the group G on the curve S . But

$$d = 13 + g - q \geq 2g - 1,$$

by (4.12), which implies that $g \leq 14 - q$. Hence $g \in \{3, 8, 10\}$ by Lemma 2.16.

Let us show that $g = 3$.

Suppose that $g \neq 3$. Then $g \in \{8, 10\}$. It follows from (4.12) that

$$q \leq 14 - g \leq 6,$$

which implies that $q = 1$, because $q \in \{1, 7, 8\}$. Hence, it follows from (4.12) that $d \in \{20, 22\}$, which is impossible by Lemmas 2.19 and 2.20.

Thus, we have $g = 3$, so that $d = 16 - q$ by (4.12), where $q \in \{1, 7, 8\}$.

By Theorem 2.18, there is a G -invariant line bundle $\theta \in \text{Pic}(S)$ of degree 2 such that

$$\text{Pic}^G(S) = \langle \theta \rangle,$$

which implies that $q = 8$ and $H|_S \sim 2K_S$, because $d = 16 - q$. Thus $d = 8$.

Let Q be a surface in X that is spanned by lines. Then Q is G -invariant. If $S \not\subset Q$, then

$$|Q \cap S| \leq 2\deg(S) = 16$$

which is impossible by Lemma 2.15. Thus, we see that $S \subset Q \cap F$.

By Lemma 4.7, the surface Q is irreducible, $Q \sim -2K_X$ and Q is birational to $\mathfrak{C} \times \mathbb{P}^1$.

Let P be a general point in S . There exists a line $L \subset X$ such that $P \in L$. If $L \not\subset F$, then

$$1 = \deg(L) = F \cdot L \geq \text{mult}_P(F)\text{mult}_P(L) \geq \text{mult}_S(F) \geq 2,$$

which simply means $L \subset F$. Since $g \neq 0$, the generality of the point $P \in S$ implies that the surface F is swept out by lines, which is impossible since $Q \not\subset F$. \square

Remark 4.13. Let Σ be a G -invariant subset of the threefold X such that we have $|\Sigma| \in \{8, 14\}$, and let F be the unique G -invariant surface in $|-K_X|$. Then it follows from [36] that

$$\Sigma \cap F = \emptyset,$$

because F is smooth by Lemma 4.10. This is easy to prove, since G acts symplectically on F .

Now we are going to prove Theorem 4.8. Put

$$\square = X \setminus \left\{ \Gamma \in \text{Hilb}_6(\check{\mathbb{P}}^2) \mid \Gamma \text{ is polar to the curve } C \right\} \subset \text{Hilb}_6(\check{\mathbb{P}}^2).$$

Lemma 4.14. Let Σ be a G -orbit in \square such that $|\Sigma| \leq 20$. Then $|\Sigma| = 8$ and Σ is unique.

Proof. It follows from [25, § 2.3] that there exists an effective one-cycle

$$m_1 L_1 + \dots + m_r L_r \sim \mathcal{O}_{\mathbb{P}^2}(6)$$

on \mathbb{P}^2 that corresponds to a point in Σ , where L_i are lines in \mathbb{P}^2 and m_i are positive integers.

Without loss of generality, we may assume that $m_1 \geq \dots \geq m_r$. Then

$$(m_1, \dots, m_r) \in \left\{ (2, 1, 1, 1, 1), (3, 1, 1, 1), (2, 2, 1, 1), (2, 2, 2), (4, 2) \right\}$$

by [25, Theorem 1.1] (cf. [24, Theorem A]). Thus, it follows from $|\Sigma| \leq 20$ that

$$(m_1, \dots, m_r) = (2, 2, 2),$$

because the smallest G -invariant subset in \mathbb{P}^2 has at least 21 points by Lemma B.6.

Let us consider the lines L_1, L_2, L_3 as points in the dual projective plane $\check{\mathbb{P}}^2$ with a natural action of the group G , and let $\hat{\mathfrak{C}}$ be the unique G -invariant quartic curve in $\check{\mathbb{P}}^2$. Then

$$\hat{\mathfrak{C}} \cap \{L_1, L_2, L_3\} \neq \emptyset$$

by [25, Proposition 3.13], which easily implies that $|\Sigma| = 8$ and Σ is unique by Lemma 2.16. \square

Therefore, to prove Theorem 4.8 it is enough to consider only G -orbits of points in $X \setminus \square$, which are points in X that can be represented by polar hexagons to the curve \mathfrak{C} .

Suppose that the assertion of Theorem 4.8 is false. Let us derive a contradiction.

By Lemmas 4.14 and B.1, there are six lines L_1, \dots, L_6 on \mathbb{P}^2 such that

$$(4.15) \quad F(x, y, z) = L_1^4(x, y, z) + L_2^4(x, y, z) + L_3^4(x, y, z) + L_4^4(x, y, z) + L_5^4(x, y, z) + L_6^4(x, y, z),$$

where $L_i(x, y, z)$ is a linear form such that L_i is given by $L_i(x, y, z) = 0$, and the stabilizer subgroup of the hexagon $\sum_{i=1}^6 L_i$ in the group G is isomorphic either to $\mathbb{Z}_7 \times \mathbb{Z}_3$ or to S_4 .

Remark 4.16. The lines L_1, \dots, L_6 are different by [10, Theorem 6.12.2] and Remark 4.2.

Lemma 4.17. Let $g \in G$ be an element of order 7. Then $\sum_{i=1}^6 L_i$ is not g -invariant.

Proof. Suppose that $\sum_{i=1}^6 L_i$ is g -invariant. Then

$$g(L_i) = L_i$$

for every $i \in \{1, \dots, 6\}$. Take $M \in \mathrm{SL}_3(\mathbb{C})$ such that $\psi(M) = g^{-1}$ and $M^7 = 1$. Then

$$L_i\left((x, y, z)M\right) = \lambda_i L_i(x, y, z)$$

for some $\lambda_i \in \mathbb{C}$ for every $i \in \{1, \dots, 6\}$, i. e. $\lambda_1, \dots, \lambda_6$ are eigenvalues of the matrix M .

Since g has order 7, we have $\lambda_1^7 = \dots = \lambda_6^7 = 1$. We may assume that $\lambda_1 = \lambda_2 = \lambda_3 = \lambda_4 = \lambda_5$, since L_1, \dots, L_6 are different lines. Then $\lambda_6 \neq \lambda_1$ by (4.15). Thus

$$|L_1 \cap L_2 \cap L_3 \cap L_4 \cap L_5| = 1$$

and $L_1 \cap L_2 \cap L_3 \cap L_4 \cap L_5 \notin L_6$. Therefore

$$\sum_{i=1}^6 L_i^4(x, y, z) = F\left((x, y, z)M\right) = \lambda_6^4 L_6^4(x, y, z) + \lambda_1^4 \sum_{i=1}^5 L_i^4(x, y, z),$$

by (4.15), which easily implies that $\lambda_1^4 = \lambda_6^4 = 1$. Hence $\lambda_1 = \lambda_6 = 1$, which is a contradiction. \square

Let Γ be a stabilizer subgroup in G of the hexagon $\sum_{i=1}^6 L_i$. Then

$$\Gamma \cong \mathrm{S}_4 \cong \psi\left(\langle A, B, C \rangle\right) \cong \langle A, B, C \rangle$$

by Lemma 4.17. We may assume that $\Gamma = \langle \psi(A), \psi(B), \psi(C) \rangle$, because G has two subgroups isomorphic to S_4 up to a conjugation (see Lemma B.1), which are switched by an outer automorphism in $\mathrm{Aut}(G)$ that can be realized as a complex conjugation in appropriate coordinates.

Lemma 4.18. The group Γ acts transitively on the lines $L_1, L_2, L_3, L_4, L_5, L_6$.

Proof. Since the action of the group Γ on \mathbb{P}^2 comes from an irreducible three-dimensional representation of the group Γ , the Γ -orbit of every line in \mathbb{P}^2 contains at least 3 lines. But

$$xyz = 0 \subset \mathbb{P}^2 \cong \mathrm{Proj}\left(\mathbb{C}[x, y, z]\right),$$

gives the only Γ -invariant 3-tuple of lines in \mathbb{P}^2 . \square

Let Ω be a stabilizer subgroup in Γ of the line L_1 . Then either $\Omega \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ or $\Omega \cong \mathbb{Z}_4$.

Lemma 4.19. The group Ω is not isomorphic to \mathbb{Z}_4 .

Proof. Suppose that $\Omega \cong \mathbb{Z}_4$. Then we may assume that Ω is generated $\psi(CB)$. One has

$$CB = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix},$$

and the eigenvalues of the matrix CB are 1 and $\pm\sqrt{-1}$.

Note that $(0, 1, 0)$ is the eigenvector of the matrix CB that corresponds to the eigenvalue 1, and $(\mp\sqrt{-1}, 0, 1)$ is its eigenvector that corresponds to the eigenvalue $\pm\sqrt{-1}$. Thus

$$L_1(x, y, z) = \mu\left(x \pm \sqrt{-1}z\right),$$

where μ is a non-zero complex number. The hexagon $\sum_{i=1}^6 L_i$ is given by

$$(x - \sqrt{-1}z)(x + \sqrt{-1}z)(y + \sqrt{-1}x)(y - \sqrt{-1}x)(z + \sqrt{-1}y)(z - \sqrt{-1}y) = 0,$$

which implies that the quartic form $x^4 + y^4 + z^4 + 3\epsilon(x^2y^2 + x^2z^2 + y^2z^2)$ is equal to $\mu_1(x - \sqrt{-1}z)^4 + \mu_2(x + \sqrt{-1}z)^4 + \mu_3(y + \sqrt{-1}x)^4 + \mu_4(y - \sqrt{-1}x)^4 + \mu_5(z + \sqrt{-1}y)^4 + \mu_6(z - \sqrt{-1}y)^4$ for some $(\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6) \in \mathbb{C}^6$. In particular, we obtain a system of linear equations

$$\begin{cases} 4\mu_3 + 4\mu_4 = 1 - \sqrt{-7}, \\ 4\mu_1 + 4\mu_2 = 1 - \sqrt{-7}, \\ 4\mu_5 + 4\mu_6 = 1 - \sqrt{-7}, \\ \mu_1 + \mu_2 + \mu_3 + \mu_4 = 1, \end{cases}$$

which is inconsistent. \square

Thus, we see that $\Omega \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.

Lemma 4.20. The subgroup Ω is not contained in a subgroup of the group G isomorphic to A_4 .

Proof. Suppose that Ω is contained in a subgroup of the group G that is isomorphic to A_4 . Then

$$\Omega = \left\{ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\},$$

which implies that the G -orbit of the line L_1 is the curve that is given by

$$xyz = 0 \subset \mathbb{P}^2 \cong \text{Proj}(\mathbb{C}[x, y, z]),$$

which is impossible by Lemma 4.18. \square

By Lemma 4.20, we may assume that Ω is generated by $\psi(AB)$ and $\psi(C)$. Therefore

$$\Omega = \left\{ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right\},$$

and the equation of the hexagon $\sum_{i=1}^6 L_i$ is

$$(x - z)(x + z)(y + x)(y - x)(z + y)(z - y) = 0.$$

Arguing as in the proof of Lemma 4.19, we obtain a contradiction.

The assertion of Theorem 4.8 is proved.

5. PROOF OF THEOREM 1.12

Throughout this section we use the assumptions and notation of Section 3.

Suppose that Theorem 1.12 is false. Let us derive a contradiction.

Lemma 5.1. There is a G -invariant linear system \mathcal{M} without fixed components on \mathbb{P}^3 such that

$$\text{NCS}(\mathbb{P}^3, \lambda\mathcal{M}) \neq \emptyset \neq \text{NCS}(\mathbb{P}^3, \lambda'\tau(\mathcal{M})),$$

where λ and λ' are positive rational numbers such that $\lambda\mathcal{M} \sim_{\mathbb{Q}} \lambda'\tau(\mathcal{M}) \sim_{\mathbb{Q}} \mathcal{O}_{\mathbb{P}^3}(4)$.

Proof. The required assertion is well-known (see [7], [2, Theorem A.16], [2, Corollary A.22]). \square

The assertion of Lemma 5.1 is known as the Noether–Fano inequality.

Lemma 5.2. Either $\text{mult}_{C_6}(\mathcal{M}) \leq 1/\lambda$ or $\text{mult}_{C_6}(\tau(\mathcal{M})) \leq 1/\lambda'$.

Proof. This is easy (cf. the proof of [2, Lemma B.15]). \square

Without loss of generality, we may assume that $\text{mult}_{C_6}(\mathcal{M}) \leq 1/\lambda$. Then $C_6 \notin \text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$.

Remark 5.3. The set $\text{NLCS}(\mathbb{P}^3, 2\lambda\mathcal{M})$ contains every center in $\text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$ by Lemma 2.8.

Let H be a sufficiently general hyperplane in \mathbb{P}^3 .

Lemma 5.4. Let Λ be a union of all curves contained in $\text{NLCS}(\mathbb{P}^3, 2\lambda\mathcal{M})$. Then $\deg(\Lambda) \leq 15$.

Proof. Let M_1 and M_2 be general surfaces in \mathcal{M} . Then

$$16/\lambda^2 = H \cdot M_1 \cdot M_2 \geq \sum_{P \in \Lambda \cap H} \text{mult}_P(M_1 \cdot M_2) > \deg(\Lambda)/\lambda^2$$

by Theorem 2.5, which implies that $\deg(\Lambda) \leq 15$. \square

Note that $\text{NLCS}(\mathbb{P}^3, 2\lambda\mathcal{M})$ may be non-empty even if $\text{NCS}(\mathbb{P}^3, \lambda\mathcal{M}) = \emptyset$ (see Example 3.11).

Lemma 5.5. The set $\text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$ does not contain any point in $\Sigma_8 \sqcup C_6$.

Proof. Let M_1 and M_2 be general surfaces in \mathcal{M} , and let Q be a general surface in \mathcal{Q} . Then

$$32/\lambda^2 = Q \cdot M_1 \cdot M_2 \geq \sum_{O_i \in \Sigma_8} \text{mult}_{O_i}(M_1 \cdot M_2) = 8 \text{mult}_{O_i}(M_1 \cdot M_2),$$

which implies that $\text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$ does not contain any point in Σ_8 by Theorem 2.7.

Suppose that $\text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$ contains a point $P \in C_6$. Put

$$M_1 \cdot M_2 = \nu C_6 + \Upsilon,$$

where μ is a non-negative integer, and Υ is an effective one-cycles such that $C_6 \not\subset \text{Supp}(\Upsilon)$. Then

$$\text{mult}_P(\nu C_6 + \Upsilon) \geq 4/\lambda^2$$

by Theorem 2.7. Let Θ be the G -orbit of the point P . Then $|\Theta| \geq 24$ by Lemma 2.15.

Let S be a general cubic surface in \mathbb{P}^3 such that $C_6 \subset S$. Then

$$48/\lambda^2 - 18\nu = S \cdot \Upsilon \geq \sum_{O \in \Theta} \text{mult}_O(\Upsilon) = |\Theta| \text{mult}_O(\Upsilon) > 24(4/\lambda^2 - \nu),$$

because C_6 is a scheme-theoretic intersection of cubic surfaces in \mathbb{P}^3 . Thus $\nu > 8/\lambda^2$. But

$$16/\lambda^2 = H \cdot M_1 \cdot M_2 = \nu H \cdot C_6 + H \cdot \Upsilon \geq \nu H \cdot C_6 = 6\nu,$$

which implies that $\nu \leq 8/(3\lambda^2)$, which is a contradiction, since $\nu > 8/\lambda^2$. \square

Lemma 5.6. The set $\text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$ does not contain the curve C_{14} .

Proof. Suppose that $C_{14} \in \text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$. Put $\mu = \text{mult}_{C_{14}}(\mathcal{M})$ and $m = \text{mult}_{O_1}(\mathcal{M})$. Then

$$m = \text{mult}_{O_2}(\mathcal{M}) = \text{mult}_{O_3}(\mathcal{M}) = \dots = \text{mult}_{O_8}(\mathcal{M}),$$

since \mathcal{M} is G -invariant. But $\mu > 1/\lambda$ because $C_{14} \in \text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$.

Let us find an upper bound for μ . Let M_1 and M_2 be general surfaces in \mathcal{M} . Then

$$16/\lambda^2 = H \cdot M_1 \cdot M_2 \geq \sum_{P \in C_{14} \cap H} \text{mult}_{C_{14}}(M_1 \cdot M_2) \geq 14\mu^2,$$

which implies that $\mu \leq \sqrt{8/7}/\lambda$. Thus, we have $1/\lambda < \mu \leq \sqrt{8/7}/\lambda$.

Lets us find a lower and an upper bound for m .

Let \bar{C}_{14} and $\bar{\mathcal{M}}$ be the proper transforms of the curve C_{14} and the linear system \mathcal{M} on the threefold U , respectively, let C be a general conic in $E_1 \cong \mathbb{P}^2$ such that $\bar{C}_{14} \cap E_1 \subset C$. Then

$$|\bar{C}_{14} \cap E_1| = 3,$$

and the points of the set $\bar{C}_{14} \cap E_1$ are non-coplanar (see the proof of Lemma 2.11), which implies that the conic C is not contained in the base locus of the linear system $\bar{\mathcal{M}}$. Hence

$$2m = C \cdot \bar{\mathcal{M}} \geq \sum_{P \in \bar{C}_{14} \cap E_1} \text{mult}_P(\bar{\mathcal{M}}) \geq \sum_{P \in \bar{C}_{14} \cap E_1} \text{mult}_{\bar{C}_{14}}(\bar{\mathcal{M}}) = 3\mu,$$

which implies that $m \geq 3\mu/2$. Thus, we see that $m > 3/(2\lambda)$, since $\mu > 1/\lambda$.

Let us find an upper bound for m (a trivial upper bound $m \leq 2/\lambda$ follows by Lemma 5.5).

Let \bar{M}_1 and \bar{M}_2 be the proper transforms on U of the surfaces M_1 and M_2 , respectively, and let \bar{Q} be the proper transform of a general surface in the linear system \mathcal{Q} . Then

$$32/\lambda^2 - 8m^2 = \bar{Q} \cdot \bar{M}_1 \cdot \bar{M}_2 \geq \sum_{P \in \bar{C}_{14} \cap \bar{Q}} \text{mult}_{\bar{C}_{14}}(\bar{M}_1 \cdot \bar{M}_2) = 4\text{mult}_{\bar{C}_{14}}(\bar{M}_1 \cdot \bar{M}_2) \geq 4\mu^2 \geq 4/\lambda^2,$$

which implies that $m \leq \sqrt{7/2}/\lambda$. Thus, we have $3/(2\lambda) < m \leq \sqrt{7/2}/\lambda$.

Let $v: W \rightarrow U$ be the blow up of the smooth curve \bar{C}_{14} , let F be the v -exceptional divisor, let $\hat{\mathcal{M}}$ and $\hat{\mathcal{D}}$ be the proper transforms of \mathcal{M} and \mathcal{D} (see Lemma 3.16) on W , respectively. Then

$$(5.7) \quad \begin{cases} \hat{\mathcal{M}} \sim_{\mathbb{Q}} (\pi \circ v)^* \left(\frac{4}{\lambda} H \right) - m \sum_{i=1}^8 v^*(E_i) - \mu F, \\ \hat{\mathcal{D}} \sim_{\mathbb{Q}} (\pi \circ v)^* (5H) - 2 \sum_{i=1}^8 v^*(E_i) - F. \end{cases}$$

Let \hat{D} be a general surface in $\hat{\mathcal{D}}$. Then \hat{D} is nef by Lemma 3.16 and (5.7).

Let \hat{M}_1 and \hat{M}_2 be the proper transforms on W of the surfaces M_1 and M_2 , respectively. Then

$$80/\lambda^2 - 2(17\mu^2 + 4m^2 + 56\mu/\lambda - 24m\mu) = \hat{D} \cdot \hat{M}_1 \cdot \hat{M}_2 \geq 0,$$

because \hat{D} is nef and $F^3 = -12$. Put $\bar{\mu} = \mu\lambda$ and $\bar{m} = m\lambda$. One has

$$\begin{cases} 17\bar{\mu}^2 + 4\bar{m}^2 + 56\bar{\mu} - 24\bar{m}\bar{\mu} \geq 40, \\ 1 < \bar{\mu} \leq \sqrt{\frac{8}{7}}, \\ \frac{3}{2} < \bar{m} \leq \sqrt{\frac{7}{2}}, \end{cases}$$

which easily leads to a contradiction. \square

Lemma 5.8. The set $\text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$ does not contain any point in C_{14} .

Proof. Suppose that $\text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$ contains a point $P \in C_{14}$. By Lemma 5.5, we have $P \notin \Sigma_8$, which implies that P is a smooth point of the curve C_{14} , by Lemma 3.13.

Let M_1 and M_2 be general surfaces in the linear system \mathcal{M} . Put

$$M_1 \cdot M_2 = \mu C_{14} + \Upsilon + \Xi,$$

where μ is a non-negative integer, and Υ and Ξ are effective one-cycles such that $C_{14} \not\subset \text{Supp}(\Upsilon)$, every irreducible component of the curve $\text{Supp}(\Upsilon)$ intersects the curve C_{14} by a non-empty set, and none of the irreducible components of the curve $\text{Supp}(\Xi)$ intersects C_{14} . Then

$$\text{mult}_P(\mu C_{14} + \Upsilon) \geq 4/\lambda^2$$

by Theorem 2.7. On the other hand, we have

$$16/\lambda^2 = H \cdot M_1 \cdot M_2 = \mu H \cdot C_{14} + H \cdot (\Upsilon + \Xi) \geq \mu H \cdot C_{14} = 14\mu,$$

which implies that $\mu \leq 8/(7\lambda^2)$.

Let Θ be the G -orbit of the point P . Then $|\Theta| \geq 42$ by Lemma 2.15.

Let D be a general quintic surface in the linear system \mathcal{D} (see Lemma 3.16). Then

$$80/\lambda^2 - 70\mu \geq 80/\lambda^2 - 70\mu - D \cdot \Xi = D \cdot \Upsilon \geq \sum_{O \in \Theta} \text{mult}_O(\Upsilon) = |\Theta| \text{mult}_O(\Upsilon) > 42 \left(4/\lambda^2 - \mu\right),$$

which immediately leads to a contradiction. \square

Thus, the set $\text{NLCS}(\mathbb{P}^3, 2\lambda\mathcal{M})$ contains a center that is not contained in $C_{14} \sqcup C_6$.

Lemma 5.9. Suppose that $\text{NLCS}(\mathbb{P}^3, 2\lambda\mathcal{M})$ contains a curve C such that $C_6 \neq C \neq C_{14}$. Then

$$C_{14} \notin \text{NLCS}(\mathbb{P}^3, 2\lambda\mathcal{M}),$$

and if $C_6 \in \text{NLCS}(\mathbb{P}^3, \lambda\mathcal{M})$, then $C_6 \cap C = \emptyset$ and $\deg(C) \leq 9$.

Proof. Let Z be a G -orbit of the curve C . If $C_{14} \in \text{NLCS}(\mathbb{P}^3, 2\lambda\mathcal{M})$, then

$$\deg(Z) + 14 = \deg(Z) + \deg(C_{14}) \leq 15$$

by Lemma 5.4, which is a contradiction, because there are no G -invariant lines in \mathbb{P}^3 .

Suppose that $C_6 \in \text{NLCS}(\mathbb{P}^3, \lambda\mathcal{M})$. Then $\deg(C) \leq \deg(Z) \leq 9$ by Lemma 5.4.

To complete the proof, we must show that $C_6 \cap Z = \emptyset$.

Suppose that $C_6 \cap Z \neq \emptyset$. Then $|C_6 \cap Z| \geq 24$ by Lemma 2.15.

Let \bar{Z} be a proper transform of the curve Z on the threefold V (see (3.9)), and let \bar{S} be a proper transform on V of a general cubic surface in \mathbb{P}^3 that passes through the curve C_6 . Then

$$3 \geq 3\deg(Z) - |C_6 \cap Z| \geq \bar{S} \cdot \bar{Z} = \deg(\beta(\bar{Z})),$$

which implies that \bar{Z} is contracted by β by Lemma 3.7, since $\beta(\bar{Z})$ is G -invariant.

The only curves contracted by the morphism β are proper transforms of the lines in \mathbb{P}^3 that are trisecants of the curve C_6 . Then $\beta(\bar{Z}) \subset C_6$, the subset $\beta(\bar{Z})$ is G -invariant and $|\beta(\bar{Z})| \leq 9$, which is impossible by Lemma 2.15. \square

It follows from Lemmas 5.5, 5.6 and 5.8 that the set $\text{NLCS}(\mathbb{P}^3, 2\lambda\mathcal{M})$ contains an irreducible subvariety that is not contained in $C_6 \cup C_{14}$. In fact, we can say a little bit more than this.

Lemma 5.10. There are $\mu \in \mathbb{Q}$ and $S \in \text{LCS}(\mathbb{P}^3, \mu\mathcal{M})$ such that

- the inequalities $0 < \mu < 2\lambda$ holds,
- the log pair $(\mathbb{P}^3, \mu\mathcal{M})$ is log canonical along S ,
- the subvariety S is a minimal center in $\text{LCS}(\mathbb{P}^3, \mu\mathcal{M})$,
- the subvariety S is not a point of the set Σ_8 ,
- the subvariety S is neither the curve C_6 nor the curve C_{14} ,
- exactly one of the following six cases is possible:
 - (A) the log pair $(\mathbb{P}^3, \mu\mathcal{M})$ has log canonical singularities,
 - (B) $\text{NCS}(\mathbb{P}^3, \mu\mathcal{M}) = C_{14}$ and S is a point such that $S \notin C_{14}$,
 - (C) $\text{NCS}(\mathbb{P}^3, \mu\mathcal{M}) = \Sigma_8$ and $S \cap \Sigma_8 = \emptyset$,
 - (D) $\text{NCS}(\mathbb{P}^3, \mu\mathcal{M}) = C_6$ and S is a curve such that $S \cap C_6 = \emptyset$ and $\deg(S) \leq 9$,
 - (E) $\text{NCS}(\mathbb{P}^3, \mu\mathcal{M}) = C_6$ and S is a point such that $S \notin \Sigma_8$,
 - (F) $\text{NCS}(\mathbb{P}^3, \mu\mathcal{M}) = C_6 \cup \Sigma_8$ and S is a curve such that $S \cap (C_6 \cup \Sigma_8) = \emptyset$ and $\deg(S) \leq 9$,
 - (G) $\text{NCS}(\mathbb{P}^3, \mu\mathcal{M}) = C_6 \cup \Sigma_8$ and S is a point such that $S \notin C_6 \cup \Sigma_8$.

Proof. Let us show how to find μ and S in several steps. Put

$$\mu_1 = \sup \left\{ \epsilon \in \mathbb{Q} \mid \text{the log pair } (\mathbb{P}^3, \epsilon\mathcal{M}) \text{ is log canonical} \right\},$$

and let S_1 be a minimal center in $\text{LCS}(\mathbb{P}^3, \mu_1\mathcal{M})$. Then $\mu_1 < 2\lambda$.

If S_1 is a curve, then $S_1 \cap \Sigma_8 = \emptyset$ by Lemmas 3.14 and 5.4, since S_1 is smooth by Theorem 2.10.

If $S_1 \cap \Sigma_8 = \emptyset$ and $S_1 \neq C_6$, then we have the case **(A)** by putting $\mu = \mu_1$ and $S = S_1$.
 To complete the proof, we may assume that either $S_1 \cap \Sigma_8 \neq \emptyset$ or $S_1 = C_6$.
 Let us consider the mutually excluding cases $S_1 \cap \Sigma_8 \neq \emptyset$ and $S_1 = C_6$ separately.
 Suppose that $S_1 = C_6$. Put

$$\mu_2 = \sup \left\{ \epsilon \in \mathbb{Q} \mid \text{the log pair } \left(\mathbb{P}^3, \epsilon \mathcal{M} \right) \text{ is log canonical outside of the curve } C_6 \right\} < 2\lambda,$$

let T_2 be a center in the set $\text{LCS}(\mathbb{P}^3, \mu_2 \mathcal{M})$ such that $T_2 \neq C_6$, and let S_2 be a minimal center in the set $\text{LCS}(\mathbb{P}^3, \mu_2 \mathcal{M})$ such that $S_2 \subseteq T_2$. If T_2 is a curve, then $\deg(T_2) \leq 9$ and

$$T_2 \cap \Sigma_8 = T_2 \cap C_6 = \emptyset$$

by Lemmas 3.14, 5.4 and 5.9, since $C_{14} \cap C_6 = \emptyset$ by Lemma 3.13 and $C_6 \in \text{NLCS}(\mathbb{P}^3, 2\lambda \mathcal{M})$.

If T_2 is a curve and $S_2 = T_2$, then we have the case **(D)** by putting $\mu = \mu_2$ and $S = S_2$.

If T_2 is a curve and S_2 is a point, then we have the case **(E)** by putting $\mu = \mu_2$ and $S = S_2$.

If $T_2 = S_2$ is a point not in Σ_8 , then we have the case **(E)** by putting $\mu = \mu_2$ and $S = S_2$.

To finish the case $S_1 = C_6$, we may assume that $S_2 = T_2$ is a point in Σ_8 . Put

$$\mu_3 = \sup \left\{ \epsilon \in \mathbb{Q} \mid \text{the log pair } \left(\mathbb{P}^3, \epsilon \mathcal{M} \right) \text{ is log canonical outside of the set } C_6 \cup \Sigma_8 \right\} < 2\lambda,$$

let T_3 be a center in the set $\text{LCS}(\mathbb{P}^3, \mu_3 \mathcal{M})$ such that $T_3 \not\subseteq C_6 \cup \Sigma_8$, and let S_3 be a minimal center in the set $\text{LCS}(\mathbb{P}^3, \mu_3 \mathcal{M})$ such that $S_3 \subseteq T_3$. If T_3 is a curve, then $\deg(T_3) \leq 9$ and

$$T_3 \cap \Sigma_8 = T_3 \cap C_6 = \emptyset$$

by Lemmas 3.14, 5.4 and 5.9, since $C_{14} \cap C_6 = \emptyset$ by Lemma 3.13 and $C_6 \in \text{NLCS}(\mathbb{P}^3, 2\lambda \mathcal{M})$.

If T_3 is a curve and $S_3 = T_3$, then we have the case **(F)** by putting $\mu = \mu_2$ and $S = S_2$.

If T_3 is a curve and S_3 is a point, then we have the case **(G)** by putting $\mu = \mu_2$ and $S = S_2$.

If $T_3 = S_3$ is a point, then we have the case **(G)** by putting $\mu = \mu_2$ and $S = S_2$.

Therefore, the case $S_1 = C_6$ is done, and we may assume that $S_1 \cap \Sigma_8 \neq \emptyset$, which implies that the subvariety S_1 is a point in Σ_8 by Theorem 2.10 and Lemmas 3.14 and 5.4. Put

$$\mu'_2 = \sup \left\{ \epsilon \in \mathbb{Q} \mid \text{the log pair } \left(\mathbb{P}^3, \epsilon \mathcal{M} \right) \text{ is log canonical outside of the set } \Sigma_8 \right\} < 2\lambda,$$

let T'_2 be a center in the set $\text{LCS}(\mathbb{P}^3, \mu'_2 \mathcal{M})$ such that $T'_2 \not\subseteq \Sigma_8$, and let S'_2 be a minimal center in the set $\text{LCS}(\mathbb{P}^3, \mu'_2 \mathcal{M})$ such that $S'_2 \subseteq T'_2$. Note that if S'_2 is a point in Σ_8 , then

$$T'_2 \neq S'_2$$

which implies that T'_2 must be the curve C_{14} by Lemmas 3.14 and 5.4.

If T'_2 is a point, then $S'_2 = T'_2$, and we have the case **(C)** by putting $\mu = \mu'_2$ and $S = S'_2$.

To complete the proof, we may assume that T'_2 is an irreducible curve.

If $S'_2 = T'_2$, then S'_2 is a smooth curve such that $\deg(S'_2) \leq 15$ by Theorem 2.10 and Lemma 5.4, which immediately implies that $T'_2 \cap \Sigma_8 = \emptyset$ by Lemma 3.14.

If $S'_2 = T'_2 \neq C_6$, then we have the case **(C)** by putting $\mu = \mu'_2$ and $S = S'_2$.

If S'_2 is a point not in Σ_8 , then we have the case **(C)** by putting $\mu = \mu'_2$ and $S = S'_2$.

To complete the proof, we may assume that either S'_2 is a point of the set Σ_8 and $T'_2 = C_{14}$, or we have $S'_2 = T'_2 = C_6$. Let us consider these cases separately.

Suppose that S'_2 is a point of the set Σ_8 and $T'_2 = C_{14}$. Put

$$\mu'_3 = \sup \left\{ \epsilon \in \mathbb{Q} \mid \text{the log pair } \left(\mathbb{P}^3, \epsilon \mathcal{M} \right) \text{ is log canonical outside of the curve } C_{14} \right\} < 2\lambda,$$

let T'_3 be a center in the set $\text{LCS}(\mathbb{P}^3, \mu'_3 \mathcal{M})$ such that $T'_3 \not\subseteq C_{14}$. Then T'_3 is a point by Lemma 5.9, which implies that we have the case **(B)** by putting $\mu = \mu'_3$ and $S = T'_3$.

To complete the proof, we may assume that $S'_2 = T'_2 = C_6$. Put

$$\mu''_3 = \sup \left\{ \epsilon \in \mathbb{Q} \mid \text{the log pair } \left(\mathbb{P}^3, \epsilon \mathcal{M} \right) \text{ is log canonical outside of the set } C_6 \cup \Sigma_8 \right\} < 2\lambda,$$

let T_3' be a center in the set $\text{LCS}(\mathbb{P}^3, \mu_3''\mathcal{M})$ such that $T_3'' \not\subset C_6 \cap \Sigma_8$, and let S_3'' be a minimal center in the set $\text{LCS}(\mathbb{P}^3, \mu_3''\mathcal{M})$ such that $S_3'' \subseteq T_3''$. Note that $C_{14} \notin \text{LCS}(\mathbb{P}^3, 2\lambda\mathcal{M})$ by Lemma 5.9.

If T_3'' is a point, then $S_3'' = T_3''$, and we have the case **(G)** by putting $\mu = \mu_3''$ and $S = S_3''$.

We may assume that T_3'' is a curve. Then $\deg(T_3'') \leq 9$ and

$$T_3'' \cap C_6 = T_3'' \cap \Sigma_8 = \emptyset$$

by Lemmas 3.14, 5.4 and 5.9, since $C_6 \in \text{LCS}(\mathbb{P}^3, 2\lambda\mathcal{M}) \not\cong C_{14}$.

Finally, we put $\mu = \mu_3''$ and $S = S_3''$. Then we have the case **(G)** if the center S_3'' is a point, and we have the case **(F)** if the center S_3'' is a curve, which completes the proof. \square

Let ϵ be any rational number such that $\mu < \epsilon\mu < 2\lambda$. Then it follows from Lemma 2.13 that there is a G -invariant linear system \mathcal{B} on \mathbb{P}^3 such that \mathcal{B} does not have fixed components, and there are positive rational numbers ϵ_1 and ϵ_2 such that $1 \geq \epsilon_1 \gg \epsilon_2 \geq 0$ and

$$\text{LCS}(\mathbb{P}^3, \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B}) = \left(\bigsqcup_{g \in G} \{g(S)\} \right) \bigsqcup \text{NLCS}(\mathbb{P}^3, \mu\mathcal{M}),$$

the log pair $(\mathbb{P}^3, \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B})$ is log canonical at every point of $g(Z)$ for every $g \in G$, and

$$\epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B} \sim_{\mathbb{Q}} \epsilon\mu\mathcal{M}.$$

Let Z be the G -orbit of the subvariety S . Then one of the following cases is possible:

- (A) $\text{LCS}(\mathbb{P}^3, \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B}) = Z$,
- (B) $\text{LCS}(\mathbb{P}^3, \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B}) = Z \sqcup C_{14}$ and Z is finite set,
- (C) $\text{LCS}(\mathbb{P}^3, \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B}) = Z \sqcup \Sigma_8$,
- (D) $\text{LCS}(\mathbb{P}^3, \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B}) = Z \sqcup C_6$ and Z is a curve such that $\deg(Z) \leq 9$,
- (E) $\text{LCS}(\mathbb{P}^3, \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B}) = Z \sqcup C_6$ and Z is a finite set such that $Z \cap \Sigma_8 = \emptyset$,
- (F) $\text{LCS}(\mathbb{P}^3, \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B}) = Z \sqcup C_6 \sqcup \Sigma_8$ and Z is a curve such that $\deg(Z) \leq 9$,
- (G) $\text{LCS}(\mathbb{P}^3, \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B}) = Z \sqcup C_6 \sqcup \Sigma_8$ and Z is a finite set.

Put $D = \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B}$. Let \mathcal{L} be the union of all connected components of the log canonical singularity subscheme $\mathcal{L}(\mathbb{P}^3, D)$ whose supports contains no components of the subvariety Z . Then

$$\text{Supp}(\mathcal{L}) = \begin{cases} \emptyset & \text{in the case (A),} \\ C_{14} & \text{in the case (B),} \\ \Sigma_8 & \text{in the case (C),} \\ C_6 & \text{in the cases (D) and (E),} \\ C_6 \sqcup \Sigma_8 & \text{in the cases (F) and (G),} \end{cases}$$

Let $\mathcal{I}(\mathbb{P}^3, D)$ be the multiplier ideal sheaf of the log pair (\mathbb{P}^3, D) . Then

$$(5.11) \quad h^0(\mathcal{O}_{\mathcal{L}} \otimes \mathcal{O}_{\mathbb{P}^3}(4)) + h^0(\mathcal{O}_Z \otimes \mathcal{O}_{\mathbb{P}^3}(4)) = 35 - h^0(\mathcal{O}_{\mathbb{P}^3}(4) \otimes \mathcal{I}(\mathbb{P}^3, D)) \leq 35$$

by Theorem 2.2.

Corollary 5.12. Suppose that Z is a finite set. Then

- the subset Z contains at most 35 points,
- the subset Z imposes independent linear conditions on quartic surfaces in \mathbb{P}^3 .

Lemma 5.13. Suppose that S is a point. Then $|Z| \neq 24$.

Proof. Suppose that $|Z| = 24$. Then $Z = \Sigma_{24}$ by Lemma 3.2, and there is $F \in |\mathcal{O}_{\mathbb{P}^3}(4)|$ such that

$$Z \setminus S \subset F \not\supset S$$

by Corollary 5.12. Thus, there is a unique point $P \in C_6$ such that $P \neq S$ and

$$F|_{C_6} = P + Z \setminus S \sim F_4|_{C_6} = Z,$$

which implies that $P \sim S$ on C_6 , which is impossible, since C_6 is a smooth curve of genus 3. \square

Lemma 5.14. The subvariety S is a curve.

Proof. Suppose that S is a point. By Lemmas 3.2 and 5.13 and Corollary 5.12, we have $|Z| = 28$, because $Z \neq \Sigma_8$. Without loss of generality, we may assume that $Z = \Sigma_{28}$.

Let M_1 and M_2 be general surfaces in \mathcal{M} . Put

$$M_1 \cdot M_1 = \Xi + \Lambda,$$

where Ξ and Λ are effective cycles such that $\text{Supp}(\Xi) \cap \text{Supp}(\Lambda)$ consists of finitely many points, and $\text{Supp}(\Xi) \supset \Sigma_{28} \not\subset \text{Supp}(\Lambda)$. If $S \in \text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$, then $\text{mult}_S(\Xi) > 4/\lambda^2$ by Theorem 2.7.

Let R be a general surface in \mathcal{R} (see Lemma 3.19). Then $\text{Supp}(\Xi) \cap R$ consists of at most finitely many points by Lemma 3.19 and Corollary 5.12. Thus, we must have

$$64/\lambda^2 \geq 64/\lambda^2 - R \cdot \Lambda = R \cdot \Xi \geq \sum_{P \in \Sigma_{28}} \text{mult}_P(\Xi) = 28 \text{mult}_S(\Xi),$$

which implies that $\text{mult}_S(\Xi) \leq 16/(7\lambda^2)$. So $g(S) \notin \text{NCS}(\mathbb{P}^3, \lambda\mathcal{M})$ for every $g \in G$.

Note that $\Sigma_{28} \not\subset C_6 \sqcup C_{14}$ by Lemma 2.15, since C_6 is smooth and $\text{Sing}(C_{14}) = \Sigma_8$.

It follows from Lemmas 5.5, 5.6 and 5.8 that the set $\text{NLCS}(\mathbb{P}^3, 2\lambda\mathcal{M})$ contains an irreducible subvariety that is not contained in $C_6 \cup C_{14} \cup \Sigma_{28}$. Put

$$\bar{\mu} = \sup \left\{ \epsilon \in \mathbb{Q} \mid \text{the log pair } (\mathbb{P}^3, \epsilon\mathcal{M}) \text{ is log canonical outside of the set } C_6 \cup C_{14} \cup \Sigma_{28} \right\}.$$

The set $\text{LCS}(\mathbb{P}^3, \bar{\mu}\mathcal{M})$ contains every point in Σ_{28} . If $\bar{\mu} > \mu$, then $\text{NLCS}(\mathbb{P}^3, \bar{\mu}\mathcal{M})$ also contains every points in Σ_{28} . It follows from Lemma 2.8 that $\bar{\mu} < 2\lambda$. Note that $\bar{\mu} \geq \mu$.

Let Ω be a center $\text{LCS}(\mathbb{P}^3, \bar{\mu}\mathcal{M})$ such that $\Omega \not\subset C_6 \cup C_{14} \cup \Sigma_{28}$. Note that Ω does exist.

Let us choose a center $\Gamma \in \text{LCS}(\mathbb{P}^3, \bar{\mu}\mathcal{M})$ in the following way:

- if Ω is a point, then we put $\Gamma = \Omega$,
- if Ω is a curve that is a minimal center in $\text{LCS}(\mathbb{P}^3, \bar{\mu}\mathcal{M})$, then we put $\Gamma = \Omega$,
- if Ω is a curve that is not a minimal center in $\text{LCS}(\mathbb{P}^3, \bar{\mu}\mathcal{M})$,
then let Γ be a point in Ω that is also a center in $\text{LCS}(\mathbb{P}^3, \bar{\mu}\mathcal{M})$.

Let Δ be a G -orbit of the center Γ . Then $\Delta \cap \Sigma_8 = \Delta \cap \Sigma_{28} = \emptyset$ by Lemmas 3.14, 3.17, 5.9, 5.4.

Let $\bar{\epsilon}$ be a rational number such that $\bar{\mu} < \bar{\epsilon}\bar{\mu} < 2\lambda$. Arguing as in the proof of Lemma 2.13, we obtain a G -invariant linear system \mathcal{B}' on \mathbb{P}^3 such that \mathcal{B}' does not have fixed components, and there are positive rational numbers $\bar{\epsilon}_1$ and $\bar{\epsilon}_2$ such that $1 \geq \bar{\epsilon}_1 \gg \bar{\epsilon}_2 \geq 0$ and

$$\text{LCS}(\mathbb{P}^3, \bar{\epsilon}_1\bar{\mu}\mathcal{M} + \bar{\epsilon}_2\mathcal{B}') = \begin{cases} \left(\bigsqcup_{g \in G} \{g(\Gamma)\} \right) \sqcup \left(\bigsqcup_{P \in \Sigma_{28}} \{P\} \right) \sqcup \text{NLCS}(\mathbb{P}^3, \bar{\mu}\mathcal{M}) & \text{if } \bar{\mu} = \mu, \\ \left(\bigsqcup_{g \in G} \{g(\Gamma)\} \right) \sqcup \text{NLCS}(\mathbb{P}^3, \bar{\mu}\mathcal{M}) & \text{if } \bar{\mu} > \mu. \end{cases}$$

Put $\bar{D} = \bar{\epsilon}_1\bar{\mu}\mathcal{M} + \bar{\epsilon}_2\mathcal{B}'$. Then Γ is a connected component of the subscheme $\mathcal{L}(\mathbb{P}^3, \bar{D})$.

Let $\bar{\mathcal{L}}$ be the union of all connected components of the subscheme $\mathcal{L}(\mathbb{P}^3, \bar{D})$ whose supports contain no components of the subvariety Δ . Then $h^0(\mathcal{O}_{\bar{\mathcal{L}}} \otimes \mathcal{O}_{\mathbb{P}^3}(4)) \geq 28$, since $\Sigma_{28} \subseteq \text{Supp}(\bar{\mathcal{L}})$.

Let $\mathcal{I}(\mathbb{P}^3, \bar{D})$ be the multiplier ideal sheaf of the log pair (\mathbb{P}^3, \bar{D}) . Then

$$h^0(\mathcal{O}_{\Delta} \otimes \mathcal{O}_{\mathbb{P}^3}(4)) = 35 - h^0(\mathcal{O}_{\mathbb{P}^3}(4) \otimes \mathcal{I}(\mathbb{P}^3, \bar{D})) - h^0(\mathcal{O}_{\bar{\mathcal{L}}} \otimes \mathcal{O}_{\mathbb{P}^3}(4)) \leq 7$$

by Theorem 2.2, which implies that Γ is not a point by Lemma 3.2. We see that Δ is a curve.

By Theorem 2.10, the curve Δ is a smooth curve in \mathbb{P}^3 of degree d and genus $g \leq 2d$. Then

$$\frac{\deg(\Delta)}{\deg(\Gamma)}(2d+1) \leq \frac{\deg(\Delta)}{\deg(\Gamma)}(4d-g+1) \leq 7,$$

which gives $d \leq 3$. Thus, we have $\Delta \neq \Gamma$, so that $\deg(\Delta) \geq 7\deg(\Gamma)$ by Corollary B.2. We have

$$21 \leq 7(2d+1) \leq (2d+1)\frac{\deg(\Delta)}{\deg(\Gamma)}(2d+1) \leq \frac{\deg(\Delta)}{\deg(\Gamma)}(4d-g+1) \leq 7,$$

which is a contradiction. \square

By Theorem 2.10, the curve S is a smooth curve of degree d and genus g such that $g \leq 2d$, which implies, in particular, that the case **(B)** is not possible by Lemma 5.9.

Let G_S be the stabilizer subgroup in G of the subvariety S . Put $p = h^0(\mathcal{L} \otimes \mathcal{O}_{\mathbb{P}^3}(4))$, put

$$q = h^0(\mathcal{O}_{\mathbb{P}^3}(4) \otimes \mathcal{I}(\mathbb{P}^3, D)),$$

and let r be the number of irreducible components of the curve Z . Then

$$(5.15) \quad r(4d-g+1) = 35 - q - p$$

by (5.11), the Riemann–Roch theorem and Remark 2.12.

Lemma 5.16. The equality $r = 1$ holds.

Proof. Suppose that $r \neq 1$. Then $r \geq 7$ by Corollary B.2, which implies that

$$4d-g+1 = \frac{35-q-p}{r} \leq 5$$

by (5.15). But $g \leq 2d$. Then $4d-g+1 \leq 5$, which implies that $g = p = q = 0$, $d = 1$ and $r = 7$.

The induced action of the group $G_S \cong S_4$ on the line S is faithful, which implies that $S \subset F_4$, since G_S -invariant subsets in S have at least 6 points. Then $Z \subset F_4$, which contradicts $q = 0$. \square

Therefore, we see that $Z = S$.

Remark 5.17. Let I be the trivial representation of the group G . By Lemma B.8 we have

$$H^0(\mathcal{O}_{\mathbb{P}^3}(4)) \cong I \oplus W_6 \oplus W_6 \oplus W_7 \oplus W_7 \oplus W_8,$$

where W_i is an irreducible representation of the group $G \cong \mathrm{PSL}_2(\mathbb{F}_7)$ of dimension i . Then

$$p+q \notin \{2, 3, 4, 5, 10, 11, 17\},$$

because p is divisible by 8. If $S \subset F_4$, then $p+q \notin \{0, 2, 3, 4, 5, 6, 10, 11, 12, 17\}$.

Note that there is a natural faithful action of the group G on the curve S .

Lemma 5.18. We have $g \in \{3, 8, 10, 15, 17, 22, 24, 29\}$.

Proof. This follows from Lemmas 2.16 and 3.2, because $g \leq 2d \leq 30$ by Lemma 5.4. \square

Lemma 5.19. Suppose that $d \neq 6$ and $d \neq 12$. Then $S \subset F_4$.

Proof. Suppose that $S \not\subset F_4$. Then $F_4 \cap S$ is union of some G -orbits $\Lambda_1, \dots, \Lambda_s$. Thus

$$\sum_{i=1}^s n_i |\Lambda_i| = 4d$$

for some positive integers n_1, n_2, \dots, n_s .

Using Lemma 2.16, we obtain that $d = 14$, $s = 1$, $n_1 = 1$ and $|\Lambda_1| = 56$. Then

$$g = 22 + q + p \leq 2d = 28,$$

which implies that $g = 22$ by Lemma 5.18 and Remark 5.17.

By Lemma 2.16, we have $|\Lambda_1| \neq 56$, which is a contradiction. \square

Note that $d \in \{6, 7, 8, 9\}$ in the cases **(D)**, **(E)**, **(F)**, **(G)**.

Lemma 5.20. The equality $d = 6$ is impossible.

Proof. Suppose that $d = 6$. Then it follows from Lemma 5.18 that

$$3 \leq g = q + p - 10 \leq 2d = 12,$$

which implies that $g \in \{3, 8, 10\}$. But $g \leq 4$ by Theorem 2.14. Then $S = C_6$ by Lemma 3.7, which is a contradiction, because $S \neq C_6$ by the choice of the center $S \in \text{LCS}(\mathbb{P}^3, \mu\mathcal{M})$. \square

Lemma 5.21. The equality $d = 7$ is impossible.

Proof. Suppose that $d = 7$. Then $g \leq 7$ by Theorem 2.14 and $g = 3$ by Lemma 5.18.

By Lemma 5.19, we see that $S \subset F_4$. Arguing as in the proof of Lemma 5.18, we see that the curve S is contained in the intersection $F_4 \cap F_6 \cap F_{14}$, which is impossible by Lemma 3.6. \square

Lemma 5.22. The equality $d = 8$ is impossible.

Proof. Suppose that $d = 8$. Then $g \leq 9$ by Theorem 2.14 and $g \in \{3, 8\}$ by Lemma 5.18.

By Lemma 5.19, we see that $S \subset F_4$. Arguing as in the proof of Lemma 5.18, we see that the curve S is contained in $F_4 \cap F_6 \cap F'_8$ if $g = 8$. Then $g = 3$ by Lemma 3.6. Then

$$\mathcal{O}_{\mathbb{P}^3}(1) \Big|_C \sim 2K_S$$

by Theorem 2.18, which is impossible, since the vector space $H^0(\mathcal{O}_S(2K_S))$ is an irreducible six-dimensional representation of the group $G \cong \text{PSL}_2(\mathbb{F}_7)$ by Lemma B.8. \square

Lemma 5.23. The equality $d = 9$ is impossible.

Proof. Suppose that $d = 9$. Then $g \leq 12$ by Theorem 2.14 and $g \in \{3, 8, 10\}$ by Lemma 5.18.

By Lemma 5.19, we have $S \subset F_4$. Arguing as in the proof of Lemma 5.18, we see that

$$S \subset F_4 \cap F_6 \cap F'_8,$$

if $g = 8$. Similarly, we see that $S \subset F_4 \cap F_6 \cap F_{14}$ if $g = 3$. Thus $g = 10$ by Lemma 3.6.

The curve S is a complete intersection of two cubic surfaces in \mathbb{P}^3 (see [14, Example 6.4.3]), which is impossible, because there are no G -invariant pencils of cubic surfaces by Remark 3.4. \square

Thus, the cases **(D)**, **(E)**, **(F)**, **(G)** are not possible.

Lemma 5.24. The equality $d = 10$ is impossible.

Proof. Suppose that $d = 10$. Then $p + q \notin \{2, 4, 11\}$ by Lemma 5.19 and Remark 5.17. But

$$g = 6 + q + p \leq 2d = 20,$$

which implies that $g \in \{8, 10, 15, 17\}$ by Lemma 5.18. Thus, we see that $g = 15$ and $p + q = 9$.

By Lemma 5.19, we see that $S \subset F_4$. Arguing as in the proof of Lemma 5.18, we see that the curve S is contained in the intersection $F_4 \cap F_6 \cap F'_8$, which is impossible by Lemma 3.6. \square

Lemma 5.25. The equality $d = 11$ is impossible.

Proof. Suppose that $d = 11$. Then $p + q \notin \{0, 5, 12\}$ by Lemma 5.19 and Remark 5.17. But

$$g = 10 + q + p \leq 2d = 22,$$

which implies that $g \in \{10, 15, 17, 22\}$ by Lemma 5.18. Thus, we see that $g = 17$ and $p + q = 7$.

By Lemma 5.19, we see that $S \subset F_4$. Arguing as in the proof of Lemma 5.18, we see that the curve S is contained in the intersection $F_4 \cap F_6 \cap F'_8$, which is impossible by Lemma 3.6. \square

Lemma 5.26. The equality $d = 12$ is impossible.

Proof. Suppose that $d = 12$. Then $p + q \notin \{3, 5, 10\}$ by Remark 5.17. But

$$g = 14 + q + p \leq 2d = 24,$$

which implies that $g \in \{15, 17, 22, 24\}$ by Lemma 5.18. Thus either $g = 15$ or $g = 22$.

Arguing as in the proof of Lemma 5.18 and using Lemma 2.16, we see that

$$S \subset F_4 \cap F_6 \cap F'_8,$$

which is impossible by Lemma 3.6. □

Lemma 5.27. The equality $d = 13$ is impossible.

Proof. Suppose that $d = 13$. Then $p + q \notin \{4, 6\}$ by Lemma 5.19 and Remark 5.17. But

$$g = 18 + q + p \leq 2d = 26,$$

which implies that $g \in \{22, 24\}$ by Lemma 5.18. Thus $p + q \in \{4, 6\}$, which is a contradiction. □

Lemma 5.28. The equality $d = 14$ is impossible.

Proof. Suppose that $d = 14$. Then $p + q \neq 0$ by Lemma 5.19 and Remark 5.17. But

$$g = 22 + q + p \leq 2d = 28,$$

which implies that $g = 22$ by Lemma 5.18. Thus $p = q = 0$, which is a contradiction. □

By Lemmas 5.20, 5.21, 5.22, 5.23, 5.24, 5.25, 5.26, 5.27, 5.28 and Theorem 2.14, one has $d \geq 15$, which implies that $d = 15$ by Lemma 5.4. Then $p + q \neq 3$ by Lemma 5.19 and Remark 5.17. But

$$g = 26 + q + p \leq 2d = 30,$$

which implies that $g = 29$ by Lemma 5.18. Thus, we have $p + q = 3$, which is a contradiction.

The assertion of Theorem 1.12 is proved.

6. PROOF OF THEOREM 1.13

Throughout this section we use assumptions and notation of Theorem 1.13, and we identify the threefold X with its anticanonical image in \mathbb{P}^{13} (cf. Theorem 4.5).

Suppose that Theorem 1.13 is false. Let us derive a contradiction.

Lemma 6.1. There is a G -invariant linear system \mathcal{M} without fixed components on X such that

$$\text{NCS}(X, \lambda\mathcal{M}) \neq \emptyset,$$

where λ is a positive rational number such that $-K_X \sim_{\mathbb{Q}} \lambda\mathcal{M}$.

Proof. The required assertion is well-known (see [7], [2, Corollary A.18]). □

Let H be a general surface in the linear system $|-K_X|$.

Lemma 6.2. Let Λ be a union of all curves in $\text{NLCS}(X, 2\lambda\mathcal{M})$. Then $\deg(\Lambda) \leq 21$.

Proof. Let M_1 and M_2 be general surfaces in \mathcal{M} . By Theorem 2.5, we have

$$22/\lambda^2 = H \cdot M_1 \cdot M_2 \geq \sum_{P \in \Lambda \cap H} \text{mult}_P(M_1 \cdot M_2) > (H \cdot \Lambda)/\lambda^2 = \deg(\Lambda)/\lambda^2,$$

which implies that $\deg(\Lambda) \leq 21$. □

By Lemma 2.8, the set $\text{NLCS}(X, 2\lambda\mathcal{M})$ contains every center in $\text{NCS}(X, \lambda\mathcal{M})$. Put

$$\hat{\mu} = \sup \left\{ \epsilon \in \mathbb{Q} \mid \text{the log pair } (X, \epsilon\mathcal{M}) \text{ is log canonical} \right\} < 2\lambda,$$

let Σ_8 be the unique G -invariant subset of the threefold X consisting of 8 points (see Lemma 6.3), and let F be the unique G -invariant surface in the linear system $|-K_X|$ (see Corollary 4.6).

Lemma 6.3. Suppose that $\Sigma_8 \subset \text{LCS}(X, \hat{\mu}\mathcal{M})$. Then

$$\Sigma_8 \not\subset \text{NCS}(X, \lambda\mathcal{M}),$$

and the set $\text{NLCS}(X, 2\lambda\mathcal{M})$ does not contain curves that pass through a point in Σ_8 .

Proof. Take $\hat{\epsilon} \in \mathbb{Q}$ such that $\hat{\mu} < \hat{\epsilon}\hat{\mu} < 2\lambda$. By Lemma 2.13, there is a G -invariant linear system without fixed components $\hat{\mathcal{B}}$ on X , and there are $\hat{\epsilon}_1 \in \mathbb{Q} \ni \hat{\epsilon}_2$ such that $1 \geq \hat{\epsilon}_1 \gg \hat{\epsilon}_2 \geq 0$ and

$$\text{LCS}(X, \hat{\epsilon}_1\hat{\mu}\mathcal{M} + \hat{\epsilon}_2\hat{\mathcal{B}}) = \Sigma_8,$$

the log pair $(X, \hat{\epsilon}_1\hat{\mu}\mathcal{M} + \hat{\epsilon}_2\hat{\mathcal{B}})$ is log canonical, and $\hat{\epsilon}_1\hat{\mu}\mathcal{M} + \hat{\epsilon}_2\hat{\mathcal{B}} \sim_{\mathbb{Q}} \hat{\epsilon}\hat{\mu}\mathcal{M}$. Put $\hat{D} = \hat{\epsilon}_1\hat{\mu}\mathcal{M} + \hat{\epsilon}_2\hat{\mathcal{B}}$.

Let $\mathcal{I}(X, \hat{D})$ is the multiplier ideal sheaf of the log pair (X, \hat{D}) . The sequence of groups

$$0 \rightarrow H^0(\mathcal{O}_X(H) \otimes \mathcal{I}(X, \hat{D})) \rightarrow H^0(\mathcal{O}_X(H)) \rightarrow H^0(\mathcal{O}_{\Sigma_8}) \rightarrow 0$$

is exact by Theorem 2.2. Then Σ_8 imposes independent linear conditions on surfaces in $|-K_X|$.

Let \mathcal{D} be a linear subsystem of the linear system $|-K_X|$ that consists of all surfaces passing through Σ_8 . Then it follows from Theorem 4.5 that \mathcal{D} is the unique five-dimensional G -invariant linear subsystem of the linear system $|-K_X|$ and $F \notin \mathcal{D}$.

It is clear that the base locus of the linear system \mathcal{D} does not contain surfaces.

Suppose that the base locus of the linear system \mathcal{D} does not contain any curve that passes through a point in Σ_8 . Let us show that this assumption implies everything we have to prove.

Let M_1 and M_2 be general surfaces in the linear system \mathcal{M} . Put

$$M_1 \cdot M_1 = \Xi + \Lambda,$$

where Ξ and Λ are effective one-cycles such that

$$\text{Supp}(\Xi) \supset \Sigma_8 \not\subset \text{Supp}(\Lambda)$$

and $\text{Supp}(\Xi) \cap \text{Supp}(\Lambda)$ consists of at most finitely many points.

Let D be a general surface in the linear system \mathcal{D} . Since $|\text{Supp}(\Xi) \cap D| < +\infty$, one has

$$22/\lambda^2 \geq 22/\lambda^2 - D \cdot \Lambda = D \cdot \Xi \geq \sum_{P \in \Sigma_8} \text{mult}_P(\Xi) = 8\text{mult}_P(\Xi),$$

for every point $P \in \Sigma_8$. Then $\text{NCS}(X, \lambda\mathcal{M})$ contains no points in Σ_8 by Theorem 2.7.

Let Z be a G -orbit of an irreducible curve in X such that $\Sigma_8 \subset Z$. Then

$$\deg(Z) = D \cdot Z \geq \sum_{P \in \Sigma_8} \text{mult}_P(D)\text{mult}_P(Z) \geq \sum_{P \in \Sigma_8} \text{mult}_P(Z) \geq 24$$

by Lemma 2.11. Hence $\text{NLCS}(X, 2\lambda\mathcal{M})$ contains no components of the curve Z by Lemma 6.2, which implies that $\text{NLCS}(X, 2\lambda\mathcal{M})$ does not contain curves that pass through a point in Σ_8 .

To complete the proof, we must show that the base locus of the linear system \mathcal{D} contains no curves that contain a point in Σ_8 . Suppose that this is not true. Let us derive a contradiction.

The base locus of the linear system \mathcal{D} contains a curve C that is a G -orbit of an irreducible curve such that $\Sigma_8 \subset C$. Then $\text{mult}_P(C) \geq 3$ for every $P \in \Sigma_8$ by Lemma 2.11.

Let Q_1, Q_2, Q_3 and Q_4 be general points in X , and let \mathcal{H} be a linear subsystem in $|-K_X|$ that consists of all surfaces that contain the set $\{Q_1, Q_2, Q_3, Q_4\}$. Then $\mathcal{H} \cap \mathcal{D}$ is a pencil, and the base locus of the linear system \mathcal{H} contains no curves.

Let H be a general surface in \mathcal{H} , let D_1 and D_2 be general surfaces in the pencil $\mathcal{H} \cap \mathcal{D}$. Then

$$22 = H \cdot D_2 \cdot D_2 \geq \sum_{i=1}^4 \text{mult}_{Q_i}(H) \text{mult}_{Q_i}(D_1) \text{mult}_{Q_i}(D_2) + |H \cap C| \geq 4 + \deg(C),$$

since $\deg(C) = |H \cap C|$. Thus, we see that $\deg(C) \leq 18$.

Note that $C \not\subset F$, since $\Sigma_8 \not\subset F$. Therefore, we have

$$\sum_{P \in F \cap C} \text{mult}_P(C) \leq |F \cap C| \leq F \cdot C = \deg(C) \leq 18,$$

because $C \not\subset F$. Applying Theorem 4.8, we see that

$$\deg(C) = F \cdot C = |F \cap C| = 14$$

and C is smooth at every point of the set $F \cap C$. Then C is reducible by Lemma 2.15.

Put $C = \sum_{i=1}^r C_i$, where C_i is an irreducible curve and $r \in \mathbb{Z}_{>0}$. Then $r = 14/\deg(C_1)$.

By Corollary B.2, either $\deg(C_1) = 1$ or $\deg(C_1) = 2$. On the other hand, we know that

$$\deg(C_1) + 1 \geq |C_1 \cap \Sigma_8|,$$

since Σ_8 imposes independent linear conditions on surfaces in $|-K_X|$. Hence, we must have

$$\frac{14}{\deg(C_1)} = r \geq \frac{8 \cdot 3}{|C_1 \cap \Sigma_8|},$$

because $\text{mult}_P(C) \geq 3$ for every point $P \in \Sigma_8$. Therefore $|C_1 \cap \Sigma_8| = 1 + \deg(C_1)$.

Note that $\deg(C_1) \neq 1$. Indeed, if $\deg(C_1) = 1$, then $|C_1 \cap \Sigma_8| = 2$ and $r = 28$, because the group G acts doubly transitive on the set Σ_8 (see [12, p. 173]), which is a contradiction.

Thus, we see that C_1 is an irreducible conic and $r = 7$, so that $|C_i \cap \Sigma_8| = 3$.

Let Π_1 be a plane in \mathbb{P}^{13} such that $C_1 \subset \Pi_1$. Then Π_1 contains 3 lines L_1^1, L_1^2 and L_1^3 such that

$$|L_1^1 \cap \Sigma_8| = |L_1^2 \cap \Sigma_8| = |L_1^3 \cap \Sigma_8| = 2,$$

and the G -orbit of the line L_1^1 consists of 28 different lines, since G acts doubly transitive on Σ_8 .

Now one can easily see that the G -orbit of the plane Π_1 consists of at least $28/3 > 9$ planes, which is impossible since the G -orbit of Π_1 consists of at most $r = 7$ planes. \square

By Lemma 6.3, the set $\text{NLCS}(X, 2\lambda\mathcal{M})$ contains a center that is not contained in Σ_8 . Put

$$\mu = \sup \left\{ \epsilon \in \mathbb{Q} \mid \text{the log pair } (X, \epsilon\mathcal{M}) \text{ is log canonical outside of the subset } \Sigma_8 \right\}$$

in the case when $\Sigma_8 \subset \text{LCS}(X, \hat{\mu}\mathcal{M})$. If $\Sigma_8 \not\subset \text{LCS}(X, \hat{\mu}\mathcal{M})$, then put $\mu = \hat{\mu}$.

Lemma 6.4. There exist a minimal center in $\text{LCS}(X, \mu\mathcal{M})$ that is not a point of the set Σ_8 .

Proof. This immediately follows from Lemma 6.3. \square

Let S be a minimal center in $\text{LCS}(X, \mu\mathcal{M})$ such that $S \not\subset \Sigma_8$, and let Z be its G -orbit.

Remark 6.5. If $Z = S$ and S is a curve, then $Z \cap \Sigma_8 = \emptyset$ by Theorem 2.10 and Lemma 2.15.

Let ϵ be any rational number such that $\mu < \epsilon\mu < 2\lambda$. Then it follows from Lemma 2.13 that there exists a G -invariant linear system \mathcal{B} on X such that \mathcal{B} does not have fixed components, and there exist positive rational numbers ϵ_1 and ϵ_2 such that $1 \geq \epsilon_1 \gg \epsilon_2 \geq 0$ and

$$\text{LCS}(X, \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B}) = \left(\bigsqcup_{g \in G} \{g(S)\} \right) \bigsqcup \text{NLCS}(X, \mu\mathcal{M}),$$

the log pair $(X, \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B})$ is log canonical at every point of the subvariety Z , and

$$\epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B} \sim_{\mathbb{Q}} \epsilon\mu\mathcal{M}.$$

Put $D = \epsilon_1\mu\mathcal{M} + \epsilon_2\mathcal{B}$. By Lemma 6.4, we may have the following cases:

- (A) $\text{LCS}(X, D) = Z$ and the log pair (X, D) is log canonical,
- (B) $\text{LCS}(X, D) = Z \sqcup \Sigma_8$ and Z is a finite set,
- (C) $\text{LCS}(X, D) = Z \sqcup \Sigma_8$ and Z is a curve.

Let \mathcal{L} be the union of all connected components of the subscheme $\mathcal{L}(X, D)$ whose supports do not contain any component of the subvariety Z . Then

$$\text{Supp}(\mathcal{L}) = \begin{cases} \emptyset & \text{in the case (A),} \\ \Sigma_8 & \text{in the cases (B) and (C).} \end{cases}$$

Let $\mathcal{I}(X, D)$ be the multiplier ideal sheaf of the log pair (X, D) . Then

$$(6.6) \quad h^0(\mathcal{O}_Z \otimes \mathcal{O}_X(H)) = 14 - h^0(\mathcal{O}_X(H) \otimes \mathcal{I}(X, D)) - h^0(\mathcal{O}_{\mathcal{L}}) \leq 14$$

by Theorem 2.2, since \mathcal{L} is at most a zero-dimensional subscheme. We have $h^0(\mathcal{O}_{\mathcal{L}}) \in \{0, 8\}$.

Corollary 6.7. If Z is a finite set, then

$$|Z| \leq 14,$$

and the points of Z impose independent linear conditions on surfaces in $|-K_X|$.

Lemma 6.8. The subvariety S is a curve.

Proof. Suppose that S is not a curve. Then $|Z| = 14$ by Theorem 4.8 and Corollary 6.7, because we know that $Z \neq \Sigma_8$. Note that Z is not contained in any surface in $|-K_X|$ by Corollary 6.7.

Let \mathcal{R} be a linear subsystem of the linear system $|-2K_X|$ that consists of all surfaces in $|-2K_X|$ that pass through the set Z . Then its base locus consists of the set Z by [13, Theorem 2].

Let M_1 and M_2 be general surfaces in the linear system \mathcal{M} . Put

$$M_1 \cdot M_1 = \Xi + \Lambda,$$

where Ξ and Λ are effective one-cycles such that

$$\text{Supp}(\Xi) \supset Z \not\subset \text{Supp}(\Lambda)$$

and $\text{Supp}(\Xi) \cap \text{Supp}(\Lambda)$ consists of at most finitely many points. If $S \in \text{NCS}(X, \lambda\mathcal{M})$, then

$$\text{mult}_S(\Xi) > 4/\lambda^2$$

by Theorem 2.7. Let R be a general surface in the linear system \mathcal{R} . Then

$$44/\lambda^2 \geq 44/\lambda^2 - R \cdot \Lambda = R \cdot \Xi \geq \sum_{P \in Z} \text{mult}_P(\Xi) = 14 \text{mult}_S(\Xi),$$

which implies that $g(S) \notin \text{NCS}(X, \lambda\mathcal{M})$ for every $g \in G$.

By Lemma 6.3, the set $\text{NLCS}(X, 2\lambda\mathcal{M})$ contains a center not contained in $\Sigma_8 \cup Z$. Put

$$\bar{\mu} = \sup \left\{ \epsilon \in \mathbb{Q} \mid \text{the log pair } (X, \epsilon\mathcal{M}) \text{ is log canonical outside of the set } \Sigma_8 \cup Z \right\}.$$

Note that $\bar{\mu} \geq \mu$ and $S \in \text{LCS}(X, \bar{\mu}\mathcal{M})$. Moreover, if $\bar{\mu} > \mu$, then

$$S \in \text{NLCS}(X, \bar{\mu}\mathcal{M}),$$

because $S \in \text{LCS}(X, \mu\mathcal{M})$. It follows from Lemma 2.8 that $\bar{\mu} < 2\lambda$.

Let Γ be a center in $\text{LCS}(X, \bar{\mu}\mathcal{M})$ such that $\Gamma \not\subset \Sigma_8 \sqcup Z$, and let Δ be its G -orbit. Then

$$\Delta \cap \Sigma_8 = \emptyset$$

by Lemma 6.3. Note that $\Delta \cap Z = \emptyset$ if Γ is a point and $\Gamma \in \text{NLCS}(X, 2\lambda\mathcal{M})$.

Suppose that the set $\text{NLCS}(X, 2\lambda\mathcal{M})$ does not contain curves in X that have a non-empty intersection with the set Z . Let us use this assumption to derive a contradiction.

Let $\bar{\epsilon}$ be a rational number such that $\bar{\mu} < \bar{\epsilon}\bar{\mu} < 2\lambda$. Arguing as in the proof of Lemma 2.13, we obtain a G -invariant linear system \mathcal{B}' on X such that \mathcal{B}' does not have fixed components, and there are positive rational numbers $\bar{\epsilon}_1$ and $\bar{\epsilon}_2$ such that $1 \geq \bar{\epsilon}_1 \gg \bar{\epsilon}_2 \geq 0$ and

$$\mathbb{LCS}(X, \bar{\epsilon}_1\bar{\mu}\mathcal{M} + \bar{\epsilon}_2\mathcal{B}') = \begin{cases} \left(\bigsqcup_{g \in G} \{g(\Gamma)\} \right) \bigsqcup \left(\bigsqcup_{g \in G} \{g(S)\} \right) \bigsqcup \text{NLCS}(X, \bar{\mu}\mathcal{M}) & \text{if } \bar{\mu} = \mu, \\ \left(\bigsqcup_{g \in G} \{g(\Gamma)\} \right) \bigsqcup \text{NLCS}(X, \bar{\mu}\mathcal{M}) & \text{if } \bar{\mu} > \mu. \end{cases}$$

Put $\bar{D} = \bar{\epsilon}_1\bar{\mu}\mathcal{M} + \bar{\epsilon}_2\mathcal{B}'$. Then Γ is a connected component of the subscheme $\mathcal{L}(X, \bar{D})$.

Let $\bar{\mathcal{L}}$ be the union of all connected components of the subscheme $\mathcal{L}(X, \bar{D})$ whose supports do not contain any component of the subvariety Δ . Then $Z \subseteq \text{Supp}(\bar{\mathcal{L}})$, which implies that

$$h^0(\mathcal{O}_{\bar{\mathcal{L}}} \otimes \mathcal{O}_X(H)) \geq 14,$$

because $\text{NLCS}(X, 2\lambda\mathcal{M})$ does not contain curves that have a non-empty intersection with Z .

Let $\mathcal{I}(X, \bar{D})$ be the multiplier ideal sheaf of the log pair (X, \bar{D}) . Then

$$0 \leq h^0(\mathcal{O}_{\Delta} \otimes \mathcal{O}_X(H)) = 14 - h^0(\mathcal{O}_X(H) \otimes \mathcal{I}(X, \bar{D})) - h^0(\mathcal{O}_{\bar{\mathcal{L}}} \otimes \mathcal{O}_X(H)) \leq 0,$$

by Theorem 2.2. Thus, we have $h^0(\mathcal{O}_{\Delta} \otimes \mathcal{O}_X(H)) = 0$, which implies that Γ is not a point.

It follows from Theorem 2.10 that Γ is a smooth curve of genus g such that $H \cdot \Gamma \geq 2g - 1$.

By Remark 2.12, the curve Δ is a disjoint union of smooth curves isomorphic to Γ . Then

$$0 = h^0(\mathcal{O}_{\Delta} \otimes \mathcal{O}_X(H)) \geq H \cdot \Gamma - g + 1 > 0,$$

which is a contradiction.

Thus, there is a curve $C_1 \in \text{NLCS}(X, 2\lambda\mathcal{M})$ such that $C_1 \cap Z \neq \emptyset$.

Let C be the G -orbit of the curve C_1 . Then $\text{NLCS}(X, 2\lambda\mathcal{M})$ contains every irreducible component of the curve C , which implies that $\deg(C) \leq 21$. Hence, we have $Z \subset C$. But

$$\text{mult}_P(C) \geq 3$$

for every $P \in \Sigma_8$ by Lemma 2.11. Thus, we have

$$42 \geq 2\deg(C) = R \cdot C \geq \sum_{P \in Z} \text{mult}_P(R) \text{mult}_P(C) \geq 42,$$

where R is a general surface in \mathcal{R} . Therefore $\deg(C) = 21$ and $\text{mult}_P(C) = 3$ for every $P \in \Sigma_8$.

Note that $C \not\subset F$, since $Z \not\subset F$. Thus

$$\text{mult}_P(C) |F \cap C| \leq \text{mult}_P(F \cdot C) |F \cap C| = F \cdot C = \deg(C) = 21,$$

for every point $P \in Z$. So the curve C is smooth at every point of the set $F \cap C$ by Theorem 4.8, which immediately implies that C is reducible by Lemma 2.15.

Put $C = \sum_{i=1}^r C_i$, where C_i is an irreducible curve and $r \in \mathbb{Z}_{>0}$. Then $\deg(C_1) \in \{1, 3\}$. But

$$\deg(C_1) + 1 \geq |C_1 \cap Z|,$$

because the points of Z impose independent linear conditions on surfaces in $|-K_X|$. We have

$$\frac{21}{\deg(C_1)} = r \geq \frac{14 \cdot 3}{|C_1 \cap Z|},$$

because $\text{mult}_P(C) \geq 3$ for every point $P \in Z$. Thus $\deg(C_1) = 1$, $r = 21$ and $|C_1 \cap Z| = 2$.

Since irreducible components of the curve C are lines and $\text{mult}_P(C) = 3$ for every point $P \in Z$, we can easily see that each connected component of the curve C must have at least 4 components, so that C has at most 6 connected components. Then C is connected by Corollary B.2.

Let P be a point in Z , let G_P be its stabilizer subgroup in G . Then $G_P \cong A_4$ by Lemma B.1, and there are exactly 3 irreducible components of the curve C containing P , since $\text{mult}_P(C) = 3$.

Without loss of generality, we may assume that $P = C_1 \cap C_2 \cap C_3$.

The group G_P naturally acts on the set $\{C_1, C_2, C_3\}$, which implies that there is a subgroup

$$G'_P \subset G_P \cong A_4$$

such that G'_P acts trivially on $\{C_1, C_2, C_3\}$ and $G'_P \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. Note that C_1 is G'_P -invariant.

Let \bar{P} be the point in $Z \cap C_1$ such that $\bar{P} \neq P$, let $G_{\bar{P}}$ be its stabilizer subgroup in G . Then

$$\mathbb{Z}_2 \times \mathbb{Z}_2 \cong G'_P \subset G_{\bar{P}} \cong A_4,$$

because $Z \cap C_1 = \{P, \bar{P}\}$. But the group $G_{\bar{P}}$ contains a unique subgroup that is isomorphic to G'_P , which very easily (almost immediately) implies that every point of the set Z is G'_P -invariant, which is impossible, since the group G acts faithfully on Z , because the group G is simple. \square

By Theorem 2.10, the curve S is a smooth curve of genus g such that $\deg(S) \geq 2g - 1$.

Remark 6.9. By Remark 2.12, the curve Z is a disjoint union of smooth curves isomorphic to S .

Let r be the number of connected components of the curve Z . Put $d = \deg(S)$. Then

$$(6.10) \quad r(d - g + 1) = 14 - h^0(\mathcal{O}_X(H) \otimes \mathcal{I}(X, D)) - h^0(\mathcal{O}_{\mathcal{L}})$$

by (6.6). Note that $h^0(\mathcal{O}_{\mathcal{L}}) = 0$ if and only if $\mathcal{L}(X, D) = Z$. Put $q = h^0(\mathcal{O}_X(H) \otimes \mathcal{I}(X, D))$.

Lemma 6.11. One has $\mathcal{L}(X, D) = Z$.

Proof. Suppose that $Z \subsetneq \mathcal{L}(X, D)$. Then $h^0(\mathcal{O}_{\mathcal{L}}) = 8$. It follows from (6.10) that

$$r(d - g + 1) = 6 - q \leq 6,$$

since $h^0(\mathcal{O}_{\mathcal{L}}) \neq 0$. But $d - g + 1 \geq 1$, which implies that $r = 1$. Therefore

$$g \leq d - g + 1 = 6 - q \leq 6,$$

which implies that $g = 3$ by Lemma 2.16.

If $S \not\subset F$, then $|F \cap S| \leq 21$ by Lemma 6.2, which contradicts Lemma 2.15.

We see that $S \subset F$. But $q \leq 6$, so that $q = 1$ by Lemmas 4.5. Then $d = 7$.

There is a natural faithful action of the group G on the curve S such that every G -invariant divisor on S has even degree by Theorem 2.18. Hence d is even, which is a contradiction. \square

In particular, we see that the cases **(B)** and **(C)** are impossible.

Lemma 6.12. Suppose that $\deg(S) = 1$. Then $r \neq 7$.

Proof. Suppose that $r = 7$. Then $q = 0$ by (6.10). In particular, we have $S \not\subset F$. Then

$$|F \cap Z| \leq F \cdot Z = 7,$$

which contradicts Theorem 4.8, because $F \cap Z$ is G -invariant. \square

Lemma 6.13. The equality $r = 1$ holds.

Proof. Suppose that $r \geq 2$. Then $r \geq 7$ by Corollary B.2. If $r \geq 8$, then $d - g + 1 = 1$ by (6.10), which implies that $g = d \geq 2g - 1$, which leads to a contradiction.

We see that $r = 7$, so that $1 \leq d - g + 1 \leq 2$ by (6.10). Hence

$$d - g + 1 = 2,$$

since the equality $d - g + 1 = 1$ leads to a contradiction. Therefore, we have $g = d - 1 \geq 2g - 2$, which gives $g \leq 2$ and $d \leq 3$. Therefore $g = 0$ and $d = 1$, which is impossible by Lemma 6.12. \square

Thus, there is a natural faithful action of the group G on the curve S . We have

$$d = 13 + g - q \geq 2g - 1,$$

which implies that $q \leq 14 - g$. Note that $g \leq 14 - q \leq 14$. Then $g \in \{3, 8, 10\}$ by Lemma 2.16.

Lemma 6.14. The curve S is contained in the surface F .

Proof. If $S \not\subset F$, then $|F \cap S| \leq 21$ by Lemma 6.2, which is impossible by Lemma 2.15. \square

Applying Lemmas 4.5 and 6.14, we see that $q \in \{1, 7, 8\}$.

Lemma 6.15. The equality $g = 3$ holds.

Proof. Suppose that $g \neq 3$. Then $g \in \{8, 10\}$. It follows from (6.10) that

$$q \leq 14 - g \leq 6,$$

which implies that $q = 1$. Then $d \in \{20, 22\}$, which is impossible by Lemmas 2.19 and 2.20. \square

Thus, we have $d = 16 - q$, where $q \in \{1, 7, 8\}$.

Lemma 6.16. The equality $d = 8$ holds.

Proof. By Theorem 2.18, there is a G -invariant line bundle $\theta \in \text{Pic}(S)$ of degree 2 such that

$$\text{Pic}^G(S) = \langle \theta \rangle,$$

which implies that d is even. But $d = 16 - q$ and $q \in \{1, 7, 8\}$, so that $d = 8$. \square

Let Q be a surface in the threefold X that is swept out by lines. Then

$$Q \sim -2K_X$$

and the surface Q is irreducible by Lemma 4.7. Note that Q is G -invariant.

Lemma 6.17. The curve S is contained in $Q \cap F$.

Proof. By Lemma 6.14, we have $S \subset F$. If $S \not\subset Q$, then

$$|Q \cap S| \leq 2\deg(S) = 16$$

which is impossible by Lemma 2.15. Thus, we see that $S \subset Q \cap F$. \square

The surface F is a smooth $K3$ surface by Lemma 4.10. Then $S \cdot S = 0$ on the surface X . Put

$$Q|_F = mS + \Delta,$$

where $m \in \mathbb{Z}_{>0}$, and Δ is curve such that $S \not\subset \text{Supp}(\Delta)$. Then

$$(6.18) \quad 16 = 2\deg(S) = (mS + \Delta) \cdot S = mS \cdot S + \Delta \cdot S = 4m + \Delta \cdot S,$$

which implies that $|\Delta \cap S| \leq 12$. Hence, it follows from Lemma 2.15 that

$$\text{Supp}(\Delta) \cap S = \emptyset,$$

since Δ is G -invariant. Then $\Delta = \emptyset$, because $\text{Supp}(\Delta) \cup S$ is connected (see [14, Corollary 7.9]).

It follows from (6.18) that $m = 4$, which immediately leads to a contradiction, since

$$44 = Q \cdot F \cdot H = 4H \cdot S = 32,$$

where H is a general surface in $|-K_X|$.

The obtained contradiction completes the proof of Theorem 1.13.

In this section we describe Mukai's constructions of prime Fano threefolds of degree 22. Let $F(x, y, z)$ be a quartic form, let C be a curve in \mathbb{P}^2 defined by

$$F(x, y, z) = 0 \subset \mathbb{P}^2 \cong \text{Proj}(\mathbb{C}[x, y, z]),$$

and let $L_1, L_2, L_3, L_4, L_5, L_6$ be lines in \mathbb{P}^2 .

Definition A.1. We say that $\sum_{i=1}^6 L_i$ is a hexagon in \mathbb{P}^2 .

Let $L_i(x, y, z)$ is a linear form such that the line L_i is given by $L_i(x, y, z) = 0$.

Definition A.2 ([10, Definition 4.1]). The hexagon $\sum_{i=1}^6 L_i$ is polar to the curve C if

$$F(x, y, z) = \sum_{i=1}^6 L_i^4(x, y, z).$$

Consider $\sum_{i=1}^6 L_i$ as an element of the Hilbert scheme of points in the dual plane $\check{\mathbb{P}}^2$.

Definition A.3 ([27], [28], [32]). The variety of polar hexagons to the curve C is

$$\text{VSP}(C, 6) = \overline{\left\{ \Gamma \in \text{Hilb}_6(\check{\mathbb{P}}^2) \mid \Gamma \text{ is polar to the curve } C \right\}} \subset \text{Hilb}_6(\check{\mathbb{P}}^2).$$

Let $H(x, y, z)$ be the Hessian form of the form $F(x, y, z)$. Put $X = \text{VSP}(C, 6)$.

Theorem A.4 ([27, Theorem 5], [28, Theorem 11]). Suppose that $H(x, y, z) \neq 0$. Then

- the variety X is a normal threefold,
- the divisor $-K_X$ is an ample Cartier divisor such that $(-K_X)^3 = 22$,
- if no polar hexagon to C is a complete quadrangle (see [10, Definition 6.11]), then

$$\text{Pic}(X) = \mathbb{Z}[-K_X]$$

and X is smooth.

Suppose that $H(x, y, z) \neq 0$ (cf. [10, Theorem 6.12.2]). Put $W = \text{Spec}(\mathbb{C}[x, y, z]) \cong \mathbb{C}^3$. Then

$$F(x, y, z) \in \text{Sym}^4(W^\vee),$$

and the partial derivatives of the form $F(x, y, z)$ give an embedding $\phi: W \rightarrow \text{Sym}^3(W^\vee)$.

Suppose that C is not degenerate (see [10, Definition 2.8]). Then every hexagon $\Gamma \in X$ defines a six-dimensional subspace $W_\Gamma \subset \text{Sym}^3(W^\vee)$ such that $\phi(W) \subset W_\Gamma$. This gives a rational map

$$X \dashrightarrow \text{Gr}(3, U_7),$$

where $U_7 \cong \text{Sym}^3(W^\vee)/W$. The constructed rational map $X \dashrightarrow \text{Gr}(3, U_7)$ can be extended to an embedding $X \hookrightarrow \text{Gr}(3, U_7)$. This gives an embedding $X \hookrightarrow \mathbb{P}(\Lambda^3(U_7))$.

There is a natural sequence of maps

$$\begin{aligned} \Lambda^2(W) \otimes \text{Sym}^4(W) &\xrightarrow{\alpha} (W \otimes W) \otimes (\text{Sym}^2(W) \otimes \text{Sym}^2(W)) \xrightarrow{\beta} \\ &\xrightarrow{\beta} \text{Sym}^3(W) \otimes \text{Sym}^3(W) \xrightarrow{\gamma} \Lambda^2(\text{Sym}^3(W)), \end{aligned}$$

and it follows from [10, Section 2.3] that the quartic form $F(x, y, z)$ defines a natural map

$$\text{Sym}^2(W) \xrightarrow{\delta_F} \text{Sym}^2(W^\vee).$$

Since the quartic C is not degenerate, the map δ_F is invertible. Therefore, there is a natural choice of a non-zero element $\xi\delta_F^{-1} \in \text{Sym}^4(W)$ via the natural map

$$\text{Hom}\left(\text{Sym}^2(W^\vee), \text{Sym}^2(W)\right) \cong \text{Sym}^2(W) \otimes \text{Sym}^2(W) \xrightarrow{\xi} \text{Sym}^4(W),$$

which implies that the composition $\gamma \circ \beta \circ \alpha$ gives a map

$$W^\vee \cong \Lambda^2(W) \xrightarrow{\zeta} \Lambda^2(\text{Sym}^3(W)).$$

Lemma A.5. Let ω be an element in $\text{im}(\zeta)$ considered as a skew form on

$$\text{Sym}^3(W^\vee),$$

and let $\Pi \subset \text{Sym}^3(W^\vee)$ be the kernel of the form ω . Then $\text{im}(\phi) \subset \Pi$.

Proof. This is a straightforward computation. □

By Lemma A.5, the map ζ gives us a map $W^\vee \rightarrow \Lambda^2(U_7^\vee)$. Therefore, one has

$$W^\vee \otimes U_7^\vee \xrightarrow{\sigma} \Lambda^2(U_7^\vee) \otimes U_7^\vee \xrightarrow{\nu} \Lambda^3(U_7^\vee)$$

so that $\nu \circ \sigma$ is a monomorphism. Put

$$U_{14} = \Lambda^3(U_7) / (W^\vee \otimes U_7^\vee) \cong \mathbb{C}^{14},$$

and consider $\text{im}(\nu \circ \sigma)$ as a 21-dimensional linear system of hyperplanes in $\mathbb{P}(\Lambda^3(U_7))$ vanishing on the image of the threefold X . This gives us a natural embedding $X \hookrightarrow \mathbb{P}(U_{14})$.

Theorem A.6 (cf. [27], [28], [32]). The embedding $X \hookrightarrow \mathbb{P}(U_{14})$ is the anticanonical embedding.

Note that X is smooth by Theorem A.4 in the case when C is sufficiently general.

APPENDIX B. REPRESENTATION THEORY

In this section we collect some facts about the groups $\text{PSL}_2(\mathbb{F}_7)$ and $\text{SL}_2(\mathbb{F}_7)$.

Lemma B.1 ([6]). Let Γ be a maximal subgroup in $\text{PSL}_2(\mathbb{F}_7)$. Then

- either $\Gamma \cong \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ and Γ is unique up to a conjugation,
- or $\Gamma \cong \text{S}_4$ and $\text{PSL}_2(\mathbb{F}_7)$ contains two subgroups isomorphic to Γ up to a conjugation.

Corollary B.2. If $\text{PSL}_2(\mathbb{F}_7)$ acts transitively on a finite set Σ such that $|\Sigma| \leq 41$, then

$$|\Sigma| \in \{1, 7, 8, 14, 21, 24, 28\}.$$

The group $\text{PSL}_2(\mathbb{F}_7)$ has exactly six non-isomorphic irreducible representations (see [6]), which we denote by $I, W_3, W_3^\vee, W_6, W_7, W_8$. The values of their characters are listed in the table:

	id	(2)	(3)	(4)	(7)	(7')
‡	1	21	56	42	24	24
I	1	1	1	1	1	1
W_3	3	-1	0	1	ϵ	$\bar{\epsilon}$
W_3^\vee	3	-1	0	1	$\bar{\epsilon}$	ϵ
W_6	6	2	0	0	-1	-1
W_7	7	-1	1	-1	0	0
W_8	8	0	-1	0	1	1

where we use the following notation:

- the first row represents the conjugacy classes in the group $\text{PSL}_2(\mathbb{F}_7)$ such that
 - the symbol id denotes the identity element,
 - the symbol (n) denotes a class of elements of order n ,

- the symbols (7) and (7') denote two classes of elements of order 7 such that
 - * if $g \in (7)$, then $g^2 \in (7)$ and $g^4 \in (7)$,
 - * if $g \in (7)$, then $g^3 \in (7')$, $g^5 \in (7')$ and $g^6 \in (7')$,
- the second row lists the number of elements in each conjugacy class,
- the next six rows list the values of the characters of irreducible representations,
- the symbol ϵ denotes the complex number $-1/2 + \sqrt{-7}/2$,
- the symbol $\bar{\epsilon}$ denotes the complex conjugate of the number ϵ .

Looking at the above table, one easily obtains the following corollaries.

Corollary B.3. Let Γ be a subgroup of the group $\mathrm{PSL}_2(\mathbb{F}_7)$ such that $\Gamma \cong A_4$. Then

- the representation W_3 is an irreducible Γ -representation,
- the representation W_6 is a sum of two irreducible three-dimensional Γ -representations.

Corollary B.4. The following isomorphisms of the representations of the group $\mathrm{PSL}_2(\mathbb{F}_7)$ hold:

$$\Lambda^2(W_3^\vee) \cong W_3, \quad \Lambda^4(W_7) \cong \Lambda^3(W_7)^\vee \cong \Lambda^3(W_7), \quad \mathrm{Sym}^3(W_3^\vee) \cong W_7 \oplus W_3,$$

$$W_7 \otimes W_3^\vee \cong W_6 \oplus W_7 \oplus W_8, \quad \Lambda^3(W_7) \cong I \oplus W_6 \oplus W_7 \oplus W_6 \oplus W_7 \oplus W_8.$$

Let ζ be a primitive seventh root of unity, let \hat{G} be a subgroup in $\mathrm{SL}_3(\mathbb{C})$ such that

$$\hat{G} = \left\langle \left(\begin{array}{ccc} \zeta & 0 & 0 \\ 0 & \zeta^2 & 0 \\ 0 & 0 & \zeta^4 \end{array} \right), \left(\begin{array}{ccc} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{array} \right), \frac{1}{\sqrt{-1}} \left(\begin{array}{ccc} \zeta^4 - \zeta^3 & \zeta^2 - \zeta^5 & \zeta - \zeta^5 \\ \zeta^2 - \zeta^5 & \zeta - \zeta^6 & \zeta^4 - \zeta^3 \\ \zeta - \zeta^6 & \zeta^4 - \zeta^3 & \zeta^2 - \zeta^5 \end{array} \right) \right\rangle,$$

and let $\phi: \mathrm{SL}_3(\mathbb{C}) \rightarrow \mathrm{Aut}(\mathbb{P}^2)$ be a natural projection. Put $G = \phi(\hat{G})$. Then $G \cong \hat{G} \cong \mathrm{PSL}_2(\mathbb{F}_7)$.

Lemma B.5 ([37, Section 2.10]). Let C be a G -invariant curve in \mathbb{P}^2 . Then

- $\deg(C) \notin \{1, 2, 3, 5\}$,
- if $\deg(C) = 4$, then C is the quartic curve described in Example 1.10,
- if $\deg(C) = 6$, then C is the Hessian curve of the quartic curve described in Example 1.10.

Lemma B.6. There are no G -invariant subsets in \mathbb{P}^2 consisting of at most 20 points.

Proof. Restricting W_3 to subgroups of the group $\mathrm{PSL}_2(\mathbb{F}_7)$, we obtain the required assertion, because we may assume that W_3 is given by the monomorphism $\hat{G} \hookrightarrow \mathrm{SL}_3(\mathbb{C})$. \square

Let Γ be a subgroup in $\mathrm{SL}_2(\mathbb{F}_7)$, let $\pi: \mathrm{SL}_2(\mathbb{F}_7) \rightarrow \mathrm{PSL}_2(\mathbb{F}_7)$ be a natural epimorphism. Then

$$\Gamma \cong 2 \cdot \pi(\Gamma)$$

if $\pi(\Gamma)$ is isomorphic to $\mathrm{PSL}_2(\mathbb{F}_7)$, S_4 , A_4 , $\mathbb{Z}_7 \rtimes \mathbb{Z}_3$, D_4 , or S_3 .

The following table contains the character values of one four-dimensional and one eight-dimensional irreducible representations of the group $\mathrm{SL}_2(7)$ and some information about its subgroups:

	id	-id	(3) ₃	(3) ₆	(7) ₇	(7) ₇	(7) ₁₄	(7) ₁₄	(2) ₄	(4) ₈	(4) ₈
G	1	1	56	56	24	24	24	24	42	42	42
$2 \cdot S_4$	1	1	8	8	0	0	0	0	18	6	6
$2 \cdot A_4$	1	1	8	8	0	0	0	0	6	0	0
$2 \cdot (\mathbb{Z}_7 \rtimes \mathbb{Z}_3)$	1	1	14	14	3	3	3	3	0	0	0
$2 \cdot D_4$	1	1	0	0	0	0	0	0	10	2	2
$2 \cdot S_3$	1	1	2	2	0	0	0	0	6	0	0
U_4	4	-4	1	-1	$\bar{\alpha}$	α	$-\alpha$	$-\bar{\alpha}$	0	0	0
U_8	8	-8	-1	1	1	1	-1	-1	0	0	0

where we use the following notations:

- the first row represents the conjugacy classes in $\mathrm{SL}_2(7)$ such that

- the symbol id denotes the identity element,
- the symbol $-\text{id}$ denotes the element different from id such that $\pi(-\text{id}) \in \ker(\pi)$,
- the symbol $(n)_k$ denotes elements of order k such that $\pi((n)_k)$ have order n ,
- the next six rows list the number of conjugacy class elements that lie in the corresponding subgroup of the group $\text{SL}_2(7)$ isomorphic to a group listed in the first cell of the row,
- the next two rows list the values of the characters of two irreducible representations,
- the symbol α denotes the complex number $-(\zeta^3 + \zeta^5 + \zeta^6)$,
- the symbol $\bar{\alpha}$ denotes the complex conjugate of the number α .

Lemma B.7. Suppose that $\Gamma \cong 2.(\mathbb{Z}_7 \rtimes \mathbb{Z}_3)$. Then $U_4 \cong T \oplus J$ and

$$U_8 \cong T \oplus T_1 \oplus J_1 \oplus J_2$$

as representations of the group Γ , where J , J_1 and J_2 are pairwise non-isomorphic one-dimensional representations, while T and T_1 are irreducible three-dimensional representations.

Proof. Let χ_4 and χ_8 be the characters of the representations U_4 and U_8 , respectively. Then

$$\langle \chi_4, \chi_4 \rangle = 2,$$

which immediately implies that $U_4 \cong J \oplus T$ for some one-dimensional representations J and some irreducible three-dimensional representation T of the group Γ , because irreducible representations of the group Γ are either one-dimensional or three-dimensional. Similarly, we have

$$\langle \chi_8, \chi_8 \rangle = 4,$$

which implies that $U_8 \cong J_1 \oplus J_2 \oplus T_1 \oplus T_2$ for some one-dimensional representations $J_1 \not\cong J_2$ and some irreducible three-dimensional representations $T_1 \not\cong T_2$ of the group Γ .

We may assume that $T_2 \cong T$, because there exist exactly two three-dimensional representations of the group Γ with a non-trivial action of its center. But

$$\langle \chi_4, \chi_8 \rangle = 1,$$

which implies that that neither J_1 nor J_2 is isomorphic to J . □

Note that we can consider $\text{PSL}_2(\mathbb{F}_7)$ -representations also as $\text{SL}_2(\mathbb{F}_7)$ -representations.

Lemma B.8. One has $\text{Sym}^2(U_6) \cong W_6$ and

$$\text{Sym}^4(U_6) \cong I \oplus W_6 \oplus W_6 \oplus W_7 \oplus W_7 \oplus W_8$$

as representations of the group $\text{PSL}_2(\mathbb{F}_7)$ or the group $\text{SL}_2(\mathbb{F}_7)$.

Proof. This follows from elementary and explicit computations (see also [8, Appendix 1]). □

Lemma B.9. As a representation of the group Γ , the representation U_4 splits as

- a sum of two irreducible two-dimensional subrepresentations if $\Gamma \cong 2.S_4$,
- a sum of two irreducible two-dimensional subrepresentations if $\Gamma \cong 2.A_4$,
- a sum of two irreducible two-dimensional subrepresentations if $\Gamma \cong 2.D_4$,
- a sum of an irreducible two-dimensional and two non-isomorphic one-dimensional subrepresentations if $\Gamma \cong 2.S_3$.

Proof. Let χ_4 be the character of the representations U_4 . If $\Gamma \cong 2.D_4$, then

$$\langle \chi_4, \chi_4 \rangle = 2,$$

which easily implies that U_4 splits as a sum of two irreducible two-dimensional subrepresentations, because $2.D_4$ has no odd-dimensional non-trivial irreducible representations.

Since all irreducible representations of the group $2.A_4$ with a non-trivial action of its center are two-dimensional, the representation U_4 splits as a sum of two irreducible two-dimensional subrepresentations of the group Γ if $\Gamma \cong 2.A_4$ or $\Gamma \cong 2.S_4$.

To complete the proof, we may assume that $\Gamma \cong 2.S_3$. Then there is an epimorphism $\Gamma \rightarrow \mathbb{Z}_4$.

Let U be the standard unitary two-dimensional irreducible representation of the group Γ , let J and J_1 be one-dimensional representations of the group Γ that arise from the faithful non-isomorphic one-dimensional representations of the group \mathbb{Z}_4 . Then

$$U_4 \cong U \oplus J \oplus J_1$$

as can be seen from the following table that lists the character values of these representations:

	id	-id	$(3)_3$	$(3)_6$	$(2)_4$
$2.S_3$	1	1	2	2	6
U_4	4	-4	1	-1	0
U	2	-2	-1	1	0
J	4	-4	1	-1	$\sqrt{-1}$
J_1	4	-4	1	-1	$-\sqrt{-1}$

where we used notation similar to the ones used in the table above. □

REFERENCES

- [1] H. Blichfeldt, *Finite collineation groups*
University of Chicago Press, Chicago, 1917
- [2] I. Cheltsov, *Two local inequalities*
Russian Mathematical Surveys, to appear
- [3] I. Cheltsov, C. Shramov, *Log canonical thresholds of smooth Fano threefolds*
Russian Mathematical Surveys **63** (2008), 73–180
- [4] I. Cheltsov, C. Shramov, *On exceptional quotient singularities*
arXiv:math/0909.0918 (2009)
- [5] I. Cheltsov, C. Shramov, *Five embeddings of one simple group*
arXiv:math/0910.1783 (2009)
- [6] J. Conway, R. Curtis, S. Norton, R. Parker, R. Wilson, *Atlas of finite groups*
Clarendon Press, Oxford, 1985
- [7] A. Corti, *Singularities of linear systems and threefold birational geometry*
L.M.S. Lecture Note Series **281** (2000), 259–312
- [8] I. Dolgachev, *Invariant stable bundles over modular curves $X(p)$*
Contemporary Mathematics **224** (1999), 65–99
- [9] I. Dolgachev, V. Iskovskikh, *Finite subgroups of the plane Cremona group*
Birkhauser Boston, Progress in Mathematics **269** (2009), 443–548
- [10] I. Dolgachev, V. Kanev, *Polar covariants of plane cubics and quartics*
Advances in Mathematics **98** (1993), 216–301
- [11] S. Donaldson, *Kähler geometry on toric manifolds, and some other manifolds with large symmetry*
Handbook of geometric analysis, Advanced Lectures in Mathematics **1** (2008), 29–75
- [12] W. Edge, *The Klein group in three dimensions*
Acta Mathematica **79** (1947), 153–223
- [13] D. Eisenbud, J.-H. Koh, *Remarks on points in a projective space*
MSRI Publications **15**, Springer, New York, 157–172
- [14] R. Hartshorne, *Algebraic geometry*
Graduate Texts in Mathematics **52** (1977) Springer-Verlag, New York-Heidelberg
- [15] M. Gross, S. Popescu, *Calabi-Yau threefolds and moduli of abelian surfaces I*
Compositio Mathematica **127** (2001), 169–228
- [16] V. Iskovskikh, Yu. Prokhorov, *Fano varieties*
Encyclopaedia of Mathematical Sciences **47** (1999) Springer, Berlin

- [17] Y. Kawamata, *On Fujita's freeness conjecture for 3-folds and 4-folds*
Mathematische Annalen **308** (1997), 491–505
- [18] Y. Kawamata, *Subadjunction of log canonical divisors II*
American Journal of Mathematics **120** (1998), 893–899
- [19] J. Kollár, *Singularities of pairs*
Proceedings of Symposia in Pure Mathematics **62** (1997), 221–287
- [20] R. Lazarsfeld, *Positivity in algebraic geometry II*
Springer-Verlag, Berlin, 2004
- [21] S. Levy (editor), *The eightfold way. The beauty of Klein's quartic curve*,
MSRI Publications **35** (1999), Cambridge University Press
- [22] C. Mallows, N. Sloane, *On the invariants of a linear group of order 336*
Proceedings of the Cambridge Philosophical Society **74** (1973), 435–440
- [23] N. Manolache, F.-O. Schreyer, *Moduli of $(1, 7)$ -polarized abelian surfaces via syzygies*
Mathematische Nachrichten **226** (2001), 177–203
- [24] A. Marini, *On a family of $(1, 7)$ -polarised abelian surfaces*
Mathematica Scandinavica **95** (2004), 181–225
- [25] F. Melliez, K. Ranestad, *Degenerations of $(1, 7)$ -polarized abelian surfaces*
Mathematica Scandinavica **97** (2005), 161–187
- [26] S. Mukai, *Finite groups of automorphisms of K3 surfaces and the Mathieu group*
Inventiones Mathematicae **94** (1988), 183–221
- [27] S. Mukai, *Biregular classification of Fano 3-folds and Fano manifolds of coindex 3*
Proceedings of the National Academy of Sciences of the United States of America **9** (1989), 3000–3002
- [28] S. Mukai, *Fano 3-folds*
London Mathematical Society Lecture Note Series **179** (1992), 255–263
- [29] K. Oguiso, D.-Q. Zhang, *The simple group of order 168 and K3 surfaces*
Complex geometry, Gottingen (2002), 165–184
- [30] Yu. Prokhorov, *Exotic Fano varieties*
Moscow University Mathematics Bulletin **45** (1990), 36–38
- [31] Yu. Prokhorov, *Simple finite subgroups of the Cremona group of rank 3*
arXiv:math/0908.0678 (2009)
- [32] F.-O. Schreyer, *Geometry and algebra of prime Fano 3-folds of genus 12*
Compositio Mathematica **127** (2001), 297–319
- [33] T. Springer, *Invariant theory*
Lecture Notes in Mathematics **585** (1977), Springer-Verlag, Berlin-New York
- [34] G. Tian, *On Kähler–Einstein metrics on certain Kähler manifolds with $c_1(M) > 0$*
Inventiones Mathematicae **89** (1987), 225–246
- [35] G. Tian, *Kähler–Einstein metrics with positive scalar curvature*
Inventiones Mathematicae **130** (1997), 1–37
- [36] G. Xiao, *Galois covers between K3 surfaces*
Annales de l'Institut Fourier **46** (1996), 73–88
- [37] S. Yau, Y. Yu, *Gorenstein quotient singularities in dimension three*
Memoirs of the American Mathematical Society **505** (1993), Providence

UNIVERSITY OF EDINBURGH, EDINBURGH EH9 3JZ, UK, I.CHELTSOV@ED.AC.UK

STEKLOV INSTITUTE OF MATHEMATICS, MOSCOW 119991, RUSSIA, SHRAMOV@MCCME.RU