

# Analytic combinatorics of the transfinite: generalized Mahler partitions and natural Gödel numberings for ordinals

Andreas Weiermann  
Vakgroep Zuivere Wiskunde en Computeralgebra  
Krijgslaan 281 Gebouw S22  
9000 Ghent  
Belgium  
e-mail: weierman@math.uu.nl

## Abstract

We study the asymptotic of certain count functions which are connected with a certain well-ordered subset of Hardy's orders of infinity. As special cases we find Mahler partitions and its iterated versions as well as their multiplicative counterparts.

## 1 Introduction

We determine the asymptotics of certain count functions for a natural subclass of Hardy's order of infinity. These results find applications in classifying phase transitions for Gödel incompleteness and logical limit laws. The class of functions  $\mathbb{E}$  in question is defined as follows. Each member  $f \in \mathbb{E}$  is a function from  $\mathbb{N} \rightarrow \mathbb{N}$ . Let  $\alpha_0(x) := 0$  be the function constant zero defined for non negative integers. We put  $\alpha_0 \in \mathbb{E}$ . Now assume that  $f, g$  are elements of  $\mathbb{E}$ . Let  $h(x) := x^{f(x)} + g(x)$ . Then  $h$  is put into  $\mathbb{E}$ . Now we define  $\mathbb{E}$  to be the least set containing  $\alpha_0$  which is closed under this formation rule.

The set  $\mathbb{E}$  comes equipped with a natural order  $\prec$  of eventual domination which is defined as follows:  $f \prec g$  iff there is a  $k \in \mathbb{N}$  such that  $f(x) < g(x)$  for all  $x > k$ . Then e.g. Hardy [?] has shown that the ordering  $\prec$  is a linear ordering on  $\mathbb{E}$ . Moreover there will not exist any infinite descending chain  $f_0 \succ f_1 \succ f_2 \succ \dots$  of elements in  $\mathbb{E}$ , i.e.  $\prec$  is a well-ordering on  $\mathbb{E}$ . Moreover the well-orderedness of  $\prec$  with respect to arithmetical sets cannot be proved from the Peano axioms for the natural numbers.

There are several canonical complexity measures which can be assigned to members of  $\mathbb{E}$ . The desired property of such a measure  $c : \mathbb{E} \rightarrow \mathbb{N}$  is that for any  $k \in \mathbb{N}$  and any  $f \in \mathbb{E}$  the number of elements in  $\{g \prec f : c(g) \leq k\}$  is finite. A canonical choice for  $c$  is given by evaluation. We may put  $c(g) := g(k)$  for some fixed  $k$ . In this case we put

$$M_f^k(n) := \#\{g \prec f : g(k) = n\} \quad (1)$$

and one may ask for the asymptotic of  $M_f^k(n)$  as  $n \rightarrow \infty$ . For a certain choice of  $f$  there is much information on this problem available from the literature about Mahler partitions. Let  $f(x) = x^x$  (which is an element from  $\mathbb{E}$  since  $f(x) = x^{x^{c_0(x)} + c_0(x)} + c_0(x)$ ). Then  $M_f^k(n)$  is the number of Mahler partitions of  $n$  in sums of exponentials with base  $k$ . For example it is well known that

$$\ln(M_f^k(n)) \sim \frac{1}{2\ln(k)}(\ln(n))^2$$

as  $n \rightarrow \infty$ . In the sequel we stick to the case  $k = 2$  when we consider this type of investigation. The case for general  $k$  means that one has to replace in the corresponding results  $\ln(2)$  by  $\ln(k)$ . Therefore we drop in the sequel the upper index in  $m_f^k$  and assume that this index is equal to 2.

Another natural complexity functions emerges from a natural Gödel numbering of  $\mathbb{E}$ . The idea is to assign to each object  $f$  in  $\mathbb{E}$  a unique natural number such that effective operations on  $\mathbb{E}$  translate into elementary recursive operations on the corresponding Gödel numbers. To this end note that every non constant zero element  $f$  in  $\mathbb{E}$  has a unique representation  $f = id^{f_1} + \dots + id^{f_d}$  with  $f_1 \succeq \dots \succeq f_d$ . We further let  $(p_i)_{i=1}^\infty$  denote the enumeration of the primes starting with  $p_1 = 2$ .

We put  $[c_0] := 1$  and if  $f \in \mathbb{E}$  has the representation  $f = id^{f_1} + \dots + id^{f_d}$  with  $f_1 \succeq \dots \succeq f_d$  then we put  $[f] := p_1^{[f_1]} \cdot \dots \cdot p_d^{[f_d]}$ . There are different choices of the Gödel numbering possible but we have chosen one which typically appears in textbooks on recursion theory. We put

$$G_f(n) := \#\{g \prec f : [g] \leq n\}. \quad (2)$$

For  $f = id^{id}$  we get a multiplicative analogue of the Mahler partition function as we will see in a minute. Getting non trivial bounds on  $G_f$  seems even more diffi cult then for  $M_f$ . But luckily large machinery from analytic combinatorics has already been developed and a seminal paper by Parameswaran allows to obtain weak asymptotics for  $M_f$  as well as for  $G_f$ . These results are strong enough for the intended proof theoretic applications. It seems that even better bounds are available by applying the saddle point method a la Dumas and Flajolet [5] but we leave this for the experts in the field. Parameswaran's result is as follows.

**Theorem 1 (Parameswaran [8]).** *Suppose that the following conditions hold.*

1.  $L(u)$  and  $P(u)$  are functions on the non negative reals such that  $\int_0^R L(u)du$  and  $\int_0^R P(u)du$  exist in the Lebesgue sense for every positive  $R$ .
2.  $\exp(s \int_0^\infty \frac{e^{-su}}{1-e^{-su}} L(u)du) = s \int_0^\infty P(u)e^{-su} du$  for all positive  $s$ ,
3.  $\langle M, M^* \rangle$  form a pair of conjugate slowly varying functions,
4.  $M$  is non decreasing,
5.  $\int_0^u \frac{L(t)}{t} dt \sim M(u)$  as  $u \rightarrow \infty$ , and
6.  $P(u)$  is non decreasing.

Then  $\log P(u) \sim \frac{1}{M^*(u)}$  as  $u \rightarrow \infty$ .

We now state our main results. For a compact presentation we use the following notations. We put

$$\ln_{d+1}(x) := \ln(\ln_d(x))$$

where  $\ln_1(x) = \ln(x)$ . Moreover we put  $id(x) := x$  and let

$$id_{d+1}(k)(x) := x^{id_d(k)(x)}$$

where  $id_1(k)(x) = x^k$ . In addition we put

$$\exp_{d+1}(x) := \exp(\exp_d(x))$$

where  $\exp_1(x) = \exp(x) = \sum_{i=0}^{\infty} \frac{x^i}{i!}$ .

**Theorem 2.** 1. If  $f = id_1(k)$  then there exist explicitly calculable constants  $C_1, C_2$  such that

$$M_f(x) \sim C_1 \cdot x^{k-1} \quad (3)$$

$$G_f(x) \sim C_1 \cdot \left(\frac{\ln(x)}{\ln(\ln(x))}\right)^k \quad (4)$$

2. If  $f = id_2(k)$  then there exist explicitly calculable constants  $C_3, C_4$  such that

$$\ln(M_f(x)) \sim C_3 \cdot (\ln(x))^{k+1} \quad (5)$$

$$\ln(G_f(x)) \sim C_4 \cdot \ln(\ln(x)) \cdot \left(\frac{\ln \ln(x)}{\ln(\ln(\ln(x)))}\right)^k \quad (6)$$

3. If  $f = id_d(k)$  and  $d \geq 2$  then with the same constants  $C_3, C_4$  as in the previous item

$$\ln_{d-1}(M_f(x)) \sim C_3 \cdot (\ln_{d-1}(x))^{k+1} \quad (7)$$

$$\ln_{d-1}(G_f(x)) \sim C_4 \cdot \ln_d(x) \cdot \left(\frac{\ln_d(x)}{\ln(\ln_d(x))}\right)^k \quad (8)$$

*Proof.* We prove the results for the generalized Mahler norms. The asymptotic (3) is well known. Indeed we may consider  $\{g \prec id^k\}$  as a generalized additive number system generated from the additive primes  $id^l$  for  $0 \leq l < k$ . By Theorem 2.48 in Burris [4] we therefore obtain

$$M_{id^k}(x) \sim \frac{1}{(k-1)!} \frac{1}{\prod_{l < k} 2^l} x^{k-1} \quad (9)$$

and assertion (3) follows.

Let us now prove assertion (5). By remark 2.32 and Theorem 2.48 in Burris [4] we obtain

$$m(x) := \#\{g \prec id^k : g(2) \leq x\} \sim \frac{1}{k!} \frac{1}{\prod_{l < k} 2^l} x^k \quad (10)$$

Let

$$n(u) = \sum_{2^l \leq u} M_{id^k}(l) = m\left(\frac{\ln(u)}{\ln(2)}\right).$$

Then

$$n(u) \sim \frac{1}{k!(\prod_{l < k} 2^l)(\ln(2))^k} (\ln(u))^k =: L(u).$$

Let

$$C := \frac{1}{k!(\prod_{l < k} 2^l)(\ln(2))^k}.$$

Let

$$M(u) := \int_a^u \frac{L(t)}{t} dt$$

where  $a >$  is arbitrary but fixed. Then by de l'Hospital's rule

$$M(u) \sim \frac{C}{k+1} (\ln(u))^{k+1}.$$

Let

$$P(u) := \sum_{l \leq u} M_{id^k}(l).$$

By Theorem 1 of Parameswaran we obtain

$$\ln(P(u)) \sim \frac{C}{k+1} (\ln(u))^{k+1}.$$

Moreover this yields  $\ln(M_{id^k}(u)) \sim \frac{C}{k+1} (\ln(u))^{k+1}$  as indicated on the last page of Parameswaran. So we may put  $C_3 := \frac{C}{k+1}$ .

Let us now prove assertion 7 by induction on  $d$ . Put

$$m(x) := \#\{g < M_{id_d}(k) : g(2) \leq x\}. \quad (11)$$

Let

$$n(u) = \sum_{2^l \leq u} M_{id_d}(k)(l) = m\left(\frac{\ln(u)}{\ln(2)}\right) =: L(u).$$

The induction hypothesis yields

$$\log_{d-1}(m(x)) \sim C \cdot (\log_{d-1}(x))^{k+1}. \quad (12)$$

for  $C = C_3$ . Let

$$M(u) := \int_a^u \frac{L(t)}{t} dt$$

for some arbitrary fixed  $a > 0$  Let

$$P(u) := \sum_{l \leq u} M_{id_d}(k)(l).$$

By Theorem 1 of Parameswaran we obtain

$$\ln(P(u)) \sim M(u).$$

We claim that

$$\ln_d(P(u)) \sim C \cdot (\ln_d(x))^{k+1}.$$

Proof: Pick

$$\varepsilon > 0.$$

Then (12) yields

$$m(u) \leq \exp_{d-1}\left(\left(1 + \frac{\varepsilon}{2}\right)C(\ln_{d-1}(u))^k\right)$$

for large enough  $u$ . Hence

$$L(u) \leq \exp_{d-1}\left(\left(1 + \frac{\varepsilon}{2}\right)C(\ln_{d-1}\left(\frac{\ln(u)}{\ln(2)}\right))^{k+1}\right).$$

Thus

$$M(u) \leq \int_a^u \frac{\exp_{d-1}\left(\left(1 + \frac{\varepsilon}{2}\right)C(\ln_{d-1}\left(\frac{\ln(u)}{\ln(2)}\right))^{k+1}\right)}{u} du.$$

Put

$$N(u) := \exp_{d-1}\left(\left(1 + \varepsilon\right)C(\ln_{d-1}(\ln(u)))^{k+1}\right).$$

Then de l'Hospital's rule yields

$$M(u) = o(N(u))$$

as  $u \rightarrow \infty$ . In particular we obtain that  $M(u) \leq N(u)$  for large enough  $u$ . Therefore

$$\ln(P(u)) \sim M(u) \leq N(u)$$

for large enough  $u$ . Hence

$$P(u) \leq \exp_d\left(\left(1 + \varepsilon\right)C(\ln_d(u))^{k+1}\right).$$

By a similar argument we obtain

$$P(u) \geq \exp_d\left(\left(1 - \varepsilon\right)C(\ln_d(u))^{k+1}\right).$$

Thus

$$\log_d(P(u)) \sim C(\ln_d(u))^{k+1}.$$

Further

$$\ln_d(M_{id_{d+1}(k)}) \sim C(\ln_d(u))^{k+1}(u)$$

as indicated on the last page of Parameswaran. □

The intended proof-theoretic applications are as follows. Let small Greek letters range over elements of  $\mathbb{E}$ . With  $\omega$  we denote the identity function *id*. Elements in  $\mathbb{E}$  of the form  $\alpha + \omega^{c_0}$  are called successors. Non zero elements of  $\mathbb{E}$  which are not successors are called limits. For a limit  $\lambda \in \mathbb{E}$  let  $\lambda[x]$  be the  $x$ -th element of the canonical fundamental sequence for  $\lambda$ . This means that if  $\lambda = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$  where  $\alpha_1 \succeq \dots \succeq \alpha_n$  and  $\alpha_n = \gamma + \omega^{c_0}$  then  $\lambda[x] := \omega^{\alpha_1} + \dots + \omega^\gamma \cdot x$  and that if  $\lambda = \omega^{\alpha_1} + \dots + \omega^{\alpha_n}$  where  $\alpha_1 \succeq \dots \succeq \alpha_n$  and  $\alpha_n$  is a limit then  $\lambda[x] := \omega^{\alpha_1} + \dots + \omega^{\alpha_n[x]}$ . Then we have that for all  $\beta \prec \lambda$  there is an  $x$  such that  $\beta \prec \lambda[x]$  so that  $\lambda[x]$  converges to  $\lambda$  as  $x \rightarrow \infty$ . It is convenient to introduce a top element for the elements of  $\mathbb{E}$ . We call this  $\varepsilon_0$  and we write  $\alpha \prec \varepsilon_0$  in place of  $\alpha \in \mathbb{E}$ . For  $\varepsilon_0$  the fundamental sequence is defined via  $\varepsilon_0[x] := \omega_x = id_x$ . We then can define  $F_\alpha$  for  $\alpha \preceq \varepsilon_0$  as follows by recursion on  $\prec$ .

$$\begin{aligned} F_0(x) &:= x + 1, \\ F_{\alpha+1}(x) &:= F_\alpha^{(x)}(x) \text{ where the upper index denotes number of iterations,} \\ F_\lambda(x) &:= F_{\lambda[x]}(x) \text{ where } \lambda \text{ is a limit.} \end{aligned}$$

Let  $c$  be a complexity measure for the elements of  $\mathbb{E}$ . Let  $\text{SWO}(\beta, f, c)$  be the statement

$$(\forall K)(\exists L)(\forall \alpha_0, \dots, \alpha_L \prec \beta) ((\forall i \leq L)[c(\alpha_i) \leq K + f_\alpha(i)] \rightarrow (\exists i < L)[\alpha_i \preceq \alpha_{i+1}]).$$

**Theorem 3.** *Let*

$$f_\alpha(i) := \exp_{F_\alpha^{-1}(i)}(\sqrt{\ln_{F_\alpha^{-1}(i)}(i)}).$$

*Then the following phase transition result holds for  $\text{SWO}(\varepsilon_0, f, M)$ .*

1. *If  $\alpha \prec \varepsilon_0$  then*

$$\text{PA} \vdash \text{SWO}(\varepsilon_0, f_\alpha, M).$$
2. *If  $\alpha = \varepsilon_0$  then*

$$\text{PA} \not\vdash \text{SWO}(\varepsilon_0, f_\alpha, M).$$

Let  $\mathbb{I}\Sigma_d$  be the fragment of PA where the induction scheme is restricted to formulas with at most  $d$  quantifiers.

**Theorem 4.** *Let  $d \geq 1$ . Let*

$$f_\alpha(i) := \exp_d(\overset{F_\alpha^{-1}(i)}{\sqrt{\ln_d(i)}}).$$

*Then the following phase transition result holds for  $\text{SWO}(\omega_{d+1}, f, M)$ .*

1. *If  $\alpha \prec \omega_{d+1}$  then*

$$\mathbb{I}\Sigma_d \vdash \text{SWO}(\omega_{d+1}, f_\alpha, M).$$
2. *If  $\alpha = \omega_{d+1}$  then*

$$\mathbb{I}\Sigma_d \not\vdash \text{SWO}(\omega_{d+1}, f_\alpha, M).$$

In the multiplicative situation the following phase transition results are obtained.

**Theorem 5.** *Let*

$$f_\alpha(i) := \exp\left(\exp_{F_\alpha^{-1}(i)}\left(\sqrt{\ln_{F_\alpha^{-1}(i)}(i)}\right)\right).$$

*Then the following phase transition result holds for  $\text{SWO}(\varepsilon_0, f, [\cdot])$ .*

1. *If  $\alpha \prec \varepsilon_0$  then*

$$\text{PA} \vdash \text{SWO}(\varepsilon_0, f_\alpha, [\cdot]).$$

2. *If  $\alpha = \varepsilon_0$  then*

$$\text{PA} \not\vdash \text{SWO}(\varepsilon_0, f_\alpha, [\cdot]).$$

Let  $\mathbb{I}\Sigma_d$  be the fragment of PA where the induction scheme is restricted to formulas with at most  $d$  quantifiers.

**Theorem 6.** *Let  $d \geq 1$ . Let*

$$f_\alpha(i) := \exp\left(\exp_d\left({}^{\varepsilon_\alpha^{-1}(i)}\sqrt{\ln_d(i)}\right)\right).$$

*Then the following phase transition result holds for  $\text{SWO}(\omega_{d+1}, f, [\cdot])$ .*

1. *If  $\alpha \prec \omega_{d+1}$  then*

$$\mathbb{I}\Sigma_d \vdash \text{SWO}(\omega_{d+1}, f_\alpha, [\cdot]).$$

2. *If  $\alpha = \omega_{d+1}$  then*

$$\mathbb{I}\Sigma_d \not\vdash \text{SWO}(\omega_{d+1}, f_\alpha, [\cdot]).$$

In a sequel paper we will exploit our investigations to prove (joint project with A.R. Woods) logical limit laws for segments of  $\mathbb{E}$ . We plan to investigate further properties of  $M_f$  and  $G_f$  with J.P. Bell.

## References

- [1] Toshiyasu Arai: *On the slowly well-orderedness of  $\varepsilon_0$* . *Mathematical Logic Quarterly* 48 (2002) 125-139.
- [2] Nicholas H. Bingham, Charles M. Goldie and Jozef L. Teugels: *Regular variation*. *Encyclopedia of Mathematics and its applications* 27. Cambridge University Press, 1989.
- [3] N.G. de Bruijn: *Pairs of slowly oscillating functions occurring in asymptotic problems concerning the Laplace transform*, *Nieuw Arch. Wiskunde* 7 (19659), 20-26.
- [4] Stanley N. Burris: *Number Theoretic Density and Logical Limit Laws*. *Mathematical Surveys and Monographs* 86. American Mathematical Society 2001.
- [5] P. Dumas, P. Flajolet, Philippe: *Asymptotique des recurrences mahleriennes: le cas cyclotomique*. *J. Theor. Nombres Bordeaux* 8 (1996), no. 1, 1–30.

- [6] Harvey Friedman and Michael Sheard: *Elementary descent recursion and proof theory*. Annals of Pure and Applied Logic 71 (1995), 1-45.
- [7] G.H. Hardy: *Orders of infinity. The Infinitärcalcul of Paul du Bois-Reymond*. Reprint of the 1910 edition. Cambridge Tracts in Mathematics and Mathematical Physics, No. 12. Hafner Publishing Co., New York, 1971. v+62 pp.
- [8] S. Parameswaran. *Partition functions whose logarithms are slowly oscillating*. Transactions of the American Mathematical Society 100 (1961), 217-240.
- [9] Andreas Weiermann. *An application of graphical enumeration to PA*. The Journal of Symbolic Logic 68 (2003), 5-16.