SURJECTIVITY OF MOD ℓ REPRESENTATIONS ATTACHED TO ELLIPTIC CURVES AND CONGRUENCE PRIMES

IMIN CHEN

ABSTRACT. For a modular elliptic curve $E/\mathbb{Q}$, we show a number of links between the primes ℓ for which the mod ℓ representation of $E/\mathbb{Q}$ has projective dihedral image and congruence primes for the newform associated to $E/\mathbb{Q}$.

1. INTRODUCTION

Let $E/\mathbb{Q}$ be an elliptic curve. Denote by $\overline{\rho}_{E/\mathbb{Q},\ell} : G_{\mathbb{Q}} \to \mathrm{GL}_2(\mathbb{F}_\ell)$ its mod ℓ representation, i.e. the representation obtained by the action of the absolute Galois group $G_{\mathbb{Q}}$ of $\mathbb{Q}$ on the ℓ-torsion points of $E/\mathbb{Q}$ for ℓ prime. Let $S_{E/\mathbb{Q}} = \left\{ \ell \text{ prime } \mid \overline{\rho}_{E/\mathbb{Q},\ell} \text{ is not surjective} \right\}$.

Theorem 1.1 (Serre, [9]). *The set $S_{E/K}$ is finite if E/K does not have complex multiplication.*

In the same paper [9], the following question was asked.

Question 1.2. *Is $S_{\mathbb{Q}} = \bigcup_{E/\mathbb{Q}} S_{E/\mathbb{Q}}$ finite as $E/\mathbb{Q}$ runs through elliptic curves without complex multiplication?*

This question is usually analyzed according to the nature of the image of $\overline{\rho}_{E/\mathbb{Q},\ell}$. If $\overline{\rho}_{E/\mathbb{Q},\ell}$ is not surjective, then by a classification of the subgroups of $\mathrm{GL}_2(\mathbb{F}_\ell)$ we have that $\mathrm{im}\,\overline{\rho}_{E/\mathbb{Q},\ell}$ is contained the normalizer N' or N of a non-split or split Cartan subgroup, a Borel subgroup B, or a subgroup D with projective image S_4. The former three subgroups can be conjugated into one of the following standard forms (under the assumption ℓ is odd in case of N'), respectively,

$$N' = \left\{ \begin{pmatrix} \alpha & \lambda\beta \\ \beta & \alpha \end{pmatrix}, \begin{pmatrix} \alpha & \lambda\beta \\ -\beta & -\alpha \end{pmatrix} \mid \alpha,\beta \in \mathbb{F}_\ell, (\alpha,\beta) \neq (0,0) \right\}$$
$$N = \left\{ \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}, \begin{pmatrix} 0 & b \\ a & 0 \end{pmatrix} \mid a,b \in \mathbb{F}_\ell \right\}$$
$$B = \left\{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \mid a,d \in \mathbb{F}_\ell^\times, b \in \mathbb{F}_\ell \right\},$$

where λ is a non-square in $\mathbb{F}_\ell^\times$.

Date: 1 August 2000.

1991 *Mathematics Subject Classification.* Primary: 11G05, Secondary: 11F80; Torsion points of elliptic curves, Galois representations, congruence primes, Serre tori, grossencharacters, non-split Cartan.

This research was supported by CICMA (1998-99) and MPIM (2000).

Let $S^H_{E/\mathbb{Q}} = \left\{\ell \text{ prime } \mid \operatorname{im}(\overline{\rho}_{E/\mathbb{Q},\ell}) \subset H\right\}$. For H being conjugate to one of N', N, B, D, one can ask whether $S^H_{\mathbb{Q}} = \bigcup_{E/\mathbb{Q}} S^H_{E/\mathbb{Q}}$ is finite as $E/\mathbb{Q}$ runs through elliptic curves without complex multiplication.

Mazur's [3] results on rational isogenies of prime degree show that

$$S^B_{\mathbb{Q}} \subset \{p \text{ prime} \mid p \leq 37\}.$$

Momose shows [6] that an $E/\mathbb{Q}$ with $\operatorname{im}(\overline{\rho}_{E/\mathbb{Q},\ell})$ contained in a conjugate of N has potentially good reduction at all odd primes if $\ell > 13$. These results rely on studying the associated modular curves and bounding their $\mathbb{Q}$-rational points via the arithmetic and geometry of their jacobians. Finally, Serre shows that $S^D_{\mathbb{Q}} \subset \{p \text{ prime} \mid p \leq 13\}$ using local methods (c.f. [3] p. 36).

The case of N' is the most difficult to study using jacobians of modular curves because the jacobians in question do not have a non-trivial quotient with finitely-many $\mathbb{Q}$-rational points.

In this paper, we investigate more carefully the sets $S^N_{E/\mathbb{Q}}$ and $S^{N'}_{E/\mathbb{Q}}$ for a fixed elliptic curve $E/\mathbb{Q}$ from the point of view of modular forms. We briefly recall one such connection implicit in work of Ribet [7] and Kraus [2].

Suppose $E/\mathbb{Q}$ is such that $\operatorname{im}(\overline{\rho}_{E/\mathbb{Q},\ell})$ is contained in H where $H = N', N$ and ℓ is odd. Let C' and C denote the split and non-split Cartan subgroups which are normalized by N' and N, respectively.

Let $\epsilon_{E/\mathbb{Q},\ell}$ be the character obtained by composing $\overline{\rho}_{E/\mathbb{Q},\ell}$ with the map to the quotients $N/C \cong N'/C' \cong \{\pm 1\}$. The character $\epsilon_{E/\mathbb{Q},\ell}$ is non-trivial in the case $H = N'$ as complex conjugation cannot be sent to an element in C' under $\overline{\rho}_{E/\mathbb{Q},\ell}$. In the case $H = N$, we may assume without loss of generality that $\epsilon_{E/\mathbb{Q},\ell}$ is non-trivial or else we are back in the $H = B$ case. Thus, the character $\epsilon_{E/\mathbb{Q},\ell}$ cuts out a quadratic extension K of $\mathbb{Q}$ which is imaginary in the case $H = N'$.

The representation $\overline{\rho}_{E/\mathbb{Q},\ell} \cong \operatorname{Ind}^{\mathbb{Q}}_K \chi$ is induced from a character $\chi : G_K \to \mathbb{F}^\times$, where $\mathbb{F} = \mathbb{F}_{\ell^2}$ or $\mathbb{F}_\ell$ in the cases $H = N'$ or N, respectively. It thus has the property $\overline{\rho}_{E/\mathbb{Q},\ell} \otimes \epsilon_{E/\mathbb{Q},\ell} \cong \overline{\rho}_{E/\mathbb{Q},\ell}$. The following lemma can then be shown.

Lemma 1.3. *Let $E/\mathbb{Q}$ be a modular elliptic curve whose associated newform is $f \in S_2(\Gamma_0(N_E))$. Suppose $\operatorname{im}(\overline{\rho}_{E/\mathbb{Q},\ell}) \subset H$ with $H = N', N$ and ℓ is odd. Let E' be the twist of E by $\epsilon_{E/\mathbb{Q},\ell}$, and let f' be the corresponding twist of $f \in S_2(\Gamma_0(N_E))$. Then $N_{E'} = N_E$ and $f' \in S_2(\Gamma_0(N_E))$ is a newform.*

Proof. Kraus shows in [2] that the type of reduction (good, multiplicative, additive) of E and E' are the same, i.e. the tame exponents ϵ_p of E and E' are the same. On the other hand, the wild exponent δ_p of E depends only on the restriction of $\overline{\rho}_{E/\mathbb{Q},\ell}$ and $\overline{\rho}_{E',\ell} = \overline{\rho}_{E/\mathbb{Q},\ell} \otimes \epsilon_{E/\mathbb{Q},\ell}$ to the wild inertia group at p. For $p \geq 3$, the restrictions are the same as $\epsilon_{E/\mathbb{Q},\ell}$ is trivial on the wild inertia at p. For $p = 2$, the restrictions still have the same image. $\square$

We say that two eigenforms $f, g \in S_2(\Gamma_0(N))$ are congruent modulo λ if $a_p(f) \equiv a_p(g) \pmod{\lambda}$ for $p \nmid \ell N$ where λ is a prime above ℓ of $\mathbb{Q}(a_p(f), a_p(g))$. We say that ℓ is a congruence prime for newform $f \in S_2(\Gamma_0(N))$, if there exists an eigenform g in the (Petersson) orthogonal complement of f such that g is congruent to f modulo λ above ℓ.

The property that $\rho_{E'/\mathbb{Q},\ell} \cong \overline{\rho}_{E/\mathbb{Q},\ell} \otimes \epsilon_{E/\mathbb{Q},\ell} \cong \overline{\rho}_{E/\mathbb{Q},\ell}$ implies the two newforms f, f' are congruent modulo ℓ. Thus the following proposition holds.

Proposition 1.4. *Let $E/\mathbb{Q}$ be a modular elliptic curve whose associated newform is $f \in S_2(\Gamma_0(N_E))$. Suppose ℓ is odd and $\operatorname{im}(\overline{\rho}_{E/\mathbb{Q},\ell})$ is contained in N', or N but not C. Then ℓ is a congruence prime for f.*

In this paper, we will show that there are additional congruences between f and CM-forms in the case N' and discuss how the character of these CM-forms can be controlled under certain hypotheses.

Theorem 1.5. *Let $E/\mathbb{Q}$ be a modular elliptic curve whose associated newform is $f \in S_2(\Gamma_0(N_E))$. Suppose $\operatorname{im}(\overline{\rho}_{E/\mathbb{Q},\ell}) \subset N'$ for $3 < \ell \nmid N_E$. Then there exists a newform $g \in S_2(\Gamma_1(M))$ which is induced from a grossencharacter on K and is congruent to f modulo λ a prime above ℓ where $M \mid N_E$ is the Artin conductor of $\overline{\rho}_{E/\mathbb{Q},\ell}$.*

Theorem 1.6. *Let $E/\mathbb{Q}$ be a modular elliptic curve whose associated newform is $f \in S_2(\Gamma_0(N_E))$ and $16 \nmid N_E$. Suppose $\operatorname{im}(\overline{\rho}_{E/\mathbb{Q},\ell}) \subset N'$ for $3 < \ell \nmid N_E$. Then there exists a newform $g \in S_2(\Gamma_0(M))$ which is induced from a grossencharacter on K and is congruent to f modulo λ a prime above ℓ where $M \mid N_E$ is the Artin conductor of $\overline{\rho}_{E/\mathbb{Q},\ell}$.*

I would like to thank F. Momose for mentioning to me the connection between elliptic curves with $\operatorname{im}(\overline{\rho}_{E/\mathbb{Q},\ell}) \subset N'$ and grossencharacters on imaginary quadratic fields (c.f. also the paper [5] from which the case of prime power N_E in Theorem 1.6 follows).

2. Congruences with CM-forms

2.1. **Algebraic characters.** Fix an algebraic closure $\overline{\mathbb{Q}}$ of $\mathbb{Q}$ and an algebraic closure $\overline{\mathbb{Q}_\ell}$ of $\mathbb{Q}_\ell$. Let $K \subset \overline{\mathbb{Q}} \subset \overline{\mathbb{Q}_\ell}$ be a number field and denote by D_K the set of embeddings of K into $\overline{\mathbb{Q}}$.

Let $T/\mathbb{Q} = \operatorname{Res}^K_{\mathbb{Q}}(\mathbb{G}_m/K)$ be the restriction of scalars of $\mathbb{G}_m/K$ to $\mathbb{Q}$. This is a commutative algebraic group over $\mathbb{Q}$, isomorphic over $\overline{\mathbb{Q}}$ to $\mathbb{G}_m^{[K:\mathbb{Q}]}$, with the following properties.

1. $T(\mathbb{Q}) = K^\times$ and $T(\mathbb{Q}_\ell) = (K \otimes \mathbb{Q}_\ell)^\times = \prod_{v|\ell} K_v$
2. For all $\sigma \in D_K$, there is an algebraic character $[\sigma] : T/\overline{\mathbb{Q}} \to \mathrm{GL}_1/\overline{\mathbb{Q}}$ such that composite
$$K^\times = T(\mathbb{Q}) \subset T(\overline{\mathbb{Q}}) \xrightarrow{[\sigma]} \mathrm{GL}_1(\overline{\mathbb{Q}}) = \overline{\mathbb{Q}}^\times$$
is given by the embedding σ.
3. Every algebraic homormorphism $f : T/\overline{\mathbb{Q}} \to \mathrm{GL}_1/\overline{\mathbb{Q}}$ is of the form $f = \prod_{\sigma \in D_K} [\sigma]^{n(\sigma)}$ where $n(\sigma) \in \mathbb{Z}$. The element $\sum_{\sigma \in D_K} n(\sigma)\sigma \in \mathbb{Z}[D_K]$ is called the weight of f and completely determines f. Given a weight $k \in \mathbb{Z}[D_K]$, let $[k]$ denote the algebraic homomorphism determined by k.

2.2. **Grossencharacters of type A_0.** Let $K \subset \overline{\mathbb{Q}}$ be a number field. For a place v of K, let K_v be the completion of K at v, π_v a uniformizer of K_v, and $\mathcal{O}_v$ the ring of integers of K_v in the case v is a finite place. Let J_K be the ideles of K and $C_K = J_K/K^\times$ the idèle class group. For a modulus $\mathfrak{m}$ of K let $U_\mathfrak{m} = \prod_v U_{\mathfrak{m},v}$ where
$$U_{\mathfrak{m},v} = \begin{cases} \ker(\mathcal{O}_v^\times \to (\mathcal{O}_v/\mathfrak{m}\mathcal{O}_v)^\times) & \text{if } v \nmid \infty \\ \text{the connected component of } 1 & \text{if } v \mid \infty \end{cases}$$

Let $E_{\mathfrak{m}} = \ker(K^\times \to J_K/U_{\mathfrak{m}})$ denote the units congruent to 1 modulo $\mathfrak{m}$, and $C_{\mathfrak{m}} = C/U_{\mathfrak{m}}K^\times$ be the ray class group of modulus $\mathfrak{m}$.

Let $\chi : C_K \to \overline{\mathbb{Q}_\ell}^\times$ be a continuous character. This can be written in the form $\chi = \prod_v \chi_v$ where $\chi_v \mid_{\mathcal{O}_v^\times} = 1$ for all but finitely-many v. The homormorphism $\chi : C_K \to \overline{\mathbb{Q}_\ell}$ is said to be locally algebraic of weight $k \in \mathbb{Z}[D_K]$ if $\chi_\ell = \prod_{v\mid\ell} \chi_v$ coincides with the algebraic character $[-k] : \prod_{v\mid\ell} K_v = T(\overline{\mathbb{Q}_\ell}) \to \mathrm{GL}_1(\overline{\mathbb{Q}_\ell}) = \overline{\mathbb{Q}_\ell}^\times$ of weight $-k$ on the subgroup $\prod_{v\mid\ell} U_{\mathfrak{m},v}$. We say χ has modulus $\mathfrak{m}$ if χ_ℓ coincides with $[-k]$ on $\prod_{v\mid\ell} U_{\mathfrak{m},v}$ and $\chi_v \mid_{U_{\mathfrak{m},v}} = 1$ for $v \nmid \ell$. The smallest modulus for χ is called the conductor of χ.

When $\ell = \infty$, a locally algebraic character χ of modulus $\mathfrak{m}$ and weight k coincides with the notion of a grossencharacter of type A_0 of modulus $\mathfrak{m}$ and weight k.

Theorem 2.1 (Weil, [11]). *Let χ be a grossencharacter of type A_0. The extension $\mathbb{Q}(\chi(\pi_v) \mid v \nmid \infty\mathfrak{m})$ is a finite extension of $\mathbb{Q}$ called the field generated by χ.*

Proposition 2.2. *Let $k \in \mathbb{Z}[D_K]$ be a weight. There exists a non-trivial grossencharacter of type A_0 of weight k and modulus $\mathfrak{m}$ if and only if $[k](E_{\mathfrak{m}}) = 1$. If this holds then there are $h_{\mathfrak{m}}$ such grossencharacters where $h_{\mathfrak{m}}$ is the order of the class group $C_{\mathfrak{m}}$.*

There is a natural grossencharacter of type A_0 of conductor $\mathcal{O}_K$ and weight $\sum_{\sigma\in D_K} \sigma$. This is given by

$$\begin{aligned} \omega_K : C_K &\to \mathbb{R}^{>0} \subset \mathbb{C}^\times \\ x &\mapsto \prod_v ||x_v|| \end{aligned}$$

where $||x_v|| = |x_v|^{[K_v:\mathbb{Q}_p]}$ and for p finite, $|\pi_v| = 1/p^{1/e_v}$, and e_v is the ramification index of $v \mid p$. The character ω_K is trivial on $K^\times$ by the product formula.

2.3. **Fundamental characters.** For $v \mid \ell$ let $\overline{K_v}$ be a fixed algebraic closure of K_v. This fixes an algebraic closure $\overline{k_v}$ of the residue field k_v. Let I_{K_v} denote the inertia subgroup of G_{K_v} and $I_{K_v,t}$ its tame quotient.

A character $\overline{\chi} : I_{K_v,t} \to \overline{k_v}^\times$ is called a tame character. For all $q = \ell^n$, there is a tame character

$$\Theta_{q-1} : I_{K_v,t} \to \mathbb{F}_q^\times \subset \overline{k_v}^\times$$

called the fundamental tame character of level n which is surjective to $\mathbb{F}_q^\times$.

A tame character is said to have level n if its image is contained in $\mathbb{F}_q^\times \subset \overline{k_v}^\times$, $q = \ell^n$, but no smaller finite field. The fundamental tame character of level n has the property that any character $\overline{\chi}$ of level $\leq n$ can be expressed as a power of Θ_{q-1}.

Suppose $\overline{\chi}$ is a tame character of level n and $\overline{\chi} = \Theta_{q-1}^a$ with $0 \leq a < q-1$. Because of the assumption that $\overline{\chi}$ has level n, not all possible a arise. We may write the integer a uniquely in the form $a = a_0 + a_1\ell + \ldots + a_{n-1}\ell^{n-1}$ where $0 \leq a_i \leq \ell - 1$ and hence $\overline{\chi} = \Theta_{q-1}^{a_0}\Theta_{q-1}^{\ell a_1} \ldots \Theta_{q-1}^{\ell^{n-1}a_{n-1}}$.

Let $\overline{\chi} : G_{K_v^{\mathrm{ab}}} \to \overline{k_v}^\times$ be a character and consider its restriction to $I_{K_v^{\mathrm{ab}}}$. This restriction factors to $I_{K_{v,t}^{\mathrm{ab}}}$ to yield a tame character $\overline{\chi}$ which we also denote by $\overline{\chi}$. The local class field homomorphism $r_v : K_v^\times \to G_{K_v^{\mathrm{ab}}}$ induces an isomorphism $k_v^\times \cong I_{K_{v,t}^{\mathrm{ab}}}$ so that the tame character $\overline{\chi}$ has level $\leq n$ where $q = \ell^n = \# k_v$. We denote by $\overline{\chi}\,|_{k_v^\times}$ the character on $k_v^\times$ obtained by precomposing the tame character $\overline{\chi}$ with the local class field homomorphism. Let D_{K_v} denote the set of embeddings $\sigma_v : K_v \to \overline{K_v}$. For each such embedding σ_v, let $\overline{\sigma_v} : k_v \to \overline{k_v}$ denote the associated embedding of residue fields. We can therefore write the tame character in the form as above $\overline{\chi} = \prod_{\sigma_v \in D_{K_v}} \Theta_{q-1}^{\overline{\sigma_v} a(\sigma_v)}$ where $0 \leq a(\sigma_v) \leq \ell - 1$. The element $\sum_{\sigma_v \in D_{K_v}} a(\sigma_v)\sigma_v \in \mathbb{Z}[D_{K_v}]$ is called the optimal weight of $\overline{\chi}$ at v.

A calculation in [9] shows that the composition

$$k_v^\times \cong I_{K_v^{\mathrm{ab}}} \xrightarrow{\Theta_{q-1}} k_v^\times$$

corresponds with the character $x \mapsto x^{-1}$.

3. Adjustment to optimal level and weight

Proposition 3.1. *Let $\mathbb{Q} \subset K \subset \overline{\mathbb{Q}}$ be an imaginary quadratic field whose set of embeddings to $\overline{\mathbb{Q}}$ is denoted by $D_K = \{1, \tau\}$. Let $\overline{\chi} : G_K \to \mathbb{F}_{\ell^2}^\times$ be a continuous character with Artin conductor $\mathfrak{m}$ and let $\widetilde{\chi} : C_K \to \mathbb{C}^\times$ be its Teichmüller lift considered as a continuous character of C_K. Suppose that*

1. *ℓ is inert in K*
2. *$\overline{\chi}\,|_{k_\ell^\times} = \overline{[1]}^{-1}$.*

Then there exists a grossencharacter χ of type A_0 with conductor $\mathfrak{m}$ and weight 1 and a prime λ above ℓ in the field generated by $\widetilde{\chi}$ and χ such that $\widetilde{\chi}(\pi_v) \equiv \chi(\pi_v) \pmod{\lambda}$ for all $v \nmid \infty\ell\mathfrak{m}$.

Proof. By the global class field homomorphism $r_K : C_K \to G_{K^{\mathrm{ab}}}$ we may regard both $\overline{\chi}$ and $\widetilde{\chi}$ as continuous characters of C_K and can write $\overline{\chi} = \prod_v \overline{\chi}_v$ and $\widetilde{\chi} = \prod_v \widetilde{\chi}_v$ where $\overline{\chi}_v$ and $\widetilde{\chi}_v$ are characters of $K_v^\times$. By comparing $\overline{\chi}_v$ and $\widetilde{\chi}_v$ place by place we see that $\widetilde{\chi}$ has conductor $\mathfrak{m}\ell$ and weight 0.

Let $u \in E_{\mathfrak{m}} = K^\times \cap U_{\mathfrak{m}}$. Since $\overline{\chi}$ is trivial on $K^\times$, $\overline{\chi}(u) = 1$. On the other hand, we also have $\overline{\chi}\,|_{k_\ell^\times}(u) = \overline{[1]}^{-1}(u) = \overline{u}^{-1}$ and $\overline{\chi}\,|_{U_{\mathfrak{m},v}}(u) = 1$ for $v \neq \ell$. Thus, we have that $u \equiv 1 \pmod{\ell}$. As K is imaginary quadratic and $\ell \geq 5$, this implies $u = 1$.

Since $E_{\mathfrak{m}}$ is trivial, there exists a grossencharacter ϕ of type A_0 with modulus $\mathfrak{m}$ and weight 1. Write $\phi = \prod_v \phi_v$. As ϕ has weight 1, $\phi_\infty(z) = \overline{z}$. Let $\delta : J_K \to \overline{\mathbb{Q}_\ell}^\times$, $\delta = \prod_v \delta_v$ be defined as follows. For $v \nmid \infty\ell$, let $\delta_v = \phi_v$, and define $\delta_\infty = 1$, $\delta_\ell = \phi_v[1]^{-1}$. By construction, δ factors to a character of C_K. Let $\overline{\delta} : C_K \to \overline{\mathbb{F}_\ell}^\times$ be the reduction of δ modulo a prime λ' above ℓ of the field generated by ϕ (which is the same as the field generated by δ), and let $\widetilde{\delta} : C_K \to \mathbb{C}^\times$ be the Teichmüller lift of $\overline{\delta}$

The desired grossencharacter of type A_0 is then $\chi = \widetilde{\chi}\widetilde{\delta}^{-1}\phi$. The weight of χ is 1 and it evidently has modulus $\mathfrak{m}\ell$. In fact, χ has conductor $\mathfrak{m}$. Since $\chi_\ell = \widetilde{\chi}_\ell\widetilde{\delta}_\ell^{-1}\phi_\ell = \widetilde{[1]}^{-1}\widetilde{[1]}\widetilde{\phi_\ell}^{-1}\phi_\ell$ we see that χ_ℓ is trivial on $\mathcal{O}_\ell^\times$. Thus, χ has modulus $\mathfrak{m}$. To see that χ has conductor precisely $\mathfrak{m}$, consider the character $\chi : C_K \to \overline{\mathbb{Q}_\ell}^\times$ given by $\chi = \widetilde{\chi}\widetilde{\delta}^{-1}\delta$ which has the same conductor as χ. Since χ reduces modulo λ to $\overline{\chi}$ having Artin conductor $\mathfrak{m}$, it follows that $\mathfrak{m}$ divides the Artin conductor of χ as the Artin conductor can only decrease under reduction modulo λ.

For $v \nmid \infty\ell\mathfrak{m}$, $\chi(\pi_v) = \widetilde{\chi}_v(\pi_v)\widetilde{\delta}_v^{-1}(\pi_v)\phi_v(\pi_v) = \widetilde{\chi}_v(\pi_v)\widetilde{\phi}_v^{-1}(\pi_v)\phi_v(\pi_v) \equiv \widetilde{\chi}_v(\pi_v) \pmod{\lambda}$ where λ is a prime of the field generated by $\widetilde{\chi}$ and $\widetilde{\delta}$ above λ'. ∎

4. Proof of Theorem 1

Let $E/\mathbb{Q}$ be a modular elliptic curve whose associated newform is $f \in S_2(\Gamma_0(N_E))$. Suppose $\mathrm{im}(\overline{\rho}_{E/\mathbb{Q},\ell}) \subset N'$ for $3 < \ell \nmid N_E$. Then $\overline{\rho}_{E/\mathbb{Q},\ell} \cong \mathrm{Ind}_K^{\mathbb{Q}} \overline{\chi}$ is induced from a character $\overline{\chi} : G_K \to \mathbb{F}_{\ell^2}^\times$ on the imaginary quadratic field K associated to such a $\overline{\rho}_{E/\mathbb{Q},\ell}$. Since $\ell \nmid N_E$, the argument in [9] p. 317 shows that ℓ is inert in K.

Let us briefly recall the definition of the Serre's optimal weight attached to this particular $\overline{\rho}_{E/\mathbb{Q},\ell}$ [10]. Identifying $G_{\mathbb{Q}_\ell}$ with a decomposition subgroup at ℓ of $G_{\mathbb{Q}}$, the restriction of $\overline{\rho}_{E/\mathbb{Q},\ell}$ to $I_{\mathbb{Q}_\ell}$ factors through its tame quotient $I_{\mathbb{Q}_\ell,t}$ and is semi-simple. Since ℓ is unramified in K, $I_{\mathbb{Q}_\ell} \subset G_K$ so that

$$\overline{\rho}_{E/\mathbb{Q},\ell} \,|_{I_{\mathbb{Q}_\ell}} \cong \begin{pmatrix} \overline{\chi} & 0 \\ 0 & \overline{\chi}' \end{pmatrix}$$

where $\overline{\chi}'(g) = \overline{\chi}(\tau^{-1} g \tau)$.

Both $\overline{\chi}$ and $\overline{\chi}'$ are tame characters of level 2 so that $\overline{\chi}^\ell = \overline{\chi}'$. Write $\overline{\chi} = \Theta_{q-1}^{a_0+\ell a_1}$ where $q = \ell^2 = \# k_v^\times$ and $0 \le a_0, a_1 \le \ell - 1$. Since $\overline{\rho}_{E/\mathbb{Q},\ell}$ is induced from either $\overline{\chi}$ or $\overline{\chi}'$, up to switching $\overline{\chi}'$ for $\overline{\chi}$ we may assume $a > b$. The optimal weight is defined as $k = 1 + a_0 + \ell a_1$. Since $\ell \nmid N_E$, Proposition 4 of [10] implies that $k = 2$ and so $a_0 = 1, a_1 = 0$ in our situation. Thus, we see that $\overline{\chi} \,|_{k_v^\times} = \overline{[1]}^{-1}$.

Let $\widetilde{\chi}$ be the Teichmüller lift of $\overline{\chi}$. By Proposition 3.1, there exists a grossencharacter χ of type A_0 with conductor $\mathfrak{m}$ and weight 1 such that $\chi(\pi_v) \equiv \widetilde{\chi}(\pi_v) \pmod{\lambda}$ where λ is a prime above ℓ of the field generated by $\widetilde{\chi}$ and χ.

Let $I(\mathfrak{m})$ denote the group of fractional ideals of K prime to $\mathfrak{m}$. For an ideal $\mathfrak{a} \in I(\mathfrak{m})$ denote by $[\mathfrak{a}]$ the idèle $\prod_{v \nmid \infty\mathfrak{m}} \pi_v^{e_v}$ associated to the ideal $\mathfrak{a} = \prod_{v \nmid \infty\mathfrak{m}} \mathfrak{p}_v^{e_v}$, $\mathfrak{p}_v$ is the prime of K associated to the finite place v, and π_v is any choice of uniformizer for K_v.

Theorem 4.1 (Hecke). *Let $K \subset \overline{\mathbb{Q}}$ be an imaginary quadratic field with discriminant d_K and let $D_K = \{1, \tau\}$ denote its embeddings into $\overline{\mathbb{Q}}$. Let χ be a grossencharacter of type A_0 on K with conductor $\mathfrak{m}$ and weight $k = u \cdot 1 \in \mathbb{Z}[D_K], u > 0$. Consider $g(z) = \sum_{\mathfrak{a} \in I(\mathfrak{m})} \chi([\mathfrak{a}]) q(z)^{N_K(\mathfrak{a})}$ where $q(z) = e^{2\pi i z}$. Then g is a newform on $S_{u+1}(\Gamma_0(M), \xi)$ where $M = N_K(\mathfrak{m}) \, |d_K|$ and $\xi : (\mathbb{Z}/M\mathbb{Z})^\times \to \mathbb{C}^\times$ is defined by $\xi = \epsilon_K \frac{\chi \circ \mathrm{Ver}}{\omega_{\mathbb{Q}}^u}$. Here $\epsilon_K : C_{\mathbb{Q}} \to \{\pm 1\}$ is the character defining K and* Ver $: C_{\mathbb{Q}} \to C_K$ *is the Verlagerung map.*

Proof. c.f. Theorem 4.8.2 [4] (but note Miyake normalizes his grossencharacters so they are unitary) □

Let $g(z) = \sum_{\mathfrak{a} \in I(\mathfrak{m})} \chi([\mathfrak{a}]) q(z)^{N_K(\mathfrak{a})} \in S_2(\Gamma_1(M))$ be the newform constructed from χ as in the theorem above. Let $p \nmid \ell M$ be a prime and $\mathrm{Fr}_p \in G_{\mathbb{Q}}$ a Frobenius element at p. If p is inert in K, then $\mathrm{Fr}_p \notin G_K$ so that $a_p(f) \equiv \mathrm{tr}\, \overline{\rho}_{E/\mathbb{Q},\ell}(\mathrm{Fr}_p) \equiv \mathrm{tr}(\mathrm{Ind}_K^{\mathbb{Q}} \overline{\chi})(\mathrm{Fr}_p) \equiv 0 = a_p(g) \pmod{\lambda}$. If p is split in K with $v_i \mid p$ being the two places above p, then $\mathrm{Fr}_p \in G_K$ so that $a_p(f) \equiv \mathrm{tr}\, \overline{\rho}_{E/\mathbb{Q},\ell}(\mathrm{Fr}_p) \equiv \mathrm{tr}(\mathrm{Ind}_K^{\mathbb{Q}} \overline{\chi})(\mathrm{Fr}_p) \equiv \overline{\chi}(\mathrm{Fr}_p) + \overline{\chi}(\tau \,\mathrm{Fr}_p\, \tau^{-1}) \equiv \overline{\chi}(\pi_{v_1}) + \overline{\chi}(\pi_{v_2}) \equiv \chi(\pi_{v_1}) + \chi(\pi_{v_2}) = a_p(g) \pmod{\lambda}$. Thus, $a_p(f) \equiv a_p(g) \pmod{\lambda}$ for $p \nmid \ell M$.

Lemma 4.2. *Let ℓ be a prime which is inert in an imaginary quadratic field $K \subset \overline{\mathbb{Q}}$ with its set of embeddings denoted by $D_K = \{1, \tau\}$. Let $\overline{\chi} : G_K \to \mathbb{F}_{\ell^2}^\times$ be a character with Artin conductor $\mathfrak{m}(\overline{\chi})$ prime to ℓ and suppose $\overline{\rho} = \mathrm{Ind}_K^{\mathbb{Q}} \overline{\chi}$ is irreducible. If we* *denote by $N(\overline{\rho})$ the Artin conductor of $\overline{\rho}$, then*

$$N(\overline{\rho}) = (d_K)N_K(\mathfrak{m}(\overline{\chi})).$$

Proof. Let $\widetilde{\chi} : G_K \to L^\times \subset \mathbb{C}^\times, L = \mathbb{Q}(\zeta_n), n = \ell^2 - 1$ be the Teichmüller lift of $\overline{\chi}$, and let $\widetilde{\rho} = \mathrm{Ind}_K^{\mathbb{Q}} \widetilde{\chi}$. By [8] VI.3 Proposition 6, $N(\widetilde{\rho}) = (d_K)N_K(\mathfrak{m}(\widetilde{\chi})) = (d_K)\ell^2 N_K \mathfrak{m}(\overline{\chi})$.

Let us compare $N(\overline{\rho}) = \prod_{p \neq \ell} p^{\overline{e}_p}$ and $N(\widetilde{\rho}) = \prod_{p \neq \ell} p^{\widetilde{e}_p}$ (as $\ell \nmid N(\widetilde{\rho})$). The quantities $\overline{e}_p$ and $\widetilde{e}_p$ are defined as

$$\overline{e}_p = \sum_{i=0}^{\infty} \frac{\#\overline{\rho}(G_{p,i})}{\#\overline{\rho}(G_{p,0})}(2 - \dim \overline{\rho}^{G_{p,i}})$$

$$\widetilde{e}_p = \sum_{i=0}^{\infty} \frac{\#\widetilde{\rho}(G_{p,i})}{\#\widetilde{\rho}(G_{p,0})}(2 - \dim \widetilde{\rho}^{G_{p,i}})$$

where $G_{p,i}$ denotes the i-th ramification group of a decomposition group at p, indexed so that $G_{p,0}$ is the inertia subgroup at p.

Our aim is to show that $N(\overline{\rho})$ is the prime to ℓ-part of $N(\widetilde{\rho})$ and hence equal to $N(\widetilde{\rho}) = (d_K)N_K(\mathfrak{m}(\overline{\chi}))$. It suffices from the definitions of $\overline{e}_p$ and $\widetilde{e}_p$ to show that $\dim \overline{\rho}^H = \dim \widetilde{\rho}^H$ for any given subgroup H of $G_{\mathbb{Q}}$. Let $\widetilde{V} = L \oplus L\tau$ be the representation space of $\widetilde{\rho}$ and let $\Lambda = \mathcal{O}_L \oplus \mathcal{O}_L\tau$ be the natural $G_{\mathbb{Q}}$-invariant lattice lying inside $\widetilde{V}$. For any prime λ above ℓ of L, the $\mathbb{F}_{\ell^2}$-vector space $\Lambda/\lambda\Lambda$ is isomorphic to $\overline{\rho}$.

If H is a given subgroup of $G_{\mathbb{Q}}$, then we see from the description of $\overline{\rho}$ as a reduction of $\widetilde{\rho}$ that $\dim \widetilde{\rho}^H \leq \dim \overline{\rho}^H$. To show equality, we first show that given a non-zero $\overline{v} \in \overline{V}^H$ it is possible to find a lift $\widetilde{v} \in \Lambda^H$, i.e. $\widetilde{v} \in \Lambda^H, \widetilde{v} \equiv \overline{v} \pmod{\lambda\Lambda}$. To do this write $\overline{v} = \overline{x} + \overline{y}\tau$ and let $H_1 = H \cap G_K$ and $H_2 = H \cap \tau G_K$.

Suppose both $\overline{x}, \overline{y} \neq 0$. For every $h \in H_1$, $\overline{\rho}(h)(\overline{v}) = \overline{\chi}(h)\overline{x} + \overline{\chi}'(h)\overline{y}\tau = \overline{x} + \overline{y}\tau = \overline{v}$. It follows that $\overline{\chi}(h) = \overline{\chi}'(h) = 1$ for all $h \in H_1$, and hence $\widetilde{\rho}(h) = 1$ for all $h \in H_1$ so that any lift of $\overline{v}$ is invariant under $h \in H_1$. For every $h = \tau\sigma \in H_2$, $\overline{\rho}(h)(\overline{v}) = \overline{\chi}(\sigma)\overline{y} + \overline{\chi}'(\sigma)\overline{x}\tau = \overline{x} + \overline{y}\tau = \overline{v}$. Thus, $\overline{\chi}(\sigma)\overline{y} = \overline{x}$ and $\overline{\chi}'(\sigma)\overline{x} = \overline{y}$ for all $h = \tau\sigma \in H_2$. Note this implies that $\overline{\chi}(\sigma), \overline{\chi}'(\sigma), \widetilde{\chi}(\sigma), \widetilde{\chi}'(\sigma)$ are constant as $h = \tau\sigma$ varies in H_2. Let $\widetilde{y} \in \mathcal{O}_L$ be any lift of $\overline{y}$. Define $\widetilde{x} = \widetilde{\chi}(\sigma)\widetilde{y}$ and $\widetilde{v} = \widetilde{x} + \widetilde{y}\tau$. Then also $\widetilde{\chi}'(\sigma)\widetilde{x} = \widetilde{y}$, and hence $\widetilde{\rho}(h)(\widetilde{v}) = \widetilde{v}$ for all $h \in H_2$.

Suppose one of $\overline{x}, \overline{y} = 0$. If there exists an element $h = \tau\sigma \in H_2$, then arguing as above, we have that $\overline{\chi}(\sigma)\overline{y} = \overline{x}$ and $\overline{\chi}'(\sigma)\overline{x} = \overline{y}$. But then implies both $\overline{x}, \overline{y} = 0$ contradicting $\overline{v} \neq 0$. Hence, we must have $H \subset G_K$. Again, arguing as above, we see that $\widetilde{\rho}(h) = 1$ for all $h \in H$ and so any lift $\widetilde{v}$ of $\overline{v}$ lies in Λ^H.

The equality $\dim \overline{\rho}^H = \dim \widetilde{\rho}^H$ now follows by picking a lift as above for each element of a basis of $\overline{V}^H$ to form a basis for $\widetilde{V}^H$ of the same size. □

From the above lemma, it follows that $M = N_K(\mathfrak{m})\,|d_K|$ is the Artin conductor of $\overline{\rho}_{E/\mathbb{Q},\ell}$ which divides N_E.

5. Proof of Theorem 2

In this section, we show that the grossencharacter χ used to prove Theorem 1 can adjusted (in certain situations) so that it has the additional property that the character $\xi : (\mathbb{Z}/M\mathbb{Z})^\times \to \mathbb{C}^\times$ of the the associated newform $g(z) = \sum_{\mathfrak{a} \in I(\mathfrak{m})} \chi([\mathfrak{a}])q(z)^{N_K(\mathfrak{a})}$ is trivial.

The Verlagerung map $\mathrm{Ver} : C_{\mathbb{Q}} \to C_K$ is defined by $\mathrm{Ver} = \prod_p \mathrm{Ver}_p$, where Ver_p is the natural map $\mathbb{Q}_p^\times \to (K \otimes \mathbb{Q}_p)^\times$ (here $p = \infty$ is included). By Theorem 4.1, the character ξ is given by the formula $\xi = \epsilon_K \frac{\chi \circ \mathrm{Ver}}{\omega_{\mathbb{Q}}^u}$. If χ is a grossencharacter on K of weight $k = a_0 + a_1\tau$ and modulus $\mathfrak{m}$, then $\chi \circ \mathrm{Ver}$ is a grossencharacter on $\mathbb{Q}$ of weight $a_0 + a_1$ and modulus $N_K(\mathfrak{m})$. Thus, the expression for ξ is indeed a grossencharacter of $\mathbb{Q}$ of weight 0 and modulus M and hence factors to $C_{\mathbb{Q},M} \cong (\mathbb{Z}/M\mathbb{Z})^\times$.

Proposition 5.1. *Let $H \subset G$ be finite abelian groups. Let v be the unique valuation of $\overline{\mathbb{Q}_\ell}$ extending that of $\mathbb{Q}_\ell$. Suppose $f : H \to \overline{\mathbb{Q}_\ell}^\times$ is a character such that such that $v(f(h) - 1) > 0$ for all $h \in H$. Then there exists a character $f' : G \to \overline{\mathbb{Q}_\ell}^\times$ extending f such that $v(f'(g) - 1) > 0$ for all $g \in G$.*

Proof. The main idea of the proof is to mimic the proof of Baer's criterion (c.f. [1]). We shall write the abelian groups $H \subset G$ additively. The first step is to show the following intermediate result.

Let $f : m\mathbb{Z} \to \overline{\mathbb{Q}_\ell}^\times$ be a homomorphism such that $f(m)$ is a root of unity and $v(f(m) - 1) > 0$. Then there exists a homomorphsim $\overline{f} : \mathbb{Z} \to \overline{\mathbb{Q}_\ell}^\times$ extending f such that $\overline{f}(1)$ is a root of unity and $v(\overline{f}(1) - 1) > 0$. To show that $\overline{f}$ exists, choose a root of unity $x \in \overline{\mathbb{Q}_\ell}$ such that $x^m = f(m)$. Let $L = \mathbb{Q}_\ell(x)$, $K = \mathbb{Q}_\ell(f(m))$, with $\lambda \mid \ell$ the unique primes of L, $\mathbb{Q}_\ell$ corresponding to the restrictions of v to these fields. Let $\overline{x}$ be the reduction of x modulo λ and let $\widetilde{x}$ denote the Teichmüller lift of this reduction to $\overline{\mathbb{Q}_\ell}^\times$. Since $x^m = f(m) \equiv 1 \pmod{\lambda}$, we see that $\widetilde{x}^m = 1$, so $\widetilde{x}$ is an m-th root of unity in $\overline{\mathbb{Q}_\ell}$. Now, $(x/\widetilde{x})^m$ is also equal to $f(m)$ but $x/\widetilde{x}$ is congruent to 1 modulo λ. We define $\overline{f}$ by $\overline{f}(1) = x/\widetilde{x}$.

Let $f : H \to \overline{\mathbb{Q}_\ell}^\times$ be given such that $v(f(h) - 1) > 0$ for all $h \in H$. There exists a maximal extension $\overline{f} : \overline{H} \to \overline{\mathbb{Q}_\ell}^\times$ extending $f : H \to \overline{\mathbb{Q}_\ell}^\times$ such that $v(\overline{f}(\overline{h}) - 1) > 0$ for all $\overline{h} \in \overline{H}$. If $\overline{H} = G$, then we are done. If $\overline{H}$ is strictly contained in G, then let $a \in G$ such that $a \notin \overline{H}$. Consider the ideal $\mathfrak{a} = \{r \in \mathbb{Z} : ra \in \overline{H}\}$ of $\mathbb{Z}$. We define a homomorphism $f_0 : \mathfrak{a} \to \overline{\mathbb{Q}_\ell}^\times$ by $f_0(r) = \overline{f}(ra)$. By the intermediate result above, there is an extension $\overline{f}_0 : \mathbb{Z} \to \overline{\mathbb{Q}_\ell}^\times$ such that $v(\overline{f}_0(1) - 1) > 0$. Let $u = \overline{f}_0(1)$. We now define $f'(x + ra) = \overline{f}(x) \cdot u^r$, where $x \in \overline{H}$, and $r \in \mathbb{Z}$. This is well-defined since if $x + ra = 0$, then $r \in \mathfrak{a}$, and hence $\overline{f}(x) \cdot ru = \overline{f}(x) \cdot \overline{f}(ra) = \overline{f}(x + ra) = 0$. Now, f' extends $\overline{f}$ to $H' = <\overline{H}, a>$ still keeping the property $v(f'(h') - 1) > 0$ for all $h' \in H'$, contradicting the maximality of $\overline{f}$. □

Let $\overline{\mathrm{Ver}} : C_{\mathbb{Q},M} \to C_{K,\mathfrak{m}}$ be the homomorphism induced by Ver on ray class groups.

Corollary 5.2. *Let $\xi : C_{\mathbb{Q},M} \to \mathbb{C}^\times$ be a character such that ξ is trivial on the kernel of $\overline{\mathrm{Ver}}$ and $\xi \equiv 1 \pmod{\lambda}$ for λ a prime above ℓ of the field generated by ξ. Then there exists a character $\psi : C_{\mathbb{Q},\mathfrak{m}} \to \mathbb{C}^\times$ such that $\psi \circ \overline{\mathrm{Ver}} = \xi$ and $\psi \equiv 1 \pmod{\lambda}$.*

Let χ be as in the proof of Theorem 1 and let $\xi = \epsilon_K \frac{\chi \circ \mathrm{Ver}}{\omega_{\mathbb{Q}}}$. Assume ξ^{-1} satisfies the requirements of the corollary above and let ψ be the character extending the character ξ^{-1} from the corollary. The character $\chi' = \chi\psi$ also satisfies the requirements for Theorem 1, but now the character of the associated newform g'

becomes

$$\xi' = \epsilon_K \frac{\chi\psi \circ \mathrm{Ver}}{\omega_{\mathbb{Q}}} = \epsilon_K \frac{\chi \circ \mathrm{Ver}}{\omega_{\mathbb{Q}}} \psi \circ \mathrm{Ver} = \xi\xi^{-1} = 1.$$

Since $\overline{\rho}_{f,\ell} \cong \overline{\rho}_{g,\lambda} \pmod{\lambda}$, it follows that $\xi \equiv 1 \pmod{\lambda}$ as the character of f is trivial. Thus, to prove Theorem 2, we need only verify that ξ is trivial on the kernel of $\overline{\mathrm{Ver}}$.

Let $\pi_N : C_{\mathbb{Q}} \to C_{\mathbb{Q},M}$ denote the quotient map. Let $x \in C_{\mathbb{Q}}$ such that $\overline{\mathrm{Ver}}(\pi_N(x)) = 1$. This means that $\mathrm{Ver}(x) = u \cdot k \in U_{K,\mathfrak{m}} \cdot K^{\times}$. Now, $\chi(u \cdot k) = \chi(u) = \chi_\infty(u_\infty) = u_\infty$. On the other hand, $\omega_{\mathbb{Q}} = \omega_K^{1/2}$ and $\omega_K(u \cdot k) = \omega_{\mathbb{Q}}(u) = \omega_{K,\infty}(u_\infty) = u_\infty^2$. Hence, $\frac{\chi \circ \mathrm{Ver}}{\omega_{\mathbb{Q}}}$ considered as a character of $C_{\mathbb{Q},M}$ is trivial on the kernel of $\overline{\mathrm{Ver}}$.

Thus, it remains to show that ϵ_K is trivial on the kernel of $\overline{\mathrm{Ver}}$. The class group $C_{\mathbb{Q},M} \cong (\mathbb{Z}/M\mathbb{Z})^\times$. Given an element $g \in C_{\mathbb{Q},M}$, there exist infinitely many primes q such that $\overline{q} = g \pmod{M}$ by the Chebaterov density theorem. The character ϵ_K considered as a character of $C_{\mathbb{Q},M}$ can then be described by $q \mapsto \left(\frac{d_K}{q}\right)$. Let $g \in C_{\mathbb{Q},M}$ be such that $\overline{\mathrm{Ver}}(g) = 1$ and let us represent $g = \overline{q}$ for an odd prime q. The property $\overline{\mathrm{Ver}}(\overline{q}) = 1$ implies that $q \equiv 1 \pmod{p}$ for every prime $p \nmid N_K(\mathfrak{m})$.

Assume now that $2 \nmid d_K$ so that $d_K \equiv 1 \pmod{4}$ is square-free. From [9] §5.8, we deduce that

1. The character $\epsilon_K = \epsilon_{E/\mathbb{Q},\ell}$ is unramified outside $p \mid N_E$ because of the condition $3 < \ell \nmid N_E$.
2. Furthermore, if $p \nmid d_K$, then $p^2 \mid N_E$.

Thus, since d_K is square-free, it follows that if $p \mid d_K$, then $p \mid N_E/d_K$. But then $p \mid N_K(\mathfrak{m})$ as only semi-stable primes can be stripped from N_E (c.f. [10] §4.6). Thus, we have

$$\epsilon_K(q) = \left(\frac{d_K}{q}\right) = (-1)^{\frac{q-1}{2}\frac{d_K-1}{2}} \left(\frac{q}{d_K}\right) = 1$$

as $d_K \equiv 1 \pmod{4}$ and $q \equiv 1 \pmod{p}$ for each $p \mid N_K(\mathfrak{m})$. Given the following lemma, Theorem 2 is now proved.

Lemma 5.3. *Let $E/\mathbb{Q}$ be an elliptic curve with conductor N_E and suppose $\mathrm{im}(\overline{\rho}_{E/\mathbb{Q},\ell}) \subset N'$ for ℓ odd. Let K be the imaginary quadratic field associated to such $\overline{\rho}_{E/\mathbb{Q},\ell}$. If $16 \nmid N_E$ then $2 \nmid d_K$.*

Proof. Let $\Phi_2 = \overline{\rho}_{E/\mathbb{Q},\ell}(I_2)$ be the image of inertia at 2. Then Φ_2 can be considered as a subgroup of $\mathrm{SL}_2(\mathbb{F}_3)$ with order restricted to $1, 2, 3, 4, 6, 8, 24$ [9]. If $\#\Phi_2 = 1, 2, 3, 6$, then under the assumption ℓ odd, we have that $2 \nmid d_K$ by [9] §5.8. In fact, if $\#\Phi_2 = 24$, then $2 \nmid d_K$ as $\mathrm{SL}_2(\mathbb{F}_3)$ cannot be embedded into the normalizer of a non-split Cartan subgroup N'.

If $\#\Phi_2 = 4$ and $2 \mid d_K$ then $\#\overline{\rho}_{E/\mathbb{Q},\ell}(G_{2,0}) = \#\overline{\rho}_{E/\mathbb{Q},\ell}(G_{2,1}) = 4$ which implies the Artin exponent e_p of $\overline{\rho}_{E/\mathbb{Q},\ell}$ is ≥ 4. Similarly, if $\#\Phi_2 = 8$, then also $e_p \geq 4$. □

6. Acknowledgements

I would like to thank CICMA and the MPIM for their support and hospitality during my stay there 1998-99 and 2000 respectively.

References

[1] P.M. Cohn. *Algebra*, volume 2. John Wiley & Sons, second edition, 1982.

[2] A. Kraus. Une remarque sur les points de torsion des courbes elliptiques. *C. R. Acad. Sci. Paris, t. 321, Série I*, pages 1143–1146, 1995.

[3] B. Mazur. Rational isogenies of prime degree. *Inventiones mathematicae*, 44:129–162, 1978.

[4] T. Miyake. *Modular Forms.* Springer-Verlag, 1989.

[5] F. Momose. Galois action on some ideal section points of the abelian variety associated with a modular form and its application. *Nagoya Mathematical Journal*, 91:19–36, 1983.

[6] F. Momose. Rational points on the modular curves $X_{\mathrm{split}}(p)$. *Compositio Mathematica*, 52:115–137, 1984.

[7] K. Ribet. On ℓ-adic representations attached to modular forms. *Inventiones Mathematicae*, 28:245–275, 1975.

[8] J.P. Serre. *Corps locaux.* Number VIII in Publications de l'Université de Nancago. Hermann, Paris, deuxième edition, 1968.

[9] J.P. Serre. Propriétés galoisiennes des points d'ordre fini des courbes elliptiques. *Inventiones Mathematicae*, 15:259–331, 1972.

[10] J.P. Serre. Sur les représentations modulaires de degré 2 de $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. *Duke Mathematical Journal*, 54(1):179–230, 1987.

[11] A. Weil. On a certain type of characters of the idèle-class groups of an algebraic number-field. In *Proceedings of the international symposium of algebraic number theory, Tokyo & Nikko, 1955*, pages 1–7, Tokyo, 1956. Science Council of Japan.

Max-Planck-Institut-für-Mathematik, Vivatsgasse 7, P.O. Box 7280, D-53072 Bonn, Germany

E-mail address: `chen@mpim-bonn.mpg.de`