


Max-Planck-Institut für Mathematik Bonn

Generalized presentations of infinite groups, in
particular of $\text{Aut}(F_\omega)$

by

O. Bogopolski
W. Singhof



Generalized presentations of infinite groups, in particular of $\text{Aut}(F_\omega)$

O. Bogopolski
W. Singhof

Max-Planck-Institut für Mathematik
Vivatsgasse 7
53111 Bonn
Germany

Institute of Mathematics of Siberian Branch of
Russian Academy of Sciences
Novosibirsk
Russia

Düsseldorf University
Germany

Generalized presentations of infinite groups, in particular of $\text{Aut}(F_\omega)$

O. Bogopolski

Institute of Mathematics of Siberian
Branch of Russian Academy of Sciences,
Novosibirsk, Russia
and Düsseldorf University, Germany
e-mail: Oleg_Bogopolski@yahoo.com

W. Singhof

Düsseldorf University, Germany
e-mail: singhof@math.uni-duesseldorf.de

December 9, 2010

Abstract

We develop a theory of generalized presentations of groups. We give generalized presentations of the symmetric group $\Sigma(X)$ and of the automorphism group of the free group of countable rank, $\text{Aut}(F_\omega)$.

1 Introduction

According to the classical group theory, a group G is generated by a subset A , if every element of G is a *finite product* of elements of $A \cup A^{-1}$. In particular, if G is uncountable, it cannot be generated by a countable set. However, if we allow appropriate *infinite products* this becomes possible for at least the two types of groups mentioned in the abstract. We show that $\Sigma(X)$ and $\text{Aut}(F_\omega)$ can be generated in this generalized sense by transpositions and by elementary Nielsen automorphisms, respectively. Moreover, we describe *generalized presentations* of these groups on these sets of generalized generators.

In the classical group theory, the free groups serve as universal objects, in our theory the universal objects are the so called *generalized free groups*, which include the *big free groups*. On the basis of these topological groups, we develop in Section 2 a theory of generalized presentations of groups.

In Section 3 we describe two generalized presentations of the group $\Sigma(X)$, see Theorems 3.4.2 and 3.5.2. In Section 4 we describe a generalized presentation of the group $\text{Aut}(F_\omega)$, see Theorem 4.8.11. A more algebraic description is given in Theorems 4.8.6 and 4.8.13.

Whereas in the case of the symmetric groups $\Sigma(X)$, we admit arbitrary sets X , for the free groups $F(X)$ we have to assume that X is countable. For bigger sets X , the corresponding questions on $F(X)$ remain open.

2 Generalized free groups and generalized presentations

2.1 Big free groups

We begin with a few recollections concerning big free groups [9]. Let Λ be a set. By $\mathcal{T}(\Lambda)$ we denote the set of all maps

$$f : S \rightarrow \Lambda$$

with the following properties:

- S is a totally ordered set.
- For each $\lambda \in \Lambda$, the set $f^{-1}(\lambda)$ is finite.

We identify two elements $f : S \rightarrow \Lambda$ and $f' : S' \rightarrow \Lambda$ of $\mathcal{T}(\Lambda)$ if there is an order preserving bijection $\varphi : S \rightarrow S'$ with $f = f' \circ \varphi$.

For the remainder of Section 2, we assume that we are given a free involution $\lambda \mapsto \lambda^{-1}$ on Λ .

Definition. A subset I of a linearly ordered set S is called an *interval* of S if the following holds: for $s, s', s'' \in S$ with $s < s'' < s'$ and $s, s' \in I$, we have $s'' \in I$.

We define

$$[s, s'] := [s, s']_S := \{x \in S \mid s \leq x \leq s'\}.$$

Other types of intervals such as $]s, s'[$ or $]s, \infty[$ are defined accordingly.

Given $f, g \in \mathcal{T}(\Lambda)$, we say that g is obtained from f by *cancellation* and write $f \searrow g$, if the following holds:

- If f is of the form $f : S \rightarrow \Lambda$, there is $T \subseteq S$ such that $g = f|_{S \setminus T}$.
- There is an involution $*$ on T such that for all $t \in T$ we have

$$f(t^*) = f(t)^{-1} \tag{1}$$

$$[t, t^*] \subseteq T \tag{2}$$

$$([t, t^*])^* = [t, t^*]. \tag{3}$$

Let \approx be the equivalence relation on $\mathcal{T}(\Lambda)$ generated by $f \searrow g$ and let

$$\text{BF}(\Lambda) := \mathcal{T}(\Lambda) / \approx .$$

By $[f] \in \text{BF}(\Lambda)$ we denote the class of $f \in \mathcal{T}(\Lambda)$. Then $\text{BF}(\Lambda)$ becomes a group by $[f][f'] := [ff']$ where ff' is the concatenation of f and f' . We call $\text{BF}(\Lambda)$ the *big free group* generated by Λ .

We consider Λ as a subset of $\text{BF}(\Lambda)$; the subgroup of $\text{BF}(\Lambda)$ generated by Λ will be denoted by $\mathcal{F}(\Lambda)$. If Λ^+ is a subset of Λ containing, for every $\lambda \in \Lambda$, exactly one of the two elements λ, λ^{-1} , then $\mathcal{F}(\Lambda)$ is the free group $F(\Lambda^+)$ with basis Λ^+ . Following common usage, one should rather call $\text{BF}(\Lambda)$ the big free group generated by Λ^+ .

2.2 Admissible sets and generalized free groups

For certain questions concerning infinite groups, big free groups are a more appropriate tool than free groups. They present, however, new problems: Whereas subgroups of free groups are again free groups, subgroups of big free groups need not be big free groups.

Definition. Any group G with $\mathcal{F}(\Lambda) \leq G \leq \text{BF}(\Lambda)$ is called a *generalized free group* generated by Λ .

The subgroups of $\text{BF}(\Lambda)$ can be described by specifying certain subsets of $\mathcal{T}(\Lambda)$ which we define next:

Definition. A subset \mathcal{S} of $\mathcal{T}(\Lambda)$ is called *admissible* if it has the following properties:

- $f, f' \in \mathcal{S} \Rightarrow ff' \in \mathcal{S}$.
- $f \in \mathcal{S} \Rightarrow \bar{f} \in \mathcal{S}$.
(For a totally ordered set S , let \bar{S} be the set S with the reverse ordering. Given $f : S \rightarrow \Lambda$, we define $\bar{f} : \bar{S} \rightarrow \Lambda$ by $\bar{f}(s) := f(s)^{-1}$.)
- For $f \in \mathcal{S}$ and $g \in \mathcal{T}(\Lambda)$ with $f \searrow g$, we have $g \in \mathcal{S}$.

Given an admissible subset \mathcal{S} of $\mathcal{T}(\Lambda)$, we obtain a group $\text{BF}(\Lambda; \mathcal{S}) := \mathcal{S} / \approx$ where \approx is the equivalence relation on \mathcal{S} generated by $f \searrow g$. To see that $\text{BF}(\Lambda; \mathcal{S})$ is a subgroup of $\text{BF}(\Lambda)$, recall the following definition from [9]:

Definition. An element $f \in \mathcal{T}(\Lambda)$ is called *reduced* if no element of $\mathcal{T}(\Lambda)$ except f itself can be obtained from f by cancellation.

As shown in [9, Theorem 3.9'], every element of $\text{BF}(\Lambda)$ admits a unique reduced representing element in $\mathcal{T}(\Lambda)$. From this fact, it follows easily that $\text{BF}(\Lambda; \mathcal{S})$ is a subgroup of $\text{BF}(\Lambda)$ if \mathcal{S} is admissible. Conversely, any subgroup of $\text{BF}(\Lambda)$ is of the form $\text{BF}(\Lambda; \mathcal{S})$ with an admissible set \mathcal{S} .

With this notation, we have $\text{BF}(\Lambda) = \text{BF}(\Lambda; \mathcal{T}(\Lambda))$. For an admissible subset \mathcal{S} of $\mathcal{T}(\Lambda)$, the group $\text{BF}(\Lambda; \mathcal{S})$ is a generalized free group generated by Λ if and only if $\Lambda \subseteq \mathcal{S}$.

2.3 Big free groups as topological groups

The groups $\text{BF}(\Lambda)$, and hence also their subgroups $\text{BF}(\Lambda; \mathcal{S})$, carry the structure of topological groups. This is what we must explain next.

For a subset A of Λ with $A^{-1} = A$ there is a map

$$\Phi_A : \mathcal{T}(\Lambda) \rightarrow \mathcal{T}(A)$$

which sends $f : S \rightarrow \Lambda$ to its restriction to $f^{-1}(A)$. The map Φ_A induces a homomorphism

$$\varphi_A : \text{BF}(\Lambda) \rightarrow \text{BF}(A) .$$

Since $\text{BF}(A) = \mathcal{F}(A)$ for finite sets A , we obtain a homomorphism

$$\varphi : \text{BF}(\Lambda) \longrightarrow \varinjlim_{A \text{ finite}} \mathcal{F}(A)$$

which is injective by [9, Theorem 3.10]. We get a topology on $\text{BF}(\Lambda)$ which has the subgroups $\ker \varphi_A$ for finite subsets A of Λ as a basis of the neighborhoods of the neutral element.

Let us call this topology on $\text{BF}(\Lambda)$ and on its subgroups $\text{BF}(\Lambda; \mathcal{S})$ the *natural topology*. In the natural topology, the free group $\mathcal{F}(\Lambda)$ is dense in any generalized free group generated by Λ .

Unfortunately, the natural topology is too coarse for many purposes. Depending on the situation, we have to consider topologies which belong to the following class:

Definition. Given an admissible subset \mathcal{S} of $\mathcal{T}(\Lambda)$ with $\Lambda \subseteq \mathcal{S}$, a topology \mathfrak{T} on $\text{BF}(\Lambda; \mathcal{S})$ is called *admissible* if it has the following three properties:

- With \mathfrak{T} , the group $\text{BF}(\Lambda; \mathcal{S})$ becomes a topological group.
- The topology \mathfrak{T} is finer than (i.e. contains at least as many open sets as) the natural topology.
- The free group $\mathcal{F}(\Lambda)$ is dense in $\text{BF}(\Lambda; \mathcal{S})$ with respect to \mathfrak{T} .

2.4 Infinite products in big free groups

In the groups $\text{BF}(\Lambda)$ and, more generally, $\text{BF}(\Lambda; \mathcal{S})$, one can form certain infinite products: Suppose that T is a totally ordered set and that for every $t \in T$ there is given an element $x_t \in \text{BF}(\Lambda)$. Let f_t be the reduced representative of x_t , and suppose that, for each $\lambda \in \Lambda$, the sets $f_t^{-1}(\lambda)$ are empty for all but finitely many values of t . Then we can form, in an obvious manner, the element

$$\prod_{t \in T} f_t \in \mathcal{T}(\Lambda)$$

and can define the infinite product

$$\prod_{t \in T} x_t := \left[\prod_{t \in T} f_t \right] \in \text{BF}(A).$$

For each finite subset A of T , we can of course form the finite product

$$x_A := \prod_{t \in A} x_t.$$

These elements x_A form a net in $\text{BF}(A)$ which converges to $\prod_{t \in T} x_t$ in the standard topology (see [24, Chapter 2]).

All this can be appropriately generalized to the subgroups $\text{BF}(A; \mathcal{S})$.

2.5 Generalized presentations

Now we come to the main definitions of the present paper.

Definition. Let G be a group and A a subset of G . We say that G is *generated by A in the generalized sense* if there exist a set Λ with a free involution, an admissible subset \mathcal{S} of $\mathcal{T}(\Lambda)$ with $\Lambda \subseteq \mathcal{S}$, and an epimorphism $p : \text{BF}(\Lambda; \mathcal{S}) \twoheadrightarrow G$ with $p(\Lambda) \subseteq A$.

Definition. Let G be a group. A *generalized presentation* of G is a tuple $(\Lambda, \mathcal{S}, \mathfrak{T}, R)$ with the following properties:

- Λ is a set with a free involution.
- \mathcal{S} is an admissible subset of $\mathcal{T}(\Lambda)$ with $\Lambda \subseteq \mathcal{S}$.
- \mathfrak{T} is an admissible topology on $\text{BF}(\Lambda; \mathcal{S})$.
- R is a subset of the generalized free group $\text{BF}(\Lambda; \mathcal{S})$.
- There is an epimorphism $p : \text{BF}(\Lambda; \mathcal{S}) \twoheadrightarrow G$ such that $\ker p$ is the smallest normal subgroup which contains the set R and is closed with respect to \mathfrak{T} .

Definition. Given a group G and cardinal number c , we say that G *admits a generalized presentation of type c* if there is a generalized presentation $(\Lambda, \mathcal{S}, \mathfrak{T}, R)$ of G with $|\Lambda| \leq c$ and $|R| \leq c$.

3 Generalized presentations for infinite symmetric groups

For a set X , let $\Sigma(X)$ be the symmetric group consisting of all bijections $\sigma : X \rightarrow X$. The element obtained from x by applying σ will be denoted by $x \cdot \sigma$. In

the present section, we will find two different generalized presentations for $\Sigma(X)$. Both of them are of type $|X|$ if X is an infinite set.

For $x, y \in X$ with $x \neq y$, let $\tau_{x,y} \in \Sigma(X)$ be the transposition interchanging x and y and leaving all other elements of X fixed. We will show that $\Sigma(X)$ is, in the generalized sense introduced in Section 2, generated by these transpositions and that the usual relations between them are actually defining generalized relations.

3.1 An example

In $\Sigma(\mathbb{Z})$, consider the shift $\sigma : n \mapsto n + 1$. In a self-explaining way, we can write, for instance,

$$\sigma = \dots \tau_{2,3} \tau_{1,2} \tau_{0,1} \tau_{-1,0} \dots \quad (4)$$

or

$$\sigma = \tau_{0,1} \tau_{0,-1} \tau_{-1,2} \tau_{-1,-2} \tau_{-2,3} \tau_{-2,-3} \dots \quad (5)$$

There is an important difference between these two ways of writing σ as an infinite product: Consider e.g. the “subword”

$$\tau_{1,2} \tau_{0,1} \tau_{-1,0} \dots$$

of (4). This does not represent an element of $\Sigma(\mathbb{Z})$: Indeed, if it would represent $\rho \in \Sigma(\mathbb{Z})$, what would $\rho(2)$ be?

On the other hand, each subword (i.e. each finite or infinite string of consecutive letters) of (5) defines an element of $\Sigma(\mathbb{Z})$.

In the first of our two generalized presentations of $\Sigma(\mathbb{Z})$, both (4) and (5) will be legal ways of writing σ . In the second presentation, (4) will be illegal, but (5) will remain legal.

3.2 The admissible set \mathcal{S}

We return to the general case and will now describe the set Λ and the admissible set \mathcal{S} of a generalized presentation $(\Lambda, \mathcal{S}, \mathfrak{T}, R)$ of $\Sigma(X)$. Let

$$\Lambda := \Lambda(X) := \{T_{x,y} \mid x, y \in X, x \neq y\};$$

the free involution on Λ sends $T_{x,y}$ to $T_{y,x}$.

The definition of \mathcal{S} requires a certain amount of notation.

With every map $f : S \rightarrow \Lambda$, where S is a totally ordered set, we associate two maps $f_1 : S \rightarrow X$ and $f_2 : S \rightarrow X$ by the following rule: if $s \in S$ and $f(s) = T_{x,y}$, we set $f_1(s) = x$ and $f_2(s) = y$. For $z \in X$ we set

$$U(z, f) := \{s \in S \mid z \in \{f_1(s), f_2(s)\}\} .$$

Let us write

$$\mathcal{S}_0 := \{(f : S \rightarrow \Lambda) \mid |U(x, f)| < \infty \forall x \in X\} \subseteq \mathcal{T}(\Lambda) .$$

We will describe a subset \mathcal{S} of \mathcal{S}_0 .

Let $f : S \rightarrow \Lambda$ be an element of \mathcal{S}_0 which is fixed for the moment; we have to define what it means that f belongs to \mathcal{S} . For each $x \in X$, we will define inductively four sequences

$$x_0^+, \quad x_1^+, \quad x_2^+, \quad \dots \tag{6}$$

$$s_1^+(x), \quad s_2^+(x), \quad \dots \tag{7}$$

$$x_0^-, \quad x_1^-, \quad x_2^-, \quad \dots \tag{8}$$

$$s_1^-(x), \quad s_2^-(x), \quad \dots \tag{9}$$

The sequences (6) and (8) will consist of elements of X , the sequences (7) and (9) of elements of S .

These sequences may be finite or infinite. The sequence (6) will be finite iff (7) is finite, and if this is the case, both will end with the term with the same index, say $n^+(x)$. Here, we allow that $n^+(x) = 0$; this is to mean that (6) is the 1-term sequence x_0^+ and (7) is the empty sequence. If (6) and (7) are infinite sequences, let us put $n^+(x) := \infty$. So, for any $x \in X$, we will have

$$n^+(x) \in \mathbb{N} \cup \{0, \infty\} .$$

Similarly, for the sequences (8) and (9); so we will obtain also

$$n^-(x) \in \mathbb{N} \cup \{0, \infty\} .$$

Now we come to the actual definition of the four sequences:

$$x_0^+ := x_0^- := x .$$

If $U(x, f) = \emptyset$, let $n^+(x) := n^-(x) = 0$.

If $U(x, f) \neq \emptyset$, let

$$s_1^+(x) := \min U(x, f) ,$$

$$s_1^-(x) := \max U(x, f) .$$

Now suppose inductively that for some $n \in \mathbb{N}$, we have already defined

$$x_0^+, \dots, x_{n-1}^+, s_1^+(x), \dots, s_n^+(x)$$

and that $x_{k-1}^+ \in \{f_1(s_k^+(x)), f_2(s_k^+(x))\}$ for all $k = 1, \dots, n$. Then we define x_n^+ by requiring

$$\{x_{n-1}^+, x_n^+\} = \{f_1(s_n^+(x)), f_2(s_n^+(x))\}.$$

Let $s_{n+1}^+(x)$ be the smallest element of S which is contained in $U(x_n^+, f)$ and is bigger than $s_n^+(x)$, assuming that such an element exists. If there is no such element, let $n^+(x) := n$.

This completes the definition of the sequences (6) and (7), and it should be obvious how, by symmetry, the sequences (8) and (9) are defined. Observe that

$$\begin{aligned} s_1^+(x) &< s_2^+(x) < \dots, \\ s_1^-(x) &> s_2^-(x) > \dots. \end{aligned}$$

Now we define the subset \mathcal{S} of \mathcal{S}_0 by declaring that f belongs to \mathcal{S} if and only if, for all $x \in X$, we have

$$n^+(x) < \infty \quad , \quad n^-(x) < \infty.$$

Lemma 3.2.1. *The subset \mathcal{S} of $\mathcal{T}(A)$ is admissible.*

Proof. We have to show that an element of $\mathcal{T}(A)$ is contained in \mathcal{S} if it is obtained from an element of \mathcal{S} by cancellation. This amounts to the following: Suppose we are given an element $(f : S \rightarrow A) \in \mathcal{S}$ and a subset T of S with an involution $*$ such that, for all $t \in T$, we have, in addition to the conditions (2) and (3) of Section 2, that

$$f(t^*) = T_{x,y} \Leftrightarrow f(t) = T_{y,x}. \tag{1'}$$

Then we have to show that $g := f|_{S \setminus T}$ belongs to \mathcal{S} . Let $x \in X$. As in the definition of \mathcal{S} , we have the four finite sequences (x_n^\pm) and $(s_n^\pm(x))$ associated with f . We have to consider the corresponding sequences associated with g . We denote them by (ξ_n^\pm) and $(\sigma_n^\pm(x))$ with $\xi_n^\pm \in X$ and $\sigma_n^\pm(x) \in S \setminus T$. We have to show that they are finite sequences. We will show that (ξ_n^+) is a subsequence of (x_n^+) and $(\sigma_n^+(x))$ is a subsequence of $(s_n^+(x))$. To abbreviate, let us write $s_n := s_n^+(x)$ and $\sigma_n := \sigma_n^+(x)$. We have $\xi_0^+ = x = x_0^+$. We will show that σ_1 is a term in the sequence (s_n) and that ξ_1^+ is the corresponding term in the sequence (x_n^+) . We have

$$s_1 = \min\{s \in S \mid x \in \{f_1(s), f_2(s)\}\}, \tag{10}$$

$$\sigma_1 = \min\{s \in S \setminus T \mid x \in \{f_1(s), f_2(s)\}\}. \tag{11}$$

If $s_1 \in S \setminus T$, then obviously $\sigma_1 = s_1$ and $\xi_1 = x_1$. Therefore we can assume that $s_1 \in T$. Then we conclude from (1') and (2) that

$$s_1 < s_1^* < \sigma_1 .$$

By (3) and the definition of s_2 , there are two possibilities:

Either $s_2 = s_1^*$ or

$$s_1 < s_2 < s_2^* < s_1^* .$$

Iterating this argument, we see that there is a number m such that the finite sequence (s_n) begins with the terms

$$s_1 < s_2 < \dots < s_m < s_m^* < \dots < s_2^* < s_1^*$$

which all lie in T . From (1'), we conclude that the sequence (x_n^+) begins with the terms

$$x_0^+, x_1^+, \dots, x_{m-1}^+, x_m^+, x_{m-1}^+, \dots, x_1^+, x_0^+ .$$

Since $s_{2m} = s_1^* < \sigma_1$, we see that $n^+(x) > 2m$. If $s_{2m+1} \in S \setminus T$, we have $\sigma_1 = s_{2m+1}$ and $\xi_1^+ = x_{2m+1}^+$. If $s_{2m+1} \in T$, we can repeat the above argument; we find that the sequence (s_n) begins with terms of the form

$$\begin{aligned} s_{11}, \dots, s_{1,m_1}, s_{1,m_1}^*, \dots, s_{11}^*, \\ s_{21}, \dots, s_{2,m_2}, s_{2,m_2}^*, \dots, s_{21}^*, \dots, \sigma_1 \end{aligned} \tag{12}$$

and that the sequence (x_n^+) begins with terms of the form

$$\begin{aligned} x, x_{11}, \dots, x_{1,m_1-1}, x_{1,m_1}, x_{1,m_1-1}, \dots, x_{11}, x, \\ x_{21}, \dots, x_{2,m_2-1}, x_{2,m_2}, x_{2,m_2-1}, \dots, x_{21}, x, \dots, \xi_1^+ . \end{aligned} \tag{13}$$

Furthermore, the part of the sequence (12) lying between σ_1 and σ_2 has the same form as the part preceding σ_1 , and so on. \square

For later use, we state a fact which is clear from the proof of Lemma 3.2.1.

Lemma 3.2.2. *Given $f \in \mathcal{S}$ and $x \in X$, denote by $x_\infty(f)$ the last element in the finite sequence x_0^+, x_1^+, \dots . If $f \searrow g$, then $x_\infty(f) = x_\infty(g)$. \square*

3.3 The admissible topology \mathfrak{T} and the homomorphism p

Since \mathcal{S} is an admissible subset of $\mathcal{T}(\Lambda)$ containing Λ , the group $\text{BF}(\Lambda; \mathcal{S})$ contains the free group $\mathcal{F}(\Lambda)$. In order to continue with the description of a generalized presentation of $\Sigma(X)$, we will now define an admissible topology \mathfrak{T} on $\text{BF}(\Lambda; \mathcal{S})$ and a homomorphism $p : \text{BF}(\Lambda; \mathcal{S}) \rightarrow \Sigma(X)$.

On $\Sigma(X)$, there is a natural topology making $\Sigma(X)$ into a topological group. A basis for the neighborhoods of 1 consists of the subgroups

$$U_C := \{\sigma \in \Sigma(X) \mid c \cdot \sigma = c \ \forall c \in C\},$$

where C goes through the set of finite subsets of X .

Using the notation explained in Section 2.3, we have for each finite subset C of X the homomorphism

$$\psi_C := \varphi_{\Lambda(C)} \mid_{\text{BF}(\Lambda; \mathcal{S})} : \text{BF}(\Lambda; \mathcal{S}) \rightarrow \text{BF}(\Lambda(C)) = \mathcal{F}(\Lambda(C)) \hookrightarrow \mathcal{F}(\Lambda).$$

The kernels of these homomorphisms form a basis of neighborhoods of 1 for the natural topology on $\text{BF}(\Lambda; \mathcal{S})$. We define a finer topology \mathfrak{T} by requiring that a basis of neighborhoods of 1 is given by the subgroups

$$W_C := \ker \psi_C \cap V_C$$

where

$$V_C := \{[\gamma] \in \text{BF}(\Lambda; \mathcal{S}) \mid x_\infty(\gamma) = x \ \forall x \in C\}.$$

Here $x_\infty(\gamma)$ is the element introduced in Lemma 3.2.1. It is easy to verify that \mathfrak{T} is an admissible topology.

Lemma 3.3.1. *If we endow $\text{BF}(\Lambda, \mathcal{S})$ with the topology \mathfrak{T} , there is a unique continuous homomorphism $p : \text{BF}(\Lambda, \mathcal{S}) \rightarrow \Sigma(X)$ with $p(T_{x,y}) = \tau_{x,y}$ for all $x, y \in X, x \neq y$.*

For $F = [f] \in \text{BF}(\Lambda, \mathcal{S})$ with $f \in \mathcal{S}$ and $x \in X$, the element $x \cdot p(F) \in X$ is given by

$$x \cdot p(F) = x_\infty(f).$$

Equivalently, we can describe p as follows: There is a unique homomorphism $p : \mathcal{F}(\Lambda) \rightarrow \Sigma(X)$ sending $T_{x,y}$ to $\tau_{x,y}$. We have to extend p to all of $\text{BF}(\Lambda, \mathcal{S})$. Given F, f and x as above, consider the finite sequence (x_n^+) assigned to x and f . Let C be a finite subset of X containing the elements x_n^+ . Then $\psi_C(F) \in \mathcal{F}(\Lambda)$ is a finite word in $T_{y,z}$ with $y, z \in C$, and we have

$$x \cdot p(F) = x \cdot p(\psi_C(F)).$$

□

Lemma 3.3.2. *The homomorphism $p = p_X : \text{BF}(\Lambda, \mathcal{S}) \rightarrow \Sigma(X)$ is surjective.*

Proof. Let $\sigma \in \Sigma(X)$. Suppose that X is the disjoint union of subsets X_β such that $\sigma(X_\beta) = X_\beta$ for all β , and denote by $\sigma_\beta \in \Sigma(X_\beta)$ the restriction of σ to X_β . If $\sigma_\beta \in \text{im } p_{X_\beta}$ for all β , then, obviously, $\sigma \in \text{im } p_X$.

Therefore, to show that $\sigma \in \text{im } p_X$, it suffices to assume that the group generated by σ acts transitively on X . Then we are either in the trivial situation that X is finite, or we are in the situation of the Example in Section 3.1. □

3.4 A generalized presentation of $\Sigma(X)$

Now we complete the description of the generalized presentation $(\Lambda, \mathcal{S}, \mathfrak{T}, R)$ of $\Sigma(X)$.

Let R be the subset of $\text{BF}(\Lambda; \mathcal{S})$ consisting of the elements

- $T_{x,y}^2$,
- $[T_{x,y}, T_{z,w}]$ for $|\{x, y, z, w\}| = 4$,
- $T_{x,y} T_{x,z} T_{x,y}^{-1} T_{y,z}^{-1}$ for $|\{x, y, z\}| = 3$.

For finite X , $\langle \Lambda \mid R \rangle$ is a presentation of $\Sigma(X)$. Indeed, this presentation can be easily obtained from the classical one (see [5, Theorem 7.1 in Chapter 2]) with the help of Tietze transformations.

Let $N = N_X$ be the smallest closed (with respect to \mathfrak{T}) normal subgroup of $\text{BF}(\Lambda; \mathcal{S})$ containing R .

Lemma 3.4.1. $\ker p = N$.

Proof. Given $g \in \ker p$, we have to show that every neighborhood gW_B of g , where $W_B = \ker \psi_B \cap V_B$, contains an element of the normal closure $\langle\langle R \rangle\rangle$ of R . It suffices to show that for each finite subset B of X there is an element $h \in \langle\langle R \rangle\rangle$ with $g^{-1}h \in \ker \psi_B$. Indeed, the inclusion $g^{-1}h \in V_B$ will follow trivially from $g \in \ker p$ and $h \in \langle\langle R \rangle\rangle \subseteq \ker p \subseteq V_B$.

Let $g = [\gamma]$ with $\gamma \in \mathcal{S}$. There is a finite subset C of X which contains, for every $x \in B$, all elements x_n^+ , formed with respect to γ . Then $p \circ \psi_C(g)$ is a permutation of X which is the identity on $X \setminus C$ and on B since $g \in \ker p$. Hence there exists an element $a \in \mathcal{F}(\Lambda(C \setminus B))$ such that $p \circ \psi_C(g) = p(a)$, that is, $p \circ \psi_C(a^{-1}g) = 1$. Since C is finite, we conclude that

$$h := \psi_C(a^{-1}g) \in N_C \subseteq \langle\langle R \rangle\rangle,$$

hence $\psi_C(h^{-1}a^{-1}g) = 1$, and therefore

$$\psi_B(h^{-1}g) = \psi_B(h^{-1}a^{-1}g) = 1.$$

□

The last two lemmas show

Theorem 3.4.2. *For any set X , the triple $(\Lambda, \mathcal{S}, \mathfrak{T}, R)$ is a generalized presentation of $\Sigma(X)$. In particular, if X is infinite, the group $\Sigma(X)$ admits a generalized presentation of type $|X|$.* □

3.5 Another generalized presentation of $\Sigma(X)$

As already indicated in Section 3.1, there is a second generalized presentation of $\Sigma(X)$ with an admissible set \mathcal{S}' which is smaller than \mathcal{S} .

We continue to consider the admissible subset \mathcal{S} of $\mathcal{T}(\Lambda)$ introduced in Section 3.2. Let \mathcal{S}' be the subset of $\mathcal{T}(\Lambda)$ consisting of all maps $f : S \rightarrow \Lambda$ which satisfy the following condition:

For each interval I of S , we have $f \upharpoonright I \in \mathcal{S}$.

Note that the second representation of the shift σ on \mathbb{Z} in Example 3.1 belongs to \mathcal{S}' . Fortunately we have the following lemma and theorem.

Lemma 3.5.1. *\mathcal{S}' is an admissible subset of $\mathcal{T}(\Lambda)$ and the relative topology \mathfrak{T}' on \mathcal{S}' defined by \mathfrak{T} is an admissible topology.* □

Theorem 3.5.2. *$(\Lambda, \mathcal{S}', \mathfrak{T}', R)$ is a generalized presentation of $\Sigma(X)$.* □

4 A generalized presentation of $\text{Aut}(F_\omega)$

We consider a set X and the free group $F(X)$ with basis X . We introduce the set $X^\pm = X \cup X^{-1}$ with the free involution $x \mapsto x^{-1}$. With the notation introduced in Section 2.1, we have $F(X) = \mathcal{F}(X^\pm)$. We will study the automorphism group $\text{Aut } F(X)$.

For $w \in F(X)$ and $\varphi \in \text{End } F(X)$, we write $w\varphi$ for the element obtained from w by applying φ . For $x, y \in X^\pm$ with $x \neq y^{\pm 1}$, let $E_{xy} \in \text{Aut}(F(X))$ be the automorphism which sends x to xy and keeps all elements of $X^\pm \setminus \{x, x^{-1}\}$ fixed. We will show that, for a countably infinite set X , the group $\text{Aut}(F(X))$ is, in our generalized sense, generated by the E_{xy} and that the usual relations are defining generalized relations for $\text{Aut}(F(X))$. In particular, we will find a generalized presentation $(\mathcal{E}, \mathcal{S}, \mathfrak{T}, R)$ for $\text{Aut}(F(X))$ of type \aleph_0 if X is a countably infinite set.

For the first steps of our argument, X is allowed to be an arbitrary set.

4.1 The admissible set \mathcal{S}

Let \mathcal{E} be the set of all elementary Nielsen automorphisms E_{xy} of X , where $x, y \in X^\pm$ and $y \neq x^{\pm 1}$. This set will play the role of the set called Λ in the previous two sections. In the spirit of these sections, it would be more consequent to consider instead of the automorphism E_{xy} the abstract pair (x, y) ; we think that it is more suggestive to use E_{xy} . Of course, we have to consider certain infinite products of the E_{xy} .

Example 4.1.1. Let $X = \{x_1, x_2, \dots\}$ be a countably infinite set.

- 1) The infinite product $E_{x_1x_2}E_{x_3x_4}E_{x_5x_6} \dots$ can be interpreted as the automorphism of $F(X)$ which fixes x_i for even i and sends x_i to x_ix_{i+1} for odd i .
- 2) The infinite product $\dots E_{x_4x_3}E_{x_3x_2}E_{x_2x_1}$ determines an automorphism of $F(X)$ which sends x_i to $x_i \dots x_2x_1$.
- 3) The infinite word $E_{x_1x_2}E_{x_1x_3}E_{x_1x_4} \dots$ does not determine an endomorphism of $F(X)$ since it would send x_1 to the infinite word $x_1(\dots x_4x_3x_2)$.

Given a totally ordered set S and a map $f : S \rightarrow \mathcal{E}$, we obtain two maps $f_1, f_2 : S \rightarrow X^\pm$ by

$$f(s) = E_{f_1(s)f_2(s)}.$$

The last example suggests that for a map $(f : S \rightarrow \mathcal{E}) \in \mathcal{T}(\mathcal{E})$ to be admissible we should at least require that f belongs to

$$\mathcal{S}_0 := \{f \in \mathcal{T}(\mathcal{E}) \mid f_1^{-1}(x) \text{ is finite for each } x \in X^\pm\}.$$

Example 4.1.2. The infinite word $E_{x_1x_2}E_{x_2x_3}E_{x_3x_4} \dots$ is defined by an element of \mathcal{S}_0 but still does not determine an endomorphism of $F(X)$ since x_1 would be sent to the infinite word $x_1x_2x_3 \dots$.

This example suggests that we should also require that f belongs to

$$\mathcal{S}_1 := \{(f : S \rightarrow \mathcal{E}) \in \mathcal{T}(\mathcal{E}) \mid \text{there is no injective and order preserving map } \varphi : \mathbb{N} \rightarrow S \text{ with } f_1(\varphi(n+1)) = (f_2(\varphi(n)))^{\pm 1}\}.$$

Example 4.1.3. The infinite word $\dots E_{x_3x_4}E_{x_2x_3}E_{x_1x_2}$ is defined by an element of $\mathcal{S}_0 \cap \mathcal{S}_1$ and it determines the endomorphism α of $F(X)$ which sends x_i to x_ix_{i+1} for all i . However, α is not invertible since its image consists of words of even length.

So we are finally led to the definition

$$\boxed{\mathcal{S} := \{f \in \mathcal{T}(\mathcal{E}) \mid f \in \mathcal{S}_0 \text{ and } f, \bar{f} \in \mathcal{S}_1\}}.$$

Recall how \bar{f} was defined in Section 2.2: The condition $\bar{f} \in \mathcal{S}_1$ means that there is no injective and order preserving map $\psi : (-\mathbb{N}) \rightarrow S$ with $f_1(\psi(n-1)) = (f_2(\psi(n)))^{\pm 1}$ for all $n \in (-\mathbb{N})$.

The set \mathcal{S} will be the (obviously admissible) set used in our generalized presentation of $\text{Aut } F(X)$.

4.2 A bigger admissible set

As a technical device, we have to introduce another admissible subset $\tilde{\mathcal{S}} \subseteq \mathcal{S}_0$ which is very similar to the one used for $\Sigma(X)$.

Let us fix an element $f \in \mathcal{S}_0$. We want to define what it means that f belongs to $\tilde{\mathcal{S}}$. Let us also fix for the moment an element $x \in X^\pm$. We define inductively two sequences

$$A_0, A_1, A_2, \dots \quad (14)$$

$$s_1, s_2, \dots \quad (15)$$

The sequence (14) consists of subsets of X^\pm , the sequence (15) consists of elements of S . These sequences may be finite or infinite. The sequence (14) will be finite iff (15) is finite, and if this is the case, both will end with the term with the same index, say $n^+ = n^+(f, x)$. We allow $n^+ = 0$; this is to mean that (14) is the 1-term sequence A_0 and (15) is the empty sequence. If (14) and (15) are infinite sequences, let us put $n^+(f, x) := \infty$.

Now we come to the actual definition of the two sequences:

$$A_0 := \{x, x^{-1}\}.$$

If $f_1^{-1}(A_0) = \emptyset$, let $n^+(f, x) = 0$. Otherwise, let

$$\begin{aligned} s_1 &:= \min f_1^{-1}(A_0), \\ A_1 &:= A_0 \cup \{f_2(s_1), f_2(s_1)^{-1}\}, \\ s_2 &:= \min \{s \in S \mid s > s_1 \text{ and } f_1(s) \in A_1\} \end{aligned}$$

if the latter set is non-empty; otherwise put $n^+ = 1$. Observe that the minimum exists since $f \in \mathcal{S}_0$. Put

$$A_2 := A_1 \cup \{f_2(s_2), f_2(s_2)^{-1}\}$$

and so on. Obviously,

$$\tilde{\mathcal{S}} := \{f \in \mathcal{S}_0 \mid n^+(f, x) < \infty \text{ and } n^+(\bar{f}, x) < \infty \text{ for all } x \in X^\pm\}$$

is an admissible subset of $\mathcal{T}(\mathcal{E})$.

Lemma 4.2.1. $\mathcal{S} \subseteq \tilde{\mathcal{S}}$.

Proof. Let $f \in \mathcal{S}$ and $x \in X^\pm$. It suffices to show that $n^+(f, x) < \infty$. Consider the corresponding sequence (s_k) in S and the sequence (A_k) of finite subsets of X^\pm . Let $V_k := A_k / \sim$ where the equivalence relation \sim is given by $a \sim a^{\pm 1}$. Then we have $V_0 \subseteq V_1 \subseteq V_2 \subseteq \dots$, where the set V_0 consists of one element, say v_0 . Let $V := V_0 \cup V_1 \cup \dots$. It suffices to show that V is a finite set.

Let K be the set of all natural numbers k for which V_k is bigger than V_{k-1} . For $k \in K$, denote by v_k the element of $V_k \setminus V_{k-1}$. Then $V = \{v_k \mid k \in K \cup \{0\}\}$.

We define a graph Γ as follows: Let V be the set of vertices of Γ . For each $k \in K$, we introduce an edge e_k beginning in the class of $f_1(s_k)$ and ending in the class of $f_2(s_k)$. Therefore e_k begins in V_{k-1} and ends in v_k . Since $f \in \mathcal{S}_0$, there can start only finitely many edges in each vertex. Therefore Γ is a tree to which we can apply König's lemma if Γ is infinite. This leads to an immediate contradiction to the condition $f \in \mathcal{S}_1$. \square

4.3 The homomorphism $\Psi : \mathbf{BF}(\mathcal{E}; \mathcal{S}) \rightarrow \mathbf{Aut}(F(X))$

Suppose we are given a map $f : S \rightarrow \mathcal{E}$ belonging to the admissible set $\tilde{\mathcal{S}}$. Let us write

$$f(s) = E_{x_s y_s}.$$

It should be intuitively clear that we obtain an endomorphism α_f of $F(X)$ by

$$\alpha_f := \prod_{s \in S} E_{x_s y_s}.$$

The objective of the present subsection is to make this definition precise, to show that α_f is an automorphism and that by assigning α_f to f we obtain a homomorphism Ψ from $\mathbf{BF}(\mathcal{E}; \tilde{\mathcal{S}})$ and hence from $\mathbf{BF}(\mathcal{E}; \mathcal{S})$ to $\mathbf{Aut}(F(X))$.

Fix $f \in \tilde{\mathcal{S}}$ as above. For any finite subset $C := \{s_1, \dots, s_n\}$ of S with $s_1 < \dots < s_n$, we obtain the automorphism

$$E_C := E_{x_{s_1} y_{s_1}} \dots E_{x_{s_n} y_{s_n}}.$$

For $x \in X^\pm$, we have defined in Section 4.2 the finite subset

$$\{s_1, \dots, s_{n^+}\} =: C(f, x)$$

with $n^+ = n^+(f, x)$. Observe that $C(f, x) = C(f, x^{-1})$. Hence we obtain an endomorphism α_f of $F(X)$ by

$$x\alpha_f := xE_{C(f, x)} \text{ for } x \in X^\pm.$$

Lemma 4.3.1. *Let $f, g \in \tilde{\mathcal{S}}$ with $f \searrow g$. Then $\alpha_f = \alpha_g$.*

Proof. There is a subset T of S with $g = f|S \setminus T$ and there is an involution $*$ on T satisfying (1), (2), (3). We can write the ordered set $C(f, x)$ in the form

$$C(f, x) = T_1 S_1 \dots T_r S_r$$

with subsets T_i of T and subsets S_i of $S \setminus T$; of these subsets, only T_1 and S_r are allowed to be possibly empty. Then we have

$$C(g, x) = S'_1 \dots S'_r$$

where S'_i is a possibly empty subset of S_i . As in the proof of Lemma 3.2.1, we see that each T_i is a concatenation of subsets of the form

$$\sigma_1 < \sigma_2 < \dots < \sigma_m < \sigma_m^* < \dots < \sigma_2^* < \sigma_1^*.$$

Therefore $E_{T_i} = \text{id}$, and it is clear that the elements of S_i which are not in S'_i do not affect x . Hence we have

$$xE_{C(f,x)} = xE_{C(g,x)}. \quad \square$$

Observe that for any finite subset C' of S containing $C(f, x)$, we have

$$x\alpha_f = xE_{C'}.$$

Hence, for each $w \in F(X)$ and each finite subset C' of S containing the sets $C(f, x)$ for all the letters x of the word w , we have

$$w\alpha_f = wE_{C'}.$$

It is not difficult to conclude:

Lemma 4.3.2. *For $f, f' \in \tilde{\mathcal{S}}$, we have $\alpha_{ff'} = \alpha_f\alpha_{f'}$.* \square

As an immediate consequence of Lemmas 4.3.1 and 4.3.2, we get:

Proposition 4.3.3. *For each $f \in \tilde{\mathcal{S}}$, the endomorphism α_f is an automorphism. By $\Psi[f] := \alpha_f$ we obtain a homomorphism Ψ from $\text{BF}(\mathcal{E}; \tilde{\mathcal{S}})$ to $\text{Aut}(F(X))$.* \square

4.4 Refined Nielsen's method

Here we introduce a *complexity of a word* and prove Theorem 4.4.2 which is a refinement of the classical Nielsen method for simplifying tuples of elements in free groups (see [25, Chapter 1, Proposition 2.2]). Condition (1) of this theorem is contained (in a similar form) in Nielsen's method, while Condition (2) is new. Both conditions are present in Corollary 4.4.3, which will be used later in the proof of Theorem 4.5.3.

First we give some definitions. Let F be a free group with a basis X . We identify the elements of F with reduced words in the alphabet X^\pm . For any element $w \in F$, we denote by $|w|$ the length of w with respect to X .

We define four types of transformations on an arbitrary countable tuple of elements $U = (u_1, u_2, \dots)$ of F .

- (I_i) replace u_i by u_i^{-1} (*inversion*);
- (R_{ij}) replace u_i by $u_i u_j$ where $i \neq j$ (*right multiplication*);
- (L_{ij}) replace u_i by $u_j u_i$ where $i \neq j$ (*left multiplication*);

(D_{*i*}) delete u_i if $u_i = 1$ (*deletion*).

In all cases it is understood that the u_k for $k \neq i$ remain unchanged. These transformations are called *elementary Nielsen transformations*.

A tuple $U = (u_1, u_2, \dots)$ of elements of F is called *Nielsen reduced* if for any three elements v_1, v_2, v_3 of the form $u_i^{\pm 1}$, where $u_i \in U$, the following conditions hold:

(N1) $v_1 \neq 1$;

(N2) if $v_1 v_2 \neq 1$, then $|v_1 v_2| \geq |v_1|, |v_2|$;

(N3) if $v_1 v_2 \neq 1$ and $v_2 v_3 \neq 1$, then $|v_1 v_2 v_3| > |v_1| - |v_2| + |v_3|$.

Condition (N2) means that in the product $v_1 v_2$ not more than half of each factor cancels. Condition (N3) means that in the product $v_1 v_2 v_3$ at least one letter of v_2 remains uncanceled.

Now we will introduce some notations. Suppose that the set $X \cup X^{-1}$ is totally ordered. This order induces the *graded lexicographical order* \preceq on the set of all reduced words in the alphabet X^{\pm} by the following rule.

Let u and v be two reduced words in the alphabet X^{\pm} . Denote by w their maximal common initial segment. We write $u \preceq v$ if either $|u| < |v|$ or $|u| = |v|$ and the letter of u following w (if exists) occurs earlier in the ordering than the letter of v following w .

We write $u \prec v$ if $u \preceq v$ and $u \neq v$. Note that $u \prec v$ implies that $uw \prec vw$ for any $w \in F$, provided that the words uw and vw are reduced. For any $w \in F$, let $\phi(w)$ denote the cardinality of the set $\{z \mid z \preceq w\}$.

From now on and to the end of this subsection let X be a finite set. Then $u \prec v \iff \phi(u) < \phi(v)$ and $\phi(u) < \phi(v) \iff \phi(uw) < \phi(vw)$ provided the words uw and vw are reduced.

Let $v \in F$ be a reduced word. By $L(v)$ we denote the initial segment of v of length $\lfloor (|v| + 1)/2 \rfloor$. The *weight* $W(v)$ of the word v is defined to be $W(v) = \phi(L(v)) + \phi(L(v^{-1}))$. Obviously, $W(v) = W(v^{-1})$ and there exists only a finite number of words with weight not exceeding a given natural number. Note also, that $W(u) = W(v)$ do not imply $u = v$.

The *complexity* of v is defined to be the pair of nonnegative integer numbers $(|v|, W(v))$ denoted $\text{Compl}(v)$. We will write $\text{Compl}(v) <_{lex} \text{Compl}(u)$ if either $|v| < |u|$ or $|v| = |u|$ and $W(v) < W(u)$. Obviously, $\text{Compl}(u) = \text{Compl}(u^{-1})$. Note that $\text{Compl}(v) = \text{Compl}(u)$ does not imply $v = u^{\pm 1}$ in general.

Lemma 4.4.1. *Let u, v be two reduced words in F , such that $uv \neq 1$ and $|uv| < |u| = |v|$. Then $W(u) \neq W(v)$.*

Proof. The condition $|uv| < |u| = |v|$ implies, that $L(u^{-1}) = L(v)$. Suppose $W(u) = W(v)$. Then $\phi(L(u)) + \phi(L(u^{-1})) = \phi(L(v)) + \phi(L(v^{-1}))$ and hence $L(u) = L(v^{-1})$. The conditions $L(u^{-1}) = L(v)$ and $L(u) = L(v^{-1})$ imply $uv = 1$, a contradiction. \square

Theorem 4.4.2. *Let F be a free group of finite rank. Let $U = (u_1, \dots, u_m)$ be a finite tuple of elements of F with $\text{rank}\langle U \rangle = m$, which is not Nielsen reduced. Then there exists a Nielsen transformation \mathcal{N} of the form $(R_{ij})^{\pm 1}$ or $(L_{ij})^{\pm 1}$, which carries U to another tuple $U' = (u'_1, \dots, u'_m)$, such that the following holds:*

- (1) $\text{Compl}(u'_i) <_{\text{lex}} \text{Compl}(u_i)$ and $u'_k = u_k$ for all $k \in \{1, \dots, m\} \setminus \{i\}$,
- (2) $\text{Compl}(u_j) <_{\text{lex}} \text{Compl}(u_i)$.

Proof. Because of the assumption on the rank, the condition (N1) is satisfied. We will use the following easy observation: *it is sufficient to prove the theorem for a tuple $(u_1^{\epsilon_1}, \dots, u_m^{\epsilon_m})$ with some choice of $\epsilon_1, \dots, \epsilon_m \in \{1, -1\}$.* (*)

a) Suppose that the condition (N2) is not satisfied. Then there are v_i, v_j of the form $u_i^{\pm 1}$, where $u_i \in U$, and such that $v_1 v_2 \neq 1$ and additionally $|v_1 v_2| < |v_1|$ or $|v_1 v_2| < |v_2|$.

Using (*), we may assume that $v_1, v_2 \in U$, say $v_1 = u_i$ and $v_2 = u_j$. Without loss of generality, we may also assume that $|u_i u_j| < |u_i|$. Indeed, if $|u_i u_j| < |u_j|$, then $|u_j^{-1} u_i^{-1}| < |u_j^{-1}|$ and using (*) again, we can reduce to the previous situation.

Consider what happens if we apply (R_{ij}) or (L_{ji}) to the relevant part of U .

$$\begin{aligned} (u_i, u_j) &\xrightarrow{(R_{ij})} (u_i u_j, u_j) \\ (u_i, u_j) &\xrightarrow{(L_{ji})} (u_i, u_i u_j) \end{aligned}$$

If $|u_j| < |u_i|$, we choose $\mathcal{N} = (R_{ij})$. Then Condition (2) will be satisfied automatically and Condition (1) will be satisfied by our assumption $|u_i u_j| < |u_i|$.

If $|u_i| < |u_j|$, we choose $\mathcal{N} = (L_{ji})$. Then Condition (2) will be satisfied automatically and Condition (1) will be satisfied, since $|u_i u_j| < |u_i| < |u_j|$.

If $|u_j| = |u_i|$, then both (R_{ij}) and (L_{ji}) satisfy Condition (1). By Lemma 4.4.1, we can choose one of them as \mathcal{N} so that Condition (2) will be satisfied.

b) Suppose that Condition (N2) is satisfied, but Condition (N3) is not. Then there are v_1, v_2, v_3 of the form $u_i^{\pm 1}$, where $u_i \in U$, such that $v_1 v_2 \neq 1$, $v_2 v_3 \neq 1$, and

$$|v_1 v_2 v_3| \leq |v_1| - |v_2| + |v_3|. \quad (16)$$

Let $v_1 = ap^{-1}$ and $v_2 = pb$, where p is the maximal initial segment of v_2 canceling in the product $v_1 v_2$. Similarly, we write $v_2 = cq^{-1}$ and $v_3 = qd$, where q^{-1} is the maximal terminal segment of v_2 cancelling in the product $v_2 v_3$. By Condition (N2), we have $|p|, |q| \leq |v_2|/2$. Then $v_2 = prq^{-1}$ for some r . Assuming that $r \neq 1$ we would have

$$|v_1 v_2 v_3| = |v_1| - |v_2| + |v_3| + 2|r|,$$

a contradiction to (16). Therefore, $r = 1$, $v_2 = pq^{-1}$ and $|p| = |q| = |v_2|/2$. Since $v_2 \neq 1$, we obtain $p \neq q$.

Case 1. Suppose that $\phi(p) < \phi(q)$. Using $(*)$, we may assume that $v_2, v_3 \in U$, say $v_2 = u_i, v_3 = u_j$. As above, we will look, what happens if we apply (R_{ij}) or (L_{ji}) to the relevant part of U :

$$\begin{aligned}(u_i, u_j) &= (pq^{-1}, qd) \xrightarrow{(R_{ij})} (pd, qd) = (u_i u_j, u_j) \\ (u_i, u_j) &= (pq^{-1}, qd) \xrightarrow{(L_{ji})} (pq^{-1}, pd) = (u_i, u_i u_j)\end{aligned}$$

Note that $|u_j| \geq |u_i|$, since $|u_j| = |u_i u_j| \geq |u_i|$, where the last inequality follows from Condition (N2). Thus, we consider two subcases.

Case 1.1. Suppose $|u_i| < |u_j|$. In this case we choose $\mathcal{N} = (L_{ji})$. Then Condition (2) is trivially satisfied. Condition (1) is also satisfied, since $|u_i u_j| = |u_j|$ and $W(u_i u_j) < W(u_j)$ because of $\phi(p) < \phi(q)$.

Case 1.2. Suppose $|u_j| = |u_i|$. Then $|d| = |p| = |q|$ and so

$$|u_i u_j| = |u_i| = |u_j|. \quad (17)$$

First we show that $W(u_i) \neq W(u_j)$. Assume the contrary. Then from

$$\begin{aligned}W(u_i) &= \phi(p) + \phi(q), \\ W(u_j) &= \phi(q) + \phi(d^{-1})\end{aligned}$$

follows that $\phi(p) = \phi(d^{-1})$. This implies that $p = d^{-1}$, and hence $v_2 v_3 = 1$, a contradiction. Thus, it remains to consider two subcases.

Case 1.2.1. Suppose that $W(u_j) < W(u_i)$. In this case we choose $\mathcal{N} = (R_{ij})$. Then Condition (2) is trivially satisfied. Now we show, that Condition (1) is satisfied. In view of (17), we shall prove that $W(u_i u_j) < W(u_i)$. Since $W(u_j) < W(u_i)$, we have $\phi(q) + \phi(d^{-1}) < \phi(p) + \phi(q)$ and hence $\phi(d^{-1}) < \phi(p)$. Recall, that we consider Case 1 where $\phi(p) < \phi(q)$. Then

$$W(u_i u_j) = W(pd) = \phi(p) + \phi(d^{-1}) < \phi(p) + \phi(q) = W(u_i)$$

and we are done.

Case 1.2.2. Suppose that $W(u_i) < W(u_j)$. In this case we choose $\mathcal{N} = (L_{ji})$. Then Condition (2) is trivially satisfied. Now we show, that Condition (1) is satisfied. In view of (17), we shall prove, that $W(u_i u_j) < W(u_j)$. The later is valid:

$$W(u_i u_j) = W(pd) = \phi(p) + \phi(d^{-1}) < \phi(q) + \phi(d^{-1}) = W(u_j).$$

Case 2. Suppose that $\phi(p) > \phi(q)$. Then we consider the pair (v_2^{-1}, v_1^{-1}) instead of the pair v_2, v_3 and reduce to Case 1. \square

Corollary 4.4.3. *Let F be a free group of finite rank with a basis X . Every finite tuple $U = (u_1, \dots, u_m)$ of elements of F with $\text{rank}\langle U \rangle = m$ can be carried to a Nielsen reduced tuple $V = (v_1, \dots, v_m)$ by a finite number of Nielsen's right and left multiplications, so that at each step Conditions (1) and (2) of Theorem 4.4.2 are satisfied.*

If x_1, \dots, x_r, x_{r+1} is a part of the basis X , such that U contains x_1, \dots, x_r , up to inversions, on some places i_1, \dots, i_r and $x_{r+1} \in \langle U \rangle$, then the Nielsen transformations do not change the elements on places i_1, \dots, i_r of the involved tuples and the last tuple V will additionally contain x_{r+1} up to inversion.

Proof. For every tuple $W = (w_1, \dots, w_m)$ we define its complexity by the rule $\text{Compl}(U) = \sum_{i=1}^m \text{Compl}(w_i)$, where we summate the complexities as 2-dimensional vectors. By Theorem 4.4.2, if the tuple W is not Nielsen reduced, we can decrease its complexity by applying an appropriate Nielsen transformation. Since the complexity cannot infinitely decrease, if we start from U , after a finite number of steps described in Theorem 4.4.2 we get a Nielsen reduced tuple. The elements on places i_1, \dots, i_r will not be changed because of Conditions (1), (2). The last claim follows from the fact, that V is Nielsen reduced and from $x_{r+1} \in \langle U \rangle = \langle V \rangle$ (see [25, Corollary 2.4 in Chapter I] with $w = x_{r+1}$). \square

Lemma 4.4.4. *Let F be a free group of finite rank, and let $u, v \in F$. Suppose that for some $\epsilon \in \{-1, 1\}$ the following holds:*

- (1) $|u| = |v|$,
- (2) $\text{Compl}(uv) <_{lex} \text{Compl}(u)$,
- (3) $\text{Compl}(v) <_{lex} \text{Compl}(u)$.

Then $L(v) = L(u^{-1})$ and $L(v^{-1}) \prec L(u)$.

Proof. From Condition (2) we have $|uv| \leq |u|$. Together with $|u| = |v|$ this implies that $L(v) = L(u^{-1})$. From Conditions (3) and $|u| = |v|$ we have $W(v) < W(u)$, i.e. $\varphi(L(v)) + \varphi(L(v^{-1})) < \varphi(L(u)) + \varphi(L(u^{-1}))$. Hence $\varphi(L(v^{-1})) < \varphi(L(u))$ and so $L(v^{-1}) \prec L(u)$. \square

Using the fact that $\text{Compl}(w) = \text{Compl}(w^{-1})$ one can easily deduce the following lemma from the previous one.

Lemma 4.4.5. *Let F be a free group of finite rank, and let $u, v \in F$. Suppose that for some $\epsilon, \tau \in \{-1, 1\}$ the following holds:*

- (1) $|u| = |v|$,
- (2) $\text{Compl}(u^\epsilon v^\tau) <_{lex} \text{Compl}(u)$,
- (3) $\text{Compl}(v) <_{lex} \text{Compl}(u)$.

Then $L(v^\tau) = L(u^{-\epsilon})$ and $L(v^{-\tau}) \prec L(u^\epsilon)$.

Notation 4.4.6. *It is convenient to unify four different notations into one: For $\epsilon, \tau \in \{-1, 1\}$ and $i, j \in \mathbb{N}$ with $i \neq j$ we define*

$$T_{i^\epsilon j^\tau} = \begin{cases} R_{ij}, & \text{if } \epsilon = \tau = 1 \\ R_{ij}^{-1}, & \text{if } \epsilon = 1, \tau = -1 \\ L_{ij}, & \text{if } \epsilon = -1, \tau = -1 \\ L_{ij}^{-1}, & \text{if } \epsilon = -1, \tau = 1. \end{cases}$$

4.5 The first step towards the surjectivity of $\Psi : \text{BF}(\mathcal{E}, \mathcal{S}) \rightarrow \text{Aut}(F_\omega)$

An automorphism of $F(X)$ is called *monomial*, if it maps X^\pm onto itself. Let $\mathcal{M}(X)$ be the group of monomial automorphisms of $F(X)$. For short, we denote by F_n the free group with the finite basis $X_n = \{x_1, \dots, x_n\}$ and by F_ω the free group with the infinite countable basis $X_\omega = \{x_1, x_2, \dots\}$.

In this section we prove Theorem 4.5.3 which states that, up to a monomial automorphism, every automorphism of F_ω lies in $\text{im } \Psi$. In the next section we show that every monomial automorphism of F_ω lies in $\text{im } \Psi$. With that the surjectivity of $\Psi : \text{BF}(\mathcal{E}, \mathcal{S}) \rightarrow \text{Aut}(F_\omega)$ will be proven.

Orders. We will use the following order on X_n^\pm : $x_1^{-1} \prec x_1 \prec \dots \prec x_n^{-1} \prec x_n$. The union of these orders for $n \in \mathbb{N}$ gives the order on X_ω^\pm . The corresponding graded lexicographical orders on F_n and F_ω are denoted by \prec_n and \prec_ω . The complexity of a word $v \in F_n$ in the group F_m with $m \geq n$ is denoted by $\text{Compl}_m(v)$. We stress that the complexity of the same word in different groups may be different.

Remark 4.5.1. 1) If $u \prec_n v$ for some $u, v \in F_n$, then $u \prec_m v$ for every $m > n$ and, moreover, $u \prec_\omega v$.

2) For any $w \in F_n$, the set $\{z \in F_n \mid z \prec_n w\}$ is finite. If $|w| \geq 2$ and the first letter of w differs from x^{-1} , then the set $\{z \in F_\omega \mid z \prec_\omega w\}$ is infinite.

3) Every decreasing chain $\dots \prec_\omega z_3 \prec_\omega z_2 \prec_\omega z_1$ in F_ω , where all z_i have the same length, is finite.

In the proof of Theorem 4.5.3 we will indirectly use the following density lemma (see [25, Proposition 4.1 in Chapter I]).

Lemma 4.5.2. *For every $\alpha \in \text{Aut}(F_\omega)$ and for every n , there exists $m \geq n$ and an automorphism $\beta \in \text{Aut}(F_m)$, such that $\alpha|_{X_n} = \beta|_{X_n}$.*

Consider the set $\mathcal{E} = \{E_{xy} \mid x, y \in X^\pm, y \neq x, x^{-1}\}$ of elementary automorphisms of F_ω . Recall that with every map $f : S \rightarrow \mathcal{E}$ we associate two maps $f_1 : S \rightarrow X^\pm$ and $f_2 : S \rightarrow X^\pm$ by the following rule: if $s \in S$ and $f(s) = E_{xy}$, we set $f_1(s) = x$ and $f_2(s) = y$.

Theorem 4.5.3. *For every automorphism α of F_ω there exists a monomial automorphism $\sigma \in \mathcal{M}(X_\omega)$, such that α can be written in the form $\sigma\alpha_f$ for some $f : (-\mathbb{N}) \rightarrow \mathcal{E}$ with the following properties.*

- (i) *For every $x \in X^\pm$ the set $\{n \in (-\mathbb{N}) \mid f(n) = E_{xy} \text{ for some } y \in X^\pm\}$ is finite.*
- (ii) *There is no injective and order preserving map $\psi : (-\mathbb{N}) \rightarrow (-\mathbb{N})$ with conditions $f_1(\psi(n-1)) = (f_2(\psi(n)))^{\pm 1}$ for all $n \in (-\mathbb{N})$.*

Proof. Below in part **a)** we define some auxiliary elementary Nielsen transformations, which will be used in part **b)** for construction the function f which satisfy this theorem. Denote $w_i^{(0)} = x_i\alpha^{-1}$ and consider the infinite tuple $W^{(0)} = (w_1^{(0)}, w_2^{(0)}, \dots)$.

a) We will inductively define Nielsen transformations \mathcal{N}_t and tuples $W^{(t)} = (w_1^{(t)}, w_2^{(t)}, \dots)$, $t \geq 1$, such that they satisfy the following conditions:

- 1) \mathcal{N}_t is a right or a left multiplication,
- 2) $W^{(t)} = (\mathcal{N}_t \circ \dots \circ \mathcal{N}_1) W^{(0)}$,
- 3) if \mathcal{N}_{t+1} , where $t \geq 0$, equals to $R_{ij}^{\pm 1}$ or $L_{ij}^{\pm 1}$, then
 - a) $w_k^{(t+1)} = w_k^{(t)}$ for every $k \neq i$,
 - b) There exists a natural number $n(t)$, such that

$$\begin{aligned} \text{Compl}_{n(t)}(w_i^{(t+1)}) &<_{lex} \text{Compl}_{n(t)}(w_i^{(t)}) \\ \text{Compl}_{n(t)}(w_j^{(t)}) &<_{lex} \text{Compl}_{n(t)}(w_i^{(t)}) \end{aligned}$$

- 4) $\lim_{t \rightarrow \infty} W^{(t)} = (x_1, x_2, \dots)\sigma^{-1}$ for some $\sigma \in \mathcal{M}(X_\omega)$.

Now we show how to define the transformations \mathcal{N}_t . We set $\mathcal{N}_0 = id$. Suppose that $\mathcal{N}_1, \dots, \mathcal{N}_p$ are already defined and $W^{(p)}$ contains x_1, \dots, x_r , up to inversions, but not x_{r+1} .

Since α is an automorphism of F_ω , there exists m such that $x_{r+1} \in \langle w_1^{(p)}, \dots, w_m^{(p)} \rangle$ and x_1, \dots, x_r are contained in $(w_1^{(p)}, \dots, w_m^{(p)})$ up to inversions. By Corollary 4.4.3, there exist elementary Nielsen transformations $\mathcal{N}_{p+1}, \dots, \mathcal{N}_q$, such that they satisfy Conditions 1)–3) and the tuple $W^{(q)} = (\mathcal{N}_q \circ \dots \circ \mathcal{N}_{p+1}) W^{(p)}$ contains x_1, \dots, x_{r+1} up to inversions. Moreover, these transformations do not change the places of the involved tuples with $x_1^{\pm 1}, \dots, x_r^{\pm 1}$. This recursive definition of \mathcal{N}_t will give us $W^{(t)}$, $t \geq 0$, which obviously satisfy 4).

Remark. Fix a natural number n . By 4), there exists p , such that $W^{(p)}$ contains a letter of X^\pm on the place n . Then the elementary Nielsen transformations \mathcal{N}_t , $t > p$, do not have the form $R_{nj}^{\pm 1}$ or $L_{nj}^{\pm 1}$.

b) Now we show that α can be written in the form $\sigma\alpha_f$ for some $f : (-\mathbb{N}) \rightarrow \mathcal{E}$ satisfying Conditions (i) and (ii). We will use σ and the sequence of Nielsen

transformations \mathcal{N}_t , $t \in \mathbb{N}$, which was defined in **a**). Condition 4) can be written as $(\dots \mathcal{N}_2 \circ \mathcal{N}_1) W^{(0)} = X\sigma^{-1}$. Since $W^{(0)} = X\alpha^{-1}$, we have

$$X\sigma^{-1}\alpha = (\dots \mathcal{N}_2 \circ \mathcal{N}_1)X = X(\dots \mathcal{A}_{-2} \mathcal{A}_{-1}),$$

where \mathcal{A}_{-t} are elementary automorphisms of F_ω , defined by the rule: $\mathcal{A}_{-t} = E_{x_i^\epsilon x_j^\tau}$, if $\mathcal{N}_t = T_{i^\epsilon j^\tau}$ for $i, j \in \mathbb{N}$ and $\epsilon, \tau \in \{-1, 1\}$. Then

$$\sigma^{-1}\alpha = \dots \mathcal{A}_{-2} \mathcal{A}_{-1}$$

and we will show that the theorem is satisfied for the function $f : (-\mathbb{N}) \rightarrow \mathcal{E}$, defined by the rule $f(-t) = \mathcal{A}_{-t}$, $t \in \mathbb{N}$.

Condition (i) follows from Remark before **b**).

Now we verify Condition (ii). If it is not satisfied, then there exists an infinite sequence of increasing natural numbers $t_1 < t_2 < t_3 < \dots$, such that $\dots \mathcal{A}_{-t_3}, \mathcal{A}_{-t_2}, \mathcal{A}_{-t_1}$ have the following form:

$$\dots \mathcal{A}_{-t_3} = E_{x_k^\rho x_l^\theta}, \quad \mathcal{A}_{-t_2} = E_{x_j^\delta x_k^\mu}, \quad \mathcal{A}_{-t_1} = E_{x_i^\epsilon x_j^\tau}.$$

Then

$$\dots \mathcal{N}_{t_3} = T_{k^\rho l^\theta}, \quad \mathcal{N}_{t_2} = T_{j^\delta k^\mu}, \quad \mathcal{N}_{t_1} = T_{i^\epsilon j^\tau}.$$

By Condition 3) we have

$$\dots \leq |w_k^{(t_2-1)}| \leq |w_j^{(t_2-1)}| \leq |w_j^{(t_1-1)}| \leq |w_i^{(t_1-1)}|.$$

Since the length cannot infinitely decrease, we may assume, that all these lengths coincide. Then we can apply Lemma 4.4.5 to the following pairs of words and exponents:

$$\begin{aligned} & (w_i^{(t_1-1)}, w_j^{(t_1-1)}), \quad (\epsilon, \tau), \\ & (w_j^{(t_2-1)}, w_k^{(t_2-1)}), \quad (\delta, \mu), \\ & (w_k^{(t_3-1)}, w_l^{(t_3-1)}), \quad (\rho, \theta), \\ & \dots \end{aligned}$$

Note, that the second word in every pair of words coincides with the first word in the next pair. So, we simplify and unify notations:

$$\begin{aligned} & (u_1, u_2), \quad (\sigma_1, \tau_1), \\ & (u_2, u_3), \quad (\sigma_2, \tau_2), \\ & (u_3, u_4), \quad (\sigma_3, \tau_3), \\ & \dots \end{aligned}$$

By applying Lemma 4.4.5 and Remark 4.5.1.1) to pairs (u_i, u_{i+1}) , (σ_i, τ_i) we obtain

$$L(u_{i+1}^{\tau_i}) = L(u_i^{-\sigma_i}) \text{ and } L(u_{i+1}^{-\tau_i}) \prec_\omega L(u_i^{\sigma_i})$$

for all $i \in \mathbb{N}$.

Claim. The set $\bigcup_{i \in \mathbb{N}} \{L(u_i^1), L(u_i^{-1})\}$ contains an infinite subset $\{z_j \mid j \in \mathbb{N}\}$, such that

$$\cdots \prec_\omega z_3 \prec_\omega z_2 \prec_\omega z_1.$$

Proof. We construct a graph \mathcal{G} with oriented edges colored in red and blue. The vertex set of \mathcal{G} is $\bigcup_{i \in \mathbb{N}} \{U_i^1, U_i^{-1}\}$, where U_i^1, U_i^{-1} are formal symbols. The vertices

$U_i^{-\sigma_i}$ and $U_{i+1}^{\tau_i}$ are connected by an oriented red edge and the vertices $U_i^{\sigma_i}$ and $U_{i+1}^{-\tau_i}$ are connected by an oriented blue edge.

There are two infinite paths in this graph starting at U_1^1 and at U_1^{-1} respectively, say p and q . We consider these paths as subgraphs of \mathcal{G} . Since $\mathcal{G} = p \cup q$ and \mathcal{G} contains infinitely many blue edges, we may assume that p contains infinitely many blue edges. Let $(U_i^{\epsilon_i})_{i \in \mathbb{N}}$ be the sequence of vertices of p , where $\epsilon_i \in \{-1, 1\}$. Then the corresponding sequence $(L(u_i^{\epsilon_i}))_{i \in \mathbb{N}}$ of elements of $F(X)$ contains an infinite subsequence $(z_j)_{j \in \mathbb{N}}$, such that $\cdots \prec_\omega z_3 \prec_\omega z_2 \prec_\omega z_1$. \square

Since the words u_i have the same length, the words z_j have the same length too and we have a contradiction to Remark 4.5.1.3). \square

4.6 The second step towards

the surjectivity of $\Psi : \text{BF}(\mathcal{E}, \mathcal{S}) \rightarrow \text{Aut}(F_\omega)$

We fix a basis $X_\omega = \{x_1, x_2, \dots\}$ of F_ω . Recall that $\mathcal{M}(X_\omega)$ is the group of monomial automorphisms of F_ω . Let $\mathcal{I}(X_\omega)$ be the subgroup of $\mathcal{M}(X_\omega)$ consisting of the automorphisms, which invert a part of X_ω and fix the complementary part of X_ω .

Theorem 4.6.1. *Every monomial automorphism of F_ω lies in the image of the homomorphism $\Psi : \text{BF}(\mathcal{E}, \mathcal{S}) \rightarrow \text{Aut}(F_\omega)$.*

Proof. Every monomial automorphism $\alpha \in \text{Aut}(F_\omega)$ can be expressed in the form $\alpha = \beta\gamma$ for some $\beta \in \mathcal{I}(X_\omega)$, $\gamma \in \Sigma(X_\omega)$.

We can express γ as a product of independent countable cycles: $\gamma = \prod_{j \in J} \gamma_j$. By Nielsen, every finite cycle γ_j can be expressed in the form $\gamma_j = \nu_j \gamma'_j$, where ν_j is the identity or the inversion of an element of X_ω , and γ'_j is a finite product of \mathcal{E} -automorphisms. If γ_j is an infinite countable cycle, it is similar to σ in (5). Recall that

$$\sigma = \prod_{i=0}^{\infty} (\tau_{-i, i+1} \tau_{-i, -(i+1)}). \quad (18)$$

One can check, that

$$\begin{aligned} & \tau_{-i,i+1} \tau_{-i,-(i+1)} = \\ & = E_{x_{-(i+1)},x_{-i}} E_{x_{-(i+1)},x_{-i}}^{-1} E_{x_{-(i+1)},x_{i+1}}^{-1} E_{x_{i+1},x_{-(i+1)}}^{-1} \cdot \\ & \quad E_{x_{-(i+1)},x_{i+1}}^{-1} E_{x_{i+1},x_{-i}}^{-1} E_{x_{-i},x_{i+1}}^{-1} E_{x_{i+1},x_{-i}}^{-1}. \end{aligned}$$

So, σ lies in the Ψ -image of an infinite product of \mathcal{E} -letters. It is not hard to check, that this product is admissible, i.e. lies in \mathcal{S} .

Thus, we have to consider an element $\rho \in \mathcal{I}(X_\omega)$. If the support of ρ is infinite, then ρ is an infinite product of some $\rho_j \in \mathcal{I}(X_\omega)$ with disjoint supports of cardinality 2. Each ρ_j is a finite product of \mathcal{E} -automorphisms. So, ρ lies in the Ψ -image of an admissible infinite product of \mathcal{E} -letters. If the support of ρ is finite, then we can write $\rho = \rho' \rho''$ for some $\rho' \rho'' \in \mathcal{I}(X_\omega)$ with infinite supports and so we have reduced to the previous case. \square

Theorem 4.6.2. *The homomorphism $\Psi : \text{BF}(\mathcal{E}, \mathcal{S}) \rightarrow \text{Aut}(F_\omega)$ is surjective.*

Proof. The proof follows immediately from Theorems 4.5.3 and 4.6.1.

4.7 Three topologies on $\text{BF}(\mathcal{E}, \mathcal{S})$

We consider $\text{Aut}(F(X))$ as a topological group with the basis for the neighborhoods of 1 consisting of the subgroups

$$\text{St}(Y) := \{\alpha \in \text{Aut}(F(X)) \mid y\alpha = y \forall y \in Y\},$$

where Y goes over the set of finite subsets of X . The preimage of this topology with respect to the homomorphism $\Psi : \text{BF}(\mathcal{E}, \mathcal{S}) \rightarrow \text{Aut}(F(X))$ gives us a topology on $\text{BF}(\mathcal{E}, \mathcal{S})$; we denote it by $\mathfrak{T}_{\text{stab}}$ and call the *stabilizer topology*.

Recall, that the group $\text{BF}(\mathcal{E}, \mathcal{S})$ also possesses the natural topology (see Section 2.3); we denote it by $\mathfrak{T}_{\text{nat}}$.

Let \mathfrak{T} be the topology on $\text{BF}(\mathcal{E}, \mathcal{S})$ generated by $\mathfrak{T}_{\text{nat}}$ and $\mathfrak{T}_{\text{stab}}$. To be more precise, a basis for the neighborhoods of 1 in \mathfrak{T} consists of the sets

$$\mathcal{U}_{Y,A} = \Psi^{-1}(\text{St}(Y)) \cap \ker \varphi_A,$$

where Y runs through the finite subsets of X and A runs through the finite subsets of \mathcal{E} . Recall that the map $\varphi_A : \text{BF}(\mathcal{E}) \rightarrow \text{BF}(A)$ sends every map $f : S \rightarrow \mathcal{E}$ from $\text{BF}(\mathcal{E})$ to its restriction to $f^{-1}(A)$.

Note that $\mathcal{U}_{Y,A}$ are subgroups of $\text{BF}(\mathcal{E}, \mathcal{S})$ and that $\text{BF}(\mathcal{E}, \mathcal{S})$ is a topological group with respect to \mathfrak{T} .

4.8 A generalized presentation of $\text{Aut}(F_\omega)$

In this section we formulate our main Theorems 4.8.6 and 4.8.11. The first theorem describes the kernel of the epimorphism $\Psi : \text{BF}(\mathcal{E}, \mathcal{S}) \rightarrow \text{Aut}(F_\omega)$ algebraically, while the second one topologically. We deduce Theorem 4.8.11 from Theorem 4.8.6. The proof of Theorem 4.8.6 is based on technical Sections 4.9 and 4.10, and it will be given in Section 4.11. We conclude this section with Theorem 4.8.13 which is a compact reformulation of Theorem 4.8.6.

We fix a basis $X = \{x_1, x_2, \dots\}$ of F_ω . Given a group G and a subset $\mathcal{R} \subseteq G$ we denote by $\langle\langle \mathcal{R} \rangle\rangle_G$ the normal closure of \mathcal{R} in G .

Now we will work with the group $G = \text{BF}(\mathcal{E}, \mathcal{S})$, where $\mathcal{E} = \{E_{xy} \mid x, y \in X^\pm, y \neq x, x^{-1}\}$ and \mathcal{S} is the set of classes of admissible maps defined in Section 4.1. Let G_n be the subset of G , consisting of all classes of maps $f : S \rightarrow \mathcal{E}$ from G , such that $f_1(s) \in \{x_n, x_n^{-1}, x_{n+1}, x_{n+1}^{-1}, \dots\}$ for every $s \in S$. Note that G_n is a subgroup of G and we have $G = G_1 \supset G_2 \supset \dots$, and $\bigcap_{n=1}^{\infty} G_n = 1$. The sequence of subgroups $(G_n)_{n \in \mathbb{N}}$ is called the *filtration on G* .

Remark 4.8.1. *Every neighborhood $\mathcal{U}_{Y,A}$ contains some G_n . Indeed, we can take*

$$n = \max\{k \mid x_k \in Y\} + \max\{k \mid E_{x_k^{\pm 1} *} \in A\} + 1.$$

Definition 4.8.2. A product $\prod_{i \in I} g_i$, where I is a linearly ordered set and $g_i \in G$, $i \in I$, is called *admissible with respect to the filtration on G* , if for every natural n , the reduced forms of all but finitely many g_i lie in G_n . In particular, this product is a well defined element of $\text{BF}(\mathcal{E})$. To stress that the product is admissible, we will write

$$\prod_{\substack{(G_n)\text{-adm.} \\ i \in I}} g_i.$$

Note, that this product may lie outside of G .

By Theorem 4.6.2, there is an epimorphism

$$\Psi : \text{BF}(\mathcal{E}, \mathcal{S}) \rightarrow \text{Aut}(F_\omega).$$

Our aim is to describe a subset $\mathcal{R} \subset \ker \Psi$, which in some sense generates $\ker \Psi$ (see Theorem 4.8.11). For that it is convenient to consider the elements of $\text{BF}(\mathcal{E}, \mathcal{S})$ as (infinite) words in the alphabet \mathcal{E} .

Before we describe some infinite words of \mathcal{R} , we present here their finite analogon. Let $x, y, z \in X^\pm$, such that $y \notin \{x, x^{-1}\}$ and $z \notin \{x, x^{-1}, y, y^{-1}\}$. Then the following relations hold in $\text{Aut}(F_\omega)$:

$$E_{xy}E_{zx} = (E_{zy^{-1}}E_{zx})E_{xy},$$

$$E_{xy}E_{zx^{-1}} = (E_{zx^{-1}}E_{zy})E_{xy},$$

$$E_{xy}E_{zt} = E_{zt}E_{xy}, \quad t \notin \{x, x^{-1}\}.$$

Now we describe an “infinite” relation.

Definition 4.8.3. Let $\alpha = E_{xy}$ be a word consisting of one \mathcal{E} -letter (clearly $\alpha \in \mathcal{S}$). A word $\beta \in \mathcal{S}$ is called α -admissible, if for each of its letters E_{zt} we have $z \notin \{x, x^{-1}, y, y^{-1}\}$.

Let β be an α -admissible word, where $\alpha = E_{xy}$. We denote by β_α the word, obtained from β by the following replacements of each occurrence of E_{zx} and $E_{zx^{-1}}$:

$$E_{zx} \quad \rightsquigarrow \quad E_{zy^{-1}}E_{zx},$$

$$E_{zx^{-1}} \quad \rightsquigarrow \quad E_{zx^{-1}}E_{zy}.$$

It is easy to check, that $\beta_\alpha \in \mathcal{S}$ and that $\Psi(\alpha\beta) = \Psi(\beta_\alpha\alpha)$ holds in $\text{Aut}(F_\omega)$. Thus the following words lie in $\ker \Psi$:

$(\mathcal{R}_{\alpha\beta}) \quad \alpha\beta\alpha^{-1}\beta_\alpha^{-1}$, where $\alpha \in \mathcal{E}$ and β is an α -admissible (infinite) word from G .

Now, let F_n be the free group of finite rank n with basis $\{x_1, \dots, x_n\}$. Consider the canonical epimorphism $\text{Aut}(F_n) \rightarrow \text{GL}_n(\mathbb{Z})$. The full preimage of $\text{SL}_n(\mathbb{Z})$ with respect to this epimorphism is a subgroup of index 2 in $\text{Aut}(F_n)$, which is denoted by $\text{SAut}(F_n)$. Gersten obtained the following presentation for this subgroup.

Theorem 4.8.4. [20]. *A presentation for $\text{SAut}(F_n)$ is given by generators $\{E_{ab} \mid a, b \in X_n^\pm, a \neq b, b^{-1}\}$ subject to relations:*

$$(\mathcal{R}1) \quad E_{ab}^{-1} = E_{ab^{-1}},$$

$$(\mathcal{R}2) \quad [E_{ab}, E_{cd}] = 1 \text{ for } a \neq c, d, d^{-1} \text{ and } b \neq c, c^{-1},$$

$$(\mathcal{R}3) \quad [E_{ab}, E_{bc}] = E_{ac} \text{ for } a \neq c, c^{-1},$$

$$(\mathcal{R}4) \quad w_{ab}^{-1}E_{cd}w_{ab} = E_{\sigma(a)\sigma(b)},$$

where w_{ab} is defined to be $E_{ba}E_{a^{-1}b}E_{b^{-1}a}^{-1}$, and σ is the monomial map, determined by w_{ab} , i.e. $a \mapsto b^{-1}, b \mapsto a$.

$$(\mathcal{R}5) \quad w_{ab}^4 = 1.$$

Definition 4.8.5. We say that a word $W \in \text{BF}(\mathcal{E}, \mathcal{S})$ is of *type* $(\mathcal{R}i)$, if it can be written in the form UV^{-1} , where $U = V$ is the relation $(\mathcal{R}i)$ for some $n \in \mathbb{N}$.

We say that W is of *type* $(\mathcal{R}_{\alpha\beta})$ if it can be written as $\alpha\beta\alpha^{-1}\beta_\alpha^{-1}$, where β is α -admissible. Let \mathcal{R}_{fin} be the set of all words of types $(\mathcal{R}1) - (\mathcal{R}5)$.

Theorem 4.8.6. For $G = \text{BF}(\mathcal{S}, \mathcal{E})$ the kernel of the epimorphism

$$\Psi : G \rightarrow \text{Aut}(F_\omega)$$

coincides with the set of all products UV , where $U, V \in G$ have the form

$$U = \prod_{\substack{(G_n)\text{-adm.} \\ i \in \mathbb{N}}} f_i^{-1} r_i^{\epsilon_i} f_i, \quad V = \prod_{\substack{(G_n)\text{-adm.} \\ i \in (-\mathbb{N})}} f_i^{-1} r_i^{\epsilon_i} f_i, \quad (19)$$

where $f_i \in G$, $r_i \in \mathcal{R}$, $\epsilon_i \in \{-1, 1\}$, and \mathcal{R} is the set of all words of G of types $(\mathcal{R}1) - (\mathcal{R}5)$ and $(\mathcal{R}_{\alpha\beta})$.

Corollary 4.8.7. Let $g \in \ker \Psi$. Then for every natural m there exist elements $h_m \in \langle\langle \mathcal{R} \rangle\rangle_G$ and $g_m \in G_m$, such that $g = h_m g_m$.

Proof. We write $g = UV$ for U, V as in (19). For every natural k we write $U = U_k U'_k$ and $V = V'_k V_k$, where

$$U_k = \prod_{i=1}^k f_i^{-1} r_i^{\epsilon_i} f_i, \quad V_k = \prod_{i=-k}^{-1} f_i^{-1} r_i^{\epsilon_i} f_i.$$

Since the products in (19) are (G_n) -admissible, there exists k such that $U'_k, V'_k \in G_m$. We set $h_m = U_k \cdot (U'_k V'_k) V_k (U'_k V'_k)^{-1}$ and $g_m = U'_k V'_k$. \square

Lemma 4.8.8. Every word $g \in G$ of type $(\mathcal{R}_{\alpha\beta})$ lies in the closure of $\langle\langle \mathcal{R}_{\text{fin}} \rangle\rangle_G$ in the topology \mathfrak{T} .

Proof. Let $g = \alpha\beta\alpha^{-1}\beta_\alpha^{-1}$, where $\alpha = E_{xy}$ and $\beta \in G$ is an α -admissible (infinite) word. We show that an arbitrary neighborhood $g\mathcal{U}$ of g in the topology \mathfrak{T} contains an element $gu \in \langle\langle \mathcal{R}_{\text{fin}} \rangle\rangle_G$. We may assume that $\mathcal{U} = \mathcal{U}_{Y,A}$ for some finite subsets $Y \subset X$ and $A \subset \mathcal{E}$. Let A' be the minimal subset of \mathcal{E} , such that

- 1) $A \cup \{E_{xy}\} \subseteq A'$,
- 2) if $E_{zx^\epsilon} \in A'$ for some $z \in X^{\pm 1}$ and $\epsilon \in \{-1, 1\}$, then $E_{zy^{-\epsilon}} \in A'$.

Clearly, A' is finite. By the choice of A' , we have $\varphi_{A'}(g) = \alpha\beta'\alpha^{-1}(\beta')_\alpha^{-1}$ for $\beta' = \varphi_{A'}(\beta)$. We set $u = g^{-1}\varphi_{A'}(g)$. Obviously, $gu \in \langle\langle \mathcal{R}_{\text{fin}} \rangle\rangle_G$. Since $g \in \ker \Psi$, this implies $u \in \ker \Psi$, and since $\varphi_A \circ \varphi_{A'} = \varphi_A$, we have $u \in \ker \varphi_A$. Hence $u \in \mathcal{U}$. \square

Lemma 4.8.9. $\langle\langle \mathcal{R} \rangle\rangle_G$ lies in the closure of $\langle\langle \mathcal{R}_{\text{fin}} \rangle\rangle_G$ in the topology \mathfrak{T} .

Proof. Let $g \in \langle\langle \mathcal{R} \rangle\rangle_G$ and $\mathcal{U} = \mathcal{U}_{Y,A}$ be a neighborhood of 1 in the group G . Then $g = \prod_{i=1}^k f_i^{-1} r_i f_i$ for some natural k , some $r_i \in \mathcal{R}$ and $f_i \in G$. By Lemma 4.8.8,

$r_i = r'_i u_i$ for some $r'_i \in \langle\langle \mathcal{R}_{\text{fin}} \rangle\rangle_G$ and $u_i \in \mathcal{U}$. Note that $u_i = (r'_i)^{-1} r_i \in \mathcal{U} \cap \ker \Psi$. Since $\mathcal{U} \cap \ker \Psi = \ker \varphi_A \cap \ker \Psi$ is normal in G , we have

$$g \in g'(\mathcal{U} \cap \ker \Psi) \subseteq \langle\langle \mathcal{R}_{\text{fin}} \rangle\rangle_G \mathcal{U}$$

for $g' = \prod_{i=1}^k f_i^{-1} r'_i f_i$. Since this holds for every \mathcal{U} , the proof is completed. \square

Lemma 4.8.10. *$\ker \Psi$ lies in the closure of $\langle\langle \mathcal{R}_{\text{fin}} \rangle\rangle_G$ in the topology \mathfrak{T} .*

Proof. Let $g \in \ker \Psi$ and $\mathcal{U} = \mathcal{U}_{Y,A}$ be a neighborhood of 1 in G . By Remark 4.8.1, there exists m , such that $G_m \subseteq \mathcal{U}$. By Corollary 4.8.7 and Lemma 4.8.9 we have

$$g \in \langle\langle \mathcal{R} \rangle\rangle_G G_m \subseteq \langle\langle \mathcal{R}_{\text{fin}} \rangle\rangle_G \mathcal{U} G_m = \langle\langle \mathcal{R}_{\text{fin}} \rangle\rangle_G \mathcal{U}.$$

\square

Theorem 4.8.11. *For $G = \text{BF}(\mathcal{E}, \mathcal{S})$ the kernel of the epimorphism*

$$\Psi : G \rightarrow \text{Aut}(F_\omega)$$

coincides with the closure of $\langle\langle \mathcal{R}_{\text{fin}} \rangle\rangle_G$ in the topology \mathfrak{T} on G :

$$\ker \Psi = \overline{\langle\langle \mathcal{R}_{\text{fin}} \rangle\rangle_G}^{\mathfrak{T}}.$$

Here \mathcal{R}_{fin} is the set of all words of G of types (R1) – (R5).

In particular, $(\mathcal{E}, \mathcal{S}, \mathfrak{T}, \mathcal{R}_{\text{fin}})$ is a generalized presentation of $\text{Aut}(F_\omega)$ of type \aleph_0 .

Proof. In view of Lemma 4.8.10, it is sufficient to prove that every $g \in \overline{\langle\langle \mathcal{R} \rangle\rangle_G}^{\mathfrak{T}}$ lies in $\ker \Psi$. Fix a natural n . Since $g\mathcal{U}_{\{x_n\}, \emptyset}$ is a neighborhood of g , there exists $u \in \mathcal{U}_{\{x_n\}, \emptyset}$, such that $gu \in \langle\langle \mathcal{R} \rangle\rangle_G$. Then $\Psi(g) = \Psi(u^{-1})$ stabilizes x_n , and since this holds for every n , we have $g \in \ker \Psi$. \square

Remark 4.8.12. *$\ker \Psi$ is properly contained in the topological closure of \mathcal{R} in $\text{BF}(\mathcal{S}, \mathcal{E})$, with respect to the topology $\mathfrak{T}_{\text{nat}}$ defined in Section 4.7. Indeed, the sequence of elements of $\ker \Psi$*

$$E_{x_1 x_2} E_{x_1 x_t} E_{x_t x_2^{-1}} E_{x_1 x_t^{-1}} E_{x_t x_2} \quad (t \in \mathbb{N})$$

converges to $E_{x_1 x_2} \notin \ker \Psi$, when $t \rightarrow \infty$.

Finally we reformulate Theorem 4.8.6 in a compact form.

Theorem 4.8.13. *For $G = \text{BF}(\mathcal{E}, \mathcal{S})$ the kernel of the epimorphism*

$$\Psi : G \rightarrow \text{Aut}(F_\omega)$$

consists of conjugate to elements of the set

$$\left\{ \prod_{\substack{(G_n)\text{-adm.} \\ i \in \mathbb{Z}}} f_i^{-1} r_i^{\pm 1} f_i \mid f_i \in G, r_i \in \mathcal{R} \right\} \cap G, \quad (20)$$

where \mathcal{R} is the set of words of G of types (R1) – (R5) and $(\mathcal{R}_{\alpha\beta})$.

Proof. Let $g \in \ker \Psi$. Then $g = UV$ for U, V as in Theorem 4.8.6, and we can write $g = U(VU)U^{-1}$, where VU lies in the set (20). The inverse inclusion follows from the next lemma. \square

Lemma 4.8.14. *Let I be a linearly ordered set and $g_i \in \ker \Psi$ for any $i \in I$. If*

$$g = \prod_{\substack{(G_n)\text{-adm.} \\ i \in I}} g_i, \quad (21)$$

and $g \in G$, then $g \in \ker \Psi$.

Proof. Fix $x_r \in X$. Since the product (21) is (G_n) -admissible, the set $\{i \in I \mid g_i \notin G_{r+1}\}$ is finite. Let $i_1 < i_2 < \dots < i_k$ be its elements. Denote

$$h_0 = \prod_{\substack{(G_n)\text{-adm.} \\ \{i \in I \mid i < i_1\}}} g_i, \quad h_k = \prod_{\substack{(G_n)\text{-adm.} \\ \{i \in I \mid i_k < i\}}} g_i, \quad h_l = \prod_{\substack{(G_n)\text{-adm.} \\ \{i \in I \mid i_l < i < i_{l+1}\}}} g_i,$$

where $l = 1, \dots, k-1$. Then $g = h_0 g_{i_1} h_1 g_{i_2} \dots g_{i_k} h_k$. Since $g \in G$ and every (infinite) subword of a word from G also lies in G , we have $h_j \in G$ for $j = 0, \dots, k$. Then all h_i belong to G_{r+1} , and hence $\Psi(h_i)$ stabilize x_r . Moreover, $\Psi(g_i)$ is the identity for any $i \in I$ by assumption. Therefore $\Psi(g)$ stabilizes x_r . Since this holds for every r , we have $g \in \ker \Psi$. \square

Remark 4.8.15. The following example shows that the assumption $g \in G$ in Lemma 4.8.14 cannot be omitted:

$$g = \prod_{i \in \mathbb{N}} [E_{x_{4i-1}, x_{4i}}, E_{x_{4i+1}, x_{4i+2}}].$$

Indeed, $[E_{x_{4i-1}, x_{4i}}, E_{x_{4i+1}, x_{4i+2}}] \in \ker \Psi$ for every $i \in \mathbb{N}$, but $g \notin G$, and so $g \notin \ker \Psi$.

4.9 An inequality for the length of chains

Definition 4.9.1. Let $f : S \rightarrow \mathcal{E}$ be an element of $\text{BF}(\mathcal{E})$ and let $x \in X$. A sequence of elements of S , (\dots, s_2, s_1) , is called a *backward x -chain for f* , if the following conditions hold:

- 1) $\dots \prec s_2 \prec s_1$,
- 2) $f_1(s_1) \in \{x, x^{-1}\}$,
- 3) $f_1(s_{i+1}) \in \{f_2(s_i), f_2(s_i)^{-1}\}$ for $i \geq 1$.

We denote by $M(f, x)$ the supremum of lengths of backward x -chains for f . Note that $M(f, x) \leq n^+(f, x)$; hence $M(f, x)$ is finite for every f in \mathcal{S} with the notation from Section 4.2.

Proposition 4.9.2. *Let $f = \delta\alpha\beta\gamma$ be a word from \mathcal{S} , such that β is α -admissible; let $f' = \delta\beta_\alpha\alpha\gamma$. Then $M(f', x) \leq M(f, x)$ for every $x \in X^\pm$.*

To facilitate the understanding of the proof look at the following example.

Example 4.9.3.

$$f = (\dots E_{x_2x_5} \dots) \overbrace{E_{x_1x_2}}^\alpha \left(\overbrace{\dots E_{x_3x_1} \dots E_{x_3x_1^{-1}} \dots}^\beta \right) (\dots E_{x_4x_1^{-1}} \dots E_{x_4x_3^{-1}} \dots).$$

$$f' = (\dots E_{x_2x_5} \dots) \left(\overbrace{\dots E_{x_3x_2^{-1}} E_{x_3x_1} \dots E_{x_3x_1^{-1}} E_{x_3x_2} \dots}^{\beta_\alpha} \right) \overbrace{E_{x_1x_2}}^\alpha (\dots E_{x_4x_1^{-1}} \dots E_{x_4x_3^{-1}} \dots).$$

We write down 3 backward x_4 -chains in f :

$$C : \quad (a, b_2, c_2), (d, a, b_1, c_2), (d, a, c_1),$$

and 3 backward x_4 -chains in f' :

$$C' : \quad (b_2, c_2), (d, b'_1, c_2), (d, a', c_1).$$

Proof of Proposition 4.9.2. Let $f : (D \sqcup \{a\} \sqcup B \sqcup C) \rightarrow \mathcal{E}$ be the map corresponding to the product $f = \delta\alpha\beta\gamma$ and let $f' : (D \sqcup B' \sqcup \{a'\} \sqcup C) \rightarrow \mathcal{E}$ be the map corresponding to the product $f' = \delta\beta_\alpha\alpha\gamma$. As in the example above, we assume, that B' is obtained from B by inserting some b' -letters.

Let C' be a backward x -chain for f' . It is sufficient to indicate a backward x -chain \mathcal{C} for f , which is not shorter than C' . Note that if C' contains a b'_i -letter, then all smaller letters of this chain lie in D .

Case 1. Suppose that C' does not contain a' and does not contain any b'_i -letter. Then we set $\mathcal{C} = C'$.

Case 2. Suppose that C' does not contain a' , but contains some b'_i -letter. Then we define \mathcal{C} to be the chain obtained from C' by replacing b'_i by two letters a, b_i .

Case 3. Suppose that C' contains a' .

In this case C' does not contain elements of B' . Then we define \mathcal{C} to be the chain obtained from C' by replacing a' by a . \square

4.10 Transformation of elements of $\text{BF}(\mathcal{E}, \mathcal{S})$ to elements of $\text{BF}_{-\mathbb{N}}(\mathcal{E}, \mathcal{S})$ modulo $\ker \Psi$

We denote by $\text{BF}_{-\mathbb{N}}(\mathcal{E}, \mathcal{S})$ the subset of $G = \text{BF}(\mathcal{E}, \mathcal{S})$ consisting of classes of all maps $g : -\mathbb{N} \rightarrow \mathcal{E}$. The main aim of this section is to prove that any element

$f \in \text{BF}(\mathcal{E}, \mathcal{S})$ can be represented in the form $f = RA$ for some $R \in \ker \Psi$ and $A \in \text{BF}_{-\mathbb{N}}(\mathcal{E}, \mathcal{S})$ (see Proposition 4.10.4).

In Section 4.8 we introduced the chain of subgroups $G = G_1 \supseteq G_2 \supseteq \dots$, such that $\bigcap_{n=1}^{\infty} G_n = 1$. Now we decompose each difference $G_n \setminus G_{n+1}$ into smaller subsets. By Lemma 4.2.1, the number $n^+(f, x_n)$ is finite. Now, for every $m \geq 1$ we set

$$G_{n,m} = \{f \in G_n \setminus G_{n+1} \mid n^+(f, x_n) = m\}.$$

Clearly $G_n \setminus G_{n+1} = \bigsqcup_{m \geq 1} G_{n,m}$.

Definition 4.10.1. (*split and derived forms of f*) Suppose that $f \in G_n$ for some n . Denote $n^+ = n^+(f, x_n)$ and set

$$S_1 = \{s \in S \mid s \prec n^+\}, \quad S_2 = \{s \in S \mid n^+ \prec s\}.$$

Then we can write f in the form $f = \delta\alpha\beta$, where $\delta = f|_{S_1}$, $\alpha = f|_{n^+}$ and $\beta = f|_{S_2}$. This form will be called the *split form* of f .

Note that β is α -admissible (see Definition 4.8.3). Using the definition of β_α given after this definition, we rewrite f in the form $f = r\bar{f}\alpha$, where $r = \delta(\alpha\beta\alpha^{-1}\beta_\alpha^{-1})\delta^{-1}$ and $\bar{f} = \delta\beta_\alpha$. The expression $f = r\bar{f}\alpha$ will be called the *derived form* of f .

For a subgroup B of a group C we denote by B^C the normal closure of B in C .

Lemma 4.10.2. *Let $f \in G_{n,m}$ and let $f = r\bar{f}\alpha$ be the derived form of f . Then the following properties are satisfied.*

- P1.** $r \in (G_n \cap \mathcal{R})^{G_n}$.
- P2.** $\bar{f} \in G_{n,m-1}$ if $m > 1$ and $\bar{f} \in G_{n+1}$ if $m = 1$.
- P3.** $\alpha \in G_n$.

Proof. The proof is straightforward. □

We can continue and rewrite the element \bar{f} in the derived form. We want to do that countably many times. So, we put $f^{(1)} = f$ and define three sequences of elements of G : $(f^{(i)})_{i \geq 1}$, $(r^{(i)})_{i \geq 1}$ and $(\alpha^{(i)})_{i \geq 1}$, such that

$$f^{(i)} = r^{(i)}\overline{f^{(i)}}\alpha^{(i)} \tag{22}$$

is the derived form of $f^{(i)}$ and

$$f^{(i+1)} = \overline{f^{(i)}}.$$

We have

$$f = (r^{(1)}r^{(2)} \dots r^{(i)})f^{(i+1)}(\alpha^{(i)} \dots \alpha^{(2)}\alpha^{(1)}). \tag{23}$$

Lemma 4.10.3. *Let $f = f^{(1)} \in G_{n,m}$. Then the following properties are satisfied for every $1 \leq i \leq m$.*

P4. $f^{(i)} \in G_{n,m-i+1}$; moreover $f^{(m+1)} \in G_{n+1}$.

P5. $r^{(i)} \in (G_n \cap \mathcal{R})^{G_n}$; moreover $r^{(m+1)} \in (G_{n+1} \cap \mathcal{R})^{G_{n+1}}$.

P6. $\alpha^{(i)} \in G_n$; moreover $\alpha^{(m+1)} \in G_{n+1}$.

Proof. Property P4 follows by induction from P2. Properties P5 and P6 follow from P4 and (22). \square

Proposition 4.10.4. *For every element $f \in \text{BF}(\mathcal{E}, \mathcal{S})$ the following claims hold.*

1) *The sequence $(f^{(i)})_{i \geq 1}$ converges to 1 in the topology $\mathfrak{T}_{\text{nat}}$ on $\text{BF}(\mathcal{E}, \mathcal{S})$, when $i \rightarrow \infty$.*

2) *The product $R = \prod_{i \in \mathbb{N}} r^{(i)}$ is well defined, i.e. it belongs to $\text{BF}(\mathcal{E})$.*

3) *The product $A = \prod_{i \in -\mathbb{N}} \alpha^{(-i)}$ belongs to $\text{BF}_{-\mathbb{N}}(\mathcal{E}, \mathcal{S})$.*

4) $f = RA$.

5) *The product $R = \prod_{i=1}^{\infty} r^{(i)}$ is admissible with respect to the filtration $(G_n)_{n \geq 1}$ on $\text{BF}(\mathcal{E}, \mathcal{S})$ and it belongs to $\text{BF}(\mathcal{E}, \mathcal{S})$.*

6) $R \in \ker \Psi$.

Proof. Claims 1) and 2) follow from P4 and P5. Now we prove the most difficult Claim 3). Clearly A is a map from $(-\mathbb{N})$ to \mathcal{E} . By Property P6 we have $A \in \text{BF}(\mathcal{E})$. We have to verify, that for every $x \in X$, the cardinal number $M(A, x)$ is finite.

Let $f^{(k)} = \delta^{(k)} \alpha^{(k)} \beta^{(k)}$ be the split form of $f^{(k)}$. By definition, we have

$$f^{(k+1)} = \overline{f^{(k)}} = \delta^{(k)} (\beta^{(k)})_{\alpha^{(k)}}.$$

Denote $\gamma^{(k)} = \alpha^{(k-1)} \dots \alpha^{(1)}$. Then we have

$$f^{(k)} \gamma^{(k)} = \delta^{(k)} \alpha^{(k)} \beta^{(k)} \gamma^{(k)}$$

and

$$f^{(k+1)} \gamma^{(k+1)} = \delta^{(k)} (\beta^{(k)})_{\alpha^{(k)}} \alpha^{(k)} \gamma^{(k)}.$$

Applying Proposition 4.9.2 to these two words we get

$$M(f^{(k+1)} \gamma^{(k+1)}, x) \leq M(f^{(k)} \gamma^{(k)}, x).$$

for every $x \in X^{\pm 1}$. Since $f = f^{(1)} \gamma^{(1)}$, we have by induction

$$M(f^{(k)} \gamma^{(k)}, x) \leq M(f, x)$$

and so $M(\gamma^{(k)}, x) \leq M(f, x)$ for every $k \geq 1$. This implies $M(A, x) \leq M(f, x)$.

Claim 4) follows from Equation (23) by taking the limit with respect to the natural topology on $\text{BF}(\mathcal{E})$ and using Claims 1)-3).

5) It is sufficient to prove that $R \in \text{BF}(\mathcal{E}, \mathcal{S})$. But this follows from $R = fA^{-1}$ by the assumption on f and by Claim 3).

6) This follows from 5) and Lemma 4.8.14. \square

4.11 Proof of Theorem 4.8.6 about $\ker \Psi$

The following technical lemma is used in Lemma 4.11.2.

Lemma 4.11.1. *Let $W : (-\mathbb{N}) \rightarrow \mathcal{E}$ be an infinite word from $G_k \cap \ker \Psi$. Then there exists an infinite word $W_2 : (-\mathbb{N}) \rightarrow \mathcal{E}$ from $G_{k+1} \cap \ker \Psi$, and a finite word $W_1 \in G_k \cap \ker \Psi$, such that $W = W_2W_1$.*

Proof. We write $W = BC$, where C is a finite subword of W of minimal length which contains all letters E_{xy} with $x \in \{x_k, x_k^{-1}\}$. Clearly $B \in G_{k+1}$, in particular $\Psi(B)$ stabilizes F_k . Let

$$n = \max\{i, j \mid E_{x_i x_j} \text{ is a letter of } C\} + k.$$

Obviously, $n \geq k$. Then $\Psi(C)$ is an automorphism of F_n , which stabilizes F_k (this follows from $W = BC$, $\Psi(W) = 1$, and the fact that $\Psi(B)$ stabilizes F_k). By Proposition 4.12.6, there exists a finite word C' in letters E_{xy} with $x \in \{x_{k+1}, \dots, x_n\}^{\pm 1}$ and $y \in \{x_1, \dots, x_n\}^{\pm 1}$, such that $\Psi(C) = \Psi(C')$ in $\text{Aut}(F_n)$. We set $W_2 = BC'$ and $W_1 = (C')^{-1}C$. Then $W_1 \in G_k \cap \ker \Psi$. Since $W = W_2W_1 \in \ker \Psi$, we conclude that $W_2 \in G_{k+1} \cap \ker \Psi$. \square

Lemma 4.11.2. *Let $A : (-\mathbb{N}) \rightarrow \mathcal{E}$ be an infinite word from $\ker \Psi$. Then it can be written as*

$$A = \prod_{\substack{(G_n)\text{-adm.} \\ i \in (-\mathbb{N})}} f_i^{-1} r_i^{\pm 1} f_i, \quad (24)$$

for some finite words f_i and r_i in the alphabet \mathcal{E} , where r_i 's are words of types $(\mathcal{R}1) - (\mathcal{R}5)$.

Proof. By Lemma 4.11.1, we can write A as $A = (\dots A_3 A_2 A_1]$ for some finite words $A_k \in G_k \cap \ker \Psi$. By Proposition 4.12.6, A_k can be written in the form

$$A_k = \prod_{j \in J(k)} f_j^{-1} r_j^{\pm 1} f_j,$$

where $J(k)$ is a finite set, f_j and r_j are finite words over \mathcal{E} , which belong to G_k , and r_j 's are words of types $(\mathcal{R}1) - (\mathcal{R}5)$. This completes the proof. \square

Proof of Theorem 4.8.6. Let $f \in \ker \Psi$. We show that f can be written in the form $f = UV$ with U, V satisfying the conclusion of Theorem 4.8.6. First, we write $f = RA$, where R and A as in Proposition 4.10.4. In particular, $A : (-\mathbb{N}) \rightarrow \mathcal{E}$ is an infinite word from G and

$$R \in \left\{ \prod_{\substack{(G_n)\text{-adm.} \\ i \in \mathbb{N}}} f_i^{-1} r_i^{\pm 1} f_i \mid f_i \in G, r_i \in \mathcal{R} \right\} \cap G.$$

Since $R \in \ker \Psi$ by Proposition 4.10.4, we have $A \in \ker \Psi$. Then we can apply Lemma 4.11.2 and write A in the form (24). So, we can set $U = R, V = A$.

Conversely, if $f = UV$ with U, V satisfying the conclusion of Theorem 4.8.6, then $f \in \ker \Psi$ by Lemma 4.8.14. \square

4.12 Appendix: Finite presentations for some stabilizers in $\text{SAut}(F_n)$

Let F_n be the free group with the basis $X_n = \{x_1, \dots, x_n\}$, where n is finite. If $a, b \in X_n^\pm$ and $a \neq b, b^{-1}$, the *Nielsen map* E_{ba} is defined by the rule:

$$\begin{aligned} b &\mapsto ba, \\ x &\mapsto x \quad \text{if } x \in X_n^\pm \setminus \{b, b^{-1}\}. \end{aligned}$$

An automorphism of F_n is called *monomial*, if it permutes the set $X_n \cup X_n^{-1}$. Let \mathcal{M}_n be the group of monomial automorphisms. If $a, b \in X_n^\pm$ and $a \neq b, b^{-1}$, the monomial automorphism w_{ab} is defined by the rules:

$$\begin{aligned} a &\mapsto b^{-1} \\ b &\mapsto a \\ x &\mapsto x \quad \text{if } x \in X_n^\pm \setminus \{a, b, a^{-1}, b^{-1}\}. \end{aligned}$$

If $A \subseteq X_n^\pm$, and $a \in A, a^{-1} \notin A$, the *Whitehead automorphism* (A, a) is defined by the rules

$$\begin{aligned} a &\mapsto a, \\ x &\mapsto xa \quad \text{if } x \neq a, x \in A, x^{-1} \notin A, \\ x &\mapsto a^{-1}x \quad \text{if } x \neq a, x^{-1} \in A, x \notin A, \\ x &\mapsto a^{-1}xa \quad \text{if both } x \text{ and } x^{-1} \text{ are in } A, \\ x &\mapsto x \quad \text{if neither } x \text{ nor } x^{-1} \text{ is in } A. \end{aligned}$$

Let \mathcal{W}_n denote the set of all Whitehead automorphisms (A, a) .

Observe that E_{ba} coincides with the Whitehead map $(\{b, a\}, a)$ and that

$$(A, a) = \prod_{\substack{b \in A \\ b \neq a}} E_{ba}. \tag{25}$$

We now recall McCool's presentation for $\text{Aut}(F_n)$.

Theorem 4.12.1. [28] *A finite presentation for $\text{Aut}(F_n)$ is given by generators $\mathcal{M}_n \cup \mathcal{W}_n$ and relations:*

- (M0) *A defining set of relations for \mathcal{M}_n .*
- (M1) $(A, a)^{-1} = (A - a + a^{-1}, a^{-1})$.
- (M2) $(A, a)(B, a) = (A \cup B, a)$ where $A \cap B = \{a\}$.
- (M3) $(A, a)(B, b) = (B, b)(A, a)$ where $A \cap B = \emptyset$, $a^{-1} \notin B$, $b^{-1} \notin A$.
- (M4) $(A, a)(B, b) = (B, b)(A + B - b, a)$ where $A \cap B = \emptyset$, $a^{-1} \notin B$, $b^{-1} \in A$.
- (M5) $(A, a)(A - a + a^{-1}, b) = w_{ab}(A - b + b^{-1}, a)$ with $w_{ab} \in \mathcal{M}_n$ as above, $b \in A$, $b^{-1} \notin A$, $a \neq b$.
- (M6) $\sigma^{-1}(A, a)\sigma = (A\sigma, a\sigma)$ for $\sigma \in \mathcal{M}_n$.

Using Expression (25) and rewriting McCool's relations in terms of Nielsen automorphisms E_{ba} , Gersten deduced in [20, Theorem 1.2] the following finite presentations of $\text{Aut}(F_n)$.

Theorem 4.12.2. [20]. *A presentation for $\text{Aut}(F_n)$ is given by generators $\{E_{ab} \mid a, b \in X_n^\pm, a \neq b, b^{-1}\} \cup \mathcal{M}_n$ subject to relations:*

- (S0) *A defining set of relations for \mathcal{M}_n ,*
- (S1) $E_{ab}^{-1} = E_{ab^{-1}}$,
- (S2) $[E_{ab}, E_{cd}] = 1$ for $a \neq c, d, d^{-1}$ and $b \neq c, c^{-1}$,
- (S3) $[E_{ab}, E_{bc}] = E_{ac}$ for $a \neq c, c^{-1}$,
- (S4) $w_{ab} = E_{ba}E_{a^{-1}b}E_{b^{-1}a^{-1}}$
- (S5) $\sigma^{-1}E_{ab}\sigma = E_{\sigma(a)\sigma(b)}$, $\sigma \in \mathcal{M}_n$.

By applying the Reidemeister-Schreier method, Gersten obtained in [20, Theorem 1.4] the following presentation for $\text{SAut}(F_n)$.

Theorem 4.12.3. [20]. *A presentation for $\text{SAut}(F_n)$ is given by generators $\{E_{ab} \mid a, b \in X_n^{\pm 1}, a \neq b, b^{-1}\}$ subject to relations:*

- (R1) $E_{ab}^{-1} = E_{ab^{-1}}$,
- (R2) $[E_{ab}, E_{cd}] = 1$ for $a \neq c, d, d^{-1}$ and $b \neq c, c^{-1}$,
- (R3) $[E_{ab}, E_{bc}] = E_{ac}$ for $a \neq c, c^{-1}$,

(R4) $w_{ab}^{-1}E_{cd}w_{ab} = E_{\sigma(a)\sigma(b)}$,
 where w_{ab} is defined to be $E_{ba}E_{a^{-1}b}E_{b^{-1}a}^{-1}$, and σ is the monomial map,
 determined by w_{ab} , i.e. $a \mapsto b^{-1}, b \mapsto a$.

(R5) $w_{ab}^4 = 1$.

Notation 4.12.4. Let F_n be the free group with basis $X_n = \{x_1, \dots, x_n\}$. For a subset $Y \subseteq F_n$ we denote by $\text{St}_{\text{Aut}(F_n)}(Y)$ the pointwise stabilizer of Y in $\text{Aut}(F_n)$ and by $\text{St}_{\text{SAut}(F_n)}(Y)$ the pointwise stabilizer of Y in $\text{SAut}(F_n)$.

In [29], McCool proved that the group $\text{St}_{\text{Aut}(F_n)}(Y)$ is finitely presented for every finite $Y \subseteq F_n$ and gave an algorithm for finding such a presentation (see also Proposition 5.7 in Chapter 1 of [25] and a remark after it).

We are specially interested in finding presentations of $\text{St}_{\text{Aut}(F_n)}(X_k)$ and $\text{St}_{\text{SAut}(F_n)}(X_k)$ for $X_k = \{x_1, x_2, \dots, x_k\}$, where $1 \leq k \leq n$. Denote

$$\mathcal{M}_{n,k} = \mathcal{M}_n \cap \text{St}_{\text{Aut}(F_n)}(X_k)$$

and

$$\mathcal{W}_{n,k} = \mathcal{W}_n \cap \text{St}_{\text{Aut}(F_n)}(X_k).$$

So, $\mathcal{W}_{n,k}$ consists of those $(A, a) \in \mathcal{W}_n$, for which

$$A \setminus \{a\} \subseteq \{x_{k+1}, \dots, x_n\}^{\pm}. \quad (26)$$

Proposition 4.12.5. *Let $X_n = \{x_1, \dots, x_n\}$. For every $1 \leq k \leq n$, a finite presentation for $\text{St}_{\text{Aut}(F_n)}(X_k)$ is given by generators $\mathcal{M}_{n,k} \cup \mathcal{W}_{n,k}$ subject to the relations of (M0) – (M6) which contain only these generators.*

Proof. By the result of McCool, $\text{St}_{\text{Aut}(F_n)}(X_k)$ is isomorphic to the fundamental group of the following 2-dimensional simplicial complex \mathcal{K} .

The vertices of \mathcal{K} are k -element ordered subsets of X_n^{\pm} , i.e. they have the form $(x_{i_1}^{\epsilon_1}, \dots, x_{i_k}^{\epsilon_k})$, where $x_{i_j} \in X_n$ are different and $\epsilon_{i_j} \in \{1, -1\}$. We distinguish the vertex $\mathbf{x} = (x_1, \dots, x_k)$. Two vertices (u_1, \dots, u_k) and (v_1, \dots, v_k) are joint by an edge with label α if α is a Whitehead automorphism of type (A, a) or a monomial automorphism from \mathcal{M}_n , such that $v_j = u_j \alpha$ for $j = 1, \dots, k$. The inverse edge is labeled by α^{-1} . The initial vertex of an edge e is denoted by $i(e)$ and the terminal one by $t(e)$.

Let $\phi(e)$ denote the label of an edge e . This labeling can be obviously extended to paths in \mathcal{K} : if $p = e_1 e_2 \dots e_m$ is a path in \mathcal{K} , then $\phi(p) = \phi(e_1) \phi(e_2) \dots \phi(e_m)$. By a loop we understand a cyclically ordered sequence of edges (e_1, e_2, \dots, e_m) , such that $i(e_{j+1}) = t(e_j)$, $j = 1, \dots, m$, where the indexes are added modulo m . We glue a 2-cell along a loop $p = (e_1, e_2, \dots, e_m)$ in \mathcal{K} if there exists a relation $r = s$ from (M0) – (M6), such that the cyclic sequence $(\phi(e_1), \phi(e_2), \dots, \phi(e_m))$ coincides with the cyclic word rs^{-1} .

Clearly, if p and p' are homotopic paths in \mathcal{K} , then $\phi(p) = \phi(p')$. Moreover, if p is a closed path at \mathbf{x} in \mathcal{K} , then $\phi(p)$ is an element of $\text{St}_{\text{Aut}(F_n)}(X_k)$. Thus, ϕ induces a homomorphism $\Phi : \pi_1(\mathcal{K}, \mathbf{x}) \rightarrow \text{St}_{\text{Aut}(F_n)}(X_k)$. By the cited general result of McCool, Φ is an isomorphism.

To describe $\pi_1(\mathcal{K}, x)$, we choose a maximal subtree T in \mathcal{K} in the following way. For every vertex $y \neq \mathbf{x}$ of \mathcal{K} , we choose an edge e_y from \mathbf{x} to y , such that $\phi(e_y) \in \mathcal{M}_n$. Let T be the maximal subtree in the 1-skeleton of \mathcal{K} , consisting of all vertices and all these edges. For convenience we introduce the formal symbol $e_{\mathbf{x}}$, which we identify with \emptyset .

The generators of $\pi_1(\mathcal{K}, \mathbf{x})$ are the homotopy classes $\gamma(f)$ of paths $e_{i(f)} f e_{t(f)}^{-1}$, where f runs through the edges of \mathcal{K} .

For every 2-cell in \mathcal{K} with the boundary $f_1 f_2 \dots f_m$ we write the corresponding relation $\gamma(f_1) \gamma(f_2) \dots \gamma(f_m)$. These relations together with the trivial relations $\gamma(f)^{-1} = \gamma(f^{-1})$ form the complete set of defining relations for the chosen set of generators of $\pi_1(\mathcal{K}, \mathbf{x})$.

By applying Φ , we get generators and defining relations for $\text{St}_{\text{Aut}(F_n)}(X_k)$. Now we describe precisely these generators and relations.

Generators. Let f be an edge. Denote $\sigma = \phi(e_{i(f)})$ and $\tau = \phi(e_{t(f)})$. By definition of T we have $\sigma, \tau \in \mathcal{M}_n$.

a) Suppose that $\phi(f) = (A, a) \in \mathcal{W}_n$. Then the initial and the terminal vertices of f coincide. It follows that $\phi(\gamma(f)) = \sigma \cdot (A, a) \cdot \sigma^{-1} = (A\sigma^{-1}, a\sigma^{-1})$ is an automorphism from $\mathcal{W}_{n,k}$.

b) Suppose that $\phi(f) = \nu \in \mathcal{M}_n$. Then $\phi(\gamma(f)) = \sigma \nu \tau^{-1} \in \mathcal{M}_{n,k}$.

Thus, $\mathcal{M}_{n,k} \cup \mathcal{W}_{n,k}$ is the generator set for $\text{St}_{\text{Aut}(F_n)}(X_k)$.

Relations. Let Δ be a 2-cell in \mathcal{K} with the boundary $f_1 f_2 \dots f_m$. By construction, $\phi(f_1) \phi(f_2) \dots \phi(f_m)$ is one of the relations **(M0)** – **(M6)** (up to a rewriting of kind $r = s \rightsquigarrow r s^{-1}$), say **(Mi)**. The corresponding relation for $\text{St}_{\text{Aut}(F_n)}(X_k)$ is $\phi(\gamma(f_1)) \phi(\gamma(f_2)) \dots \phi(\gamma(f_m))$, which is obviously a word of length m in elements of $\mathcal{M}_{n,k} \cup \mathcal{W}_{n,k}$. We claim, that it also has the form **(Mi)**.

For instance, consider a 2-cell with boundary $f_1 f_2 f_3 f_4^{-1}$, such that the equation

$$\phi(f_1) \phi(f_2) \phi(f_3) = \phi(f_4)$$

has the form **(M6)**: $\sigma^{-1} \cdot (A, a) \cdot \sigma = (A\sigma, a\sigma)$.

As explained above, f_2 and f_4 are loops. Denoting $\tau = \phi(e_{i(f_1)})$, $\delta = \phi(e_{i(f_2)})$, we can write the corresponding relation

$$\phi(\gamma(f_1)) \cdot \phi(\gamma(f_2)) \cdot \phi(\gamma(f_3)) = \phi(\gamma(f_4))$$

in $\text{St}_{\text{Aut}(F_n)}(X_k)$:

$$\tau \sigma^{-1} \delta^{-1} \cdot (\delta(A, a) \delta^{-1}) \cdot \delta \sigma \tau^{-1} = \tau(A\sigma, a\sigma) \tau^{-1}.$$

This is exactly

$$\tau\sigma^{-1}\delta^{-1} \cdot (A\delta^{-1}, a\delta^{-1}) \cdot \delta\sigma\tau^{-1} = (A\sigma\tau^{-1}, a\sigma\tau^{-1})$$

and it has the form **(M6)**. The other cases can be considered similarly. \square

Proposition 4.12.6. *Let $X_n = \{x_1, \dots, x_n\}$. For every $1 \leq k \leq n$, a finite presentation for $\text{St}_{\text{SAut}(F_n)}(X_k)$ is given by generators*

$$\{E_{ab} \mid a \in \{x_{k+1}, \dots, x_n\}^\pm, b \in X_n^\pm, a \neq b, b^{-1}\} \quad (27)$$

*subject to those relations **(R1)** – **(R5)** which contain only these generators.*

Proof. We deduce this proposition from Proposition 4.12.5. Using (25) we conclude, that $\text{St}_{\text{Aut}(F_n)}(X_k)$ is generated by $\mathcal{M}_{n,k}$ and all E_{ba} with

$$b \in \{x_{k+1}, \dots, x_n\}^\pm. \quad (28)$$

Now we can rewrite the relations in Proposition 4.12.5 in terms of these generators. We should do that exactly as Gersten in his proof of [20, Theorem 1.2]. As a result, we deduce that $\text{St}_{\text{Aut}(F_n)}(X_k)$ has the presentation as in Theorem 4.12.2 with the only restriction, that all generators of type E_{ba} satisfy Condition (28). Finally, we apply Reidemeister-Schreier method to obtain a presentation for the subgroup $\text{St}_{\text{SAut}(F_n)}(X_k)$. Arguing as in the proof of [20, Theorem 1.2], we complete our proof. \square

5 Acknowledgements

We thank M. Bridson, A. Klyachko, D. Segal and P. Zalesski for helpful discussions. The first named author thanks the MPIM at Bonn for its support and excellent working conditions during the fall 2010, while this research was finished.

References

- [1] W. A. Bogley and A. J. Sieradski, *Weighted combinatorial group theory and wild metric complexes*, Groups-Korea 98, edited by Y.G. Baik, D.L. Johnson, and A.C. Kim (de Gruyter, Berlin, New York, 2000), 53-80.
- [2] W. A. Bogley and A. J. Sieradski, *Omega-groups II: Weighted presentations for omega-groups*, preprint.
- [3] W. A. Bogley and A. J. Sieradski, *Omega groups III: Inverse limits and profinite groups*, preprint.

- [4] W. A. Bogley and A. J. Sieradski, *Universal path spaces*, preprint. Available at: <http://people.oregonstate.edu/~bogleyw/research/ups.pdf>
- [5] O. Bogopolski, *Introduction to group theory*, EMS: Zürich, 2008.
- [6] O. Bogopolski, A. Zastrow, *An infinite commutator-product is not automatically trivial in the homology*, Max-Planck-Institute of Mathematics Preprint Series, no. **125** (2008), 24 pages.
- [7] R.M. Bryant, D.M. Evans, *The small index property for free groups and relatively free groups*, J. London Math. Soc. (2) **55** (1997), 363-369.
- [8] R.G. Burns, L. Pi, *Generators for the bounded automorphisms of infinite-rank free nilpotent groups*, Bull. Austral. Math. Soc., **40** (1989), 175-187.
- [9] J.W. Cannon and G.R. Conner, *The combinatorial structure of the Hawaiian earring group*, Topology Appl., **106** (3) (2000), 225-271.
- [10] J.W. Cannon and G.R. Conner, *On the fundamental groups of one-dimensional spaces*, Topology Appl., **153** (2006), 2648-2672.
- [11] J.W. Cannon and G.R. Conner, *The big fundamental group, big Hawaiian earrings, and the big free product*, Topology Appl., **106** (3) (2000), 273-291.
- [12] J.M. Cohen, *Aspherical 2-complexes*, J. Pure Appl. Algebra **12** (1978), 101-110.
- [13] R. Cohen, *Classes of automorphisms of free groups of infinite rank*, Trans. Am. Math. Soc., **177**, (1973), 99-120.
- [14] R. Diestel and P. Sprüssel, *The fundamental group of a locally finite graph with ends*, Preprint 2008. Available at: <http://www.math.uni-hamburg.de/home/diestel/>
- [15] K. Eda, *Free σ -products and noncommutatively slender groups*, J. Algebra, **148** (1992), 243-263.
- [16] K. Eda, *Free σ -products and fundamental groups of subspaces of the plane*, Topology Appl. **84** (1998), 283-306.
- [17] K. Eda, *The fundamental groups of one-dimensional spaces and spatial homomorphisms*, Topology Appl. **123** (3) (2002), 479-505.
- [18] K. Eda and K. Kawamura, *The fundamental group of one-dimensional spaces*, Topology Appl., **87** (3) (1998), 163-172.
- [19] D.M. Evans, *Subgroups of small index in infinite general linear groups*, Bull. London Math. Soc. **18** (1986), 587-590.

- [20] S.M. Gersten, *A presentation for the special automorphism group of a free group*, J. Pure Appl. Algebra **33**, (3) (1984), 269-279.
- [21] C.K. Gupta, W. Holubovski, *Automorphisms of a free group of infinite rank*, Algebra and Analysis, **19** (2) (2007), 74-85.
- [22] G. Higman, *Unrestricted free products and varieties of topological groups*, J. London Math. Soc., (2) **27** (1952), 73-81.
- [23] W. Hodges, I. Hodkinson, D. Laskar, S. Shelah *The small index property for ω -stable ω -categorical structures and for the random graph*, J. London Math. Soc. (2) **48** (1993), 204-218.
- [24] J.L. Kelley, *General Topology*, Van Nostrand, 1955; reprinted by Springer as Vol. **27** in Graduate Texts in Mathematics.
- [25] R.C. Lyndon, P.E. Schupp, *Combinatorial group theory*, Springer, 1977.
- [26] O. Macedonska-Nosalska, *Note on automorphisms of a free abelian group*, Canad. Math. Bull., **23** (1980), 111-113.
- [27] G. Maxwell, *Infinite general linear groups over rings*, TAMS, **151** (1970), 371-375.
- [28] J. McCool, *A presentation for the automorphism group of a free group of finite rank*, J. London Math. Soc., **8** (1974), 259-266.
- [29] J. McCool, *Some finitely presented subgroups of the automorphism group of a free group*. J. Algebra, **35** (1975), 205-213.
- [30] A. Rosenberg, *The structure of the infinite general linear groups*, Ann. of Math., II. Ser. **68** (1958), 278-294.
- [31] A. J. Sieradski, *Omega-groups*, preprint. Available at <http://people.oregonstate.edu/~bogleyw/research/og.pdf>
- [32] V. Tolstykh, *On Bergmann's property for the automorphism groups of relatively free groups*, J. of the London Math. Soc. **73** (3) (2006), 669-680.