


Max-Planck-Institut für Mathematik Bonn

Singularity analysis and integrability of a Burgers-type
system by Foursov

by

Sergei Sakovich



Singularity analysis and integrability of a Burgers-type system by Foursov

Sergei Sakovich

Max-Planck-Institut für Mathematik
Vivatsgasse 7
53111 Bonn
Germany

Institute of Physics
National Academy of Sciences
220072 Minsk
Belarus

SINGULARITY ANALYSIS AND INTEGRABILITY OF A BURGERS-TYPE SYSTEM BY FOURSOV

SERGEI SAKOVICH

ABSTRACT. We apply the Painlevé test for integrability of partial differential equations to a system of two coupled Burgers-type equations found by Foursov, which was recently shown by Sergyeyev to possess infinitely many commuting local generalized symmetries without any recursion operator. The Painlevé analysis easily detects that this is a typical C-integrable system in the Calogero sense and rediscovers its linearizing transformation.

1. INTRODUCTION

The system of two coupled Burgers-type equations

$$\begin{aligned} w_t &= w_{xx} + 8ww_x + (2 - 4\alpha)zz_x, \\ z_t &= (1 - 2\alpha)z_{xx} - 4\alpha zw_x + (4 - 8\alpha)wz_x - (4 + 8\alpha)w^2z - (2 - 4\alpha)z^3, \end{aligned} \quad (1)$$

where α is a parameter, was discovered by Foursov [1] as a nonlinear system which possesses generalized symmetries of orders three through at least eight but apparently has no recursion operator for a generic value of α . Foursov [1] noted that two systems equivalent to the cases $\alpha = 0$ and $\alpha = 1$ of (1) had already appeared in [2] and [3], respectively, and found a recursion operator for the system (1) with $\alpha = 1/2$. Very recently, Sergyeyev [4] proved that the system (1) does possess an infinite commutative algebra of local generalized symmetries but the existence of a recursion operator—of a reasonably “standard” form—for a generic value of α is disallowed by the structure of symmetries. Sergyeyev [4] found that the algebra of generalized symmetries of (1) is generated by a nonlocal two-term recursion relation rather than a recursion operator.

In the present paper, we find it interesting to see what the Painlevé test for integrability, in its formulation for partial differential equations [5, 6, 7], can tell about the integrability of this unusual system (1) with $\alpha \neq 1/2$, which possesses infinitely many higher symmetries without any recursion operator. The Painlevé test easily detects that this is a typical C-integrable system, in the terminology of Calogero [8]. In Section 2, we show that the singularity analysis of the Burgers-type system (1) naturally suggests to introduce the new dependent variable $s(x, t)$,

$$s = z^2, \quad (2)$$

to improve the dominant behavior of solutions. The system (1) in the variables w and s passes the Painlevé test for integrability successfully: positions of resonances are integer in all branches, and there are no nontrivial compatibility conditions at the resonances. In Section 3, we show that the truncation of singular expansions straightforwardly produces the transformation

$$w = \frac{\phi_x}{4\phi}, \quad s = \frac{a^2}{(4 - 8\alpha)\phi} \quad (3)$$

to the new dependent variables $\phi(x, t)$ and $a(x, t)$ satisfying the triangular linear system

$$a_t = (1 - 2\alpha)a_{xx}, \quad \phi_t = \phi_{xx} + a^2. \quad (4)$$

This linearizing transformation was found in an inverse form in [9] and used in a form close to (2)–(3) in [4]. Section 4 contains concluding remarks.

2. SINGULARITY ANALYSIS

First of all, let us note that the cases $\alpha = 1/2$ and $\alpha \neq 1/2$ of the system (1) are essentially different, at least because the total order of the system's equations is different in these cases, and the general solution of this two-dimensional system contains different numbers of arbitrary functions of one variable in these cases, three and four, respectively. When $\alpha = 1/2$, the system (1) is the triangular system

$$w_t = w_{xx} + 8ww_x, \quad z_t = -2zw_x - 8w^2z, \quad (5)$$

where the first equation is the linearizable Burgers equation possessing the Painlevé property [5], whereas the second equation simply defines a function $z(x, t)$ by the relation $z = f(x) \exp \int (-2w_x - 8w^2) dt$, with $f(x)$ being arbitrary, for any solution $w(x, t)$ of the Burgers equation. Thus, integrability of this case is trivial.

In the generic case of the Burgers-type system (1) with $\alpha \neq 1/2$, we substitute into (1) the expansions

$$\begin{aligned} w &= w_0(t)\phi^\sigma + \dots + w_r(t)\phi^{\sigma+r} + \dots, \\ z &= z_0(t)\phi^\tau + \dots + z_r(t)\phi^{\tau+r} + \dots, \end{aligned} \quad (6)$$

where $\phi_x(x, t) = 1$, in order to determine the dominant behavior of solutions near a movable non-characteristic manifold $\phi(x, t) = 0$ and the corresponding positions of resonances. In this way, we obtain the following four branches, omitting the ones corresponding to the Taylor expansions governed by the Cauchy–Kovalevskaya theorem:

$$\sigma = \tau = -1, \quad w_0 = \frac{1}{2}, \quad z_0 = \pm \sqrt{\frac{1}{4\alpha - 2}}, \quad r = -2, -1, 1, 2; \quad (7)$$

$$\sigma = \tau = -1, \quad w_0 = 1, \quad z_0 = \pm \sqrt{\frac{3}{2\alpha - 1}}, \quad r = -4, -3, -1, 2; \quad (8)$$

$$\sigma = -1, \quad \tau = -\frac{1}{2}, \quad w_0 = \frac{1}{4}, \quad \forall z_0(t), \quad r = -1, 0, 1, 2; \quad (9)$$

$$\sigma = -1, \quad \tau = \frac{1}{2}, \quad w_0 = \frac{1}{4}, \quad \forall z_0(t), \quad r = -1, -1, 0, 2. \quad (10)$$

We see that the system (1) does not possess the Painlevé property because of the non-integer values of τ in the branches (9) and (10). Nevertheless, the positions of resonances are integer in all branches, and we can improve the dominant behavior of solutions by a simple power-type transformation of the dependent variable z , like we did for the Golubchik–Sokolov system in [10]. We introduce the new dependent variable s given by (2), and this brings the Burgers-type system (1) into the form

$$\begin{aligned} w_t &= w_{xx} + 8ww_x + (1 - 2\alpha)s_x, \\ ss_t &= (1 - 2\alpha)ss_{xx} - \frac{1}{2}(1 - 2\alpha)s_x^2 - 8\alpha s^2 w_x + (4 - 8\alpha)wss_x \\ &\quad - (8 + 16\alpha)w^2s^2 - (4 - 8\alpha)s^3. \end{aligned} \quad (11)$$

This form is hardly simpler than the original one, but the studied system (1) in this form (11) will pass the Painlevé test.

We substitute into (11) the expansions

$$\begin{aligned} w &= w_0(t)\phi^\sigma + \dots + w_r(t)\phi^{\sigma+r} + \dots, \\ s &= s_0(t)\phi^\rho + \dots + s_r(t)\phi^{\rho+r} + \dots, \end{aligned} \quad (12)$$

with $\phi_x(x, t) = 1$, and find the following four branches:

$$\sigma = -1, \quad \rho = -2, \quad w_0 = \frac{1}{2}, \quad s_0 = \frac{1}{4\alpha - 2}, \quad r = -2, -1, 1, 2; \quad (13)$$

$$\sigma = -1, \quad \rho = -2, \quad w_0 = 1, \quad s_0 = \frac{3}{2\alpha - 1}, \quad r = -4, -3, -1, 2; \quad (14)$$

$$\sigma = \rho = -1, \quad w_0 = \frac{1}{4}, \quad \forall s_0(t), \quad r = -1, 0, 1, 2; \quad (15)$$

$$\sigma = -1, \quad \rho = 1, \quad w_0 = \frac{1}{4}, \quad \forall s_0(t), \quad r = -1, -1, 0, 2. \quad (16)$$

Now the exponents of the dominant behavior of solutions, as well as the positions of resonances, are integer in all branches.

The next step of the Painlevé analysis is to derive from (11) and (12) the recursion relations for the coefficients w_n and s_n ($n = 0, 1, 2, \dots$) and then to check the compatibility conditions arising at the resonances. Omitting tedious computational details of this, we give here only the result. The compatibility conditions turn out to be satisfied identically at the resonances of all branches (13)–(16), hence there is no need to introduce logarithmic terms into the expansions (12) representing solutions of the system (11). The function $\psi(t)$ in $\phi = x + \psi(t)$ remains arbitrary in all branches. Also the following functions remain arbitrary: $s_1(t)$, and either $s_2(t)$ if $\alpha = 1$ or $w_2(t)$ if $\alpha \neq 1$, in the branch (13); either $s_2(t)$ if $\alpha = 3/2$ or $w_2(t)$ if $\alpha \neq 3/2$, in the branch (14); $s_0(t)$, $s_1(t)$ and $w_2(t)$ in the branch (15); and $s_0(t)$ and $w_2(t)$ in the branch (16). The generic branch is (15): the expansions (12) contain four arbitrary functions of one variable in this case, thus representing the general solution of the system (11).

Consequently, the Burgers-type system (1) in its equivalent form (11) has passed the Painlevé test for integrability.

3. TRUNCATION TECHNIQUE

There is a strong empirical evidence that any nonlinear differential equation which passed the Painlevé test must be integrable. The test itself, however, does not tell whether the equation is C-integrable (solvable by quadratures or exactly linearizable) or S-integrable (solvable by an inverse scattering transform technique). Often some additional information on integrability of the studied equation, such as its linearizing transformation, Lax pair, Bäcklund transformation, etc., can be obtained by truncation of the Laurent-type expansion representing the equation's general solution [5, 11, 12, 13, 14, 15, 16, 17].

Let us apply the truncation technique to the system (11). We make the truncation in the generic branch (15) which corresponds to the general solution. In what follows, the simplifying reduction $\phi = x + \psi(t)$, $w_n = w_n(t)$ and $s_n = s_n(t)$ ($n = 0, 1, \dots$) is not used. We substitute the truncated expansions

$$w = \frac{w_0(x, t)}{\phi(x, t)} + w_1(x, t), \quad s = \frac{s_0(x, t)}{\phi(x, t)} + s_1(x, t) \quad (17)$$

to the coupled equations (11), equate to zero the sums of terms with equal degrees of ϕ , and in this way obtain the definitions

$$w_0 = \frac{\phi_x}{4}, \quad s_0 = \frac{\phi_t - \phi_{xx} - 8w_1\phi_x}{4 - 8\alpha} \quad (18)$$

for the coefficients w_0 and s_0 , as well as a system of four nonlinear partial differential equations for three functions, $w_1(x, t)$, $s_1(x, t)$ and $\phi(x, t)$. Two of the four equations of that system are the same initial equations (11) with w and s replaced by w_1 and s_1 , respectively, which means that the obtained system and the relations (17) and (18) constitute a so-called Painlevé–Bäcklund transformation relating a

solution (w_1, s_1) of (11) with a solution (w, s) of (11). The other two equations of the obtained system are fourth-order polynomial partial differential equations—let us simply denote them as E_1 and E_2 because it is easy to obtain them by computer algebra tools but not so easy to put them onto a printed page—which involve the functions w_1 , s_1 and ϕ and contain, respectively, 55 and 110 terms.

Fortunately, there is no need to study the obtained complicated system of four nonlinear equations for compatibility in its full form. Instead, let us see what will happen if we take $w_1 = 0$ and $s_1 = 0$, which means that we apply the obtained Painlevé–Bäcklund transformation to the trivial zero solution of the system (11). The reason to do so consists in the following empirically observed difference between C-integrable equations and S-integrable equations, which, as far as we know, has never been formulated explicitly in the literature. A Painlevé–Bäcklund transformation of a C-integrable equation, being applied to a single trivial solution of the equation, produces the whole general solution of the equation at once. Examples of this are the Burgers equation [5] and the Liouville equation in its polynomial form $uu_{xy} = u_x u_y + u^3$ [12]. On the contrary, numerous examples in the literature show that a Painlevé–Bäcklund transformation of a S-integrable equation, being applied to a single trivial solution of the equation, produces only a class of special solutions of the equation, usually a rational solution or a one-soliton solution with some arbitrary parameters.

Taking $w_1 = 0$ and $s_1 = 0$, we find that the equation we denoted as E_1 is satisfied identically, whereas the equation we denoted as E_2 is reduced to

$$\begin{aligned} (\phi_t - \phi_{xx})(\phi_t - \phi_{xx})_t + \frac{1}{2}(1 - 2\alpha)(\phi_t - \phi_{xx})_x^2 \\ - (1 - 2\alpha)(\phi_t - \phi_{xx})(\phi_t - \phi_{xx})_{xx} = 0. \end{aligned} \quad (19)$$

The general solution of this fourth-order equation contains four arbitrary functions of one variable, which is exactly the degree of arbitrariness of the general solution of the system (11). For this reason, we conclude that the system (11) must be C-integrable. Now it only remains to notice that, if we introduce the new dependent variable $a(x, t)$ such that

$$\phi_t - \phi_{xx} = a^2, \quad (20)$$

the equation (19) becomes linear:

$$a_t = (1 - 2\alpha)a_{xx}. \quad (21)$$

Finally, combining the relations (17), (18), (20), (21) and $w_1 = s_1 = 0$, we obtain the exact linearization (3) and (4) for the system (11).

4. CONCLUSION

In the present paper, we used the Painlevé test for integrability of partial differential equations to study the integrability of a system of two coupled Burgers-type equations discovered by Foursov, which possesses an unusual algebra of generalized symmetries as was shown by Sergyeyev. The Painlevé analysis easily detected that the studied Burgers-type system is a typical C-integrable system in the Calogero sense and rediscovered its linearizing transformation. As a by-product, we obtained a new example confirming the empirically observed difference between the Painlevé–Bäcklund transformations of C-integrable equations and S-integrable equations. In our opinion, the Painlevé test deserves to be used more widely to search for new integrable nonlinear equations, because with its help one can discover new equations possessing such new properties which look unusual from the point of view of other integrability tests.

ACKNOWLEDGEMENTS

This work was partially supported by the BRFFR grant Φ 10-117. The author also thanks the Max Planck Institute for Mathematics for hospitality and support.

REFERENCES

- [1] M.V. Foursov. On integrable coupled Burgers-type equations. *Phys. Lett. A* 272 (2000) 57–64.
- [2] S.I. Svinolupov. On the analogues of the Burgers equation. *Phys. Lett. A* 135 (1989) 32–36.
- [3] P.J. Olver, V.V. Sokolov. Integrable evolution equations on associative algebras. *Commun. Math. Phys.* 193 (1998) 245–268.
- [4] A. Sergyeyev. Infinitely many local higher symmetries without recursion operator or master symmetry: Integrability of the Foursov–Burgers system revisited. *Acta Appl. Math.* 109 (2010) 273–281; arXiv:0804.2020.
- [5] J. Weiss, M. Tabor, G. Carnevale. The Painlevé property for partial differential equations. *J. Math. Phys.* 24 (1983) 522–526.
- [6] M. Tabor. *Chaos and Integrability in Nonlinear Dynamics: An Introduction*. Wiley, New York, 1989.
- [7] A. Ramani, B. Grammaticos, T. Bountis. The Painlevé property and singularity analysis of integrable and non-integrable systems. *Phys. Rep.* 180 (1989) 159–245.
- [8] F. Calogero. Why are certain nonlinear PDEs both widely applicable and integrable? In: *What Is Integrability?* (Ed. V.E. Zakharov) 1–62. Springer, New York, 1991.
- [9] T. Tsuchida, T. Wolf. Classification of polynomial integrable systems of mixed scalar and vector evolution equations: I. *J. Phys. A: Math. Gen.* 38 (2005) 7691–7733; arXiv:nlin/0412003.
- [10] S.Yu. Sakovich. A system of four ODEs: The singularity analysis. *J. Nonlinear Math. Phys.* 8 (2001) 217–219; arXiv:nlin/0003039.
- [11] J. Weiss. The Painlevé property for partial differential equations. II: Bäcklund transformations, Lax pairs, and the Schwarzian derivative. *J. Math. Phys.* 24 (1983) 1405–1413.
- [12] W.-H. Steeb, M. Kloke, B.M. Spieker. Liouville equation, Painlevé property and Bäcklund transformation. *Z. Naturforsch. A* 38 (1983) 1054–1055.
- [13] M. Musette, R. Conte. Algorithmic method for deriving Lax pairs from the invariant Painlevé analysis of nonlinear partial differential equations. *J. Math. Phys.* 32 (1991) 1450–1457.
- [14] A. Karasu-Kalkanlı, S.Yu. Sakovich. Bäcklund transformation and special solutions for the Drinfeld–Sokolov–Satsuma–Hirota system of coupled equations. *J. Phys. A: Math. Gen.* 34 (2001) 7355–7358; arXiv:nlin/0102001.
- [15] A. Karasu-Kalkanlı, A. Karasu, A. Sakovich, S. Sakovich, R. Turhan. A new integrable generalization of the Korteweg–de Vries equation. *J. Math. Phys.* 49 (2008) 073516; arXiv:0708.3247.
- [16] R. Conte, M. Musette. *The Painlevé Handbook*. Springer, Dordrecht, 2008.
- [17] A.N.W. Hone. Painlevé tests, singularity structure and integrability. In: *Integrability* (Ed. A.V. Mikhailov) 245–277. Springer, Berlin, 2009. ArXiv:nlin/0502017.

INSTITUTE OF PHYSICS, NATIONAL ACADEMY OF SCIENCES, 220072 MINSK, BELARUS
 MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY
E-mail address: saks@tut.by