

A Note on the Geography of Symplectic Manifolds

András Stipsicz

Department of Mathematics
University of California
Irvine, CA 92717
USA

Max-Planck-Institut
für Mathematik
Gottfried-Claren-Str. 26
53225 Bonn
GERMANY



A Note on the Geography of Symplectic Manifolds

András Stipsicz

July 26, 1995

1 Introduction

Based on recent developments in gauge theory – the introduction of Seiberg-Witten invariants and results of Taubes – our understanding of the differential topology of symplectic manifolds improved by a margin in the past year. In this note we would like to discuss some existence problems of minimal simply connected symplectic manifolds; in particular we would like to compare the "geography" of symplectic manifolds and complex surfaces.

Let us first briefly recall the geography of simply connected compact complex surfaces. Since X is simply connected, b^+ is odd (by the Noether formula $12 \mid c_1^2(X) + c_2(X)$), and the holomorphic Euler characteristic $\chi(X)$ is $\frac{1+b^+}{2}$. Also note that $c_1^2(X) = 3\sigma(X) + 2e(X)$, here $\sigma(X)$ denotes the signature, $e(X)$ the Euler characteristic of X .

To a complex surface X let us associate this two integers

$$X \rightarrow (\chi(X), c_1^2(X)).$$

For example $(\chi(\mathbb{C}\mathbb{P}^2), c_1^2(\mathbb{C}\mathbb{P}^2)) = (1, 9)$; $(\chi(S^2 \times S^2), c_1^2(S^2 \times S^2)) = (1, 8)$ and $(\chi(E(n)), c_1^2(E(n))) = (n, 0)$ ($E(n)$ is the regular elliptic surface with a section and $e(E(n)) = 12n$, in particular $E(2)$ is the $K3$ surface).

Note also that if X' is the blow up of X , then $(\chi(X'), c_1^2(X')) = (\chi(X), c_1^2(X) - 1)$.

By the classification result of Kodaira we know that a simply connected compact complex surface is either rational, elliptic or a surface of general type. If X is rational (meaning birationally equivalent to $\mathbb{C}\mathbb{P}^2$), then $b^+ = \chi(X) = 1$, and the simply connected minimal rationals are diffeomorphic to $\mathbb{C}\mathbb{P}^2$, $S^2 \times S^2$ or $\mathbb{C}\mathbb{P}^2 \# \bar{\mathbb{C}}\mathbb{P}^2$ (the Hirzebruch-surfaces).

If X is minimal elliptic (so X admits a holomorphic map $\pi : X \rightarrow \mathbb{C}\mathbb{P}^1$ with a smooth elliptic curve as a generic fiber), then $(\chi(X), c_1^2(X)) = (n, 0)$ for some $n \in \mathbb{N}$. For surfaces

of general type we know that $c_1^2(X) > 0$, and the two famous inequalities (the Noether inequality and the Bogomolov-Miyaoka-Yau inequality) give constraints for $c_1^2(X)$ in terms of $\chi(X)$:

$$2\chi(X) - 6 \leq c_1^2(X) \leq 9\chi(X).$$

Most of the points of this region (like $2\chi(X) - 6 \leq c_1^2(X) \leq 4\chi(X)$) is known to correspond to a minimal surface of general type (see [P] or [BPV] for further details).

The same geography question makes sense for symplectic manifolds as well – namely which points $(a, b) \in \mathbb{Z}^2$ can be realized as $(\chi(X) = \frac{1+b^+}{2}, c_1^2(X) = 3\sigma(X) + 2e(X))$ of a minimal simply connected symplectic manifold X . Note that $12 \mid c_1^2(X) + c_2(X)$ holds for an almost complex manifold, and so in particular for a symplectic manifold as well, so b_X^+ is odd for a symplectic manifold X . Also blow up and blow down of a symplectic (-1) -sphere makes sense in the symplectic category, so minimality can be defined for symplectic manifolds as well.

A simply connected complex surface is Kähler, hence symplectic; so the regions populated by complex surfaces are already covered by symplectic manifolds as well. In the followig we will show, that a big part of the region under the Noether-line can be populated by minimal symplectic manifolds, more precisely if $D = \{(a, b) \in \mathbb{Z}^2 \mid 0 < b < 2a - 6\}$, then

Theorem 1.1 *If $(a, b) \in D$ and b is even, then there is a minimal symplectic manifold X such that $(\chi(X), c_1^2(X)) = (a, b)$.*

Remark 1.2 • *Note that – by recent result of Taubes – $c_1^2(X) \geq 0$ for a minimal symplectic manifold.*

- *The region D above was already populated by examples of Gompf ([G]) which were symplectic, but it is not clear yet whether those examples are minimal – although they very likely are. Fintushel and Stern gave irreducible examples covering the region D as well; since the rational blow down process (defined in [FS2]) is not proved to keep the symplectic structure, it is not clear whether those examples carry symplectic structure – very likely they do. Our theorem represents only points $(a, b) \in D$ with even b as $(\chi(X), c_1^2(X))$ of a minimal symplectic manifold X , although most probably the same argument works for every point in D .*
- *Note also that the examples given by Theorem 1.1 do not carry complex structure.*

Acknowledgement: We would like to thank the Max-Planck-Institute for their hospitality and Bob Gompf for the many helpful discussions.

2 Donaldson series

Let us briefly recall the rudiments of Donaldson series (see also [KM], [DK]).

For a simply connected manifold X with $b^+ \geq 3$ and odd an analytic function

$$\mathbb{D}_{X,c} : H_2(X; \mathbb{R}) \rightarrow \mathbb{R}$$

can be defined. The definition of $\mathbb{D}_{X,c}$ uses the ASD equation for connections on auxiliary principal $\mathrm{SO}(3)$ -bundles P over X with $w_2(P) \equiv c \pmod{2}$.

Remark 2.1 *To define $\mathbb{D}_{X,c}$ one needs an additional property of X – it has to be of simple type (see [KM]). Also the definition of $\mathbb{D}_{X,c}$ needs a choice of $c \in H_2(X; \mathbb{Z})$ and a homology orientation of X (see [D]).*

The beautiful structure theorem of Kronheimer-Mrowka and Fintushel-Stern ([KM], [FS1]) states that

$$\mathbb{D}_{X,c} = \exp\left(\frac{Q}{2}\right) \cdot \sum_{i=1}^s a_i e^{K_i}$$

where $a_i \in \mathbb{Q} \setminus \{0\}$ and $K_i \in H^2(X; \mathbb{Z})$ ($i = 1, \dots, s$). $\{K_i\}_{i=1}^s$ is the set of (KM)-basic classes of the manifold X , these classes satisfy the following properties:

- $K_i \equiv w_2(X) \pmod{2}$;
- if K is a basic class, then $-K$ is a basic class as well;
- if $\Sigma \subset X$ is a smoothly embedded surface with $[\Sigma]^2 \geq 0$ and genus $g(\Sigma)$, then for any basic class K

$$2g(\Sigma) - 2 \geq [\Sigma]^2 + |K([\Sigma])|.$$

Theorem 2.2 *(Blow up formula)*

If $\{K_i\}_{i=1}^s$ is the set of basic classes for X , then $\{K_i \pm E\}_{i=1}^s$ is the set of basic classes for $X \# \mathbb{C}\mathbb{P}^2$ (E is the Poincare dual of the exceptional fiber).

More generally if $X = X_1 \# X_2$ where $b^+(X_2) = 0$ (so the intersection form of X_2 is $n\langle -1 \rangle$ spanned by $\{e_1, \dots, e_n\}$), then the set of basic classes of X is $\{K_i \pm E_1 \pm \dots \pm E_n\}$ where $\{K_i\}$ is the set of basic classes of X_1 (and E_i is the Poincare dual of e_i).

By the connected sum theorem of Donaldson we know, that if X has non-zero series, then X cannot admit a decomposition $X = X_1 \# X_2$ with $b^+(X_i) > 0$ ($i = 1, 2$). A decomposition with $b^+(X_2) = 0$ however is possible, so irreducibility doesn't follow directly from the non-vanishing of the invariants.

Proposition 2.3 *Assume that the set of basic classes $\{K_i\}_{i=1}^s$ of the manifold X satisfies*

$$(K_i - K_j)^2 \neq -4 \text{ for all } 1 \leq i, j \leq s.$$

In this case X is irreducible.

Proof: The existence of basic classes insure, that $\mathbb{D}_X \neq 0$, so if X is reducible, then $X = X_1 \# X_2$ with $b^+(X_2) = 0$ is the only possibility. By the previous remark however in this case there are basic classes K_i, K_j such that $K_i - K_j = 2E_1$, so $(K_i - K_j)^2 = -4$ contradicting our assumption.

Assume that the manifold X has only 2 basic classes $\pm K \in H^2(X; \mathbb{Z})$ and $K^2 > 0$. Assume also that X contains a torus f with square 0 lying in a cusp neighborhood. In this case one can take the fiber sum of X with the regular elliptic surface $E(n)$ along f .

Proposition 2.4 *$X \#_f E(n)$ is an irreducible manifold.*

Proof: Applying the computations presented in [S] (Proposition 3.3), the set of basic classes of $X \#_f E(n)$ is

$$\{\pm K + k \cdot F \mid k \equiv n \pmod{2}, |k| \leq n\}$$

(F is the Poincare dual of the homology class represented by f). The difference of two basic classes is either $k_1 \cdot F$ or $\pm(2K + k_2 \cdot F)$; the squares of these elements are at least 0 so by Proposition 2.3 $X \#_f E(n)$ is irreducible.

3 Irreducible symplectic manifolds

Let us take the set Ξ of simply connected symplectic manifolds X having the following properties:

1. X has exactly two basic classes $(\pm K)$ and $K^2 > 0$;
2. X contains a torus f with $f^2 = 0$ such that f is lying in a cusp neighborhood and f is a symplectic or lagrangian submanifold of X .

By the construction of Gompf $X \#_f E(n)$ is symplectic; by Proposition 2.4 it is irreducible as well. Note that $(\chi(X \#_f E(n)), c_1^2(X \#_f E(n))) = (\chi(X) + n, c_1^2(X))$. So to prove Theorem 1.1 we only have to show, that for every even $b > 0$ Ξ contains an element X such that

$(\chi(X), c_1^2(X)) = (a, b)$ with $b \geq 2a - 6$. As Fintushel and Stern observed ([FS]), complete intersections, Moishezon surfaces and Salvetti surfaces are elements of Ξ (note that in these cases the torus f is a lagrangian submanifold). Also by analyzing the effect of rational blowdown, Fintushel and Stern realized ([FS2]) that surfaces on the Noether-line $c_1^2 = 2\chi - 6$ (the Horikawa surfaces) can be constructed by blowing down rationally elliptic surfaces $E(n)$. Since $E(n)$ contains lagrangian tori disjoint from the configurations one blows down to get the Horikawa surfaces, we have

Theorem 3.1 *The Horikawa surfaces constructed by rationally blowing down the elliptic surfaces $E(n)$ are in Ξ .*

In this way we have an element of Ξ with $c_1^2 = 2\chi - 6$ for every even c_1^2 , and this proves Theorem 1.1.

Remark 3.2 • *By performing a logarithmic transformation of multiplicity 2 on f — which is known to be a symplectic operation — we can turn a spin manifold into a non-spin one; the resulting manifold remains irreducible.*

- *Most probably the surfaces on the "next Horikawa line" $c_1^2 = 2\chi - 5$ contain also the required symplectic or lagrangian torus in the cusp neighborhood, so we can relax the assumption on the parity of b in Theorem 1.1. This issue will be discussed elsewhere.*

References

- [BPV] W. Barth, C. Peters, A. Van de Ven *"Compact Complex Surfaces"* *Ergebnisse der Mathematik, Springer-Verlag, Berlin 1984*
- [D] S.K. Donaldson *Polynomial Invariants for Smooth 4-Manifolds*, *Topology* 29 (1990) 257-315
- [DK] S.K. Donaldson, P. Kronheimer *"The Geometry of Four-Manifolds"* *Oxford Mathematical Monography, Oxford University Press, Oxford 1990*
- [FS] R. Fintushel, R. Stern *Surgery in Cusp Neighborhoods and the Geography of Irreducible 4-Manifolds*, to appear
- [FS1] R. Fintushel, R. Stern *Donaldson invariants of 4-manifolds with simple type*, preprint
- [FS2] R. Fintushel, R. Stern *Rational blowdowns of smooth 4-manifolds*, preprint
- [G] R. Gompf *A New Construction of Symplectic Manifolds*, preprint

- [KM] P. Kronheimer, T. Mrowka *Embedded surfaces and the structure of Donaldson's polynomial invariants, to appear in Journal of Differential Geometry*
- [P] U. Persson *Chern invariants of surfaces of general type, Compositio Mathematica, Vol 43 (1981) 3-58*
- [S] A. Stipsicz *Donaldson series and (-1) -tori, to appear*

Department of Mathematics,
University of California
Irvine, CA 92717

e-mail: astipsic@math.uci.edu