

**Equivariant Reidemeister torsion on
symmetric spaces**

Kai Köhler

Max-Planck-Institut für Mathematik
Gottfried-Claren-Straße 26
53225 Bonn

Germany

**THE HOMEOMORPHISM TYPES
OF CONTRACTIBLE PLANAR
POLYHEDRA**

Hans-Joachim Baues*
Antonio Quintero**

**

Departamento de Algebra, Computacion,
Geometria y Topologia
Universidad de Sevilla
Apdo 1160
41080 - Sevilla

Spain

*

Max-Planck-Institut für Mathematik
Gottfried-Claren-Straße 26
53225 Bonn

Germany

Equivariant Reidemeister torsion on symmetric spaces

KAI KÖHLER

Max-Planck-Institut für Mathematik

Gottfried-Claren-Str. 26

D-53225 Bonn

e-mail: Koehler@MPIM-Bonn.MPG.De

April 21, 1994

Abstract

We calculate explicitly the equivariant Ray-Singer torsion for all symmetric spaces G/K of compact type with respect to the action of G . We show that it equals zero except for the odd-dimensional Grassmannians and the space $\mathbf{SU}(3)/\mathbf{SO}(3)$. As a corollary, we classify up to diffeomorphism all isometries of these spaces which are homotopic to the identity; in particular, we classify the diffeomorphism types of their quotients by finite group actions.

Contents

1	Introduction	3
2	Equivariant Ray-Singer metrics	4
3	Homogeneous and symmetric spaces	5
4	The zeta function for symmetric spaces	8
5	The torsion for symmetric spaces	10

1 Introduction

In 1935, Franz and Reidemeister established the following classification of lens spaces:

Theorem 1 ([5]) *Let Γ_1, Γ_2 be cyclic groups acting isometrically and freely on the spheres S^{2n-1} , $n > 1$. Then the quotients $S^{2n-1}/\Gamma_1, S^{2n-1}/\Gamma_2$ are diffeomorphic iff they are isometric, i.e. if Γ_1 and Γ_2 are conjugated in $O(2n)$.*

To prove this theorem they invented a real-valued combinatorial invariant of CW-complexes which is fine enough to distinguish the lens spaces, the Reidemeister torsion. Their result was generalized by de Rham in 1964:

Theorem 2 ([4]) *Two isometries of S^n are diffeomorphic iff they are isometric.*

Here, two transformations g_1, g_2 of a manifold M are called diffeomorphic (resp. isometric) iff there exists a diffeomorphism (resp. an isometry) ϕ of M with $\phi g_1 = g_2 \phi$. The first purpose of this article is to prove the following result:

Theorem 3 *Two isometries homotopic to the identity of an odd-dimensional Grassmannian $G_{2m, 2p-1}(\mathbf{R}) = \mathbf{SO}(2m)/\mathbf{SO}(2p-1) \times \mathbf{SO}(2m-2p+1)$ or of $\mathbf{SU}(3)/\mathbf{SO}(3)$ are diffeomorphic iff they are isometric.*

As $\mathbf{SU}(3)/\mathbf{SO}(3)$ and the Grassmannians except the circle are simply connected, this classifies in particular the quotients of these symmetric spaces by finite group actions. The proof is given by calculating explicitly the equivariant Ray-Singer torsion for all compact symmetric spaces. The Ray-Singer torsion is defined as the derivative at zero of a certain zeta function associated to the spectrum of the Laplace operator on differential forms on a compact Riemannian manifold [12]. It has been determined by Ray for the lens spaces in an extensive calculation by determining first the eigenvalues and eigenspaces of the Laplacian on spheres. He found that the Reidemeister and Ray-Singer torsions are equal for these spaces. Using this result, Cheeger and Müller proved independently in 1978 the equality of the Reidemeister torsion and the Ray-Singer torsion. The second aim of our paper is to give a new, shorter proof of Ray's result.

The equivariant Ray-Singer torsion associated to an isometry g acting on M has been investigated by Lott and Rothenberg. They compared it with an equivariant Reidemeister torsion for finite group actions. Using Ray's calculation, they found that the equivariant torsion for spheres is mainly given by sums of the digamma function; this enabled them to give a new proof of theorem 2 for orientation-preserving actions on odd-dimensional spheres. We shall apply their method to deduce theorem 3 from our result for the torsion.

Also, we shall show that the equivariant torsion equals zero for all symmetric spaces G/K with respect to the action of any $g \in G$, except for products of $G_{2m,2p-1}(\mathbf{R})$ or $\mathbf{SU}(3)/\mathbf{SO}(3)$ with some G'/K' so that G' and K' have the same rank. A similar result has been shown by Moscovici and Stanton for locally symmetric spaces of the compact type [10].

Our method to obtain the value of the torsion is similar to the one used in a previous paper on holomorphic Ray-Singer torsion on Hermitian symmetric spaces [8]. For a symmetric space G/K an eigenvalue of the Laplacian is determined by its eigenspace as a G -representation. This reduces the problem of determining the zeta function to a problem in finite-dimensional representation theory. Nevertheless, there are big differences between the real and the holomorphic situation: In the complex case, the equivariant torsion is always non-trivial and depends in a rather subtle way on the fixed-point set of the isometry $g \in G$, in sharp contrast to our result theorem 10.

2 Equivariant Ray-Singer metrics

Let F be a complex flat hermitian vector bundle over a compact oriented Riemannian manifold M . Let

$$d : \Gamma(\Lambda^q T^* M \otimes F) \rightarrow \Gamma(\Lambda^{q+1} T^* M \otimes F)$$

denote the de Rham operator with coefficients in F and let d^* denote its formal adjoint with respect to the L^2 -metric. Consider the Hodge-Laplacian $\Delta_q := (d + d^*)^2$ acting on q -forms with coefficients in F . We denote the eigenspace of Δ_q corresponding to an eigenvalue $\lambda \in \text{Spec } \Delta_q$ by $\text{Eig}_\lambda(\Delta_q)$. Let g be an isometry of M preserving the hermitian bundle F . Consider the

zeta function

$$Z_g(s) := \sum_{q>0} (-1)^q q \sum_{\substack{\lambda \in \text{Spec} \Delta_q \\ \lambda \neq 0}} \lambda^{-s} \text{Tr} g|_{\text{Eig}_\lambda(\Delta_q)^*}$$

for $\Re s > \dim M/2$. Classically, this zeta function has a meromorphic continuation to the complex plane which is holomorphic at zero.

Definition 1 *The equivariant analytic torsion is defined as*

$$\tau_g(M, F) := e^{-\frac{1}{2}Z'_g(0)}.$$

This object has been defined by Ray [11]. We shall denote the torsion with coefficients in the trivial line bundle by $\tau_g(M)$. Ray showed the following property of τ_g : Consider a fixed point free action of a finite group Γ on M . Let $\rho : \Gamma \rightarrow \mathbf{U}(1)$ be an unitary representation, thus defining a flat hermitian line bundle F on M/Γ . Then the usual non-equivariant Ray-Singer torsion with coefficients in F is given by

$$\log \tau(M/\Gamma, F) = \frac{1}{\#\Gamma} \sum_{g \in \Gamma} \bar{\rho}(g) \log \tau_g(M).$$

The equivariant torsion has been investigated by Lott and Rothenberg [9] for flat metrics on F . They showed that it equals zero on even dimensional manifolds or for orientation reversing actions on odd dimensional manifolds. They proved the result

Theorem 4 ([9]) *Assume that g is homotopic to the trivial action. Choose a sequence $(f_\nu) \in \prod_{\nu \in \mathbf{Z}} \mathbf{R}$, $\sum f_\nu = 0$. Then the torsion of powers of g weighed with (f_ν)*

$$\sum f_\nu \tau_{g^\nu}(M)$$

is a smooth invariant.

This result has been generalized strongly by Bismut and Zhang [2].

3 Homogeneous and symmetric spaces

Let G be a connected compact Lie group and let K be a connected compact subgroup. Let $T_G \supset T_K$ denote the maximal tori of G and K . We denote

the Lie algebras of G , K , T_G and T_K by \mathfrak{g} , \mathfrak{k} , \mathfrak{t}_G and \mathfrak{t}_K , respectively. We fix compatible orderings on \mathfrak{t}_G^* and \mathfrak{t}_K^* . The action of K on the homogeneous space G/K induces a representation $\text{Ad}_{G/K}$ on the tangent space of G/K at the class of $[1] \in G$, i.e. on $\mathfrak{g}/\mathfrak{k}$. Let Ψ denote the set of weights of this representation, the isotropy representation, and let Δ_G and Δ_K be the sets of roots of G resp. K . Then the weights of the adjoint representation of G on \mathfrak{g} are given by Δ_G and the weight $\{0\}$ with multiplicity the rank of G . The weights of the action of K on \mathfrak{g} are given by Δ_K , Ψ and $\text{rk } K$ -times the $\{0\}$, thus

$$(\Delta_G \cup \{0\} \cdot \text{rk } G)|_{T_K} = \Delta_K \cup \Psi \cup \{0\} \cdot \text{rk } K \quad (1)$$

(counted with multiplicity). In particular, the dimension $\#\Psi$ of G/K is odd-dimensional iff $\dim G - \dim K$ is so. The space of forms $\Gamma(\Lambda^q T^*G/K)$ is an infinite dimensional G -representation which contains the space of its irreducible subrepresentations (V_π, π) as a L^2 -dense subspace. Thus,

$$\Gamma(\Lambda^q T^*G/K) \stackrel{\text{dense}}{\supset} \bigoplus_{\pi} \text{Hom}_G(V_\pi, \Gamma(\Lambda^q T^*G/K)) \otimes V_\pi. \quad (2)$$

In this imbedding, the tensor product $\text{Hom}_G(V_\pi, \Gamma(\Lambda^q T^*G/K)) \otimes V_\pi$ is the direct sum of $\dim \text{Hom}_G(V_\pi, \Gamma(\Lambda^q T^*G/K))$ copies of the representations (V_π, π) . By a Frobenius law due to Bott [3], there is a canonical isomorphism

$$\text{Hom}_G(V_\pi, \Gamma(\Lambda^q T^*G/K)) \cong \text{Hom}_K(V_\pi, \Lambda^q \text{Ad}_{G/K}) \quad (3)$$

(Note that $(\mathfrak{g}/\mathfrak{k} \otimes \mathbf{C})^* \cong \mathfrak{g}/\mathfrak{k} \otimes \mathbf{C}$ via the metric). In particular, the representations (V_π, π) which occur are finite dimensional.

Let (X_1, \dots, X_N) be an orthonormal basis of \mathfrak{g} with respect to the negative Killing form. The Casimir operator of \mathfrak{g} is defined as the following element of the universal enveloping algebra of \mathfrak{g}

$$\text{Cas} := - \sum X_j \cdot X_j. \quad (4)$$

Ikeda and Taniguchi proved the following beautiful result [6]:

Theorem 5 (Ikeda, Taniguchi) *Assume that G/K is a symmetric space equipped with the metric induced by the Killing form. Then the Laplacian Δ acts on the V_π 's as $-\text{Cas}$ with respect to the imbedding (2).*

The Casimir is known to act by multiplication with a constant on irreducible representations. Thus, the eigenspaces of the Laplacian correspond to the irreducible representations π with multiplicity $\dim \text{Hom}_K(V_\pi, \Lambda^q \text{Ad}_{G/K})$ and its eigenvalue there depends only on π .

Let $\rho_G := \frac{1}{2} \sum_{\alpha \in \Delta^+} \alpha$ be half the sum of the positive roots of G and let W_G be the Weyl group of G . Let $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$ denote the metric and the norm on \mathfrak{t}_G^* induced by the Killing form. We denote the sign of an element $w \in W_G$ by $(-1)^w$. As usual, we define

$$(\alpha, \rho_G) := \frac{2\langle \alpha, \rho_G \rangle}{\|\alpha\|^2} \quad (5)$$

for any weight α . For an irreducible representation π we denote by b_π the sum of its highest weight and ρ_G . Then, classically, the action of the Casimir is given by

$$\pi(\text{Cas}) = \|\rho_G\|^2 - \|b_\pi\|^2.$$

To abbreviate we set

$$\text{Alt}_G\{b\} := \sum_{w \in W_G} (-1)^w e^{2\pi i w b}.$$

Then the Weyl character formula for the character χ_{b_π} of the representation evaluated at $t \in T$ may be written as

$$\chi_{b_\pi}(t) = \frac{\text{Alt}_G\{b_\pi\}(t)}{\text{Alt}_G\{\rho_G\}(t)}.$$

This formula provides the definition of the so-called virtual (or formal) character χ_b for any b equal to ρ_G +some weight. This extends to all of G by setting χ_b to be invariant under the adjoint action. The corresponding virtual representation shall be denoted by V_b . Occasionally we shall use the notation $\chi_b^G, \chi_b^K, V_b^G, V_b^K$ to distinguish G - and K -representations. From now on we shall consider the irreducible symmetric space G/K as being equipped with any G -invariant metric $\langle \cdot, \cdot \rangle_\circ$. All these metrics are proportional to the metric induced by the Killing form [1, Th. 7.44]. We shall denote the dual metric and norm on \mathfrak{t}_G^* by $\langle \cdot, \cdot \rangle_\circ, \| \cdot \|_\circ$, too.

4 The zeta function for symmetric spaces

By theorem 5, the equivariant zeta function defining the torsion is given by

$$Z(s) = \sum_{q=1}^n (-1)^q q \sum_{\pi} \frac{\dim \operatorname{Hom}_K(V_{\pi}, \Lambda^q \operatorname{Ad}_{G/K})}{(\|b_{\pi}\|_{\circ}^2 - \|\rho_G\|_{\circ}^2)^s} \chi_{b_{\pi}}. \quad (6)$$

In the case $K = \{1\}$, we observe that

$$\begin{aligned} Z(s) &= \left(\sum_{q=1}^n (-1)^q q \dim \Lambda^q \mathfrak{g} \right) \sum_{\pi} \frac{\dim V_{\pi} \cdot \chi_{b_{\pi}}}{(\|b_{\pi}\|_{\circ}^2 - \|\rho_G\|_{\circ}^2)^s} \\ &= n(1-1)^{n-1} \sum_{\pi} \frac{\dim V_{\pi} \cdot \chi_{b_{\pi}}}{(\|b_{\pi}\|_{\circ}^2 - \|\rho_G\|_{\circ}^2)^s}. \end{aligned}$$

Thus the torsion $\tau_g(G)$ equals zero for all compact Lie groups except the circle. Our key result is the following

Lemma 6 *Let G/K be a n -dimensional homogeneous space.*

- *If $\operatorname{rk} G > \operatorname{rk} K + 1$ then the virtual representation*

$$\sum_{q=1}^n (-1)^q q \Lambda^q \operatorname{Ad}_{G/K}$$

is trivial.

- *Assume that $\operatorname{rk} G = \operatorname{rk} K + 1$ and let L denote the line of those weights of G which are zero on \mathfrak{t}_K . If (V_{π}, π) is an irreducible G -representation then the sum*

$$\sum_{q=1}^n (-1)^q q \dim \operatorname{Hom}_K(V_{\pi}, \Lambda^q \operatorname{Ad}_{G/K}) \cdot \chi_{b_{\pi}}$$

equals the sum of $\chi_{\rho_G + w\alpha}$ over those $[w] \in W_G/W_K$ and $\alpha \in L$ such that b_{π} lies in the W_G -orbit of $\rho_G + w\alpha$.

By this lemma and theorem 5 we get the following expression for the equivariant zeta function Z :

Lemma 7 For any odd-dimensional symmetric space G/K , the zeta function $Z(s)$ is given by

$$Z(s) = \sum_{\substack{\{w\} \in W_G/W_K \\ \alpha \in L \\ \langle w\alpha, w\alpha + 2\rho_G \rangle > 0}} \frac{\chi_{\rho_G + w\alpha}}{\langle w\alpha, w\alpha + 2\rho_G \rangle^s} = \frac{1}{\#W_K} \sum_{\substack{w \in W_G \\ \alpha \in L \\ \langle w\alpha, w\alpha + 2\rho_G \rangle > 0}} \frac{\chi_{\rho_G + w\alpha}}{\langle w\alpha, w\alpha + 2\rho_G \rangle^s}$$

if $\text{rk } G = \text{rk } K + 1$ and zero otherwise.

In particular, one observes the following consequence:

Corollary 8 Let $G/K_1, G/K_2$ be two symmetric spaces with G -conjugate tori T_{K_1} and T_{K_2} . Then the associated zeta functions Z_1 and Z_2 are proportional by the factor $\#W_{K_1}/\#W_{K_2}$.

This shows already the desired classification of $\mathbf{SO}(2m)$ -actions on odd-dimensional Graßmannians $G_{2m,2p-1}(\mathbf{R})$.

Theorem 9 Two homotopically trivial isometries of the Graßmannian $G_{2m,2p-1}$ are conjugate by a diffeomorphism iff they are conjugate by an isometry in $\mathbf{O}(2m)$. In particular, two quotients of $G_{2m,2p-1}$ by finite groups $\Gamma_1, \Gamma_2 \subset \mathbf{SO}(2m)$ are diffeomorphic iff Γ_1 and Γ_2 are conjugate in $\mathbf{O}(2m)$.

Proof The maximal tori of $\mathbf{SO}(2p-1) \times \mathbf{SO}(2m-2p+1)$ are conjugate in $\mathbf{SO}(2m)$ for all p . By corollary 8, the equivariant torsion of $G_{2m,2p-1}$ at $g \in \mathbf{SO}(2m)$ equals the torsion of S^{2m-1} at g up to a non-zero constant. In [9], Lott and Rothenberg proved that an element of $\mathbf{SO}(2m)$ is determined up to conjugacy in $\mathbf{O}(2m)$ by certain linear combinations of this torsion (see the next section). Thus, the result for the spheres extends to all Graßmannians. \square

Proof of lemma 6 Let χ^K denote the virtual character in the representation ring of K given by

$$\chi^K := \sum_{q=1}^n (-1)^q q \chi(\Lambda^q \text{Ad}_{G/K})$$

where $\chi(\Lambda^q \text{Ad}_{G/K})$ denotes the character of the K -representation $\Lambda^q \text{Ad}_{G/K}$. By classical representation theory, one knows

$$\sum_{q=1}^n (-1)^q q \dim \text{Hom}_K(V_\pi, \Lambda^q \text{Ad}_{G/K}) = \int_K \overline{\chi^K} \cdot \chi_\pi \, d\text{vol}_K. \quad (7)$$

Using the Weyl integral formula this transforms to

$$\frac{1}{\#W_K} \int_{T_K} \overline{\text{Alt}_K\{\rho_K\}} \text{Alt}_K\{\rho_K\} \chi^K \cdot \chi_\pi \, \text{dvol}_{T_K}$$

(where T_K is identified with the quotient of \mathfrak{t}_K by the integral lattice). Classically the restriction of $\text{Alt}_G\{\rho_G\}$ and $\text{Alt}_K\{\rho_K\}$ to \mathfrak{t}_K is given by

$$\text{Alt}_G\{\rho_G\}|_{\mathfrak{t}_K} = \prod_{\alpha \in \Delta_G^+} 2i \sin \pi \alpha|_{\mathfrak{t}_K}$$

and

$$\text{Alt}_K\{\rho_K\} = \prod_{\alpha \in \Delta_K^+} 2i \sin \pi \alpha.$$

Equation (1) shows that the restriction of χ^K is given by

$$\chi^K = \frac{\partial}{\partial s|_{s=1}} \det(1 - s \text{Ad}_{G/K})|_{T_K} \quad (8)$$

$$= \frac{\partial}{\partial s|_{s=1}} (1 - s)^{\text{rk } G - \text{rk } K} \prod_{\alpha \in \Delta_G - \Delta_K} (1 - s e^{2\pi i \alpha}) \quad (9)$$

$$= \begin{cases} -\prod_{\alpha \in \Delta_G - \Delta_K} (1 - e^{2\pi i \alpha}) & \text{if } \text{rk } G = \text{rk } K + 1 \\ 0 & \text{if } \text{rk } G > \text{rk } K + 1 \end{cases} \quad (10)$$

This shows the first part of lemma 6. Assume now that $\text{rk } G = \text{rk } K + 1$. Then

$$\chi^K|_{T_K} = -\frac{\prod_{\alpha \in \Delta_G} 2i \sin \pi \alpha|_{T_K}}{\prod_{\alpha \in \Delta_K} 2i \sin \pi \alpha} = -\frac{\overline{\text{Alt}_G\{\rho_G\}} \text{Alt}_G\{\rho_G\}}{\overline{\text{Alt}_K\{\rho_K\}} \text{Alt}_K\{\rho_K\}},$$

hence equation (7) yields

$$\sum_{q=1}^n (-1)^q q \dim \text{Hom}_K(V_\pi, \Lambda^q \text{Ad}_{G/K}) = -\frac{1}{\#W_K} \int_{T_K} \overline{\text{Alt}_G\{\rho_G\}} \text{Alt}_G\{b_\pi\} \, \text{dvol}_{T_K}$$

which finishes the proof of lemma 6. \square

5 The torsion for symmetric spaces

Classically, a compact symmetric space decomposes as a Riemannian manifold into a product of irreducible symmetric spaces G_ν/K_ν , where the metric

on G_ν/K_ν is induced by a negative real number times the Killing form of G_ν . By lemma 7 the equivariant torsion is non-zero only if $\text{rk } G = \text{rk } K + 1$. Also it is zero for all Lie groups except the circle. By the classification of irreducible symmetric spaces, among them only the odd-dimensional Graßmannians

$$G_{2m,2p-1}(\mathbf{R}) = \mathbf{SO}(2m)/\mathbf{SO}(2p-1) \times \mathbf{SO}(2m-2p+1) \quad (m, p \in \mathbf{N}, m \geq p).$$

and the 5-dimensional space $\mathbf{SU}(3)/\mathbf{SO}(3)$ may have non-zero torsion. Thus, the torsion is zero except for the spaces

$$G_{2m,2p-1} \times G'/K' \text{ and } \mathbf{SU}(3)/\mathbf{SO}(3) \times G'/K', \quad (T)$$

where G'/K' is an arbitrary symmetric space with $\text{rk } G' = \text{rk } K'$. We imbed $\mathbf{SO}(2p-1) \times \mathbf{SO}(2m-2p+1)$ in $\mathbf{SO}(2m)$ as $K = \left\{ \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \mid A \in \mathbf{SO}(2p-1), B \in \mathbf{SO}(2m-2p+1) \right\}$. To diagonalize the standard maximal torus, we imbed $\mathbf{SO}(3)$ in $\mathbf{SU}(3)$ after conjugation by the matrix $\begin{pmatrix} -i & 1 & 0 \\ 0 & 0 & \sqrt{2} \end{pmatrix} / \sqrt{2}$. As maximal tori of G we choose

$$\mathfrak{t}_{\mathbf{SO}(2m)} := \left\{ 2\pi \begin{pmatrix} 0-\lambda_1 & & & \\ \lambda_1 & 0 & & \\ & & \ddots & \\ & & & 0-\lambda_m \\ & & & \lambda_m & 0 \end{pmatrix} \mid \lambda_1, \dots, \lambda_m \in \mathbf{R} \right\}$$

and

$$\mathfrak{t}_{\mathbf{SU}(3)} := \left\{ 2\pi i \begin{pmatrix} \lambda_1 & & \\ & \lambda_2 & \\ & & \lambda_3 \end{pmatrix} \mid \lambda_1 + \lambda_2 + \lambda_3 = 0 \right\}.$$

Let $e_\nu \in \mathfrak{t}_{\mathbf{SO}(2m)}$, $1 \leq \nu \leq m$ resp. $f_\nu \in \mathfrak{t}_{\mathbf{SU}(3)}$, $1 \leq \nu \leq 3$ denote the weight mapping one of the above matrices to λ_ν , ordered according to their index. We set \mathfrak{t}_K as the kernel of e_p resp. f_3 ; thus, these weights generate $L \cong \mathbf{Z}$. They shall be denoted by α_0 . The orbit of α_0 under W_G/W_K is then given by $\{\pm e_\nu\}_{\nu=1}^m$ and $\{f_\nu\}_{\nu=1}^3$, respectively.

Let $\psi : \mathbf{R}/\mathbf{Z} \rightarrow \mathbf{R}$ denote the map

$$[x] \mapsto 2C + \psi([x]) + \psi([-x])$$

where C denotes Euler's constant and ψ is the digamma function evaluated on the fundamental domain $]0, 1]$, i.e. $\psi([x]) := \psi(x + [1 - x])$. We show now the following formula for the equivariant Ray-Singer torsion:

Theorem 10 *Let $G/K \neq S^1$ be a space of type (T) and let α_0 be a generator of L . Then the logarithm of the equivariant torsion for $t \in T_G$ is given by*

$$-\frac{1}{2}Z'_i(0) = -\log \prod_{[w] \in W_G/W_K} \frac{\langle w\alpha_0, \rho_G \rangle_\circ}{\pi} + \frac{1}{2} \sum_{[w] \in W_G/W_K} \tilde{\psi}(w\alpha_0(t))$$

and for general $g \in G$ it is obtained by conjugating g into T_G .

Using the formula for the values of ψ at rational numbers this reproves Ray's result for the spheres [11]. Before proving the theorem we show two auxiliary lemmas. Let S_β denote the reflection of \mathfrak{t}_G^* in the hyperplane orthogonal to $\beta \in \mathfrak{t}_G^*$. We need the following symmetry:

Lemma 11 *Let G/K be a space of the type (T). Then, for all $[w] \in W_G/W_K$ there exists $\tilde{w} \in W_G$ with $(-1)^{\tilde{w}} = 1$ such that*

$$S_{w\alpha_0}\rho_G = \tilde{w}\rho_G.$$

In particular, $(w\alpha_0, \rho_G) = -(\tilde{w}^{-1}w\alpha_0, \rho_G)$ is an integer and the map $[w] \mapsto [\tilde{w}^{-1}w] \in W_G/W_K$ is bijective.

Proof The proof reduces to that for the cases $G_{2m, 2p-1}$ and $\mathbf{SU}(3)/\mathbf{SO}(3)$. Classically, $\rho_{\mathbf{SO}(2m)} = \sum_{\nu=1}^m (m-\nu)e_\nu$ and $\rho_{\mathbf{SU}(3)} = f_1 - f_3$. Hence, $S_{w\alpha_0}\rho_G$ is a weight and thus given by $\tilde{w}\rho_G$ for some unique $\tilde{w} \in W_G$. One observes immediately that $(-1)^{\tilde{w}} = 1$. By the invariance of $\langle \cdot, \cdot \rangle$ under reflections, one shows

$$\langle w\alpha_0, \rho_G \rangle = \langle S_{w\alpha_0}w\alpha_0, S_{w\alpha_0}\rho_G \rangle = -\langle w\alpha_0, \tilde{w}\rho_G \rangle = -\langle \tilde{w}^{-1}w\alpha_0, \rho_G \rangle;$$

as $S_{w\alpha_0}\rho_G = \rho_G - (w\alpha_0, \rho_G)w\alpha_0$, one notices that $(w\alpha_0, \rho_G)$ is an integer. As $S_{\tilde{w}^{-1}w\alpha_0}\rho_G = \tilde{w}^{-1}S_{w\alpha_0}(\tilde{w}\rho_G) = \tilde{w}^{-1}\rho_G$, the map $[w] \mapsto [\tilde{w}^{-1}w]$ is an involution. \square

Also, we need a lemma about values of zeta functions at zero. For $p, n \in \mathbf{Z}$ and $h, \phi \in \mathbf{R}$, let $\tilde{\zeta}_{n,p,h}(s, \phi)$ denote the zeta function

$$\tilde{\zeta}_{n,p,h}(s, \phi) := \sum_{\substack{k \in \mathbf{Z} \\ k(k+p) > 0}} \frac{k^n e^{ik\phi} (h + \log |k|)}{(k(k+p))^s} + \sum_{\substack{k \in \mathbf{Z} \setminus \{0\} \\ k(k+p) \leq 0}} \frac{k^n e^{ik\phi} (h + \log |k|)}{|k|^{2s}}$$

for $\Re s > \frac{n+1}{2}$. This zeta function has a meromorphic extension to the complex plane.

Lemma 12 *The value at zero of $\tilde{\zeta}$ is independent of p for $n \in \mathbf{N}_0$. For $n = 0$ it takes the value*

$$\tilde{\zeta}_{n,p,h}(0, \phi) = \log 2\pi + \tilde{\psi}\left(\frac{\phi}{2\pi}\right) - h.$$

Proof The proof may be given easily by applying the more general result of [8, Lemma 7]; instead, for the case $\phi \not\equiv 0 \pmod{2\pi}$ we shall give an alternative proof which is better adapted to this particular situation. Choose $\phi \in]0, 2\pi[$. First, observe that in this case

$$\tilde{\zeta}_{n,p,h}(s, \phi) = i^{-n-1} \left(\frac{\partial}{\partial \phi} \right)^{n+1} \tilde{\zeta}_{-1,p,h}(s, \phi).$$

The series defining $\tilde{\zeta}_{-1,p,h}(s, \phi)$ converges at $s = 0$. Using Kummer's Fourier series for the logarithm of the Gamma function [7, sect. 5] one finds

$$\begin{aligned} \tilde{\zeta}_{-1,p,h}(0, \phi) &= \sum_{k>0} \frac{2i \sin k\phi \cdot (h + \log k)}{k} \\ &= i(C + \log 2\pi - h)(\phi - \pi) + i\pi \log \Gamma\left(\frac{\phi}{2\pi}\right) - i\pi \log \Gamma\left(1 - \frac{\phi}{2\pi}\right). \end{aligned}$$

To prove the lemma for the case $\phi \equiv 0$ one may use the Taylor expansion for $|k+p|^{-s} = |k|^{-s}|1+p/k|^{-s}$ as in [7, sect. 6]. \square

Proof of theorem 10 The derivative of $Z(s)$ is given by

$$\begin{aligned} Z'(s) &= - \sum_{\substack{[w] \in W_G/W_K \\ k \in \mathbf{Z} \\ k(k+(w\alpha_0, \rho_G)) > 0}} \frac{\chi_{\rho_G+k w\alpha_0} (\log |k| + \log |k + (w\alpha_0, \rho_G)| + \log \|\alpha_0\|_{\diamond}^2)}{(k(k + (w\alpha_0, \rho_G)) \|\alpha_0\|_{\diamond}^2)^s} \\ &= - \sum_{\substack{[w] \in W_G/W_K \\ k \in \mathbf{Z} \\ k(k+(w\alpha_0, \rho_G)) > 0}} \frac{\chi_{\rho_G+k w\alpha_0} \log |k|}{(k(k + (w\alpha_0, \rho_G)) \|\alpha_0\|_{\diamond}^2)^s} \\ &\quad - \sum_{\substack{[w] \in W_G/W_K \\ k \in \mathbf{Z} \\ k(k-(w\alpha_0, \rho_G)) > 0}} \frac{\chi_{\rho_G+(k-(w\alpha_0, \rho_G))w\alpha_0} \log |k|}{(k(k - (w\alpha_0, \rho_G)) \|\alpha_0\|_{\diamond}^2)^s} - Z(s) \log \|\alpha_0\|_{\diamond}^2. \quad (11) \end{aligned}$$

By lemma 11 one obtains

$$\rho_G + (k - (w\alpha_0, \rho_G))w\alpha_0 = S_{w\alpha_0}(\rho_G - kw\alpha_0) = \tilde{w}(\rho_G + k\tilde{w}^{-1}w\alpha_0)$$

with $(-1)^{\tilde{w}} = 1$. Hence the second sum in line (11) equals

$$- \sum_{\substack{[w] \in W_G/W_K \\ k \in \mathbf{Z} \\ k(k + (\tilde{w}^{-1}w\alpha_0, \rho_G)) > 0}} \frac{\chi_{\rho_G + k\tilde{w}^{-1}w\alpha_0} \log |k|}{(k(k + (\tilde{w}^{-1}w\alpha_0, \rho_G)) \|\alpha_0\|_0^2)^s}.$$

By the bijectivity of $[w] \mapsto [\tilde{w}^{-1}w]$ one finds

$$Z'(s) = \sum_{\substack{[w] \in W_G/W_K \\ k \in \mathbf{Z} \\ k(k + (w\alpha_0, \rho_G)) > 0}} \frac{\chi_{\rho_G + kw\alpha_0} (-2 \log |k| - \log \|\alpha_0\|_0^2)}{(k(k + (w\alpha_0, \rho_G)) \|\alpha_0\|_0^2)^s}.$$

Fix $g \in G$ and consider the character $\chi_{\rho_G + kw\alpha_0}(g)$ as a function in $k \in \mathbf{Z}$. One shows easily that $\chi_{\rho_G + kw\alpha_0}(g)$ is a linear combination of functions of the type $k \mapsto k^n e^{i\phi k}$ ($n \in \mathbf{Z}$, $\phi \in \mathbf{R}$) [8, eq. 81]. The characters $\chi_{\rho_G + kw\alpha_0}$ are zero for $0 > k(k + (w\alpha_0, \rho_G)) = (\|\rho_G + k\alpha_0\|_0^2 - \|\rho_G\|_0^2) / \|\alpha_0\|_0^2$. For $k = -(w\alpha_0, \rho_G)$ we have by lemma 6 the result

$$\chi_{\rho_G - (w\alpha_0, \rho_G)w\alpha_0} = \chi_{S_{w\alpha_0}\rho_G} = (-1)^{\tilde{w}} = 1.$$

Thus, we may write $Z'(s)$ as

$$Z'(s) = \sum_{[w] \in W_G/W_K} \left(\sum_{\substack{k \in \mathbf{Z} \\ k(k + (w\alpha_0, \rho_G)) > 0}} \frac{\chi_{\rho_G + kw\alpha_0} (-2 \log |k| - \log \|\alpha_0\|_0^2)}{(k(k + (w\alpha_0, \rho_G)) \|\alpha_0\|_0^2)^s} \right. \\ \left. + \sum_{\substack{k \in \mathbf{Z} \setminus \{0\} \\ k(k + (w\alpha_0, \rho_G)) \leq 0}} \frac{\chi_{\rho_G + kw\alpha_0} (-2 \log |k| - \log \|\alpha_0\|_0^2)}{\|k\alpha_0\|_0^{2s}} + \frac{\log(w\alpha_0, \rho_G)^2 \|\alpha_0\|_0^2}{|(w\alpha_0, \rho_G)|^{2s} \|\alpha_0\|_0^{2s}} \right).$$

Lemma 8 states that the value at zero of $Z'(s)$ equals the value at zero of the zeta function

$$\hat{Z}(s) = \sum_{[w] \in W_G/W_K} \left(\sum_{k \in \mathbf{Z} \setminus \{0\}} \frac{\chi_{\rho_G + kw\alpha_0} (-2 \log |k| - \log \|\alpha_0\|_0^2)}{\|k\alpha_0\|_0^{2s}} + \frac{\log(w\alpha_0, \rho_G)^2 \|\alpha_0\|_0^2}{|(w\alpha_0, \rho_G)|^{2s} \|\alpha_0\|_0^{2s}} \right).$$

The sum of the characters over W_G/W_K equals

$$\sum_{[w] \in W_G/W_K} \chi_{\rho_G + kw\alpha_0} = \frac{1}{\#W_K} \sum_{w \in W_G} \frac{\sum_{w' \in W_G} (-1)^{w'} e^{2\pi i(w'\rho_G + kw'w\alpha_0)}}{\text{Alt}_G\{\rho_G\}}$$

$$\begin{aligned}
&= \frac{1}{\#W_K} \frac{\sum_{w \in W_G} e^{2\pi i k w \alpha_0} \sum_{w' \in W_G} (-1)^{w'} e^{2\pi i w' \rho_G}}{\text{Alt}_G\{\rho_G\}} \\
&= \sum_{[w] \in W_G/W_K} e^{2\pi i k w \alpha_0}.
\end{aligned}$$

Hence we may apply the formula in lemma 8 to $\tilde{Z}(s)$. This proves the theorem. \square

To proof theorem 3 for the case $\mathbf{SU}(3)/\mathbf{SO}(3)$ we apply the method used by Lott and Rothenberg for the spheres. In [9, Prop. 32] they proved

Lemma 13 *Let $x_\nu \in \mathbf{R}/\mathbf{Z}$, $1 \leq \nu \leq N$ be elements of the circle. Then the x_ν are determined up to order and sign by the sequence*

$$\left(\sum_{\nu=1}^N \tilde{\psi}(n x_\nu) \right)_{n \in \mathbf{Z}}.$$

This lemma has been proven by Franz for rational x_ν . For the action of a torus element $t := 2\pi i \text{diag}(\lambda_1, \lambda_2, \lambda_3)$ this means that λ_1, λ_2 and λ_3 are determined up to order and sign by the torsion τ_n of powers of t . As $\lambda_1 + \lambda_2 + \lambda_3 = 0$, all λ_ν have to change sign if one of them does so. Now a change of the order is an element of $W_{\mathbf{SU}(3)}$, thus given by conjugation with an element of $\mathbf{SU}(3)$. A change of the sign of all λ_ν is obtained by the symmetry around $[1] \in \mathbf{SU}(3)/\mathbf{SO}(3)$ composed with interchanging λ_1 and λ_2 (recall our non-standard imbedding of $\mathbf{SO}(3)$). Thus, all isometries which have the same torsion are actually isometric.

Acknowledgement *During the preparation of this paper I have been supported by the Max-Planck-Institut in Bonn.*

References

- [1] A. L. Besse, *Einstein manifolds*, Springer, 1987.
- [2] J.-M. Bismut and Weiping Zhang, *Milnor and Ray-Singer metrics on the equivariant determinant of a flat vector bundle*, GAFA **4** (1994), 136–212.

- [3] R. Bott, *The index theorem for homogeneous differential operators*, Differential and Combinatorial Topology (S. S. Cairns, ed.), Princeton university press, 1965.
- [4] G. de Rham, *Reidemeister's torsion invariant and rotations of S^n* , Differential Analysis, Tata Institute and Oxford University Press, 1964, pp. 27–36.
- [5] W. Franz, *über die Torsion einer Überdeckung*, J. Reine Angew. Math. **173** (1935), 245–254.
- [6] A. Ikeda and Y. Taniguchi, *Spectra and eigenforms of the Laplacian on S^n and $\mathbf{P}^n\mathbf{C}$* , Osaka J. Math. **15** (1978), 515–546.
- [7] K. Köhler, *Equivariant analytic torsion on $\mathbf{P}^n\mathbf{C}$* , Math. Ann. **297** (1993), 553–565.
- [8] ———, *Holomorphic torsion on Hermitian symmetric spaces*, preprint 1, MPI, 1994.
- [9] J. Lott and M. Rothenberg, *Analytic torsion for group actions*, J. Diff. Geom. **34** (1991), 431–481.
- [10] H. Moscovici and R. J. Stanton, *R-torsion and zeta functions for locally symmetric manifolds*, Inv. math. **105** (1991), 185–216.
- [11] D. B. Ray, *Reidemeister torsion and the Laplacian on lens spaces*, Adv. in Math. **4** (1970), 109–126.
- [12] D. B. Ray and I. M. Singer, *Analytic torsion for complex manifolds*, Ann. Math. **98** (1973), 154–177.