

# Locally constant $n$ -operads as higher braided operads

M.A. Batanin\*

Macquarie University, NSW 2109, Australia

e-mail: mbatanin@ics.mq.edu.au

May 5, 2008

## Abstract

We introduce a category of *locally constant  $n$ -operads* which can be considered as the category of higher braided operads. For  $n = 1, 2, \infty$  the homotopy category of locally constant  $n$ -operads is equivalent to the homotopy category of classical nonsymmetric, braided and symmetric operads correspondingly.

1991 Math. Subj. Class. 18D20 , 18D50, 55P48

## 1 Introduction

It is well known that contractible nonsymmetric operads detect 1-fold loop spaces, contractible braided operads detect 2-fold loop spaces and that contractible symmetric operads detect  $\infty$ -fold loop spaces. A natural question arises : is there a sequence of groups  $G^{(n)} = \{G_k^{(n)}\}_{k \geq 0}$  together with a notion of  $G^{(n)}$ -operad, which we would call  *$n$ -braided operad*, such that the algebras of a contractible such operad are  $n$ -fold loop spaces? With some natural minor assumptions one can prove that the answer on the above question is negative. This is because for such an operad  $A$  the quotient  $A_k/G_k^{(n)}$  is a  $K(G_k^{(n)}, 1)$ -space. One can show, however, that such a quotient must have a homotopy type of the space of unordered configurations of  $k$  points in  $\mathbb{R}^n$ , which is a  $K(\pi, 1)$ -space only for  $n = 1, 2, \infty$ .

In this paper we show that there is a category of operads which we can think of as a correct replacement for the nonexistent category of  $G^{(n)}$ -operads in all dimensions. We call them *locally constant  $n$ -operads*. For  $n = 1, 2, \infty$  the homotopy category of locally constant  $n$ -operads is equivalent to the homotopy category of classical nonsymmetric, braided and symmetric operads correspondingly.

---

\*The author holds the Scott Russell Johnson Fellowship in the Centre of Australian Category Theory at Macquarie University

## 2 Symmetric and braided operads

For a natural number  $n$  we will denote by  $[n]$  the ordinal

$$0 < 1 < \dots < n.$$

A morphism from  $[n] \rightarrow [k]$  is any function between underlying sets. It can be order preserving or not. It is clear that we then have a category. We denote this category by  $\Omega^s$ . Of course,  $\Omega^s$  is equivalent to the category of finite sets. In particular, the symmetric group  $S_{n+1}$  is the group of automorphisms of  $[n]$ .

Let  $\sigma : [n] \rightarrow [k]$  be a morphism in  $\Omega^s$  and let  $0 \leq i \leq k$ . Then the preimage  $\sigma^{-1}(i)$  has a linear order induced from  $[n]$ . Hence, there exists a unique object  $[n_i] \in \Omega^s$  and a unique order preserving bijection  $[n_i] \rightarrow \sigma^{-1}(i)$ . We will call  $[n_i]$  the *fiber* of  $\sigma$  over  $i$  and will denote it  $\sigma^{-1}(i)$  or  $[n_i]$ .

Analogously, given a composite of morphisms in  $\Omega^s$  :

$$[n] \xrightarrow{\sigma} [l] \xrightarrow{\omega} [k] \tag{1}$$

we will denote  $\sigma_i$  the *i-th fiber* of  $\sigma$ ; i.e. the pullback

$$\begin{array}{ccccc} \sigma^{-1}(\omega^{-1}(i)) & \xrightarrow{\sigma_i} & \omega^{-1}(i) & \longrightarrow & [1] \\ \downarrow & & \downarrow & & \downarrow \xi_i \\ [n] & \xrightarrow{\sigma} & [l] & \xrightarrow{\omega} & [k] . \end{array}$$

Let  $\mathcal{S}$  be the subcategory of bijections in  $\Omega^s$ . This is a strict monoidal groupoid with tensor product  $\oplus$  given by ordinal sum and with  $[-1]$  as its unital object.

A *right symmetric collection* in a symmetric monoidal category  $V$  is a functor  $A : \mathcal{S}^{op} \rightarrow V$ . The value of  $A$  on an object  $[n]$  will be denoted  $A_n$ . Notice, that this is not a standard operadic notation. Classically, the notation for  $A_{[n]}$  is  $A_{n+1}$  to stress the fact that  $A_{n+1}$  is the space of operations of arity  $n+1$ .

The following definition is classical May definition [6] of symmetric operad.

**Definition 2.1** *A (right) symmetric operad in  $V$  is a right symmetric collection  $A$  equipped with the following additional structure:*

- a morphism  $e : I \rightarrow A_0$
- for every order preserving map  $\sigma : [n] \rightarrow [k]$  in  $\Omega^s$  a morphism :

$$\mu_\sigma : A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) \longrightarrow A_n,$$

where  $[n_i] = \sigma^{-1}(i)$ .

*They must satisfy the following identities:*

1. for any composite of order preserving morphisms in  $\Omega^s$

$$[n] \xrightarrow{\sigma} [l] \xrightarrow{\omega} [k],$$

the following diagram commutes

$$\begin{array}{ccc}
 A_k \otimes A_{l_\bullet} \otimes A_{n_0^\bullet} \otimes \dots \otimes A_{n_i^\bullet} \otimes \dots \otimes A_{n_k^\bullet} & \xrightarrow{\cong} & A_k \otimes A_{l_1} \otimes A_{n_0^\bullet} \otimes \dots \otimes A_{l_i} \otimes A_{n_i^\bullet} \otimes \dots \otimes A_{l_k} \otimes A_{n_k^\bullet} \\
 \downarrow & & \downarrow \\
 A_l \otimes A_{n_0^\bullet} \otimes \dots \otimes A_{n_i^\bullet} \otimes \dots \otimes A_{n_k^\bullet} & & A_k \otimes A_{n_\bullet} \\
 & \searrow & \swarrow \\
 & A_n & 
 \end{array}$$

Here

$$A_{l_\bullet} = A_{l_0} \otimes \dots \otimes A_{l_k},$$

$$A_{n_i^\bullet} = A_{n_i^0} \otimes \dots \otimes A_{n_i^{m_i}}$$

and

$$A_{n_\bullet} = A_{n_0} \otimes \dots \otimes A_{n_k};$$

2. for an identity  $\sigma = id : [n] \rightarrow [n]$  the diagram

$$\begin{array}{ccc}
 A_n \otimes A_0 \otimes \dots \otimes A_0 & \longleftarrow & A_n \otimes I \otimes \dots \otimes I \\
 \downarrow & \swarrow id & \\
 A_n & & 
 \end{array}$$

commutes;

3. for the unique morphism  $[n] \rightarrow [0]$  the diagram

$$\begin{array}{ccc}
 A_0 \otimes A_n & \longleftarrow & I \otimes A_n \\
 \downarrow & \swarrow id & \\
 A_n & & 
 \end{array}$$

commutes.

The following equivariance conditions are also required:

1. For any order preserving  $\sigma : [n] \rightarrow [k]$  and any bijection  $\rho : [k] \rightarrow [k]$  the following diagram commutes:

$$\begin{array}{ccc}
A_k \otimes (A_{n_{\rho(0)}} \otimes \dots \otimes A_{n_{\rho(k)}}) & \xrightarrow{\mu_\sigma} & A_n \\
\uparrow A(\rho) \otimes \tau(\rho) & & \uparrow A(\pi) \\
A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) & \xrightarrow{\mu_\sigma} & A_n \quad ,
\end{array}$$

where  $\tau(\rho)$  is the symmetry in  $V$  which corresponds to permutation  $\rho$  and  $\pi = \Gamma_S(\rho; 1, \dots, 1)$  is the permutation, which permutes the fibers  $[n_0], \dots, [n_k]$  according to  $\rho$  and whose restriction on each fiber is an identity.

2. For any order preserving  $\sigma : [n] \rightarrow [k]$  and any set of bijections  $\rho_i : [n_i] \rightarrow [n_i], 0 \leq i \leq k$ , the following diagram commutes

$$\begin{array}{ccc}
A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) & \xrightarrow{\mu_\sigma} & A_n \\
\uparrow id \otimes A(\rho_0) \otimes \dots \otimes A(\rho_k) & & \uparrow A(\rho_0 \oplus \dots \oplus \rho_k) \\
A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) & \xrightarrow{\mu_\sigma} & A_n \quad ,
\end{array}$$

We can give an alternative definition of symmetric operad [2].

**Definition 2.2** A (right) symmetric operad in  $V$  is a right symmetric collection  $A$  equipped with the following additional structure:

- a morphism  $e : I \rightarrow A_0$
- for every order preserving map  $\sigma : [n] \rightarrow [k]$  in  $\Omega^s$  a morphism:

$$\mu_\sigma : A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) \longrightarrow A_n,$$

where  $[n_i] = \sigma^{-1}(i)$ .

They must satisfy the same conditions as in the definition 2.1 with respect to order preserving maps and identities but the equivariance conditions are replaced by the following:

1. For every commutative diagram in  $\Omega^s$

$$\begin{array}{ccc}
[n'] & \xrightarrow{\sigma'} & [k'] \\
\pi \downarrow & & \downarrow \rho \\
[n] & \xrightarrow{\sigma} & [k]
\end{array}$$

whose vertical maps are bijections and whose horizontal maps are order preserving the following diagram commutes:

$$\begin{array}{ccc}
A_{k'} \otimes (A_{n'_{\rho(0)}} \otimes \dots \otimes A_{n'_{\rho(k)}}) & \xrightarrow{\mu_{\sigma'}} & A_{n'} \\
\uparrow A(\rho) \otimes \tau(\rho) & & \uparrow A(\pi) \\
A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) & \xrightarrow{\mu_{\sigma}} & A_n
\end{array},$$

where  $\tau(\rho)$  is the symmetry in  $V$  which corresponds to permutation  $\rho$ .

2. For every commutative diagram in  $\Omega^s$

$$\begin{array}{ccc}
[n''] & \xrightarrow{\sigma'} & [n'] \\
\sigma \downarrow & & \downarrow \eta' \\
[n] & \xrightarrow{\eta} & [k]
\end{array}$$

where  $\sigma, \sigma'$  are bijections and  $\eta, \eta'$  are order preserving maps, the following diagram commutes

$$\begin{array}{ccc}
A_k \otimes (A_{n'_0} \otimes \dots \otimes A_{n'_k}) & \xrightarrow{\mu_{\eta'}} & A_{n'} \\
\downarrow 1 \otimes A(\sigma'_0) \otimes \dots \otimes A(\sigma'_k) & & \downarrow A(\sigma') \\
A_k \otimes (A_{n''_0} \otimes \dots \otimes A_{n''_k}) & & A_{n''} \\
\uparrow 1 \otimes A(\sigma_0) \otimes \dots \otimes A(\sigma_k) & & \uparrow A(\sigma) \\
A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) & \xrightarrow{\mu_{\eta}} & A_n
\end{array}$$

**Proposition 2.1** *The definition 2.1 and 2.2 are equivalent.*

We leave this proposition as an exercise for the reader.

Let  $\mathcal{B}r$  be the groupoid of braid groups. We will think that the objects of  $\mathcal{B}r$  are ordinals. There is a monoidal structure on  $\mathcal{B}r$  given by ordinal sum on objects and concatenation of braids on morphism. The ordinal  $[-1]$  is the unital object.

The following is the definition of braided operad from [4]. A *right braided collection* in a symmetric monoidal category  $V$  is a functor  $A : \mathcal{B}r^{op} \rightarrow V$ . The value of  $A$  on an object  $[n]$  will be denoted  $A_n$ .

**Definition 2.3** A *right braided operad* in  $V$  is a right braided collection  $A$  equipped with the following additional structure:

- a morphism  $e : I \rightarrow A_0$
- for every order preserving map  $\sigma : [n] \rightarrow [k]$  in  $\Omega^s$  a morphism :

$$\mu_\sigma : A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) \longrightarrow A_n,$$

where  $[n_i] = \sigma^{-1}(i)$ .

They must satisfy the identities (1-3) from the definition 2.1 and the following two equivariancy conditions:

1. For any order preserving  $\sigma : [n] \rightarrow [k]$  and any braid  $\rho : [k] \rightarrow [k]$  the following diagram commutes:

$$\begin{array}{ccc} A_k \otimes (A_{n_{\rho(0)}} \otimes \dots \otimes A_{n_{\rho(k)}}) & \xrightarrow{\mu_\sigma} & A_n \\ \uparrow A(\rho) \otimes \tau(\rho) & & \uparrow A(\pi) \\ A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) & \xrightarrow{\mu_\sigma} & A_n \end{array} ,$$

where  $\tau(\rho)$  is the symmetry in  $V$  which corresponds to the braid  $\rho$  and  $\pi = \Gamma_B(\rho; 1, \dots, 1)$  is a braid obtained from  $\rho$  by replacing the  $i$ -th strand of  $\rho$  by  $n_i$  parallel strands for each  $i$ .

2. For any order preserving  $\sigma : [n] \rightarrow [k]$  and any set of braids  $\rho_i : [n_i] \rightarrow [n_i], 0 \leq i \leq k$ , the following diagram commutes

$$\begin{array}{ccc} A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) & \xrightarrow{\mu_\sigma} & A_n \\ \uparrow id \otimes A(\rho_0) \otimes \dots \otimes A(\rho_k) & & \uparrow A(\rho_0 \oplus \dots \oplus \rho_k) \\ A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) & \xrightarrow{\mu_\sigma} & A_n \end{array} ,$$

### 3 The category of $n$ -ordinals and of quasibijections

**Definition 3.1** Let  $T$  be a finite set equipped with  $n$  binary relations  $<_0, \dots, <_{n-1}$ .  $T$  is called an  $n$ -ordinal if these relations satisfy the following properties

1.  $<_p$  is nonreflexive;
2. for every pair  $a, b \in T$ , there exists exactly one  $p$  such that

$$a <_p b \text{ or } b <_p a;$$

3. if  $a <_p b$  and  $b <_q c$  then  $a <_{\min(p,q)} c$ .

Every  $n$ -ordinal can be represented as a pruned planar tree with  $n$  levels or as  $n$ -dimensional globular graph (see [1] for a discussion).

**Definition 3.2** A map of  $n$ -ordinals

$$\sigma : T \rightarrow S$$

is a map  $\sigma : T \rightarrow S$  of underlying sets such that

$$i <_p j \text{ in } T$$

implies that

1.  $\sigma(i) <_r \sigma(j)$  for some  $r \geq p$  or
2.  $\sigma(i) = \sigma(j)$  or
3.  $\sigma(j) <_r \sigma(i)$  for  $r > p$ .

For every  $i \in S$  the preimage  $\sigma^{-1}(i)$  (the fiber of  $\sigma$  over  $i$ ) has a natural structure of an  $n$ -ordinal.

We denote by  $Ord(n)$  the skeletal category of  $n$ -ordinals. The category  $Ord(n)$  is monoidal. The monoidal structure  $\oplus$  is defined as follows. For two  $n$ -ordinals  $S$  and  $T$  the  $n$ -ordinal  $S \oplus T$  has as an underlying set the union of underlying sets of  $S$  and  $T$ . The orders  $<_k$  restricted to the elements of  $S$  and  $T$  coincide with respective orders on  $S$  and  $T$ . and  $a <_0 b$  if  $a \in S$  and  $b \in T$ . The unital object for this monoidal structure is empty  $n$ -ordinal.

A total order on an  $n$ -ordinal  $T$  is a linear order  $<$  on the underlying set of  $T$  defined as follows:

$$a < b \text{ iff } a <_r b \text{ for some } 0 \leq r \leq n - 1 .$$

We will denote by  $[T]$  the set  $T$  with its total linear order. In this way we have a monoidal functor

$$[-] : Ord(n) \rightarrow \Omega^s .$$

We also introduce the category of  $\infty$ -ordinals  $Ord(\infty)$ .

**Definition 3.3** Let  $T$  be a finite set equipped with a sequence of binary relations  $<_0, <_{-1} \dots, <_p, <_{p-1} \dots$  for all integers  $p \leq 0$ . The set  $T$  is called an  $\infty$ -ordinal if these relations satisfy the following properties:

1.  $<_p$  is nonreflexive;
2. for every pair  $a, b \in T$ , there exists exactly one  $p$  such that
$$a <_p b \text{ or } b <_p a;$$
3. if  $a <_p b$  and  $b <_q c$  then  $a <_{\min(p,q)} c$ .

The definition of morphisms between  $\infty$  ordinals coincides with the Definition 3.2. The category  $Ord(\infty)$  is the skeletal category of  $\infty$ -ordinals. As for  $Ord(n)$  we have a functor of total order

$$[-] : Ord(\infty) \rightarrow \Omega^s.$$

For a  $k$ -ordinal  $R$ ,  $k \leq n$  we consider its  $(n - k)$ -th vertical suspension  $S^{n-k}R$  which is an  $n$ -ordinal with the underlying set  $R$ , and the order  $<_m$  equal the order  $<_{m-k}$  on  $R$  (so  $<_m$  are empty for  $0 \leq m < n - k$ .) We also can consider the horizontal  $(n - 1)$ -suspension  $T^{n-k}R$  which is a  $n$ -ordinal with the underlying set  $R$ , and the order  $<_m$  equal the order on  $R$  (so  $<_m$  are empty for  $k - 1 < m \leq n - 1$ .)

The vertical suspension provides us with a functor  $S : Ord(n) \rightarrow Ord(n+1)$ . We also define an  $\infty$ -suspension functor  $Ord(n) \rightarrow Ord(\infty)$  as follows. For an  $n$ -ordinal  $T$  its  $\infty$ -suspension is an  $\infty$ -ordinal  $S^\infty T$  whose underlying set is the same as the underlying set of  $T$  and  $a <_p b$  in  $S^\infty T$  if  $a <_{n+p-1} b$  in  $T$ . It is not hard to see that the sequence

$$Ord(0) \xrightarrow{S} Ord(1) \xrightarrow{S} Ord(2) \longrightarrow \dots \xrightarrow{S} Ord(n) \longrightarrow \dots \xrightarrow{S^\infty} Ord(\infty),$$

exhibits  $Ord(\infty)$  as a colimit of  $Ord(n)$ .

The category  $Ord(\infty)$  is not monoidal, however. But in practice, calculations in  $Ord(\infty)$  amount to the calculations inside one of  $Ord(n)$  with finite  $n$ , and we often can use the monoidal structure in  $Ord(n)$ .

**Definition 3.4** A map of  $n$ -ordinals is called quasibijection if it is a bijection of the underlying sets.

Let  $Q_n$ ,  $1 \leq n \leq \infty$  be the subcategory of quasibijections of  $Ord(n)$ . The total order functor induces then a functor which we will denote by the same symbol:

$$[-] : Q_n \rightarrow \mathcal{S}.$$

**Definition 3.5** A map  $\sigma$  of  $n$ -ordinals  $1 \leq n \leq \infty$  is called order preserving if it preserves the total orders in the usual sense or equivalently only conditions 1 and 2 from the Definition 3.2 hold for  $\sigma$ .

**Lemma 3.1** For every morphism  $\sigma : T \rightarrow S$  in  $Ord(n)$   $1 \leq n \leq \infty$  there exists a factorisation

$$T \xrightarrow{\pi} T' \xrightarrow{\nu} S$$

where  $\pi$  is a quasibijection, and  $\nu$  is an order preserving and  $\pi$  preserves total order on fibers of  $\nu$ .



**Proof.** For  $n = 1$  this factorisation is trivial, since all maps of 1-ordinals are order preserving.

Let  $n = 2$ . Let  $\sigma : T \rightarrow S$  be a map of 2-ordinals and let  $S = S[k]$  be a suspension of 1-ordinal  $[k]$ . Let  $T'$  be the 2-ordinal whose underlying set is the same as of  $T$ , whose only nonempty order is  $<_1$  and whose total order coincides with  $[T]$ . So  $T'$  itself is a vertically suspended 1-ordinal. Now, one can factorise the map  $[\sigma] : [T] \rightarrow [S]$  in  $\Omega^s$

$$[T] \xrightarrow{\pi} [T'] \xrightarrow{\nu} [S]$$

with  $\nu$  being total order preserving and  $\pi$  a bijection which preserves the order on fibers of  $\sigma$  [2]. Obviously,  $\nu$  can be considered as a map of 2-ordinals and it is order preserving. Let us check that  $\pi$  is also a map of 2-ordinals. Indeed, if  $i, j$  are from the same fiber of  $\sigma$  then  $\pi$  preserves their order. If  $i <_0 j$  in  $T$  and they are from different fibers then there is no restriction on  $\pi$  since  $T'$  is a suspended 1-ordinal. Finally, if  $i <_1 j$  in  $T$  and they are from different fibers then  $\sigma(i) <_1 \sigma(j)$ , so  $\pi(i) <_1 \pi(j)$  because  $\nu$  is order preserving.

Finally, if  $S$  is an arbitrary 2-ordinal then  $S = S_1 \oplus \dots \oplus S_k$  for some suspended 1-ordinals  $S_1, \dots, S_k$  and moreover,

$$\sigma = \sigma_1 \oplus \dots \oplus \sigma_k : T = T_1 \oplus \dots \oplus T_k \rightarrow S_1 \oplus \dots \oplus S_k.$$

By applying the previous result to each  $\sigma_k$  we obtain a required factorisation of  $\sigma$ .

The factorisation for  $n > 2$  can be obtained similarly. □

## 4 Quasisymmetric $n$ -operads.

We now recall the definition of pruned  $(n - 1)$ -terminal  $n$ -operad [1]. Since we do not need other types of  $n$ -operads in this paper we will call them simply  $n$ -operads. The notation  $U_n$  means the terminal  $n$ -ordinal.

Let  $V$  be a symmetric monoidal category. For a morphism of  $n$ -ordinals  $\sigma : T \rightarrow S$  the  $n$ -ordinal  $T_i$  is the fiber  $\sigma^{-1}(i)$ .

**Definition 4.1** *An  $n$ -operad in  $V$  is a collection  $A_T$ ,  $T \in \text{Ord}(n)$  of objects of  $V$  equipped with the following structure :*

- a morphism  $e : I \rightarrow A_{U_n}$  (the unit);
- for every morphism  $\sigma : T \rightarrow S$  in  $\text{Ord}(n)$ , a morphism

$$m_\sigma : A_S \otimes A_{T_0} \otimes \dots \otimes A_{T_k} \rightarrow A_T \quad (\text{the multiplication}).$$

*They must satisfy the following identities:*

- for any composite  $T \xrightarrow{\sigma} S \xrightarrow{\omega} R$ , the associativity diagram

$$\begin{array}{ccc}
A_R \otimes A_{S_\bullet} \otimes A_{T_0^\bullet} \otimes \dots \otimes A_{T_i^\bullet} \otimes \dots \otimes A_{T_k^\bullet} & \simeq & A_R \otimes A_{S_0} \otimes A_{T_1^\bullet} \otimes \dots \otimes A_{S_i} \otimes A_{T_i^\bullet} \otimes \dots \otimes A_{S_k} \otimes A_{T_k^\bullet} \\
\downarrow & & \downarrow \\
A_S \otimes A_{T_1^\bullet} \otimes \dots \otimes A_{T_i^\bullet} \otimes \dots \otimes A_{T_k^\bullet} & & A_R \otimes A_{T_\bullet} \\
& \searrow & \swarrow \\
& A_T &
\end{array}$$

commutes, where

$$A_{S_\bullet} = A_{S_0} \otimes \dots \otimes A_{S_k},$$

$$A_{T_i^\bullet} = A_{T_i^0} \otimes \dots \otimes A_{T_i^{m_i}}$$

and

$$A_{T_\bullet} = A_{T_0} \otimes \dots \otimes A_{T_k};$$

- for an identity  $\sigma = id : T \rightarrow T$  the diagram

$$\begin{array}{ccc}
A_T \otimes A_{U_n} \otimes \dots \otimes A_{U_n} & \longleftarrow & A_T \otimes I \otimes \dots \otimes I \\
\downarrow & \swarrow id & \\
A_T & &
\end{array}$$

commutes;

- for the unique morphism  $T \rightarrow U_n$  the diagram

$$\begin{array}{ccc}
A_{U_n} \otimes A_T & \longleftarrow & I \otimes A_T \\
\downarrow & \swarrow id & \\
A_T & &
\end{array}$$

commutes.

Let  $\sigma : T \rightarrow S$  be a quasibijection and  $A$  be a pruned  $n$ -operad. Since a fiber of  $\sigma$  is the terminal  $n$ -ordinal  $U_n$ , the multiplication

$$\mu_\sigma : A_S \otimes (A_{U_n} \otimes \dots \otimes A_{U_n}) \longrightarrow A_T$$

in composition with the morphism

$$A_S \rightarrow A_S \otimes (I \otimes \dots \otimes I) \rightarrow A_S \otimes (A_{U_n} \otimes \dots \otimes A_{U_n})$$

induces a morphism

$$A(\sigma) : A_S \rightarrow A_T.$$

It is not hard to see that in this way  $A$  becomes a contravariant functor on  $Q_n$ .

**Definition 4.2** We call a pruned  $n$ -operad  $A$  quasisymmetric if for every quasibijection  $\sigma : T \rightarrow S$  the morphism

$$A(\sigma) : A_S \rightarrow A_T$$

is an isomorphism.

Desymmetrisation functor from symmetric to  $n$ -operads for finite  $n$  was defined in [2] using pulling back along the functor  $[-] : Ord(n) \rightarrow \Omega^s$ . It was shown that this functor has a left adjoint which we call symmetrisation. We can obviously extend these definitions to  $n = \infty$ . By construction the desymmetrisation of a symmetric operad is a quasisymmetric  $n$ -operad for any  $n$ .

let  $\Pi Q_n$  be the fundamental groupoid of  $Q_n$ . A quasisymmetric operad provides, therefore, a contravariant functor on  $\Pi Q_n$ .

**Definition 4.3** A  $Q_n$ -collection is a contravariant functor on  $Q_n$ . A  $\Pi Q_n$ -collection is a contravariant functor on  $\Pi Q_n$ .

**Definition 4.4** A  $Q_n$ -operad is a  $\Pi Q_n$ -collection  $A$  together with the following structure

- for every order preserving map  $\sigma : T \rightarrow S$  the usual operadic map:

$$\mu_\sigma : A_S \otimes (A_{T_0} \otimes \dots \otimes A_{T_k}) \longrightarrow A_T.$$

This collection of maps must satisfy the usual associativity and unitarity conditions plus two equivariancy conditions:

- For every commutative diagram

$$\begin{array}{ccc} T' & \xrightarrow{\sigma'} & S' \\ \downarrow & & \downarrow \\ T & \xrightarrow{\sigma} & S \end{array}$$

where vertical maps are quasibijections and horizontal maps are order preserving the diagram

$$\begin{array}{ccc} A_S \otimes (A_{T_0} \otimes \dots \otimes A_{T_k}) & \longrightarrow & A_T \\ \downarrow & & \downarrow \\ A_{S'} \otimes (A_{T'_0} \otimes \dots \otimes A_{T'_k}) & \longrightarrow & A_{T'} \end{array}$$

commutes

- For every commutative diagram

$$\begin{array}{ccc}
T & \xrightarrow{\sigma'} & T' \\
\sigma \downarrow & & \downarrow \eta' \\
T'' & \xrightarrow{\eta} & S
\end{array}$$

where  $\sigma, \sigma'$  are quasibijections and  $\eta, \eta'$  are order preserving, the diagram

$$\begin{array}{ccc}
A_S \otimes (A_{T'_0} \otimes \dots \otimes A_{T'_k}) & \longrightarrow & A_{T'} \\
\downarrow & & \downarrow \\
A_S \otimes (A_{T_0} \otimes \dots \otimes A_{T_k}) & & A_T \\
\uparrow & & \uparrow \\
A_S \otimes (A_{T''_0} \otimes \dots \otimes A_{T''_k}) & \longrightarrow & A_{T''}
\end{array}$$

commutes.

**Theorem 4.1** *The category of  $Q_n$ -operads is equivalent to the category of quasisymmetric  $n$ -operads.*

**Proof.** Obviously, every quasisymmetric  $n$ -operad is a  $Q_n$ -operad. Let us construct an inverse functor. Given a  $Q_n$ -operad  $C$  we define a quasisymmetric operad  $A$  on an  $n$ -ordinal  $T$  to be equal to  $C_T$ . We have to define  $A$  on an arbitrary map of  $n$ -ordinals  $\sigma : T \rightarrow S$ .

let us choose a factorisation of  $\sigma$  according to Lemma 3.1.


Now we can define operadic multiplication by the following commutative diagram

$$\begin{array}{ccc}
A_S \otimes (A_{T_0} \otimes \dots \otimes A_{T_k}) & \xrightarrow{\mu_\sigma} & A_T \\
1 \otimes (\alpha_{\pi_1}^{-1} \otimes \dots \otimes \alpha_{\pi_k}^{-1}) \downarrow & & \uparrow \alpha_\pi \\
A_S \otimes (A_{T'_0} \otimes \dots \otimes A_{T'_k}) & \xrightarrow{\mu_\nu} & A_{T'}
\end{array}$$


The second equivariancy axiom implies that this definition does not depend on a chosen factorisation. Suppose now we have a composite

$$T \xrightarrow{\sigma} S \xrightarrow{\omega} R.$$

It generates the following factorization diagram



which in its turn generates the following huge diagram



In this diagram we omit the symbol  $\otimes$  to shorten the notations. Then the central region of the diagram commutes because of associativity of  $A$  with respect to order preserving maps of  $n$ -ordinals. Other regions commute either by one of equivariancy conditions either by naturality either by functoriality. The commutativity of this diagram means associativity of  $A$  with respect to composition of maps of  $n$ -ordinals. □

## 5 The category of quasibijections and configuration spaces.

It is clear that the category  $Q_n$  is the union of connected components  $Q_n(k)$  where  $k$  is the cardinality of the  $n$ -ordinals.

**Theorem 5.1** • *For a finite  $n$  the space  $N(Q_n(k))$  has homotopy type of unordered configuration spaces of  $k$ -points in  $\mathfrak{R}^n$ ;*

- *The localisation functors*

$$l_2 : Q_2 \rightarrow \Pi Q_2,$$

*induces a weak equivalence of the nerves;*

- *The groupoid  $\Pi Q_2$  is equivalent to the groupoid of braids;*
- *The localisation functors*

$$l_\infty : Q_\infty \rightarrow \Pi Q_\infty,$$

*induces a weak equivalence of the nerves;*

- *the groupoids  $\Pi Q_n$ ,  $3 \leq n \leq \infty$  are equivalent to the symmetric groups groupoid.*

**Proof.** We give a sketch of the proof. A detailed discussion can be found in [1, 3]. Consider the configuration space of ordered  $k$ -points in  $\mathfrak{R}^n$  :

$$\text{Conf}_k(\mathfrak{R}^n) = \{(x_1, \dots, x_k) \in (\mathfrak{R}^n)^k \mid x_i \neq x_j \text{ if } i \neq j\}$$

It admits a so called Fox-Neuwirth stratification.

Let  $S_+^{n-p-1}$  denote the open  $(n-p-1)$ -hemisphere in  $\mathfrak{R}^n$ ,  $0 \leq p \leq n-1$ :

$$\overset{\circ}{S}_+^{n-p-1} = \left\{ x \in \mathfrak{R}^n \mid \begin{array}{l} x_1^2 + \dots + x_n^2 = 1 \\ x_{p+1} > 0 \text{ and } x_i = 0 \text{ if } 1 \leq i \leq p \end{array} \right\}$$

Similarly,

$$\overset{\circ}{S}_-^{n-p-1} = \left\{ x \in \mathfrak{R}^n \mid \begin{array}{l} x_1^2 + \dots + x_n^2 = 1 \\ x_{p+1} < 0 \text{ and } x_i = 0 \text{ if } 1 \leq i \leq p \end{array} \right\} .$$

Let  $u_{ij} : \text{Conf}_k(\mathfrak{R}^n) \rightarrow S^{n-1}$  be a function

$$u_{ij}(x_1, \dots, x_k) = \frac{x_j - x_i}{\|x_j - x_i\|}$$

The Fox-Neuwirth cell corresponding to an  $n$ -ordinal  $T$  with  $[T] = [k-1]$  is a subspace of  $\text{Conf}_k(\mathfrak{R}^n)$

$$FN_T = \left\{ x \in \text{Conf}_k(\mathfrak{R}^n) \left| \begin{array}{l} u_{ij}(x) \in \overset{\circ}{S}_+^{n-p-1} \quad \text{if } i <_p j \text{ in } T \\ u_{ij}(x) \in \overset{\circ}{S}_-^{n-p-1} \quad \text{if } j <_p i \text{ in } T \end{array} \right. \right\} .$$

Each Fox-Neuwirth cell is an open convex subspace of  $(\mathfrak{R}^n)^k$ . We also have

$$\text{Conf}_k(\mathfrak{R}^n) = \bigcup_{[T]=[k-1], \pi \in S_k} \pi FN_T.$$

Here  $\pi FN_T$  means a space obtained from  $FN_T$  by renumbering points according to the permutation  $\pi$ .

Let  $J_n(k)$  be the Milgram poset of all possible  $n$ -ordinal structures on the set  $\{0, \dots, k-1\}$  [1]. The group  $S_k$  acts on  $J_n(k)$  and the quotient  $J_n(k)/S_k$  is isomorphic to  $Q_n(k)$ .

One can think of an element from  $J_n(k)$  as a pair  $(T, \pi)$  where  $T$  is an  $n$ -ordinal and  $\pi$  is a permutation from  $S_k$  and  $(T, \pi) > (S, \xi)$  in  $J_n(k)$  when there exists a quasibijection  $\sigma : T \rightarrow S$  and  $\xi \cdot \pi = \sigma$ .

We also can associate a convex subspace of configuration space  $FN(T, \pi) = \pi FN_T$  with every element of  $J_n(k)$ . Moreover, if  $(T, \pi) > (S, \xi)$  then  $FN(S, \xi)$  is on the boundary of the closure of  $FN(T, \pi)$ . Let us define

$$\overline{FN}(T, \pi) = \bigcup_{(S, \xi) \leq (T, \pi)} FN(S, \xi).$$

The spaces  $\overline{FN}(T, \pi)$  are contractible and, moreover, we have a functor

$$\overline{FN} : J_n^{op}(k) \rightarrow Top.$$

We then have the following zig-zag of weak equivalences

$$N(J_n^{op}(k)) \leftarrow hocolim \overline{FN} \rightarrow colim \overline{FN} \simeq \text{Conf}_k(\mathfrak{R}^n).$$

The first statement of the theorem follows then from the quotient of the zig-zag above by the action of the symmetric group. The second and the third statements are the consequences of the fact that the space  $\text{Conf}_k(\mathfrak{R}^2)$  is the  $K(Br_k, 1)$ -space. The fifth statement follows from simply connectedness of  $\text{Conf}_k(\mathfrak{R}^n)$ ,  $n > 3$ . Finally the fourth statement can be obtained using the formula  $Q_\infty = colim_n Q_n$ .

□

We need to make the equivalence of the groupoids  $\Pi Q_2$  and  $\mathcal{B}r$  more explicit.

The total order functor  $[-] : Q_2 \rightarrow \mathcal{S}$  induces by universal property a functor  $s_2 : \Pi Q_2 \rightarrow \mathcal{S}$ .

Let  $p : \mathcal{B}r \rightarrow \mathcal{S}$  be the canonical functor. The map  $p$  admits a section  $q$ , which is not a homomorphism. For  $\sigma \in S_n$  we construct a braid  $q(\sigma)$  which for  $i < j$  such that  $\sigma(i) > \sigma(j)$  has a strand from  $i$  to  $\sigma(i)$  which goes over the strand from  $j$  to  $\sigma(j)$  and there is no crossing if  $\sigma$  preserves the order of  $i$  and  $j$ .

**Lemma 5.1** • *The composite*

$$Q_2 \xrightarrow{[-]} \mathcal{S} \xrightarrow{q} \mathcal{B}r$$

*is a functor;*

- *The functor induced by universal property of  $\Pi Q_2$*

$$b : \Pi Q_2 \rightarrow \mathcal{B}r$$

*is an equivalence of groupoids.*

**Proof.** To prove that  $q[-]$  is a functor we have to prove that it preserves composition. We observe that in a composite of quasibijections of 2-ordinals  $T \xrightarrow{\sigma} S \xrightarrow{\xi} R$  if  $\sigma$  reverses the total order of two elements  $i, j \in T$  then  $\xi$  can not reverse the order of  $\sigma(i)$  and  $\sigma(j)$ . So, the resulting overcrossings in the composite  $q[\sigma]q[\xi]$  are the same as in  $q[\sigma \cdot \xi]$ .

To prove the second claim it is sufficient to check that the induced morphism of groups

$$b : \Pi Q_2(S[n-1], S[n-1]) \rightarrow \mathcal{B}r_n$$

is an isomorphism.

It is obviously an epimorphism. So we have to prove that it is also a monomorphism.

For this it will be enough to prove that if a zig-zag

$$z : S[n-1] \leftarrow T[n-1] \rightarrow S[n-1] \leftarrow \dots \rightarrow S[n-1],$$

where each arrow is given by a permutation of two consecutive elements or an identity permutation, is such that the corresponding braid  $b(z)$  is trivial then  $z$  is trivial in  $\Pi Q_2$ .

This can be done if we prove that the morphisms in  $\Pi Q_2(S[n-1], S[n-1])$


$$\bar{\sigma}_i : S[n-1] \xleftarrow{1} T[n-1] \xrightarrow{\sigma_i} S[n-1],$$

where the left arrow is given by an identity and the right arrow is given by permutation  $\sigma_i$  which change the order of  $i$  and  $i+1$ , satisfy the classical Artin's




braid relations. Then we can prove triviality of  $z$  using the same rewriting process as for  $b(z)$ .

Let  $j > i + 1$  and choose  $m, l$  such that  $[m - 1] \oplus [l - 1] = [n - 1]$  and  $i \in [m - 1], j = m + 1$ . The following commutative diagram in  $Q_2$  proves that  $\bar{\sigma}_i \bar{\sigma}_j = \bar{\sigma}_j \bar{\sigma}_i$ :



In this diagram all unnamed morphisms are identities on the underlying sets. The morphism  $(\sigma_i, \sigma_j)$  acts as  $\sigma_i$  on  $[m - 1]$  and as  $\sigma_0$  on  $[l - 1]$ .

For the proof of Yang-Baxter relations  $\bar{\sigma}_i \bar{\sigma}_{i+1} \bar{\sigma}_i = \bar{\sigma}_{i+1} \bar{\sigma}_i \bar{\sigma}_{i+1}$  we should consider the following commutative diagram in  $Q_2$  which express the morphism  $\bar{\sigma}_{i+1} \bar{\sigma}_i \bar{\sigma}_{i+1}$



Analogous diagram (the mirror image of the above diagram) can be written for  $\bar{\sigma}_i \bar{\sigma}_{i+1} \bar{\sigma}_i$ . The relation follows from it immediately.  $\square$

So, we have a commutative diagram of categories and functors

$$\begin{array}{ccccc}
Q_2 & \longrightarrow & \Pi Q_2 & \xrightleftharpoons[b]{c} & Br \\
[-] \searrow & & s_2 \downarrow & \swarrow p & \\
& & S & & 
\end{array}$$

where  $c$  is an adjoint equivalence to  $b$ . Notice that all functors in this diagram are strict monoidal functors.

**Lemma 5.2** *Let*

$$z : S \xleftarrow{\sigma} T \xrightarrow{\eta} R$$

*be a zig-zag of quasibijections of  $n$ -ordinals such that*

$$s_2(z) = \tau_1 \oplus \dots \oplus \tau_k.$$

*Then there exist braids  $b_i$ ,  $1 \leq i \leq k$  such that  $p(b_i) = \tau_i$ ,  $1 \leq i \leq k$  and*

$$b(z) = b_1 \oplus \dots \oplus b_k.$$

**Proof.** We will prove that there exist quasibijections  $\sigma_i : T_i \rightarrow S_i = S[n_i]$ ,  $\eta_i : T_i \rightarrow R_i = S[n_i]$   $1 \leq i \leq k$ , two quasibijections  $\xi : \oplus_i S_i \rightarrow S$ ,  $\zeta : \oplus_i R_i \rightarrow R$  and a quasibijection  $\kappa : \oplus_i T_i \rightarrow T$ , such that the following diagram commutes

$$\begin{array}{ccccc}
\oplus_i S_i & \xleftarrow{\oplus_i \sigma_i} & \oplus_i T_i & \xrightarrow{\oplus_i \eta_i} & \oplus_i R_i \\
\xi \downarrow & & \kappa \downarrow & & \zeta \downarrow \\
S & \xleftarrow{\sigma} & T & \xrightarrow{\eta} & R
\end{array}$$

and  $b(\xi) = b(\zeta) = \Gamma_B(\pi; 1, \dots, 1)$  for a braid  $\pi$  on  $k$  strands. Then the result will follow from an elementary observation that the braid

$$b(S) \xrightarrow{b(\xi)^{-1}} \oplus_i b(S_i) \xrightarrow{\oplus_i b(\sigma_i)^{-1}} \oplus_i b(T_i) \xrightarrow{\oplus_i b(\eta_i)} \oplus_i b(R_i) \xrightarrow{b(\xi)} b(R)$$

is equal to

$$\oplus_i b(S_i) \xrightarrow{\oplus_i b(\sigma_i^{-1})} \oplus_i b(T_i) \xrightarrow{\oplus_i b(\eta_i)} \oplus_i b(R_i).$$

It is enough to proof the lemma for  $k = 2$ . The rest will follow by induction. Also without loss of generality we can assume that  $S = S[n]$  and  $T = T[n]$ . Now,  $p(S)$  is the ordinal sum  $[l] \oplus [m]$ ,  $n = m + 1 + 1$  and the image of the restriction of the map  $\sigma^{-1}\eta$  on  $\{0, \dots, l\}$  is  $\{0, \dots, l\}$  and the image of the restriction on  $\{l + 1, \dots, m + l + 1\}$  is  $\{l + 1, \dots, m + l + 1\}$ .

We put  $S_1 = S[l]$ ,  $T_1 = T[l]$  and  $S_2 = S[m]$ ,  $T_1 = T[m]$ . We have to construct quasibijections

$$\sigma_i, \eta_i : T_i \rightarrow S_i \quad i = 1, 2,$$

and also quasibijections

$$\xi, \zeta : S_1 \oplus S_2 \rightarrow S, \kappa : T_1 \oplus T_2 \rightarrow T,$$

which make the diagram

$$\begin{array}{ccccc} S_1 \oplus S_2 & \xleftarrow{\sigma_1 \oplus \sigma_2} & T_1 \oplus T_2 & \xrightarrow{\eta_1 \oplus \eta_2} & S_1 \oplus S_2 \\ \downarrow & & \downarrow & & \downarrow \\ S & \xleftarrow{\sigma} & T & \xrightarrow{\eta} & S \end{array} \quad (2)$$

commutative.

The quasibijection  $\kappa$  is simply the identity. Let us describe  $\sigma_1$ . Let  $\sigma([l])$  be the image of the set  $\{0, \dots, l\}$  in the ordinal  $[n]$ . This image gets an induced order from  $[n]$  which makes it isomorphic to  $[l]$ . Let  $\phi_1 : \sigma([l]) \rightarrow [l]$  be this unique isomorphism. We define  $\sigma_1$  as the composite

$$[l] \rightarrow \sigma([l]) \xrightarrow{\phi_1} [l].$$

Similarly, we define  $\sigma_2$  as the composite

$$[m] \rightarrow \sigma([m]) \xrightarrow{\phi_2} [m],$$

where  $\sigma([m])$  is the image of  $\{l+1, \dots, m+l+1\}$  and we give analogous definitions for  $\eta_1$  and  $\eta_2$ .

Finally, we define  $\xi$  by the formula

$$\xi(x) = \begin{cases} \phi_1^{-1}(x) & \text{if } x \in \{0, \dots, l\} \\ \phi_2^{-1}(x) & \text{if } x \in \{l+1, \dots, m+l+1\} \end{cases}$$

We use a similar argument to define  $\zeta$ . The commutativity of the diagram (2) follows from the definition.  $\square$

## 6 Quasisymmetric $n$ -operads vs symmetric and braided operads.

**Theorem 6.1** *The category of quasisymmetric 2-operads and the category of braided operads are equivalent.*

**Proof.** We first prove that the category of quasisymmetric 2-operads is equivalent to the category whose objects are *mixed 2-operads* in the sense of the definition below and whose morphisms are multiplications and units preserving morphisms of the underlying braided collections.

**Definition 6.1** A mixed 2-operad in  $V$  is a right braided collection  $A$  equipped with the following additional structure:

- a morphism  $e : I \rightarrow A_0$
- for every order preserving map  $\sigma : [n] \rightarrow [k]$  in  $\Omega^s$  a morphism :

$$\mu_\sigma : A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) \longrightarrow A_n,$$

where  $[n_i] = \sigma^{-1}(i)$ .

They must satisfy the identities (1-3) from the definition of symmetric operad and the following two equivariance conditions:

1. For any two quasibijections of 2-ordinals  $\pi, \rho$  and two order preserving maps  $\sigma, \sigma' \in \Omega^s$  such that the following diagram commutes in  $\Omega^s$

$$\begin{array}{ccc} [T'] & \xrightarrow{\sigma'} & [S'] \\ [\pi] \downarrow & & \downarrow [\rho] \\ [T] & \xrightarrow{\sigma} & [S] \end{array}$$

the following induced diagram commutes:

$$\begin{array}{ccc} A_{k'} \otimes (A_{n'_{\rho(0)}} \otimes \dots \otimes A_{n'_{\rho(k)}}) & \xrightarrow{\mu_{\sigma'}} & A_{n'} \\ A(b(\rho)) \otimes \tau(\rho) \uparrow & & \uparrow A(b(\pi)) \\ A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) & \xrightarrow{\mu_\sigma} & A_n \end{array},$$

where  $\tau(\rho)$  is the symmetry in  $V$  which corresponds to the permutation  $[\rho]$ .

2. For any two quasibijections  $\sigma, \sigma'$  and two order preserving maps  $\eta, \eta' \in \Omega^s$  such that the following diagram commutes in  $\Omega^s$

$$\begin{array}{ccc} [T''] & \xrightarrow{[\sigma']} & [T'] \\ [\sigma] \downarrow & & \downarrow \eta' \\ [T] & \xrightarrow{\eta} & [S] \end{array}$$

the following diagram commutes

$$\begin{array}{ccc}
A_k \otimes (A_{n'_0} \otimes \dots \otimes A_{n'_k}) & \xrightarrow{\mu_{\eta'}} & A_{n'} \\
\downarrow 1 \otimes A(b(\sigma'_0)) \otimes \dots \otimes A(b(\sigma'_k)) & & \downarrow A(b(\sigma')) \\
A_k \otimes (A_{n''_0} \otimes \dots \otimes A_{n''_k}) & & A_{n''} \\
\uparrow 1 \otimes A(b(\sigma_0)) \otimes \dots \otimes A(b(\sigma_k)) & & \uparrow A(b(\sigma)) \\
A_k \otimes (A_{n_0} \otimes \dots \otimes A_{n_k}) & \xrightarrow{\mu_\eta} & A_n
\end{array}$$

For a quasisymmetric 2-operad  $A$  we define a mixed 2-operad  $B$  by pulling back along the equivalence  $c : Br \rightarrow \Pi Q_2$ . And vice versa, we produce a quasisymmetric 2-operad from a mixed 2-operad by pulling back along  $b : \Pi Q_2 \rightarrow Br$ . It is not hard to check that this indeed gives the necessary equivalence of the corresponding operadic categories.

Now, we will prove that the category of mixed 2-operads is equivalent to the category of braided operads. Let  $A$  be an operad in the sense of 6.1. We have to check that  $A$  also satisfies the Fiedorowicz's equivariance conditions. Let us start from the second condition.

For each  $\rho_i$  let us choose a zigzag of  $2L$  morphisms in  $Q_2$ , such that

$$\rho_i = b(T_i \xleftarrow{\tau_1} R_i^1 \xrightarrow{\tau_2} R_i^2 \leftarrow \dots \leftarrow R_i^{2L} \xrightarrow{\tau_{2k}} S_i).$$

Obviously, such a zig-zag exists and  $L$  can be chosen independently on  $i$ . Then the following square commutes for each odd  $j$  :

$$\begin{array}{ccc}
[n] & \xrightarrow{[\oplus_i \tau_i^{j+1}]} & [n] \\
[\oplus_i \tau_i^j] \downarrow & & \downarrow \sigma \\
[n] & \xrightarrow{\sigma} & [k]
\end{array}$$

Hence, the application of the second equivariance condition of definition 6.1  $L$  times gives the second Fiedorowicz's equivariance condition.

For the first equivariance condition we do an analogous construction by choosing a presentation of the braid  $\rho$  as an image of a zigzag.

Let  $A$  be an operad in the sense of 2.3. We construct an operad  $B$  in the sense of 6.1 as follows. As a braided collection  $B$  coincides with  $A$ . Its multiplication is the same as in  $A$  also. The only nontrivial statement to check is that  $B$  satisfies the equivariance conditions from Definition 6.1. To prove the second condition we use Lemma 5.2.

It is obvious also that the first equivariance condition is satisfied in the following special case. Let  $\sigma' : T \rightarrow S'$  be an order preserving map and let

$\rho : S' \rightarrow S$  be a quasibijection. Apply Lemma 3.1 to produce a quasibijection  $\pi(\rho, \sigma') : T' \rightarrow T$  and order preserving map  $\sigma(\rho, \sigma') : T \rightarrow S$  such that  $\sigma' \cdot \rho = \pi(\rho, \sigma') \cdot \sigma(\rho, \sigma')$ . Then  $b(\pi(\rho, \sigma')) = \Gamma_B(b(\rho); 1, \dots, 1)$  and we can apply the first equivariance Fiedorowicz's condition.

Then the first equivariance condition is satisfied in general because of the second equivariance condition of the Definition 6.1 applied to the commutative diagram

$$\begin{array}{ccc}
 [T'] & \xrightarrow{[\pi(\rho, \sigma')]} & [T] \\
 [\pi] \downarrow & & \downarrow [\sigma] \\
 [T] & \xrightarrow{[\sigma]} & [S]
 \end{array}$$

□

**Theorem 6.2** *The category of  $Q_n$ -operads  $3 \leq n \leq \infty$  and the category of symmetric operads are equivalent.*

**Proof.** The proof is a repetition of the above proof with a simplification that  $s_n : \Pi Q_n \rightarrow \mathcal{S}$  for  $3 \leq n \leq \infty$  is an equivalence.

□

## 7 Locally constant $n$ -operads.

The quasisymmetric  $n$ -operads are defined in any symmetric monoidal category  $V$ . But according to Theorems 6.1 and 6.2 they are different from symmetric operads only when  $n = 2$ . If we, however, allow  $V$  to have also symmetric monoidal model structure (which is the case for most applications) we can define a new interesting class of operads.

As we have seen before the main reason why quasisymmetric operads collapse to symmetric operads for  $n > 2$  is that configuration space  $Conf_k(\mathbb{R}^n)$  is simply connected and so localising with respect to quasibijections can only produce a groupoid equivalent to  $\mathcal{S}$ . The correct procedure, therefore, should be to take the weak  $\omega$ -groupoid  $\Pi_\infty Q_n$  and consider presheafs on it with values in  $V$  as category of collections. There are, however, considerable technical difficulties with this approach.

Fortunately, the results of Cisinski [5] show a way around this problem by considering as the category of collections the category of *locally constant* functors from  $Q_n^{\text{op}}$  to  $V$ . Pursuing this idea we give the following definition.

**Definition 7.1** *Let  $V$  be a symmetric monoidal model category. A locally constant  $n$ -operad in  $V$  is an  $n$ -operad  $A$  in  $V$  such that for every quasibijection  $\sigma : T \rightarrow S$  the morphism  $A(\sigma) : A_S \rightarrow A_T$  is a weak equivalence.*

An example of an interesting locally constant  $n$ -operad, which is not a quasisymmetric  $n$ -operad is the Getzler-Jones  $n$ -operad  $GJ^n$  constructed in [1] for

all  $1 < n < \infty$ . One can also construct an  $\infty$ -version  $GJ^\infty$  by the formula  $GJ_T^\infty = GJ_{\bar{T}}^n$ , where  $<_{-n}$  is the minimal nonempty relation in the  $\infty$ -ordinal  $T$ , the  $n$ -ordinal  $\bar{T}$  has the same underlying set as  $T$  and the relation  $<_{n-p-1}$  in  $\bar{T}$  coincides with the relation  $<_{-p}$  in  $T$ .

Let us describe the relations between different sorts of operads we deal with in this paper. We fixed our base symmetric monoidal model category  $V$ . Moreover, we will assume that  $V$  satisfies the conditions from Section 5 of [1], which means that it is transferable to the category of operads and also is a simplicial model category (see [1] for the details). We introduce the following notations:

- $SO$  is the category of symmetric operads;
- $BO$  is the category of braided operads;
- $O_n$  is the category of  $n$ -operads;
- $QO_n$  is the full subcategory of  $O_n$  of quasisymmetric  $n$ -operads;
- $LCO_n$  is the full subcategory of  $O_n$  of locally constant  $n$ -operads.

A morphism of operads (in any of the categories above) is a weak equivalence if it is a termwise weak equivalence of the collections. The homotopy category of operads is the category of operads localised with respect to the class of weak equivalences.

For  $n = 1$  the relationships between these categories are very simple. The following categories are isomorphic to the category of nonsymmetric operads

$$O_1 \simeq LCO_1 \simeq QO_1$$

and we have a classical adjunction between nonsymmetric operads and symmetric operads. All this is true on the level of homotopy categories.

For  $n = 2$  we have the following diagram of categories and right and left adjoint functors:

$$\begin{array}{ccc}
 O_2 & \begin{array}{c} \xleftarrow{Des_2} \\ \xrightarrow{Sym_2} \end{array} & SO \\
 \uparrow I_2 & \searrow L_2 & \uparrow U_2 \\
 LCO_2 & \begin{array}{c} \xleftarrow{J_2} \\ \xrightarrow{K_2} \end{array} & QO_2 \begin{array}{c} \xleftarrow{A_2} \\ \xrightarrow{B_2} \end{array} BO \\
 & & \uparrow F_2
 \end{array}$$

In this diagram the functor  $Des_2$  is right adjoint to  $Sym_2$  (see [1, 2] for the construction). The functors  $I_2$  and  $J_2$  are natural inclusions. The functor  $K_2$  is left adjoint to  $J_2$  and  $L_2$  is left adjoint to the composite  $J_2 \cdot I_2$ . Using the theory of internal operads from [2] one can show that  $L_2$  on the level of collections is given by the left Kan extension along the localisation functor  $l_2 : Q_2 \rightarrow \Pi Q_2$  :

$$L_2(A) = Lan_{l_2}(A). \quad (3)$$

We have also the same formula for  $K_2$ . The functor  $A_2$  is a right adjoint and  $B_2$  is a left adjoint part of the equivalence constructed in the section 6. Finally,  $U_2$  is the functor which produces a braided operad from symmetric operad by pulling back along the functor  $p : \mathcal{B}r \rightarrow \mathcal{S}$  and  $F_2$  is its left adjoint given by quotient with respect to pure braid groups action.

**Theorem 7.1** • *The homotopy category of locally constant 2-operads and the homotopy category of quasisymmetric 2-operads and the homotopy category of braided operads are equivalent.*

- *The functor of symmetrisation  $Sym_2$  can be factorised as  $L_2 \cdot B_2 \cdot F_2$ .*
- *A base space  $X$  is a 2-fold loop space (up to group completion) if and only if it is an algebra of a contractible 2-operad, if and only if it is an algebra of a contractible braided operad (Fiedorowicz's recognition principle [4]).*

**Proof.** Since  $\Pi Q_2$  is a groupoid, the localisation functor  $l_2$  is locally constant in the sense of [5, 1.14]. By Formal Serre spectral sequence [5, Prop. 1.15] we get that the homotopy left Kan extension along  $l_2$  is a left adjoint to the restriction functor between homotopy categories of collections. The functor  $l_2$  induces a weak equivalence of the nerves and so, by Quillen theorem B, it is also aspherical in the sense of [5, 1.4]. So, by [5, Prop. 1.16] the homotopy left Kan extension along  $l_2$  is an equivalence of homotopy categories of collections.

Taking into account the formula (3) we see that to prove the equivalence of homotopy categories of operads it is enough to show that for an  $n$ -operad  $A$  ( $1 \leq n \leq \infty$ ) there exists a cofibrant replacement  $B(A)$  such that the underlying  $Q_n$ -collection of  $B(A)$  is cofibrant in the projective model structure.

Recall [1], that  $\mathbf{ph}^n$  is the categorical symmetric operad representing the 2-functor of internal pruned  $n$ -operads. In particular an  $n$ -operad  $A$  is represented by an operadic functor  $\tilde{A} : \mathbf{ph}^n \rightarrow V^\bullet$ . If we forget about operadic structures then for any  $k \geq 0$  we will have a functor  $\tilde{A}_k : \mathbf{ph}_k^n \rightarrow V$ . Take the bar-resolution  $B(L, L, C(\tilde{A}))$ , where  $(L, \mu, \epsilon)$  is the monad on the functor category  $[d(\mathbf{ph}^n), V]$  generated by restriction and left Kan extension along the inclusion of discretisation  $d(\mathbf{ph}_k^n)$  of  $\mathbf{ph}_k^n$  to  $\mathbf{ph}_k^n$  and  $C(A)$  is the termwise cofibrant replacement of the underlying  $n$ -collection of  $A$ . These functors for all  $k \geq 0$  form an operadic functor  $\mathbf{B}(A) : \mathbf{ph}^n \rightarrow V^\bullet$  and, hence, determines an  $n$ -operad  $B(A)$  which is a cofibrant replacement for  $A$  [2].

Since,  $\mathbf{B}_k(A)$  is a bar-construction on cofibrant collection it is cofibrant in the projective model category of functors. Recall also that there is a symmetric categorical operad  $\mathbf{rh}^n$  representing the 2-functor of internal reduced  $n$ -operads [1] and a projection  $p : \mathbf{ph}^n \rightarrow \mathbf{rh}^n$ . A typical fiber (in a strict sense) of this projection over an object  $w \in \mathbf{rh}^n$  is a category with a terminal object  $s(w)$ . The map  $s$  assembles to the (nonoperadic) functor  $s : \mathbf{rh}^n \rightarrow \mathbf{ph}^n$ , which is by definition a section of  $p$  and it is also a right adjoint to  $p$ . The counit of this adjunction is identity and the unit is the unique map to the terminal object  $s(w)$ .



Now, the simple calculations with this adjunction shows that the restriction functor  $s^*$  maps preserves the cofibrant objects for projective model structures and so  $s^*(\mathbf{B}_k(A))$  is cofibrant.

There is also an inclusion  $j : J_n^{op} \rightarrow \mathbf{rh}^n$  [1]. It is not hard to see also that the categories  $J_n^{op}(k)$  and  $\mathbf{rh}_k^n$  are Reedy categories. Recall, that the objects of  $\mathbf{rh}_k^n$  are planar trees decorated by pruned  $n$ -trees (i.e.  $n$ -ordinals). One can choose the total number of edges of  $n$ -trees in a decorated planar tree as a degree function and see that each morphism decreases strictly this function.<sup>1</sup>

It follows from these considerations that the functor  $s^*(\mathbf{B}_k(A))$  satisfies the following property characterising cofibrant objects in the projective model categories for functor categories over Reedy categories:

$$\text{colim}(s^*(\mathbf{B}_k(A))(w)) \rightarrow s^*(\mathbf{B}_k(A))(T) \quad (4)$$

is a cofibration. Here colimit is taken over the category of all  $w \rightarrow T$ ,  $w \neq T$  in  $\mathbf{rh}_k^n$ . It was proved in [1] that  $J_n^{op}(k)$  is cofinal in  $\mathbf{rh}_k^n$ . Exactly the same argument show that in the colimit (4) one can replace  $w \in \mathbf{rh}_k^n$  by the objects from  $J_n^{op}(k)$ . And, therefore the restriction of  $j^*s^*(\mathbf{B}_k(A))$  is cofibrant as well.

Finally, the quotient functor  $q : J_n^{op}(k) \rightarrow Q_n^{op}(k)$  induces the restriction functor  $q^*$  on functor categories which is fully faithful. It follows from this that  $q^*$  reflects cofibrations. Finally, we observe that

$$q^*(u(B(A))) = j^*s^*(\mathbf{B}_k(A))$$

and so  $u(B(A))$  is cofibrant. Hence the first statement of the theorem is proved.

The statement about symmetrisation is obvious since  $Des_2 = U_2 \cdot A_2 \cdot J_2 \cdot I_2$ . Finally, a contractible operad is locally constant so the third statement follows from the first statement, Theorem 8.6 from [1] and the fact that the functors  $U_2, A_2, J_2, I_2$  preserve endomorphism operads.  $\square$

For  $3 \leq n \leq \infty$  the corresponding diagram is:

$$\begin{array}{ccc} O_n & \begin{array}{c} \xleftarrow{Des_n} \\ \xrightarrow{Sym_n} \end{array} & SO \\ & \searrow L_n & \uparrow A_n \\ I_n \uparrow & & \downarrow B_n \\ LCO_n & \begin{array}{c} \xrightarrow{J_n} \\ \xleftarrow{K_n} \end{array} & QO_n \end{array}$$

**Theorem 7.2** • For  $3 \leq n \leq \infty$  the category of symmetric operads is equivalent to the category of quasisymmetric  $n$ -operads;

- For  $3 \leq n < \infty$  a base space  $X$  is an  $n$ -fold loop space (up to group completion) if and only if it is an algebra of a contractible  $n$ -operad;

<sup>1</sup>In fact,  $\mathbf{rh}_k^n$  is a poset but we did not provide a proof of this fact in [1].

- The homotopy category of locally constant  $\infty$ -operads, the homotopy category of quasisymmetric  $\infty$ -operads and the homotopy category of symmetric operads are equivalent.
- A base space  $X$  is an infinite loop space (up to group completion) if and only if it is an algebra of a contractible  $\infty$ -operad if and only if it is an algebra of a contractible symmetric operad (May's recognition principle [6]).

**Proof.** The proof is analogous to the proof of Theorem 7.1. □

An interesting question which we do not consider here is the existence of model structures on the various categories of operads. The results of [5] indicate that this might be possible. But it is a subject for a future paper.

**Acknowledgements.** I would like to thank Denis-Charles Cisinski for his nice answers [5] to my some time naive questions. I wish to express my gratitude to C.Berger, I.Galvez, E.Getzler, V.Gorbunov, A.Davydov, R.Street, A.Tonks, M.Weber for many useful discussions.

I also gratefully acknowledge the financial support of Scott Russel Johnson Memorial Foundation, Max Plank Institut für Mathematik and Australian Research Council (grant No. DP0558372).

## Bibliography.

- [1] Batanin M.A., *The symmetrisation of  $n$ -operads and compactification of real configuration spaces*, Adv. Math. **211** (2007), pp. 684-725.
- [2] Batanin M.A., *The Eckmann-Hilton argument and higher operads*, Adv. Math. **217**(2008), pp. 334-385.
- [3] Berger C., *Combinatorial models for real configuration spaces and  $E_n$ -operads*, *Contemp. Math.* **202** (1997), 37-52.
- [4] Fiedorowicz Z., *The symmetric bar construction*, (1992) preprint, available from the author's home page.
- [5] Cisinski D.C., *Locally constant functors*. arXiv:0803.4342 .
- [6] May P., *The Geometry of Iterated Loop Spaces*, *Lecture Notes in Math.*, **271**, Springer Verlag, Berlin, (1972).