

**Finiteness of Mordell-Weil
groups of Kuga fiber spaces
of Abelian varieties**

Masa-Hiko Saito

Department of Mathematics
Faculty of Science
Kyoto University
Kyoto, 606

Japan

Max-Planck-Institut für Mathematik
Gottfried-Claren-Straße 26
D-5300 Bonn 3

Germany

FINITENESS OF MORDELL-WEIL GROUPS OF KUGA FIBER SPACES OF ABELIAN VARIETIES

BY
MASA-HIKO SAITO
Kyoto University

§0 INTRODUCTION.

Let $f : \mathcal{X} \rightarrow M$ be a projective abelian scheme over an arithmetic quotient of a hermitian symmetric domain $M = \Gamma \backslash \mathcal{D}$, constructed from a symplectic representation of the associated algebraic group. Such fiber spaces of abelian varieties have been studied by Kuga, Shimura, Satake, Mumford, et al. Following Satake [S1], Ch. IV, we call such a fiber space a Kuga fiber space (of abelian varieties). Let η be the generic point of M and \mathcal{X}_η denotes the generic fiber of f . Then \mathcal{X}_η can be considered as an abelian variety defined over the rational function field $K = \mathbb{C}(M)$, so define the Mordell-Weil group to be the group $\mathcal{X}_\eta(K)$ of K -rational points, or equivalently, the group of rational sections of $f : \mathcal{X} \rightarrow M$, and denote it by $MW(\mathcal{X}/M)$. In this paper, we shall study Mordell-Weil groups $MW(\mathcal{X}/W)$ of Kuga fiber spaces, and prove a finiteness theorem for them.

Historically, Shioda first showed that the Mordell-Weil groups of the elliptic modular surfaces corresponding to arithmetic subgroups $\Gamma \subset SL_2(\mathbb{Z})$ are finite in [Sd]. Generalizing Shioda's result, Silverberg [Si1] proved the finiteness of the Mordell-Weil groups of those Kuga fiber spaces which are characterized by an endomorphism algebra with positive involution and a polarization, introduced by Shimura in [Sh1] and [Sh2]. She later obtained in [Si2] a cohomological criterion for the finiteness, which covered the most of her former results.

Denote by $\mathbf{R}_1 f_* \mathbb{C}_{\mathcal{X}}$ the local system of the first homology groups of the fibers of f . Then the local system $\mathbf{R}_1 f_* \mathbb{C}_{\mathcal{X}}$ is induced by a representation $\Gamma \rightarrow GL(W_{\mathbb{C}})$, and we have natural isomorphisms $H^q(M, \mathbf{R}_1 f_* \mathbb{C}_{\mathcal{X}}) \cong H^q(\Gamma, W_{\mathbb{C}})$ where $H^q(\Gamma, W_{\mathbb{C}})$ denotes the Eilenberg-MacLane cohomology group. The criterion of Silverberg says that if $\dim M > 1$ or M is compact and $H^q(\Gamma, W_{\mathbb{C}}) = 0$ for $q = 0, 1$ then the Mordell-Weil group $MW(\mathcal{X}/M)$ is finite.

This criterion directly works for the cases when the algebraic group $G_{\mathbb{Q}}$ defined over \mathbb{Q} under consideration has rational rank ≥ 2 , or the rational rank = 0 (i.e. $\Gamma \subset G_{\mathbb{R}}$ is cocompact) and $G_{\mathbb{R}}$ has no compact factor and no factor isomorphic to $SU(n, 1)$ (see Th. 6 and Th.7 in [Si2], or [B-W]). (When the rational rank = 1, see Theorem 7 of [Si2]).

On the other hand, there are examples of Kuga fiber spaces for which one can not apply these vanishing theorem directly, and in some cocompact cases, we do have examples with

$H^1(\Gamma, \mathbf{W}_{\mathbf{C}}) \neq 0$. (See §5). But we can still expect the finiteness of the Mordell-Weil group (see [Si1], [Si3]).

As far as the classification of Kuga fiber spaces is concerned, Satake studied deeply \mathbb{Q} -symplectic representations, and classified all \mathbb{Q} -primary symplectic representations with a very mild additional condition ([S2], see also IV, §6, [S1]), and every \mathbb{Q} -symplectic representation is a sum of primary representations. They consist of the standard one which is constructed from the pair of a D -module V with a D -skew hermitian or a D -hermitian form h where D is a division algebra over \mathbb{Q} with center F_1 , and the non-standard one obtained from exterior product and spin representations. In the standard case, the \mathbb{Q} -algebraic group is given by $R_{F_1/\mathbb{Q}}(SU(V, h))$, which is obtained from the F_1 -algebraic group $SU(V, h)$ by Weil's restriction of the scalars. We remark that the the standard representations include the cases which were studied by Shimura in [Sh3].

In this paper, we will only consider the standard \mathbb{Q} -symplectic representation. Also, we will exclude the following case from our consideration (cf. (3.42)):

$$(0.1) \quad \text{Case (R2, -1), } n = 2: G_{\mathbb{R}} \cong SU_2(\mathbb{H})^- \times \cdots \times SU_2(\mathbb{H})^- \times SO_4(\mathbb{R}) \times \cdots \times SO_4(\mathbb{R}),$$

because the reducibility of $SU_2(\mathbb{H})^-$ forces annoying distinctions about the nature of Γ . (For the notation, see §3, (3.23) and (3.31)).

Then the main theorem in this paper can be stated as follows.

(0.2) Theorem. ((4.23), (5.8) and (6.25)). *Let $f : \mathcal{X} \rightarrow M$ be a Kuga fiber space associated to a standard \mathbb{Q} -primary representation not isomorphic to the case (0.1). Assume that $\dim M \geq 1$. Then the Mordell-Weil group $MW(\mathcal{X}/M)$ is finite.*

The main idea of our proof is a generalization of Silverberg's method in [Si2] by introducing the L^2 -cohomology and the Hodge theory, which can be outlined as follows.

If the codimension of the singular locus of the Satake compactification M^* of M is greater than 1, then for $q \leq 1$, $H^q(\Gamma, \mathbf{W}_{\mathbf{C}}) \cong H^q(M, \mathbf{W}_{\mathbf{C}})$ is isomorphic to the middle perversity intersection cohomology $\mathbf{IH}^q(M^*, \mathbf{W}_{\mathbf{C}})$. Then by the Zucker conjecture proved in [L] and [Sa-St], these are also isomorphic to L^2 -cohomology groups. By Borel-Casselman [B-C], the L^2 -cohomology is calculated by (\mathfrak{g}, K) -cohomology, and hence we can apply the Borel-Wallach vanishing theorem in [B-W] even in the case when Γ is not cocompact, and deduce that $H_{(2)}^q(M, \mathbf{W}_{\mathbf{C}}) = 0$ if $q < \text{rk}_{\mathbb{R}} G_{\mathbb{R}}$. So if $\text{rk}_{\mathbb{R}} G_{\mathbb{R}} \geq 2$, we always have $H^q(M, \mathbf{W}_{\mathbf{C}}) = 0$ for $q = 0, 1$. In case when $\text{rk}_{\mathbb{R}} G_{\mathbb{R}} = 1$, we will separate the proof into two cases, that is, the cases where $M = \Gamma \backslash \mathcal{D}$ is compact or non-compact.

If M is compact, we can use Deligne-Zucker Hodge theory on $H^q(M, \mathbf{W}_{\mathbf{C}})$, because $\mathbf{W}_{\mathbf{C}}$ admits a variation of polarized Hodge structure. It is proved that the Mordell-Weil group $MW(\mathcal{X}/M)$ is isomorphic to $H^1(M, \mathbf{W}_{\mathbf{Z}}) \cap (H^{0,0})$ in this case. Since $\mathbf{W}_{\mathbb{Q}}$ has a structure of a local system of F_1 -vector spaces, we have a decomposition of $\mathbf{W}_{\mathbf{C}}$ according to the distinct embeddings of F_1 into \mathbb{C} . We can see from Satake's classification that this decomposition is compatible with the Hodge structure. Though in this case it is possible that $H^1(M, \mathbf{W}_{\mathbf{C}})^{0,0} \neq 0$, we can use the decomposition of $H^1(M, \mathbf{W}_{\mathbf{C}})$ to conclude that $H^1(M, \mathbf{W}_{\mathbb{Q}})^{0,0} = 0$.

If M is not compact and $rk_{\mathbb{R}} G_{\mathbb{R}} = 1$, we can take a smooth toroidal compactification $j : M \hookrightarrow \overline{M}$ such that $D = \overline{M} - M$ is a smooth divisor and consider the cohomology group $H^1(\overline{M}, j_* W_{\mathbb{Z}})$. Then by a result due to Cattani-Kaplan-Schmid [C-K-S] and Kashiwara-Kawai [K-K], this admits polarized Hodge structure of weight 0. On the other hand, we can extend the Kuga fiber space $f : \mathcal{X} \rightarrow M$ to a semi-abelian scheme $\overline{f} : \overline{\mathcal{X}} \rightarrow \overline{M}$. And in this case one can prove that $H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{an}(\overline{\mathcal{X}})) = H^1(\overline{M}, j_* W_{\mathbb{Z}})^{0,0}$, where $H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{an}(\overline{\mathcal{X}}))$ denote the group of holomorphic sections of \overline{f} . By using the theory of Néron model, it can be shown that there is an injective homomorphism $r : H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{an}(\overline{\mathcal{X}})) \hookrightarrow MW(\mathcal{X}/M)$ with finite cokernel. Now by using the description of Hodge structure due to Yuji Shimizu [ShzY], we calculate the Hodge component and we can finally prove that $H^1(\overline{M}, j_* W_{\mathbb{Q}})^{0,0} = 0$.

The organization of this paper as follows. In §1, we introduce \mathbb{Q} -symplectic representations and Kuga fiber spaces. In §2, we introduce the Mordell-Weil groups of Kuga fiber spaces and recall some results due to Silverberg [Si1], [Si2]. We also review a Hodge theory of the cohomology group to give a slight refinement of Silverberg's results. In §3, we summarize the basic fact on Satake's classification of \mathbb{Q} -symplectic representations. In §4, we recall some results from Borel-Casselman [B-C] and Borel-Wallach [B-W], and prove the desired vanishing theorem when the \mathbb{R} -rank of $G_{\mathbb{R}} \geq 2$, even if $G_{\mathbb{R}}$ has compact factors. In §5, we shall deal with the case when the \mathbb{R} -rank of $G_{\mathbb{R}}$ is 1 and $M = \Gamma \backslash \mathcal{D}$ is compact. We will check that the decomposition (see (5.10)) is compatible with the Hodge structure, and we calculate the first Gauss-Manin complex whose H^1 is the space of $(0, 0)$ -elements. In §6, we shall deal with the case when the \mathbb{R} -rank of $G_{\mathbb{R}}$ is 1 and M is non-compact.

The author would like to thank Professor Steven Zucker for very useful discussions about Hodge theory, L^2 -cohomology and intersection cohomology. He would also like to thank Professor Alice Silverberg for reading the preliminary version of this paper and giving useful comments. He would like to express his gratitude to JAMI in Johns Hopkins University for its hospitality during academic year 1990/91.

After I have finished the preliminary version of this paper, the author was informed that Ngaiming Mok obtained the similar result to one in this paper independently. It is announced in his preprint "Aspect of Kähler geometry on Arithmetic Varieties". The author believes that the method in this paper is different from his and it is worth while publishing this paper.

NOTATION. Let T be a complex vector space. For a complex endomorphism I and $\alpha \in \mathbb{C}$, we set $T(\alpha, I) = \{u \in T | I(u) = \alpha \cdot u\}$, the eigen space of I . We denote by $\mathbb{H} = \mathbb{R} + \mathbb{R} \cdot i + \mathbb{R} \cdot j + \mathbb{R} \cdot k$ the field of Hamilton quaternions.

§1 \mathbb{Q} -SYMPLECTIC REPRESENTATIONS AND KUGA FIBER SPACES.

Let $G_{\mathbb{Q}}$ be a \mathbb{Q} -algebraic group such that its \mathbb{R} -valued point $G_{\mathbb{R}}$ is a Zariski connected semisimple \mathbb{R} -group of hermitian type. Let K be a maximal compact subgroup of $G_{\mathbb{R}}$ and $\mathcal{D} = G_{\mathbb{R}}/K$ the corresponding Hermitian bounded symmetric space. We denote by $\mathfrak{g}, \mathfrak{k}$ Lie algebras of $G_{\mathbb{R}}$ and K respectively, and by \mathfrak{p} the orthogonal complement of \mathfrak{k} in \mathfrak{g} with respect to the Killing form. Then the complex structure of \mathcal{D} is induced by an element $H_0 \in \text{Cent}(\mathfrak{k})$ such that $(\text{ad}_{\mathfrak{p}}(H_0))^2 = -1_{\mathfrak{p}}$. A pair $(G_{\mathbb{Q}}, H_0)$ consisting of the above $G_{\mathbb{Q}}$ and H_0 is called a \mathbb{Q} -hermitian pair.

(1.1) Definition. A \mathbb{Q} -symplectic representation of a \mathbb{Q} -hermitian pair $(G_{\mathbb{Q}}, H_0)$ is a quadruples $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ consisting of

- (i) a \mathbb{Q} -vector space $W_{\mathbb{Q}}$ of dimension n ,
- (ii) a non-degenerate symplectic bilinear form $A_{\mathbb{Q}}$ on $W_{\mathbb{Q}} \times W_{\mathbb{Q}}$,
- (iii) a faithful representation $\rho_{\mathbb{Q}} : G_{\mathbb{Q}} \rightarrow Sp(W_{\mathbb{Q}}, A_{\mathbb{Q}})$ and
- (iv) a complex structure $I \in \mathcal{D}(W_{\mathbb{R}}, A_{\mathbb{R}})$ satisfying the condition

$$(1.2) \quad [d\rho_{\mathbb{R}}(H_0) - (1/2)I, d\rho_{\mathbb{R}}(X)] = 0 \quad \text{for all } X \in \mathfrak{g}_{\mathbb{R}},$$

where $\mathcal{D}(W_{\mathbb{R}}, A_{\mathbb{R}})$ denotes

$$(1.3) \quad \{I \in \text{End}(W_{\mathbb{R}}) \mid I^2 = -1_{W_{\mathbb{R}}}, A_{\mathbb{R}}(x, Iy) \text{ is a positive-definite } \mathbb{R}\text{-symmetric form}\}.$$

(See (3.11)).

Next we introduce a *Kuga fiber space of abelian varieties* induced from a \mathbb{Q} -symplectic representation. Let $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ be a \mathbb{Q} -symplectic representation of a \mathbb{Q} -hermitian pair $(G_{\mathbb{Q}}, H_0)$. By a *lattice* in $W_{\mathbb{Q}}$, we mean a free \mathbb{Z} -submodule $W_{\mathbb{Z}}$ in $W_{\mathbb{Q}}$ such that $W_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Q} \cong W_{\mathbb{Q}}$. Considering $G_{\mathbb{Q}}$ as a subgroup in $GL(W_{\mathbb{Q}})$ through the representation $\rho_{\mathbb{Q}} : G_{\mathbb{Q}} \rightarrow Sp(W_{\mathbb{Q}}, A_{\mathbb{Q}})$, for each lattice $W_{\mathbb{Z}}$ in $W_{\mathbb{Q}}$, we set

$$(1.4) \quad G_{W_{\mathbb{Z}}} = \{g \in G_{\mathbb{Q}} \mid gW_{\mathbb{Z}} = W_{\mathbb{Z}}\}.$$

Then $G_{W_{\mathbb{Z}}} \subset G_{\mathbb{Q}}$ becomes a discrete subgroup of $G_{\mathbb{R}}$.

(1.5) Definition-Proposition. ([S1], Ch. IV, /S 7). A discrete subgroup Γ of $G_{\mathbb{R}}$ commensurable to $G_{W_{\mathbb{Z}}}$ for some lattice $W_{\mathbb{Z}}$ is called an arithmetic subgroup of $G_{\mathbb{R}}$. The quotient space $\Gamma \backslash G_{\mathbb{R}}$ is of finite measure with respect the measure induced from the Haar measure of $G_{\mathbb{R}}$, and there always exists a normal subgroup Γ' of Γ of finite index such that Γ' is torsion-free.

(1.6) Definition. A 5-tuple $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I, W_{\mathbb{Z}})$ is said to be a *Kuga 5-tuple* if $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ is a \mathbb{Q} -symplectic representation of a \mathbb{Q} -hermitian pair $(G_{\mathbb{Q}}, H_0)$ and $W_{\mathbb{Z}}$ is lattice of $W_{\mathbb{Q}}$ such that

$$(1.7) \quad A_{\mathbb{Q}}(W_{\mathbb{Z}}, W_{\mathbb{Z}}) \subset \mathbb{Z}.$$

From a Kuga 5-tuple, we obtain a fiber space of abelian varieties as follows. Let K be the maximal compact subgroup of $G_{\mathbb{R}}$ determined by H_0 , and denote by $\mathcal{D} = G_{\mathbb{R}}/K$ the corresponding hermitian symmetric space. Set $W_{\mathbb{R}} = W_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{R}$, $W_{\mathbb{C}} = W_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C}$. We have a complex structure $I_0 \in \mathcal{D}(W_{\mathbb{R}}, A_{\mathbb{R}})$ (cf. (1.3)) satisfying (1.2). For an element $g \in G_{\mathbb{R}}$, define

$$I_g = \rho^{-1}(g) \cdot I \cdot \rho(g).$$

Then, by definition, we have $I_g \in \mathcal{D}(W_{\mathbb{R}}, A_{\mathbb{R}})$, and from (1.2), $I_g = I_0$ for $g \in K$. Hence we define, for each point $z = [g] \in \mathcal{D} = G_{\mathbb{R}}/K$,

$$I_z = I_g \in \mathcal{D}(W_{\mathbb{R}}, A_{\mathbb{R}}).$$

Setting $W_z^+ = \{u \in W_{\mathbb{C}} | I_z u = \sqrt{-1}u\}$, we can obtain a holomorphic vector bundle $\tilde{\mathcal{F}}^0 = \cup_{z \in \mathcal{D}} W_z^+$ over \mathcal{D} such that the following diagram commutes.

$$(1.8) \quad \begin{array}{ccc} \mathcal{D} \times W_{\mathbb{C}} & \hookrightarrow & \tilde{\mathcal{F}}^0 \\ \downarrow & & \swarrow \\ \mathcal{D} & & \end{array}$$

Let Γ be a torsion-free arithmetic subgroup of $G_{\mathbb{R}}$ such that $\Gamma \subset G_{W_{\mathbb{Z}}}$. Then the quotient space $M = \Gamma \backslash \mathcal{D}$ is a complex manifold, which is known to be a quasi-projective variety ([Ba-B]). Denote by $\mathbf{W}_{\mathbb{Z}}$ the local system of free \mathbb{Z} -modules on M induced by the flat bundle $(\mathcal{D} \times W_{\mathbb{Z}} / \sim)$, where \sim denotes the equivalence relation given by

$$(1.9) \quad (z, w) \sim (\gamma \cdot z, \rho(\gamma) \cdot w) \quad \text{for } \gamma \in \Gamma.$$

We also denote by $\mathbf{W}_{\mathbb{Q}}, \mathbf{W}_{\mathbb{R}}, \mathbf{W}_{\mathbb{C}}$ the local systems on M corresponding to $W_{\mathbb{Q}}, W_{\mathbb{R}}, W_{\mathbb{C}}$ respectively. The $G_{\mathbb{Q}}$ -invariant form $A_{\mathbb{Q}}$ induces a flat symplectic bilinear form \mathbf{A} on $\mathbf{W}_{\mathbb{Q}}$. A holomorphic vector bundle $\tilde{\mathcal{F}}^0$ on \mathcal{D} descends to M and we denote by \mathcal{F}^0 the corresponding locally free sheaf on M . Now we have the following

(1.10) Definiton-Proposition. The triple $(\mathbf{W}_{\mathbb{Z}}, \mathbf{A}, \mathcal{F}^0)$ constructed above becomes a *variation of polarized Hodge structure (VPHS, for short) of weight -1, and of types (-1,0), (0,-1) over $M = \Gamma \backslash \mathcal{D}$, i.e.,*

- (i) \mathbf{A} is a flat \mathbb{Z} -valued non-degenerate symplectic form on $\mathbf{W}_{\mathbb{Z}}$,
- (ii) $\mathcal{F}^0 \subset \mathbf{W}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathcal{O}_M$ defines a Hodge filtration of weight -1, and of types (-1,0), (0,-1), i.e.

$$0 = \mathcal{F}^1 \subset \mathcal{F}^0 \subset \mathcal{F}^{-1} = \mathbf{W}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathcal{O}_M,$$

such that

$$\mathcal{F}^0 \oplus \overline{\mathcal{F}^0} \cong \mathbf{W}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathcal{O}_M.$$

- (iii) \mathbf{A} satisfies the Hodge-Riemann bilinear relations, i.e. for a non-zero local section $u \in \mathcal{F}^0$, we have

$$\begin{aligned} \mathbf{A}(u, u) &= 0, \\ -(\sqrt{-1})\mathbf{A}(u, \bar{u}) &> 0. \end{aligned}$$

As explained in [D2] (4.4.3), we have an equivalence between the category of polarized abelian schemes over M and the category of variations of polarized Hodge structure over M of weight -1, and of type (-1,0), (0,-1), so we obtain a fiber space $f : \mathcal{X} \rightarrow M$ of abelian varieties over M .

(1.11) Definition-Proposition. (Ch. IV, §8, [S1] or 3.10, [Sh2]) A fiber space of abelian varieties $f : \mathcal{X} \rightarrow M = \Gamma \backslash \mathcal{D}$ obtained from a Kuga 5-tuple $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I, W_{\mathbb{Z}})$ and a torsion-free arithmetic subgroup $\Gamma \subset G_{W_{\mathbb{Z}}}$ of $G_{\mathbb{R}}$ is called a Kuga fiber space (of abelian varieties). The total space \mathcal{X} is a smooth quasi-projective variety and f is a smooth projective morphism.

§2 A CRITERION OF SILVERBERG AND A GENERALIZATION.

In this section, we review a criterion of the finiteness of Mordell-Weil group of Kuga fiber spaces due to Silverberg [Si2], and give a slight generalization.

First of all, we introduce the Mordell-Weil group of a fiber space of abelian varieties. Let M be a connected smooth quasi-projective variety. By a fiber space of abelian varieties over M we mean a polarized smooth abelian scheme $f : \mathcal{X} \rightarrow M$. Consider the generic fiber \mathcal{X}_{η} of f . Then \mathcal{X}_{η} is considered as an abelian variety over the field $K = \mathbb{C}(M)$ of the rational functions on M . Then the Mordell-Weil group of f is defined to be the group of K -rational points $\mathcal{X}_{\eta}(K)$, and is denoted by $MW(\mathcal{X}/M)$. There exists a natural isomorphism

$$(2.1) \quad MW(\mathcal{X}/M) = \{\text{a rational section } s : M \cdots \rightarrow \mathcal{X} \text{ of } f\}.$$

Now let $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I, W_{\mathbb{Z}})$ be a Kuga 5-tuple for a \mathbb{Q} -hermitian pair $(G_{\mathbb{Q}}, H_0)$, $\Gamma \subset G_{W_{\mathbb{Z}}}$, $M = \Gamma \backslash \mathcal{D}$ as in §1, and $f : \mathcal{X} \rightarrow M$ the associated Kuga fiber space (see (1.11)). Let $\mathcal{O}_M(\mathcal{X})$ (resp. $\mathcal{O}_M^{an}(\mathcal{X})$) denote the sheaf of germs of regular algebraic (resp. holomorphic) sections with values in \mathcal{X} (resp. \mathcal{X}^{an}). The cohomology group $H^0(M, \mathcal{O}_M(\mathcal{X}))$ is isomorphic to the group of regular algebraic sections of f . A rational section $s \in MW(\mathcal{X}/M)$ always extends to a regular algebraic section in case of a Kuga fiber space (see Prop. 2.1 [Si1]). So we have

(2.2) Proposition. For a Kuga fiber space $f : \mathcal{X} \rightarrow M$, we have an isomorphism

$$(2.3) \quad MD(\mathcal{X}/M) \cong H^0(M, \mathcal{O}_M(\mathcal{X})).$$

(2.4) REMARK: From the construction, there exists a natural map

$$(2.5) \quad H^0(M, \mathcal{O}_M(\mathcal{X})) \rightarrow H^0(M, \mathcal{O}_M^{an}(\mathcal{X})).$$

In general, there exists a holomorphic section of f which is not algebraic, (e.g. consider the case where M is a non-compact curve.) Assume that Γ is irreducible in $G_{\mathbb{R}}$ (see (4.4)). Then, if either $\dim(\mathcal{D}) > 1$, or M is compact, one can show that (2.5) must be an isomorphism (see [Si2, §1], [Ba-B, §10]).

For $\mathbb{K} = \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$, let $H^*(\Gamma, W_{\mathbb{K}})$ denote the Eilenberg-MacLane cohomology groups induced by the representation $\rho_{\mathbb{Q}}$ and an arithmetic group Γ . Since \mathcal{D} is contractible, we have natural isomorphisms for $\mathbb{K} = \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$

$$(2.6) \quad H^*(\Gamma, W_{\mathbb{K}}) \cong H^*(M, \mathbf{W}_{\mathbb{K}})$$

where $\mathbf{W}_{\mathbb{K}}$ denote the local system on M associated to $W_{\mathbb{K}}$, (see (1.9)).

Now we can state the Silverberg's criterion of the finiteness of $MD(\mathcal{X}/M)$ ([Si2], Theorem 5).

(2.7) Theorem. Assume that Γ is irreducible (cf. (4.4)) and $\dim \mathcal{D} > 1$ or $M = \mathcal{D}/\Gamma$ is compact. If

$$(2.8) \quad H^0(\Gamma, W_{\mathbf{C}}) = H^1(\Gamma, W_{\mathbf{C}}) = 0,$$

the Mordell-Weil group $MW(\mathcal{X}/M)$ is finite, and isomorphic to $H^1(\Gamma, W_{\mathbf{Z}}) \cong H^1(M, W_{\mathbf{Z}})$.

(2.9) L^2 -cohomology.

Let $f : \mathcal{X} \rightarrow M = \Gamma \backslash \mathcal{D}$ be a Kuga fiber space as above, and $(W_{\mathbf{Z}}, \mathbf{A}, \mathcal{F}^0)$ the corresponding VPHS of type $(0, -1), (-1, 0)$ as in (1.10).

The local system $W_{\mathbf{C}} = W_{\mathbf{Z}} \otimes_{\mathbf{Z}} \mathbf{C}$ has a flat symmetric bilinear form $\mathbf{A}_{\mathbf{C}}$, and if we denote by C_z the Weil operator, (or the complex structure) of a fiber $W_{\mathbf{C}}$, the form $T_z(x, y) := \mathbf{A}_{\mathbf{C}}(x, C_z \bar{y})$ becomes a positive-definite hermitian form, so it induces a metric on $W_{\mathbf{C}}$. From the construction of $W_{\mathbf{C}}$ and $\mathbf{A}_{\mathbf{C}}$, this metric is nothing but the one induced by the admissible inner product on $W_{\mathbf{C}}$ ([M-M], p375). The base space $M = \Gamma \backslash \mathcal{D}$ is endowed with a complete metric induced by the Bergman metric on \mathcal{D} . Hence, we can give a norm on each term of the complex $A(M, W_{\mathbf{C}})^{\infty}$ of $W_{\mathbf{C}}$ -valued C^{∞} exterior forms on M . Let $L_{(2)}(M, W_{\mathbf{C}})^{\infty}$ denote its subcomplex consisting of square-integrable elements whose exterior derivative are also square-integrable. We define the L^2 -cohomology group for $W_{\mathbf{C}}$ by

$$(2.10) \quad H_{(2)}(M, W_{\mathbf{C}}) := H(L_{(2)}(M, W_{\mathbf{C}})^{\infty}).$$

Let M^* denote the Baily-Borel, Satake compactification of M . It is known that M^* is a normal projective variety which has a stratification by complex subvarieties. Following [G-M], we can define the *middle perversity intersection cohomology group* $\mathbf{IH}(M^*, W_{\mathbf{C}})$. The following theorem is a direct consequence of the result, which was known as the Zucker conjecture, proved by Looijenga [L] and Saper-Stern [Sa-St].

(2.11) Theorem. Under the notation and assumption as above, we have isomorphisms

$$H_{(2)}(M, W_{\mathbf{C}}) \cong \mathbf{IH}(M^*, W_{\mathbf{C}}).$$

(2.12) Corollary. If $\text{codim}_{\mathbf{C}}(M^* - M) = i$ in M^* , then we have isomorphisms

$$H_{(2)}^q(M, W_{\mathbf{C}}) \cong H^q(M, W_{\mathbf{C}}) \quad \text{for } q < i.$$

PROOF: From the definition of the intersection cohomology group [G-M, §3, 3.1], one can easily deduce that

$$\mathbf{IH}^q(M, W_{\mathbf{C}}) \cong H^q(M, W_{\mathbf{C}}) \quad \text{for } q < i,$$

hence (2.11) implies the assertion.

If Γ is irreducible in $G_{\mathbb{R}}$ (see (4.4)) and $\dim \mathcal{D} > 1$, one has $\text{codim}_{\mathbf{C}}(M^* - M) \geq 2$ in M^* . Hence, thanks to (2.12), we have the following

(2.13) Corollary. Assume that $\Gamma \subset G_{\mathbb{R}}$ is irreducible and $\dim \mathcal{D} > 1$, or $M = \Gamma \backslash \mathcal{D}$ is compact. Then there exist isomorphisms

$$(2.14) \quad H_{(2)}^q(M, \mathbf{W}_{\mathbf{C}}) \cong H^q(M, \mathbf{W}_{\mathbf{C}}), \quad \text{for } q \leq 1.$$

(2.15) Hodge theory in case M is compact.

We recall that the triple $(\mathbf{W}_{\mathbf{Z}}, \mathbf{A}, \mathcal{F}^0)$ constructed in §1 is a VPHS of weight (-1) of types $(0,-1), (-1,0)$ (see (1.10)). In particular, the sheaf $\mathbf{W}_{\mathcal{O}} := \mathbf{W}_{\mathbf{Z}} \otimes_{\mathbf{Z}} \mathcal{O}_M$ has a Hodge filtration

$$0 = \mathcal{F}^1 \subset \mathcal{F}^0 \subset \mathcal{F}^{-1} = \mathbf{W}_{\mathcal{O}}.$$

Assume now that $M = \Gamma \backslash \mathcal{D}$ is compact. Then we have an isomorphism

$$(2.16) \quad H^n(M, \mathbf{W}_{\mathbf{C}}) \cong H_{(2)}^n(M, \mathbf{W}_{\mathbf{C}}) \quad \text{for all } n.$$

In this case, from the L^2 -harmonic theory, the right hand side of (2.16) can be expressed as a space of $\mathbf{W}_{\mathbf{C}}$ -valued L^2 -harmonic forms. Deligne showed that, as in the classical Hodge theory, there exists a decomposition

$$(2.17) \quad H^n(M, \mathbf{W}_{\mathbf{C}}) \cong H_{(2)}^n(M, \mathbf{W}_{\mathbf{C}}) = \bigoplus_{p+q=n-1} H^{p,q}$$

such that $\overline{H^{p,q}} \cong H^{q,p}$ (see [Z1]). Moreover the associated Hodge filtration on $H^n(M, \mathbf{W}_{\mathbf{C}})$ is given as follows. Let $\Omega_M(\mathbf{W}_{\mathbf{C}})$ denote the holomorphic deRham complex with values in $\mathbf{W}_{\mathbf{C}}$, with differential ∂_M . If we define the filtration $(F^r \Omega_M(\mathbf{W}_{\mathbf{C}}))$ by

$$F^r \Omega_M^p(\mathbf{W}_{\mathbf{C}}) = \Omega_M^p \otimes \mathcal{F}^{r-p},$$

Griffiths' transversality (see e.g. [Z1]) implies that they actually become subcomplexes of $\Omega_M(\mathbf{W}_{\mathbf{C}})$. The holomorphic Poincaré lemma implies that

$$H^*(M, \mathbf{W}_{\mathbf{C}}) \cong H^*(\Omega_M(\mathbf{W}_{\mathbf{C}})),$$

and the above filtration induces a filtration on the cohomology.

(2.18) Theorem. Under the above notation, we have the following.

(i) The spectral sequence

$$(2.19) \quad E_1^{p,q} = H^{p+q}(M, Gr_F^p \Omega_M(\mathbf{W}_{\mathbf{C}})) \implies H^{p+q}(M, \mathbf{W}_{\mathbf{C}})$$

degenerates at E_1 .

(ii) The filtration induced by $\{F^p \Omega_M(\mathbf{W}_{\mathbf{C}})\}$ on $H^n(M, \mathbf{W}_{\mathbf{C}})$ coincides with the Hodge filtration induced from the decomposition (2.17).

(iii) There is a natural identification

$$H^{p,q} \cong H^n(M, Gr_F^p \Omega_M(\mathbf{W}_{\mathbf{C}}))$$

for $p + q = n - 1$.

(iv) The cohomology group $H^n(M, \mathbf{W}_{\mathbf{Z}})/\text{torsion}$ is a \mathbf{Z} -structure of $H^n(M, \mathbf{W}_{\mathbf{C}})$, and has a natural polarization \mathbf{B} , i.e. a \mathbf{Z} -valued bilinear form satisfying the Hodge-Riemann bilinear relations.

For example, $H^0(M, \mathbf{W}_{\mathbf{C}})$ has a 2-step filtration $0 = F^1 \subset F^0 \subset F^{-1}$ whose successive quotients are:

$$\begin{aligned} H^{0,-1} &= Gr_{F^0}^0 = F^0 = \mathbf{H}^0(\mathcal{F}^0 \longrightarrow \Omega_M \otimes Gr_{\mathcal{F}^{-1}}^{-1}), \\ H^{-1,0} &= Gr_{F^1}^{-1} = F^{-1}/F^0 = \mathbf{H}^0(Gr_{\mathcal{F}^{-1}}^{-1}). \end{aligned}$$

where $Gr_{\mathcal{F}^{-1}}^{-1} = \mathcal{F}^{-1}/\mathcal{F}^0$. $H^1(M, \mathbf{W}_{\mathbf{C}})$ has a 3-step filtration $0 = F^2 \subset F^1 \subset F^0 \subset F^{-1} = H^1$ whose successive quotients are:

$$(2.20) \quad H^{1,-1} = Gr_{F^1}^1 = F^1 = \mathbf{H}^1(0 \longrightarrow \Omega_M^1 \otimes \mathcal{F}^0 \longrightarrow \Omega_M^2 \otimes Gr_{\mathcal{F}^{-1}}^{-1}),$$

$$(2.21) \quad H^{0,0} = Gr_{F^0}^0 = F^0/F^1 = \mathbf{H}^1(\mathcal{F}^0 \longrightarrow \Omega_M^1 \otimes Gr_{\mathcal{F}^{-1}}^{-1}),$$

$$(2.22) \quad H^{-1,1} = Gr_{F^2}^{-1} = F^{-1}/F^0 = \mathbf{H}^1(Gr_{\mathcal{F}^{-1}}^{-1}).$$

Considering $H^1(M, \mathbf{W}_{\mathbf{Q}})$ as a lattice of $H^1(M, \mathbf{W}_{\mathbf{C}})$, we set

$$(2.23) \quad H^1(M, \mathbf{W}_{\mathbf{Q}})^{0,0} = H^1(M, \mathbf{W}_{\mathbf{Q}}) \cap H^{0,0}.$$

Let $p_n : H^n(M, \mathbf{W}_{\mathbf{C}}) \longrightarrow H^{-1,n} = H^n(M, Gr_{\mathcal{F}^{-1}}^{-1})$ be the natural projection map induced by the spectral sequence (2.19). Set also

$$(2.24) \quad X_{const} = \text{coker}\{p_0 : H^0(M, \mathbf{W}_{\mathbf{Z}}) \rightarrow H^0(Gr_{\mathcal{F}^{-1}}^{-1})\},$$

$$(2.25) \quad H^1(M, \mathbf{W}_{\mathbf{Z}})^{0,0} = \ker\{p_1 : H^1(M, \mathbf{W}_{\mathbf{Z}}) \rightarrow H^1(M, Gr_{\mathcal{F}^{-1}}^{-1})\}.$$

Then by Hodge theory (2.18), one has

$$(2.26) \quad H^1(M, \mathbf{W}_{\mathbf{Q}})^{0,0} = H^1(M, \mathbf{W}_{\mathbf{Z}})^{0,0} \otimes_{\mathbf{Z}} \mathbf{Q}.$$

Under these notations, we can state the following theorem which gives a very natural description of $MW(\mathcal{X}/M)$. (Cf. [Z1], Cor. 10.2)

(2.27) Theorem. Assume that $M = \Gamma \backslash \mathcal{D}$ is compact. Then

(i) X_{const} in (2.24) is an abelian variety over \mathbf{C} , and

(ii) we have a natural exact sequence of abelian group

$$(2.28) \quad 0 \longrightarrow X_{const} \longrightarrow MW(\mathcal{X}/M) \longrightarrow H^1(M, \mathbf{W}_{\mathbf{Z}})^{0,0} \longrightarrow 0.$$

PROOF: The assertion (i) is an immediate consequence of (2.18). Since M is projective, all holomorphic sections become algebraic, so by (2.5), we have an isomorphism $MW(\mathcal{X}/M) \cong H^0(M, \mathcal{O}_M^{an}(\mathcal{X}))$. The relative exponential map for an abelian scheme $f : \mathcal{X} \longrightarrow M$ yields the following exact sequence of sheaf on M^{an}

$$(*) \quad 0 \longrightarrow \mathbf{R}_1 f_* \mathbf{Z} \longrightarrow Lie(\mathcal{X}) \longrightarrow \mathcal{O}_M^{an}(\mathcal{X}) \longrightarrow 0,$$

where $\mathbf{R}_1 f_* \mathbb{Z}$ denote the local system of the first homology of fibers of f . From the construction of a Kuga fiber space, we have isomorphisms $\mathbf{W}_{\mathbb{Z}} \cong \mathbf{R}_1 f_* \mathbb{Z}$ and $Lie(\mathcal{X}) \cong Gr_{\mathcal{F}}^{-1}$, hence (*) can be written as

$$(2.29) \quad 0 \longrightarrow \mathbf{W}_{\mathbb{Z}} \longrightarrow Gr_{\mathcal{F}}^{-1} \longrightarrow \mathcal{O}_M^{an}(\mathcal{X}) \longrightarrow 0.$$

This yields an exact sequence of cohomology group

$$(2.30) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H^0(M, \mathbf{W}_{\mathbb{Z}}) & \xrightarrow{p_0} & H^0(M, Gr_{\mathcal{F}}^{-1}) & \longrightarrow & H^0(M, \mathcal{O}_M^{an}(\mathcal{X})) \\ & & \longrightarrow & H^1(M, \mathbf{W}_{\mathbb{Z}}) & \xrightarrow{p_1} & H^1(M, Gr_{\mathcal{F}}^{-1}) & \longrightarrow 0 \end{array},$$

from which (2.28) follows. q.e.d.

As a corollary, we have the following generalization of Silverberg's result (2.7).

(2.31) Theorem. *Assume that $\Gamma \backslash \mathcal{D}$ is compact. The Mordell-Weil group $MD(\mathcal{X}/\mathcal{M})$ of a Kuga fiber space is finite if and only if*

$$H^0(M, Gr_{\mathcal{F}}^{-1}) = H^1(M, \mathbf{W}_{\mathbb{Q}})^{0,0} = 0.$$

§3 SATAKE'S CLASSIFICATION OF \mathbb{Q} -SYMPLECTIC REPRESENTATIONS.

In this section, we will summarize the Satake's work of classification of \mathbb{Q} -symplectic representations. The main references are [S1], [S2].

(3.1) Preliminary.

Let F be a field of characteristic zero and D a division algebra over F . Denoting by F_1 the center of D , we set

$$(3.2) \quad [F_1 : F] = d, \quad [D : F_1] = r^2.$$

Consider a finite dimensional F -vector space V with a structure of a *right* D -module, and set $n = rank_D V$. We set:

$$\begin{aligned} GL(V/D) &= \{g \in \text{End}_D(V) \mid g \text{ is invertible} \}, \\ SL(V/D) &= \{d \in GL(V/D) \mid N(d) = 1\}, \end{aligned}$$

where N denote the reduced norm of $\text{End}_D(V)$. The corresponding matrix group are denoted by $GL_n(D)$ and $SL_n(D)$ respectively.

Let ι be an involution on D and let $\epsilon = \pm 1$. A (D, ϵ) -hermitian form h on V with respect to ι is by definition a F -bilinear mapping $h : V \times V \longrightarrow D$ satisfying the following conditions:

$$(3.3) \quad h(v, v'\alpha) = h(v, v')\alpha,$$

$$(3.4) \quad h(v', v) = \epsilon h(v, v')^{\iota} \quad \text{for all } v, v' \in V, \alpha \in D.$$

A (D, ϵ) -hermitian form h is called *non-degenerate* if a intersection matrix $T = (h(e_i, e_j))$ for a D -basis (e_i) of V is invertible. Fix an involution ι on D . For a non-degenerate (D, ϵ) -hermitian form h on V with respect to ι , we define the *unitary group* and the *special unitary group* for h by

$$(3.5) \quad U(V, h) = \{g \in GL(V/D) | h(gv, gv') = h(v, v'), \quad (v, v' \in V)\}$$

$$(3.6) \quad SU(V, h) = U(V, h) \cap SL(V/D),$$

and the corresponding matrix group are denoted by $U_n(D, h)$ and $SU_n(D, h)$ respectively.

The groups $GL_n(D)$, $SL_n(D)$, $U_n(D, h)$ and $SU_n(D, h)$ can be viewed as algebraic group defined over F_1 . For a general F_1 -group G , we denote by $R_{F_1/F}(G)$ the F -group obtained by scalar restriction (Weil [W, 1.3]).

(3.7) Classical groups over \mathbb{R} and classical domains.

If $F = \mathbb{R}$, we can define the classical groups and classical domains of type (I), (II), (III). A division algebra D over \mathbb{R} must be either \mathbb{R} , \mathbb{C} , or \mathbb{H} , and here let ι be the standard involution of D .

Let h be a non-degenerate skew-hermitian form on V (i.e. $(D, -1)$ -hermitian form) with respect to ι . We can find a D -basis (e_i) for V such that the corresponding matrix $T = (h(e_i, e_j)) \in M_n(D)$ is in the following form:

(i) $D = \mathbb{R}$; n is an even integer

$$T = J_{n/2} = \begin{pmatrix} 0 & 1_{n/2} \\ -1_{n/2} & 0 \end{pmatrix}.$$

(ii) $D = \mathbb{C}$; (p, q) is a pair of non-negative integers such that $p + q = n$.

$$T = -i1_{pq} = \begin{pmatrix} -i1_p & 0 \\ 0 & i1_q \end{pmatrix}.$$

(iii) $D = \mathbb{H}$;

$$T = j1_n.$$

Hence the corresponding special unitary groups $SU_n(D, h)$ are given by the following matrix groups:

(i)' $D = \mathbb{R}$; n is even.

$$(3.8) \quad SU_n(\mathbb{R}, h) = Sp_{n/2}(\mathbb{R}) = \{g \in SL_n(\mathbb{R}) | {}^t g J_{n/2} g = J_{n/2}\}.$$

(ii)' $D = \mathbb{C}$; $p + q = n$.

$$(3.9) \quad SU_n(\mathbb{C}, h) = SU(p, q, \mathbb{C}) = \{g \in SL_n(\mathbb{C}) | {}^t \bar{g} 1_{pq} g = 1_{pq}\}.$$

(iii)' $D = \mathbb{H}$

$$(3.10) \quad SU_n(\mathbb{H}, h) = SU_n(\mathbb{H})^- = \{g \in SL_n(\mathbb{H}) | {}^t g' (j1_n) g = j1_n\}.$$

These groups are \mathbb{R} -algebraic groups, which are of non-compact hermitian type unless $G = SU(n, 0, \mathbb{C}) \cong SU(0, n, \mathbb{C}) \cong SU(n, \mathbb{C})$ or $SU_1(\mathbb{H})^-$. Moreover these groups are \mathbb{R} -simple except for the case where $G = SU_2(\mathbb{H})^-$ (see (4.12), or [S1], Appendix, §1).

These groups act on bounded symmetric domains as follows. Consider the following set of complex structures on V

$$(3.11) \quad \mathcal{D}(V, h) = \{I \in \text{End}_{\mathbb{R}}(V) \mid I^2 = -1_V, h(x, Iy) \text{ is a positive-definite } D\text{-hermitian}\}.$$

Then the special unitary group $SU_n(D, h)$ acts on $\mathcal{D}(V, h)$ transitively, and $\mathcal{D}(V, h)$ becomes an irreducible hermitian symmetric domain and is isomorphic to a homogeneous space $SU_n(D, h)/K$ where K is a maximal compact subgroup of $SU_n(D, h)$. A bounded symmetric domain $\mathcal{D}(V, h)$ obtained as above is called a classical domain and isomorphic to one of the following bounded symmetric domains.

$$(3.12) \quad (I)_{pq} = \{Z \in M(p, q, \mathbb{C}) \mid 1_q - {}^t \bar{Z} Z \gg 0\},$$

$$(3.13) \quad (II)_n = \{Z \in M_n(\mathbb{C}) \mid {}^t Z = -Z, 1_n - {}^t \bar{Z} Z \gg 0\},$$

$$(3.14) \quad (III)_m = \{Z \in M_m(\mathbb{C}) \mid {}^t Z = Z, 1_m - {}^t \bar{Z} Z \gg 0\}.$$

The relations between $SU(V, h)$ and $\mathcal{D}(V, h)$ and the \mathbb{R} -rank of $SU(V, h)$ are shown in the following table.

	D	$G = SU(V, h)$	$\mathcal{D} = \mathcal{D}(V, h)$	$\dim_{\mathbb{C}} \mathcal{D}$	\mathbb{R} -rank
(3.15)	\mathbb{R}	$Sp_{n/2}(\mathbb{R})$	$(III)_{n/2}$	$(n/2)(n/2 + 1)/2$	$n/2$
	\mathbb{C}	$SU(p, q, \mathbb{C})$	$(I)_{pq}$	$p \cdot q$	$\min(p, q)$
	\mathbb{H}	$SU_n(\mathbb{H})^-$	$(II)_n$	$n(n - 1)/2$	$[n/2]$

(3.16) Satake's classification.

A \mathbb{Q} -symplectic representation $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ of a \mathbb{Q} -hermitian pair $(G_{\mathbb{Q}}, H_0)$ (cf. (1.1)) is called \mathbb{Q} -primary if $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}})$ is a sum of $G_{\mathbb{Q}}$ -stable subspaces isomorphic to an irreducible \mathbb{Q} -representation $\rho_1 : G_{\mathbb{Q}} \rightarrow GL(V/\mathbb{Q})$.

In this section, we review the classification of \mathbb{Q} -primary *standard* symplectic representations. In order to classify \mathbb{Q} -primary symplectic representations, the following proposition is fundamental. For a proof, see [S1], Ch. IV.

(3.17) Proposition. *Let $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ be a \mathbb{Q} -primary symplectic representation of a \mathbb{Q} -hermitian pair $(G_{\mathbb{Q}}, H_0)$, and $\rho : G_{\mathbb{Q}} \rightarrow GL(V)$ an irreducible representation containing in $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}})$. Setting*

$$D = \text{End}_{G_{\mathbb{Q}}}(V), \quad F_1 = \text{Cent } D, \quad U = \text{Hom}_{G_{\mathbb{Q}}}(V, W_{\mathbb{Q}}),$$

we have the following.

(i) D is a division algebra over \mathbb{Q} , and V (resp. U) becomes a left D -module (resp. a right D -module).

(ii) There exists a canonical isomorphism

$$(3.18) \quad W_{\mathbb{Q}} \cong U \otimes_D V.$$

(iii) There exist a natural involution ι on D , a (D, ϵ) -hermitian form h on V and a $(D, -\epsilon)$ -hermitian form h' on U with respect to the involution ι such that

$$(3.19) \quad A_{\mathbb{Q}} = \text{tr}_{D/\mathbb{Q}}(h' \otimes h).$$

(iv) The form h on V is $G_{\mathbb{Q}}$ -invariant. In particular, ρ is reduced to a natural representation over F_1

$$(3.20) \quad \rho_1 : G_{\mathbb{Q}} \longrightarrow SU(V, h)$$

(with $\text{End}_{G_{\mathbb{Q}}}(V) = D$).

(3.21) Definition. A \mathbb{Q} -primary representation $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ of a \mathbb{Q} -hermitian pair $(G_{\mathbb{Q}}, H_0)$ is said to be *standard* if $G_{\mathbb{Q}} = R_{F_1/\mathbb{Q}}(SU(V, h))$ and ρ in (3.20) is induced by the universal homomorphism of the scalar restriction (cf. [W, 1.13]).

(3.22) Remark. Satake [S2] determined all \mathbb{Q} -primary symplectic representation under an reasonable additional condition. Besides the standard one, there exist few non-standard representations involving skew-symmetric representations and spin representations. But there exist also a \mathbb{Q} -primary symplectic representation which does not satisfy his condition (see p195 [S1] for references). In this paper, we will not deal with non-standard case.

A standard representation is determined only by the data D, ι, V, U, h, h' in proposition (3.17). First we have the following

(3.23) Proposition. ([S1], Ch. IV, §6). Let $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ be a \mathbb{Q} -primary symplectic representation (not necessarily standard) of a \mathbb{Q} -hermitian pair $(G_{\mathbb{Q}}, H_0)$, and $D, F_1, \iota, V, h, U, h'$ be as in lemma (3.17). Then one of the following cases occurs.

(R1) $D = F_1$ is a totally real algebraic number field and $\iota = \text{identity}$, and h is a symplectic form on V ($\epsilon = -1$).

(R2, ϵ) D is a quaternion algebra over a totally real algebraic number field F_1 and ι is the standard involution, h is a (D, ϵ) -hermitian form V with respect to ι , where $\epsilon = \pm 1$.

(C) F_1 is a CM field, i.e. a purely imaginary quadratic extension of a totally real algebraic number field F_{10} , D is a central division algebra over F_1 , ι is an involution of D of the second kind, and h is a (D, ϵ) -hermitian form with respect to ι where $\epsilon = \pm 1$.

Let D, F_1, ι be as in Proposition (3.23). If we set $F_1^+ = \{z \in F_1 \mid z' = z\}$, then F_1^+ is a totally real algebraic number field. Setting $t = [F_1^+ : \mathbb{Q}]$, let $\{\tau_i : F_1^+ \hookrightarrow \mathbb{R}, 1 \leq i \leq t\}$ be

the set of t -distinct embeddings of F_1^+ into \mathbb{R} . For each $\tau_i : F_1^+ \hookrightarrow \mathbb{R}$, we put

$$(3.24) \quad F_1^{(i)} = F_1 \otimes_{F_1^+, \tau_i} \mathbb{R},$$

$$(3.25) \quad D^{\tau_i} = D \otimes_{F_1^+, \tau_i} \mathbb{R},$$

$$(3.26) \quad W^{\tau_i} = W_{\mathbb{Q}} \otimes_{F_1^+, \tau_i} \mathbb{R},$$

$$(3.27) \quad V^{\tau_i} = V \otimes_{F_1^+, \tau_i} \mathbb{R},$$

$$(3.28) \quad U^{\tau_i} = U \otimes_{F_1^+, \tau_i} \mathbb{R}.$$

The algebra D^{τ_i} becomes a central simple algebra over $F_1^{(i)}$, so there exists a division algebra $D^{(i)}$ over $F_1^{(i)}$ such that

$$D^{\tau_i} \cong M_s(D^{(i)}).$$

Fixing an above isomorphism, we denote by $\epsilon_{\nu\mu}^i$ the corresponding matrix unit in D^{τ_i} . We moreover set:

$$(3.29) \quad V^{(i)} := \epsilon_{11}^i V^{\tau_i}, \quad U^{(i)} = U^{\tau_i} \epsilon_{11}^i.$$

Then $V^{(i)}$ (resp. $U^{(i)}$) are left (resp. right) $D^{(i)}$ -modules and we have an isomorphism (cf. [S1], p189),

$$(3.30) \quad W^{\tau_i} = U^{(i)} \otimes_{D^{(i)}} V^{(i)}.$$

Note that from (3.23), $F_1^{(i)}$ is isomorphic to \mathbb{R} or \mathbb{C} , corresponding to the case **(R1)**, **(R2)**, **(e)** or **(C)**, so $D^{(i)}$ is isomorphic to \mathbb{R} , \mathbb{H} , or \mathbb{C} .

Under these notations, we can state the following theorem.

(3.31) Theorem. ([S1], Ch. IV. §6). *Let $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ be a standard \mathbb{Q} -primary symplectic representation, and $D, \iota, F_1, V, h, U, h', W_{\mathbb{Q}} = U \otimes_D V, A_{\mathbb{Q}} = \text{tr}_{D/\mathbb{Q}}(h' \otimes h)$ be as in (3.17). Then we have the following.*

(i) *There exists a decomposition*

$$(3.32) \quad W_{\mathbb{R}} := W_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{R} = \bigoplus_{i=1}^t W^{\tau_i} \cong \bigoplus_{i=1}^t U^{(i)} \otimes_{D^{(i)}} V^{(i)}.$$

(ii) *For each $i, 1 \leq i \leq t$, h (resp. h') induces a $(D^{(i)}, \epsilon\eta_i)$ -hermitian form $h^{(i)}$ on $V^{(i)}$ (resp. $(D^{(i)}, -\epsilon\eta_i)$ -hermitian form $h'^{(i)}$ on $U^{(i)}$), where $\eta_i = \pm 1$. We have a decomposition of $A_{\mathbb{R}} := A_{\mathbb{Q}} \otimes \mathbb{R} = \bigoplus_{i=1}^t A^{(i)}$ corresponding to (3.32), where one set*

$$(3.33) \quad A^{(i)} := \text{tr}_{D^{(i)}/F^{(i)}}(h'^{(i)} \otimes h^{(i)}).$$

(iii) *The \mathbb{R} -valued points $G_{\mathbb{R}}$ of $G_{\mathbb{Q}} = R_{F_1/\mathbb{Q}}(SU(V, h))$ has a canonical decomposition*

$$(3.34) \quad G_{\mathbb{R}} = R_{F_1/\mathbb{Q}}(SU(V, h))_{\mathbb{R}} = \prod_{i=1}^t SU(V^{(i)}, h^{(i)}),$$

and, for each i , the natural representation $\rho_1 : G_{\mathbb{Q}} \longrightarrow SU(V, h)$ induces a representation

$$(3.35) \quad \rho_1^{(i)} : G_{\mathbb{R}} = R_{F_1/\mathbb{Q}}(SU(V, h))_{\mathbb{R}} \longrightarrow SU(V^{(i)}, h^{(i)}),$$

where $\rho_1^{(i)}$ can be written in the form

$$(3.36) \quad \rho_1^{(i)} = 1 \otimes \cdots \otimes 1 \otimes id_{V^{(i)}} \otimes 1 \cdots \otimes 1.$$

according to the decomposition (3.34).

Moreover, for each case in (3.23), we have the following

(3.37) Theorem. ([S1], Ch. IV, §6). Under the notation in Proposition (3.23), we have the following explicit descriptions of $F_1^{(i)}$, D^{τ_i} , $D^{(i)}$, $V^{(i)}$, $h^{(i)}$, $U^{(i)}$, $G_{\mathbb{R}}$ for the cases of **(R1)**, **(R2, ϵ)**, **(C)** respectively.

(R1) ($\epsilon = -1$) $D = F_1 = F_1^+$. Set $\dim_{F_1} V = n$, $\dim_{F_1} U = m$. Then one has:

$$F^{(i)} \cong D^{\tau_i} \cong D^{(i)} \cong \mathbb{R}, \quad V^{(i)} \cong \mathbb{R}^n, \quad U^{(i)} \cong \mathbb{R}^m,$$

$$h^{(i)} : \mathbb{R}\text{-symplectic form on } V^i, \quad (\eta_i = 1) \text{ for } 1 \leq i \leq t = d,$$

$$(3.38) \quad G_{\mathbb{R}} \cong \underbrace{Sp_{n/2}(\mathbb{R}) \times \cdots \times Sp_{n/2}(\mathbb{R})}_d.$$

(R2, ϵ) We have $F_1 = F_1^+$, and D is a quaternion algebra over F_1 . Set $\text{rank}_D V = n$, $\text{rank}_D U = m$. Then one has $F^{(i)} = \mathbb{R}$. After a suitable renumbering of $\{\tau_i\}$, we may assume that for some t' , $0 \leq t' \leq t$,

$$D^{\tau_i} \cong \begin{cases} \mathbb{H} & 1 \leq i \leq t' \\ M_2(\mathbb{R}) & t' + 1 \leq i \leq t, \end{cases} \quad D^{(i)} \cong \begin{cases} \mathbb{H} & 1 \leq i \leq t' \\ \mathbb{R} & t' + 1 \leq i \leq t. \end{cases}$$

Then one has:

$$V^{(i)} \cong \begin{cases} \mathbb{H}^n \\ \mathbb{R}^{2n} \end{cases}, \quad U^{(i)} \cong \begin{cases} \mathbb{H}^m \\ \mathbb{R}^{2m} \end{cases}, \quad W^{\tau_i} \cong \begin{cases} \mathbb{H}^n \otimes_{\mathbb{H}} \mathbb{H}^m & 1 \leq i \leq t' \\ \mathbb{R}^{2n} \otimes_{\mathbb{R}} \mathbb{R}^{2m} & t' + 1 \leq i \leq t. \end{cases}$$

($\epsilon = 1$)

$$h^{(i)} = \begin{cases} \text{positive-definite } \mathbb{H}\text{-symmetric form } (\eta_i = 1) & 1 \leq i \leq t', \\ \mathbb{R}\text{-symplectic form } (\eta_i = -1) & t' + 1 \leq i \leq t, \end{cases}$$

$$(3.39) \quad G_{\mathbb{R}} = \underbrace{SU_n(\mathbb{H}) \times \cdots \times SU_n(\mathbb{H})}_{t' \times \text{compact}} \times \underbrace{Sp_n(\mathbb{R}) \times \cdots \times Sp_n(\mathbb{R})}_{(t-t') \times (III)_n}.$$

($\epsilon = -1$)

$$(3.40) \quad h^{(i)} = \begin{cases} \mathbb{H}\text{-symplectic form } (\eta_i = 1) & 1 \leq i \leq t', \\ \text{positive-definite } \mathbb{R}\text{-symmetric form } (\eta_i = -1) & t' + 1 \leq i \leq t, \end{cases}$$

$$G_{\mathbb{R}} = \underbrace{SU_n(\mathbb{H})^- \times \cdots \times SU_n(\mathbb{H})^-}_{t' \times (11)_n} \times \underbrace{SO_{2n}(\mathbb{R}) \times \cdots \times SO_{2n}(\mathbb{R})}_{(t-t') \times \text{compact}}.$$

(C) ($\epsilon = \pm 1$). F_1 is a purely imaginary quadratic extension of F_1^+ , so $t = \frac{1}{2}[F_1 : \mathbb{Q}]$. We set $[D : F_1] = r^2$, $\text{rank}_D V = n$, and $\text{rank}_D U = m$. Then one has:

$$F_1^{(i)} \cong D^{(i)} \cong \mathbb{C}, \quad D^{r_i} \cong M_r(\mathbb{C}),$$

$$V^{(i)} \cong \mathbb{C}^{nr}, \quad U^{(i)} \cong \mathbb{C}^{mr}, \quad W^{r_i} \cong \mathbb{C}^{mr} \otimes_{\mathbb{C}} \mathbb{C}^{nr}.$$

We may assume that for $t', 0 \leq t' \leq t$,

$$h^{(i)} = \begin{cases} \mathbb{C}\text{-symplectic form with the signature } (p_i, q_i) & 1 \leq i \leq t' \quad (p_i \geq q_i), \\ \text{positive-definite } \mathbb{C}\text{-hermitian form} & t' + 1 \leq i \leq t, \end{cases}$$

$$(3.41) \quad G_{\mathbb{R}} \cong \prod_{i=1}^{t'} \underbrace{SU(p_i, q_i, \mathbb{C})}_{(1)_{p_i, q_i}} \times \underbrace{SU_{nr}(\mathbb{C}) \times \cdots \times SU_{nr}(\mathbb{C})}_{(t-t') \times \text{compact}}.$$

(3.42) Proposition. A \mathbb{Q} -algebraic group $G_{\mathbb{Q}} = R_{F_1/\mathbb{Q}}(SU(V, h))$ in (3.37) is Zariski connected. Assume that $G_{\mathbb{R}}$ is non-compact, i.e., $\dim \mathcal{D} \geq 1$. Then $G_{\mathbb{Q}}$ is \mathbb{Q} -simple except for the case (R2,-1), $n = 2$.

PROOF: See [S1], Appendix, §1.

§4. VANISHING THEOREM AND THE CASE $rk_{\mathbb{R}} G_{\mathbb{R}} \geq 2$.

Let G be a connected semi-simple real Lie group with finite center of hermitian type, K a maximal compact subgroup of G , so that a quotient space $\mathcal{D} = G/K$ becomes a hermitian symmetric bounded domain. Let Γ a discrete subgroup of G of a finite covolume with respect to the Haar measure. If Γ is torsion-free, the quotient space $M = \Gamma \backslash \mathcal{D}$ becomes a smooth quasi-projective variety. For a finite dimensional complex representation $\rho : G \rightarrow GL(W_{\mathbb{C}})$, we denote by $W_{\mathbb{C}}$ the associated local system on $M = \Gamma \backslash \mathcal{D}$. Let $L_{(2)}(M, W_{\mathbb{C}})$ be as in (2.9), and $H_{(2)}(M, W_{\mathbb{C}})$ the L^2 -cohomology group for it. Let $L^2(\Gamma \backslash \mathcal{D})^{\infty}$ denote the set of C^{∞} square-integrable function on $\Gamma \backslash \mathcal{D}$, and view it as a unitary G -module under the right translation. Since it is a (\mathfrak{g}, K) -module, we may consider the relative Lie algebra complex $C^*(\mathfrak{g}, K; L^2(\Gamma \backslash G)^{\infty} \otimes W_{\mathbb{C}})$, whose cohomology yields the relative Lie algebra cohomology (cf. [B-W]).

First, we recall the following

(4.1) **Theorem.** ([B], [B-C]). *There exists a quasi-isomorphism*

$$C^*(\mathfrak{g}, K; L^2(\Gamma \backslash G)^\infty \otimes W_{\mathbf{C}}) \longrightarrow L_{(2)}^*(M, W_{\mathbf{C}})^\infty.$$

In particular, we have isomorphisms

$$Ext_{(\mathfrak{g}, K)}(W^*, L^2(\Gamma \backslash G)^\infty) \cong H_{(2)}(\Gamma \backslash \mathcal{D}, \mathbf{W}).$$

Write $L_2(\Gamma \backslash G)^\infty$ as the direct sum of the discrete spectrum $L_2(\Gamma \backslash G)_d^\infty$ and its orthogonal complement, the so-called continuous spectrum $L_2(\Gamma \backslash G)_{ct}^\infty$.

The following theorem is a special case of results in [B-C].

(4.2) **Theorem.** (see [B-C], Prop. 4.4 and Th. 4.5) *Under the assumption as above, we have the following.*

- (i) $H_{(2)}(M, W_{\mathbf{C}})$ is finite dimensional¹,
- (ii) there exists a finite set $(H_i), (i \in S)$ of mutually orthogonal closed irreducible G -invariant subspaces of $L^2(\Gamma \backslash G)_d$ such that

$$(4.3) \quad H_{(2)}(M, W_{\mathbf{C}}) = Ext_{(\mathfrak{g}, K)}(W_{\mathbf{C}}^*, L^2(\Gamma \backslash G)_d^\infty) = \bigoplus_{i \in S} Ext_{(\mathfrak{g}, K)}(W_{\mathbf{C}}^*, H_i),$$

(4.4) **Definition.** Let G be as above. We say that G has *no compact factor* if it has no infinite normal compact subgroup. A discrete subgroup Γ of G is said to be *irreducible* if the image of Γ under any surjective morphism $G \rightarrow G'$ with non-trivial image and non-compact kernel is non-discrete.

We can prove the following vanishing theorem of L^2 -cohomology group.

(4.5) **Theorem.** *Let G be as above. Assume that G has no compact factor and Γ is an irreducible discrete subgroup of G with a finite covolume. If $(\rho, W_{\mathbf{C}})$ is a non-trivial finite complex representation of G , we have*

$$H_{(2)}^q(\Gamma \backslash \mathcal{D}, W_{\mathbf{C}}) = 0 \quad \text{for } q < rk_{\mathbb{R}} G.$$

where $rk_{\mathbb{R}} G$ denote the \mathbb{R} -rank of G .

PROOF: If Γ is cocompact, then this is nothing but Proposition 6.4 in Ch. VII of Borel-Wallach [B-W]. Thanks to (4.3), their proof works even if $\Gamma \backslash G$ is not compact.

(4.6) **Vanishing theorem.**

Now we apply this theorem for standard \mathbb{Q} -primary symplectic representations. Let $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ be a standard \mathbb{Q} -symplectic representation, $D, \iota, F_1, V, U, h, h'$ as in (3.17), and $G_{\mathbb{Q}} = R_{F_1/\mathbb{Q}}(SU(V, h))$.

¹Of course, this also follows from the Zucker conjecture (2.11)

We take a lattice $V_{\mathbb{Z}}$ in V (see §1), and set $D_{\mathbb{Z}} = \{m \in D \mid mV_{\mathbb{Z}} \subset V_{\mathbb{Z}}\}$. Then $D_{\mathbb{Z}}$ becomes a \mathbb{Z} -subalgebra of D such that $D_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Q} \cong D$, which is called an *order* of D . Taking a $D_{\mathbb{Z}}$ -right submodule $U_{\mathbb{Z}}$ of U , we set

$$(4.7) \quad W_{\mathbb{Z}} = U_{\mathbb{Z}} \otimes_{D_{\mathbb{Z}}} V_{\mathbb{Z}}.$$

Then $W_{\mathbb{Z}}$ becomes a lattice in $W_{\mathbb{Q}}$, and we may assume that $W_{\mathbb{Z}}$ satisfies the condition (1.7), i.e., $A_{\mathbb{Q}}(W_{\mathbb{Z}}, W_{\mathbb{Z}}) \subset \mathbb{Z}$. From definition (1.4) and the above construction, we have an isomorphism of discrete groups

$$G_{W_{\mathbb{Z}}} \cong G_{V_{\mathbb{Z}}}.$$

Take a torsion-free arithmetic subgroup $\Gamma \subset G_{V_{\mathbb{Z}}}$.

Let $G_{\mathbb{Q}} = R_{F_1/\mathbb{Q}}(SU(V, h))$ be as above. Then from (3.37) and (3.42), except for the case $(\mathbf{R2}, -1)$, $\mathfrak{n} = 2$, we can write

$$(4.8) \quad G_{\mathbb{R}} = G_1 \times \cdots \times G_l \times U,$$

where $G_i = SU(V^{(i)}, h^{(i)})$ is a \mathbb{R} -simple non-compact Lie group of hermitian type for $1 \leq i \leq l$ and U is a compact group.

(4.9) Proposition. *Assume that (V, h) is not in the case $(\mathbf{R2}, -1)$, $\mathfrak{n} = 2$. For any torsion-free arithmetic subgroup $\Gamma \subset G_{\mathbb{R}}$, let Γ' denote the image of Γ under the projection $G_{\mathbb{R}} \rightarrow G'_{\mathbb{R}} = G_1 \times \cdots \times G_l$ (cf. (4.8)). Then Γ' is an irreducible torsion-free discrete subgroup with finite covolume.*

PROOF: It is easy to see that Γ' is a discrete subgroup in $G'_{\mathbb{R}}$ with finite covolume. Let $\rho_1^{(i)} : G_{\mathbb{R}} \rightarrow G_i = SU(V^{(i)}, h^{(i)})$ be the representation in (3.35) for $1 \leq i \leq l$. Then from the construction we can see that $\rho_{i|\Gamma'}$ induces an isomorphism $\Gamma \cong \rho_i(\Gamma)$. By a corollary in No. 4 of [Shz], Γ' is irreducible in $G'_{\mathbb{R}}$. Since the projection map $\Gamma \rightarrow \Gamma'$ is injective, Γ' is also torsion-free.

Let K be a maximal compact subgroup of $G_{\mathbb{R}} = G_1 \times \cdots \times G_l \times U$, and write K as $K_1 \times \cdots \times K_l \times U$, so that the corresponding hermitian symmetric space $\mathcal{D} = G_{\mathbb{R}}/K$ has a decomposition as

$$(4.10) \quad \mathcal{D} = \mathcal{D}_1 \times \cdots \times \mathcal{D}_l,$$

where $\mathcal{D}_i = G_i/K_i$ are irreducible symmetric spaces. We have a natural isomorphism

$$(4.11) \quad M := \Gamma \backslash \mathcal{D} \cong \Gamma' \backslash \mathcal{D}$$

(4.12) Remark. We have an isomorphism $SU_2(\mathbb{H})^- \cong SU(2, \mathbb{C}) \times SL_2(\mathbb{R})$.

Now we state our main theorem in this section.

(4.13) Theorem. *Let $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ be a standard \mathbb{Q} -primary symplectic representation, which is not the case $(\mathbf{R2}, -1)$, $n=2$, and (V, h) , $\Gamma \subset G_{\mathbb{R}}$ as above. Assume that $rk_{\mathbb{R}} G_{\mathbb{R}} \geq 2$. Then we have*

$$(4.14) \quad H^q(M, \mathbf{W}_{\mathbb{Q}}) = 0, \quad q \leq 1.$$

Even if $rk_{\mathbb{R}} G_{\mathbb{R}} = 1$, we have $H^0(M, \mathbf{W}_{\mathbb{Q}}) = 0$.

PROOF: From (3.17), $W_{\mathbb{Q}}$ is a vector space over a field $F_1 = \text{Cent}(D)$. The field F_1 is a totally real field, or a CM field (see (3.23)). Set $t = [F_1 : \mathbb{Q}]$. Let $\{\sigma_i : F_1 \hookrightarrow \mathbb{C}\}_{i=1}^t$ denote the set of t -distinct embeddings of F_1 into \mathbb{C} . For an embedding $\sigma_i : F_1 \hookrightarrow \mathbb{C}$, we put

$$(4.15) \quad W^{\sigma_i} = W_{\mathbb{Q}} \otimes_{F_1, \sigma_i} \mathbb{C}, \quad \mathbf{W}^{\sigma_i} = \mathbf{W}_{\mathbb{Q}} \otimes_{F_1, \sigma_i} \mathbb{C}.$$

By the universal coefficient theorem, we have an isomorphism

$$(4.16) \quad H^q(M, \mathbf{W}_{\mathbb{Q}}) \otimes_{F_1, \sigma_i} \mathbb{C} \cong H^q(M, \mathbf{W}^{\sigma_i}).$$

Note that \mathbf{W}^{σ_i} is a local system on M associated to a representation

$$(4.17) \quad (\rho_{\mathbb{Q}})^{(i)} : G_{\mathbb{R}} \longrightarrow GL(W^{\sigma_i}/\mathbb{C}).$$

induced by $\rho_{\mathbb{Q}}$. From the assumption and (4.9), an arithmetic group $\Gamma \subset G_{\mathbb{R}}$ is irreducible, so from (2.13), we have isomorphisms

$$(4.18) \quad H_{(2)}^q(M, \mathbf{W}^{\sigma_i}) \cong H^q(M, \mathbf{W}^{\sigma_i}) \quad \text{for } q \leq 1.$$

From (4.16) and (4.18), in order to show (4.14), it suffices to show that

$$(4.19) \quad H_{(2)}^q(M, \mathbf{W}^{\sigma_i}) = 0 \quad \text{for } q \leq 1.$$

Recall that we have an isomorphism $W_{\mathbb{Q}} = U \otimes_D V$ (see (3.17)). Set $U^{\sigma_i} := U \otimes_{F_1, \sigma_i} \mathbb{C}$, $V^{\sigma_i} := V \otimes_{F_1, \sigma_i} \mathbb{C}$, and $D^{\sigma_i} := D \otimes_{F_1, \sigma_i} \mathbb{C}$. Choosing an isomorphism $D^{\sigma_i} \cong M_s(\mathbb{C})$, let $\epsilon_{\mu\nu}^i$ denote the matrix unit in D^{σ_i} . Then, as in (3.29) and (3.30), setting $U_{\mathbb{C}}^{(i)} := U^{\sigma_i} \epsilon_{11}^i$, $V_{\mathbb{C}}^{(i)} := \epsilon_{11}^i V^{\sigma_i}$, we have an isomorphism

$$(4.20) \quad W^{\sigma_i} \cong U_{\mathbb{C}}^{(i)} \otimes_{\mathbb{C}} V_{\mathbb{C}}^{(i)}.$$

Assume that F_1 is totally real. Then, the representation $\rho_1 : G_{\mathbb{Q}} \longrightarrow SU(V, h)$ induces a representation

$$\rho_{1\mathbb{C}}^{(i)} : G_{\mathbb{R}} \longrightarrow SU(V_{\mathbb{C}}^{(i)}, h_{\mathbb{C}}^{(i)})$$

which is obtained by a scalar extension of (3.35) from \mathbb{R} to \mathbb{C} . Hence, from (3.36), $\rho_{1\mathbb{C}}^{(i)}$ can be written in form

$$\rho_{1\mathbb{C}}^{(i)} = 1 \otimes \cdots \otimes 1 \otimes (id_{V^{(i)}})_{\mathbb{C}} \otimes 1 \cdots \otimes 1.$$

Write $G_{\mathbb{R}} = G_1 \times \cdots \times G_l \times U$ as in (4.8) and take i such that $1 \leq i \leq l$. Then since $\rho_{1\mathbb{C}}^{(i)}$ is trivial on the compact factor U , it descends to a representation of $G'_{\mathbb{R}} = G_1 \times \cdots \times G_l$. Let Γ' be as in (4.9). Then we can apply Theorem (4.5) for $G'_{\mathbb{R}}$, $\rho_{1\mathbb{C}}^{(i)}$, $V_{\mathbb{C}}^{(i)}$, Γ' to deduce that

$$(4.21) \quad H_{(2)}^q(M, V_{\mathbb{C}}^{(i)}) = 0 \text{ for } q < rk_{\mathbb{R}} G'_{\mathbb{R}}.$$

By the assumption that $rk_{\mathbb{R}} G'_{\mathbb{R}} \geq 2$, one has

$$H_{(2)}^q(M, V_{\mathbb{C}}^{(i)}) = 0 \text{ for } q \leq 1.$$

Hence the assertion (4.19) (so (4.14)) follows from this and the following isomorphism.

$$H_{(2)}^q(M, W^{\sigma_i}) \cong U_{\mathbb{C}}^{(i)} \otimes_{\mathbb{C}} H_{(2)}^q(M, V^{(i)}) \quad \text{by (4.20)}.$$

The proof for the case when F_1 is a CM field is similar, so we omit it.

(4.22) Remark. Note that we have the isomorphism $SU_3(\mathbb{H})^- \cong SU(3, 1, \mathbb{C})$.

By virtue of Silverberg's criterion (2.7), as a corollary of (4.13), we obtain the following

(4.23) Theorem. *The Mordell-Weil group $MW(\mathcal{X}/M)$ of a Kuga fiber space $f : \mathcal{X} \rightarrow M$ associated to a standard \mathbb{Q} -primary symplectic representation is finite whenever $rk_{\mathbb{R}} G_{\mathbb{R}} \geq 2$.*

§5 \mathbb{R} -RANK 1 AND Γ COCOMPACT.

(5.1) In this section, we shall deal with the cases where the \mathbb{R} -rank of $G_{\mathbb{R}}$ is 1 and Γ is cocompact. For technical reasons, we exclude the case **(R2, -1)**, $\mathfrak{n} = 2$.

From the Satake's classification (cf. Theorem (3.37)), the cases where $G_{\mathbb{R}}$ has the \mathbb{R} -rank 1 are listed as follows:

$$(5.2) \quad \text{Case (R2, -1), } \mathfrak{n} = 3 \quad G_{\mathbb{R}} \cong SU_3(\mathbb{H})^- \times \underbrace{SO_6(\mathbb{R}) \times \cdots \times SO_6(\mathbb{R})}_{\text{possibly}=\{1\}},$$

$$(5.3) \quad \text{Case (C)} \quad G_{\mathbb{R}} \cong SU(nr - 1, 1) \times \underbrace{SU_{nr}(\mathbb{C}) \times \cdots \times SU_{nr}(\mathbb{C})}_{\text{possibly}=\{1\}}.$$

and

$$(5.4) \quad \dim \mathcal{D} = 1.$$

In the above case, we can no more expect the vanishing of the $H^1(M, W_{\mathbb{C}})$ in general, though we have the vanishing of $H^0(M, W_{\mathbb{C}})$ (see (4.13)). In fact, in the case (5.3) when

$r = 1$ and $t \geq 2$, there is an arithmetic subgroup $\Gamma \subset G_{W_{\mathbf{z}}}$ such that $H^1(M, \mathbf{W}_{\mathbf{C}}) \neq 0$ (See Ch. VIII, §5, [B-W]). Hence we should consider the Hodge decomposition of $H^1(M, \mathbf{W}_{\mathbf{C}})$, and appeal to Theorem (2.31). In this section, we always assume that $\Gamma \backslash \mathcal{D}$ is compact. Note that $\Gamma \backslash \mathcal{D}$ is compact whenever $G_{\mathbb{R}}$ has a compact factor.

(5.5) Let $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ be a standard \mathbb{Q} -primary symplectic representation, $W_{\mathbf{z}} \subset W_{\mathbb{Q}}$ a lattice, $\Gamma \subset G_{W_{\mathbf{z}}} \subset G_{\mathbb{R}}$ a torsion free arithmetic subgroup. Let $(\mathbf{W}_{\mathbf{z}}, \mathbf{A}, \mathcal{F}^0)$ denote the corresponding VPHS over the smooth manifold $\Gamma \backslash \mathcal{D}$ (see (1.10)). The main result in this section is the following

(5.6) **Theorem.** *Under the notation as above, we have*

$$(5.7) \quad H^1(M, \mathbf{W}_{\mathbb{Q}})^{0,0} = 0$$

in the cases (5.2), (5.3), and (5.4).

As a corollary of this theorem, we have the following

(5.8) **Corollary.** *The Mordell-Weil groups of the Kuga fiber spaces associated to a standard \mathbb{Q} -primary symplectic representation is finite when $rk_{\mathbb{R}} G_{\mathbb{R}} = 1$ and $\Gamma \backslash \mathcal{D}$ is compact.*

PROOF: Since we always have $H^0(M, \mathbf{W}_{\mathbf{C}}) = 0$, by (2.31), Theorem (5.6) implies the assertion.

(5.9) **A reduction.**

We keep the notation in (5.5). Let F_1, D be as in (3.17). Denote by $\{\sigma_1, \dots, \sigma_d\}$ the set of all embeddings F_1 into \mathbb{C} where $d = [F_1 : \mathbb{Q}]$. Considering $W_{\mathbb{Q}}$ as a F_1 -vector, we set $W^{\sigma_i} = W_{\mathbb{Q}} \otimes_{F_1, \sigma_i} \mathbb{C}$ and $\mathbf{W}^{\sigma_i} = \mathbf{W}_{\mathbb{Q}} \otimes_{F_1, \sigma_i} \mathbb{C}$. Then we have the decompositions

$$(5.10) \quad \mathbf{W}_{\mathbf{C}} := \mathbf{W}_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{C} = \bigoplus_{i=1}^d \mathbf{W}^{\sigma_i},$$

$$(5.11) \quad H^1(M, \mathbf{W}_{\mathbf{C}}) = \bigoplus_{i=1}^d H^1(M, \mathbf{W}^{\sigma_i}).$$

Let $\nabla : \mathcal{O}_M(\mathbf{W}_{\mathbf{C}}) \rightarrow \Omega_M^1 \otimes \mathbf{W}_{\mathbf{C}}$ denote the Gauss-Manin connection on $\mathbf{W}_{\mathbf{C}}$. From the horizontality, we have the complex

$$(5.12) \quad \nabla : \mathcal{F}^0 \rightarrow \Omega_M^1 \otimes Gr_{\mathcal{F}}^{-1}$$

whose \mathbf{H}^1 is isomorphic to $H^1(M, \mathbf{W}_{\mathbf{C}})^{0,0}$ (see (2.21)). We have the following

(5.13) **Lemma.** *Assume that the Hodge filtration \mathcal{F}^0 and the Gauss-Manin connection ∇ on $\mathbf{W}_{\mathbf{C}}$ is compatible with the decomposition (5.10). Then if for at least one $\sigma_i : F_1 \hookrightarrow \mathbb{C}$*

$$(5.14) \quad H^1(M, \mathbf{W}^{\sigma_i})^{0,0} = 0,$$

we have $H^1(M, \mathbf{W}_{\mathbb{Q}})^{0,0} = 0$.

PROOF: From the construction of the Hodge structure in (2.15), under the assumption, we have the decomposition

$$H^1(M, \mathbf{W}^{\sigma_i}) = \bigoplus_{p+q=0} H_{\sigma_i}^{p,q}$$

such that

$$H^1(M, \mathbf{W}_{\mathbb{C}})^{p,q} = \bigoplus_{i=1}^d H_{\sigma_i}^{p,q}.$$

Let $\pi_i : H^1(M, \mathbf{W}_{\mathbb{Q}}) \longrightarrow H^1(M, \mathbf{W}^{\sigma_i})$ be the natural projection map. Then we have

$$H^1(M, \mathbf{W}_{\mathbb{Q}})^{0,0} = \bigcap_{i=1}^d \pi_i^{-1}(H_{\sigma_i}^{0,0}).$$

Since the map π_i is injective, this implies the assertion.

(5.15) Gauss-Manin complex.

Let $(G_{\mathbb{Q}}, H_0)$ be the \mathbb{Q} -hermitian pair corresponding to the \mathbb{Q} -symplectic representation in (5.5), and K the maximal compact subgroup of $G_{\mathbb{R}}$ corresponding to H_0 . We also denote by $\mathfrak{g}_{\mathbb{R}}, \mathfrak{k}$ the Lie algebras of $G_{\mathbb{R}}$ and K respectively, and by \mathfrak{p} the orthogonal complement of \mathfrak{k} in $\mathfrak{g}_{\mathbb{R}}$ with respect to the Killing form. Let us set $W_{\mathbb{C}}^{\pm} = W_{\mathbb{C}}(\pm i, I_{\mathbb{C}})$, $\mathfrak{p}^{\pm} = \mathfrak{p}_{\mathbb{C}}(\pm i, \text{ad}_{\mathfrak{p}}(H_0))$. Then, by the condition (1.2), the spaces $W_{\mathbb{C}}^{\pm}$ and \mathfrak{p}^{\pm} are stable under the action of K , hence they become representations of K .

For any representation T of K , we can define a holomorphic vector bundle, or a locally free sheaf \mathcal{T} on $M = \Gamma \backslash \mathcal{D}$ as in §2 in [Z2]. In the notation in §1, the representations $W_{\mathbb{C}}^+$ (resp. $W_{\mathbb{C}}^-$) defines a Hodge bundle \mathcal{F}^0 (resp. $Gr_{\mathcal{F}}^{-1}$) and \mathfrak{p}^- defines the cotangent sheaf Ω_M^1 on M .

We call the natural complex

$$(5.16) \quad \nabla : \mathcal{F}^0 \longrightarrow \Omega_M^1 \otimes Gr_{\mathcal{F}}^{-1}$$

the (first) Gauss-Manin complex. Then the Gauss-Manin complex in this case is induced by the following homomorphism of the representations of K :

$$(5.17) \quad W_{\mathbb{C}}^+ \longrightarrow \mathfrak{p}^- \otimes W_{\mathbb{C}}^-.$$

(5.18) Proof of Theorem (5.6) in the case (5.3).

In this case, since F_1 is a CM field, we can denote by $\{\sigma_1, \dots, \sigma_t, \overline{\sigma_1}, \dots, \overline{\sigma_t}\}$ the set of all embeddings of F_1 into \mathbb{C} such that $\sigma_i|_{F_1^+}$ is an extension of $\tau_i : F_1^+ \hookrightarrow \mathbb{R}$. Since $G_{\mathbb{R}} = \prod_{i=1}^t SU(V^{(i)}, h^{(i)}) \cong SU(nr-1, 1, \mathbb{C}) \times SU_{nr}(\mathbb{C}) \times \dots \times SU_{nr}(\mathbb{C})$, $(V^{(1)}, h^{(1)})$ is a \mathbb{C} -vector space with a skew-hermitian form $h^{(1)}$ such that the signature of $ih^{(1)}$ is $(nr-1, 1)$. Recalling that the decomposition $W_{\mathbb{R}} = \bigoplus_{i=1}^t W^{\tau_i} = \bigoplus_{i=1}^t U^{(i)} \otimes_{\mathbb{C}} V^{(i)}$, we can write the complex structure $I \in \mathcal{D}(W_{\mathbb{R}}, A_{\mathbb{R}})$ as

$$I = 1_{U^{(1)}} \otimes I_{(1)} + \sum_{i=2}^t I'_{(i)} \otimes 1_{V^{(i)}},$$

for some $I_{(1)} \in \mathcal{D}(V^{(1)}, h^{(1)}) \cong (I)_{nr-1,1}$ and $I'_{(i)} \in \mathcal{D}(U^{(i)}, h'^{(i)})$. (See [S1], Ch. IV or [S2]). If we set

$$H'_0 = I_{(1)} - i \frac{nr-1}{nr+1} 1_{V^{(1)}}, \quad H_0 = H'_0 + \sum_{i=2}^t 1_{V^{(i)}},$$

we can check that I and H_0 satisfy the condition (1.2). The corresponding maximal compact subgroup K in $G_{\mathbb{R}}$ can be written in the form $K = K_1 \times \prod_{i=2}^t SU(V^{(i)}, h^{(i)})$ where $K_1 \subset G_1 := SU(V^{(1)}, h^{(1)})$ is the maximal compact subgroup corresponding to H_0' .

Let $\mathfrak{g}_1, \mathfrak{k}_1$ denote the Lie algebras of G_1, K_1 , and \mathfrak{p} the orthogonal complement of \mathfrak{k}_1 in \mathfrak{g}_1 . Then we have the decompositions

$$\mathfrak{g}_1 = \mathfrak{k}_1 \oplus \mathfrak{p}, \quad \mathfrak{g}_{\mathbb{R}} = \mathfrak{k} \oplus \mathfrak{p},$$

and an isomorphism

$$\mathcal{D}(V^{(1)}, h^{(1)}) \cong G_{\mathbb{R}}/K \cong G_1/K_1.$$

We have the expression

$$W^{\sigma_i} = U_{\mathbb{C}}^{(i)} \otimes_{\mathbb{C}} V_{\mathbb{C}}^{(i)}$$

as in (4.20), and in this case, we have the decomposition

$$V^{(1)} \otimes_{\mathbb{R}} \mathbb{C} = V_{\mathbb{C}}^{(1)} \oplus \overline{V_{\mathbb{C}}^{(1)}}.$$

We may assume that the natural projection $V^{(1)} \rightarrow V_{\mathbb{C}}^{(1)}$ becomes a \mathbb{C} -linear isomorphism. Then if we set $V_{\mathbb{C}}^{(1)\pm} = V_{\mathbb{C}}^{(1)}(\pm i, I_{(1)})$, we have $\dim V_{\mathbb{C}}^{(1)+} = nr - 1$, $\dim V_{\mathbb{C}}^{(1)-} = 1$, and $W^{\sigma_1\pm} = U_{\mathbb{C}}^{(1)} \otimes V_{\mathbb{C}}^{(1)\pm}$. From the description as above, the homomorphism (5.17) of representation of K is compatible with decomposition (5.10) and the (σ_1) -part of the homomorphism is given by

$$(5.19) \quad \begin{aligned} W^{\sigma_1,+} &\longrightarrow \mathfrak{p}^- \otimes W^{\sigma_1,-} \\ &\cong U_{\mathbb{C}}^{(1)} \otimes [V_{\mathbb{C}}^{(1)+} \longrightarrow \mathfrak{p}^- \otimes V_{\mathbb{C}}^{(1)-}]. \end{aligned}$$

(5.20) Lemma. *The homomorphism (5.19) of the representations of K and K_1 is an isomorphism.*

PROOF: It suffices to show that $V_{\mathbb{C}}^{(1)+} \rightarrow \mathfrak{p}^- \otimes V_{\mathbb{C}}^{(1)-}$ is an isomorphism of K_1 -modules. Since $V_{\mathbb{C}}^{(1)+}$ and $\mathfrak{p} \otimes V_{\mathbb{C}}^{(1)-}$ are irreducible representations of K_1 of dimension $nr - 1$ and the homomorphism is not trivial, it must be an isomorphism.²

The following corollary shows Theorem (5.6) for the case (5.3).

(5.21) Corollary. *In case (5.3), we have*

$$H^1(M, \mathbf{W}^{\sigma_1})^{0,0} = 0,$$

so in particular $H^1(M, \mathbf{W}_{\mathbb{Q}})^{0,0} = 0$.

PROOF: Let ∇_{σ_i} denote the Gauss-Manin connection restricted to \mathbf{W}^{σ_i} . Then the corresponding Gauss-Manin complex

$$\nabla_{\sigma_1} : \mathcal{F}_{\sigma_1}^{-1} \longrightarrow \Omega_M^1 \otimes Gr_{\mathcal{F}_{\sigma_1}}^{-1}$$

²Considering the Harish-Chandra embedding $(I)_{nr-1,1} \hookrightarrow \mathbf{P}_{\mathbb{C}}^{nr-1}$, we can easily see that $\mathfrak{p}^- \cong V_{\mathbb{C}}^{(1)+} \otimes (V_{\mathbb{C}}^{(1)-})^*$

is induced by the homomorphism (5.19). Then by (5.20), this ∇_{σ_1} becomes an isomorphism. Hence we have $H^1(\mathbf{W}^{\sigma_1})^{0,0} \cong \mathbf{H}^1(\nabla_{\sigma_1}) = 0$. The last assertion follows from this and Lemma (5.13).

(5.22) Remark. If $t \geq 2$ and $H^1(M, \mathbf{W}_{\mathbb{C}}) \neq 0$, we can show that $H^1(M, \mathbf{W}^{\sigma_i})^{0,0} \neq 0$ for $i \geq 2$. Therefore from the example with non-vanishing $H^1(M, \mathbf{W}_{\mathbb{C}})$ mentioned in (5.1), we have examples with non-vanishing $H^1(M, \mathbf{W}_{\mathbb{C}})^{0,0}$, but still we have (5.21).

(5.23) Proof of Theorem (5.6) in the case of (5.2).

In this case, F_1 is a totally real field, and D is a quaternion algebra over F_1 . We denote by $\sigma_i : F_1 \hookrightarrow \mathbb{C}$ the embedding which is the extension of τ_i . Since $G_{\mathbb{R}} = \prod_{i=1}^t SU(V^{(i)}, h^{(i)}) \cong SU_3(\mathbb{H}) \times SO_6(\mathbb{R}) \times \cdots \times SO_6(\mathbb{R})$, $(V^{(1)}, h^{(1)})$ is a left \mathbb{H} -module of rank 3 with a \mathbb{H} -skew-hermitian form $h^{(1)}$. Recall that the expression $W^{\sigma_1} \cong U_{\mathbb{C}}^{(1)} \otimes_{\mathbb{C}} V_{\mathbb{C}}^{(1)}$ as in (4.23). Let us take a complex structure $I_{(1)} \in \mathcal{D}(V^{(1)}, h^{(1)}) \cong (II)_3$ and define $V_{\mathbb{C}}^{(1)\pm} = V_{\mathbb{C}}^{(1)}(\pm i, I_{(1)})$. Then we have the decomposition $V_{\mathbb{C}}^{(1)} = V_{\mathbb{C}}^{(1)+} \oplus V_{\mathbb{C}}^{(1)-}$. Setting $H_0 = (1/2)I_{(1)} + \sum_{i=2}^t 1_{V^{(i)}}$, we obtain the associated maximal compact subgroup $K = K_1 \times \prod_{i=2}^t SU(V^{(i)}, h^{(i)})$ of $G_{\mathbb{R}} = G_1 \times \prod_{i=2}^t SU(V^{(i)}, h^{(i)})$. Then as in (5.19), we have the homomorphism of representations of K and K_1 :

$$(5.24) \quad \begin{aligned} W^{\sigma_1, +} &\longrightarrow \mathfrak{p}^- \otimes W^{\sigma_1, -} \\ &\cong U_{\mathbb{C}}^{(1)} \otimes [V_{\mathbb{C}}^{(1)+} \longrightarrow \mathfrak{p}^- \otimes V_{\mathbb{C}}^{(1)-}] \end{aligned}$$

In this case, we have the isomorphism $SU(3, 1, \mathbb{C}) \cong SU_3(\mathbb{H})^-$, which is induced as follows. Let (T, h) be a complex vector space of dimension 4 with a hermitian form h of signature (3,1), and set $G = SU(T, h) \cong SU(3, 1, \mathbb{C})$. Let $I' \in \mathcal{D}(T, ih)$, and set $T^{\pm} = T(\pm i, I')$. Note that $\dim T^+ = 3$ and $\dim T^- = 1$. Then the space $\Lambda^2 T$ has a hermitian form h' induced by h , and the decomposition

$$\Lambda^2 T = \Lambda^2 T^+ \oplus (T^+ \otimes T^-)$$

corresponds to an element $I'' \in \mathcal{D}(\Lambda^2 T, h')$. It is known that $\mathcal{D}(\Lambda^2 T, ih') \cong (II)_3$ and the correspondence $T^+ \mapsto \Lambda^2 T^+$ induces an isomorphism $(I)_{3,1} \cong (II)_3$ (cf. §5, IV, [S1]), which can be lifted to a group isomorphism $SU(3, 1, \mathbb{C}) \cong SU_3(\mathbb{H})^-$. Thus the homomorphism $V_{\mathbb{C}}^{(1)+} \longrightarrow \mathfrak{p}^- \otimes V_{\mathbb{C}}^{(1)-}$ in (5.24) is isomorphic to

$$(5.25) \quad \Lambda^2 T^+ \longrightarrow \mathfrak{p}^- \otimes (T^+ \otimes T^-)$$

as a homomorphism of representation of K_1 (and K). Since we have an isomorphism $\mathfrak{p}^- \cong T^+ \otimes (T^-)^*$ as K_1 -modules (cf. (5.20)), the homomorphism (5.25) is isomorphic to

$$(5.26) \quad \Lambda^2 : \Lambda^2 T^+ \longrightarrow T^+ \otimes T^+.$$

Hence it is trivial that the homomorphism Λ^2 is injective and

$$\text{coker}(\Lambda^2) \cong S^2(T^+).$$

Let \mathcal{T} denote the locally free sheaf on M corresponding to the representation T^+ . Then, from (5.24), we have the isomorphism

$$(5.27) \quad \text{coker} \nabla_{\sigma_1} \cong U_{\mathbb{C}}^{(1)} \otimes S^2(\mathcal{T}).$$

Now we have the following result which implies Theorem (5.6) in the case (5.2).

(5.28) Proposition. *In the case (5.2), we have*

$$H^1(M, \mathbf{W}^{\sigma_1})^{0,0} = 0.$$

PROOF: Since from (5.27)

$$H^1(\nabla_{\sigma_1}) \cong H^0(\text{coker}(\nabla_{\sigma_1})) \cong U_{\mathbf{C}}^{(1)} \otimes H^0(M, S^2(\mathcal{T})),$$

we only have to show that $H^0(M, S^2(\mathcal{T})) = 0$. Let $\mathbf{T}_{\mathbf{C}}$ denote the local system on M induced by \mathcal{T} . Since we have the natural inclusion $\mathcal{T} \hookrightarrow \mathcal{O}_M(\mathbf{T}_{\mathbf{C}})$, we also have the inclusion

$$(5.29) \quad H^0(M, S^2(\mathcal{T})) \hookrightarrow H^0(M, S^2(\mathbf{T}_{\mathbf{C}})).$$

Then since the right hand side of (5.29) vanishes by Theorem (4.13), we have the assertion.

(5.30) Proof of Theorem (5.6) in the case (5.4).

In this case, we always have $G_{\mathbb{R}} \cong G_1 \times K_2 \times \cdots \times K_t$ where $G_1 \cong SL_2(\mathbb{R}) \cong Sp_1(\mathbb{R}) \cong SU(1, 1)$ and K_i are compact. We also have an expression $W^{\sigma_1} \cong U_{\mathbf{C}}^{(1)} \otimes V_{\mathbf{C}}^{(1)}$ where $V_{\mathbf{C}}^{(1)}$ is a complex irreducible representation of $SL_2(\mathbb{R})$ and $U_{\mathbf{C}}^{(1)}$ is a trivial representation. Then since $M = \Gamma \backslash \mathcal{D}$ is compact, we can apply the result in [Z2] ((5.33), Example) to deduce that

$$H^1(M, \mathbf{W}^{\sigma_1})^{0,0} = 0.$$

Hence, as before, we have the assertion.

§6 \mathbb{R} -RANK 1 AND Γ NON-COCOMPACT.

(6.1) Let $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ be a standard \mathbb{Q} -symplectic representation, $W_{\mathbb{Z}} \subset W_{\mathbb{Q}}$ a \mathbb{Z} -lattice, $\Gamma \subset G_{W_{\mathbb{Z}}} \subset G_{\mathbb{R}}$ a torsion free arithmetic group. In this section, we assume that $rk_{\mathbb{R}} G_{\mathbb{R}} = 1$ and $\Gamma \subset G_{\mathbb{R}}$ is not cocompact. Again, we will not deal with the case **(R2, -1)**, $\mathbf{n} = 2$. If $\dim \mathcal{D} = 1$, we can deduce the finiteness results from Zucker's results in [Z1] (see remark (6.30)). Hence we will assume that $\dim \mathcal{D} > 1$ unless we state otherwise.

We only have to consider the following cases:

$$(6.2) \quad \text{Case (R2, -1), } \mathbf{n} = 3 \quad G_{\mathbb{R}} \cong SU_3(\mathbb{H})^- \cong SU(3, 1, \mathbb{C}),$$

$$(6.3) \quad \text{Case (C)} \quad G_{\mathbb{R}} \cong SU(nr - 1, 1, \mathbb{C}).$$

In the above cases, the bounded symmetric domain $\mathcal{D} \simeq G_{\mathbb{R}}/K$ is isomorphic to the m -dimensional unit ball $B^m \subset \mathbb{C}^m$ for some $m \geq 1$. Since $\Gamma \subset G_{\mathbb{R}}$ is a torsion free arithmetic subgroup of $G_{\mathbb{R}}$, $M = \Gamma \backslash \mathcal{D}$ is a smooth complex manifold with a finite invariant measure, but, by assumption, is not compact. The Baily-Borel-Satake compactification M^* of M

can be obtained by adding a finite number of cusps $\{p_i\}$ to M . Note that M^* is projective. Moreover, according to Hemperly [He], a resolution of singularities $\pi : \overline{M} \rightarrow M^*$ is obtained by the blowing up of the cusps $\{p_i\}$, and the inverse images $D_i = \pi^{-1}(p_i)$ are abelian varieties.

(6.4) Let $(W_{\mathbb{Q}}, \rho_{\mathbb{Q}}, A_{\mathbb{Q}}, I)$ be a standard \mathbb{Q} -symplectic representation in the case (6.2) or (6.3), $D, \iota, F_1, V, U, h, h'$ be as in (3.17). Let $f : \mathcal{X} \rightarrow M$ denote the Kuga fiber space associated to the above representation and the lattice $W_{\mathbb{Z}}$ in (6.1). Then, as in (2.29), we have the exact sequence

$$(6.5) \quad 0 \rightarrow W_{\mathbb{Z}} \rightarrow Gr_{\mathcal{F}}^{-1} \rightarrow \mathcal{O}_M^{an}(\mathcal{X}) \rightarrow 0.$$

Let us assume that the local monodromy around each D_i is unipotent. This is always possible if one replaces Γ with a normal subgroup Γ' of finite index. Then we can extend the abelian scheme $f : \mathcal{X} \rightarrow M$ to a semi-abelian scheme $\overline{f} : \overline{\mathcal{X}} \rightarrow \overline{M}$ as follows. Let $\mathcal{W} := \mathcal{O}_M \otimes W_{\mathbb{C}}$. Then we have the Gauss-Manin connection $\nabla : \mathcal{W} \rightarrow \Omega_M^1 \otimes \mathcal{W}$ which is integrable. Let $\overline{\mathcal{W}}$ denote the Deligne canonical extension of \mathcal{W} which is a locally free $\mathcal{O}_{\overline{M}}$ -module with a logarithmic connection $\overline{\nabla} : \overline{\mathcal{W}} \rightarrow \Omega_{\overline{M}}^1(\log D) \otimes \overline{\mathcal{W}}$ such that $Res_{D_i}(\overline{\nabla})$ is nilpotent (see [D1]). Let $j : M \hookrightarrow \overline{M}$ denote the inclusion. We set:

$$(6.6) \quad \overline{\mathcal{F}}^p := j_* \mathcal{F}^p \cap \overline{\mathcal{W}}.$$

By the nilpotent orbit theorem [Sc, (4.12)], these are locally free subsheaf of $\overline{\mathcal{W}}$. As in [Z3], we can obtain a semi-abelian scheme $\overline{f} : \overline{\mathcal{X}} \rightarrow \overline{M}$ which is an extension of the original abelian scheme f and fits into the following sheaf exact sequence

$$(6.7) \quad 0 \rightarrow j_* W_{\mathbb{Z}} \rightarrow Gr_{\overline{\mathcal{F}}}^{-1} \rightarrow \mathcal{O}_{\overline{M}}^{an}(\overline{\mathcal{X}}) \rightarrow 0.$$

(6.8) **Proposition.** *Under the notations and the assumptions as above, the natural restriction map (see (2.4))*

$$r : H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{an}(\overline{\mathcal{X}})) \rightarrow H^0(M, \mathcal{O}_M(\mathcal{X})) \cong MW(\mathcal{X}/M)$$

is injective and has a finite cokernel.

PROOF: First, I remark that all sections $H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{an}(\overline{\mathcal{X}}))$ is algebraic, so r is well-defined. The injectivity of r is obvious. To prove r has a finite cokernel, we first remark that we can construct the Néron model $N(f) : N(\mathcal{X}) \rightarrow \overline{M}$ of $f : \mathcal{X} \rightarrow M$ which has the following properties.

- (i) $N(f) : N(\mathcal{X}) \rightarrow \overline{M}$ is a group scheme over \overline{M} which is an extension of f .
- (ii) Let $Y \rightarrow \overline{M}$ be a smooth morphism and $\phi : Y \cdots \rightarrow N(\mathcal{X})$ a rational map over \overline{M} . Then ϕ extends to a morphism $\phi : Y \rightarrow N(\mathcal{X})$.
- (iii) The semi-abelian scheme $\overline{\mathcal{X}}$ is a connected component of $N(\mathcal{X})$, i.e. $\overline{\mathcal{X}}$ is a subgroup scheme of $N(\mathcal{X})$ such that for each closed point $p \in \overline{M}$, $\overline{\mathcal{X}}_p$ is the connected component of $N(\mathcal{X})_p$ containing the identity.

Moreover there exists a projective manifold $\overline{N(\mathcal{X})}$ containing $N(\mathcal{X})$ as a Zariski open set and a projective morphism $\overline{N(f)} : \overline{N(\mathcal{X})} \rightarrow \overline{M}$ which is an extension of $N(f)$ such that $N(\mathcal{X})$ is the maximal open subset of $\overline{N(\mathcal{X})}$ where $\overline{N(f)}$ is smooth. The existence of the above Néron model $N(\mathcal{X})$ and its projective completion is proved as follows. It suffices to show that the existence of them over a some tubular neighborhood U of an irreducible component D_i of $D = \sum_{i=1}^i D_i$. For each point $p \in D_i$, we can take a neighborhood U_p which is isomorphic to $\Delta^n = \{(z_i) \in \mathbb{C}^n \mid |z_i| < 1\}$ and $U_p \cap D_i = \{z_1 = 0\}$. Then the Néron model of $f|_{U_p - D_i} : \mathcal{X}|_{U_p - D_i} \rightarrow U_p - D_i \cong \Delta^* \times \Delta^{n-1}$ can be constructed as in [A]. Since the Néron model has a uniqueness property, such local Néron models can be patched together and one gets a global Néron model over the tubular neighborhood U of D_i .

Now we prove that the finiteness of cokernel of r . Every algebraic section $s : M \rightarrow \mathcal{X}$ defines a rational map $\bar{s} : \overline{M} \dashrightarrow \overline{N(\mathcal{X})}$. Considering locally around D , we can show that \bar{s} must actually map to $N(\mathcal{X})$. Then by the property (ii), \bar{s} is a morphism $\bar{s} : \overline{M} \rightarrow N(\mathcal{X})$ and so it is a section of $N(f)$. This shows that $H^0(M, \mathcal{O}_M(\mathcal{X}))$ is isomorphic to $H^0(\overline{M}, \mathcal{O}_{\overline{M}}(N(\mathcal{X})))$, i.e. the group of sections of $N(f) : N(\mathcal{X}) \rightarrow \overline{M}$. Then the cokernel of r is a subgroup of $H^0(\overline{M}, N(\mathcal{X})/\overline{\mathcal{X}})$, where $N(\mathcal{X})/\overline{\mathcal{X}}$ is a finite group scheme over D . Since the fiber $N(\mathcal{X})/\overline{\mathcal{X}}$ over each component D_i is a finite group, $H^0(D, N(\mathcal{X})/\overline{\mathcal{X}})$ is also a finite group, and this completes the proof.

(6.9) Hodge theory for $j_* \mathbf{W}_{\mathbb{C}}$.

Let $(\mathbf{W}_{\mathbf{z}}, \mathbf{A}, \mathcal{F})$ be the VPHS (see (1.10)) over $\Gamma \backslash \mathcal{D}$ of weight -1 associated to the symplectic representation as in (6.4). As in (6.1), there exists a projective manifold \overline{M} and an inclusion $j : M \hookrightarrow \overline{M}$ such that $D = \overline{M} - M$ is a union of smooth hypersurfaces each of which is isomorphic to an abelian variety.

It is known that the cohomology group $H^i(\overline{M}, j_* \mathbf{W}_{\mathbf{z}})$ has a polarized pure Hodge structure of weight $i-1$. This fact can be considered as a generalization of Zucker's results in [Z1] to the cases of the higher dimensional bases, and was proved by Cattani-Kaplan-Schmid [C-K-S] and Kashiwara-Kawai [K-K] as follows.

One can see that M admits a complete Kähler metric with Poincaré singularities along D . In the above case, $j_* \mathbf{W}_{\mathbb{C}}$ equals the intersection complex $\mathcal{IC}(\overline{M}, \mathbf{W}_{\mathbb{C}})$ of Deligne and Goresky-MacPherson. Then they showed that $\mathcal{IC}(\overline{M}, \mathbf{W}_{\mathbb{C}})$ is quasi-isomorphic to the L^2 -complex $\mathcal{L}_{(2)}(M, \mathbf{W}_{\mathbb{C}})$ with respect to the above Kähler metric on M and the Hodge metric on $\mathbf{W}_{\mathbb{C}}$.³ Therefore we have the isomorphisms

$$H^i(\overline{M}, j_* \mathbf{W}_{\mathbb{C}}) \cong \mathrm{IH}^i(\overline{M}, \mathbf{W}_{\mathbb{C}}) \cong H_{(2)}^i(M, \mathbf{W}_{\mathbb{C}}).$$

Each element of L^2 -cohomology group can be represented by a harmonic form, so by using the Kähler identity between the Laplacians (cf. [Z1]), we obtain a Hodge decomposition of the cohomology group. (See also [ShzY].)

(6.10) Mixed Hodge theory.

³ Actually, they proved this result for the more general case where $\overline{M} - M$ is a divisor with normal crossings.

We will recall a more explicit description of the Hodge structure on $H^i(\overline{M}, j_* \mathbf{W}_{\mathbf{C}})$ in our case following [ShzY] (cf. [Z1]). In order to see this, we shall introduce the mixed Hodge structure on $H^i(M, \mathbf{W}_{\mathbf{Q}})$.

Since we have $H^i(M, \mathbf{W}_{\mathbf{Q}}) \cong H^i(\overline{M}, \mathbf{R}j_* \mathbf{W}_{\mathbf{Q}})$, we have the long exact sequence of cohomology groups

$$(6.11) \quad \longrightarrow H^i(\overline{M}, j_* \mathbf{W}_{\mathbf{Q}}) \longrightarrow H^i(M, \mathbf{W}_{\mathbf{Q}}) \longrightarrow H^{i-1}(\overline{M}, R^1 j_* \mathbf{W}_{\mathbf{Q}}) \xrightarrow{\delta}$$

which comes from the Leray spectral sequence for the inclusion $j : M \hookrightarrow \overline{M}$. Then it is known that $H^i(M, \mathbf{W}_{\mathbf{Q}})$ and $H^{i-1}(\overline{M}, R^1 j_* \mathbf{W}_{\mathbf{Q}})$ has a mixed Hodge structure, which makes (6.11) an exact sequence of mixed Hodge structures.

There are a weight filtration $\{W\}$ on $H^i(M, \mathbf{W}_{\mathbf{Q}})$ and the Hodge filtration $\{F\}$ on $H^i(M, \mathbf{W}_{\mathbf{C}})$ such that for each k , $Gr_k^W(H^i(M, \mathbf{W}_{\mathbf{Q}}))$ with the induced Hodge filtration F forms a polarized (pure) Hodge structure. In our case, we have 3-step weight filtration $0 = W_{-1} \subset W_0 \subset W_1 \subset W_2 = H^i(M, \mathbf{W}_{\mathbf{Q}})$, such that

$$(6.12) \quad W_0(H^i(M, \mathbf{W}_{\mathbf{Q}})) \cong \text{Im}\{H^i(\overline{M}, j_* \mathbf{W}_{\mathbf{Q}}) \rightarrow H^i(M, \mathbf{W}_{\mathbf{Q}})\},$$

$$Gr_1^W \cong \ker\{H^{i-1}(D, P_0) \rightarrow H^{i+1}(\overline{M}, j_* \mathbf{W}_{\mathbf{Q}})\},$$

$$Gr_2^W \cong \ker\{H^{i-1}(D, P_1) \rightarrow H^{i+1}(\overline{M}, j_* \mathbf{W}_{\mathbf{Q}})\},$$

where the P_k 's denote the local systems on D which underlies VPHS coming from the limit Hodge structure along D . (See [ShzY], (3.1.4)).

One can show that there is a quasi-isomorphism $\mathbf{R}j_* \mathbf{W}_{\mathbf{C}} \cong \Omega_{\overline{M}}^i(\log D) \otimes \overline{\mathcal{W}}$ (cf. [ShzY], (3.1.1)). Hence we have an isomorphism $H^i(M, \mathbf{W}_{\mathbf{C}}) \cong \mathbf{H}^i(\overline{M}, \Omega_{\overline{M}}^i(\log D) \otimes \overline{\mathcal{W}})$. The Hodge filtration $\{F\}$ on the complex $K_{\mathbf{C}} \cong \Omega_{\overline{M}}^i(\log D) \otimes \overline{\mathcal{W}}$ can be defined by

$$(6.13) \quad F^p K_{\mathbf{C}} := \Omega_{\overline{M}}^i(\log D) \otimes \overline{\mathcal{F}}^{p-i},$$

and this induces a Hodge filtration on $H^i(M, \mathbf{W}_{\mathbf{C}})$. The spectral sequence induced by this filtration

$$(6.14) \quad E_1^{p,q} = \mathbf{H}^{p+q}(\overline{M}, Gr_F^p \Omega_{\overline{M}}^i(\log D)) \implies H^{p+q}(M, \mathbf{W}_{\mathbf{C}})$$

degenerates at E_1 .

(6.15) Now we restrict our attention to H^1 . From (6.11), one has the exact sequence of the mixed Hodge structures

$$(6.16) \quad 0 \longrightarrow H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{Q}}) \longrightarrow H^1(M, \mathbf{W}_{\mathbf{Q}}) \longrightarrow H^0(D, R^1 j_* \mathbf{W}_{\mathbf{Q}}) \xrightarrow{\delta}$$

From (6.13), we have the 3-step Hodge filtration $0 = F^2 \subset F^1 \subset F^0 \subset F^{-1}$ on $H^1(M, \mathbf{W}_{\mathbf{C}})$ whose successive quotients are:

$$(6.17) \quad H^{1,-1} = Gr_F^1 = F^1 = \mathbf{H}^1(0 \rightarrow \Omega_M^1(\log D) \otimes \overline{\mathcal{F}}^0 \rightarrow \Omega_M^2(\log D) \otimes Gr_{\overline{\mathcal{F}}}^{-1}),$$

$$(6.18) \quad H^{0,0} = Gr_F^0 = F^0/F^1 = \mathbf{H}^1(\overline{\mathcal{F}}^0 \rightarrow \Omega_M^1(\log D) \otimes Gr_{\overline{\mathcal{F}}}^{-1}),$$

$$(6.19) \quad H^{-1,1} = Gr_F^{-1} = F^{-1}/F^0 = \mathbf{H}^1(Gr_{\overline{\mathcal{F}}}^{-1}).$$

(6.20) Proposition. *Let us denote by $H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{C}})^{p,q}$ the (p,q) -component of the pure Hodge structure of $H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{C}})$. Then we have*

(i) *the isomorphism*

$$H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{C}})^{-1,1} \cong H^1(\overline{M}, Gr_{\overline{\mathcal{F}}}^{-1}),$$

(ii) *and the inclusion*

$$H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{C}})^{0,0} \hookrightarrow \mathbf{H}^1(\overline{\mathcal{F}}^0 \rightarrow \Omega_M^1(\log D) \otimes Gr_{\overline{\mathcal{F}}}^{-1}).$$

PROOF: These come from (6.18), (6.19) and the fact that (6.16) is an exact sequence of mixed Hodge structures.

(6.21) Now we have the following proposition which is a generalization of (2.27) (cf. (10.2) of [Z1]).

(6.22) Proposition. *Let $f : \mathcal{X} \rightarrow M$ be a Kuga fiber space as in (6.4) and $\overline{f} : \overline{\mathcal{X}} \rightarrow \overline{M}$ the extended semi-abelian scheme. Then we have an isomorphism*

$$H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{un}(\overline{\mathcal{X}})) \cong H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{Z}})^{0,0}.$$

Here we set $H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{Z}})^{0,0} \cong i^{-1}(H^{0,0})$ where $i : H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{Z}}) \rightarrow H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{C}})$ is the natural map.

PROOF: In this case, $H^0(\overline{M}, Gr_{\overline{\mathcal{F}}}^{-1}) = 0$, because $H^0(M, \mathbf{W}_{\mathbf{C}}) = 0$ by (2.12) and (4.5). Therefore, from (6.7), we have the long exact sequence

$$0 \longrightarrow H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{un}(\overline{\mathcal{X}})) \longrightarrow H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{Z}}) \xrightarrow{p} H^1(\overline{M}, Gr_{\overline{\mathcal{F}}}^{-1}).$$

which implies that

$$H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{un}(\overline{\mathcal{X}})) \cong \ker\{p : H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{Z}}) \rightarrow H^1(\overline{M}, Gr_{\overline{\mathcal{F}}}^{-1})\}.$$

Since $H^1(\overline{M}, Gr_{\overline{\mathcal{F}}}^{-1}) \cong H^{-1,1}$ by (6.20), the map p coincides with the composite of i and the projection from $H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{C}})$ to its $(-1,1)$ -part. Let us take an element $u \in \ker p$. Since $u \in H^1(\overline{M}, j_* \mathbf{W}_{\mathbf{Z}})$ is real and u has no $(-1,1)$ -component, it has also no $(1,-1)$ -component. Thus u is of type $(0,0)$, and conversely.

(6.23) Corollary. *Let $f : \mathcal{X} \rightarrow M$ be a Kuga fiber space as in (6.4). The Mordell-Weil group $MW(\mathcal{X}/M)$ is finite if and only if*

$$(6.24) \quad H^1(\overline{M}, j_* \mathbf{W}_{\mathbb{Q}})^{0,0} = 0.$$

PROOF: By Proposition (6.8), we only have to prove that the group $H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{un}(\overline{\mathcal{X}}))$ is finite. Since $H^1(\overline{M}, j_* \mathbf{W}_{\mathbb{Z}})^{0,0} \otimes \mathbb{Q} \cong H^1(\overline{M}, j_* \mathbf{W}_{\mathbb{Q}})^{0,0}$, Proposition (6.22) implies that the condition (6.24) is equivalent to the finiteness of $H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{un}(\overline{\mathcal{X}}))$.

(6.25) Theorem. *Let $f : \mathcal{X} \rightarrow M$ be the Kuga fiber spaces associated to the \mathbb{Q} -symplectic representation of type (6.2) or (6.3). Assume that $M = \Gamma \backslash \mathcal{D}$ is not compact. Then the Mordell-Weil group $MW(\mathcal{X}/M)$ is finite.*

PROOF: We first remark that we can replace $M = \Gamma \backslash \mathcal{D}$ with its finite unramified covering. So we may assume that local monodromies around the components of D are unipotent.

We first prove the case (6.3). We shall use the notation in (5.18). In this case, F_1 is a purely imaginary quadratic field over \mathbb{Q} , so denote by $\{\sigma, \bar{\sigma}\}$ the embedding of F_1 into \mathbb{C} . We have the decomposition

$$W_{\mathbb{C}} = W^{\sigma} \oplus W^{\bar{\sigma}}$$

where we put $W^{\sigma} := W_{\mathbb{Q}} \otimes_{F_1, \sigma} \mathbb{C}$. We also have the expression

$$W^{\sigma} = U_{\mathbb{C}} \otimes_{\mathbb{C}} V_{\mathbb{C}}$$

where $V_{\mathbb{C}}$ is an nr -dimensional \mathbb{C} -vector space which has a \mathbb{C} -symplectic form $h_{\mathbb{C}}$ such that the signature of $\sqrt{-1}h_{\mathbb{C}}$ is $(nr - 1, 1)$. As in (5.18), a complex structure $I \in \mathcal{D}$ defines a decomposition $V_{\mathbb{C}} = V_{\mathbb{C}}^{+} \oplus V_{\mathbb{C}}^{-}$ where $\dim V_{\mathbb{C}}^{+} = nr - 1$ and $\dim V_{\mathbb{C}}^{-} = 1$. And setting $W^{\sigma, \pm} = U_{\mathbb{C}} \otimes V_{\mathbb{C}}^{\pm}$, we have the homomorphism of K -module

$$(6.26) \quad \begin{aligned} W^{\sigma, +} &\longrightarrow \mathfrak{p}^{-} \otimes W^{\sigma, -} \\ &\cong U_{\mathbb{C}} \otimes [V_{\mathbb{C}}^{+} \longrightarrow \mathfrak{p}^{-} \otimes V_{\mathbb{C}}^{-}]. \end{aligned}$$

which induces the σ -part of the first Gauss-Manin complex on M

$$(6.27) \quad \nabla_{\sigma} : \mathcal{F}_{\sigma}^0 \longrightarrow \Omega_M^1 \otimes Gr_{\mathcal{F}_{\sigma}}^{-1},$$

where we set $\mathcal{F}_{\sigma}^p = \mathcal{O}_M(\mathbf{W}_{\mathbb{C}}^{\sigma}) \cap \mathcal{F}^p$. From Lemma (5.20), the homomorphism (6.26) is an isomorphism of K -modules and so the sheaf homomorphism (6.27) is also an isomorphism. Now let us write $\mathcal{D} = G_{\mathbb{R}}/K$. Since K is compact, $W^{\sigma, +}$, \mathfrak{p}^{-} and $W^{\sigma, -}$ admit $G_{\mathbb{R}}$ -invariant Hermitian metrics, which induce Hermitian metrics on the locally free sheaves \mathcal{F}_{σ}^0 , Ω_M^1 and $Gr_{\mathcal{F}_{\sigma}}^{-1}$ respectively. Note that on \mathcal{F}_{σ}^0 and $Gr_{\mathcal{F}_{\sigma}}^{-1}$ these metric are constant multiple of the metric induced by the original polarization \mathbf{A} . Let E be a any locally free sheaf on $M = \Gamma \backslash \mathcal{D}$ induced by a K -representation with an above hermitian metric h . In [Mum], Mumford showed that such a E admits a canonical extension \overline{E} to a smooth toroidal compactification \overline{M} in (6.1) such that h is a singular Hermitian metric *good* on \overline{M} . (For

the definition of *goodness of a singular Hermitian metric*, see [Mum] §1.) One can see that such canonical extensions of \mathcal{F}_σ^0 and $Gr_{\mathcal{F}_\sigma}^{-1}$ coincides with $\overline{\mathcal{F}}_\sigma^0$ and $Gr_{\overline{\mathcal{F}}_\sigma}^{-1}$ defined in (6.4), that is, those induced from the Deligne's canonical extension. (For the proof of this fact, see [H], Theorem 4.2.) Moreover, the canonical extension of Ω_M^1 in the sense of Mumford is $\Omega_M^1(\log D)$. Therefore, by uniqueness of canonical extensions, the isomorphism (6.27) is extended to the isomorphism

$$(6.28) \quad \overline{\nabla}_\sigma : \overline{\mathcal{F}}_\sigma^0 \longrightarrow \Omega_M^1(\log D) \otimes Gr_{\overline{\mathcal{F}}_\sigma}^{-1},$$

over \overline{M} . Then by (ii), Proposition (6.22), we have $H^1(\overline{M}, j_* \mathbf{W}_\mathbb{C}^g)^{0,0} = 0$. From this, by the same argument as in Lemma (5.13), we deduce the vanishing condition (6.24), which implies the finiteness of the Mordell-Weil group.

Next we will deal with the case (6.2). In this case, $F_1 = \mathbb{Q}$ and $G_\mathbb{R} \cong SU_3(\mathbb{H})^- \cong SU(3, 1, \mathbb{C})$. We use the same notation as in (5.23). By the same reason as in the case (6.3), we only have to show that $H^1(\overline{M}, \overline{\nabla}) = 0$ where $\overline{\nabla}$ is the canonical extension of the Gauss-Manin complex. Over M , we have the isomorphism (5.27), so again by the uniqueness of the canonical extension, we have the isomorphism

$$(6.29) \quad \text{coker } \overline{\nabla} \cong U_\mathbb{C} \otimes S^2(\overline{\mathcal{T}})$$

where $\overline{\mathcal{T}}$ is the canonical extension of the sheaf \mathcal{T} (see (5.23)) to \overline{M} . As in proof of Proposition (5.28), we only have to show that $H^0(\overline{M}, S^2(\overline{\mathcal{T}})) = 0$. As in (5.29), we have the inclusion

$$H^0(\overline{M}, S^2(\overline{\mathcal{T}})) \hookrightarrow H^0(\overline{M}, S^2(\overline{\mathbf{T}}_\mathbb{C}))$$

where $\overline{\mathbf{T}}_\mathbb{C}$ is the canonical extension of $\mathbf{T}_\mathbb{C}$. We have the isomorphism $H^0(\overline{M}, S^2(\overline{\mathbf{T}}_\mathbb{C})) \cong H^0(M, S^2(\mathbf{T}_\mathbb{C}))$ (see (3.1.1), [ShzY]), and by (4.13) $H^0(M, S^2(\mathbf{T}_\mathbb{C})) = 0$. So we have the desired assertion.

(6.30) Remark. If $\dim \mathcal{D} = 1$ and $M = \Gamma \backslash \mathcal{D}$ is not compact, the finiteness follows from the result in [Z1]. Let $f : \mathcal{X} \longrightarrow M$ be a Kuga fiber space and $\overline{f} : \overline{\mathcal{X}} \longrightarrow \overline{M}$ the semi-abelian scheme in (6.4). By (6.8), we only have to prove that $H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{an}(\overline{\mathcal{X}}))$ is finite. Then by (6.22) (cf. Corollary (10.2) in [Z1]), we have $H^0(\overline{M}, \mathcal{O}_{\overline{M}}^{an}(\overline{\mathcal{X}})) \cong H^1(\overline{M}, j_* \mathbf{W}_\mathbb{Z})^{0,0}$. Then Lemma (12.4) in [Z1] says that $H^1(\overline{M}, j_* \mathbf{W}_\mathbb{C})^{0,0} = 0$, and hence the Mordell-Weil group is finite.

REFERENCES

- [A] Artin, *Néron Models*, "Arithmetic Geometry, Cornell, G., Silverman, J. H. eds.," Springer-Verlag, 1986, pp. 213-230.
- [Ba-B] Baily, Jr. W. L., Borel, A., *Compactification of arithmetic quotients of bounded symmetric domains*, Ann. of Math. **84** (1966), 442-528.
- [B] Borel, A., *Regularization theorems in Lie algebra cohomology. Applications*, Duke Math. J. **50** (1983), 602-624.
- [B-C] Borel, A., Casselman, W., *L_2 -cohomology of locally symmetric manifolds of finite volume*, Duke Math. J. **50**, (1983), 625-647.

- [B-W] Borel, A., Wallach, N., "Continuous cohomology, discrete subgroups and representations of reductive groups," Ann. of Math. Studies, Princeton Univ. Press, 1980.
- [C-K-S] Cattani, E., Kaplan, A., Schmid, W., *L_2 and intersection cohomologies for a polarized variation of Hodge structure*, Invent. math. **87**, 217-252.
- [D1] Deligne, P., *Equation différentielles à points singuliers réguliers*, Lecture Notes in Math. **163** (1970), Springer-Verlag, Berlin-Heidelberg-New York.
- [D2] Deligne, P., *Théorie de Hodge, II*, Publ. Math. IHES, **40** (1971), 5-57.
- [G-M] Goresky, M., MacPherson, R., *Intersection homology II*, Invent. math. **72** (1983), 77-129.
- [H] Harris, M., *Functorial properties of toroidal compactifications of locally symmetric varieties*, Proc. London Math. Soc. **59** (1989), 1-22.
- [He] Hemperly, C. J., *The parabolic contribution to the number of linear independent automorphic forms on certain bounded domain*, Amer. J. Math. **94** (1972), 1078-1100.
- [K-K] Kashiwara, K., Kawai, T., *The Poincaré lemma for variations of polarized Hodge structures*, Publ. R.I.M.S. Kyoto University **23** (1987), 345-407.
- [L] Looijenga, E., *L^2 -cohomology of locally symmetric varieties*, Compositio Math. **67** (1988), p. 3-20.
- [M-M] Matsushima, Y., Murakami, S., *On vector bundle valued harmonic forms and automorphic forms on symmetric Riemannian manifolds*, Ann. of Math. **78** (1963), 365-416.
- [Mum] Mumford, D., *Hirzebruch's Proportionality Theorem in the Non-Compact Case*, Invent. math. **42** (1977), 239-272.
- [S1] Satake, I., "Algebraic structure of symmetric domains, Publication of the Math. Soc. of Japan, 14," Iwanami Shoten and Princeton University Press, 1980.
- [S2] Satake, I., *Symplectic representations of algebraic groups satisfying a certain analyticity condition*, Acta Math. **117** (1967), 215-279.
- [Sa-St] Saper, L., Stern, M., *L_2 -cohomology of arithmetic varieties*, Ann. of Math. **132** (1990), 1-69.
- [Sc] Schmid, W., *Variation of Hodge structure: The singularities of the period mapping*, Invent. Math. **22** (1973), 211-319.
- [Sh1] Shimura, G., *On analytic families of polarized abelian varieties and automorphic functions*, Ann. of Math. **78** (1963), 149-192.
- [Sh2] Shimura, G., *Moduli and fiber systems of abelian varieties*, Ann. of Math. **83** (1966), 294-338.
- [Sh3] Shimura, G., *Discontinuous groups and abelian varieties*, Math. Ann. **168** (1967), 171-199.
- [Sd] Shioda, T., *On elliptic modular surfaces*, J. Math. Soc. Japan **24** (1972), 20-59.
- [Shz] Shimizu, H., *On discontinuous groups operating on product of the upper half planes*, Ann. of Math. **77** (1963), 33-71.
- [ShzY] Shimizu, Y., *Mixed Hodge Structures on Cohomologies with Coefficients in a Polarized Variation of Hodge Structure*, Advanced Studies in Pure Math. **10** (1985), 695-716.
- [Si1] Silverberg, A., *Mordell-Weil groups of generic Abelian varieties*, Invent. Math. **81** (1985), 71-106.

- [Si2] Silverberg, A., *Cohomology of fiber systems and Mordell-Weil groups of Abelian Varieties*, Duke Math. J. **56** (1988), 41-46.
- [Si3] Silverberg, A., *Mordell-Weil groups of generic abelian varieties in the unitary case*, Proc. of the A.M.S. **104** (1988), 723-728.
- [W] Weil, A., "Adeles and algebraic groups,," Notes by M. Demazure and T. Ono, I. A. S., Princeton, N. J., 1961.
- [Z1] Zucker, S., *Hodge theory with degenerating coefficients: L_2 -cohomology in the Poincaré metric*, Annals of Math. **109** (1979), 415-476.
- [Z2] Zucker, S., *Locally homogeneous variations of Hodge structure*, L'Ens. Math. **27** (1981), 243-276.
- [Z3] Zucker, S., *Generalized Intermediate Jacobians and the Theorem on Normal Functions*, Invent. math. **33** (1976), 185-222.

Department of Mathematics, Faculty of Science, Kyoto University, Kyoto, 606, Japan