


QUADRATIC HOMOLOGY

Hans-Joachim Baues

Max-Planck-Institut
für Mathematik
Gottfried-Claren-Str. 26
53225 Bonn

GERMANY

MPI 96-15



QUADRATIC HOMOLOGY

HANS-JOACHIM BAUES

ABSTRACT. We describe axioms for a ‘quadratic homology theory’ which generalize the classical axioms of homology. As examples we consider quadratic homology theories induced by 2-excisive homotopy functors in the sense of Goodwillie and the homology of a space with coefficients in a square group which generalizes the homology of a space with coefficients in an abelian group.

CONTENTS

§ 0 Introduction	1
§ 1 CW-spaces, CW-pairs and spectra	4
§ 2 Axioms of homology	5
§ 3 Quadratic homology	10
§ 4 Homotopy functors	17
§ 5 Homotopy functors induced by endofunctors of the category of groups	22
§ 6 Square-homology	26
§ 7 Square-homology with coefficients in a quadratic \mathbb{Z} -module	32
§ 8 Stable square homology and Steenrod squares	37
§ 9 The squaring operation Sq_M^2	42
Appendix A: A criterion for 2-excisive functors	46
Appendix B: The homology spectral sequence	48
Appendix C: A spectral sequence for square homology	50
References	52

§ 0 INTRODUCTION

The development of algebraic topology was profoundly affected by the notion of homology. Originally homology had coefficients in abelian groups and Eilenberg-Steenrod described the axioms of such an *ordinary homology theory*. Somewhat later important examples of generalized homology theories were found which led to the notion of homology with coefficients in a spectrum. The spectrum-homology can also be described as a *homology theory* satisfying all Eilenberg-Steenrod axioms except the dimension axiom concerning the value of the homology on spheres. In

1991 *Mathematics Subject Classification*. 55N35, 55Q70, 55S20.

Key words and phrases. Quadratic functors, Goodwillie calculus, Steenrod squares, EHP-sequence.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$

fact this value characterizes a homology theory in the sense that a natural transformation of homology theories which is an isomorphism on spheres is also an isomorphism on all finite CW-complexes.

In his recent work on the calculus of homotopy functors Goodwillie observed that a spectrum is equivalent to a linear homotopy functor and spectrum-homology of a space X can be equivalently described by the homotopy groups

$$(1) \quad H_n(X, L) = \pi_n L(X)$$

where L is a linear homotopy functor. We therefore call spectrum-homology also a linear homology theory. The *non-linear homology theories* are then obtained by the homotopy groups

$$(2) \quad H_n(X, D) = \pi_n D(X)$$

where D is any homotopy functor. Hence we again generalize homology by choosing now homotopy functors as coefficients. Clearly this is a very far reaching generalization since in particular homotopy groups of a space are the homology groups with coefficients in the identity functor. It is an old problem to find axioms which characterize the theory of homotopy groups in a similar way as ordinary homology theory is characterized by the Eilenberg-Steenrod axioms. However the identity functor is not linear but is still an analytic functor in the sense that there is a Taylor tower, $n \geq 1$,

$$(3) \quad X \rightarrow P_n(X) \rightarrow P_{n-1}(X) \rightarrow \dots \rightarrow P_2(X) \rightarrow P_1(X) = Q(X)$$

approximating X . Here P_1 is linear, P_2 is quadratic and more generally P_n is a reduced and n -excisive homotopy functor; compare [19, 20]. The linear functor P_1 yields the homology theory of *stable homotopy groups* $\pi_n^S(X) = H_n(X, P_1)$.

In this paper we study as a first step outside the linear world the *quadratic homology theories* $H_n(X, Q)$ obtained by a quadratic homotopy functor Q . We introduce axioms of a quadratic homology theory such that $H_n(X, Q)$ satisfies these axioms. As in the classical case the axioms characterize a quadratic homology theory in the sense that a natural transformation between theories which is an isomorphism on spheres is also an isomorphism on all finite CW-complexes. We deduce from the quadratic homology axioms various facts like the general EHP-sequence in § 3. For example quadratic homotopy groups

$$(4) \quad \pi_n^Q(X) = H_n(X, P_2)$$

defined by P_2 in the Taylor tower (3) satisfy all the axioms. This is the quadratic analogue of stable homotopy groups.

Let \mathbf{Gr} be the category of groups. Then any group functor $F : \mathbf{Gr} \rightarrow \mathbf{Gr}$ induces a homotopy functor F_{\sharp} which carries connected spaces to connected spaces. If F is

linear then $F(G) = G^{ab} \otimes A$ is given by an abelian group A and ordinary homology with coefficients in A can be described by

$$(5) \quad \tilde{H}_n(X, A) = H_n(X, F_{\sharp}).$$

If the group functor F is quadratic then we know by [10] that $F(G) = G \otimes M$ is given by a *square group* M which is the quadratic analogue of an abelian group. We prove that then F_{\sharp} is a quadratic homotopy functor. Hence the *square-homology*

$$H_n(X, M) = H_n(X, F_{\sharp})$$

with coefficients in a square group M is a quadratic homology theory which generalizes ordinary homology. Using the Taylor tower of F_{\sharp} we obtain the linearization

$$(7) \quad F_{\sharp} = P_2 F_{\sharp} \rightarrow P_1 F_{\sharp} = F_{\sharp}^{lin}$$

which gives us a linear homology theory

$$(8) \quad H_n^S(X, M) = H_n(X, F_{\sharp}^{lin})$$

termed *stable square-homology*. Here F_{\sharp}^{lin} corresponds to a spectrum E_M so that we get a functor from the category of square groups to the homotopy category of spectra which carries M to E_M . We show that E_M is always the cofiber of a map Sq_M which carries an Eilenberg-Mac Lane spectrum to a product of Eilenberg-Mac Lane spectra. We call Sq_M the *squaring operation* associated to M . For example all Steenrod squares Sq^2, Sq^3, \dots can be derived from Sq_M .

Let $\Gamma_n G$ and $\bar{\Gamma}_n G$ be the subgroups of a group G given by the lower central series and the mod-2 restricted lower central series respectively. Then we obtain the quadratic group functors

$$nil_2, \overline{nil}_2 : \mathbf{Gr} \rightarrow \mathbf{Gr}$$

which carry G to the quotients $G/\Gamma_3 G$ and $G/\bar{\Gamma}_3 G$ respectively. The corresponding square groups are \mathbb{Z}_{nil} and $\mathbb{Z}_{nil}^{4,2}$ with

$$\begin{aligned} nil_2(G) &= G \otimes \mathbb{Z}_{nil} \\ \overline{nil}_2(G) &= G \otimes \mathbb{Z}_{nil}^{4,2} \end{aligned}$$

We compute the squaring operation Sq_M for $M = \mathbb{Z}_{nil}$ and $M = \mathbb{Z}_{nil}^{4,2}$ explicitly in terms of Steenrod squares. This determines the spectrum associated to the linearization of the homotopy functor $(nil_2)_{\sharp}$, resp. $(\overline{nil}_2)_{\sharp}$; see (8.16), (8.17).

The author happily acknowledges helpful conversations with Teimuraz Pirashvili. He also thanks G. Arone and T. Goodwillie for comments concerning ‘‘Calculus’’.

§ 1 CW-SPACES, CW-PAIRS AND SPECTRA

We use the following conventions: **Top** is the category of topological spaces and **Top*** is the category of topological spaces with base point. We obtain cofibrations in **Top** by the universal homotopy extension property defined via the cylinder $I \times X$ where $I = [0, 1]$ is the unit interval. A homotopy in **Top*** is a pointed map $I(X) \rightarrow Y$ where $I(X) = I \times X / I \times \{*\}$ is the reduced cylinder. By the inclusion $(i_0, i_1) : X \vee X \subset I(X)$ we obtain the cone $CX = I(X)/i_1X$ and the suspension $\Sigma X = CX/i_0X$. Here we use the quotient space X/Y which is defined for any pair of spaces (X, Y) by the adjunction $X/Y = X \cup_Y \{*\}$. More generally an adjunction space $X \cup_Y Z$ is defined by a push out diagram

$$\begin{array}{ccc} X & \xrightarrow{\quad} & X \cup_Y Z \\ & \bar{g} \searrow & \\ \uparrow & & \uparrow \\ Y & \xrightarrow{g} & Z \end{array}$$

If (X, Y) is a pair of spaces we call $\bar{g} : (X, Y) \rightarrow (X \cup_Y Z, Z)$ an *adjunction map*.

A space X is a *CW-space* if there is a CW-complex Y together with a homotopy equivalence $Y \simeq X$ in **Top**. This is a *finite CW-space* if Y can be chosen to be a CW-complex with finitely many cells. Moreover we write $\dim(X) \leq n$ if $\dim(Y) \leq n$. A *CW-pair* (X, Y) is a pair in **Top** of CW-spaces X and Y for which the inclusion $Y \subset X$ is a cofibration. This is a *finite CW-pair* if X and Y are finite CW-spaces. A CW-space X is well pointed if the inclusion $\{*\} \rightarrow X$ is a cofibration.

Let **space** \subset **Top*** be the full subcategory of well pointed CW-spaces and pointed maps. Moreover let **pair** be the category of well pointed CW-pairs and pointed pair maps. We have functors

$$\mathbf{space} \xrightarrow{i} \mathbf{pair} \xrightarrow{q} \mathbf{space}$$

where i carries X to the pair $(X, *)$ and q carries (X, Y) to the quotient space X/Y . We also use the full subcategory **space** _{r} of $(r - 1)$ -connected objects in **space** and the full subcategory **pair** _{r} of objects (X, Y) in **pair** for which X and Y are $(r - 1)$ -connected. We point out that the suspension yields a functor

$$\Sigma : \mathbf{space}_r \rightarrow \mathbf{space}_{r+1}$$

raising the degree of connectedness.

A *spectrum* E is a sequence of maps $\varepsilon_n : \Sigma E_n \rightarrow E_{n+1}$ in **space**, $n \in \mathbb{Z}$. A map $f : E \rightarrow E'$ between spectra is a sequence of maps $f_n : E_n \rightarrow E'_n$ with $f_{n+1}\varepsilon_n = \varepsilon'_n(\Sigma f_n)$. Let **spectra** be the category of such spectra and maps.

A map $f : X \rightarrow Y$ in **Top** is a *weak equivalence* if f induces a bijection of homotopy groups $f_* : \pi_n X \approx \pi_n Y$, $n \geq 0$, for every basepoint in the domain. A map in **Top*** is a weak equivalence if it is one in **Top**. Clearly a weak equivalence in **space** or **pair** is also a homotopy equivalence. A map $f : E \rightarrow E'$ in **spectra** is a weak equivalence if it induces an isomorphism $f_* : \pi_k E \approx \pi_k E'$. Here $\pi_k E = \text{colim} \{ \pi_{n+k} E_n \}$ is an abelian group for all $k \in \mathbb{Z}$.

§ 2 AXIOMS OF HOMOLOGY

To fix notation we recall the following basic properties of a homology theory.

(2.1) *Definition.* Let $r \geq 0$. A *suspension theory* (H, s) is a sequence of covariant functors, $n \in \mathbb{Z}$,

$$H_n : \mathbf{space}_r \rightarrow \mathbf{Ab}$$

together with a sequence of natural transformations, $n \in \mathbb{Z}$,

$$s_n : H_n \rightarrow H_{n+1} \circ \Sigma$$

such that $H_n(*) = 0$ and $H_n(f_0) = H_n(f_1)$ for homotopic maps $f_0 \simeq f_1$. We also consider suspension theories which satisfy some of the following axioms.

Exactness. For $(X, Y) \in \mathbf{pair}_r$ the sequence

$$H_n(Y) \xrightarrow{i_*} H_n(X) \xrightarrow{q_*} H_n(X/Y)$$

is exact where $i : Y \rightarrow X$ is the inclusion and $q : X \rightarrow X/Y$ is the quotient map.

Suspension. For $X \in \mathbf{space}_r$ the natural map

$$s_n(X) : H_n(X) \rightarrow H_{n+1}(\Sigma X)$$

given by the transformation s_n is an isomorphism for all $n \in \mathbb{Z}$.

Colimit axiom. For each sequence $X_0 \rightarrow X_1 \rightarrow \dots$ of cofibrations in \mathbf{space}_r the induced map

$$\operatorname{colim} \{H_n(X_i)\} \rightarrow H_n(\operatorname{colim} \{X_i\})$$

is an isomorphism.

A suspension theory (H, s) is termed a *homology theory* on \mathbf{space}_r if the exactness and suspension axioms are satisfied; compare [29].

Given a suspension theory (H, s) we obtain the *stable theory* (H^S, s) associated to (H, s) . Here $H_n^S(X)$ is the colimit of

$$(2.2) \quad H_n(X) \xrightarrow{s} H_{n+1}(\Sigma X) \rightarrow \dots \rightarrow H_{n+k}(\Sigma^k X) \xrightarrow{s} \dots$$

Clearly one obtains the canonical map

$$s : H_n^S(X) \rightarrow H_{n+1}^S(\Sigma X)$$

which is an isomorphism. Hence (H^S, s) is a suspension theory satisfying the suspension axiom. If (H, s) satisfies the exactness or colimit axiom then so does (H^S, s) . Clearly for a homology theory we have $(H, s) = (H^S, s)$.

(2.3) *Definition.* Let $r \geq 0$. A *boundary theory* (H, ∂) is a sequence of covariant functors, $n \in \mathbb{Z}$,

$$H_n : \mathbf{pair}_r \rightarrow \mathbf{Ab}$$

together with a sequence of natural transformations $\partial : H_{n+1}(X, Y) \rightarrow H_n(Y)$ with $(X, Y) \in \mathbf{pair}_r$ and $H_n(Y) = H_n(Y, *)$, such that $H_n(*) = 0$ and $H_n(f_0) = H_n(f_1)$ for homotopic maps $f_0 \simeq f_1$ in \mathbf{pair}_r . Moreover ∂ -*exactness* is satisfied, that is

$$\dots \rightarrow H_{n+1}(X, Y) \xrightarrow{\partial} H_n(Y) \xrightarrow{i_*} H_n(X) \xrightarrow{j_*} H_n(X, Y) \xrightarrow{\partial} \dots$$

is exact. Here i_* and j_* are induced by $i : Y \subset X$ and $j : (X, *) \subset (X, Y)$. We also consider boundary theories with the following additional property.

Excision. For $(X, Y) \in \mathbf{pair}_r$ and $g : Y \rightarrow Z \in \mathbf{space}_r$ the adjunction map \bar{g} induces an isomorphism

$$\bar{g}_* : H_n(X, Y) \cong H_n(X \cup_Y Z, Z)$$

for all $n \in \mathbb{Z}$.

A boundary theory (H, ∂) is termed a *homology theory* on \mathbf{pair}_r if excision is satisfied. Compare [29].

Each boundary theory (H, ∂) yields a suspension theory (H, s) as follows. Let CX be the cone on X . Then $\partial : H_{n+1}(CX, X) \cong H_n(X)$ is an isomorphism. Hence we obtain the suspension map

$$(2.4) \quad s = q_* \partial^{-1} : H_n(X) \cong H_{n+1}(CX, X) \rightarrow H_{n+1}(\Sigma X)$$

where $q : (CX, X) \rightarrow (\Sigma X, *)$ is the quotient map. If (H, ∂) is a homology theory then $q_* \partial^{-1}$ is an isomorphism since excision implies that $q_* : H_n(X, Y) \rightarrow H_n(X/Y)$ is an isomorphism. This leads to the following well known lemma. For this we observe that suspension theories, resp. boundary theories form categories. Morphisms are the natural transformations compatible with s , resp. ∂ .

(2.5) Lemma. *The category of homology theories (H, ∂) on \mathbf{pair}_r and the category of homology theories (H, s) on \mathbf{space}_r are equivalent. The equivalence carries (H, ∂) to $(H, q_* \partial^{-1})$.*

These categories actually do not depend on r since the category of homology theories (H', s') on \mathbf{space}_r is equivalent to the category of homology theories (H, s) on \mathbf{space} . The equivalence carries (H', s') to (H, s) with $H_n(X) = H'_{n+r}(\Sigma^r X)$. This shows that a homology theory on \mathbf{space} is determined by its restriction to the category \mathbf{space}_r for arbitrary large $r \geq 0$. Such a statement will not be true for quadratic homology theories below.

(2.6) Example. (A) Let A be an abelian group and let $H_n(X, Y; A)$ be the *singular homology* of the pair (X, Y) with coefficients in A . This is the classical homology theory on \mathbf{pair} .

(B) Let $\pi_n(X, Y)$ be the *relative homotopy group* of the pair (X, Y) . Then (π_n, ∂) is a boundary theory on \mathbf{pair}_2 with $\pi_n(X, Y) = 0$ for $n \leq 1$. Here we use $(X, Y) \in$

\mathbf{pair}_2 in order to obtain abelian groups $\pi_n(X, Y)$, $n \in \mathbb{Z}$. The associated stable theory (π_n^S, s) is the homology theory on \mathbf{space}_2 termed *stable homotopy*.

The *spherical groups* of a homology theory H on \mathbf{pair}_r are the groups $H_n(S^r)$, $n \in \mathbb{Z}$, where S^r is the r -sphere. The suspension isomorphism $H_n(S^r) \cong H_{n+k}(S^{r+k})$ shows that the spherical groups determine the value of H on all spheres in \mathbf{space}_r . Moreover the following *uniqueness-lemma* is well known: Let $\phi : H \rightarrow H'$ be a natural transformation of homology theories on \mathbf{pair}_r such that ϕ is an isomorphism on spherical groups; that is, $\phi : H_n(S^r) \cong H'_n(S^r)$ for $n \in \mathbb{Z}$. Then ϕ is an isomorphism

$$(2.7) \quad \phi : H_n(X, Y) \cong H'_n(X, Y)$$

for all finite CW-pairs $(X, Y) \in \mathbf{pair}_r$ and $n \in \mathbb{Z}$. Compare for example (2.12) in [28]. A similar uniqueness lemma is also true for the quadratic homology theories below. Moreover we have the following *representability theorem* of E.H. Brown; compare for example I.3.8 in [28]. Let H be a homology theory on \mathbf{space}_r . Then there exists a spectrum E and a natural isomorphism

$$(2.8) \quad H_n(X) \cong \pi_n(E \wedge X) = H_n(X, E), \quad n \in \mathbb{Z},$$

for all finite CW-spaces X in \mathbf{space}_r .

Next we describe the “partial suspension” and the “cross effect suspension” of a boundary theory. To this end we have to introduce the following notation on cross effects.

(2.9) *Notation.* Let \mathbf{Gr} be the category of groups and let \mathbf{C} be a category with zero object $*$ and assume that sums (coproducts) $X \vee Y$ exist in \mathbf{C} . For objects $X, Y \in \mathbf{C}$ one has the unique zero morphism $0 : X \rightarrow * \rightarrow Y$. Maps $f : X \rightarrow Z$, $g : Y \rightarrow Z$ determine $(f, g) : X \vee Y \rightarrow Z$. We have the retraction $r_1 : X \vee Y \rightarrow X$ and $r_2 : X \vee Y \rightarrow Y$ with $r_1 = (1, 0)$ and $r_2 = (0, 1)$. Given a functor

$$F : \mathbf{C} \rightarrow \mathbf{Gr}$$

we define the *cross effect*

$$(1) \quad F(X|Y) = \text{kernel} \{(Fr_1, Fr_2) : F(X \vee Y) \rightarrow F(X) \times F(Y)\}$$

This yields the functor $F(|) : \mathbf{C} \times \mathbf{C} \rightarrow \mathbf{Gr}$ with induced maps denoted by $(f|g)_*$. The inclusion $i_{12} : F(X|Y) \subset F(X \vee Y)$ is natural in X and Y . Moreover we have the natural interchange isomorphism

$$(2) \quad T : F(X|Y) \cong F(Y|X)$$

induced by $T : X \vee Y = Y \vee X$ with $T = (i_2, i_1)$. Let $F(X \vee Y)_2$ be the kernel of $Fr_2 : F(X \vee Y) \rightarrow F(Y)$. Then we have the split short exact sequence of groups

$$(3) \quad 0 \rightarrow F(X|Y) \xrightarrow{i_{12}} F(X \vee Y)_2 \xrightarrow{Fr_1} F(X) \rightarrow 0$$

If F is a functor $\mathbf{C} \rightarrow \mathbf{Ab}$ we thus have natural isomorphisms

$$(4) \quad \begin{aligned} (Fi_1, i_{12}) : F(X) \oplus F(X|Y) &= F(X \vee Y)_2 \\ (Fi_1, Fi_2, i_{12}) : F(X) \oplus F(Y) \oplus F(X|Y) &= F(X \vee Y) \end{aligned}$$

which we use as identifications. The functor $F : \mathbf{C} \rightarrow \mathbf{Gr}$ is *linear* if $F(*) = 0$ and $F(X|Y) = 0$ for all $X, Y \in \mathbf{C}$; that is

$$(5) \quad (Fr_1, Fr_2) : F(X \vee Y) \cong F(X) \times F(Y)$$

is an isomorphism. The functor $F : \mathbf{C} \rightarrow \mathbf{Gr}$ is *quadratic* if $F(*) = 0$ and the cross effect $F(X|Y)$ as a bifunctor is linear in each variable X and Y . If F is linear then the group $F(X)$ is abelian and if F is quadratic then the group $F(X)$ has nilpotency degree 2. Moreover for a quadratic functor the subgroup $F(X|Y)$ is central in $F(X \vee Y)$. Compare [10].

Now let (H, ∂) be a boundary theory as in (2.3). Then one gets the following exact sequences from which we derive the partial suspension E . Let CY be the cone on Y so that $H_n(CY \vee X) = H_n(X)$. This yields the short exact sequence

$$0 \rightarrow H_{n+1}(CY \vee X, Y \vee X) \xrightarrow{\partial} H_n(Y \vee X) \xrightarrow{r_2} H_n(X) \rightarrow 0$$

and hence the isomorphism

$$\partial_0 : H_{n+1}(CY \vee X, Y \vee X) \cong H_n(Y \vee X)_2.$$

Moreover one gets the short exact sequence

$$0 \rightarrow H_{n+1}(Y) \rightarrow H_{n+1}(\Sigma X \vee Y) \xrightarrow{j_*} H_{n+1}(\Sigma X \vee Y, Y) \rightarrow 0$$

which shows that j_* induces the isomorphism

$$j_0 : H_{n+1}(\Sigma X \vee Y)_2 \cong H_{n+1}(\Sigma X \vee Y, Y)$$

Dividing out X yields the quotient map $q \vee 1 : (CX \vee Y, X \vee Y) \rightarrow (\Sigma X \vee Y, Y)$. Now the *partial suspension* E is the composition

$$(2.10) \quad E = j_0^{-1}(g \vee 1)_* \partial_0^{-1} : H_n(X \vee Y)_2 \rightarrow H_{n+1}(\Sigma X \vee Y)_2.$$

Compare [2, 3]. If $Y = *$ this is the suspension in (2.4). Moreover the partial suspension restricts to cross effects yielding the *cross effect suspension*

$$(2.11) \quad s : H_n(X|Y) \rightarrow H_{n+1}(\Sigma X|Y)$$

such that via (2.9) (4) the partial suspension is the composite

$$\begin{array}{ccc} H_n(X \vee Y)_2 & \xrightarrow{E} & H_{n+1}(\Sigma X \vee Y)_2 \\ \parallel & & \parallel \\ H_n(X) \oplus H_n(X|Y) & \xrightarrow{s \oplus s} & H_{n+1}(\Sigma X) \oplus H_{n+1}(\Sigma X|Y) \end{array}$$

Hence for each Y the pair $(H_n(-|Y), s)$ is a suspension theory. Using the interchange map T in (2.9) (2) also $(H_n(X|-, \bar{s}))$ is a suspension theory where \bar{s} is the composite

$$(2.12) \quad \bar{s} = TsT : H_n(X|Y) \cong H_n(Y|X) \rightarrow H_{n+1}(\Sigma Y|X) \cong H_{n+1}(X|\Sigma Y).$$

The sum or coproduct in the category \mathbf{space}_r is obtained by the one point union $X \vee Y$ of spaces.

(2.13) Lemma. *Let H be a homology theory on \mathbf{pair}_r . Then $H_n : \mathbf{space}_r \rightarrow \mathbf{Ab}$ is linear, that is $H_n(X|Y) = 0$ and $H_n(X \vee Y) = H_n(X) \oplus H_n(Y)$.*

Proof. For the CW-pair $(X \vee Y, Y) \in \mathbf{pair}_r$ we have by exactness the split short exact sequence

$$0 \rightarrow H_n(Y) \rightarrow H_n(X \vee Y) \rightarrow H_n(X \vee Y, Y) \rightarrow 0$$

where $H_n(i_2)$ is injective since we have the retraction $H_n(r_2)$. Moreover excision shows that $H_n(X) \rightarrow H_n(X \vee Y, Y)$ is an isomorphism.

q.e.d.

(2.14) Lemma. *Let (H, ∂) be a boundary theory on \mathbf{pair}_r . Then the suspension s in (2.4) makes the following diagram commute*

$$\begin{array}{ccc} H_n(X \vee Y) & \xrightarrow{((r_1)_*, (r_2)_*)} & H_n(X) \times H_n(Y) \\ \downarrow s & & \downarrow s \times s \\ H_{n+1}(\Sigma X \vee \Sigma Y) & \xleftarrow{(i_1)_* + (i_2)_*} & H_{n+1}(\Sigma X) \times H_{n+1}(\Sigma Y) \end{array}$$

This implies that the composition

$$H_n(X|Y) \xrightarrow{i_{12}} H_n(X \vee Y) \xrightarrow{s} H_{n+1}(\Sigma X \vee \Sigma Y)$$

is always trivial, $si_{12} = 0$.

Proof of (2.14). Let $\pi : CX \rightarrow \Sigma X$ be the quotient map. Then

$$C(X \vee Y) = C(X) \vee C(Y) \xrightarrow{\pi_X \vee 1} \Sigma X \vee CY \xrightarrow{1 \vee \pi_Y} \Sigma X \vee \Sigma Y = \Sigma(X \vee Y)$$

is the quotient map $\pi_{X \vee Y}$. Hence we get the following diagram in which the row is split short exact.

$$\begin{array}{ccccccc}
& & & H_{n+1}(C(X \vee Y), X \vee Y) \xrightarrow{\cong} H_n(X \vee Y) & & & \\
& & & \downarrow (\pi_{X \vee Y})_* & & & \\
0 & \longrightarrow & H_{n+1}(\Sigma X) & \longrightarrow & H_{n+1}(\Sigma X \vee CY, Y) & \xrightarrow{\partial} & H_n(Y) \longrightarrow 0 \\
& & & & \downarrow (1 \vee \pi_Y)_* & & \\
& & & & H_{n+1}(\Sigma X \vee \Sigma Y) & &
\end{array}$$

Here we have the isomorphism $(i_1)_* + (i_2)_*$

$$H_{n+1}(\Sigma X) \oplus H_{n+1}(CY, Y) \xrightarrow{\cong} H_{n+1}(\Sigma X \vee CY, Y)$$

by the exactness of the row. This yields the result.

q.e.d

§ 3 QUADRATIC HOMOLOGY

In this section we introduce new axioms of a quadratic homology. These axioms are satisfied by various examples which we describe in the following sections. We deal with some consequences of the axioms, in particular we obtain the long EHP-sequence of quadratic homology and we prove a uniqueness lemma.

(3.1) *Notation.* Let $F : \mathbf{C} \rightarrow \mathbf{Gr}$ be a functor where \mathbf{C} is a category with zero object $*$ and sums and let $F(*) = 0$. We obtain the natural homomorphism

$$P : F(X|X) \xrightarrow{i_{12}} F(X \vee X) \xrightarrow{(1,1)_*} F(X)$$

for $X \in \mathbf{C}$. Moreover if X is a cogroup in \mathbf{C} with structure maps $\mu : X \rightarrow X \vee X$ and $\nu : X \rightarrow X$ satisfying the usual identities we define the function H , with

$$F(X) \xrightarrow{H} F(X|X) \xrightarrow{i_{12}} F(X \vee X),$$

by $i_{12}H(a) = F(\mu)(a) - F(i_2)(a) - F(i_1)(a)$ where i_1, i_2 are the inclusions of X in $X \vee X$. Clearly H is natural with respect to maps between cogroups. We write $F\{X\}$ for the pair of functions

$$F\{X\} = (F(X) \xrightarrow{H} F(X|X) \xrightarrow{P} F(X))$$

It is shown in 3.6 [10] that $F\{X\}$ is a square group (see (6.3) below) if F is a quadratic functor. Clearly H is a homomorphism for a functor $F : \mathbf{C} \rightarrow \mathbf{Ab}$. If \mathbf{C} is an additive category then a quadratic functor $F : \mathbf{C} \rightarrow \mathbf{Ab}$ yields for $X \in \mathbf{C}$ a pair of homomorphisms $F\{X\} = (H, P)$ as above with $HPH = 2H$ and $PHP = 2P$; that is $F\{X\}$ is a quadratic \mathbb{Z} -module; compare [5].

(3.2) *Definition.* Let $r \geq 0$ and consider a boundary theory (Q, ∂) on \mathbf{pair}_r with the following properties (i) and (ii).

- (i) The stable theory Q^S associated to Q via the suspension (2.4) is a homology theory.

- (ii) For each $Y \in \mathbf{space}_r$ the suspension theory of cross effects $(Q(-, Y), s)$ in (2.11) is a homology theory.

A *quadratic homology theory* (Q, ∂, δ) is a boundary theory (Q, ∂) which satisfies (i) and (ii) together with a sequence of homomorphisms, $n \in \mathbb{Z}$,

$$\delta : Q_{n+1}(CA \cup_A X, X) \rightarrow Q_{n-1}(A|A)$$

which are defined for all pairs (X, A) in \mathbf{pair}_r and which are natural in (X, A) , that is: A pair map $f : (X, A) \rightarrow (Y, B) \in \mathbf{pair}_r$ induces a commutative diagram

$$\begin{array}{ccc} Q_{n+1}(CA \cup_A X, X) & \xrightarrow{\delta} & Q_{n-1}(A|A) \\ Q_n(Cg \cup f) \downarrow & & \downarrow (g|g)_* \\ Q_{n+1}(CB \cup_B Y, Y) & \xrightarrow{\delta} & Q_{n-1}(B|B) \end{array}$$

where $g : A \rightarrow B$ is the restriction of f . Moreover the following property (iii) is satisfied.

- (iii) For each pair (X, A) , the sequence

$$\dots \xrightarrow{\delta} Q_n(A|A) \xrightarrow{j_\#} Q_{n+1}(CA \vee X, A \vee X) \xrightarrow{i_\#} Q_{n+1}(CA \cup_A X, X) \xrightarrow{\delta} Q_{n-1}(A|A) \rightarrow \dots$$

is exact. Here the inclusion $i : A \rightarrow X$ yields the map $(i, 1) : A \vee X \rightarrow X$ which defines the adjunction map $\overline{(i, 1)} : (CA \vee X, A \vee X) \rightarrow (CA \cup_A X, X)$ which induces $i_\# = \overline{(i, 1)}_*$ in the sequence. Moreover $j_\#$ in the sequence is the composition

$$j_\# = \partial_0^{-1}(P, -(1|i)_*) : Q_n(A|A) \rightarrow Q_n(A) \oplus Q_n(A|X) = Q_n(A \vee X)_2 \cong Q_{n+1}(CA \vee X, A \vee X).$$

Here P is the natural map in (3.1) and $-(1|i)_*$ is the negative of the induced homomorphism $(1|i)_* = Q_n(1|i)$. We call the long exact sequence $(j_\#, i_\#, \delta)$ above the *quadratic excision sequence*.

It follows from (ii) above and (2.13) that all functors

$$Q_n : \mathbf{space}_r \rightarrow \underline{Ab}$$

are quadratic. Moreover one can check that a quadratic homology theory (Q, ∂, δ) for which these functors Q_n are linear is the same as a homology theory on \mathbf{pair}_r . This follows from (i) and (3.5) below.

We now derive from the axioms of a quadratic homology theory some consequences. First we observe that the following diagram commutes.

(3.3)

$$\begin{array}{ccccc}
Q_n(A|A) & \xrightarrow{j_!} & Q_{n+1}(CA \vee X, A \vee X) & \xrightarrow{i_!} & Q_{n+1}(CA \cup_A X, X) \\
\downarrow d_2 & & \cong \downarrow \partial_0 & & \downarrow \partial \\
& & Q_n(A \vee X)_2 & \xrightarrow{(i,1)_*} & Q_n(X) \\
& & \parallel & & \parallel \\
Q_n(A) \oplus Q_n(A|X) & \xrightarrow{d_1} & & & Q_n(X)
\end{array}$$

where $d_2 = (P, -(1|i)_*)$ and $d_1 = (i_*, P(i|1)_*)$. Here $d_1 d_2 = 0$ since P is natural; in fact,

$$d_1 d_2 = i_* P - P(i|1)_*(1|i)_* = i_* P - P(i|i)_* = 0.$$

Clearly $d_1 d_2 = 0$ also follows from $i_! j_! = 0$ and the commutativity of the diagram. Using (3.3) we can equivalently describe the excision exact sequence in (3.2) (iii) by the exact sequence in the row of the commutative diagram:

(3.4)

$$\begin{array}{ccccccc}
\dots & \xrightarrow{\delta} & Q_n(A|A) & \xrightarrow{d_2} & Q_n(A) \oplus Q_n(A|X) & \xrightarrow{\bar{d}_1} & Q_{n+1}(CA \cup_A X, X) & \xrightarrow{\delta} & \dots \\
& & & & & & \downarrow \partial & & \\
& & & & & & Q_n(X) & &
\end{array}$$

Here we set $\bar{d}_1 = i_! \partial_0^{-1}$ by (3.4). This is similar to the diagram for metastable homotopy groups in 2.4 of [4]. As a consequence of (3.4) we obtain a long exact sequence which resembles the classical EHP-sequence of James [23].

(3.5) Proposition. *Let Q be a quadratic homology theory on pair r , $r \geq 0$. Then one has the following long exact sequence which is natural in $A \in \text{space } r$.*

$$\dots \rightarrow Q_n(A|A) \xrightarrow{P} Q_n(A) \xrightarrow{s} Q_{n+1}(\Sigma A) \xrightarrow{H'} Q_{n-1}(A|A) \xrightarrow{P} Q_{n-1}(A) \rightarrow \dots$$

Proof. We consider (3.4) in case $X = CA$ is the cone on A . Then the homotopy axiom shows $Q_n(A|CA) = 0$ and $Q_n(CA) = 0$. Hence we obtain the isomorphism

$$i_* : Q_{n+1}(CA \cup_A CA) \cong Q_{n+1}(CA \cup_A CA, CA)$$

where $\Sigma A = CA \cup_A CA$. Thus (3.4) yields the commutative diagram

$$\begin{array}{ccccccc}
Q_n(A|A) & \xrightarrow{P} & Q_n(A) & \xrightarrow{s} & Q_{n+1}(\Sigma A) & \xrightarrow{H'} & Q_{n-1}(A|A) \\
\parallel & & \parallel & & \parallel & & \parallel \\
Q_n(A|A) & \xrightarrow{d_1} & Q_n(A) & \xrightarrow{d_2} & Q_{n+1}(CA \cup_A CA, CA) & \xrightarrow{\delta} & Q_{n-1}(A|A)
\end{array}$$

which defines the exact sequence in (3.5).

q.e.d.

Given a natural exact sequence as in (3.5) one obtains the associated sequence of cross effects which is also exact. Since $si_{12} = 0$ by (2.14) this yields the following commutative diagram with exact rows.

$$\begin{array}{ccccccc}
0 & \longrightarrow & Q_{n+2}(\Sigma A|\Sigma A') & \xrightarrow{\sigma'} & Q_n(A|A') \oplus Q_n(A'|A) & \xrightarrow{(1,T)} & Q_n(A|A') & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
& & Q_{n+2}(\Sigma A \vee \Sigma A') & \xrightarrow{H'} & Q_n(A \vee A'|A \vee A') & \xrightarrow{P} & Q_n(A \vee A') & &
\end{array}$$

Here the vertical arrows are the inclusions i_{12} . This shows that σ' induces the isomorphism

$$(3.6) \quad \sigma : Q_{n+2}(\Sigma A|\Sigma A') \cong Q_n(A|A')$$

where $\sigma = (pr_1)\sigma'$ is the composition of σ' and the projection pr_1 . Clearly σ is again natural in A and A' . We do not see that σ coincides up to sign with the suspension isomorphisms $\bar{s}s$ given by (3.2) (ii) and (2.11), (2.12).

(3.7) Proposition. *Let A be a cogroup in $\mathbf{space}_r / \simeq$, for example let A be a suspension. Then the operator H' in (3.5) and H in (3.1) yield the following commutative diagram*

$$\begin{array}{ccc}
& Q_{n+1}(\Sigma A) & \\
H \swarrow & & \searrow H' \\
Q_{n+1}(\Sigma A|\Sigma A) & \xrightarrow[\sigma]{\cong} & Q_{n-1}(A|A)
\end{array}$$

where σ is the isomorphism in (3.6).

Proof. Let $\mu : A \rightarrow A \vee A'$ be the comultiplication of A with $A = A'$. Then $\Sigma\mu$ is the comultiplication of the suspension ΣA . Now the naturality of H' in (3.5) shows that the following diagram commutes

$$\begin{array}{ccc}
Q_{n+1}(\Sigma A \vee \Sigma A') & \xrightarrow{H'} & Q_{n-1}(A \vee A'|A \vee A') \\
(\Sigma\mu)_* \cdot -(i_2)_* \cdot -(i_1)_* \uparrow & & \uparrow \mu_* \cdot -(i_2)_* \cdot -(i_1)_* \\
Q_{n+1}(\Sigma A) & \xrightarrow{H'} & Q_{n-1}(A|A)
\end{array}$$

Here the vertical arrows map to the cross effects; in fact the left hand side is $i_{12}H$ by definition of H and the right hand side is $i_{12}(1, T)$ by the bilinearity of the cross effect. This implies by the definition of σ' above that $\sigma'H = (1, T)H'$ and hence $\sigma H = H'$.

q.e.d.

For the metastable range of homotopy groups one has a diagram as in (3.7) where σ is actually the double suspension up to sign; compare A.6.8 in [7] and example (2.6) (B) above.

The next lemma justifies our choice of axioms of a quadratic homology theory. It is the quadratic analogue of the uniqueness lemma in (2.7) above. The *spherical groups* of a quadratic homology theory Q on **pair** r are the groups $Q_n(S^{r+k})$ and $Q_n(S^r|S^r)$, $n \in \mathbb{Z}$, $k \geq 0$. Now the following *uniqueness lemma* holds.

(3.8) Lemma. *Let $\phi : Q \rightarrow Q'$ be a natural transformation of quadratic homology theories on **pair** r compatible with ∂ and δ . Moreover assume ϕ is an isomorphism on spherical groups. Then ϕ is an isomorphism*

$$\phi : Q_n(X, Y) \cong Q'_n(X, Y)$$

for all finite CW-pairs $(X, Y) \in \mathbf{pair}_r$ and $n \in \mathbb{Z}$.

Proof. Since $Q_n(\ |)$ is a homology theory in each variable we see that ϕ induces an isomorphism $Q_n(X|Z) \cong Q'_n(X|Z)$ for all finite X, Z . Compare (2.7). This shows that for $k \geq 0$ one gets the isomorphism $\phi : Q_n(X) \cong Q'_n(X)$ for all spaces X which are finite one point unions of spheres S^{r+k} . By exactness it suffices to show that ϕ induces an isomorphism $\phi_X : Q_n(X) \cong Q'_n(X)$ for all finite X in **space** r . We proceed by induction on the dimension of X . If $\dim(X) = r$ then X is a finite one point union of spheres S^r . Now assume ϕ_X is an isomorphism for all finite X with $\dim(X) < r + k$ and let $\dim(Y) = r + k$. Then we may assume that $Y = CA \cup_A X$ where (X, A) is a pair with $\dim(X) < r + k$ and A is a finite one point union of spheres S^{r+k-1} . Hence ϕ_X and ϕ_A are isomorphisms and therefore the quadratic excision sequence shows that also

$$\phi : Q_{n+1}(Y, X) \cong Q'_{n+1}(Y, X)$$

is an isomorphism. Hence ∂ -exactness shows that ϕ_Y is an isomorphism since ϕ_X is one.

q.e.d.

A *Moore space* $M(A, n)$, $n \geq 2$, is a simply connected CW-space X with homology $H_n(X, \mathbb{Z}) = A$ and $\tilde{H}_i(X, \mathbb{Z}) = 0$ otherwise. If A is a free abelian group then $M(A, n)$ is a one point union of spheres S^n , in particular $M(\mathbb{Z}, n) = S^n$. Each homomorphism $\varphi : A \rightarrow B$ in **Ab** admits a realization $\bar{\varphi} : M(A, n) \rightarrow M(B, n)$ with $H_n(\bar{\varphi}) = \varphi$. Here $\bar{\varphi}$ is unique up to homotopy if A is free abelian. Now let (Q, ∂, δ) be a quadratic homology on **space** r and let $n \geq r, m \in \mathbb{Z}$. By (3.10) we obtain the quadratic \mathbb{Z} -module

$$(3.9) \quad Q_m\{S^n\} = (Q_m(S^n) \xrightarrow{H} Q_m(S^n|S^n) \xrightarrow{P} Q_m(S^n))$$

which describes the canonical quadratic structure of the spherical groups above. We now recall the following notation concerning quadratic \mathbb{Z} -modules; compare [5].

(3.10) Definition. Let $M = (M_e \xrightarrow{H} M_{ee} \xrightarrow{P} M_e)$ be a quadratic \mathbb{Z} -module, i.e. a pair of homomorphisms H and P with $HPH = 2H$ and $PHP = 2P$. Then

M induces functors $\mathbf{Ab} \rightarrow \mathbf{Ab}$ which carry A to $A \otimes M$, $A *' M$ and $A *'' M$ respectively. Here $A \otimes M$ is the abelian group with generators $a \otimes m, [a, b] \otimes n$ for $a, b \in A, m \in M_e, n \in M_{ee}$ and relations

$$\begin{aligned}(a + b) \otimes m &= a \otimes m + b \otimes m + [a, b] \otimes Hm \\ [a, a] \otimes n &= a \otimes P(n)\end{aligned}$$

where $a \otimes m$ is linear in m and $[a, b] \otimes n$ is linear in a, b and n . Now let

$$0 \rightarrow A_1 \xrightarrow{d} A_0 \xrightarrow{q} A \rightarrow 0$$

be a short exact sequence in \mathbf{Ab} where A_0 is free abelian. Then we obtain homomorphisms

$$A_1 \otimes A_1 \otimes M_{ee} \xrightarrow{d_2} A_1 \otimes M \oplus A_1 \otimes A_0 \otimes M_{ee} \xrightarrow{d_1} A_0 \otimes M$$

with $d_1 d_2 = 0$ as follows:

$$\begin{aligned}d_1(a \otimes m) &= (da) \otimes m, \\ d_1([a, a'] \otimes n) &= [da, a'] \otimes n, \\ d_1(a \otimes b \otimes n) &= [da, b] \otimes n, \\ d_2(a \otimes a' \otimes n) &= -a \otimes da' \otimes n + [a, da'] \otimes n\end{aligned}$$

for $a, a' \in A_1, b \in A_0, m \in M_e, n \in M_{ee}$. One can check that $\text{cok}(d_1) = A \otimes M$. Moreover we set

$$\begin{aligned}A *' M &= \ker(d_1)/\text{im}(d_2) \\ A *'' M &= \ker(d_2)\end{aligned}$$

These are the derived functors of the functor $\mathbf{Ab} \rightarrow \mathbf{Ab}$ which carries A to $A \otimes M$. An abelian group $N \in \mathbf{Ab}$ yields the quadratic \mathbb{Z} -module $N = (N \rightarrow 0 \rightarrow N)$ with $N_{ee} = 0$. In this case $A \otimes N$ is the usual tensor product and $A * N = A *' N$ is the usual torsion product. Clearly $A *'' N = 0$.

We now consider the quadratic homology of a Moore space $M(A, n)$. There is the natural homomorphism

$$(3.11) \quad \lambda : A \otimes Q_m\{S^n\} \rightarrow Q_m(M(A, n))$$

defined as follows. For $a, b \in A = \pi_n M(A, n), u \in Q_m(S^n), v \in Q_m(S^n|S^n)$ we set

$$\begin{aligned}\lambda(a \otimes u) &= Q_m(a)(u) \\ \lambda([a, b] \otimes v) &= P Q_m(a|b)(v)\end{aligned}$$

Let ${}_\lambda Q_m(M(A, n))$ be the cokernel of λ .

(3.12) Proposition. Assume that the quadratic homology (Q, ∂, δ) satisfies the colimit axiom. Then λ above is an isomorphism in case A is a free abelian group. Moreover for a general abelian group $A \in \mathbf{Ab}$ there is the natural exact sequence ($n \geq 2, m \in \mathbb{Z}$):

$$\begin{aligned} 0 \rightarrow A *' Q_m \{S^n\} \xrightarrow{e} Q_{m+1}(M(A, n)) \xrightarrow{h} A *'' Q_{m-1} \{S^n\} \\ \xrightarrow{\partial} A \otimes Q_m \{S^n\} \xrightarrow{\lambda} Q_m(M(A, n)) \rightarrow Q_m(M(A, n)) \rightarrow 0 \end{aligned}$$

This result generalizes 9.3 in [5]. If Q does not satisfy the colimit axiom then proposition (3.12) is still true for finitely generated abelian groups.

Proof. If A is free abelian we see by (3.6) [5] or (6.4) below that λ is an isomorphism. Now let $A \in \mathbf{Ab}$ and let

$$d : X = M(A_1, n) \rightarrow Y = M(A_0, n)$$

be a map which realizes d in (3.10). Then the Moore space $M(A, n)$ is the mapping cone $M(A, n) = CX \cup_d Y$. Therefore we get by (3.4) the following commutative diagram in which the column and the row are exact.

$$\begin{array}{ccccccc} & & Q_m(X|X) & & & & \\ & & \downarrow d_2 & & & & \\ & & Q_m(X) \oplus Q_m(X|Y) & \searrow & & & \\ & & \downarrow d_1 & & d_1 & & \\ \xrightarrow{i} & Q_{m+1}(M(A, n)) & \xrightarrow{j} & Q_{m+1}(M(A, n), Y) & \xrightarrow{\partial} & Q_m(Y) & \xrightarrow{i} & Q_m(M(A, n)) \rightarrow \\ & & & \downarrow \delta & & \parallel & & \uparrow \lambda \\ & & & Q_{m-1}(X|X) & & A_0 \otimes Q_m \{S^n\} & \xrightarrow{q_*} & A \otimes Q_m \{S^n\} \\ & & & \downarrow d_2 & & & & \\ & & & Q_{m-1}(X) \oplus Q_{m-1}(X|Y) & & & & \end{array}$$

The map q_* is surjective and the kernel of q_* is the image of d_1 . Moreover using the definition of d_1, d_2 in (3.4) and (3.10) we get for d_1, d_2 in the diagram

$$\begin{aligned} \ker(d_1)/\text{im}(d_2) &= A *' Q_m \{S^n\} \\ \ker(d_2) &= A *'' Q_m \{S^n\} \end{aligned}$$

We now define the operators in (3.12) as follows. The inclusion e is induced by \bar{d}_1 where $\text{cok}(i) = \text{cok}(\lambda) = \text{im}(j)$. The map h is the restriction of δ . The map ∂ in the proposition is induced by $\partial(\delta)^{-1}$ and i and j in the proposition are derived from i and j in the row of the diagram.

q.e.d.

§ 4 HOMOTOPY FUNCTORS

Following Goodwillie [17, 18, 19, 20, 21] we consider for $r \geq 0$ functors D of the form

$$(4.1) \quad \begin{aligned} D : \mathbf{space}_r &\rightarrow \mathbf{space} \quad \text{or} \\ D : \mathbf{space}_r &\rightarrow \mathbf{spectra} \end{aligned}$$

We say that D is a *homotopy functor* if D carries weak equivalences to weak equivalences and preserves filtered colimits up to homotopy. That is, for each sequence of cofibrations $X_0 \rightarrow X_1 \rightarrow \dots$ in \mathbf{space}_r the induced map

$$\mathrm{hocolim}(DX_i) \rightarrow D(\mathrm{colim} X_i)$$

is a weak equivalence. Here $\mathrm{hocolim}$ is the homotopy colimit. We say that D is *reduced* if $*$ $\rightarrow D(*)$ is a weak equivalence. We shall consider various examples of such homotopy functors below.

(4.2) *Definition.* Let D be a homotopy functor as in (4.1). We define the D -homology of a space $X \in \mathbf{space}_r$ by the homotopy group

$$H_n(X; D) = \pi_n(D(X)).$$

Moreover the *relative D -homology* is the relative homotopy group

$$H_{n+1}(X, Y; D) = \pi_{n+1}(D(X), D(Y))$$

for $(X, Y) \in \mathbf{pair}_r$. This yields the natural boundary map

$$\partial : H_{n+1}(X, Y; D) \rightarrow H_n(Y; D).$$

If D maps to $\mathbf{spectra}$ then all such homotopy groups are well defined abelian groups for $n \in \mathbb{Z}$. If D maps to \mathbf{space} then these groups are abelian only for $n \geq 2$. For $n = 1$, resp. $n = 0$, the D -homology $H_n(X, D)$ and $H_{n+1}(X, Y; D)$ is a group, resp. a pointed set. If D maps to \mathbf{space}_1 we set $H_n(X, D) = H_n(X, Y; D) = 0$ for $n \leq 0$. If D is a reduced homotopy functor we obtain as in (2.4) the suspension

$$s = q_* \partial^{-1} : H_n(X; D) \rightarrow H_{n+1}(\Sigma X; D)$$

which defines the stable theory $H_n^S(X; D)$ of *stable homology groups* with coefficients in D .

(4.3) *Remark.* We have for a reduced homotopy functor D the *linearization* D^{lin} which is the homotopy functor obtained by the homotopy colimit of

$$D(X) \rightarrow \Omega D(\Sigma X) \rightarrow \Omega^2 D(\Sigma^2 X) \rightarrow \dots$$

We clearly have the natural isomorphism

$$H_n^S(X, D) = H_n(X, D^{\mathrm{lin}}).$$

Goodwillie [17] showed that $H_n^S(X; D)$ is a homology theory if D is approximately 1-excisive. This is also true if D is n -excisive for $n \geq 1$ as follows from 3.2.4 in [20].

The following lemma is an easy consequence of the definitions.

(4.4) Lemma. *Let D be a reduced homotopy functor which maps to \mathbf{space}_2 or $\mathbf{spectra}$. Then D -homology is a boundary theory which satisfies the colimit axiom in (2.1).*

Proof. The ∂ -exact sequence is obtained by (2.6) (B). Moreover homotopic maps $f \simeq g$ induce $f_* = g_*$ in D -homology since the projection $I(X) \rightarrow X$ of the cylinder is a homotopy equivalence and hence a weak equivalence.

q.e.d.

By an n -cube of spaces (or spectra) we will mean the following. Let $\mathbf{P}(n)$ be the category whose objects are the subsets K of $\{1, 2, \dots, n\}$ and whose morphisms are the inclusion maps among the subsets. An n -cube \underline{X} in a category \mathbf{C} is a covariant functor $\underline{X} : \mathbf{P}(n) \rightarrow \mathbf{C}$. Goodwillie defines and uses particular n -cubes in \mathbf{space} or $\mathbf{spectra}$, namely Cartesian and co-Cartesian n -cubes. Let “holim” be the homotopy inverse limit and “hocolim” be the homotopy colimit as defined in Bousfield-Kan [12]. Let $P'(n)$, resp. $P''(n)$, be the full subcategory of $\mathbf{P}(n)$ consisting of all K with $K \neq \emptyset$, resp. $K \neq \{1, \dots, n\}$, and let \underline{X}' and \underline{X}'' be the restrictions of \underline{X} to $P'(n)$, resp. $P''(n)$. There are maps

$$a(\underline{X}) : \underline{X}(\emptyset) = \lim(\underline{X}) \simeq \text{holim}(\underline{X}) \rightarrow \text{holim}(\underline{X}')$$

$$b(\underline{X}) : \text{hocolim}(\underline{X}'') \rightarrow \text{hocolim}(\underline{X}) \simeq \text{colim}(\underline{X}) = \underline{X}(\{1, \dots, n\})$$

Now \underline{X} is *Cartesian*, resp. *co-Cartesian*, if $a(\underline{X})$, resp. $b(\underline{X})$, is a weak equivalence. An n -cube is *strongly co-Cartesian* if each of its 2-faces is co-Cartesian. A Cartesian 2-cube is also called a *homotopy pull back* and a co-Cartesian 2-cube is a *homotopy push out*. A basic notion in [17, 18] is the following definition.

(4.5) Definition. Let D be a homotopy functor as in (4.1). Then D is termed *n -excisive* if $D(\underline{X})$ is Cartesian for every strongly co-Cartesian $(n+1)$ -cube \underline{X} in \mathbf{space}_r . We say that D is *linear* if D is reduced and 1-excisive and we say that D is *quadratic* if D is reduced and 2-excisive.

Goodwillie [17] proved the following result:

(4.6) Theorem. *Let D be a homotopy functor as in (4.1) which maps to \mathbf{space}_1 or $\mathbf{spectra}$ and let D be linear. Then D -homology is a homology theory.*

We obtain the corresponding result for the quadratic case as follows.

(4.7) Theorem. *Let D be a homotopy functor as in (4.1) which maps to \mathbf{space}_2 or $\mathbf{spectra}$ and let D be quadratic. Then D -homology is a quadratic homology theory on \mathbf{pair}_r in the sense of (3.2).*

Here we use \mathbf{space}_2 since we want all D -homology groups to be abelian.

Proof of (4.7). We obtain (3.2) (i) by (4.3) and (3.2) (ii) by (4.9) below; see also (4.11). Moreover (3.2) (iii) is proved in (4.12) below.

q.e.d.

We need the following lemma; compare 1.18 [18]. Let $\underline{X}, \underline{Y}$ be n -cubes in \mathbf{space} and let $f : \underline{X} \rightarrow \underline{Y}$ be a map between n -cubes which may be considered to be a $(n+1)$ -cube. Let $\text{hofib}(f)$ be the n -cube obtained by taking the homotopy fibers of $f(K) : X(K) \rightarrow Y(K)$, $K \in \mathbf{P}(n)$.

(4.8) **Lemma.** Assume $Y(\phi)$ is a connected space. Then f is a Cartesian $(n+1)$ -cube if and only if $\text{hofib}(f)$ is a Cartesian n -cube.

Below we shall also consider quadratic homotopy functors D which map to \mathbf{space}_1 and not necessarily to \mathbf{space}_2 as assumed in (4.7). For such homotopy functors the following two lemmas hold.

(4.9) **Lemma.** Let D be a quadratic homotopy functor as in (4.1) which maps to \mathbf{space}_1 or $\mathbf{spectra}$. Then D -homology defines for $n \in \mathbb{Z}$ a quadratic functor

$$H_n(-; D) : \mathbf{space}_r \rightarrow \mathbf{Gr}$$

which carries X to $H_n(X, D)$. Moreover the suspension theory of cross effects $H_n(X|Y; D)$ defined as in (2.11) is a homology theory.

The lemma is a consequence of Goodwillie's results. In fact, let $D(X \vee Y)_2$ be the homotopy fiber of $D(r_2) : D(X \vee Y) \rightarrow D(Y)$ and let $D(X|Y)$ be the homotopy fiber of $D(r_1) : D(X \vee Y)_2 \rightarrow D(X)$. Then one readily checks that one has

$$(4.10) \quad H_n(X|Y, D) = \pi_n D(X|Y)$$

for any reduced homotopy functor D . If D maps to \mathbf{space}_1 then $D(X \vee Y)_2$ is always a connected space. We now repeat an argument of Goodwillie which proves (4.9).

Proof of (4.9). We first show that the functor D_X with $D_X(Y) = D(X \vee Y)_2$ is 1-excisive. In fact, let \underline{Y} be a co-Cartesian 2-cube. Then $r_2 : X \vee \underline{Y} \rightarrow \underline{Y}$ is a strongly co-Cartesian 3-cube; compare (A.1). Hence $D(r_2) = D(X \vee \underline{Y} \rightarrow \underline{Y})$ is a Cartesian 3-cube since D is 2-excisive. Therefore (4.8) shows that $D_X(\underline{Y})$ is Cartesian and hence D_X is 1-excisive. Now $D(X|Y)$ is the homotopy fiber of $D_X(Y) \rightarrow D_X(*)$. Therefore the functor $Y \mapsto D(X|Y)$ is linear since D_X is 1-excisive. This completes the proof of (4.9) by using (4.6).

q.e.d.

(4.11) *Remark.* Let D be a quadratic homotopy functor as in (4.1) which maps to \mathbf{space}_1 . Goodwillie showed that there exists a spectrum E and a natural isomorphism

$$(1) \quad H_n(X|Y; D) = \pi_n(E \wedge X \wedge Y)$$

which is compatible with the-suspension s of X and the suspension \bar{s} of Y ; see (2.11), (2.12). Moreover the isomorphism is compatible with the interchange map on both sides. Here the interchange map T on $\pi_n(E \wedge X \wedge Y)$ is introduced by a Σ_2 -action t on E and by the interchange $T_{X,Y} : X \wedge Y \approx Y \wedge X$, that is $T = \pi_n(t \wedge T_{X,Y})$. Let $(E \wedge X \wedge X)_T$ be the homotopy orbit spectrum given by T . Then the Goodwillie tower of the quadratic functor D yields a natural fibration sequence

$$(2) \quad \Omega^\infty(E \wedge X \wedge X)_T \rightarrow D(X) \rightarrow D^{lin}(X).$$

Here the linearization satisfies $D^{lin}(X) = \Omega^\infty(E' \wedge X)$ for an appropriate spectrum E' .

(4.12) **Lemma.** Let D be a quadratic homotopy functor as in (4.1) which maps to \mathbf{space}_1 or $\mathbf{spectra}$. Then for $Q_n(X, Y) = H_n(X, Y; D)$ there exists a quadratic excision sequence as in (3.2) (iii) which is natural and exact.

We point out that in the lemma $Q_1(X, Y) = H_1(X, Y; D)$ is only a pointed set if D maps to \mathbf{space}_1 .

Proof of (4.12). Let (X, A) be an object in \mathbf{pair}_r and let $f : A \rightarrow X$ be the inclusion. Then (X, A) yields a 3-cube in \mathbf{space}_r termed $Cube(X, A)$ which is obtained as the following map F between 2-cubes

$$(1) \quad \begin{array}{ccc} A \vee X & \xleftarrow{1 \vee f} & A \vee A & & X & \xleftarrow{f} & A \\ (f, 1) \downarrow & & \downarrow (1, 1) & \xrightarrow{F} & \downarrow q & & \downarrow \\ X & \xleftarrow{f} & A & & X/A & \xleftarrow{\quad} & * \end{array}$$

Here F is defined by $(0, 1) : A \vee X \rightarrow X$, $(0, 1) : A \vee A \rightarrow A$, $q : X \rightarrow X/A$ and $0 : A \rightarrow *$. One readily checks that $F = Cube(X, A)$ is well defined. Each square in $Cube(X, A)$ is a homotopy push out so that $Cube(X, A)$ is actually strongly co-Cartesian. Since D is quadratic this implies that $D(Cube(X, A))$ is Cartesian and therefore by (4.8) the following diagram of homotopy fibers is a homotopy pull back.

$$(2) \quad \begin{array}{ccc} D(A \vee A)_2 & \xrightarrow{(1 \vee f)_*} & D(A \vee X)_2 \\ (1, 1)_* \downarrow & & \downarrow (f, 1)_* \\ D(A) & \xrightarrow{f_*} & P(q_*) \end{array}$$

Here $P(q_*)$ is the homotopy fiber of $q_* : D(X) \rightarrow D(X/A)$. Let $K(A)$ be the homotopy fiber of $(1, 1)_* : D(A \vee A)_2 \rightarrow D(A)$. Then we get the fiber sequence

$$(3) \quad K(A) \xrightarrow{(1 \vee f)_*} D(A \vee X)_2 \xrightarrow{(f, 1)_*} P(q_*)$$

since (2) is a homotopy pull back. This fiber sequence of spaces or spectra induces a long exact sequence of homotopy groups. The definition of $K(A)$ yields for $n \in \mathbb{Z}$ the following commutative diagram of short exact sequences of groups.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_n K(A) & \longrightarrow & \pi_n D(A \vee A)_2 & \xrightarrow{(1, 1)_*} & \pi_n D(A) \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ 0 & \longrightarrow & \text{kernel}(1, P) & \longrightarrow & \pi_n D(A) \oplus \pi_n D(A|A) & \xrightarrow{(1, P)} & \pi_n D(A) \longrightarrow 0 \end{array}$$

Here we use the isomorphism in (4.10) and (2.4) (4) and we point out that for $D : \mathbf{space}_r \rightarrow \mathbf{space}_1$ the fiber $D(A \vee X)_2$ is connected and that $\pi_1(D(A|A))$ is central and split in $\pi_1 D(A \vee X)_2$. Now an isomorphism of abelian groups

$$(4) \quad \Theta : \pi_n D(A|A) \cong \text{kernel}(1, P) = \pi_1 K(A)$$

is obtained by mapping y to $(Py, -y)$.

The boundary operator ∂ of the long exact sequence of homotopy groups associated to (3) yields the natural transformation

$$(5) \quad \delta : H_{n+1}(CA \cup_A X, X; D) = \pi_n P(q_*) \xrightarrow{\partial} \pi_{n-1} K(A) = H_{n-1}(A|A; D)$$

This completes the proof of the quadratic excision axiom. q.e.d.

We can use (4.12) for the proof of the following result.

(4.13) Proposition. *Let $D : \mathbf{space}_1 \rightarrow \mathbf{space}_1$ be a quadratic homotopy functor. Then we obtain by (3.1) and (4.9) the square group*

$$H_1\{S^1; D\} = (H_1(S^1; D) \xrightarrow{H} H_1(S^1|S^1; D) \xrightarrow{P} H_1(S^1, D))$$

and using the tensor product in (6.3) below there is a natural isomorphism

$$H_1(Y, D) \cong \pi_1(Y) \otimes H_1\{S^1; D\}$$

for 2-dimensional CW-complexes Y .

Proof. We may assume that $Y = CA \cup_A X$ where A and X have the homotopy type of one point union of 1-spheres. We now apply (4.12) and ∂ -exactness. This gives us the exact sequence of groups, $Q_1(X) = H_1(X, D)$,


$$Q_1(A \vee X)_2 \rightarrow Q_1(X) \rightarrow Q_1(Y) \rightarrow 0.$$

Since D preserves filtered colimits up to homotopy we get by [10] with $M = H_1\{S^1; D\}$:

$$\begin{aligned} Q_1(X) &= \pi_1(X) \otimes M \\ Q_1(A \vee X)_2 &= Q_1(A) \oplus Q_1(A|X) \\ &= \pi_1(A) \otimes M \oplus \pi_1(A)^{ab} \otimes \pi_1(X)^{ab} \otimes M_{ce} \end{aligned}$$

and therefore the result follows from (8.4) [10]. For this observe that $\pi_1(A) \xrightarrow{d} \pi_1(X) \xrightarrow{q} \pi_1(Y)$ has the property that q is surjective and the normal closure of image of d is the kernel of q . q.e.d.

(4.14) Example. It follows from Goodwillie's Calculus [19, 20] that there is a sequence of functors P_n from pointed spaces to pointed spaces and natural transformations



$$P_1(X) = Q(X) = \Omega^\infty \Sigma^\infty(X)$$

Here P_n are homotopy functors satisfying n -th order excision and the maps $X \rightarrow P_n(X)$ are $(n+1)k+1$ connected, where k is the connectivity of X . The functors P_n are uniquely determined by these universal properties. According to Goodwillie the tower above is termed the Taylor tower of the identity. Here P_1 is a linear homotopy functor and the homology

$$H_n(X, P_1) = \pi_n^S(X)$$

coincides with stable homotopy groups. Therefore the quadratic homotopy functor P_2 yields a canonical quadratic homology which we call *quadratic homotopy groups*:

$$H_n(X, P_2) = \pi_n^Q(X)$$

This is the quadratic analogue of stable homotopy groups. We know by (4.7) that quadratic homotopy groups π_n^Q form a quadratic homology theory on **space**₂ satisfying all properties in section 3. The cross effect of $\pi_n^Q(X)$ is given by a natural isomorphism

$$\pi_n^Q(X|Y) = \pi_{n+1}^S(X \wedge Y)$$

where $X \wedge Y = X \times Y / X \vee Y$, compare [19], [24]. Arone-Mahowald [1] mentioned that $P_2(X)$ also can be constructed by the fiber of the stable James-Hopf map γ_2 in [13]; that is

$$P_2(X) \rightarrow Q(X) \xrightarrow{\gamma_2} Q(X \wedge X)_T$$

is a natural fiber sequence. Here $(X \wedge X)_T$ is the homotopy orbit space of the $\mathbb{Z}/2$ -action on $X \wedge X$ given by the interchange map $T_{X,X}$. In the metastable range we have considered examples of the quadratic \mathbb{Z} -modules $\pi_{n+k}^Q\{S^n\}$ in table 2 in 9.9 [5]. Such quadratic \mathbb{Z} -modules seem to be the appropriate quadratic analogue of stable homotopy groups of spheres.

§ 5 HOMOTOPY FUNCTORS INDUCED BY ENDOFUNCTORS OF THE CATEGORY OF GROUPS

Let \mathbf{Gr} be the category of groups and let $s\mathbf{Gr}$ be the category of simplicial groups. A functor $F : \mathbf{Gr} \rightarrow \mathbf{Gr}$ induces the functor

$$\bar{F} : s\mathbf{Gr} \rightarrow s\mathbf{Gr}$$

which carries the simplicial group X to the simplicial group $F \circ X$. We now consider the following composition of functors $F_{\sharp} = \beta \bar{F} \alpha$:

$$(5.1) \quad \mathbf{space}_1 \xrightarrow{\alpha} s\mathbf{Gr} \xrightarrow{\bar{F}} s\mathbf{Gr} \xrightarrow{\beta} \mathbf{space}_1$$

Here we obtain α as follows. For a pointed space X let $S(X)$ be the reduced singular set consisting of all singular simplexes $\sigma : \Delta^n \rightarrow X$ with $\sigma(v) = *$ for all vertices v of the simplex Δ^n . Then the functor α carries X to the Kan-loop group $\alpha(X) = GS(X)$ of $S(X)$; see for example [15]. Moreover for a simplicial group G let $|G|$ be the realization and let $\beta(G) = B|G|$ be the classifying space of the topological group $|G|$. Then α and β induce equivalences of homotopy categories

$$\mathbf{space}_1 / \simeq \begin{array}{c} \xrightarrow{\alpha} \\ \xrightarrow{\beta} \end{array} Ho(s\mathbf{Gr})$$

where β is the inverse of α . Here $Ho(s\mathbf{Gr})$ is the localization of $s\mathbf{Gr}$ with respect to weak equivalences.

Let \mathbf{gr} be the full subcategory of \mathbf{Gr} consisting of free groups. Since $\alpha(X)$ above is actually a free simplicial group we see that F_{\sharp} depends only on the restriction $F_0 : \mathbf{gr} \rightarrow \mathbf{Gr}$ of F . On the other hand each functor $F_0 : \mathbf{gr} \rightarrow \mathbf{Gr}$ determines a unique extension $F_1 : \mathbf{Gr} \rightarrow \mathbf{Gr}$ with the property that F_1 preserves cokernels. Therefore we may assume that F in (5.1) preserves cokernels.

Moreover it is convenient to assume that the behaviour of F_{\sharp} on infinite CW-complexes is determined by its behaviour on finite complexes. Therefore we assume that F_{\sharp} preserves filtered colimits up to homotopy. One readily checks that this is the case if and only if F preserves filtered colimits. These remarks lead to the

(5.3) *Definition.* A *group functor* is an endofunctor $F : \mathbf{Gr} \rightarrow \mathbf{Gr}$ of the category of groups which preserves cokernels and filtered colimits. Hence such a group functor is determined by its restriction to the full subcategory of finitely generated free groups.

(5.4) **Lemma.** A group functor $F : \mathbf{Gr} \rightarrow \mathbf{Gr}$ induces a functor $F_{\sharp} : \mathbf{space}_1 \rightarrow \mathbf{space}_1$ which is a reduced homotopy functor.

We point out that the lemma does not imply that the functor \bar{F} in (5.1) carries weak equivalences in $s\mathbf{Gr}$ to weak equivalences in $s\mathbf{Gr}$; this in general does not hold.

Proof of (5.4). Clearly $* \rightarrow F_{\sharp}(*)$ is a homotopy equivalence so that F_{\sharp} is reduced. We have to show that F_{\sharp} carries homotopy equivalences to weak equivalences. Let $H : f_0 \simeq f_1$ be a homotopy in \mathbf{space}_1 . Then there exists a homotopy $H' : \alpha f_0 \simeq \alpha f_1$ since $\alpha(X)$ is a free simplicial group. The functor \bar{F} carries H' to a homotopy $\bar{F}\alpha f_0 \simeq \bar{F}\alpha f_1$ in the category of simplicial sets, see (1.10) and (4.2) in [15]. This implies that $\pi_n(\bar{F}\alpha f_0) = \pi_n(\bar{F}\alpha f_1)$ and hence $\pi_n(F_{\sharp}f_0) = \pi_n(F_{\sharp}f_1)$ for $n \in \mathbb{Z}$. From this one readily derives that F_{\sharp} carries homotopy equivalences to weak equivalences. q.e.d.

(5.5) *Definition.* Let $F : \mathbf{Gr} \rightarrow \mathbf{Gr}$ be a group functor and let $X \in \mathbf{space}_1$. We define the F -homology of X ,

$$H_n(X; F) = H_n(X; F_{\sharp}) = \pi_n(F_{\sharp}X),$$

by the F_{\sharp} -homology of X in (4.2), $n \in \mathbb{Z}$. Similarly we obtain the relative F -homology $H_n(X, Y; F)$ of a pair $(X, Y) \in \mathbf{pair}_1$. Clearly F -homology has the properties of D -homology described in (4.2) and (4.4). In particular $H_1(X; F)$ is only a group; see (5.11). Moreover we obtain the *stable F -homology*

$$H_n^S(X, F) = H_n(X, F_{\sharp}^{lin})$$

Compare (4.3). Let $F, G : \mathbf{Gr} \rightarrow \mathbf{Gr}$ be group functors. Then a natural transformation $t : F \rightarrow G$ induces natural transformations

$$\begin{aligned} t_{\sharp} : F_{\sharp}(X) &\rightarrow G_{\sharp}(X) \quad \text{and} \\ t_{\star} : H_n(X, Y; F) &\rightarrow H_n(X, Y; G) \end{aligned}$$

where t_{\star} is the *coefficient homomorphism* induced by t_{\sharp} . We obtain t_{\sharp} as follows. Let $\bar{t} : \bar{F} \rightarrow \bar{G}$ be the transformation induced by F . Then we set $t_{\sharp} = \beta \bar{t}_{\alpha X}$. Clearly the coefficient homomorphism t_{\star} is compatible with the boundary map ∂ of F -homology.

(5.6) **Lemma.** Let $(X, Y) \in \mathbf{pair}_1$ be an r -connected pair. Then $H_n(X, Y; F) = 0$ for $n \leq r$. This implies that

$$H_n(X; F) = H_n(X^{n+1}; F)$$

depends only on the $(n + 1)$ -skeleton of X . Moreover if X is r -connected then so is $F_{\sharp}(X)$.

Proof. We may assume that X is a reduced CW-complex with subcomplex Y and that $X - Y$ has only cells in dimension $> r$. By a result of Kan [25] we obtain a homotopy equivalence

$$\varphi : (G_X, G_Y) \simeq (\alpha X, \alpha Y)$$

of pairs of free simplicial groups. Here G_X as a free simplicial group has generators in degree t which are exactly the $(t + 1)$ -cells of X , $t \geq 0$. Moreover G_Y is the subobject generated by the cells of Y . Hence $(G_X)_n = (G_Y)_n$ for $n < r$ and therefore also $(\bar{F}G_X)_n = (\bar{F}G_Y)_n$ for $n < r$. Since \bar{F} carries φ above to a weak equivalence we see that $H_n(X, Y; F) = 0$ for $n \leq r$.

q.e.d.

Let I be the identity functor of the category \mathbf{Gr} . Then one has the canonical natural isomorphism

$$(5.7) \quad \pi_n(X, Y) = H_n(X, Y; I)$$

which we use as an identification. For a group G let $\Gamma_k(G) \subset G$ be the subgroup of k -fold commutators. Then $G/\Gamma_2 G = \text{ab}(G) = G^{\text{ab}}$ is the abelianization of G . For $k \geq 1$ we obtain the *nilization functors*

$$(5.8) \quad \text{nil}_k : \mathbf{Gr} \rightarrow \mathbf{Gr}$$

which carry G to the quotient $\text{nil}_k(G) = G/\Gamma_{k+1}G$. Hence $\text{nil}_1 = \text{ab}$ is the abelianization functor. All functors nil_k , $k \geq 1$, are group functors in the sense of (5.3). By the result of Dold-Kan it is well known that for $(X, Y) \in \text{pair}_1$ one has the natural isomorphism

$$(5.9) \quad H_n(X, Y; \text{ab}) = H_n(X, Y)$$

where the right hand side is the *singular homology*. Moreover the natural transformation $I \rightarrow \text{ab}$ given by the quotient map $G \rightarrow \text{ab}(G)$ induces the classical *Hurewicz homomorphism*

$$\pi_n(X, Y) = H_n(X, Y; I) \rightarrow H_n(X, Y; \text{ab}) = H_n(X, Y)$$

Similarly one obtains by the natural quotient map $I \rightarrow \text{nil}_k(G)$, the *nil_k-Hurewicz homomorphism*

$$\pi_n(X, Y) = H_n(X, Y; I) \rightarrow H_n(X, Y; \text{nil}_k)$$

The following generalization of the classical Hurewicz theorem is due to Curtis [15].

(5.10) Curtis theorem. *Let $r \geq 2$ and let X be an $(r-1)$ -connected space. Then the nil_k -Hurewicz homomorphism*

$$\pi_n(X) \rightarrow H_n(X; \text{nil}_k)$$

is an isomorphism for $n < r + \{\log_2(k+1)\}$ and is surjective for $n = r + \{\log_2(k+1)\}$. Here $\{a\}$ denotest the least integer $\geq a$.

The Hurewicz theorem is the case $k = 1$ of this result since $\{\log_2(2)\} = \{1\} = 1$. For $k = 2$ we have $\{\log_2(3)\} = 2$ since $2 > \log_2(3) > 1$. Hence the nil_2 -Hurewicz homomorphism yields for an $(r-1)$ -connected space X , $r \geq 2$,

$$(5.11) \quad \begin{aligned} \pi_r(X) &= H_r(X, \text{nil}_2) \\ \pi_{r+1}(X) &= H_{r+1}(X, \text{nil}_2) \\ \pi_{r+2}(X) &\twoheadrightarrow H_{r+2}(X, \text{nil}_2) \end{aligned}$$

where \twoheadrightarrow denotes a surjection. Clearly the functor $\text{nil}_1 = \text{ab}$ is linear and the functor nil_2 is quadratic in the sense of (2.9). In the next section we consider all quadratic group functors $\mathbf{Gr} \rightarrow \mathbf{Gr}$. It would be interesting to understand the stable theory $H_n^S(X, \text{nil}_k)$, $k \geq 1$, which is a homology theory approximating for $k \rightarrow \infty$ stable homotopy $\pi_n^S(X)$. We shall determine $H_n^S(X, F)$ for any quadratic group functor in (8.5). This in particular yields an explicit spectrum E for which $H_n^S(X, \text{nil}_2) = H_n(X, E)$; compare (8.16).

The next result corresponds to (4.13) in case F is a quadratic group functor.

(5.12) Lemma. *Let $F : \mathbf{Gr} \rightarrow \mathbf{Gr}$ be a group functor. Then we have for $X \in \text{space}_1$ the natural isomorphism*

$$H_1(X, F) = F(\pi_1(X)).$$

Moreover if S is a one point union of 1-spheres we have

$$H_n(S, F) = 0 \quad \text{for } n \geq 2.$$

Proof. The group $G = \pi_1(S)$ is a free group for which the constant simplicial group \bar{G} with $\bar{G}_n = G$ for $n \geq 0$ is a free simplicial group which is homotopy equivalent to $\alpha(S)$. Hence one has a weak equivalence $\bar{F}(\bar{G}) \rightarrow \bar{F}\alpha(S)$ where $\bar{F}(\bar{G})$ is again a constant simplicial group. This shows $H_n(S, F) = \pi_{n-1}\bar{F}(\bar{G}) = 0$ for $n \geq 2$ and $H_1(S, F) = \pi_1\bar{F}(\bar{G}) = F(G)$. Now let $H = \alpha X$ with $\pi_0 H = \pi_1 X$. Then the degree 1 part of the simplicial group H

$$H_1 \rightrightarrows H_0 \rightarrow \pi_0 H$$

is a coequalizer. Since F preserves cokernels also

$$FH_1 \rightrightarrows FH_0 \rightarrow F\pi_0 H$$

is a coequalizer and therefore $\pi_0 FH = F\pi_0 H$. This shows $H_1(X, F) = F(\pi_1 X)$.

q.e.d.

If $F = ab$ is the abelianization functor we have by (5.9) and (5.11)

$$H_1(X) = H_1(X, ab) = ab(\pi_1(X))$$

This is a part of the classical Hurewicz theorem.

§ 6 SQUARE-HOMOLOGY

We show that the homology theories obtained by linear group functors $\mathbf{Gr} \rightarrow \mathbf{Gr}$ are exactly the classical “ordinary homology theories” of Eilenberg-Mac Lane. This motivates the study of homology defined by quadratic group functors $\mathbf{Gr} \rightarrow \mathbf{Gr}$. There is the following classification of linear functors in [10].

(6.1) Lemma. *The category of linear group functors $\mathbf{Gr} \rightarrow \mathbf{Gr}$ is equivalent to the category \mathbf{Ab} of abelian groups. More precisely for each linear group functor F there is an abelian group A and an isomorphism*

$$F(G) = ab(G) \otimes A$$

which is natural in $G \in \mathbf{Gr}$. The equivalence carries F to A . In particular F admits a factorization

$$F : \mathbf{Gr} \xrightarrow{ab} \mathbf{Ab} \xrightarrow{\otimes A} \mathbf{Ab} \subset \mathbf{Gr}.$$

As a consequence of this result and the Dold-Kan theorem [16] we get:

(6.2) Proposition. Let F be a linear group functor corresponding to $A \in \mathbf{Ab}$ as in (6.1). Then one has for $(X, Y) \in \mathbf{pair}_1$ the natural isomorphism, $n \in \mathbb{Z}$,

$$H_n(X, Y; F) = H_n(X, Y; A)$$

where the right hand side is the singular homology with coefficients in A .

This implies that the homology theories of linear group functors are exactly the ordinary homology theories in the sense of Eilenberg-Steenrod which satisfy the classical "dimension axiom". As a next step we consider quadratic group functors. For this recall from [10] the following notation on square groups.

(6.3) Definition. A square group

$$M = (M_e \xrightarrow{H} M_{ee} \xrightarrow{P} M_e)$$

is given by a group M_e and an abelian group M_{ee} . Both groups are written additively. Moreover P is a homomorphism and H is a quadratic function, that is the cross effect

$$(a|b)_H = H(a + b) - H(b) - H(a)$$

is linear in $a, b \in M_e$. In addition the following properties are satisfied ($x, y \in M_{ee}$).

- (1) $(Px|b)_H = 0$ and $(a|Py)_H = 0$
- (2) $P(a|b)_H = a + b - a - b$
- (3) $PHP(x) = P(x) + P(x)$

By (1) and (2) P maps to the center of M_e and by (2) the cokernel of P is abelian. Hence M_e is a group of nilpotency degree 2. Let **Square** be the category of square groups. As an example we have the square group

$$\mathbb{Z}_{nil} = (\mathbb{Z} \xrightarrow{H} \mathbb{Z} \xrightarrow{0} \mathbb{Z})$$

with $H(r) = \binom{r}{2}$ and $P = 0$; many other examples are discussed in [9, 10]. A quadratic \mathbb{Z} -module M is a square group for which H is linear and $HPH = 2H$; see (3.1).

Let G be a group and let M be a square group. Similarly as in (3.10) we define the group $G \otimes M$ by the generators $g \otimes a$ and $[g, h] \otimes x$ with $g, h \in G$, $a \in M_e$ and $x \in M_{ee}$ subject to the relations

$$\begin{aligned} (g + h) \otimes a &= g \otimes a + h \otimes a + [g, h] \otimes H(a) \\ [g, g] \otimes x &= g \otimes P(x) \end{aligned}$$

where $g \otimes a$ is linear in a and where $[g, h] \otimes x$ is central and linear in each variable g, h and x . There are obvious induced maps for this tensor product so that one gets a bifunctor

$$\otimes : \mathbf{Gr} \times \mathbf{Square} \rightarrow \mathbf{Gr}$$

One has the natural isomorphism

$$\text{nil}_2(G) = G \otimes \mathbb{Z}_{\text{nil}}.$$

In [5, 10] we obtain the following classification of quadratic group functors which is the quadratic analog of (6.1). Let \mathbf{Nil} be the category of groups of nilpotency degree 2.

(6.4) Proposition. *The category of quadratic group functors $\mathbf{Gr} \rightarrow \mathbf{Gr}$ is equivalent to the category \mathbf{Square} of square groups. More precisely for each quadratic group functor F one has the square group $M = F\{\mathbb{Z}\}$ in (3.1) and isomorphisms*

$$F(G) = G \otimes M = \text{nil}_2(G) \otimes M$$

which are natural in $G \in \mathbf{Gr}$. The equivalence carries F to M and $F = \text{nil}_2$ corresponds to $M = \mathbb{Z}_{\text{nil}}$. In particular quadratic group functors admit a factorization

$$\mathbf{Gr} \xrightarrow{\text{nil}_2} \mathbf{Nil} \xrightarrow{\otimes M} \mathbf{Nil} \subset \mathbf{Gr}$$

Moreover the quadratic group functors which admit a factorization

$$\mathbf{Gr} \xrightarrow{ab} \mathbf{Ab} \longrightarrow \mathbf{Ab} \subset \mathbf{Gr}$$

correspond exactly to quadratic \mathbb{Z} -modules M .

Using this result we identify square groups M and quadratic group functors $\mathbf{Gr} \rightarrow \mathbf{Gr}$; we write $M : \mathbf{Gr} \rightarrow \mathbf{Gr}$ for the functor which carries G to $G \otimes M$. As a quadratic analog of (6.2) we define for a square group M and $(X, Y) \in \mathbf{pair}_1$ the *square-homology with coefficients in M* , $n \in \mathbb{Z}$,

$$(6.5) \quad H_n(X, Y; M) = H_n(X, Y; M_{\sharp}) = \pi_n(M_{\sharp}X, M_{\sharp}Y).$$

Here M_{\sharp} is the homotopy functor (5.1) induced by M . The groups $H_n(X; M)$ and $H_{n+1}(X, Y; M)$ are abelian for $n \geq 2$ and of nilpotency degree 2 for $n = 1$. Moreover $H_1(X, Y; M)$ is a pointed set. For $n \leq 0$ all groups (6.5) are trivial.

(6.6) Theorem. *Let $F : \mathbf{Gr} \rightarrow \mathbf{Gr}$ be a quadratic group functor. Then F_{\sharp} in (5.1) is a quadratic homotopy functor.*

We prove this result in the appendix A. Since for a simply connected space X also $F_{\sharp}X$ is simply connected we obtain by (6.6), (6.4), (4.7) the corollary:

(6.7) Corollary. *Square-homology $H_n(X, Y; M)$ with coefficients in a square group M is a quadratic homology theory on \mathbf{pair}_2 in the sense of (3.2).*

Using (4.3) we derive from (6.6) the next corollary which yields the homology theory or spectrum associated to a square group.

(6.8) Corollary. *Stable square-homology $H_n^S(X, Y; M)$ with coefficients in a square group M is a homology theory on \mathbf{pair}_1 .*

In (6.7) we use the category \mathbf{space}_2 in order to obtain abelian groups $H_n(X, Y; M)$. The next result improves (6.7) for the category \mathbf{space}_1 .

(6.9) Theorem. *Let M be a square group. Then square-homology with coefficients in M defines a quadratic functor*

$$H_n(-; M) : \mathbf{space}_1 \rightarrow \begin{cases} \mathbf{Ab} & \text{for } n \geq 2 \\ \mathbf{Nil} & \text{for } n = 1 \end{cases}$$

with cross effect

$$H_n(X|Y; M) = H_{n+1}(X \wedge Y; M_{ee})$$

for $n \in \mathbb{Z}$. Here the right hand side is the singular homology of the smash product $X \wedge Y$ with coefficients in the abelian group M_{ee} . The isomorphism is compatible up to sign with the suspension s and \bar{s} and the interchange map on both sides. Moreover for $(X, A) \in \mathbf{pair}_1$ one has the natural exact sequence of groups:

$$\begin{array}{ccccccc} H_{n-1}(A; M) \oplus H_n(A \wedge X; M_{ee}) & & & & & & \\ \parallel & & & & & & \\ \xrightarrow{\delta} H_n(A \wedge A; M_{ee}) & \xrightarrow{j_{\sharp}} & H_n(CA \vee X, A \vee X; M) & \xrightarrow{i_{\sharp}} & H_n(CA \cup_A X, X; M) & \xrightarrow{\delta} & H_{n-1}(A \wedge A; M_{ee}) \end{array}$$

The operators i_{\sharp} and j_{\sharp} are defined as in (3.2) (iii). For $n \leq 1$ the sequence consists of trivial groups. As in (3.5) we derive from the quadratic excision sequence in (6.9) the following long exact sequence.

(6.10) Corollary. *Let M be a square group and $A \in \mathbf{space}_1$. Then one has the natural long exact sequence of groups, $n \in \mathbb{Z}$,*

$$\xrightarrow{P} H_n(A; M) \xrightarrow{s} H_{n+1}(\Sigma A; M) \xrightarrow{H'} H_n(A \wedge A; M_{ee}) \xrightarrow{P} H_{n-1}(A; M) \rightarrow$$

Proof of (6.9). In the following proof we do not use (6.6). Let $F : \mathbf{Gr} \rightarrow \mathbf{Gr}$ be given by $F(G) = G \otimes M$ where M is the square group. By [10] we have for $X, Y \in \mathbf{Gr}$ the binatural isomorphism

$$(1) \quad F(X|Y) = ab(X) \otimes ab(Y) \otimes M_{ee}$$

We now consider the following commutative cubical diagram in \mathbf{Gr} which is determined by the inclusion

$$(2) \quad f : A \rightarrow A \vee C = X$$

where X is the sum of A and C in \mathbf{Gr} . The pair (X, A) yields the three cube $G = \mathit{Cube}(X, A)$ as in (4.12):

$$(3) \quad \begin{array}{ccccc} A \vee X & \xleftarrow{1 \vee f} & A \vee A & & X & \xleftarrow{f} & A \\ (f,1) \downarrow & & \downarrow (1,1) & \xrightarrow{G} & \downarrow q & & \downarrow \\ X & \xleftarrow{f} & A & & X/A & \xleftarrow{\quad} & * \end{array}$$

Here the map G between 2-cubes is given by $(0,1) : A \vee X \rightarrow X$, $(0,1) : A \vee A \rightarrow A$, $q = (0,1) : A \vee C = X \rightarrow X/A = C$ and $0 : A \rightarrow *$. One readily checks that $Cube(X, A)$ is well defined. We now apply F to $Cube(X, A)$ and we obtain the following square consisting of the kernels of all $F(g)$ where g is one of the arrows in G .

$$(4) \quad \begin{array}{ccc} F(A \vee A)_2 & \xrightarrow{(1 \vee f)_*} & F(A \vee X)_2 \\ (1,1)_* \downarrow & & \downarrow (1,1)_* \\ F(A) & \xrightarrow{f_*} & \text{kernel}(Fq) \end{array}$$

Using the assumption that $X = A \vee C$ we see by (1) that $(1 \vee f)_*$ in (4) induces an isomorphism of the kernels of the vertical arrows in (4). Hence (4) is a pull back diagram in \mathbf{Gr} . Now let

$$(5) \quad f : A \rightarrow X$$

be a cofibration of simplicial groups. Then we know for all n that we can choose free groups C_n such that

$$(6) \quad X_n = A_n \vee C_n.$$

Hence f in (4) is of the form (2) for each n . This implies by the naturality of (3) and (4) that also (4) with f as in (5) is a pull back diagram in the category of simplicial groups. This yields the short exact sequence of simplicial groups

$$(7) \quad 0 \rightarrow K(A) \xrightarrow{(1 \vee f)_*} F(A \vee X)_2 \xrightarrow{(f,1)_*} \text{kernel}(Fq) \rightarrow 0,$$

$$(8) \quad K(A) = \text{kernel}\{(1,1)_* : F(A \vee A)_2 \rightarrow F(A)\}.$$

Using F_1 in (5.1) we see that $H_n(X'; M) = \pi_{n-1}(FX)$ where $X = \alpha X'$ is the simplicial group associated to $X' \in \mathbf{space}_r$. Hence we get by (1) the cross effect formula

$$(9) \quad \begin{aligned} H_n(X'|Y'; M) &= \pi_{n-1}(F(X|Y)) \\ &= \pi_{n-1}(ab(X) \otimes ab(Y) \otimes M_{ee}) \\ &= H_{n+1}(X' \wedge Y'; M_{ee}) \end{aligned}$$

Here the last isomorphism is given by the Eilenberg-Zilber theorem. This shows that the cross effect formula in (6.9) is satisfied. It only remains to check the quadratic excision sequence. Now a pair (X', A') in \mathbf{pair}_1 , corresponds to a cofibration $A \rightarrow X$ in the category of simplicial groups for which we have the short exact sequence (7). This sequence induces a long exact sequence of homotopy groups which is isomorphic to the quadratic excision sequence for (X', A') . The boundary ∂ for (7) determines the natural operator

$$(10) \quad \delta : H_{n+1}(CA' \cup X'; M) = \pi_{n-1} \text{kernel}(Fq) \xrightarrow{\partial} \pi_{n-2} K(A) \xrightarrow{\Theta} \pi_{n-2} F(A|A) = H_{n-1}(A'|A'; M)$$

Here the definition of $K(A)$ in (8) yields the split short exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_n K(A) & \longrightarrow & \pi_n F(A \vee A)_2 & \xrightarrow{(1,1)_*} & \pi_n F(A) \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ 0 & \longrightarrow & \text{kernel}(1, P) & \longrightarrow & \pi_n(FA) \oplus \pi_n F(A|A) & \xrightarrow{(1,P)} & \pi_n F(A) \longrightarrow 0 \end{array}$$

where we use the isomorphism in (2.9) (4). Now the isomorphism Θ in (10)

$$(11) \quad \Theta : \pi_n(F(A|A)) \cong \text{kernel}(1, P) = \pi_n K(A)$$

is obtained by mapping y to $(Py, -y)$. This corresponds to the definition of j_{\sharp} in (3.2) (iii). q.e.d.

The first non vanishing homology with coefficients in a square group is described by the following result.

(6.11) Proposition. *Let M be a square group and let $M^{add} = \text{cok}(P) \in \underline{Ab}$ be the cokernel of $P : M_{ee} \rightarrow M_e$. If X is an $(r-1)$ -connected space one has*

$$H_r(X, M) = \begin{cases} \pi_1(X) \otimes M & \text{for } r = 1 \\ \pi_r(X) \otimes M^{add} & \text{for } r > 1 \end{cases}$$

Proof. For $r = 1$ this is a consequence of (5.11) or (4.13). Now let $r \geq 2$. The EHP-sequence shows that for a one point union S of 1-spheres the sequence

$$ab(G) \otimes ab(G) \otimes M_{ee} \xrightarrow{P} G \otimes M \rightarrow H_2(\Sigma S; M) \rightarrow 0$$

is exact where $G = \pi_1(S)$. This implies

$$H_2(\Sigma S; M) = ab(G) \otimes M^{add}$$

Moreover the EHP-sequence shows that $H_2(\Sigma S; M) = H_n(\Sigma^{n-1} S; M)$ for $n \geq 2$. Now we have

$$H_r(X; M) = H_r(X^{r+1}; M).$$

Here X^{r+1} is the mapping cone of a maps $f : S' \rightarrow S''$ where S' and S'' are one point unions of r -spheres. Now the ∂ -exactness and the quadratic excision shows $H_r(X^{r+1}; M) = \pi_r(X) \otimes M^{add}$.

q.e.d.

§ 7 SQUARE HOMOLOGY WITH COEFFICIENTS IN A QUADRATIC \mathbb{Z} -MODULE

Let $s\mathbf{Ab}$ be the category of simplicial abelian groups and let \mathbf{Chain} be the category of chain complexes C of abelian groups with $C_i = 0$ for $i < 0$. It is a result of Dold and Kan that there are isomorphisms of categories (see [9])

$$(7.1) \quad s\mathbf{Ab} \underset{K}{\overset{N}{\rightleftarrows}} \mathbf{Chain}$$

where K is the inverse of the normalization N . Here $N(A)$ is also termed the Moore chain complex of A . Kan [26] 15.1 showed that for a reduced simplicial set X and its Kan loop group $G(X)$ one has the isomorphism of chain complexes

$$(7.2) \quad N(A(X)) \cong s^{-1}\tilde{C}_*X$$

Here $A(X) = abG(X)$ is the abelianization of $G(X)$ and \tilde{C}_*X is the reduced (normalized) chain complex of the simplicial set X . Recall that the *suspension* s^kC with $k \in \mathbb{Z}$ is $(s^kC)_n = C_{n-k}$ with the differential $d(s^kx) = (-1)^k s^k(dx)$. We define for $X \in \mathbf{space}_1$ the *singular chain complex*

$$(7.3) \quad \tilde{C}_*X = \tilde{C}_*(SX)$$

where SX is the reduced singular set in (5.1). Given a quadratic \mathbb{Z} -module M we obtain the induced chain functor M_{\sharp} which is the composite

$$M_{\sharp} : \mathbf{Chain} \xrightarrow{K} s\mathbf{Ab} \xrightarrow{\otimes M} s\mathbf{Ab} \xrightarrow{N} \mathbf{Chain}$$

The next result is a consequence of (7.1), (7.2) and (6.4).

(7.4) Proposition. *Let M be a quadratic \mathbb{Z} -module and $X \in \mathbf{space}_1$. Then the square homology $H_*(X, M)$ is determined by \tilde{C}_*X . More precisely there is a natural isomorphism, $n \in \mathbb{Z}$,*

$$H_n(X, M) = H_{n-1}(M_{\sharp}(s^{-1}\tilde{C}_*X))$$

Proof. Using the definitions we get

$$\begin{aligned}
H_n(X, M) &= \pi_n(M_{\sharp} X) \\
&= \pi_{n-1}((GSX) \otimes M) \\
(*) \quad &= \pi_{n-1}((GSX)^{ab} \otimes M) \\
&= \pi_{n-1}((KN(GSX)^{ab}) \otimes M) \\
&= \pi_{n-1}((K s^{-1} \tilde{C}_{\star} X) \otimes M) \\
&= H_{n-1} N((K s^{-1} \tilde{C}_{\star} X) \otimes M) \\
&= H_{n-1} M_{\sharp}(s^{-1} \tilde{C}_{\star} X)
\end{aligned}$$

In (*) we use the fact that for a group G we have $G \otimes M = G^{ab} \otimes M$ if M is a quadratic \mathbb{Z} -module; compare (6.4).

q.e.d.

(7.5) *Remark.* The homotopy type of the chain complex $\tilde{C}_{\star} X$ is determined by the homology $H_{\star}(X, \mathbb{Z})$. This shows by (7.4) that the abelian groups $H_n(X, M)$ are determined by $H_{\star}(X, \mathbb{Z})$ and M provided M is a quadratic \mathbb{Z} -module. This is not true if M is a square group. For example we obtain by (5.11)

$$0 = H_3(\mathbb{C}P_2, \mathbb{Z}_{nil}) \neq H_3(S^2 \vee S^4, \mathbb{Z}_{nil}) = \mathbb{Z}$$

with $H_{\star}(\mathbb{C}P_2, \mathbb{Z}) = H_{\star}(S^2 \vee S^4, \mathbb{Z})$. Here $\mathbb{C}P_2$ is the complex projective plane and $S^2 \vee S^4$ is the one point union of spheres.

(7.6) *Remark.* The universal coefficient formula in [11] can be used to compute $H_{\star}(X, M)$ if M is a quadratic \mathbb{Z} -module. For example if M_{ee} is torsion free one has the natural short exact sequence

$$0 \rightarrow (\bar{H}_{\star} \otimes M)_{n-1} \rightarrow H_n(X, M) \rightarrow (\bar{H}_{\star} \ast' M)_{n-2} \rightarrow 0$$

where $\bar{H}_{\star} = s^{-1} \tilde{H}_{\star}(X, \mathbb{Z})$ is the desuspended reduced homology of X . Here we use the graded tensor and torsion products defined in [11].

Now let M be a square group. Then there is the canonical short exact sequence

$$(7.7) \quad 0 \rightarrow \hat{M} \xrightarrow{i} M \xrightarrow{q} M^{add} \rightarrow 0$$

in the category **Square** where $M^{add} = \text{cok}(P)$ is an abelian group and where \hat{M} is a quadratic \mathbb{Z} -module. More precisely we obtain the commutative diagram

$$\begin{array}{ccccccc}
\hat{M} = & (im(P) & \xrightarrow{H} & M_{ee} & \xrightarrow{P} & im(P)) \\
\downarrow & \cap & & \parallel & & \cap \\
M = & (M_e & \xrightarrow{H} & M_{ee} & \xrightarrow{P} & M_e) \\
\downarrow & \downarrow & & \downarrow & & \downarrow q \\
M^{add} = & (cok(P) & \longrightarrow & 0 & \longrightarrow & cok(P))
\end{array}$$

where q is the quotient map. The definition of a square group shows readily that the restriction of H in M to $\hat{M}_e = im(P)$ is a homomorphism and that \hat{M} is a quadratic \mathbb{Z} -module. If G is a free group then (7.7) induces the short exact sequence of groups

$$(1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & G \otimes \hat{M} & \longrightarrow & G \otimes M & \longrightarrow & G \otimes M^{add} \longrightarrow 0 \\ & & \parallel & & & & \parallel \\ & & A \otimes \hat{M} & & & & A \otimes M^{add} \end{array}$$

where $A = ab(G)$. If G is a free simplicial group this sequence is still well defined and short exact. Hence the long exact sequence of homotopy groups for $G = G(SX)$ yields the following long exact sequence of square homology groups

$$(2) \quad \xrightarrow{\partial} H_n(X, \hat{M}) \xrightarrow{i_*} H_n(X, M) \xrightarrow{q_*} H_n(X, M^{add}) \xrightarrow{\partial} H_{n-1}(X, \hat{M}) \longrightarrow$$

Here $H_*(X, M^{add})$ is the usual homology of X with coefficients in the abelian group $M^{add} = cok(P : M_{ee} \rightarrow M_e)$. Moreover $H_*(X, \hat{M})$ depends by (7.5) only on the homology $H_*(X, \mathbb{Z})$ and \hat{M} . In fact $H_*(X, \hat{M})$ can be computed by the universal coefficient theorem [11] as follows; see (7.6).

Given a square group M we obtain the involution

$$T = HP - 1 : M_{ee} \rightarrow M_{ee}$$

with $TT = 1$. This yields the chain complex of groups

$$M_*^e = (M_e \xleftarrow{P} M_{ee} \xleftarrow{1-T} M_{ee} \xleftarrow{1+T} M_{ee} \xleftarrow{1-T} \dots)$$

where M_e is in degree 0. We have $1 - T = 2 - HP$ and $1 + T = HP$ so that the homology of M_*^e is

$$(7.8) \quad H_n(M_*^e) = \begin{cases} cok(P) & n = 0 \\ ker(P)/im(2 - HP) & n = 1 \\ ker(2 - HP)/im(HP) & n = 2k \geq 2 \\ ker(HP)/im(2 - HP) & n = 2k + 1 \geq 3 \end{cases}$$

Moreover we associate with M the following quadratic \mathbb{Z} -modules $Z_n M_*$, $n \geq 1$.

$$(7.9) \quad Z_n M_* = \begin{cases} ker(P) \xrightarrow{j} M_{ee} \xrightarrow{2-HP} ker(P) & n = 1 \\ ker(2 - HP) \xrightarrow{j} M_{ee} \xrightarrow{HP} ker(2 - HP) & n = 2k \geq 2 \\ ker(HP) \xrightarrow{j} M_{ee} \xrightarrow{2-HP} ker(HP) & n = 2k + 1 \geq 3 \end{cases}$$

Here j denotes the inclusion. We now compute the square-homology of a Moore space $M(A, n)$ of an abelian group A .

(7.10) Proposition. Let $n \geq 2$ and let M be a square group. Then the square homology of the Moore space $M(A, n)$, $A \in \mathbf{Ab}$, has the following properties. If A is free abelian there is a natural isomorphism

$$H_{n+k}(M(A, n); M) = \begin{cases} 0 & \text{for } k < 0 \text{ or } k \geq n \\ A \otimes H_k M_*^e & \text{for } 0 \leq k < n - 1 \\ A \otimes Z_{n-1} M_* & \text{for } k = n - 1 \end{cases}$$

In the general case one gets for $A \in \mathbf{Ab}$ the following natural isomorphisms and short exact sequences respectively.

$$H_{n+k}(M(A, n); M) = \begin{cases} 0 & \text{for } k < 0, k > n + 1 \\ A \otimes \text{cok}P & \text{for } k = 0 \\ A *' Z_{n-1} M_* & \text{for } k = n \\ A *'' Z_{n-1} M_* & \text{for } k = n + 1 \end{cases}$$

$$0 \rightarrow A \otimes Z_{n-1} M_* \rightarrow H_{2n-1}(M(A, n); M) \rightarrow A * H_{n-2} M_*^e \rightarrow 0$$

$$0 \rightarrow A \otimes H_r M_*^e \rightarrow H_{n+r}(M(A, n); M) \rightarrow A * H_{r-1} M_*^e \rightarrow 0$$

for $0 < r < n - 1$.

In (8.4) we shall see that the short exact sequence for $H_{n+r}(M(A, n); M)$, $0 < r < n - 1$, is actually naturally split.

(7.11) Remark. We now determine the spherical groups of square homology; see (3.8). Let M be a square group. For each sphere S^n we obtain by (3.1) the square group ($n \geq 1, k \in \mathbb{Z}$)

$$H_{n+k}\{S^n; M\} = (H_{n+k}(S^n; M) \xrightarrow{H} H_{n+k}(S^n|S^n; M) \xrightarrow{P} H_{n+k}(S^n; M))$$

For $k \geq 1$ this is a quadratic \mathbb{Z} -module as in (3.9). Using (5.11) we see

$$H_{1+k}\{S^1; M\} = \begin{cases} M & k = 0 \\ 0 & \text{otherwise} \end{cases}$$

Moreover by (7.10) we get for $n > 1$

$$H_{n+k}\{S^n; M\} = \begin{cases} 0 & \text{for } k < 0 \text{ or } n > 1 \\ H_k M_*^e & \text{for } 0 \leq k < n - 1 \\ Z_{n-1} M_* & \text{for } k = n - 1 \end{cases}$$

Hence only $H_{2n-1}\{S^n; M\}$ is quadratic and $H_{n+k}\{S^n; M\}$ is an abelian group for $k \neq n - 1$.

Proof of (7.10). For $k = 0$ compare (6.11). For $k > 0$ we have by the exact sequence (7.7) the formula, $X = M(A, n)$,

$$H_{n+k}(X; M) = H_{n+k}(X, \hat{M})$$

If A is free abelian the right hand side is computed in [11] by

$$H_{n+k}(X, \hat{M}) = (\bar{H}_* \otimes \hat{M})_{n+k-1}$$

Here $\bar{H}_* = s^{-1}H_*(X, \mathbb{Z})$ is concentrated in degree $n - 1$ so that the result in [11] yields the formulas of the proposition if A is free abelian. If A is not free abelian we use the exact sequence in (3.12).

q.e.d.

(7.12) *Remark.* If M is a quadratic \mathbb{Z} -module we can use (7.10) for the computation of square homology with coefficients in M . For this let $X \in \mathbf{space}_1$ and let

$$(1) \quad X_n = M(A_n, n) \quad \text{with} \quad A_n = H_n(X, \mathbb{Z})$$

be the Moore space of the $n - th$ homology of X . Then there exists a homotopy equivalence of chain complexes

$$(2) \quad C_*(X) \simeq C_*(X_1 \vee X_2 \vee \dots)$$

Hence we get by (7.4) an isomorphism

$$(3) \quad \begin{aligned} H_n(X; M) &= H_n(X_1 \vee X_2 \vee \dots; M) \\ &= \bigoplus_{i \geq 1} H_n(X_i; M) \oplus \bigoplus_{i < j} H_{n+1}(X_i \wedge X_j; M_{ee}) \end{aligned}$$

In the second row we use the cross effect formula. Since we know $H_n(X_i)$ by (7.10) we thus get a description for $H_n(X; M)$ as an abelian group. For this we point out that for $A, B, M_{ee} \in \mathbf{Ab}$

$$(4) \quad H_n(M(A, i) \wedge M(B, j); M_{ee}) = \begin{cases} 0 & \text{for } n < i + j \quad \text{and} \quad n > i + j + 2 \\ A \otimes B \otimes M_{ee} & \text{for } n = i + j \\ \text{Trp}(A, B, M_{ee}) & \text{for } n = i + j + 1 \\ A * B * M_{ee} & \text{for } n = i + j + 2 \end{cases}$$

Here Trp is the triple torsion product of Mac Lane; see Notes on page 393 in [27]. Using (3) for \hat{M} in (7.7) we thus obtain by the exact sequence in (7.7) a possibility to compute $H_n(X, M)$ for an arbitrary square group M in terms of the boundary $\partial : H_n(X, M^{add}) \rightarrow H_{n-1}(X, \hat{M})$. For example we obtain the following vanishing theorem.

(7.13) Proposition. *Let X be a connected CW-space with $\dim X = N$ and let M be a square group. Then*

$$H_n(X, M) = 0 \quad \text{for} \quad n \geq 2N$$

and

$$H_n(X, M) = H_n(X, \hat{M}) \quad \text{for} \quad N < n < 2N$$

§ 8 STABLE SQUARE-HOMOLOGY AND STEENROD SQUARES

Given a square group M we obtain the square-homology $H_n(X; M)$ and the stable square-homology $H_n^S(X; M)$ which is the stabilization of $H_n(X; M)$ as defined in (2.2). We have seen in (6.8) that the stable square-homology $H_n^S(X; M)$ is a homology theory on \mathbf{space}_1 . Hence by (2.8) there is a spectrum E_M associated to the square group M with

$$(8.1) \quad H_n^S(X; M) = H_n(X; E_M).$$

This equation also holds if X is not a finite CW-space since both sides of the equation satisfy the colimit axiom.

(8.2) *Remark.* The correspondence $M \mapsto E_M$ yields a functor

$$E : \mathbf{Square} \rightarrow Ho(\mathbf{spectra})$$

where the right hand side is the homotopy category of spectra. To obtain this functor we use Goodwillie's equivalence [17] between linear homotopy functors and spectra. Hence the functor E is obtained by the functor which carries M to M_{\sharp}^{lin} where the linear homotopy functor M_{\sharp}^{lin} determines functorially E_M by

$$M_{\sharp}^{lin}(X) = \Omega^{\infty}(E_M \wedge X).$$

Compare (4.3) and (4.11).

For an abelian group A let $K(A)$ be the *Eilenberg-MacLane spectrum* with $K(A)_n = K(A, n)$. Let $K(A)[i]$ be the shifted spectrum with $K(A)[i]_n = K(A, n + i)$ where $i \in \mathbb{Z}$. We clearly have

$$(8.3) \quad H_n(X; K(A)[i]) = \tilde{H}_{n-i}(X; A)$$

where the right hand side is the reduced singular homology of X with coefficients in A .

(8.4) Theorem. *Let M be a quadratic \mathbb{Z} -module. Then the stable square homology with coefficients in M is given by the natural isomorphism, $X \in \mathbf{space}_1$,*

$$H_n^S(X, M) = \bigoplus_{i \geq 0} \tilde{H}_{n-i}(X, H_i M_{\star}^e)$$

where M_*^e is the chain complex in (7.8). Hence by (8.3) the spectrum E_M associated to M is the product of shifted Eilenberg-MacLane spectra $K(H_i M_*^e)[i]$ with $i \geq 0$.

Proof. We know by (4.11) that there exists a spectrum E with

$$M_{\sharp}^{\text{lin}}(X) = \Omega^{\infty}(E \wedge X)$$

Hence E is determined by $M_{\sharp}^{\text{lin}}(S^n)$ where we can use a large dimension n . Now in the stable range $M_{\sharp}(S^n) \rightarrow M_{\sharp}^{\text{lin}}(S^n)$ is an equivalence. Here $M_{\sharp}(S^n)$ is given by an abelian simplicial group since M is a quadratic \mathbb{Z} -module. An abelian simplicial group, however, is equivalent to a product of Eilenberg-MacLane spaces. Hence all k -invariants of $M_{\sharp}^{\text{lin}}(S^n)$ vanish and therefore E is a product of Eilenberg-MacLane spectra.

q.e.d.

(8.5) Theorem. *Let M be a square group and let $M^{add} = \text{cok } P$ be defined by M . Then the spectrum E_M associated to M is obtained as the cofiber of a map between spectra*

$$Sq_M : K(M^{add})[-1] \rightarrow \prod_{i \geq 1} K(H_i M_*^e)[i]$$

Proof. For the proof we apply the Goodwillie calculus of functors which yields the fibre sequence in (4.11) (2). Applying this fiber sequence to M_{\sharp} and \hat{M}_{\sharp} in (7.7) gives us the rows in the following commutative diagram of homotopy functors.

$$\begin{array}{ccccc}
 & \Omega M_{\sharp}^{add} & & = & K \\
 & \downarrow & & & \downarrow \mu \\
 \hat{F} & \longrightarrow & \hat{M}_{\sharp} & \longrightarrow & \hat{M}_{\sharp}^{\text{lin}} \\
 \parallel & & \downarrow & & * \downarrow \\
 F & \longrightarrow & M_{\sharp} & \longrightarrow & M_{\sharp}^{\text{lin}} \\
 & & \downarrow & & \\
 & & M_{\sharp}^{add} & &
 \end{array}$$

Here the column in the middle is obtained by the fiber sequence in (7.7). Now (4.11) (2) and (6.9) show that the fibres F and \hat{F} coincide so that the subdiagram $*$ is a pull back. Hence the fiber K of $\hat{M}_{\sharp}^{\text{lin}} \rightarrow M_{\sharp}^{\text{lin}}$ coincides with

$$K = \Omega M_{\sharp}^{add} = K(M^{add})[-1]_{\sharp}$$

Here we denote by E_{\sharp} the homotopy functor given by a spectrum E with

$$E_{\sharp}(X) = \Omega^{\infty}(E \wedge X).$$

Now μ in the diagram yields by (8.4) the map Sq_M in the theorem since a fiber sequence of spectra is also a cofiber sequence.

q.e.d.

(8.6) *Definition.* The map Sq_M in (8.5) is a (multiple) cohomology operation which we call the *squaring operation* associated to the square group M . For $i \geq 1$ let

$$Sq_M^{i+1} : K(M^{add})[-1] \rightarrow K(H_2 M_*^e)[i]$$

be the coordinate of Sq_M of degree $i+1$. Below we show that the classical *Steenrod squares*

$$Sq^{i+1} : K(\mathbb{Z}/2)[-1] \rightarrow K(\mathbb{Z}/2)[i]$$

yield examples of such squaring operations. Moreover we compute the operation Sq_M^2 for any $M \in \mathbf{Square}$ in the next section.

It is a classical result that any short exact sequence $A \rightarrow B \rightarrow C$ of abelian groups induces a long exact sequence of homology groups:

$$\rightarrow H_n(X, A) \rightarrow H_n(X, B) \rightarrow H_n(X, C) \xrightarrow{\beta} H_{n-1}(X, A) \rightarrow$$

where β is the *Bockstein operator*. In a similar way one has for each short exact sequence $L \rightarrow M \rightarrow N$ of square groups a long exact sequence of square-homology groups:

$$(8.7) \quad \rightarrow H_n(X, L) \rightarrow H_n(X, M) \rightarrow H_n(X, N) \xrightarrow{\beta} H_{n-1}(X, L) \rightarrow$$

This sequence is obtained in the same way as the special case in (7.7) (2). In addition the stabilization of (8.7) yields the long exact sequence of stable square-homology groups:

$$(8.8) \quad \rightarrow H_n^S(X, L) \rightarrow H_n^S(X, M) \rightarrow H_n^S(X, N) \xrightarrow{\beta} H_{n-1}^S(X, L) \rightarrow$$

Clearly the sequences (8.7) and (8.8) are natural with respect to maps between short exact sequences in \mathbf{Square} . They are also natural in $X \in \mathbf{space}_1$. As a special case we obtain for $\hat{M} \rightarrow M \rightarrow M^{add}$ the commutative diagram

$$(8.9) \quad \begin{array}{ccc} H_{n+1}^S(X, M^{add}) & \xrightarrow{\beta} & H_n^S(X, \hat{M}) \\ \parallel & & \parallel \\ \tilde{H}_{n+1}(X, M^{add}) & \xrightarrow{(Sq_M)_*} & \bigoplus_{i \geq 1} \tilde{H}_{n-i}(X, H_i M_*^e) \end{array}$$

Here the bottom arrow is induced by the squaring operation Sq_M in (8.6). For this recall that a map $\phi : E \rightarrow E'$ between spectra induces a map $\phi_* : H_n(X, E) \rightarrow H_n(X, E')$ between homology groups. See (2.8) and (8.3).

It is well known that the Steenrod square Sq^1 is obtained as a Bockstein operator; that is,

$$(8.10) \quad \beta = (Sq^1)_* : H_{n+1}(X, \mathbb{Z}/2) \rightarrow H_n(X, \mathbb{Z}/2)$$

coincides with the Bockstein operator associated to $\mathbb{Z}/2 \hookrightarrow \mathbb{Z}/4 \rightarrow \mathbb{Z}/2$. The next result shows that all Steenrod squares Sq^i , $i \geq 1$, are actually obtained by a Bockstein operator of stable square-homology. For this we use the square group

$$(8.11) \quad \mathbb{Z}_{nil}^{4,2} = (\mathbb{Z}/4 \xrightarrow{H} \mathbb{Z}/2 \xrightarrow{0} \mathbb{Z}/4)$$

with $H\{n\} = \{n(n-1)/2\}$. Here $\{n\} \in \mathbb{Z}/k$ denotes the coset of $n \in \mathbb{Z}$. For \mathbb{Z}_{nil} in (6.3) one has a canonical quotient map $q : \mathbb{Z}_{nil} \rightarrow \mathbb{Z}_{nil}^{4,2}$ given by $\mathbb{Z} \rightarrow \mathbb{Z}/4$ and $\mathbb{Z} \rightarrow \mathbb{Z}/2$. There is a short exact sequence in **Square**

$$(8.12) \quad (\mathbb{Z}/2)^\Gamma \hookrightarrow \mathbb{Z}_{nil}^{4,2} \rightarrow \mathbb{Z}/2$$

given by the diagram

$$\begin{array}{ccccccc} (\mathbb{Z}/2)^\Gamma & = & (\mathbb{Z}/2 & \xrightarrow{1} & \mathbb{Z}/2 & \xrightarrow{0} & \mathbb{Z}/2) \\ & & \downarrow & & \parallel & & \downarrow \\ \mathbb{Z}_{nil}^{4,2} & = & (\mathbb{Z}/4 & \longrightarrow & \mathbb{Z}/2 & \xrightarrow{0} & \mathbb{Z}/4) \\ & & \downarrow & & \downarrow & & \downarrow \\ \mathbb{Z}/2 & = & (\mathbb{Z}/2 & \longrightarrow & 0 & \longrightarrow & \mathbb{Z}/2) \end{array}$$

One readily checks that for $M = (\mathbb{Z}/2)^\Gamma$ one has $H_i M_*^c = \mathbb{Z}/2$ for $i \geq 0$. Hence one gets by (8.4) a Bockstein operator β as in the following theorem.

(8.13) Theorem. *Let β be the Bockstein operator of stable square homology associated to the short exact sequence in (8.12). Then the following diagram commutes:*

$$\begin{array}{ccc} H_{n+1}^S(X, \mathbb{Z}/2) & \xrightarrow{\beta} & H_n^S(X, (\mathbb{Z}/2)^\Gamma) \\ \parallel & & \parallel \\ \tilde{H}_{n+1}(X, \mathbb{Z}/2) & \xrightarrow{Sq} & \bigoplus_{i \geq 0} \tilde{H}_{n-i}(X, \mathbb{Z}/2) \end{array}$$

Here Sq has the coordinates $(\chi Sq^{i+1})_*$ induced by the Steenrod squares Sq^{i+1} for $i \geq 0$ where χ is the anti-automorphism of the Steenrod algebra; see 27.24 [22].

For the proof of this theorem we use the *mod-2 restricted lower central series* $\bar{\Gamma}_n(G)$ which defines the group functor

$$(8.14) \quad \overline{nil}_2 : \mathbf{Gr} \rightarrow \mathbf{Gr} \quad \text{with} \quad \overline{nil}_2(G) = G/\bar{\Gamma}_3(G)$$

This is the restricted version of nil_2 in (5.11). For a free group G we have the following facts (1), (2) and (3).

$$(1) \quad G/\bar{\Gamma}_2(G) = G^{ab} \otimes \mathbb{Z}/2 = V$$

is a $\mathbb{Z}/2$ -vector space and

$$(2) \quad \bar{\Gamma}_2(G)/\bar{\Gamma}_3(G) = \bar{L}_2(V)$$

Here $\bar{L}_2(V) \subset V \otimes V$ is generated by $[v, w] = v \otimes w + w \otimes v$ and $v \otimes v$ for $v, w \in V$. $\bar{L}_2(V)$ is the degree 2 part of the free *mod-2 restricted Lie algebra* $\bar{L}(V)$. Using 7.6 [15] one obtains (1) and (2) above. Hence one obtains the natural short exact sequence in the top row of the following commutative diagram. The bottom row is obtained by the exact sequence in (8.12) and the quadratic tensor product.

$$(3) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \bar{L}_2(G^{ab} \otimes \mathbb{Z}/2) & \longrightarrow & \overline{nil}_2(G) & \longrightarrow & G^{ab} \otimes \mathbb{Z}/2 \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ 0 & \longrightarrow & G \otimes (\mathbb{Z}/2)^\Gamma & \longrightarrow & G \otimes \mathbb{Z}_{nil}^{4,2} & \longrightarrow & G \otimes \mathbb{Z}/2 \longrightarrow 0 \end{array}$$

The vertical isomorphisms which are natural in G are induced by $G \otimes \mathbb{Z}_{nil} = nil_2(G)$ in (6.4); see 8.1 [10].

Proof of (8.19). If $G = \alpha(X)$ is the free simplicial group given by X then the exact sequence of simplicial groups in (8.14) (3) induces the Bockstein operator in (8.13). The connecting homomorphism

$$d_1 : \pi_n(G^{ab} \otimes \mathbb{Z}/2) \rightarrow \pi_{n-1}\bar{L}_2(G^{ab} \otimes \mathbb{Z}/2)$$

is computed in 8.10 [15] in terms of the Steenrod operations Sq^i which act from the right on homology since for a finite type space X we have $H_*(X, \mathbb{Z}/2) = Hom(H^*(X, \mathbb{Z}/2), \mathbb{Z}/2)$. The anti-isomorphism χ of the Steenrod algebra has the property that for $y \in H_*(X, \mathbb{Z}/2)$ one has

$$(\chi Sq^i)_*(y) = y Sq^i$$

Therefore the stabilization of the differential d_1 in 8.10 [15] yields the result. q.e.d.

As an application of (8.13) and (8.5) we obtain the following results which determine explicitly the spectra E_M associated to $M = \mathbb{Z}_{nil}$ and $M = \mathbb{Z}_{nil}^{4,2}$ respectively. For this we consider the following commutative diagram in **Square**

$$(8.15) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}^\Lambda & \longrightarrow & \mathbb{Z}_{nil} & \longrightarrow & \mathbb{Z} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & (\mathbb{Z}/2)^\Lambda & \longrightarrow & \mathbb{Z}_{nil}^{4,2} & \longrightarrow & \mathbb{Z}/4 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & (\mathbb{Z}/2)^\Gamma & \longrightarrow & \mathbb{Z}_{nil}^{4,2} & \longrightarrow & \mathbb{Z}/2 \longrightarrow 0 \end{array}$$

Here the rows are exact and the vertical arrows are quotient maps. The top row and the row in the middle are the exact sequences $\hat{M} \rightarrow M \rightarrow M^{add}$ for $M = \mathbb{Z}_{nil}$ and $M = \mathbb{Z}_{nil}^{4,2}$ respectively. We have $\mathbb{Z}^\Lambda = (0 \rightarrow \mathbb{Z} \rightarrow 0)$ and $(\mathbb{Z}/2)^\Lambda = (0 \rightarrow \mathbb{Z}/2 \rightarrow 0)$. Naturality of the Bockstein operator yields by (8.9) a relation of Sq_M for $M = \mathbb{Z}_{nil}$ and $M = \mathbb{Z}_{nil}^{4,2}$ with β in (8.13). Since the maps $\mathbb{Z}^\Lambda \rightarrow (\mathbb{Z}/2)^\Lambda \rightarrow (\mathbb{Z}/2)^\Gamma$ induce injections

$$H_*(\mathbb{Z}^\Lambda)_*^e \rightarrow H_*(\mathbb{Z}/2^\Lambda)_*^e \rightarrow H_*(\mathbb{Z}/2^\Gamma)_*^e$$

(see (7.8)) we get the following results.

(8.16) Theorem. For the square group $M = \mathbb{Z}_{nil}$ we have the spectrum $E = E_M$ with

$$H_n^S(X, nil_2) = H_n^S(X, \mathbb{Z}_{nil}) = H_n(X, E)$$

This spectrum E is the cofiber of the map

$$Sq_M : K(\mathbb{Z})[-1] \rightarrow \prod_{i \text{ odd}} K(\mathbb{Z}/2)[i]$$

Here the coordinate of degree $i+1$ is $q^*(\chi Sq^{i+1})$ where $q : \mathbb{Z} \rightarrow \mathbb{Z}/2$ is the quotient map and χ is the anti-automorphism of the Steenrod algebra.

(8.17) Theorem. For the square group $M = \mathbb{Z}_{nil}^{4,2}$ we have the spectrum $E = E_M$ with

$$H_m^S(X, \overline{nil}_2) = H_m^S(X, \mathbb{Z}_{nil}^{4,2}) = H_m(X, E).$$

This spectrum E is the cofiber of the map

$$Sq_M : K(\mathbb{Z}/4)[-1] \rightarrow \prod_{i \geq 1} K(\mathbb{Z}/2)[i]$$

Here the coordinate of degree $i+1$ is $q^*(\chi Sq^{i+1})$ where $q : \mathbb{Z}/4 \rightarrow \mathbb{Z}/2$ is the quotient map and χ is the anti-automorphism of the Steenrod algebra.

§ 9 THE SQUARING OPERATION Sq_M^2

The squaring operation Sq_M^2 associated to a square group M is an element in the following group where n is large; see (8.6).

$$\begin{aligned} & [K(M^{add})[-1], K(H_1 M_*^e)[1]] = \\ & = [K(M^{add}, n-1), K(H_1 M_*^e, n+1)] \\ (9.1) \quad & = Hom(M^{add} \otimes \mathbb{Z}/2, H_1 M_*^e) \end{aligned}$$

Here we have $M^{add} = \text{cok}(P)$ and $H_1 M_*^e = \text{ker}(P)/\text{im}(2 - HP)$ so that Sq_M^2 is a homomorphism

$$Sq_M^2 : \text{cok}(P) \otimes \mathbb{Z}/2 \rightarrow \text{ker}(P)/\text{im}(2 - HP)$$

We now consider the following diagram where $(\begin{smallmatrix} | \\ | \end{smallmatrix})_H$ is the cross effect of H in (6.3) and where Δ is the diagonal with $\Delta(x) = (x, x)$ for $x \in M_e$.

$$(9.2) \quad \begin{array}{ccccc} M_e & \xrightarrow{q' \otimes 1} & \text{cok}(P) \otimes \mathbb{Z}/2 & \xrightarrow{Sq_M^2} & \ker(P)/\text{im}(2 - HP) \\ \Delta \downarrow & & & & \uparrow q \\ M_e \times M_e & \xrightarrow{(\begin{smallmatrix} | \\ | \end{smallmatrix})_H} & M_{ee} & \supset & \ker(P) \end{array}$$

Here q is the quotient map and $q' : M_e \rightarrow \text{cok}(P)$ is the quotient map.

(9.3) Theorem. For a square group M there is a unique homomorphism Sq_M^2 such that diagram (9.2) commutes and this homomorphism coincides with the squaring operation Sq_M^2 by use of (9.1).

For example for $M = \mathbb{Z}_{\text{nil}}$ the homomorphism Sq_M^2 is the isomorphism $Sq_M^2 : \mathbb{Z} \otimes \mathbb{Z}/2 \cong \mathbb{Z}/2$. For the proof of (9.3) we have to recall algebraic models of $(n-1)$ -connected $(n+1)$ -types, $n \geq 2$, obtained in [6].

(9.4) Definition. A reduced quadratic module (ω, δ) is a diagram

$$M^{ab} \otimes M^{ab} \xrightarrow{\omega} L \xrightarrow{\delta} M$$

of homomorphisms between groups such that the following properties hold. The groups M, L have nilpotency degree 2 and the quotient map $M \rightarrow M^{ab}$ is denoted by $x \mapsto \{x\}$. Then for $x, y \in L$, $a, b \in M$ we have

$$\begin{aligned} \delta\omega(\{a\} \otimes \{b\}) &= -a - b + a + b \\ \omega(\{\delta x\} \otimes \{\delta y\}) &= -x - y + x + y \\ \omega(\{\delta x\} \otimes \{a\} + \{a\} \otimes \{\delta x\}) &= 0 \end{aligned}$$

We say that (ω, δ) is a *stable quadratic module* if in addition

$$\omega(\{a\} \otimes \{b\} + \{a\} \otimes \{b\}) = 0$$

A morphism $(\omega, \delta) \rightarrow (\omega', \delta')$ is a commutative diagram in \mathbf{Gr}

$$\begin{array}{ccc} L & \xrightarrow{l} & L' \\ \delta \downarrow & & \downarrow \delta' \\ M & \xrightarrow{m} & M' \end{array}$$

such that $l\omega = \omega'(m^{ab} \otimes m^{ab})$. This is a weak equivalence if (l, m) induces isomorphisms

$$\ker(\delta) \cong \ker(\delta'), \text{ cok}(\delta) \cong \text{cok}(\delta')$$

Let **rquad** (resp. **squad**) be the categories of reduced (resp. stable) quadratic modules and let *Horquad*, *Hosquad* be the localization with respect to weak equivalences. There are equivalences of categories for which the following diagram commutes, $n \geq 3$; see [6] and [8].

$$(9.5) \quad \begin{array}{ccc} \mathbf{types}_2^1 & \xrightarrow[\sim]{\lambda_2} & \mathbf{Horquad} \\ \Omega^{n-2} \cup & & \cup i \\ \mathbf{types}_n^1 & \xrightarrow[\sim]{\lambda_n} & \mathbf{Hosquad} \end{array}$$

Here \mathbf{types}_k^1 is the homotopy category of CW-spaces X for which $\pi_i X = 0$ for $i < k$ and $i > k + 1$. Moreover Ω^{n-2} is the iterated loop space functor and i is the inclusion functor. For $X \in \mathbf{types}_n^1$ and $(\omega, \delta) = \lambda_n(X)$ we have natural isomorphisms

$$\pi_n X = \pi_n = \text{cok}(\delta), \quad \pi_{n+1}(X) = \pi_{n+1} = \text{ker}(\delta)$$

Moreover the k -invariant of X is an element

$$(1) \quad k(X) \in H^{n+2}(K(\pi_n, n), \pi_{n+1}) = \text{Hom}(\Gamma_n^1(\pi_n), \pi_{n+1})$$

Here Γ_n^1 is the functor $\mathbf{Ab} \rightarrow \mathbf{Ab}$ which is Whitehead quadratic functor Γ for $n = 2$ and $\otimes \mathbb{Z}/2$ for $n \geq 3$. We can obtain $k(X)$ from $(\omega, \delta) = \lambda_n(X)$ as follows. Given $(\omega, \delta) \in \mathbf{rquad}$ there is a unique homomorphism k for which the following diagram commutes

$$(2) \quad \begin{array}{ccccc} \Gamma(M^{ab}) & \xrightarrow{q_*} & \Gamma(\text{cok } \delta) & \xrightarrow{k} & \text{ker } \delta \\ H \downarrow & & & & \cap \\ M^{ab} \otimes M^{ab} & \xrightarrow{\omega} & L & \xrightarrow{\delta} & M \end{array}$$

Here H is the cross effect map in (3.1) for the functor Γ and q_* is induced by the quotient map $M \rightarrow M^{ab} \rightarrow \text{cok } \delta$ which factors through M^{ab} . If (ω, δ) is stable then k admits a factorization

$$(3) \quad k : \Gamma(\text{cok } \delta) \xrightarrow{\sigma} \text{cok}(\delta) \otimes \mathbb{Z}/2 \xrightarrow{k'} \text{ker } \delta$$

Now the k -invariant $k(X)$ with $(\omega, \delta) = \lambda_n X$ is k for $n = 2$ and is k' for $n \geq 3$. Compare [6, 8].

Recall that **Square** is the category of square groups in (6.3). We now define canonical algebraic functors

$$(9.6) \quad \begin{array}{ccccc} \text{Square} & \xrightarrow{\tau} & \mathbf{rquad} & \xrightarrow{L} & \mathbf{Horquad} \\ & & \cup & & \cup \\ \text{Square} & \xrightarrow{\tau^{lin}} & \mathbf{squad} & \longrightarrow & \mathbf{Hosquad} \end{array}$$

together with a natural transformation $a : \tau \rightarrow \tau^{lin}$. Here L is the localization functor. The functor τ is given by

$$\tau(M) = (M_e^{ab} \otimes M_e^{ab} \xrightarrow{\omega} M_{ee} \xrightarrow{\delta} M_e)$$

where $\omega(\{x\} \otimes \{y\}) = (x|y)_H$ and $\delta = P$. Moreover let

$$\tau^{lin}(M) = (M_e^{ab} \otimes M_e^{ab} \xrightarrow{\omega'} M_{ee}/\text{im}(2 - HP) \xrightarrow{\delta'} M_e)$$

where ω' , resp. δ' , are induced by ω , resp. δ , in $\tau(M)$ above. We clearly have the natural map $a : \tau(M) \rightarrow \tau^{lin}(M)$ which is the identity on M_e and the quotient map on M_{ee} . One readily checks that the functors τ and τ^{lin} are well defined; for this we only prove:

(9.7) Lemma. $\tau^{lin}(M)$ is a stable quadratic module.

Proof. We have to show that for $a, b \in M_e$ we have $(a|b)_H + (b|a)_H \in \text{im}(2 - HP)$. But we know by 3.5 (4) in [10] that $\Delta : M_e \rightarrow M_{ee}$ with

$$\Delta(a) = (HP - 2)Ha + (a|a)_H$$

is a homomorphism. Hence we get

$$\begin{aligned} (a|b)_H + (b|a)_H &= (a + b|a + b)_H - (a|a)_H - (b|b)_H \\ &= \Delta(a + b) + (2 - HP)H(a + b) \\ &\quad - (\Delta(a) + (2 - HP)H(a) + \Delta(b) + (2 - HP)H(b)) \\ &= (2 - HP)(a|b)_H \end{aligned}$$

This term needs not to be trivial in M_{ee} as the example \mathbb{Z}_{nil} shows. Hence $\tau(M)$ in general is not stable. q.e.d.

We now obtain functors

$$(9.8) \quad S, S^{lin} : \mathbf{Square} \rightarrow \mathbf{types}_2^1$$

which carry the square group M to $S(M) = M_{\sharp}(S^2)$ and $S^{lin}(M) = 3\text{-type of } M_{\sharp}^{lin}(S^2)$ respectively. Here (7.10) shows that $M_{\sharp}(S^2) \in \mathbf{types}_2^1$. We have the obvious map

$$b : S(M) = M_{\sharp}(S^2) \rightarrow M_{\sharp}^{lin}(S^2) \rightarrow S^{lin}(M)$$

which is natural in M . We also observe that $S^{lin}(M)$ is an infinite loop space since $M_{\sharp}^{lin}(S^2)$ is an infinite loop space.

(9.9) Theorem. For the functors in (9.8), (9.6) and (9.5) there exist natural isomorphisms such that the following diagram commutes.

$$\begin{array}{ccc} \lambda_2 S(M) & \cong & L\tau(M) \\ b_* \downarrow & & \downarrow a_* \\ \lambda_2 S^{lin}(M) & \cong & L\tau^{lin}(M) \end{array}$$

Proof of (9.9). One readily checks that Sq_M^2 in (9.2) is the k -invariant of $\tau^{lin}(M)$. The k -invariant of $\tau^{lin}(M)$ coincides with the first non-trivial k -invariant of $M_{\natural}^{lin}(S^2)$ by (9.9). This yields the proposition in (9.3).

q.e.d.

Proof of (9.9). We have by (7.10) that

$$\begin{aligned} \pi_2(S(M)) &= \pi_2 M^{\sharp}(S^2) = \mathbb{Z} \otimes \text{cok}(P) = \text{cok}(P) \\ \pi_3(S(M)) &= \pi_3 M^{\sharp}(S^2) = \mathbb{Z} \otimes Z_1 M_* = \ker(P) \end{aligned}$$

Hence the isomorphism $\lambda_2 S(M) \cong L\tau(M)$ exists on the level of homotopy groups. An explicit isomorphism is given by the result of Conduché [14]. For this let G be the free simplicial group generated by one element in degree 1. Then we have $G \simeq \alpha(S^2)$ in (5.1) so that $M^{\sharp}(S^2) \simeq B|G \otimes M|$. Here $G \otimes M$ is a simplicial group with only two nontrivial homotopy groups π_1 and π_2 . Hence the homotopy type of $G \otimes M$ is described by its reduced 2-module in the sense of Conduché (see (2.9) [14]). But this reduced 2-module of $G \otimes M$ coincides with $\tau(M)$. This yields the isomorphism $\lambda_2 S(M) \cong L\tau(M)$. Moreover we know that a_* coincides with b_* on the level of homotopy groups since

$$\begin{array}{ccc} \pi_3(S(M)) & = & \ker(P) \\ b_* \downarrow & & \downarrow a \\ \pi_3(S^{lin}(M)) & = & \ker(P)/\text{in}(2 - HP) \end{array}$$

commutes. This shows that the k -invariant of $S^{lin}(M)$ is actually the k -invariant of $\tau^{lin}(M)$. Moreover the natural isomorphism $\lambda_2 S(M) \cong L\tau(M)$ induces the natural isomorphism $\lambda_2 S^{lin}(M) \cong L\tau^{lin}(M)$. q.e.d.

APPENDIX A: A CRITERION FOR 2-EXCISIVE FUNCTORS

We describe a criterion for 2-exciseive functors which shows that only very special strongly co-Cartesian diagrams are needed to determine a 2-exciseive functors. We use this result for the proof of theorem (6.6). It would be interesting to obtain a similar result for n -exciseive functors.

Let \underline{Y} be a 2-cube consisting of spaces Y_1, Y_2, Y_3, Y_4 and let $A \in \mathbf{space}$. Then we define $A \vee \underline{Y}$ by

$$\begin{array}{ccc}
A \vee Y_1 & \longrightarrow & A \vee Y_3 \\
\downarrow & & \downarrow \\
A \vee Y_2 & \longrightarrow & A \vee Y_4
\end{array}$$

This yields the 3-cube $(0,1) : A \vee \underline{Y} \rightarrow \underline{Y}$. Clearly if \underline{Y} is co-Cartesian then $(0,1) : A \vee \underline{Y} \rightarrow \underline{Y}$ is a strongly co-Cartesian 3-cube. Recall the definition of $\text{Cube}(X, A)$ in the proof of (4.12).

(A.1) Theorem. *Let $D : \mathbf{space}_r \rightarrow \mathbf{space}_1, r \geq 0$, be a homotopy functor. Assume that for pairs $(X, A) \in \mathbf{pair}_r$ and all homotopy push outs \underline{Y} in \mathbf{space}_r the 3-cubes $D(\text{Cube}(X, A))$ and $D((0,1) : A \vee \underline{Y} \rightarrow \underline{Y})$ are Cartesian. Then the homotopy functor D is 2-exciseive.*

The theorem shows that essentially the 3-cubes $\text{Cube}(X, A)$ suffice to determine quadratic homotopy functors. This again shows that the quadratic excision sequence essentially covers all excision properties of a 2-exciseive functor.

Proof of (6.6). Let F be a quadratic coefficient functor so that $F = \otimes M$ by (6.4). Since we can use (5.4) it is enough to prove that F_{\sharp} is 2-exciseive. For this we can use theorem (A.1) above. Using (1) in the proof of (6.9) above we see that $F_{\sharp}((0,1) : A \vee \underline{Y} \rightarrow \underline{Y})$ is Cartesian if \underline{Y} is co-Cartesian. Moreover using the pull back diagram (4) in the proof of (6.9) we see that $F_{\sharp}(\text{Cube}(X', A'))$ is Cartesian for $(X', A') \in \mathbf{pair}_1$. For this we use (4.8). Hence the assumptions of (A.1) are satisfied and therefore F_{\sharp} is 2-exciseive.

q.e.d

Proof of (A.1). Let \underline{Y} be a cofibrant co-Cartesian 2-cube and let $i : \underline{Y} \subset A \vee \underline{Y}$ be the inclusion then $D(i)$ is Cartesian since $D(0,1)$ in (6.8) is Cartesian and $(0,1)i = 1$. For this we use the lemmas in section 1 of [12]. Next let (Y_1, A) be a pair in \mathbf{pair}_r . Then we obtain the 3-cube $q : \underline{Y} \rightarrow \underline{Y}/A$ given by the quotient maps $q_i : Y_i \rightarrow Y_i/A$. Clearly q is strongly co-Cartesian. For this we put the cubes $\text{Cube}(Y_i, A)$ and the cube $(0,1) : A \vee \underline{Y} \rightarrow \underline{Y}$ together to form a large cube with boundary $q : \underline{Y} \rightarrow \underline{Y}/A$. We apply D to this large cube and then we take homotopy fibers which form the following commutative diagram where Q_i is the homotopy fiber of $(q_i)_* : D(Y_i) \rightarrow D(Y_i/A)$.

$$(1) \quad \begin{array}{ccc}
Q_1 & \longrightarrow & Q_2 \\
\downarrow \swarrow & & \searrow \downarrow \\
D(A \vee Y_1)_2 & \longrightarrow & D(A \vee Y_2)_2 \\
\downarrow & & \downarrow \\
D(A \vee Y_3)_2 & \longrightarrow & D(A \vee Y_4)_2 \\
\downarrow \swarrow & & \searrow \downarrow \\
Q_3 & \longrightarrow & Q_4
\end{array}$$

Here the inside square is Cartesian since $D(0, 1)$ is Cartesian by the assumption in (A.1). Moreover, since $D(\text{Cube}(Y_i, A))$ is Cartesian by the assumption in (A.1) we see that the four boundary subdiagrams are Cartesian by (4.8). For this we use the following diagram; compare the proof of (4.9).

$$(2) \quad \begin{array}{ccccc} K(A) & \longrightarrow & K' & \longrightarrow & K'' \\ \downarrow & & \downarrow & & \downarrow \\ D(A \vee A)_2 & \longrightarrow & D(A \vee Y_1)_2 & \longrightarrow & D(A \vee Y_2)_2 \\ \downarrow & & \downarrow & * & \downarrow \\ D(A) & \longrightarrow & Q_1 & \longrightarrow & Q_2 \end{array}$$

Here the columns are fiber sequences. We know that $K(A) \rightarrow K'$ and $K(A) \rightarrow K''$ are homotopy equivalences. Hence also $K' \rightarrow K''$ is a homotopy equivalence and therefore the subdiagram $*$ is Cartesian. Now all subdiagrams of (1) being Cartesian we see that also the boundary diagram of (1) is Cartesian. This completes by (4.8) the proof that $D(q : \underline{Y} \rightarrow Y/A)$ is Cartesian. Now let $\underline{Y} \cup CA$ be the 2-cube given by $Y_i \cup_A CA$. Since the quotient map $Y_i \cup CA \rightarrow Y_i/A$ is a homotopy equivalence we see that also the inclusion $j : \underline{Y} \subset \underline{Y} \cup CA$ is a 3-cube for which $D(j)$ is Cartesian. Now let \underline{X} and \underline{Y} be 2-cubes and let $f : \underline{Y} \rightarrow \underline{X}$ be a strongly co-Cartesian 3-cube. We may assume up to equivalence that f is cofibrant, that is all maps in the 3-cube f are cofibrations and all subsquares are actually push outs. Using CW-decomposition we can filter f by an infinite sequence

$$f : \underline{Y} = \underline{X}_0 \subset \underline{X}_1 \subset \underline{X}_2 \subset \dots \subset \text{colim}(\underline{X}_i) \simeq \underline{X}$$

Here all $\underline{X}_i \subset \underline{X}_{i+1}$ are cofibrant co-Cartesian 3-cubes for which there exist $A_i \in \mathbf{space}_r$ such that $\underline{X}_1 = \underline{Y} \vee A_1$ and $\underline{X}_{i+1} = \underline{X}_i \cup CA_i$, $i \geq 1$. We have seen above that $D(\underline{X}_i \rightarrow \underline{X}_{i+1})$ is Cartesian for all $i \geq 0$. This implies by I.4b.5 in [3] that also $D(\underline{Y} \rightarrow \underline{X})$ is Cartesian since we assume that D preserves filtered colimits.

q.e.d.

APPENDIX B: THE HOMOLOGY SPECTRAL SEQUENCE

Let D be a reduced homotopy functor. We describe a spectral sequence which converges to the homology with coefficients in D . If D is linear this is the Atiyah-Hirzebruch spectral sequence.

(B.1) *Definition.* Let $D : \mathbf{space}_1 \rightarrow \mathbf{space}_1$ be a reduced homotopy functor and let X be a connected CW-complex. Let $U = C(X)$ be the reduced cone on X which is filtered by $X = U^0 \subset U^1 \subset \dots \subset U$ with $U^n = X \cup C(X^{n-1})$, $n \geq 1$. For $(Y, B) \in \mathbf{pair}_1$ let

$$(1) \quad H_n(Y, B) = H_n(Y, B; D)$$

be the D -homology in (4.2). Then the filtered space U gives rise to the following spectral sequence. First we can form the long exact sequence

$$(2) \quad \dots \rightarrow H_2(U^n) \xrightarrow{j} H_2(U^n, U^{n-1}) \xrightarrow{\partial} H_1(U^{n-1}) \xrightarrow{i} H_1(U^n) \xrightarrow{j} H_1(U^n, U^{n-1}) \rightarrow 0$$

where the last object is a set with basepoint 0 and all the other objects are groups. We can form the r -th derived homotopy sequences of (2) for $r \geq 0$, $n \leq 0$,

$$(3) \quad \dots \rightarrow H_2^{(r)}(n-2r-1) \rightarrow F_1^{(r)}(n-r) \rightarrow H_1^{(r)}(n-r) \rightarrow H_1^{(r)}(n-r-1) \rightarrow F_0^{(r)}(n) \rightarrow 0$$

Here we set for $q \in \mathbb{Z}$, $n = -m$,

$$H_q^{(r)}(n) = \text{image}(H_q(U^{m-r}) \rightarrow H_q(U^m))$$

$$F_q^{(r)}(n) = \text{kernel}(H_{q+1}(U^{m+1}, U^m) \rightarrow H_q(U^m)/H_q^{(r)}(n)) / \text{action of kernel}(H_{q+1}(U^{m+1}) \rightarrow H_{q+1}(U^{m+r+1}))$$

For $r = 0$ the sequence (3) coincides with (2). One can check that (3) is well defined and has the same properties as (2). Let

$$(4) \quad E_r^{s,t} = F_{t-s}^{(r-1)}(s) \quad \text{for} \quad t \in \mathbb{Z}, s \leq 0, r \geq 1,$$

and let the differential $d_r : E_r^{s,t} \rightarrow E_r^{s+r, t+r-1}$ of degree $(r, r-1)$ be the composition

$$F_{(t-s)}^{(r-1)}(s) \rightarrow H_{t-s}^{(r-1)}(s) \rightarrow F_{t-s-1}^{(r-1)}(s+r)$$

where we use the operators from (3). Clearly we have for $q \geq 0$, $s \geq 0$

$$(5) \quad E_1^{-s, q-s} = F_q^{(0)}(-s) = H_{q+1}(U^{s+1}, U^s)$$

and d_1 is the composition

$$(6) \quad d_1 = j\partial : H_{q+1}(U^{s+1}, U^s) \rightarrow H_q(U^s) \rightarrow H_q(U^s, U^{s-1})$$

Assume that the pairs (U^m, U^{m-1}) have the property that there exist $0 \leq N_0 \leq N_1 \leq \dots$ with $\lim \{N_m\} = \infty$ such that

$$(7) \quad H_i(U^{m+1}, U^m) = 0 \quad \text{for} \quad i < N_m.$$

Then we can find for each $q \geq 0$ a bound $r = r(q) < \infty$ such that

$$(8) \quad E_r^{s, q+s} = E_{r+1}^{s, q+1} = \dots = E_\infty^{s, q+s}$$

We obtain a filtration of $H_q(X)$ for $q \geq 1$, $s \geq 0$,

$$(9) \quad \begin{aligned} 0 &= K_{0,q} \subset K_{1,q} \subset \dots \subset K_{s,q} \subset \dots \subset H_q(X) \\ K_{s,q} &= \text{kernel}(H_q(X) \rightarrow H_q(X/X^s)) \end{aligned}$$

Since D is reduced we see that $\cup_{s \geq 0} K_{s,q} = H_q(X)$ and

$$(10) \quad E_{\infty}^{-s, q-s} = K_{s,q} / K_{s+1,q}.$$

A similar spectral sequence is available for reduced homotopy functors which map to **spectra**; compare (4.1).

(B.2) *Remark.* The spectral sequence $E_r^{s,t}$ above coincides with the spectral sequence in (III.10.4) [3] by considering the relative homotopy groups of the filtered space $\{D(U^n)\}$. The conventions for indexing $E_r^{s,t}$ arises from the comparison with the Bousfield-Kan spectral sequence [12]; compare the discussion in (III.10.2) [3]. We can alter the indexing by defining

$$E_{s,t}^r = E_r^{-s,t}, \quad s \geq 0, \quad t \in \mathbb{Z}.$$

Then the differential for $E_{s,t}^r$ has degree $(-r, r-1)$ as in the spectral sequence discussed by G.W. Whithead, chapter XIII page 614 [29]. Now assume that D in (B.1) is a linear homotopy functor. Then $H_n(Y, B; D) = H_n(Y, B)$ is a homology theory on pair_1 in the sense of (2.2) and we get for $s \geq 0$

$$\begin{aligned} E_{s, q-s}^1 &= E_1^{-s, q-s} = H_{q+1}(U^{s+1}, U^s) \\ &= H_{q+1}(X \cup CX^s, X \cup CX^{s-1}) \\ &= H_{q+1}(\Sigma X^s, \Sigma X^{s-1}) = H_q(X^s, X^{s-1}) \end{aligned}$$

Moreover by the exact sequence

$$H_q(X^s) \rightarrow H_q(X) \rightarrow H_q(X/X^s)$$

We see that $K_{-s,q}$ in (B.1) (9) coincides with $I_{s, q-s}$ where $I_{s, q-s}$ is the image of $H_q(X^s) \rightarrow H_q(X)$ as on page 613 [29]. This shows that the spectral sequence (B.1) yields as a special case the Atiyah-Hirzebruch spectral sequence in XIII.3.3 [29].

APPENDIX C: A SPECTRAL SEQUENCE FOR SQUARE-HOMOLOGY

In this section we apply the spectral sequence of Appendix B to square homology. Let X be a CW-complex with trivial 0-skeleton $X^0 = *$ and let M be a square group. We consider the filtration, $s \geq 0$,

$$0 = K_{0,q} \subset K_{1,q} \subset \dots \subset K_{s,q} \subset \dots \subset H_q(X; M)$$

of square homology given by

$$(C.1) \quad K_{s,q} = \text{kernel}(H_q(X; M) \rightarrow H_q(X/X^s))$$

where $X \rightarrow X/X^s$ is the quotient map. Then the spectral sequence $(E_r^{-s,q-s}, d_r)$ defined in the Appendix converges and satisfies

$$(C.2) \quad E_{\infty}^{-s,q-s} = K_{s,q}/K_{s-1,q}$$

We now determine the E_1 -term of this spectral sequence. Let

$$U^s = X \cup CX^{s-1} \simeq X/X^{s-1}$$

and let $A^s = X^s/X^{s-1}$. Then we have

$$(C.3) \quad U^{s+1} = X \cup CX^{s+1} = U^s \cup CA^s$$

where the attaching map $A^s \rightarrow U^s$ is the composite $f : A^s \subset X/X^{s-1} \simeq U^s$ given by the inclusion. Clearly $A_s = M(C_s, s)$ is the Moore space of the free abelian group $C_s + H_s(X^s, X^{s-1}; \mathbb{Z})$ which is part of the cellular chain complex $C_*X = (C_*, d)$. We can compute the E_1 -term

$$(C.4) \quad E_1^{-s,q-s} = H_{q+1}(U^{s+1}, U^s; M)$$

by applying diagram (3.4) to (C.3). This yields the following commutative diagram in which the row and the column are exact.

$$(C.5) \quad \begin{array}{ccc} H_{q+1}(A_s \wedge A_s; M_{ee}) & & \\ \downarrow (P, -(1, f)_*) & & \\ H_q(A_s; M) \oplus H_{q+1}(A_s \wedge X_s; M_{ee}) & \searrow (f_*, P(f, 1)_*) & \\ \downarrow & \xrightarrow{\partial} & H_q(U^s, M) \\ H_{q+1}(U^{s+1}, U^s; M) & & \\ \downarrow & & \\ H_q(A_s \wedge A_s; M_{ee}) & & \\ \downarrow (P, -(1, f)_*) & & \\ H_{q-1}(A_s; M) \oplus H_q(A_s \wedge X_s; M_{ee}) & & \end{array}$$

Using (7.10) this yields the following result on $H_{q+1}(U^{s+1}, U^s; M)$. Let $B_s = \text{image}(d : C_{s+1} \rightarrow C_s)$ be the group of boundaries in the cellular chain complex C_*X and let $q : C_s \rightarrow C_s/B_s$ be the quotient map.

(C.6) Proposition. The E_1 -term is given by the groups (C.4) which can be described by the following isomorphisms and exact sequences respectively.

$$H_{q+1}(U^{s+1}, U^s; M) = \begin{cases} 0 & \text{for } q < s \\ C_s \otimes H_{q-s} M_*^c & \text{for } s \leq q < 2s - 1 \\ C_s \otimes H_{q-s+1}(X; M_{ee}) & \text{for } q > 2s \end{cases}$$

For $q = 2s - 1$ and $q = 2s$ one has the exact sequence

$$\begin{aligned} 0 \rightarrow C_s \otimes H_{s+1}(X; M_{ee}) \xrightarrow{j} H_{2s+1}(U^{s+1}, U^s; M) \rightarrow C_s \otimes C_s \otimes M_{ee} \\ \xrightarrow{(P, -q^*)} \bar{C}_s \otimes Z_{s-1} M_* \oplus C_s \otimes C_s / B_s \otimes M_{ee} \rightarrow H_{2s}(U^{s+1}, U^s; M) \rightarrow 0 \end{aligned}$$

Here $\bar{C}_s = C_s$ for $s > 1$ and $\bar{C}_1 = \pi_1(X^1)$ is the fundamental group of the 1-skeleton X^1 for $s = 1$ and $Z_0 M_* = M$.

A cross effect argument shows that the inclusion j in the exact sequence of (C.6) is split injective. Moreover the differential d_1 of the spectral sequence is of the form $d \otimes 1$ (with d given by $C_* X$) for the groups in the first part of (C.6) with $q \neq 2s - 1, 2s$. It is interesting to compare (C.6) with the results in section 7. This shows that the spectral sequence for $q > s$ depends only on \hat{M} and hence is determined in this range by $C_* X$.

REFERENCES

1. Arone, G. and Mahowald, M., *The Goodwillie tower of the identity functor and the unstable periodic homotopy of spheres*, Preprint (1995), Northwestern University.
2. Baues, H.J., *Obstruction theory*, LNM 628, Springer Verlag (1977).
3. Baues, H.J., *Algebraic homotopy*, Cambridge University Press (1989).
4. Baues, H.J., *Metastable homotopy and homology*, Quaestiones Mathematicae 14 (1991), 161-178.
5. Baues, H.J., *Quadratic functors and metastable homotopy*, J. pure appl. Algebra 91 (1994), 49-107.
6. Baues, H.J., *Combinatorial Homotopy and 4-dimensional Complexes*, de Gruyter, Berlin (1991).
7. Baues, H.J., *Homotopy type and Homology*, Oxford Mathematical Monographs, 500 pages, Oxford University press, to appear.
8. Baues, H.J. and Conduché, D., *On the 2-type of an iterated loop space*, Preprint Max-Planck-Institut für Mathematik 95-112.
9. Baues, H.J. and Hartl, M. and Pirashvili, T., *Quadratic categories and square rings*, To appear in J. pure appl. Algebra.
10. Baues, H.J. and Pirashvili, T., *Quadratic endofunctors of the category of groups*, Preprint MPI für Math. 95-55.
11. Baues, H.J. and Pirashvili, T., *The universal coefficient theorem for quadratic functors*, Preprint MPI für Math. 95-57.
12. Bousfield, A.K. and Kan, D.M., *Homotopy Limits, Completions and Localizations*, LNM 304 Springer Verlag (1972).
13. Cohen, F.R. and May, J.P. and Taylor, L.R., *Splitting of certain spaces CX* , Math. Proc. Camb. Phil. Soc., (1978), 84, 465-496.
14. Conduché, D., *Modules croisés généralisés de longueur 2*, J. pure and appl. Algebra 34 (1984), 155-178.

15. Curtis, E.B., *Simplicial Homotopy theory*, Advances in Math. 6 (1971), 107-209.
16. Dold, A. and Puppe, D., *Homologie nicht-additiver Funktoren, Anwendungen*, Ann. Inst. Fourier, Grenoble 11 (1961), 201-312.
17. Goodwillie, T.G., *Calculus I: the first derivative of pseudoisotopy theory*, *K-Theory* 4 (1990), 1-27.
18. Goodwillie, T.G., *Calculus II: analytic functors*, *K-Theory* 5 (1992), 295-332.
19. Goodwillie, T.G., *Calculus III: the Taylor series of a homotopy functor*, in preparation.
20. Goodwillie, T.G., *The calculus of functors*, Preprint 1986 Brown University.
21. Goodwillie, T.G., *Calculus I. The first derivative of a homotopy functor*, Preprint Brown University.
22. Gray, B., *Homotopy theory*, Academic Press (1975).
23. James, J., *On the suspension triad*, Ann. of Math. (2) 63 (1956), 191-247.
24. Johnson, B., *The derivatives of homotopy theory*, Trans. AMS 347 (1995), 1295-1310.
25. Kan, D.M., *A relation between CW-complexes and free C.s.s. groups*, Amer. J. Math. 81 (1959), 521-528.
26. Kan, D.M., *A combinatorial definition of homotopy groups*, Annals of Math. 67 (1958), 282-312.
27. Mac Lane, S., *Homology*, Academic Press (1963).
28. Whitehead, G.W., *Recent advances in homotopy theory*, AMS Regional conference series in math. 5 (1971).
29. Whitehead, G.W., *Elements of homotopy theory*, Graduate text in math. 61, Springer Verlag (1978).