

# ELLIS HORWOOD LIMITED

International Publishers in Science and Technology

Market Cross House  
Cooper Street  
Chichester  
West Sussex  
PO19 1EB  
England

Telephone  
0244 (Chichester) 239412  
7447 46-02 14044 000  
Cable Horwood Chichester

Directors  
Ellis Horwood Chairman  
F. M. Horwood  
Care Horwood  
Michael Horwood  
James Gillman  
S. G. Gibbons



## ELLIS HORWOOD TITLES IN MATHEMATICS AND ITS APPLICATIONS

Ellis Horwood series Mathematics and its Applications

as Editor: G. M. BELL, Professor of Mathematics  
Ellis College, University of London

### GLOBAL RIEMANNIAN GEOMETRY

Edited by T. J. WILLMORE, Professor of Pure Mathematics, University of Durham, and  
HITCHIN, St. Catherine's College, University of Oxford

Outstanding developments have taken place in various aspects of Global Riemannian Geometry since the first Durham Symposium on the subject in 1974. This 1982 symposium concentrates on three of these expanding and closely inter-related areas.

#### and Conjecture and its Consequences

Includes interesting results on real Einstein manifolds and Ricci-flat manifolds, and the existence of Riemannian manifolds in a special homotopy group. In the complex domain there are important repercussions including Severi's conjecture.

Reviews the classification of Riemannian manifolds by their curvature properties, including important recent work by Munn about almost flat manifolds, and manifolds with negative (or positive) curvature. Other recent work concerns Einstein metrics and their existence on a given manifold.

#### harmonic Maps

Outstanding branches of differential geometry, where recent attention has been focused on applications involving harmonic maps. Important research areas include: (a) under what conditions is a harmonic map between Kähler manifolds holomorphic? the problem of rigidity in the complex domain, and (c) the problem of characterizing harmonic maps of  $S^2$  into  $CP^n$ .

Abstracts: Mathematicians concerned with differential geometry for its own sake, and for those interested in its applications to topology and algebraic geometry. Theoretical physicists.

0-85312-526-0 (Library edition)

In Preparation

The L. M. S. Symposium on Global Differential Geometry  
Durham, July 8 - 10, 1982

## HERMITIAN RIEMANN METRICS

Andrzej Derdziński

§ 0. Introduction.

Let  $(M, g)$  be an oriented four-dimensional Riemannian manifold. We shall say that  $(M, g)$  is locally conformally Kählerian if every point  $x$  of  $M$  has a neighbourhood  $U$  with a function  $f > 0$  on  $U$  such that  $f g$  is a Kähler metric for some complex structure in  $U$ . Compatible with the selected orientation.

The aim of the present note is to discuss some properties and examples of flexible manifolds which are locally conformally Kählerian. In the case of orientable Kählerian four-manifolds, this condition does not seem too unnatural. It holds, e.g. (for some orientation), if the manifold has sufficiently many symmetries with fixed points (see (3.5)), as well as for all known examples of compact orientable Kählerian four-manifolds ([7], §5). The list of these examples (up to finite isometric coverings) consists of:

- 1) products of surfaces, 2) locally irreducible locally symmetric spaces,
- 3) compact complex Kähler surfaces having  $c_1 < 0$  or  $c_1 = 0$  with Kähler-Kähler metrics (cf. the existence theorem of B.-T. You and T. Aubin, [10], [2], [15]), 4) the complex surface obtained by blowing up a point in  $CP^2$  (homeomorphic to  $CP^2 \setminus \{pt\} \cup CP^1$ ) with the  $U(2)$ -invariant Hermitian Kähler metric constructed by D. Pappas ([14], [4]).

In §1 and §2 we list some properties of locally conformally Kählerian Kählerian four-manifolds. Some examples of rigid manifolds of this type are described in §3; these examples are either locally conformal to products of surfaces ((3.3) i) by (2.5)), there is no chance of finding essentially new compact examples in this way), or half-flat ((3.4)). An interesting feature of our examples is that, for many of them, the conformally related Kähler metrics are essentially non-complete.

For a given Riemannian metric  $g$ , we shall use the symbols  $\nabla$ ,  $Ric$ ,  $Scal$ ,  $\nabla \cdot \nabla$  and  $\Delta = -g^{ij} \nabla_i \nabla_j$  for the Riemannian connection, Ricci tensor, scalar curvature, Weyl conformal tensor and Laplace operator.

§ 1. Kähler manifolds of real dimension four which are locally conformally Kählerian.

Before discussing the Kählerian four-manifolds which are locally conformally Kählerian, let us consider the corresponding (conformally conformally Kählerian) Kähler four-manifolds. The characterization of the latter is, generically, very simple (equation (1) below). Conditions, characterizing locally conformally Kählerian manifolds among arbitrary Riemannian ones, were found by H. W. Brinkmann [5].

For a Riemannian manifold  $(M, g)$ , the Ricci tensor  $R = R(g)$

$$(1) \quad \text{First studied by H. Bach, [1]} \text{ is defined by } R_{ij} = R^k{}_{ljk} =$$

$\cdot \{ Ric^p{}_{qij} \}$ . Under a conformal change of metric in dimension four, the Bach tensor transforms like  $R(g) = R(g)/r$ . If  $g$  is an Kählerian metric, we have  $R^p{}_{qij} = 0$ ,  $(Ric^p{}_{qij}) = 0$  and hence  $R(g) = 0$ . Consequently, conformally Kählerian  $\Leftrightarrow R(g) = 0$  is necessary in order that a Riemannian four-manifold  $(M, g)$  be locally conformally Kählerian.

(1.1) A Kähler four-manifold  $(M, J, g)$  satisfies  $R(g) = 0$  if and only if

$$(1) \quad 2P^2 Scal = Scal \cdot Ric = (Scal)^2/4 - Abscal^2/2|g$$

([7], Lemma 5). Conversely, for a Kähler four-manifold  $(M, J, g)$  satisfying (1), the conformally related metric  $\bar{g} = g/Scal^2$  (defined wherever  $Scal \neq 0$ ) is Kählerian. This Kählerian metric conformal to  $g$  is essentially unique and, if  $Scal$  is not constant,  $\bar{g}$  is not locally symmetric (since  $2Ric(\bar{g}) \cdot \bar{g}(\bar{g}) = Scal^5$ , cf. [7], Proposition 3). Therefore, the only locally conformally Kählerian four-manifolds are locally conformally Kählerian  $\Leftrightarrow$  and only  $\Leftrightarrow$  it satisfies (1).

(1.2) For a Kähler four-manifold  $(M, J, g)$  satisfying (1),  $J(Scal)$  is a holomorphic Killing vector field ([7], Proposition 4). Thus,  $g$

§ 2. Locally conformally Kählerian four-dimensional

Einstein manifolds.

Let M be an oriented smooth four-manifold. If a Riemannian metric g (or a conformal structure) is chosen in M, the bundle A^2M of 2-forms on M can be written as the Whitney sum A^2M + A^2M, where the 3-plane bundles A^2M consist of all 2-forms l with e l = i l. The Weyl tensor W of (M,g), viewed as an endomorphism of A^2M, always commutes with \* ([16]) and hence it leaves A^2M invariant; we shall denote by V^2 = V^2(g) : A^2M -> A^2M the resulting restrictions of W.

(2.1) In a Kähler four-manifold (M,J,g) endowed with the natural orientation (so that the Kähler form w is a section of A^2M), the eigenvalues of V^2 at any point are ( Scal/6, -Scal/12, -Scal/12 ) and w is an eigenvector of V^2 for the "simple" eigenvalue Scal/6 (cf. [11]). By the conformal invariance of W, this implies that the condition Spec V^2 = 2 ( " V^2 has less than three distinct eigenvalues at any point" ) is necessary for an oriented Riemannian four-manifold to be locally conformally Kählerian (in the sense of §0).

(2.2) Uniqueness of a Kähler metric within a given conformal class: Let (M,J,g) be a Kähler four-manifold, U an open subset of M with V^2 != 0 everywhere in U and F > 0 a function on U such that g = - Fg is a Kähler metric for some complex structure J compatible with the orientation determined by J. Then F is constant and J = iJ. In fact, in view of (2.1), the Kähler form w of g (resp., w of g) is determined (up to a sign) by V^2(g) (resp., by V^2(g) = V^2(g)/F), at any point of U. Thus, w = iFw and hence J = iJ; since dw = -dw = 0, F must be constant.

(2.3) Let (M,g) be an oriented Einstein four-manifold with Spec V^2 = 2.

a) If V^2 = 0 identically, then (M,g) is called an anti-self-dual

Einstein manifold ([1]). In the case where M is compact and Scal >= 0, these manifolds have been classified by M. Hitchin ([12], [13]); see also [10].

b) If V^2 does not vanish identically, then V^2 != 0 everywhere. The global conformal change g = |V^2|^{2/3} g gives rise to a Kähler metric g on M or on some two-fold cover of M (the Kähler form for g is determined by g only up to a sign) and the scalar curvature of g is non-zero everywhere ([7], Proposition 5).

(2.4) Let (M,g) be a compact oriented Einstein four-manifold such that Spec V^2 = 2 and V^2 is not parallel. Then the universal covering manifold N of M is diffeomorphic to S^2 x S^2 or to a connected sum CP^2 # (-kCP^2), 0 <= k <= 8, while the pull-back of the metric |V^2|^{2/3} g to N is a Kähler metric with positive non-constant scalar curvature ([7], Theorem 2).

From (2.3) it follows that, for oriented Einstein four-manifolds with V^2 != 0, local conformal equivalence to a Kähler manifold implies global one (at least for a two-fold cover). The only known example of a compact manifold of this type which is not (locally) Kählerian (i.e., satisfies the hypothesis of (2.4)) is CP^2 # (-CP^2) with the Page metric (f0). The following argument shows that no new compact Einstein four-manifolds can be obtained by conformal deformations of products of surfaces, even without insisting that the underlying manifold be globally diffeomorphic to a product.

(2.5) PROPOSITION. Let (M,g) be a compact Einstein four-manifold. If, for some non-void connected open subset U of M and a function F > 0 on U, the Riemannian manifold (U,Fg) is isometric to a product of surfaces, then either g is a metric of constant curvature, or F is constant and (M,g) is isometrically covered by a product of surfaces.

PROOF: By a result of D. DeTurck and J. Kazdan [9], (M,g) is analytic.

If  $g$  is not of constant curvature, we may assume that  $M$  is oriented so that  $W \neq 0$  somewhere. However, a manifold conformal to a product of surfaces satisfies  $\frac{d}{dt} \text{Spec } W \leq 2$  for both orientations, so that  $W \neq 0$  everywhere by (2.3)b). Combining (2.3)b) with (2.2) we see that  $|W|^{2/3}/P$  is constant, so that analyticity together with the de Rham decomposition theorem implies that the universal covering space  $(\tilde{M}, \tilde{G})$  of  $(M, |W|^{2/3}g)$  is isometric to a product of surfaces. If  $|W|$  were not constant, (2.4) would imply that  $(\tilde{M}, \tilde{G})$  is isometric to  $(S^2, g_1) \times (S^2, g_2)$  for some metrics  $g_1, g_2$  on  $S^2$ , one of which has non-constant Gauss curvature. On the other hand,  $\tilde{G}$  is an extremal Kähler metric (the gradient of its scalar curvature is holomorphic, cf. (1.1), (1.2)) and so the same would hold for  $g_1$  and  $g_2$ . By a theorem of E. Calabi ([6], p. 276), both  $g_1, g_2$  would have constant curvatures. This contradiction completes the proof.

§ 3. Examples.

By (2.3) and (1.1), the only way of obtaining locally conformally Kählerian Kinstein four-manifolds with  $W \neq 0$  is, essentially, to take a Kähler four-manifold  $(M, J, g)$  satisfying (1) and define the Kinstein metric by  $\bar{g} = g/\text{Scal}^2$ . Although this procedure is possible only in the set  $U$  where  $\text{Scal} \neq 0$ , it always gives rise to COMPLETE Kinstein metrics in the components of  $U$ , provided that  $g$  is complete and  $\text{Scal}$  is bounded. Actually, even weaker conditions are sufficient for completeness of  $\bar{g}$ . For convenience, we shall now consider Riemannian manifolds with boundary (empty or not); they are, naturally, metric spaces and their completeness is equivalent to the existence of end-points for any curve of finite length.

(3.1) **LBOW.** Let  $f$  be a bounded  $C^\infty$  function on a complete Riemannian manifold  $(M, g)$  (with boundary). Set  $\bar{g} = g/f^2$  wherever  $f \neq 0$ .

Then, for any component  $Q$  of the set  $M \setminus f^{-1}(0)$ ,  $(Q, \bar{g})$  is a complete Riemannian manifold (with boundary).

**PROOF:** We claim that for any  $C^1$  curve  $\gamma: [a, b] \rightarrow Q$  with  $a < b \leq \infty$ ,  $g(\dot{\gamma}, \dot{\gamma}) = 1$  and of finite  $\bar{g}$ -length  $L = \int_a^b |d\gamma/dt|_{\bar{g}} dt$ , there exists a limit  $\gamma(b) \in Q$ . If  $b = \infty$ , then  $L = \infty$ , since  $f$  is bounded. For  $b < \infty$ , completeness of  $(M, g)$  implies the existence of  $\gamma(b) \in M$ . Note that  $d(f(\gamma(t)))/dt = \kappa(\dot{\gamma}, \dot{\gamma})$  is bounded for  $t < b$ , since  $\gamma([a, b])$  is compact. If we had  $\gamma(b) \in f^{-1}(0)$ , so that  $f(\gamma(b)) = 0$ , then this would give  $|f(\gamma(t))| \leq A(b-t)$  for some  $A > 0$  and all  $t \in [a, b]$  and hence  $L \geq \infty$ , contradicting our assumption.

(3.2) **PROPOSITION.** Let a Kähler four-manifold  $(M, J, g)$  with non-constant scalar curvature satisfy (1).

- i) Every (non-empty) component  $N$  of the set  $\text{Scal}^{-1}((0, \infty))$  (or  $\text{Scal}^{-1}((-\infty, 0])$  which is not a single point, is a four-dimensional submanifold of  $M$  (possibly with boundary).  $(M, \bar{g})$  is complete if so is  $(M, g)$ .

ii) For  $N$  as in (i), if  $(M, g)$  is complete,  $\text{Scal}$  is bounded on  $N$  and  $Q$  is the subset of  $N$  given by  $\text{Scal} \neq 0$ , then  $(Q, g/\text{Scal}^2)$  is a complete Kinstein four-manifold (without boundary), which is not locally symmetric.

In fact, by (1.2) and (1),  $\text{Scal}^{-1}((0))$  is a union of disjoint hypersurfaces and isolated points (note that  $\text{Scal}(x) = 0$  and  $\nabla \text{Scal}(x) = 0$  implies  $\text{Kess } \text{Scal}(x) = -\Delta \text{Scal}(x) = g(x, x)/4 \neq 0$ , since the Killing field  $J(\nabla \text{Scal})$  is determined by its 1-jet at  $x$ ). Applying (1.1) and (3.1) with  $M$  as in i) and  $f = \text{Scal}$ , we obtain ii).

Using (3.2), we shall now describe various explicit examples of complete conformally Kählerian Kinstein four-manifolds.

(3.3) EXAMPLE. Let our Kähler four-manifold be a product of surfaces. (This case was studied by Y. Tashiro, [17].) Relation (1) holds if and only if both surfaces satisfy

$$(2) \quad 2\psi^2 \kappa = (c - \kappa^2)g,$$

$g, \psi$  and  $\kappa$  being now the metric, the connection and the Gauss curvature of the surface, while  $c$  is a common constant; by rescaling  $g$ , we shall assume that  $c \in \{-1, 0, 1\}$ . For a surface  $(S, g)$  satisfying (2),

we have  $|\psi \kappa|^2 = -\kappa^3/3 + \psi \kappa - r = P_{\epsilon, r}(\kappa) \geq 0$  for some real  $r$  and so the integral curves of  $\psi \kappa$  are geodesics. The length of such a geodesic  $\gamma : (t_1, t_2) \rightarrow S$ , containing no critical points of  $\kappa$ , is given by

$$(3) \quad L(\gamma) = \int_{t_1}^{t_2} \sqrt{P_{\epsilon, r}(\kappa)}^{-1/2} dt, \quad \kappa_i = \lim_{t \rightarrow t_i} \kappa(\gamma(t)).$$

If  $S$  is oriented, the complex structure tensor  $J$  gives rise to the Killing field  $J(\psi \kappa)$ . It is now easy to verify that a complete local description of such surfaces at points with  $\psi \kappa \neq 0$  is

$$(4) \quad g = dt^2 + s(dt/dt)^2 dx^2$$

in suitable local coordinates  $(t, x)$ , where  $a > 0$  and  $\kappa = \kappa(t)$  is any solution of the equation  $2d^2\kappa/dt^2 = c - \kappa^2$ , i.e., of

$$(5) \quad (d\kappa/dt)^2 = -\kappa^3/3 + \psi \kappa - r = P_{\epsilon, r}(\kappa)$$

for some  $r$ . The essentially distinct local types of surfaces satisfying (2) with  $\psi \kappa \neq 0$  are, thus, parametrized by  $c \in \{-1, 0, 1\}$  and  $r \in \mathbb{R}$ ; the parameter  $a$  in (4), locally irrelevant ( $adr^2 = d(a^{1/2}x)^2$ ), will later have global meaning. We can now describe some examples of such surfaces  $(S, g)$  with "as much completeness as possible". Our  $g = g_{\epsilon, r, a}$  will be the  $S^1$ -invariant metric, defined by (4) and (5) on the product  $(\inf \kappa, \sup \kappa) \times S^1$  (sometimes "completed" by adding a point), where the second coordinate  $x \in S^1 = \mathbb{R}/2\pi\mathbb{Z}$ , while the first one is parametrized by  $t$ . For brevity, we shall say that  $(S, g)$  is half-complete from above (resp., from below) if, for all  $q \in (\inf \kappa, \sup \kappa)$ ,

$\kappa^{-1}(\{q, \infty\})$  (resp.,  $\kappa^{-1}(\{-\infty, q\})$ ) is a complete surface with boundary.

Let  $\kappa_0 = \kappa_0(t, r)$  be the lowest real root of  $P = P_{\epsilon, r}$ . Note that  $P$  may have three distinct real roots (if  $c = 1$  and  $|\psi| < 2/3$ ), a simple

and a double one (if  $c = 1, |\psi| = 2/3$ ), a triple one ( $c = r = 0$ ), or only one, simple real root (otherwise). Let  $\kappa = \kappa_{\epsilon, r}$  be a solution of (5) with range  $(-\infty, \kappa_0)$ .

A) If  $(\epsilon, r)$  is none of  $(0, 0), (1, 2/3), (1, -2/3)$ , then  $\kappa_0^2 \neq c$  and, for  $a = 4/(c - \kappa_0^2)^2$ , our  $S^1$ -invariant metric  $g = g_{\epsilon, r, a}$  on  $(-\infty, \kappa_0) \times S^1 \cong \mathbb{R}^2 \setminus \{0\}$ , defined by (4), can be extended to a smooth  $S^1$ -invariant metric  $g'$  on  $\mathbb{R}^2$  with  $\kappa(0) = \kappa_0$ . The resulting surface  $A_{\epsilon, r} = (\mathbb{R}^2, g')$  is half-complete from above (since  $\kappa$  is a proper function on  $\mathbb{R}^2$ ).

B) If  $c = 1$  and  $r = -2/3$ , our metric  $g = g_{\epsilon, r, a}$  on  $S = (-\infty, \kappa_0) \times S^1$  is half-complete from above for each  $a$  (which one can easily verify using (3)) and its Gauss curvature  $\kappa \leq \kappa_0 = -1$ . Notation:  $(S, g) = B_a$ .

In the case where all roots of  $P = P_{\epsilon, r}$  are real and the highest one,  $\kappa_1$ , is simple (i.e.,  $c = 1$  and  $-2/3 \leq r < 2/3$ ), denote by  $\kappa_2$  the largest root with  $\kappa_2 < \kappa_1$ . Clearly,  $-1 \leq \kappa_2 < 1 < \kappa_1 \leq 2$ . Take a solution  $\kappa = \kappa_{\epsilon, r}$  of (5) with range  $(\kappa_2, \kappa_1)$ .

C) If  $P'(\kappa_2) \neq 0$  (i.e.,  $(\epsilon, r) \neq (1, -2/3)$ ), the choice of  $a = 4/(1 - \kappa_1^2)^2$  (resp.,  $a = 4/(1 - \kappa_2^2)^2$ ) allows to attach a point  $x_0$  to  $(\kappa_2, \kappa_1) \times S^1$  so that our metric  $g_{\epsilon, r, a}$  can be extended to an  $S^1$ -invariant metric  $g'$  on a disc  $D^2$  with  $\kappa(x_0) = \kappa_1$  (resp.,  $\kappa(x_0) = \kappa_2$ ). The surface  $(D^2, g')$  (resp.,  $(D^2, g') = C_r^+$ ) is half-complete from above (resp., from below).

D) If  $P'(\kappa_2) = 0$  ( $(\epsilon, r, \kappa_1, \kappa_2) = (1, -2/3, 2, -1)$ ), our surface  $D_a = ((\kappa_2, \kappa_1) \times S^1, g)$  is half-complete from below (by (3)) and, for  $a = 4/9$ , one can attach a point  $x_0$  to  $D_a$  so that  $g$  has an extension to a metric  $g'$  on a disc  $D^2$  with  $\kappa(x_0) = \kappa_1$ ; the surface  $(D^2, g') = D$  is complete.

Finally, we have the standard examples:

E) Any complete surface  $(S, g)$  of constant curvature  $c \in \{1, 0, -1\}$  satisfies (2) with  $\kappa = |c|$ . By abuse of notation

here  $(S, g) = E_c$  and associate with  $E_c$  the parameter  $r = 2c/3$ .

Among the above examples, complete ones occur only in D) and E). However, even for the non-complete surfaces of A) - C), certain additional conditions imply, for some of their four-dimensional products, that the subset of the product defined by  $Scal \geq 0$  (or by  $Scal \leq 0$ ) is compacte (which follows from half-completeness of our surfaces) and  $Scal$  is bounded on it. Therefore, by (3.2)ii) they give rise to complete, conformally Kählerian, open Einstein four-manifolds, which have negative scalar curvature (equal to  $48(r+r')$ , cf. notations below) and are not locally symmetric (unless both surfaces are of type E)). The details are presented in the following table (with notational conventions like  $K_0 = K_0(e, r)$ ,  $K'_0 = K_0(e, r')$  etc.). Each of the Einstein four-manifolds described here has an isometry group of positive dimension.

product manifold	additional condition	relation defining a complete 4-submanifold with boundary	topology of the associated complete Einstein 4-manifold
$A_{e, r} \times A_{e, r'}$	$K_0 + K'_0 > 0$	$Scal \geq 0$	$D^4$
$A_{1, r} \times B_a$	$K_0 > 2$	"	$S^1 \times D^3$
$A_{1, r} \times C'_2$	$K'_2 < -K_0 < K'_1$	"	$D^4$
$A_{1, r} \times D$	$K_0 + 2 > 0$	"	"
$A_{g, r} \times E_c$	$c =  c , K_0 > -c$	"	$D^2 \times E_c$
$B_a \times C'_2$	none	"	$S^1 \times D^3$
$B_a \times D$	none	"	"
$C'_2 \times E_{-1}$	none	$Scal \geq 0$ (resp., $Scal \leq 0$ )	$D^2 \times E_{-1}$
$C'_2 \times C'_2$	$K'_2 + K'_2 < 0$	$Scal \leq 0$	$D^4$
$C'_2 \times D_a$	none	"	$S^1 \times D^3$

$C'_2 \times D$	none	"	$S^1 \times D^3$
$D_a \times D_a$	none	"	$T^2 \times D^2$
$D_a \times D$	none	"	$T^2 \times D^2$
$D_a \times E_{-1}$	none	"	$S^1 \times D^1 \times E_{-1}$
$D \times D$	none	$Scal \geq 0$ $Scal \leq 0$	$D^4$ $S^3 \times D^1$
$D \times E_c$	$ c  = 1$	$Scal \geq 0$ $Scal \leq 0$	$D^2 \times E_c$ $S^1 \times D^1 \times E_c$

(3.4) EXAMPLE. Consider a four-dimensional Lie algebra with basis  $e_1, \dots, e_4$  such that  $[e_1, e_\alpha] = 0$  ( $\alpha \geq 2$ ),  $[e_2, e_3] = 2(p-q^2)e_4$ ,  $[e_2, e_4] = 2se_3$ ,  $[e_3, e_4] = 2(p-q^2)e_2$ , where  $p, q \in \mathbb{R}$ ,  $p \neq q^2$  and  $c = \pm 1$ . Let  $\sigma$  be any solution of the equation

$$(6) \quad d\sigma/dt = 2(q - \sigma)(\sigma^2 - p)$$

defined on an interval  $(a, b)$  and such that  $c(\sigma^2 - p) > 0$ ,  $c(q - \sigma) > 0$ .

The open subset  $M = (a, b) \times \mathbb{H}$  of a Lie group  $R \times \mathbb{H}$  associated to our Lie algebra ( $\mathbb{H}$  locally isomorphic to  $SU(2)$  or to  $SL(2, \mathbb{R})$ ) can now be endowed with the Riemannian metric  $g$  given by  $g(e_1, e_1) = g(e_2, e_2) = (q - \sigma)(\sigma^2 - p)$ ,  $g(e_2, e_2) = g(e_4, e_4) = c(q - \sigma)$ ,  $g(e_\alpha, e_\beta) = 0$  ( $\alpha \neq \beta$ ), where  $\sigma$  depends on  $t \in (a, b)$  and  $e_1 = 3/4t$ ,  $e_2, e_3, e_4$  are viewed as right-invariant vector fields on  $R \times \mathbb{H}$ . This metric is preserved by the right action of  $\mathbb{H}$  on  $M$  and it is Kählerian for the complex structure  $J$  given by  $J e_1 = e_3$ ,  $J e_2 = e_4$ . Moreover,  $(M, J, g)$  satisfies (1) and  $Scal = 48\sigma$  is not constant. By (1.2),  $J(\nabla Sc)$  is a non-trivial  $\mathbb{H}$ -invariant Killing field on  $M$ , so that the isometry group of  $(M, g)$  is four-dimensional (locally isomorphic to  $S^1 \times \mathbb{H}$ ). Another property of  $g$  is that it is self-dual ( $\bar{W} = 0$  for the natural orientation); for details, see [9]. Combining this construction with (3.2)ii), we shall now describe some examples of complete Einstein four-manifolds. Take  $c = -1$ ,  $p > 0$ ,  $-1/2 < q < 0$ ,  $\mathbb{H} = SU(2)$  and a solution  $\sigma$  of (6) such that

an interval  $(a,b)$  and having range  $(q,p)^{1/2}$ . The  $U(2)$ -invariant metric  $g$  constructed as above on the manifold  $(a,b) \times S^3$  (which we identify with a pointed ball  $B^4 \setminus \{0\}$  in  $C^2$ ) can be extended to a Kähler metric  $g'$  on  $B^4$  with  $\alpha(0) = \text{Scal}(0)/48 = q$  ([8], §2). In  $(B^4, g')$ , relation  $\text{Scal} < 0$  defines a ball  $Q$  with compact closure. Thus, by (3.2)ii),  $(Q, g'/\text{Scal}^2) = M_{p,q}$  is a complete Hermitian Einstein four-manifold, of negative scalar curvature  $2 \cdot 12 \cdot 3 \cdot p q$ , which is conformally Kählerian, not locally symmetric and self-dual for the natural orientation. The four-dimensional isometry group of  $M_{p,q}$  has three-dimensional principal orbits; various examples of Einstein manifolds with the similar property of "principal cohomogeneity one" have been constructed by L. Bérard Bergery, [4].

(3.5) REMARK. Let  $(M,g)$  be an orientable Einstein four-manifold such that for each  $x \in M$  there exists a non-trivial Killing field defined near  $x$  and vanishing at  $x$  (this happens, e.g., if all orbits of the isometry group  $I(M,g)$  are of dimensions less than  $\dim I(M,g)$ ). By Lemma 9 of [7], we have  $\text{Spec } \hat{W} \leq 2$  for both orientations and so, by (2.3),  $(M,g)$  is locally conformally Kählerian (for an appropriate orientation). However, if  $(M,g)$  is not locally symmetric, one can easily prove, using the Killing field mentioned in (1.2), that the Lie algebra of germs of Killing fields at any point is four-dimensional and has principal orbits of dimension three; a general existence theorem for (incomplete) Einstein metrics of cohomogeneity one is due to N. Koiso and L. Bérard Bergery ([4], Proposition 3).

References

1. M.F. Atiyah, M.J. Hitchin and I.M. Singer: Self-duality in four-dimensional Riemannian geometry. Proc. Roy. Soc. London A 362 (1978), 425-461.
2. T. Aubin: Equations du type de Monge-Ampère sur les variétés kählériennes compactes. C. R. Acad. Sci. Paris A 283 (1976), 119-121.
3. R. Bach: Zur Weylschen Relativitätstheorie und der Weylschen Erweiterung des Krümmungstensorbegriffs. Math. Zeitschr. 9 (1921), 110-135.
4. L. Bérard Bergery: Sur de nouvelles variétés riemanniennes d'Einstein.

5. H.W. Brinkmann: Riemann spaces conformal to Einstein spaces. Math. Ann. 91 (1924), 269-278.
6. E. Calabi: Extremal Kähler metrics. Seminar on Differential Geometry (ed. by S.-T. Yau), Princeton (1982), 259-290.
7. A. Derziński: Self-dual Kähler manifolds and Einstein manifolds of dimension four. Compos. Math. (to appear).
8. A. Derziński: Exemples de métriques de Kähler et d'Einstein auto-duales sur le plan complexe. Géométrie riemannienne en dimension 4 (Séminaire Arthur Besse 1978/79), Paris (1981), 334-346.
9. D. DeTurck and J.L. Kazdan: Some regularity theorems in Riemannian geometry. Ann. Sci. Ec. Norm. Sup. (4) 14 (1981), 249-260.
10. T. Friedrich and H. Kurke: Compact four-dimensional self-dual Einstein manifolds with positive scalar curvature. Math. Nachr. (to appear).
11. A. Gray: Invariants of curvature operators of four-dimensional Riemannian manifolds. Proc. 13th Biennial Seminar Canadian Math. Congress (1971), vol. 2, 42-65.
12. N. Hitchin: Compact four-dimensional Einstein manifolds. J. Differential Geometry 9 (1974), 435-441.
13. M. Hitchin: Kählerian twistor spaces. Proc. London Math. Soc. 43 (1981), 133-150.
14. D. Page: A compact rotating gravitational instanton. Phys. Lett. 79 B (1978), 235-238.
15. Séminaire Palaiseau 1978. Première classe de Chern et courbure de Ricci : preuve de la conjecture de Calabi. Astérisque, vol. 58 (1978).
16. I.M. Singer and J.A. Thorpe: The curvatures of four-dimensional Einstein spaces. Global Analysis, Papers in Honor of K. Kodaira, Princeton (1969), 355-365.
17. Y. Tashiro: On conformal diffeomorphisms of 4-dimensional Riemannian manifolds. Kodai Math. Sem. Rep. 27 (1976), 436-444.
18. S.-T. Yau: Calabi's conjecture and some new results in algebraic geometry. Proc. Natl. Acad. Sci. U.S.A. 74 (1977), 1798-1799.