

MUMFORD-TATE GROUPS AND THE THEOREM
OF THE FIXED PART

Yves André

Max-Planck-Institut
für Mathematik
Gottfried-Claren-Straße 26

D-5300 Bonn 3

MPI/87-21

CONTENTS

1. Some facts about linear algebraic groups
 2. Mumford-Tate groups
 3. Mumford-Tate groups of 1-motives
 4. Variations of mixed Hodge structure
 5. Normality
 6. Maximality
 7. Algebraic independence of Abelian integrals
-
- Appendix
8. Classification of Abelian varieties with many endomorphisms
 9. The Hodge structures H_μ over \mathbb{R}
 10. Automorphisms of $(H_\mu, h_\mu, \varphi_\mu)$

MUMFORD-TATE GROUPS AND THE THEOREM

OF THE FIXED PART

Yves André

The present paper grew out of an attempt of understanding group-theoretically the consequences of Hodge theory which are explained in Deligne [4] II 4, with an eye towards applications to algebraic independence.

After some preliminaries about representations of linear algebraic groups, we define and study Mumford-Tate groups of mixed Hodge structures over noetherian subrings R of the field \mathbb{R} of real numbers. Though in the sequel we restrict ourselves to the crucial case $R = \mathbb{Z}$, we refer to the appendix for a study of some pathologies which may occur in the case of other ground rings. We then turn to a more precise study of Mumford-Tate groups arising from 1-motives (see [4] III 10).

In the fourth paragraph a mild generalization of a result by Deligne about the monodromy of variation of Hodge structure is given; we also present our main object of study, that is Streeter-Zucker's notion of a good variation of mixed Hodge structure.

In paragraph 5, we give a group-theoretic formulation of the theorem of the fixed part proved in [12]: for almost all stalks of a given polarizable good variation of mixed Hodge structure, the connected monodromy group H_x is a normal subgroup of the derived

Mumford-Tate group $\mathcal{D}G_x$. We then state straightforward consequences about monodromy groups. In the next paragraph, we study how big can H_x be in $\mathcal{D}G_x$; we end by applying these considerations to the study of algebraic independence of Abelian integrals depending on some parameters.

Acknowledgements: first of all, I want to express my gratitude to the Max-Planck-Institut für Mathematik, Bonn, for generous hospitality which I appreciate very much. A preliminary version of this paper served as a base text for talks given at the seminar about Hodge theory in the Max-Planck-Institute; I thank the participants for their interest. The present study of Mumford-Tate groups of 1-motives (§ 3) grew out of a question of D. Bertrand; I thank him. Finally the computations in the Appendix, and some results in § 6, were worked out during my military service; I thank the authorities from E.A.M.E.A. (Cherbourg) for allowing suitable conditions to deal with such technicalities.

Note: while this work was almost completed, J.P. Winterberger has pointed to me a recent paper by G.A. Mustafin "Families of algebraic varieties and invariant cycles" Math. Izv. 27 (1986) n°2, where the author also compares H_x and G_x under a strong degeneration hypothesis; in that paper, the normality property in the "projective smooth situation" is stated, and attributed to P. Deligne.

1. Some facts about linear algebraic groups

Let K be a field of characteristic 0, and $V \cong K^N$ some K -vector space. We shall consider closed an algebraic subgroup $G \subset GL(V) = GL_N$. For non-negative integers m, n , we set $T^{m,n} = T^{m,n}(V) = V^{\otimes m} \otimes V^{\otimes n}$, where V^{\vee} denotes the dual space of V (with the contragredient action of GL_N). By "representation of G " as " G -module", we shall always mean a finite-dimensional rational one. The following three properties are well-known [13; 3.5 § 16.1], [6; I 3.1]:

- 1) every representation of G is a subquotient representation of a finite direct sum of $T^{m,n}$'s,
- 2) G is the stabilizer of some one-dimensional L in some finite direct sum $\bigoplus T^{m_i, n_i}$: $G = \text{Stab } L$,
- 3) (not used here) if G is reductive (that is, if V is a semi-simple representation of G), one can choose L so that G acts trivially on it; for l a generator of L , we then write $G = \text{Fix } l$.

For any representation W of G , and any character $\chi \in X_K(G)$ of G over K , we denote by W^G the fixed part of W under G and by W^χ the submodule of W on which G acts according to χ . We write $\text{End}_G W$ for the endomorphisms of the G -module W , so that $\text{End}_G W = (\text{End}_K W)^G$, and we denote by $Z(\text{End}_G W)$ its center.

Lemma 1. Assume that G is connected, and let $H \subset G$ be a closed subgroup. The following conditions are equivalent:

- i) $H \triangleleft G$, that is, H is normal in G ,
- ii) for every tensor space $T^{m,n}$, and for every $\chi \in X_K(H)$,
 $(T^{m,n})^\chi$ is stable under G ,
- iii) every H -isotypical component of any representation of G is stable under G .

If moreover G is reductive, these conditions imply that
 $Z(\text{End}_H V) \subset Z(\text{End}_G V)$.

Proof: iii) \Rightarrow ii) is obvious, and we shall first prove that ii) \Rightarrow i), independently of the connectedness assumption on G . We know by 2) that there exists some one-dimensional L in some $\oplus T^{m_i, n_i}$ such that $H = \text{Stab } L$. Let W be the G -module spanned by L . The line L defines a character $\chi \in X_K(H)$; we have $L \subset W^\chi$, and $W^\chi = W \cap (\oplus T^{m_i, n_i})^\chi = W$, according to the hypothesis ii). Let φ be the natural morphism $G \rightarrow GL(\text{End } W)$; it is clear that $H \subset \ker \varphi$. Conversely if $g \in \ker \varphi$, g commutes with any endomorphism of W , that is, g is scalar; this implies that g stabilizes L , so that $g \in H$. Hence $H = \ker \varphi$ is a normal subgroup.

We now prove i) \Rightarrow iii). Let W be a G -module, and W' the G -submodule of the sum of its irreducible submodules. It suffices to prove that the H -isotypical components of W' are G -stable. Let H' , G' denote the natural images of H and G respectively in $GL(W')$, so that $H' \triangleleft G'$. The normality property implies that $(\text{End } W')^{H'}$ is stable under G' , inside the G' -module $\text{End } W'$. For $w \in \text{End}_H W'$, let C_w be the kernel of the commutator map $[w, \cdot]$ in $\text{End}_H W'$. It is easy to derive the formula $gC_w = C_{gw}$,

so that $Z(\text{End}_{H,W'}) = \bigcap_{w \in \text{End}_{H,W'}} C_w$ is again a G' -module. But

$Z(\text{End}_{H,W'})$ is a finite-dimensional semi-simple algebra over K ;
by the connectedness of G' , the morphism $G' \rightarrow \text{Aut}_K(Z(\text{End}_{H,W'}))$
thus has trivial target, that is, $Z(\text{End}_{H,W'})$ is a trivial G' -module.

Now the H -isotypical components of W' are given by $p.W'$, where
 p runs among the minimal idempotents of $Z(\text{End}_{H,W'})$. We just
proved that p commutes with the action of G' on W' , and this
implies that $p.W'$ is stable under G' .

When G is reductive, we have $V' = V$, and the above proof shows
that $Z(\text{End}_H V)$ is a trivial G -module, whence an obvious imbedding
 $Z(\text{End}_H V) \subset Z(\text{End}_G V)$.

□

2. Mumford-Tate groups

We first recall some definitions. Let R be some Noetherian subring of \mathbb{R} such that $K := R \otimes \mathbb{Q}$ is a field. Let V be a noetherian R -module. A (pure R -) Hodge structure of weight $M \in \mathbb{Z}$ over V is a morphism $h : \text{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m \rightarrow \text{GL}(V_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{R})$ such that $hw(x)$ is the multiplication by x^M ; here w denotes the embedding $\mathbb{G}_m \otimes_{\mathbb{R}} \mathbb{R} \hookrightarrow \text{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m$ given by $\mathbb{R}^{\times} \subset \mathbb{C}^{\times}$. Equivalently, it is a bigraduation on $V \otimes_{\mathbb{R}} \mathbb{R} := V_{\mathbb{C}} = \bigoplus_{p+q=M} V^{p,q}$ with $\overline{V^{p,q}} = V^{q,p}$, or else a decreasing filtration F^p on $V_{\mathbb{C}}$ such that $F^p \otimes \overline{F}^{(M-p+1)} \xrightarrow{\sim} V_{\mathbb{C}}(F^p = \sum_{p' \geq p} V^{p', M-p'})$. For instance, there is one and only one Hodge structure of weight $-2M$ on $V = (2\pi\sqrt{-1})^M \mathbb{R}$, called "the Tate twist" and denoted by $R(M)$. A polarisation of the Hodge structure (V, h) of weight M is a morphism of Hodge structures (in the obvious sense) $\psi : V \otimes V \rightarrow R(-M)$ such that $(2\pi\sqrt{-1})^M \psi(\cdot, h(\sqrt{-1})\cdot)$ is a scalar product on $V_{\mathbb{R}} := V \otimes \mathbb{R}$. Elements of $T^{m,n}(V_K) := V^{\otimes m} \otimes (\text{Hom}(V, R))^{\otimes n} \otimes \mathbb{Q}$ (endowed with the natural \mathbb{Z} -K-Hodge-structure of weight $(m-n)M$) which are of type $(0,0)$ are called "Hodge tensors". In fact Hodge tensors are nothing but elements of $F^0(T^{m,n}(V_{\mathbb{C}})) \cap T^{m,n}(V_K)$. A mixed R -Hodge structure (M.H.S) is a noetherian R -module V , together with a finite filtration W of the K -space $V_K := V \otimes_{\mathbb{Z}} \mathbb{Q}$, and a finite decreasing filtration F' of $V_{\mathbb{C}}$ such that the $(\text{Gr}_n^W(V_K), \text{Gr}_n^W(F'))$ are K -Hodge structures of weight n respectively. We shall consider the category of mixed R -Hodge structures up to

isogeny, whose objects are mixed R-Hodge structures, and whose morphisms are the homomorphisms of the associated K-vector space which preserve the filtrations. We say that a M.H.S. V is of type $\epsilon \subset \mathbf{Z} \times \mathbf{Z}$ if its Hodge numbers $h^{p,q}$ are 0 for $(p,q) \notin \epsilon$.

The category of mixed R-Hodge structures up to isogeny is an Abelian K-linear tensor category [4; th 12.10] which is rigid and has an obvious exact faithful K-linear tensor functor

$\omega : (V, W, F) \mapsto V_K$. Let $\langle V \rangle$ denote the Tannakian subcategory generated by (V, W, F) , and ω_V the restriction of the fiber functor to $\langle V \rangle$. Then the functor $\underline{\text{Aut}} \otimes (\omega_V)$ is representable by some closed K-algebraic subgroup $G = G(V)$ of $GL(V_K)$, and ω_V defines an equivalence of categories $\langle V \rangle \xrightarrow{\sim} \underline{\text{Rep}}_K G$, cf. [6; II.2.11]. We call G the Mumford-Tate group of (V, W, F) .

Lemma 2. The Mumford-Tate group G is connected. Any tensor fixed by G in some $T^{m,n}$ is a Hodge tensor (an element of $F^0(T^{m,n}(V_{\mathbb{T}})) \cap T^{m,n}(V_K)$), and G is the biggest subgroup of $GL(V_K)$ which fixes Hodge tensors. If (V, W, F) arises from a pure Hodge structure (V, h) , G is the K-Zariski closure of the image of h in $GL(V_K)$, and if moreover V is polarizable, then G is reductive.

Remark: the definition of Mumford-Tate group above is slightly distinct from that given in [6; I, 3.2] in the case of pure Hodge structures; however if the weight is non-zero, this leads to an isogenous group.

Proof of the lemma: let us first prove the second and third assertions. Any invariant tensor l under G span a trivial representation L_K corresponding to a M.H.S., say L , such that $\langle L \rangle$ is equivalent to Vect_K . Thus L is a trivial M.H.S., that is to say, l is a Hodge tensor. By 1.2), we know that G is the stabilizer of some line L_K in $\otimes T^{m_i, n_i}$, which corresponds to a M.H.S. of rank one (up to isogeny), that is, to some Tate twist $L = R(N_1)$. We can assume V non-trivial (up to isogeny); thus there exists an integer N such that the weight of $\text{Det } W_N(V_K)$, say N_2 , is non-zero. Taking if necessary V_K instead of V , one can assume moreover that N_1 and N_2 have the same sign. Let r be the rank of the M.H.S. $W_N(V_K)$ over K , and let l be a generator of the one-dimensional subspace

$$L_K^{\otimes |N_2|} \otimes (\wedge^r W_N(V_K))^{\otimes 2|N_1|} \text{ inside } (\otimes T^{m_i, n_i})^{\otimes |N_2|} \otimes (\wedge^r V_K)^{\otimes 2|N_1|}.$$

Then $G = \text{Fix}(l)$. The arguments given in [6; I, 3, 4-6], with minor modifications in order to take into account the difference of definitions, prove the statements about pure Hodge structures.

To prove that G is connected in the general case, it suffices to show that any K -space V'_K on which G acts through a finite group is in fact a trivial representation. Such a V'_K correspond to some M.H.S. V' , and we have to show that V' is trivial up to isogeny. The group G acts on each quotient $\text{Gr}_n^W V'_K$ through the Mumford-Tate group G'_n of this pure Hodge structure over K , which is therefore finite. By the description of the Mumford-Tate group of a pure Hodge structure as a Zariski closure over K of some real

torus, it follows that G'_n is trivial, so that $\text{Gr}_n^W V'_K$ is a trivial Hodge structure (that is, of type $(0,0)$). This implies that $W_n V' = F^n V' = 0$ for $n \neq 0$, and finally that V'_K is a trivial representation of G by definition of the Mumford-Tate group.

□

Remark: the description of Mumford-Tate groups by their invariant tensors implies some restrictions on the groups which may occur; for example, G cannot be a Borel subgroup of $\text{GL}(V_K)$, cf. [6; I 3.2]. However, there are other restrictions on the structure of Mumford-Tate groups, as we shall see now:

Lemma 3. Let G be the Mumford-Tate group of a M.H.S. over R , say V , such that $\text{Gr}^W V$ is polarizable. Then the abelianized group $G^{\text{ab}} = G/\mathcal{D}G$ is a torus. The group of real points of its quotient $G^{\text{ab}}/G^{\text{ab}} \cap \mathbb{G}_m$ is compact.

Proof: since all morphisms in $\langle V \rangle$ are strict, one has $\text{Gr}^W V' \in \text{Ob} \langle \text{Gr}^W V \rangle$ for any $V' \in \text{Ob} \langle V \rangle$, thus $\text{Gr}^W V'$ is polarizable. Take for V' the M.H.S. corresponding to a faithful representation V'_K of the quotient U of G^{ab} by its maximal torus. We find that $G(\text{Gr}^W V') = 0$ (see lemma 2). Thus V' , which is a successive extension of trivial H.S., is also a trivial H.S., and $G(V') = U = 0$. Now let $\chi \in X_{\mathbb{C}}(G) = X_{\mathbb{C}}(G^{\text{ab}})$, and let $V''_{\mathbb{R}}$ be some real plane such that $V''_{\mathbb{C}} \simeq X \oplus \bar{X}$; after twisting à la Tate, $\text{Det } V''_{\mathbb{R}}$ corresponds to a trivial real H.S. Therefore $G^{\text{ab}}/G^{\text{ab}} \cap \mathbb{G}_m \Big|_{\mathbb{R}}$

acts trivially on $\text{Det } V_{\mathbb{R}}''$, which yields $|\chi| = 1$. All representations of $G^{\text{ab}}/G^{\text{ab}} \cap \mathbb{G}_m \Big|_{\mathbb{R}}$, are unitary, so that this torus is compact.

□

Remark. The same argument shows in the same situation that if G is nilpotent, then $G = \mathbb{G}_m \times T$ (or $G = T$ if V is pure of weight 0), where T denotes a compact torus.

3. Mumford-Tate groups of 1-motives

We recall that a 1-motive over \mathbb{C} , denoted by $M = [X \xrightarrow{u} E]$, is the following couple of data:

i) an extension $0 \rightarrow T \rightarrow E \rightarrow A \rightarrow 0$ of an Abelian variety A by a torus T ,

ii) a morphism u from a free Abelian group X to $E(\mathbb{C})$. One associates to a 1-motive a mixed Hodge structure

$V = V(M) = (V_{\mathbb{Z}}, W, F^{\bullet})$, given by:

$$\begin{aligned} V_{\mathbb{Z}} &= \{(1, x) \in \text{Lie } E \times X \mid \exp 1 = u(x)\} \\ W_0 &= V_{\mathbb{Q}} \\ W_{-1} &= H_1(E) \otimes \mathbb{Q} \text{ and } Gr_{-1} \text{ is polarizable,} \\ W_{-2} &= H_1(T) \otimes \mathbb{Q} \\ F^0 &= \text{Ker}(W_0 \otimes \mathbb{C} \rightarrow \text{Lie } E), \text{ see [4] III 10.} \end{aligned}$$

We denote by G the Mumford-Tate group of V , and by G_{-1} that of W_{-1} . Let E' be the Zariski closure of $u(X)$, and let us write $F := \text{End } E' \otimes \mathbb{Q}$.

Proposition 1. Let $H \triangleleft G$ such that $W_0^H \subseteq W_{-1}$ (for instance we may take $H = G$). Let us assume that E is a split extension. Then $U(H) := \text{Ker}(H \rightarrow G(W_{-1}))$ is canonically isomorphic to $\tilde{U} := \text{Hom}_F(F \cdot u(X); H_1(E') \otimes \mathbb{Q})$.

Proof (inspired from Kummer's theory of division points on Abelian varieties): the map $U(M) \rightarrow W_{-1} : \sigma \mapsto \sigma m$, depends only on the image (under u) of the class of $m \in W_0$ modulo W_{-1} . This map

defines therefore a G -equivariant homomorphism

$U(H) \xrightarrow{\varphi} \text{Hom}_{\mathbb{Z}}(u(X); W_{-1})$. The vanishing of $\varphi(\sigma)$ implies that σ fixes W_0 , which is a faithful representation of H ; thus $\sigma = 1$, and this shows the injectivity of φ . Because of Poincaré's complete reducibility lemma, the exact sequence of 1-motives $0 \rightarrow [X \rightarrow E'] \rightarrow [X \rightarrow E] \rightarrow [0 \rightarrow E/E'] \rightarrow 0$ splits (up to isogeny).

It follows that

$$\begin{aligned} \text{Ker}(H \rightarrow G(W_{-1})) &= \text{Ker}(H \rightarrow G(H_1(E'))) \cap \text{Ker}(H \rightarrow G(H_1(E/E'))) \\ &\subseteq \text{Ker}(H' \rightarrow G(H_1(E'))) ; \end{aligned}$$

where $H' = H \cap G(V([X \xrightarrow{u} E']))$. Thus φ factorizes through $\text{Hom}_{\mathbb{Z}}(u(X); H_1(E') \otimes \mathbb{Q})$; also it is easily seen that elements in the image of φ are F -linear in some suitable sense: $\varphi(U(H)) \subseteq \tilde{U}$.

Replacing E by E' and X by $u(X)/_{\text{torsion}}$, we may now assume that u is a dominant embedding; we identify X and $u(X)$.

Since E is a split extension, we have $F \simeq \text{End}_{G_{-1}} W_{-1}$, whence $\text{End}_{G_{-1}} \tilde{U} \simeq (\text{End}_F FX)^{\text{op}}$; also W_{-1} , whence \tilde{U} (with trivial action of G_{-1} on FX), is a semi-simple G_{-1} -module. Thus $\varphi(U(H))$ is the kernel of some G_{-1} -equivariant endomorphism ψ of \tilde{U} ; that is to say, there exists $f \in F$ such that $(\psi\varphi(\sigma)) \cdot m = \sigma fm - fm = 0$, $\forall \sigma \in U(H)$, $\forall m \in FX$. If $\varphi(U(H)) \neq \tilde{U}$, then $\psi \neq 0$, therefore we can find $x \in FX$ such that $U(H)x = x$ and $x \neq 0$.

We set $X_x = \mathbb{Z}x$, $M_x = [X_x \hookrightarrow E]$, and we denote by a subscript x the objects G_x, V_x etc. ... associated to this 1-motive. By construction the natural mappings

$$H_x = H \cap G_x \begin{array}{l} \xrightarrow{i} \text{GL}(W_{x,0}) \\ \xrightarrow{j} \text{GL}(W_{x,-1}) \end{array} \quad \text{have the same image, and } i \text{ is}$$

injective. Since E splits, $W_{x,-1} \simeq W_{-1}$ is a direct sum of polarizable pure Hodge structures, so that $j_! H_x \ll j_! G_x$ is reductive. Therefore $W_{x,-1}$ is a direct summand in the H_x -module $W_{x,0}$, which means that we could choose $x \notin W_{x,-1}$ so that $H_x x = x$: indeed, H_x acts trivially (like G_x) on $W_{x,0}/W_{x,-1}$ whose type is $(0,0)$.

Recall that $W_0^H \subseteq W_{-1}$; this implies the corresponding inclusion $W_{x,0}^H \subseteq W_{x,-1}$ since H commutes with the action of F . Therefore we get a contradiction, and deduce that $\varphi(U(H)) = \tilde{U}$.

Corollary. If E splits ($E = A \times T$), with A non trivial, one has a split exact sequence $0 \rightarrow \tilde{U} \rightarrow G \xrightarrow{\varphi} G(H_1(A)) \rightarrow 0$.

Remark: if one drops the assumption that E is split $U(G)$ can be much smaller than \tilde{U} . In "Deficient points on extensions of abelian varieties by \mathbb{G}_m " J. Number theory - (1987), O. Jacquinet and K. Ribet have build some examples (by means of endomorphisms of A which are antisymmetric with respect to a polarization) where $U(G) = 0$, corresponding to some selfdual 1-motives.

4. Variations of mixed Hodge structure

In the sequel we shall concentrate on the case $R = \mathbb{Z}$ (see the appendix for other ground rings). By a variation of M.H.S., we shall mean a finitely filtered object in the category of local systems of noetherian \mathbb{Z} -modules over a fixed connected complex manifold X ,

$$(\underline{V}_{\mathbb{Z}}, W) , W_n \underline{V}_{\mathbb{Z}} \subset W_{n+1} \underline{V}_{\mathbb{Z}} ,$$

together with a decreasing filtration of the complex bundle $V_{\mathbb{C}}^{\mathbb{C}}$ attached to $\underline{V}_{\mathbb{C}} := \underline{V}_{\mathbb{Z}} \otimes \mathbb{C}$ by subbundles F^p , such that on each fibre $\underline{V}_{\mathbb{Z},s}$, (W, F) induces a M.H.S. and that the flat covariant derivative ∇ satisfies $\nabla F^p \subset F^{p-1} \otimes \Omega_X^1$. A morphism of variation of M.H.S. is a morphism of local system which respects W and whose complexification respects the filtration F^p pointwise. This yields an abelian category (any morphism is strictly compatible with the fibrations).

We call such a variation $(\underline{V}_{\mathbb{Z}}, W, F)$ a graded-polarizable one if each of the families $Gr_n^W \underline{V}_{\mathbb{Z}}$ carries a bilinear form with values in $\mathbb{Z}(-n)_X$ which is a morphism of local system and pointwise a polarization. Any subquotient of a polarizable variation and any object isogenous to a polarizable one are polarizable.

The integral relative cohomology modules of the complement of a divisor with relatively normal crossings in a projective smooth X -scheme furnish examples of polarizable variations of M.H.S. over

the algebraic variety X (see [7; 4.3] for instance).

For a variation of M.H.S., and for a point x of X , we denote by H_x the connected monodromy group, that is the neutral component of the smallest algebraic subgroup of $GL(V_{\mathbb{Q},x})$ containing the image of $\pi_1(X,x)$. We also denote by G_x the Mumford-Tate group of the M.H.S. carried by the stalk $V_{\mathbb{Z},x}$.

Lemma 4 (see [5; 7.5]) On the complement $\overset{\circ}{X}$ of some meager subset of X , G_x is locally constant. If the variation is polarizable, then $H_x \subset G_x$ for any $x \in \overset{\circ}{X}$.

Proof: for a polarizable variation of pure Hodge structure, this is stated in loc. cit., without much detail about the proof however. So we shall write down a detailed proof, though (thanks to lemma 2) there is no new complication involved with the M.H.S.. Let \tilde{X} be the universal covering of $(X,0)$, for some base point $0 \in X$. The inverse image of the (polarized) variation of M.H.S. is a (polarized) variation of M.H.S. over \tilde{X} , whose underlying filtered local system $(\tilde{V}_{\mathbb{Z}}, \tilde{W})$ is constant. For $l \in T^{m,n}(\Gamma\tilde{V}_{\mathbb{Q}}) \cong T^{m,n}(\tilde{V}_{\mathbb{Q},0})$, we set

$$\begin{aligned} \tilde{X}(l) &:= \{x \in \tilde{X} / l_x \in T^{m,n}(\tilde{V}_{\mathbb{Q},x}) \text{ is a Hodge tensor}\} \\ &= \{x \in \tilde{X} / l_x \in F^0 T^{m,n}(\tilde{V}_{\mathbb{C},x})\}. \end{aligned}$$

Since F^0 is a subbundle, $\tilde{X}(l)$ is an analytic subvariety of \tilde{X} , and its natural projection $\pi_*\tilde{X}(l)$ on X is an analytic subvariety

too. We set $\overset{\circ}{X} = X \setminus \left(\bigcup_l \pi_* \tilde{X}(l) \right)$, which is a (dense) countable intersection of dense open subsets of X . By definition of $\overset{\circ}{X}$, any $l \in T^{m,n}(\Gamma \tilde{V}_{\mathbb{Q}})$, whose stalk at some $x_0 \in \pi^{-1} \overset{\circ}{X}$ is a Hodge tensor, is in fact a Hodge tensor at any point of \tilde{X} . For $x \in \overset{\circ}{X}$, G_x is then the biggest subgroup of $GL(V_{\mathbb{Q},x})$ which fixes the various tensors in $T^{m,n}(V_{\mathbb{Q},x})$ which lift to $F^0 T^{m,n}(\Gamma \tilde{V}_{\mathbb{Q}})$. Therefore G_x is locally constant on $\overset{\circ}{X}$.

We now assume that the variation is polarized and we shall see that $\pi_1(X,x)$ acts (through a finite group) on the spaces $HT_x^{m,n}$ of Hodge tensors in $T^{m,n}(V_{\mathbb{Q},x})$ for any $x \in \overset{\circ}{X}$; this will be sufficient to prove the lemma, since G_x can be described as $\text{Fix}(l)$, for one element l of one space $\oplus HT_x^{m_i, n_i}$. We have seen that $HT_x^{m,n}$ (for $x \in \overset{\circ}{X}$) is the subspace of $T^{m,n}(V_{\mathbb{Q},x})$ composed of tensors which lift to $F^0 T^{m,n}(\Gamma \tilde{V}_{\mathbb{Q}})$; in particular this subspace is locally constant. Hence $HT_x^{m,n}$ is the rational stalk at x associated to a sub-variation of M.H.S. $(\underline{V}_{\mathbb{Z}}, W., F^{i\cdot})$ of $(T^{m,n}(\underline{V}_{\mathbb{Z}}), T^{m,n}(W.), T^{m,n}(F^{i\cdot}))$, which is actually pure of type $(0,0)$ and which inherits a polarization. This polarization ψ on $V_{\mathbb{R},x}^i$ is a scalar product, invariant under $\pi_1(X,x)$. Thus $\pi_1(X,x)$ factors through the discrete group $\text{Aut } V_{\mathbb{Z},x}^i$ on one side, and through the compact orthogonal group $O(V_{\mathbb{R},x}^i, \psi)$ on the other side; hence the connected group H_x acts trivially on $HT_x^{m,n}$.

□

Remark: a variation of M.H.S. \underline{V} is said to be semi-simple if for any $x \in X$, the relevant category $\langle \underline{V}_x \rangle$ is semi-simple (notations of § 2). It is easily seen that a polarizable M.H.S.

is semi-simple if and only if it is a finite direct sum of variations of pure H.S. up to isogeny. Indeed, using lemma 1, we can see that both conditions imply the reductivity of G_x for any $x \in X$. Conversely, assume that for some $x \in \overset{\circ}{X}$, G_x is reductive. Then by local constancy of G_y on $\overset{\circ}{X}$, the same is true for G_y for any $y \in \overset{\circ}{X}$.

Now consider a section σ of the inclusion

$(W_m)_y \subseteq (W_{m+1})_y$ in the category $\langle V_y \rangle$, and let $\gamma_{y,z}$ be a path (up to homotopy) from y to a nearby point z in $\overset{\circ}{X}$. Then because of the horizontality of the filtration W and the local constancy of $(G_y)_{y \in \overset{\circ}{X}}$, the section $\gamma_{y,z}(\sigma)$ deduced by transporting σ along $\gamma_{y,z}$ is a section of $(W_m)_z \subseteq (W_{m+1})_z$ in the category $\langle V_z \rangle$. Thus $\underline{V}|_{\overset{\circ}{X}}$ is a direct sum of variations of pure H.S. up to isogeny, which extend to X by continuity. The semi-simplicity of \underline{V} follows from this.

We shall now recall a concept introduced by Steenbrinck-Zucker [12] (cf. also [15]). Let us now assume that X is a smooth connected algebraic variety over \mathbb{C} . The variation of mixed Hodge structure is considered good if it satisfies the following condition at infinity: there exists a compactification \bar{X} of X , for which $\bar{X} - X$ is a divisor with normal crossings, such that

i) The Hodge filtration bundles F^p extend over \bar{X} to sub-bundles \tilde{F}^p of the canonical extension $\tilde{V}_{\mathbb{C}}^C$ of $V_{\mathbb{C}}^C$, such that they induce the corresponding thing for $\text{Gr}^W(V_z, W, F^*)$,

ii) for the logarithm N_j of the unipotent part of a local monodromy transformation about a component of $\bar{X} \setminus X$, the weight

filtration of N_j relative to W exists.

The fact that these conditions are sufficient to imply those of [12] (3.13) is pointed out in [15] 1.5, and follows from [16] 4 and [12] A. The following classes of variations of M.H.S. are good:

1) polarizable semi-simple variations of M.H.S. over algebraic bases [10], [14]

2) relative cohomology modules of the complement of a divisor with relatively normal crossings in a projective smooth X-scheme, at least when X is a curve, see [12] 5.7. Moreover, the category of good variations of M.H.S. over X is stable under standard constructions of linear algebra, \otimes , \otimes , duality ... , see [12] A .

Example: smooth 1-motives.

Recall from [4] III 10.1.10 that a smooth 1-motive \underline{M} over X is the following couple of data:

$$\text{i) an extension } 0 \rightarrow \underline{T} \rightarrow \underline{E} \rightarrow \underline{A} \rightarrow 0 \text{ of a (polarizable) Abelian}$$

$$\begin{array}{c} \downarrow f \\ X \end{array}$$

scheme \underline{A} over X by a torus \underline{T} over X

ii) a morphism $\underline{u} : \underline{X} \rightarrow \underline{E}$ from a group scheme \underline{X} over X to \underline{E} ; one assumes that locally for the etale topology on X, \underline{X} is constant and defined by a free \mathbb{Z} -module of finite type.

The construction $\underline{V}(M) = (\underline{V}_{\mathbb{Z}}, W., F^{\bullet})$:

$$\underline{V}_{\mathbb{Z}} = W_0(\underline{V}_{\mathbb{Z}}) = \underline{\text{Lie}} \underline{E}/\underline{X} \times_{\underline{E}} \underline{X} \text{ defined by the exponential sequence,}$$

$$W_{-1} = \text{Ker exp} = R_1 f_{\star}^{\text{an}} \mathbb{Z}$$

$$W_{-2} = (X_{\mathbb{C}}(\underline{T}))^{\vee}$$

$$F^0 = \text{Ker} (V_{\mathbb{C}}^{\vee} \rightarrow \underline{\text{Lie}} \underline{E}/\underline{X})^{\vee},$$

which is fibrewise compatible with that of § 3, yields a polarizable variation of M.H.S. over X .

Lemma 5. Assume that X is a curve. Then the variation $V(\underline{M})$ associated to the smooth 1-motive \underline{M} is good.

(Sketch of) Proof: according to M. Raynaud [C.R.A.S. 262 (1966) 413-416], there exists a Néron model of \underline{E} over the smooth completion \bar{X} of X , such that \underline{u} extends to $\bar{u} : \bar{X} \rightarrow \bar{E}$; note that the

$$\begin{array}{ccc} \bar{X} & \longrightarrow & \bar{E} \\ & \searrow \bar{f} & \\ & & \bar{X} \end{array}$$

smooth group scheme \bar{E}/\bar{X} is not of finite type in general. Replacing \underline{X} by a subgroup-scheme of finite index, which yields an isogeneous variation of M.H.S., we may assume that $\bar{u}(\bar{X})$ lies in the neutral component \bar{E}^0 of \bar{E} . Condition i) defining good variations is fulfilled with $\tilde{F}^0 = \text{Ker} ((\text{Lie } \bar{E}^0/\bar{X} \times_{\bar{E}^0} \bar{X})^c \rightarrow \text{Lie } \bar{E}^0/\bar{X})$.

In order to verify ii), we may proceed by induction since we know that both W_{-1} (by point 2) above: the geometric situation) and W_0/W_{-2} (by duality of 1-motives and point 2) satisfy ii); the point is that $N_j W_{-1} \cap W_{-2} = 0$.

Granting ii) for W_{-1} , it follows from theorem 2.20 of [12] (formula 2.21) that ii) for W_0 reads equivalently:

$$(*) N^1 W_0 \cap W_{-1} \subset N^1 W_{-1} + (-2)^{M_{-1-1}}, \text{ for all } l > 0; \text{ here } (-2)^{M_{-1-1}}$$

is the relative weight filtration of W_{-2} , which is W_{-1-1} since the unipotent part of the local monodromy of W_{-2} is trivial (see [12] 2.14; the point is that \underline{T} is necessarily locally constant).

Therefore (*) follows from property ii) for W_0/W_{-2} .

□

5. Normality

We keep the notations of the previous paragraph. The following result is a simple consequence of the theorem of the fixed part (Griffiths-Schmid-Steenbrinck-Zucker).

Theorem 1. Let $\underline{V} = (\underline{V}_{\mathbf{Z}}, W., F')$ be a (graded-) polarizable good variation of mixed Hodge structure over a smooth connected algebraic variety X . Then for any $x \in \overset{\circ}{X}$, the connected monodromy group H_x is a normal subgroup of the derived Mumford-Tate group $\mathcal{D}G_x$.

Proof: we first prove that $H_x \triangleleft G_x$, using the implication ii) \Rightarrow i) in lemma 1. Since we already know that $H_x \subset G_x = G_x^0$, it suffices to prove ii) for H_x, G_x . Since $\pi_1(X, x)$ acts on the free \mathbf{Z} -module $T^{m,n}(\underline{V}_{\mathbf{Z}, x})/\text{torsion}$, any action of $\pi_1(X, x)$ on a line inside $T^{m,n}(\underline{V}_{\mathbf{Q}, x})$ must factor through $\{\pm 1\}$ (the only possible eigenvalues). Thus the connected group H_x has only trivial rational character. Hence it suffices to prove that for any $V' = (T^{m,n}(\underline{V}_{\mathbf{Z}}), T^{m,n}(W.), T^{m,n}(F'))$, the fixed part of $V'_{\mathbf{Q}, x}$ under H_x is the rational space $\omega_{\underline{V}_{\mathbf{Z}, x}}(V'')$ of a subobject V'' of V' in $\langle \underline{V}_{\mathbf{Z}, x} \rangle$ (notations of § 2). Replacing X by the finite covering defined by the maximal subgroup (of finite index) of $\pi_1(X, x)$ which factors through the connected component H_x of the monodromy group, we are reduced to prove that the largest

constant sub-local system of $V_{\mathbb{Z}}'$ is a (constant) sub-variation of M.H.S.. For a finite direct sum of polarizable variation of pure H.S., this is precisely the theorem of the fixed part of Griffiths-Schmid, see [3][10]. For a general polarizable good variation of M.H.S. in Steenbrinck-Zucker' sense, this is the theorem of the fixed part of these authors, see [12] 4.19. In fact, in loc. cit., this theorem is stated for a one-dimensional basis X , but we can reduce to this case by considering curves in X ; see [7] § 4.3.4.0, for the detailed argument.

So far we have proved that $H_X \triangleleft G_X$; to show that $H_X \triangleleft DG_X$, it remains to prove that $\mathcal{D}H_X = H_X$. We know that $H_X^{ab} \subseteq G_X^{ab}$ is a torus (lemma 3). Let χ a complex character of H_X . We just proved that $H_X|_{\mathbb{C}} \triangleleft G_X|_{\mathbb{C}}$, so that according to i) \Rightarrow ii) in lemma 1 for $K = \mathbb{C}$, $(T^{m,n}V_{\mathbb{C},X})^{\chi} + (T^{m,n}V_{\mathbb{C},X})^{\bar{\chi}}$ is stable under $G_X|_{\mathbb{C}}$; it is even the complexification of a real space $W_{\mathbb{R}}$ stable under $G_X|_{\mathbb{R}}$. Thus for some suitable Tate twist $\mathbb{R}(n)$, $\text{Det } W_{\mathbb{R}} \otimes \mathbb{R}(n)$ is a trivial $G_X|_{\mathbb{R}}$ -module. It follows that $\text{Det } W_{\mathbb{R}}$ is a trivial $H_X|_{\mathbb{R}}$ -module, which yields the equality: $|\chi| = 1$. Therefore all representations of H_X^{ab} are unitary; this means that $H_X^{ab}|_{\mathbb{R}}$ is a compact torus. Let $V' \subset \bigoplus T^{m_i, n_i} V_{\mathbb{Q}, X}$ a faithful representation of H_X^{ab} . A subgroup of finite index of $\pi_1(X, x)$ acts on V' through $\text{GL}(V' \cap \bigoplus T^{m_i, n_i} V_{\mathbb{Z}, X})$ which is discrete, and also through a compact torus. Because of the connectedness of H_X , it follows that V' is a trivial H_X -module, that is: $H_X = \mathcal{D}H_X$.

□

Corollary 1 (see [4; 4.2.6-9]). The local system $V_{\mathbb{Q}}$ underlying a polarizable variation of pure Hodge structure is semi-simple; each isotypical component carries a sub-variation of pure Hodge structure; the center of $\text{End}(V_{\mathbb{Q}})$ is purely of type $(0,0)$. For any $x \in X$, the connected monodromy group H_x is semi-simple.

Proof: since $H_x \triangleleft \mathcal{D}G_x$ for $x \in \overset{\circ}{X}$, and since $\mathcal{D}G_x$ is a semi-simple group (lemma 2), it follows that H_x is semi-simple; since H_x is locally constant on X , H_x is in fact semi-simple for any $x \in X$. This implies the complete reducibility of the action of $\pi_1(X,x)$ on $V_{\mathbb{Q},x}$ and the first assertion follows (the normality $H_x \triangleleft G_x$ would suffice here). By i) \Rightarrow iii) in lemma 1, applied to $H_x \triangleleft G_x$ for $x \in \overset{\circ}{X}$, we get on each stalk of each isotypical component of the local system $V_{\mathbb{Q}}|_{\overset{\circ}{X}}$ a Hodge sub-structure. By continuity, these Hodge sub-structures extend across $X \setminus \overset{\circ}{X}$ and patch together to give rise to a sub-variation of \mathbb{Q} -Hodge structure on the isotypical component of $V_{\mathbb{Q}}$. The third assertion follows from lemma 1 in the same manner.

□

Corollary 2 (see [4; 4.2.9b]). The radical of the connected monodromy group H_x associated to a polarizable variation of M.H.S. is unipotent.

Proof: let P_x be the subgroup of $GL(V_{\mathbb{Q},x})$ which respects the weight filtration W' , and N_x the subgroup of P which acts trivially on $\text{Gr}^W(V_{\mathbb{Q},x})$. Then $H_x \subset P_x$ and N_x is unipotent.

Moreover the connected monodromy group, say GrH_x , of $\text{Gr}^W(\underline{V}_x)$ at x is the image of H_x in P_x/N_x . Hence H_x is an extension of GrH_x , which (according to the previous corollary) satisfies $\text{GrH}_x = \mathcal{O} \text{GrH}_x$, by a (necessarily unipotent) subgroup of N_x .

□

Remark: a similar argument shows that the Mumford-Tate group of a M.H.S. over $R \subset \mathbb{R}$, say V , is an extension of the Mumford-Tate group of the direct sum of K-Hodge structures $\text{Gr}^W V$, by a unipotent K-group (notations of § 2). In particular if V is polarizable, $G(\text{Gr}^W V)$ is the quotient of $G(V)$ by its unipotent radical.

Remark: corollary 2 shows in particular that if G_x is solvable for some $x \in \overset{\circ}{X}$, then the variation of M.H.S. is unipotent in the sense of [15].

Remark: theorem 1 applies to the geometric situations considered in § 4 since in the course of proving it, we have made a restriction to curves.

Counterexample: we produce an example, following Deligne-Steenbrinck-Zucker (see [12] 3.16), to show that some extra hypothesis upon the variation of M.H.S. is necessary.

Consider the smooth 1-motive $\underline{M} = [\underline{\mathbb{Z}} \xrightarrow{n \mapsto x^n} \mathbb{G}_m]$ over $X = \mathbb{G}_m$. Here the set $\overset{\circ}{X}$ is $\mathbb{C}^* \setminus \mathbb{C}_{\text{tors}}^*$. The corresponding good variation of M.H.S. \underline{V} is an extension of $\underline{\mathbb{Z}}$ by $\underline{\mathbb{Z}}(1)$ inside $\underline{\mathbb{Q}}$. We

denote by ε_{-2} the generator $+i$ of $\underline{z}(1) \simeq W_{-2}$ and by ε_0 any element of $\underline{V}_{\underline{z}} \setminus W_{-2}$; then $\langle \varepsilon_0, \varepsilon_{-2} \rangle$ spans $\underline{V}_{\underline{z}}$. For some suitable determination of $\log x$ (depending on the choice of ε_0), the section $\tilde{\varepsilon}_0 := \varepsilon_0 - \frac{\log x}{2i\pi} \varepsilon_{-2}$ of $V_{\mathbb{C}}^{\mathbb{C}}$ spans F^0 and extends to a section of $\tilde{V}_{\mathbb{C}}^{\mathbb{C}}$ over \mathbb{P}^1 . We now combine notations from § 3 and § 4. For $x \in \dot{X}$, we have $U(H_x(\underline{M})) = U(G_x(\underline{M})) = \tilde{U} \simeq \mathbb{G}_a$ according to proposition 1. On the other side $H_x(\underline{M}) = U(H_x(\underline{M}))$ according to the previous corollary.

For any entire function f , let us now consider the following perturbation \underline{V}^f of \underline{V} : $(\underline{V}_{\underline{z}}^f, W_{\cdot}^f) = (\underline{V}_{\underline{z}}; W_{\cdot})$ but $(F^f)^0$ is spanned by $\tilde{\varepsilon}_0 + f \varepsilon_{-2}$. The corresponding groups $H_x(\underline{M}^f)$, $G_x(\underline{M}^f)$ admit the same description. The following assertions are easily seen to be equivalent:

- a) \underline{V}^f is good
- b) f extends analytically at ∞
- c) f is constant
- d) $\underline{V}^f \simeq \underline{V}$
- e) $V' := \text{Hom}(\underline{V}, \underline{V}^f)$ is good.

The group $H_x(\underline{V}')$ is isomorphic to \mathbb{G}_a ; viewing it as a subgroup of $GL_2 \times GL_2$ acting on $(V_{\mathbb{Q},x}^{\vee} \otimes V_{\mathbb{Q},x}^f)$, its "typical" element is of the shape

$$\begin{pmatrix} 1 & 0 \\ -a & 1 \end{pmatrix} \times \begin{pmatrix} 1 & +a \\ 0 & 1 \end{pmatrix}.$$

The "typical" element of $G_x(\underline{V}')$ is of the shape

$$\begin{pmatrix} 1/b & 0 \\ c & 1 \end{pmatrix} \times \begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix}, \text{ } a \text{ being independent of } c \text{ if}$$

(and only if) $\underline{V}^f \neq \underline{V}$. Therefore we see in this example that $H_x(\underline{V}') \triangleleft \mathcal{D}G_x(\underline{V}')$ if and only if \underline{V}' is good.

6. Maximality

Let $(\underline{V}_Z, W, F^\bullet)$ a polarizable good variation of mixed Hodge structures on X . Let $x \in \overset{\circ}{X}$ as in lemma 3. By the theorem, we know that $H_x \subset \mathcal{D}G_x$. We now study how big can H_x be in $\mathcal{D}G_x$.

Proposition 2. Assume that for some $y \in X$, G_y is nilpotent. Then for any $x \in \overset{\circ}{X}$, $H_x = \mathcal{D}G_x$.

Proof: according to the remark which follows lemma 3, G_y is actually a torus. Since the assertion is invariant under taking finite coverings of X , it suffices to show that any tensor $l \in T^{m,n}_{\mathbb{V}_{\mathbb{Q},x}}$ invariant under $\pi_1(X,x)$ spans a G_x -module W_x on which the action of G_x is abelian. It follows from the "fixed part" theorem that W_x is fixed by $\pi_1(X,x)$, and the local constancy of G_x on $\overset{\circ}{X}$, together with an argument of continuity, shows that W_x extends to a constant sub-variation of M.H.S., say $(\underline{V}', W', F'')$, of $(T^{m,n}_{\mathbb{V}_{\mathbb{Q}}}, T^{m,n}_h)$. In particular the action of G_x on $\underline{V}'_x = V'_x$ is the same as the action of G_y on \underline{V}'_y , which is abelian.

□

Before turning to applications, we propose a conjecture:

conjecture: under the general assumptions of this paragraph, assume moreover that, except possibly for one Tate twist, no Jordan-Hölder constituent of $(\underline{V}_Z, W, F^\bullet)$ is a locally constant variation of M.H.S.. Then $H_x = \mathcal{D}G_x$, for any $x \in \overset{\circ}{X}$.

Remark: this is obviously true, taking into account the normality property, if for some $x \in \overset{\circ}{X}$, the \mathbb{Q} -group $\mathcal{D}G_x$ is simple. By way of example, we consider a polarized (analytic) family of Abelian varieties with many endomorphisms over a complex algebraic base X ; by this, we mean that the generic fibre f_η of f (this makes sense since f is automatically algebraic) enjoys the following property: $(\text{End } f_\eta) \otimes_{\mathbb{Z}} \mathbb{Q}$ is a division ring which contains a commutative field of degree $\dim f_\eta$ over \mathbb{Q} . Then the derived Mumford-Tate group of the stalk $(R_1 f_* \mathbb{Q})_x$ can be computed for any Weil generic point x of X (so that $\text{End } f_\eta = \text{End } f_x$): it turns out that $\mathcal{D}G_x \cong \text{Res}_{Z^+/\mathbb{Q}} G$, where Z^+ denotes the maximal totally real subfield of the center of $(\text{End } f_\eta) \otimes_{\mathbb{Z}} \mathbb{Q}$, and G is an absolutely simple group over Z^+ (in fact $G|_{\mathbb{C}} \cong \text{SL}_2$); thus in this case $\mathcal{D}G_x$ is simple over \mathbb{Q} (see also the appendix, and [9, lemma 2.3], [1, th. 2]). However, P. Deligne has constructed some Abelian varieties whose derived Mumford-Tate groups $\mathcal{D}G$ are not simple over \mathbb{Q} (letter to the author).

Remark: in the case when G is solvable, the conjecture holds and is compatible with the results of [15]. In the reductive case, when the variation arises from geometry, the conjecture seems to be compatible with Grothendieck's conjecture about algebraic independence of periods of algebraic integrals, see [1].

7. Algebraic independence of Abelian integrals

Let \underline{M} be a smooth 1-motive over X ; its generic fibre $M := \underline{M}_\eta$ is a 1-motive over the function field $\mathbb{C}(X)$.

According to [4] III 10.1.7, there exists a universal extension M^\sharp of M by a vector group:

$$\begin{array}{ccccccc}
 & & 0 & \longrightarrow & X & = & X \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \text{Ext}^1(M, \mathbb{G}_a)^* & \longrightarrow & E^\sharp & \longrightarrow & E \longrightarrow 0
 \end{array}$$

The De Rham cohomological realization of M is by definition $H_{\text{DR}}^1(M) := \text{Co Lie } E^\sharp$. Moreover, the exact sequence

$$0 \longrightarrow \text{Hom}(X, \mathbb{G}_a) \longrightarrow \text{Ext}^1(M, \mathbb{G}_a) \longrightarrow \text{Ext}^1(E, \mathbb{G}_a) \longrightarrow 0$$

induces an exact sequence

$$(*) \quad 0 \longrightarrow \text{Hom}(X, \mathbb{G}_a) \longrightarrow H_{\text{DR}}^1(M) \longrightarrow H_{\text{DR}}^1(E) \longrightarrow 0,$$

where $H_{\text{DR}}^1(E)$ is the De Rham cohomological realization of the 1-motive $[0 \rightarrow E]$, identified with the usual first De Rham algebraic cohomology group of E .

Let K_x denote the fraction field of the local ring $\mathcal{O}_{X^{\text{an}}, x}$ at some $x \in X$. Construction [4] III 10.1.8 then yields a canonical isomorphism:

$$\text{Hom}_{\mathbb{C}(X)}(H_{\text{DR}}^1(M), K_x) = V_{\mathbb{C}, x} \otimes_{\mathbb{C}} K_x.$$

Let ∇^{an} the flat connection over $V_{\mathbb{C}}^{\mathbb{C}}$ such that $(V_{\mathbb{C}}^{\mathbb{C}})^{\nabla^{\text{an}}} = \underline{V}_{\mathbb{C}}$. According to Griffiths, $\text{Gr}^W \nabla^{\text{an}}$ has only regular singular points (see [3]). It follows that ∇^{an} itself has only regular singular points, henceforth is induced by a connection ∇ over $H_{\text{DR}}^1(M)$. In fact (*) is a sequence in the category of $\mathbb{C}(X)$ -vector spaces with $\mathbb{C}(X)/\mathbb{C}$ -connection, inducing the Gauss-Manin connection on $H_{\text{DR}}^1(E)$, and a trivial connection on $\text{Hom}(X, \mathbb{C}_a)$.

By definition of ∇ , we have

$$(**) \quad \text{Hom}_{\nabla}(H_{\text{DR}}^1(M), K_X) = V_{\mathbb{C}, X} \text{ inside } V_{\mathbb{C}, X} \otimes K_X.$$

Let us translate (*) and (**) in more down-to-earth terms, assuming that $X \xrightarrow{u} E$ is injective, and that X is constant over X . Then X may be considered as a group of sections of $\underline{E} \xrightarrow{f} X$, and \underline{V}_Z is spanned by $\langle \log_{\underline{E}X}, \text{Ker exp}_{\underline{E}} \rangle$, at least if we restrict ourselves to the subset of X where u is fibrewise injective. By means of suitable bases, a fundamental solution matrix of a Picard-Fuchs differential system of order one associated to $H_{\text{DR}}^1(M)$ can be expressed in some neighbourhood of $x_0 \in X$ by:

$$Z := \left(\begin{array}{c|c} \begin{array}{c} \phi \\ \gamma_j \\ \omega_i \end{array} & \begin{array}{c} \int_0^{\xi_k} \omega_i \\ 0 \end{array} \\ \hline \begin{array}{c} 0 \\ \text{Id} \end{array} & \end{array} \right)$$

where ω_i (resp. γ_j , resp. ξ_k) runs through some basis of $H_{\text{DR}}^1(\underline{E}/X) \otimes \mathcal{O}_{X, x_0}$ (resp. of $(\mathbb{R}_1 f_*^{\text{an}} \mathbb{C})_{x_0}$, of \underline{X}_{x_0}), so

that the entries of Z are elements of $\mathcal{O}_{X^{\text{an}}, x_0}$. In the first quadrant, we can recognize the classical

"period matrix" solution of a Picard-Fuchs differential system associated to the quotient $H_{\text{DR}}^1(E)$; such a matrix Z was already

considered by Y. Manin [17].

Our next theorem deals with a smooth 1-motive of the form

$$[0 \rightarrow \underline{E}] .$$

Theorem 2. Assume that some fibre of $\underline{E} \xrightarrow{f} X$ is split:

$$E_{x_1} = T_{x_1} \times A_{x_1} , \text{ and that } A_{x_1} \text{ is of CM type.}$$

Then the transcendence degree of the $\mathbb{C}(X)$ -extension generated by all the "periods" $\phi_{\gamma_i} \omega_i$ (γ_j, ω_i as above) equals the dimension of the "generic" derived Mumford-Tate group $\mathcal{D}G$.

Proof: by "generic", we mean the dimension δ of $\mathcal{D}(G_x(\underline{V}([0 \rightarrow \underline{E}])))$ for any $x \in \overset{\circ}{X}$. The groupe G_{x_1} is a torus, according to the CM type hypothesis. Since the variation of M.H.S. is good (at least when restricted to curves, see the example at the end of § 4), proposition 2 applies to establish the equality $\delta = \dim H_x$. Since the connection has only regular singular points, we get furthermore that δ is the dimension of the differential Galois group associated to $H_{DR}^1(E)$. But differential Galois theory tells us that this dimension is the transcendence degree of the $\mathbb{C}(X)$ -extension generated by the entries of the fundamental solution matrix Z (see [1], [2]).

□

Our last theorem is concerned with a smooth 1-motive of the form

$$[\underline{X} \xrightarrow{u} \underline{A}] , \text{ where } \underline{A} \xrightarrow{f} X \text{ is an Abelian scheme.}$$

Theorem 3. Assume that, over any finite etale covering of X , the map induced by $u : \underline{X} \rightarrow \underline{A}/\text{fixed part}$ remains injective. Then the transcendence degree of the $\mathbb{C}(X)((\phi_{\gamma_j} \omega_i)_{ij})$ - extension generated

by the germs of analytic functions $\int_0^{\xi_k} \omega_i$ (ξ_k as above), equals the dimension of the generic group \tilde{U} introduced in § 3.

Proof: using similar arguments from differential Galois theory, we can see that it is enough to show that

$$\tilde{U} \simeq \text{Ker}(H_x(\underline{V}[\underline{X} \rightarrow \underline{A}]) \longrightarrow H_x(\underline{V}[0 \rightarrow \underline{A}])) := U(H_x) .$$

According to theorem 1, we have $H_x \triangleleft G_x$; thus in order to apply proposition 1, it suffices to show that $W_{0,x}^{H_x} \subseteq W_{-1,x}$. At the cost of replacing X by a finite étale covering, we may assume that H_x is the whole monodromy group (not only its neutral component). We identify \underline{X} with its image in \underline{A} and consider it as a group of sections of f . Let $v_x \in W_{0,x}^{H_x}$; it extends to a global section v of \underline{W}_0 ; setting $\xi = \exp v \in \underline{X}$, we thus have $\nabla(d/dx) \int_0^{\xi(x)} \omega_x = 0$, for any section ω of $\Omega_{\underline{A}/X}^1$ and any derivation d/dx of $\mathbb{C}(X)$. According to Manin [17], this implies that ξ belongs to the fixed part of \underline{A} . However the hypothesis we have made upon u implies in turn that $\xi = 0$, so that $v \in \Gamma \underline{W}_{-1}$.

□

Remark: this result is the geometric variant of the "Kummer theory" on Abelian varieties, which studies the extension of the field of rationality of some torsion points, generated by the division points of some non-torsion points.

Remark: the exact sequence (*) of $\mathbb{C}(X)$ -vector spaces with connection splits if and only if $U(H_x) = 0$.

Appendix

Automorphisms of certain Hodge structure over number fields

So far we have been concerned only with polarized Hodge structures $(H_{\mathbf{Z}}, h, \psi)$ over \mathbf{Z} , and we used some variants of the argument that the automorphisms of $(H_{\mathbf{Z}}, h, \psi)$ form a finite group, say G : indeed G imbeds both into the discrete group $GL(H_{\mathbf{Z}})$ and into the compact orthogonal group $O_{\psi} = \text{Aut}(H_{\mathbf{Z}} \otimes \mathbb{R}, \psi(\cdot, h(i)\cdot))$. If \mathbf{Z} is replaced by the ring of integers R of some totally real number field, the group $GL(H_R)$ is no longer discrete in general; even if one tries to use Weil's restriction of scalars from R to \mathbf{Z} , it could happen that the "conjugates" of O_{ψ} are not compact. Here we shall study those polarized Hodge structures over R which arise naturally as pieces of the cohomology of Abelian varieties with many endomorphisms, and show how the finiteness of G involves arithmetical questions.

8. Classification of Abelian varieties with many endomorphisms

Let X a complex simple Abelian variety of dimension $g > 0$, such that $D = \text{End } X \otimes_{\mathbb{Z}} \mathbb{Q}$ contains some commutative field E of degree g over \mathbb{Q} . Since X is simple, D is a division ring whose center is denoted by Z . Any polarization ψ of X defines a positive involution $*$ over D ; this implies that the subfield Z^+ of Z fixed by $*$ is a totally real number field. After Albert's classification (cf [8] 11), four cases can occur a priori:

Type I: $Z^+ = Z = E = D$; X is then called a "Hilbert-Blumenthal" Abelian variety.

Type II: $Z^+ = Z$ and for every real place ρ of Z ,


$$D \otimes_{Z, \rho} \mathbb{R} \cong M_2(\mathbb{R}) .$$

According to [8] loc. cit., there exists $a \in D$, such that the reduced trace $\text{Tr}_{D/Z}(a)$ vanishes, and such that the involution $*$ is given by $x^* = a[\text{Tr}_{D/Z}(x) - x]a^{-1}$ for any $x \in D$. Since D is a quaternion algebra over Z , there exists $b \in D$, such that the reduced trace $\text{Tr}_{D/Z}(b)$ vanishes, and which anticommutes with a . We then have $b^* = b$. So $Z(b)$ is totally real and one can assume that $E = Z(b)$.

Type III: $Z^+ = Z$ and for every place ρ of Z , $D_{Z, \rho} \otimes \mathbb{R}$ is isomorphic to the Hamilton quaternion algebra \mathbb{H} . In fact this case does not occur under our assumptions on X . Indeed the representation of $\text{End}_{\mathbb{H}}[H_1(X^{\text{an}}, \mathbb{R}) \otimes_{Z, \rho} \mathbb{R}]$

over $H_1(X^{\text{an}}, \mathbb{R}) \otimes_{Z, \rho} \mathbb{R}$ yields, after complexification, two copies of the standard representation of SO_2 ([9, lemma 2.3]). This representation thus decomposes into four sub-representations of degree one, whose endomorphism algebra has to be $\mathbb{H} \otimes_{\mathbb{R}} \mathbb{C} \cong M_2(\mathbb{C})$: this is impossible.

Type IV: Z is a totally imaginary quadratic extension of Z^+ . Either $[Z:\mathbb{Q}] = 2g$ in which case X is said of "CM type" and we can choose $E = Z^+$, or $[Z:\mathbb{Q}] = g$ and we can assume that E is a totally imaginary quadratic extension of its subfield E^+ fixed by $*$, whence the following diagram of extensions:



since $[D:\mathbb{Q}] \leq 2g$, $[E:Z] \leq [D:E]$ (from the commutativity of E), and $[E:\mathbb{Q}] = g$, we find that $[E:Z] \leq 2$.

Except in the CM case, E is a maximal commutative subfield of D , and in any case we shall write E^+ for the subfield of E fixed by $*$, K for the Galois closure of E^+ in \mathbb{R} , and \mathcal{R} for the ring of integers of K .

9. The Hodge structures H_μ over R

Let us pick some primitive element ζ of E^+ over \mathbb{Q} in the order $(\text{End } X) \cap E^+$ of E^+ . This element acts via ζ^* on the free R -module $H^1(X^{\text{an}}, R)$, and its characteristic polynomial has rational integral coefficients and the same roots as the minimal polynomial of ζ ; that characteristic polynomial thus equals some power of this (separable) minimal polynomial, so that some essential R -submodule of $H^1(X^{\text{an}}, R)$ decomposes into a direct sum of free R -modules H_μ , the indices running among the imbeddings of E^+ into K . Let L be the compositum in \mathbb{C} of K and the image of E through some complex imbedding, so that $L = K$ except in the non-CM type IV case. Then the rank of H_μ is $2g/[E:\mathbb{Q}] = 2[L:K]$. The free R -module H_μ is naturally endowed with a structure of polarized Hodge structure (H_μ, h_μ, ψ_μ) of type $(0,1) + (1,0)$ over R , and there is an isomorphism of polarized K -Hodge structures $(H^1(X^{\text{an}}, K), h, \psi) = \bigoplus_{\mu: E^+ \rightarrow K} (H_\mu \otimes_R K, h_\mu, \psi_\mu)$. Furthermore when $L \neq K$, ψ_μ comes from a L -hermitian form φ_μ on the L -vector space $H_\mu \otimes_R K$.

10. Automorphisms of (H_μ, h_μ, ψ_μ)

Proposition 3. The group G of L -linear automorphisms of (H_μ, h_μ, ψ_μ) is infinite if and only if one of the following statements holds:

- i) $K = L$, and there exists some non-totally positive element $k \in K^X$ such that the multiple $\sqrt{k.C}$ of the Weil morphism $C = h_\mu(\sqrt{-1})$ on $H_\mu \otimes_R \mathbb{R}$ comes from an endomorphism of $H_\mu \otimes_R K$,
- ii) $K \neq L$ and the direct summand $(H_\mu \otimes_R K) \otimes_L \mathbb{C}$ of $H_\mu \otimes_R \mathbb{C}$ is bihomogeneous.

We begin the proof with the case $K = L$.

Let us choose a R -basis of H_μ such that the Riemann form $\psi_\mu = \langle \cdot, \cdot \rangle$ is represented by the matrix $\begin{pmatrix} 0 & e \\ -e & 0 \end{pmatrix}$ for some $e \in R^X$, and let us consider the matrix of C in the basis (viewed as a basis of $H_\mu \otimes_R \mathbb{R}$): since $C^2 = -1$, this matrix has the shape $\begin{pmatrix} -\beta & -\gamma \\ \alpha & \beta \end{pmatrix}$, for $(\alpha, \beta, \gamma) \in \mathbb{R}^3$ satisfying the equation $\alpha\gamma = 1 + \beta^2$. It follows that $\alpha\gamma \neq 0$. The symmetric form $\langle \cdot, C(\cdot) \rangle$ is represented by $Q = \begin{pmatrix} \alpha e & \beta e \\ \beta e & \gamma e \end{pmatrix}$. Let $\theta \in G$, so that $\theta \in \text{Aut } H_\mu \cap O(H_\mu \otimes \mathbb{R}, Q)$, and let us write $\theta_{ij} \in R$ for the coefficients of the matrix of θ . The equation ${}^t\theta Q \theta = Q$ is equivalent to the system

$$(\Sigma) \begin{cases} \alpha(\theta_{11}^2 - 1) + 2\beta \theta_{11}\theta_{21} + \gamma\theta_{21}^2 = 0 \\ \alpha\theta_{11}\theta_{12} + \beta(\theta_{12}\theta_{21} + \theta_{11}\theta_{22} - 1) + \gamma\theta_{21}\theta_{22} = 0 \\ \alpha\theta_{12}^2 + 2\beta \theta_{12}\theta_{22} + \gamma(\theta_{22}^2 - 1) = 0 . \end{cases}$$

a) Let us first deal with the case when C is defined over some totally real algebraic extension of K . Then α, β, γ are totally real algebraic numbers. Let $\sigma \in \text{Gal}(K/\mathbb{Q})$, and let $\alpha^\sigma, \beta^\sigma, \gamma^\sigma$ be conjugates (necessarily real) of α, β, γ respectively, above σ . Setting $Q^\sigma = \begin{pmatrix} \alpha^\sigma e^\sigma & \beta^\sigma e^\sigma \\ \beta^\sigma e^\sigma & \gamma^\sigma e^\sigma \end{pmatrix}$, we find ${}^t \theta^\sigma Q^\sigma \theta^\sigma = Q^\sigma$, and $\det Q^\sigma = (e^\sigma)^2 > 0$, so that θ^σ belongs to the compact orthogonal group $O_2(Q^\sigma)$. By restriction of scalars à la Weil from K to \mathbb{Q} , G imbeds into $(\text{Res}_{K/\mathbb{Q}} \text{Aut}(H_\mu \otimes_{\mathbb{R}} K))(\mathbb{Z})$ (which is discrete) and into $\prod_{\sigma} O_2(Q^\sigma)$ (which is compact), so that G is finite in this case.

Here we point out that the CM type is a special case: indeed the Hodge bigraduation of $H_\mu \otimes_{\mathbb{R}} \mathbb{C}$ comes from the CM decomposition $H_\mu \otimes_{\mathbb{R}} L' = [H^1(X^{\text{an}}, \mathbb{Q}) \otimes_{Z, \nu} L'] \oplus [H^1(X^{\text{an}}, \mathbb{Q}) \otimes_{Z, \bar{\nu}} L']$, for some complex place ν of Z over μ (here we denote by L' the compositum $K \cdot \nu(Z)$ which is a quadratic totally imaginary extension of K). Let us write $L' = K(h)$ with $h^2 = -g \in \mathbb{R}^-$; the matrix of C (in some basis adapted to the above decomposition) reads $\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$, thus C is defined over the totally real number field $K(\sqrt{g}) = K(ih)$.

- b) Let us now assume that α, β, γ span a line over K ; since $\alpha\gamma \neq 0$, we write $\beta = b\alpha$, $\gamma = c\alpha$, for some $(b, c) \in K \times K^{\times}$. This yields $\alpha^2 = \frac{1}{a-b^2} \in K \cap \mathbb{R}^+$. Getting rid of the above possibility 2), we are reduced to the case i) of the proposition, with $k = c-b^2$. Since any $\theta \in G$ commutes with $\frac{1}{\alpha}c = \begin{pmatrix} -1 & -c \\ 1 & b \end{pmatrix}$, θ has the shape $\begin{pmatrix} x & -cy \\ y & x+2by \end{pmatrix}$ for x, y, cy and $2by \in \mathbb{R}$. The set of all these matrices is an order R' in the field $K' = K(\sqrt{b^2-c}) = K(i\alpha)$, as is seen by identifying $\begin{pmatrix} x & -cy \\ y & x+2by \end{pmatrix}$ with $(x+by) + y\sqrt{b^2-c}$. Since θ is invertible, it is identified with some unit in R' . The equation ${}^+\theta Q \theta = Q$ then reads $x^2 + 2bxy + cy^2 = 1$, that is $(x+by) + y\sqrt{b^2-c} \in \text{Ker } N_{K'/K}$. But $N_{K'/K}$ has maximal rank as a morphism between unit groups $(R')^{\times} \rightarrow R^{\times}$. By assumption, K' is not totally imaginary, so that by Dirichlet's theorem $\text{rk}(R')^{\times} > \text{rk } R^{\times}$. Thus the kernel of $N_{K'/K}$ in $(R')^{\times}$ contains infinitely many elements, and so does G in this case.

- c) It remains to deal with the case when α, β, γ span a K -vector space of dimension at least 2. This implies that all minors of (Σ) vanish. In particular,

$$(1) \quad (\theta_{11}^2 - 1)(\theta_{12}\theta_{21} + \theta_{11}\theta_{22} - 1) = 2\theta_{11}^2\theta_{12}\theta_{21}$$

$$(2) \quad (\theta_{22}^2 - 1)(\theta_{12}\theta_{21} + \theta_{11}\theta_{22} - 1) = 2\theta_{12}\theta_{22}\theta_{21}^2$$

$$(3) \quad (\theta_{22}^2 - 1)(\theta_{22}^2 - 1) = \theta_{12}^2\theta_{21}^2$$

from which it follows that $(\theta_{12}\theta_{21} + \theta_{11}\theta_{22}^{-1})\theta_{12}^2\theta_{21}^2 = 2\theta_{11}\theta_{12}^2\theta_{21}^2\theta_{22}$, so that $\theta_{12}\theta_{21} = 1 + \theta_{22}\theta_{11}$ if $\theta_{12}\theta_{21} \neq 0$. Squaring, we find (using (3) again) that $\theta_{11} = -\theta_{22}$ in this case, and from (1) we get $\theta_{12}\theta_{21} = 1 - \theta_{11}^2$; that is, $\det \theta = -1$ and $\text{tr } \theta = 0$, from which it follows that $\theta^2 = 1$. If $\theta_{12}\theta_{21} = 0$, we get from the vanishing of the other minors) that $\theta_{11}^2 = \theta_{22}^2 = 1$, and moreover that $\theta_{11}\theta_{22} = -1$ if θ_{12} and θ_{21} do not vanish simultaneously; so we are reduced to the previous case where $\theta_{11} = -\theta_{22}$, except if $\theta = \pm 1$. From this description we see that any two elements of G , distinct from ± 1 , are inverse up to sign; this implies that G is finite (with at most 4 elements).

We now turn to the case $K \neq L$.

Let us choose a R -basis of H_μ such that the L -hermitian form $\varphi_\mu = \langle \cdot, \cdot \rangle$ is represented by the matrix $\begin{pmatrix} e & 0 \\ 0 & f \end{pmatrix}$, for some $(e, f) \in (R^\times)^2$. We identify $L \otimes_K \mathbb{R}$ with \mathbb{C} by means of an element h of L such that $h^2 = -g \in K \cap \mathbb{R}'$; since L is totally imaginary (like E), g is totally positive. The Weil morphism C is linear with respect to the complex structure induced by $L \otimes_K \mathbb{R}$ on $H_\mu \otimes_R \mathbb{R}$, since it commutes with L .

a) Let us first deal with the case when $(H_\mu \otimes_R K) \otimes_L \mathbb{C}$ is not bihomogeneous. Through the isomorphism $\mathbb{C} \cong L \otimes_K \mathbb{R}$, $(H_\mu \otimes_R K) \otimes_L \mathbb{C}$ can be identified with the complex 2-plan $H_\mu \otimes_R \mathbb{R}$, and C denotes the two eigenvalues $\pm i$ on

$H_\mu \otimes_{\mathbb{R}} \mathbb{R}$. Since ψ_μ is a morphism of the Hodge structure and since C is \mathbb{C} -linear, C belongs to the unitary group of φ_μ . Using this property, and the equations $C^2 = -1$ and $\text{tr } C = 0$, we get the following matrix representation for C : $\begin{pmatrix} ht & \gamma \\ -\alpha & -ht \end{pmatrix}$ for $t \in \mathbb{R}$, $(\alpha, \gamma) \in \mathbb{C}^2$, and with the following equation:

$$(*) \quad \alpha\gamma + gt^2 = 1 \quad \text{and} \quad f\bar{\alpha} = e\gamma.$$

Let us write $\alpha = v + hw$, for $(v, w) \in \mathbb{R}^2$. Taking into account (*), we find the following matrix representation for the symmetric form $\text{Re } h/g \langle \cdot, C(\cdot) \rangle$ in the real basis of $H_\mu \otimes_{\mathbb{R}} \mathbb{R}$ attached to the chosen complex basis:

$$Q_\mu = \begin{pmatrix} -et & 0 & fw & -fv \\ 0 & -get & fv & gfw \\ fw & fv & ft & 0 \\ -fv & gfw & 0 & gft \end{pmatrix} \quad \text{for } (t, v, w) \in \mathbb{R}^3.$$

Since Q_μ has maximal rank and index 0, the first main 1-minor is non zero: $t \neq 0$.

Let us first assume that $\alpha \neq 0$. Since $\theta \in G$ commutes with C , we find that θ has following matrix representation:

$$\begin{pmatrix} x & -\gamma y/\alpha \\ y & x+2hty/\alpha \end{pmatrix} = \begin{pmatrix} x & -f\bar{\alpha}y/\alpha \\ y & x+2hty/\alpha \end{pmatrix}, \quad \text{for } (x, y) \in \mathbb{L}^2.$$

Furthermore, the relation $t_{\theta^{-1}} \begin{pmatrix} e & 0 \\ 0 & f \end{pmatrix} \theta = \begin{pmatrix} e & 0 \\ 0 & f \end{pmatrix}$ yields the system

$$(\Sigma') \quad \begin{cases} x\bar{x} + f/e \cdot y\bar{y} = 1 & (1) \\ x\bar{x} + (f/e + 4gt^2/\alpha\bar{\alpha})y\bar{y} = 1 + 2ht/\alpha\bar{\alpha}(\alpha x\bar{y} + \bar{\alpha}x\bar{y}) & (2) \\ 2hty\bar{y} = \bar{\alpha}x\bar{y} - \alpha x\bar{y} & (3) \end{cases}$$

Eliminating $x\bar{x}$ between (1) and (2) and $y\bar{y}$ between (2) and (3), one obtains $\bar{x}y = 0$; reporting this equation in (1) and (3) gives $y = 0$ and $x\bar{x} = 1$. (Note that since θ is invertible, x is a unit in L).

If on the contrary $\alpha = 0$, then $\gamma = 0$ according to (*), so that θ is diagonal and $x\bar{x} = 1$ again. In both cases, to show that G is finite, it suffices to prove that the unit in $\text{Ker } N_{L/K}$ form a finite group. Since L is a totally imaginary quadratic extension of K , the unit groups U_L and U_K have the same rank $[K:\mathbb{Q}] - 1$, thus the desired statement comes from Dirichlet's theorem.

b) It remains to deal with the case ii) of the proposition. In this case C is the homothety with scale $\pm i \in L \otimes_K \mathbb{R}$ on $H_{\mu} \otimes_{\mathbb{R}} \mathbb{R}$. The matrix of the symmetric form $\text{Re } h \langle \cdot, C(\cdot) \rangle$ in the real basis of $H_{\mu} \otimes_{\mathbb{R}} \mathbb{R}$ attached to the chosen complex basis reads:

$$Q = \sqrt{g} \begin{pmatrix} e & & & \\ & \text{ge} & & \\ & & f & \\ & & & gf \end{pmatrix} .$$

Since Q has maximal rank and index 0, it follows from Sylvester's criterium that the product $\delta_1 \delta_3$ of the first and third main minors of Q is positive: $ef > 0$.

Let K' the imaginary quadratic session of K generated by $\sqrt{-e/f}$. We shall show that K' is not totally imaginary. Indeed, according to a result of Shimura [11, th 5], there exists at least one place σ_μ of K ($\sigma \in \text{Gal}(K/\mathbb{Q})$) such that H_{σ_μ} falls in case a). We apply Sylvester's criterium to the matrix

$$Q_{\sigma_\mu} = \begin{pmatrix} -e^\sigma t & 0 & f^\sigma w & -f^\sigma v \\ 0 & -g^\sigma e^\sigma t & f^\sigma v & g^\sigma f^\sigma w \\ f^\sigma w & f^\sigma v & f^\sigma t & 0 \\ -f^\sigma v & g^\sigma f^\sigma w & 0 & g^\sigma f^\sigma t \end{pmatrix} \quad \begin{array}{l} \text{considered in} \\ \text{case a) for} \\ H_{\sigma_\mu} . \end{array}$$

The product $\delta_1 \delta_3$ is $-(e^2 f)^\sigma t^2 (f^\sigma v^2 + f^\sigma g^\sigma w^2 + g^\sigma e^\sigma t^2)$. Because of the relations (*), this can be simplified: $\delta_1 \delta_3 = -(e^\sigma f^\sigma t)^2 e^\sigma f^\sigma$. We find $e^\sigma f^\sigma < 0$, so that K' is not totally imaginary.

Let $\theta \in G$ and δ its L-determinant. The relation $t_{\bar{\theta}} \begin{pmatrix} e & 0 \\ 0 & f \end{pmatrix} \theta = \begin{pmatrix} e & 0 \\ 0 & f \end{pmatrix}$ yields the shape $\begin{pmatrix} a & -f/e \bar{c} \delta \\ c & \bar{a} \delta \end{pmatrix}$ for the matrix of θ , with $\delta \bar{\delta} = 1$ and $ea\bar{a} + fc\bar{c} = e$. To show that G is infinite, it suffices to consider the case where $a, c \in R$ and $\delta = 1$. Then the set of matrices $\begin{pmatrix} a & -f/e c \\ c & a \end{pmatrix}$ with $(a, c) \in K^2$ is a field isomorphic to K' . The subring consisting of matrices with entries in R is an order R' , and the subgroup of $(R')^X$ consisting of unimodular matrices satisfying $ea^2 + fc^2 = e$ is the kernel of $N_{K'/K}$ in $(R')^X$. The same

argument as in part of the proof ($K = L$, case b), shows that this group is infinite. This completes the proof of the proposition.

□

Along the lines of [4; II 4.4.8], proposition 3 can be used to reprove the conjecture of § 6 for families of Abelian varieties with many endomorphisms. The point is that, except in case ii), the Hodge filtration of H_{μ} is locally constant if and only if the monodromy is finite. Indeed, the local constancy of F^{\bullet} implies that the monodromy group (whose neutral component is semi-simple) imbeds into the automorphism group \bar{G} which is finite except in cases i), ii) and which is a torus in case i); here \bar{G} denotes the Zariski closure of the group G determined by proposition 3.

References

- [1] ANDRE Y., "Sur certaines algèbres de Lie associées aux schémas abéliens", Note CR.A.S. t. 299 I n° 5 (1984), 137-140.

- [2] ANDRE Y., "Quatre descriptions des groupes de Galois différentiels", to appear in the proceedings of the "Séminaire d'algèbre de Paris", Springer L.N.

- [3] CORNALBA M., GRIFFITHS P., "Some transcendental aspects of algebraic geometry", Proc Symp pure Math. vol XXIX, AMS 1975, 3-110.

- [4] DELIGNE P., "Théorie de Hodge" II, Publ. Math. IHES 40 (1972), 5-57; III, Publ. Math. IHES 44 (1974) 5-78.

- [5] DELIGNE P., "La conjecture de Weil pour les surfaces K_3 ", Inv. Math. 15 (1972), 206-226.

- [6] DELIGNE P. and al., Hodge cycles, Motives and Shimura Varieties, Springer L.N: 900 (1982) I Hodge cycles on Abelian varieties 9-100; II Tannakian categories 101-228.

- [7] KATZ N., "Algebraic solutions of differential equations", Inv. Math. 18 (1972), 1-118.

- [8] MUMFORD D., Abelian Varieties, Oxford Univ. Press, Oxford (1970).

- [9] MURTY V.K., "Exceptional Hodge Classes on Certain Abelian varieties", Math. Ann. 268 (1984), 197-205.

- [10] SCHMID W., "The singularities of the period mapping", Inv. Math. 22 (1973), 211-319.

- [11] SHIMURA G., "On analytic families of polarized Abelian varieties and automorphic functions", Ann. of Math. 78 (1963), 149-192.

- [12] STEENBRINCK J., ZUCKER S., Variation of mixed Hodge structure I Inv. math. 80 (1985), 489-542.

- [13] WATERHOUSE W., Introduction to Affine Group Schemes, Springer, Heidelberg, 1979.

- [14] CATTANI E., KAPLAN A., SCHMID W., Degeneration of Hodge structure Ann. Math. 123 (1986) 457-535.

- [15] HAIN R.M., ZUCKER S., Unipotent variations of mixed Hodge structure, Inv. math. 88 (1987), 83-124.

- [16] KASHIWARA M., Variation of mixed Hodge structure (Preprint 1986).

- [17] MANIN Y., Algebraic curves over fields with differentiation, AMS Translations (2) 37, 59-78.