


**On relations of dimensions of automorphic forms  
of  $Sp(2, \mathbb{R})$  and its compact twist  $Sp(2)$  (I)**

**Tomoyoshi Ibukiyama**



**Department of Mathematics  
College of General Education  
Kyushu University  
Ropponmatsu, Fukuoka, 810 Japan**

**and**

**Sonderforschungsbereich 40/  
Max-Planck-Institut für Mathematik  
Gottfried-Claren-Straße 26  
D-5300 Bonn 3**

**SFB/MPI 84-9**

On relations of dimensions of automorphic forms  
of  $Sp(2, \mathbb{R})$  and its compact twist  $Sp(2)$  (I)

Tomoyoshi Ibukiyama

Let  $p$  be a fixed prime. In the previous paper [9], we have given some examples and conjectures on correspondence between automorphic forms of  $Sp(2, \mathbb{R})$  (size four) and  $Sp(2) = \{g \in H; g^t \bar{g} = 1_2\}$  ( $H$ : Hamilton quaternions) which preserves L-functions, where the  $p$ -adic closures of the discrete subgroups (to which automorphic forms belong) are minimal parahoric. This was an attempt to a generalization of Eichler's correspondence between  $SL_2(\mathbb{R})$  and  $SU(2)$ . Ihara raised such problem for symplectic groups and Langlands [15] has given a quite general philosophy on correspondence of automorphic forms of any reductive groups (functoriality with respect to L-groups). In this paper, we give good global dimensional relations of automorphic forms of  $Sp(2, \mathbb{R})$  and  $Sp(2)$ , when the  $p$ -adic closures of discrete subgroups in question are maximal compact. (As for similar results for other groups, see [8].) More precisely, put

$$K(p) = Sp(2, \mathbb{Q}) \cap \gamma M_4(\mathbb{Z}) \gamma^{-1}$$

$$= Sp(2, \mathbb{Q}) \cap \begin{pmatrix} * & * & 1/p & * \\ p^* & * & * & * \\ p^* & p^* & * & p^* \\ p^* & * & * & * \end{pmatrix}, \text{ where } \gamma = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & p & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

and  $*$  runs through all integers.

---

The author is partially supported by SFB 40, Univ. Bonn and Max-Planck-Institut für Mathematik.

For any  $\Gamma \subset Sp(2, R)$ , denote by  $A_k(\Gamma)$  (resp.  $S_k(\Gamma)$ ) the space of automorphic (resp. cusp) forms belonging to  $\Gamma$ . We shall calculate the dimension of  $S_k(K(p))$  for all primes  $p$  (Theorem 4 in § 4). By comparing these with those of certain automorphic forms (i.e., certain spherical functions) of  $Sp(2)$ , we shall show certain interesting relations of dimensions (Theorem 1 below). Some philosophical aspects of relations of orbital integrals have been explained in Langlands [14]. But except for the case of  $GL_n$ , or the product of its copies, as far as I know, this is the first global result concerning on the comparison of dimensions of spaces of automorphic forms belonging to different  $R$ -forms of a complex Lie group. We propose a precise conjecture on the correspondence of these spaces which is suggested by these relations (conj. 1.4). (Some examples of pairs of automorphic forms whose Euler  $\chi$ -factors fit this conjecture have been given in [9].) In a sense, the situation is fairly different from the case of  $GL_2$ , for example, it is noteworthy that, nevertheless the discrete subgroups in question are 'maximal', some 'old forms' come in these spaces. This is not because there exist some forms obtained by Saito-Kurokawa

lifting. To state the relation more explicitly, we need some more notations. Let  $B$  be the definite quaternion algebra with the prime discriminant  $p$ ,  $O$  be a maximal order of  $B$ . Put  $B_p = B \otimes_{\mathbb{Q}} \mathbb{Q}_p$  and  $O_p = O \otimes_{\mathbb{Z}} \mathbb{Z}_p$ . Put

$$G = \left\{ g \in M_2(B); g^t \bar{g} = n(g) 1_2, n(g) \in \mathbb{Q}_+^{\times} \right\} .$$

Let  $G_A$  be the adelization of  $G$ , and  $G_{\infty}$  (resp.  $G_q$ ) be the infinite (resp.  $q$ -adic) component of  $G_A$ . For any open subgroup  $U$  of  $G_A$ , denote by  $M_{\nu}(U)$  the space of automorphic forms on  $G_A$  belonging to  $U$  with 'weight  $\rho_{\nu}$ ', where  $\rho_{\nu}$  is the irreducible representation of  $Sp(2)$  which corresponds to the Young diagram

$$\begin{array}{|c|c|c|} \hline 1 & \dots & \nu \\ \hline 1 & \dots & \nu \\ \hline \end{array} . \quad (\text{cf. Ihara [13], Hashimoto [5]}) .$$

We take an open subgroup  $U_2 = G_{\infty} U_p^2 \prod_{q \neq p} U_q^1$  of  $G_A$ , where  $U_q^1 = GL_2(O_q) \cap G_q$ , and  $U_q^2$  is the unit group of the right order of a maximal left  $O_p$ -lattice in the non-principal genus in the quaternion hermitian space  $B_p^2$  with the metric  $n(x)+n(y)$  for  $(x,y) \in B_p^2$  where  $n(*)$  is the reduced norm of  $B$ . (cf. §1) Put

$$\Gamma_o(p) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z}); c \equiv 0 \pmod{p} \right\} .$$

Theorem 1      For each integer  $k \geq 5$  and each prime integer  $p$ , we have the following relation of the dimensions:

$$\dim S_k(K(p)) - 2 \dim S_k(\text{Sp}(2, Z))$$

$$= \dim M_{k-3}(U_2) - \dim A_2(\Gamma_0(p)) \times \dim S_{2k-2}(\text{SL}_2(Z)).$$


The conjectural meaning of this Theorem will be explained in §1. The dimension of  $S_k(\text{Sp}(2, Z))$  has been known by Igusa [12], and the dimension of  $M_{k-3}(U_2)$  has been given in [7] (II). So, only  $\dim S_k(K(p))$  is to be calculated. Recently, Hashimoto [6] obtained a general (but not explicit) formula of dimensions of cusp forms belonging to any discrete subgroups  $\Gamma$  of  $\text{Sp}(2, R)$ . Roughly spoken, his assertion is as follows: apparently, we have to calculate the contribution of each  $\Gamma$ -conjugacy class to the dimension, but at least for the semi-simple conjugacy classes, we can calculate everything from some data on integral property of their local conjugacy classes in  $\text{Sp}(2, Q_p)$  and  $\text{Sp}(2, R)$  (so, in these cases, we can avoid the classification of  $\Gamma$ -conjugacy classes), and besides, for all conjugacy classes, 'local data' at the infinite place can be explicitly written down. (As for the further details such as 'family', see his paper.) But in order to obtain the dimensions explicitly by using his formula, we must calculate such local data (the number of 'optimal embeddings' and some local masses) of semi-simple conjugacy classes, and classify  $K(p)$ -conjugacy classes of parabolic

type or some mixed type. (Since  $K(p)$  is not contained in  $Sp(2,2)$ , there was no known results on such classification.) These calculations are rather elaborate and have been done in somewhat lengthy case by case process similar to [7], and here, we shall often omit the proofs, or content ourselves with some sketchy proofs. (As for an expository review on results in [5], [6], [7] how to calculate dimensions in general, confer [8] §4.) In §2, we give local data of semi-simple conjugacy classes. In §3, we classify  $K(p)$ -conjugacy classes of parabolic or mixed type. In §4, we sum up them and prove Th.1.

The author would like to thank Dr. K.Hashimoto who has shown him the manuscript of his paper [6], and Dr.S.Kato who informed him the notion of the folding of the Dynkin diagrams of  $p$ -adic algebraic groups.

§1. Conjectural meaning of Theorem 1.

To explain the situation more clearly, we recall some local theory of  $p$ -adic algebraic groups (cf. Tits [18]). The extended Dynkin diagram for  $G_p$  can be obtained from the one for  $Sp(2, Q_p)$  by dividing by the non trivial graph automorphism  $\sigma$ , and each summit can be regarded as a double coset of a minimal parahoric subgroups. (See  $C_2$  and  ${}^2C_2$  in the table of [18] p.64)



These double cosets are explicitly given as follows: put

$$B(p) = \left\{ g \in Sp(2, Z); g \equiv \begin{pmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & 0 \\ 0 & 0 & * & * \end{pmatrix} \pmod{p} \right\} \quad (* : \text{integers})$$

and let  $B(p)_p$  be the  $p$ -adic closure of  $B(p)$ . Then,  $B(p)_p$  is an Iwahori subgroup of  $Sp(2, Q_p)$ . We can take

$$S_0 = B(p)_p w_0 B(p)_p, \quad S_1 = B(p)_p w_1 B(p)_p, \quad \text{and} \quad S_2 = B(p)_p w_2 B(p)_p,$$

$$\text{where } w_0 = \begin{pmatrix} 0 & 0 & -p^{-1} & 0 \\ 0 & 1 & 0 & 0 \\ p & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad w_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad \text{and}$$

$$w_2 = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

On the other hand, put

$$G_p^* = \left\{ g \in M_2(B_p) ; g \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}^t \bar{g} = n(g) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, n(g) \in Q_p^* \right\}.$$

Then,  $G_p^* \cong G_p$ . We fix such an isomorphism and regard subgroups of  $G_p$  as those of  $G_p^*$  if necessary. Put

$U^0 = \left( \begin{matrix} 0_p & 0_p \\ \pi 0_p & 0_p \end{matrix} \right)^x \cap G$ , where  $\pi$  is a prime element of  $O_p$  such that  $\pi^2 = \rho$ . Then,  $U_p^0$  is a minimal parahoric subgroup of  $G_p$ , and we can take

$$\tau_2 = U_p^0 \begin{pmatrix} 0 & -\pi^{-1} \\ \pi & 0 \end{pmatrix} U_p^0, \quad \tau_1 = U_p^0 \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} U_p^0.$$

There are three maximal compact subgroups (up to conjugation) in  $Sp(2, Q_p)$ , that is,  $K(p)_p = B(p)_p \cup S_0 \cup S_2 \cup S_0 S_2$ ,  $Sp(2, Z_p)$ , and  $\rho Sp(2, Z_p) \rho^{-1}$ , where

$$\rho = \begin{pmatrix} p & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & p & 0 & 0 \\ p & 0 & 0 & 0 \end{pmatrix}.$$

Among these, only  $K(p)_p$  is invariant by  $\sigma$ , and the group which 'corresponds' with  $K(p)_p$  by 'folding' is

$U_p^2 = U_p^0 \cup \tau_2$ . So, it is natural to consider that there exists some good correspondence between  $S_k(K(p))$  and  $M_{k-3}(U_2)$ . But, in spite that these are 'maximal' groups,



we must subtract the 'old forms' from each space. Now, we shall explain this. We intend to regard the cusp forms in  $S_k(K(p))$  obtained 'from'  $S_k(\text{Sp}(2, Z)) + S_k(\rho \text{Sp}(2, Z)\rho^{-1})$  as old forms. But  $K(p)$  is not conjugate to  $\text{Sp}(2, Z)$  or  $\rho \text{Sp}(2, Z)\rho^{-1}$ , and is not contained in, or does not contain any of these groups. So, we must define some mapping between these spaces. Define  $\text{Tr}_{K(p)/B(p)}: S_k(B(p)) \rightarrow S_k(K(p))$  by:

$$\text{Tr}_{K(p)/B(p)}(f) = \left( \sum_{\sigma \in B(p) \setminus K(p)} f | [\sigma]_k \right) / [K(p):B(p)]$$

for any  $f \in S_k(K(p))$ , where  $f | [\sigma]_k = f(\sigma z) \det(Cz+D)^{-k}$  for  $\sigma = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \text{Sp}(2, Q)$ . Denote by  $\text{Tr}$  the restriction of  $\text{Tr}_{K(p)/B(p)}$  on  $S_k(\text{Sp}(2, Z)) + S_k(\rho \text{Sp}(2, Z)\rho^{-1})$ .

We define new forms of  $S_k(K(p))$  to be the orthogonal complement of  $\text{Tr}(S_k(\text{Sp}(2, Z)) + S_k(\rho \text{Sp}(2, Z)\rho^{-1}))$  in  $S_k(K(p))$ , and denote it by  $S_k^0(K(p))$ . The map  $\text{Tr}$  does not vanish in general.

For example, we have

Lemma 1.2.      Let  $f \in S_k(\text{Sp}(2, Z))$  be an eigen form of the Hecke operators  $T(p)$  and  $T(p^2)$  with eigenvalues  $\lambda(p)$  and  $\lambda(p^2)$ , respectively. Assume that  $\lambda(p) \neq 0$  or  $\lambda(p^2) \neq p^{2k-2}$ . (For example, this is satisfied for all eigen forms of the Mass space  $M_k$ .) Then,  $\text{Tr}(f) \neq 0$ .

The proof consists of an easy argument on Fourier coefficients, and will be omitted here. In view of the Ramanujan Conjecture,

it is very plausible that the assumption of Lemma 1.2 is always satisfied. On the other hand, the map  $\text{Tr}$  is not injective in general:

Lemma 1.3.      Let  $k$  be an even integer, Then, for  
 $f \in M_k$  (the Maas space), we have  $\text{Tr}(f) = \text{Tr}(f | [L\rho]_k)$ .

The proof is easy and omitted here. It seems that, if  $k$  is odd, then  $\text{Tr}$  is injective, and if  $k$  is even, then

$\ker \text{Tr} = \{ f - f | [\rho]_k ; f \in M_k \}$ . If this is true, we have

$\dim S_k^0(K(p)) = \dim S_k(K(p)) - 2 \dim S_k(\text{Sp}(2, Z))$  for odd  $k$ , and

$\dim S_k^0(K(p)) = \dim S_k(K(p)) - 2 \dim S_k(\text{Sp}(2, Z)) + \dim S_{2k-2}(\text{SL}_2(Z))$

for even  $k$ . (Numerical examples in [9] support this.)

On the other hand, we can show that, if a common eigen form

$f \in M_\nu(U_2)$  satisfies a certain condition, then

$L(s, f) = L(s, g)L(s, h)$  for some  $g \in A_2(\Gamma_0(p))$  and

$h \in S_{2\nu+4}(\text{SL}_2(Z))$ . (This is a slight modification of

Ihara [13].) So, denote by  $M_\nu^E(U_2)$  the space spanned by

common eigen forms  $f \in M_\nu(U_2)$  such that  $L(s, f) = L(s, g)L(s, h)$

(up to Euler  $p$ -factor) for some  $g \in A_2(\Gamma_0(p))$  and

$h \in S_{2k-2}(\text{SL}_2(Z))$ . We define the space of new forms

of  $M_\nu(U_2)$  to be the orthogonal complement of  $M_\nu^E(U_2)$  in

$M_\nu(U_2)$ . Th.1 and some examples seem to suggest that

$$\dim M_\nu^0(U_2) = \dim M_\nu(U_2) - \dim \Lambda_2(\Gamma_0(p)) \times \dim S_{2\nu+4}(SL_2(\mathbb{Z}))$$

for even  $\nu$ , and

$$\dim M_\nu^0(U_2) = \dim M_\nu(U_2) - \dim S_2(\Gamma_0(p)) \times \dim S_{2\nu+4}(SL_2(\mathbb{Z}))$$

for odd  $\nu$ .

Conjecture 1.4. For any integer  $k \geq 5$ , there exists an isomorphism  $\phi$  of  $M_{k-3}^0(U_2)$  onto  $S_k^0(K(p))$  such that

$L(s, f) = L(s, \phi(f))$  (up to Euler  $p$ -factors) for any common eigen form  $f \in M_{k-3}^0(U_2)$  of all the Hecke operators  $T(n)$  ( $n \nmid p$ ).

Now, we point out one important fact. There exist some new forms of  $S_k(K(p))$  which can be obtained by lifting cusp forms in  $S_{2k-2}(\Gamma_0(p))$  (See examples in [9]). So, also in the case of  $M_\nu(U_2)$ , it seems more natural to

define new forms in the same point of view as in the case of  $S_k(K(p))$ . Put  $U_p^1 = GL_2(O_p) \cap G_p^*$ . Put

$$U_1 = G_\infty \prod_q U_q^1, \text{ and } U_0 = G_\infty U_p^0 \prod_{q \neq p} U_q^1.$$

The 'trace map'  $\text{Tr}_{U_2/U_0}$  of  $M_\nu(U_0)$  to  $M_\nu(U_2)$  can be

defined as before. Denote the orthogonal complement of  $\text{Tr}_{U_2/U_0}(M_\nu(U_1))$  in  $M_\nu(U_2)$  by  $M_\nu^1(U_2)$ . (We note here

that  $U_p^1$  is not conjugate to  $U_p^2$ , which causes the difference from the case of  $SL_2$ .) Then, it seems natural to

$$\text{expect } M_\nu^0(U_2) = M_\nu^1(U_2).$$

In the representation theoretic language,  
 our conjecture seems to be stated as follows:  
 Let  $\pi = \otimes \pi_q$  or  $\pi' = \otimes \pi'_q$  be an irreducible  
 admissible representation of  $\mathrm{GSp}(2, \mathbb{Q}_A)$  or  $G_A$ ,  
 respectively. (Here,  $\mathrm{GSp}$  means the symplectic  
 group with similitudes.) Assume that  $\pi_\infty$  corresponds  
 to  $\det^{v+3}$ ,  $\pi'_\infty$  to  $\rho_v$ ,  $\pi_q = \pi'_q$  for  $q \neq p$ , and that  
 $\pi_q (q \neq p)$  has a  $\mathrm{Sp}(2, \mathbb{Z}_q)$ -fixed vector. Assume  
 that  $\pi_p$  is an irreducible representation which has  
 a  $K(p)_p$ -fixed vector, but has no  $\mathrm{Sp}(2, \mathbb{Z}_p)$ - or  
 $\rho \mathrm{Sp}(2, \mathbb{Z}_p) \rho^{-1}$ -fixed vector, and that  $\pi'_p$  is an  
 irreducible representation of  $G_p$  which has a  $U_p^2$ -fixed  
 vector, but has no  $U_p^1$ -fixed vector. Then,  
 $L(s, \pi) = L(s, \pi')$ ?

## §2. Semi-simple conjugacy classes

In this section, we shall give 'local data' at  $p$  of semi simple conjugacy classes, then, give their contribution to  $\dim S_k(K(p))$  as Theorem 2. (The local data at  $q \neq p$  have been given in [7].) The proofs are lengthy and elaborate but similar technique can be found in [7], and we will omit them here.

We review some notations. Put

$$R = \gamma M_4(\mathbb{Z}_p) \gamma^{-1}, \quad \text{where} \quad \gamma = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & p & \\ & & & 1 \end{pmatrix},$$

and put

$$\text{GSp} = \left\{ g \in M_4(\mathbb{Q}_p); g \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} {}^t g = n(g) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\}.$$

Let  $R^\times$  be the invertible elements of  $R$ . For  $g \in \text{GSp}$ , let  $Z(g)$  be the commutator algebra of  $\mathbb{Q}_p(g)$  in  $M_4(\mathbb{Q}_p)$ .

For any  $\mathbb{Z}_p$ -order  $\Lambda_1, \Lambda_2$  of  $Z(g)$ , denote by  $\Lambda_1 \sim \Lambda_2$

when  $a^{-1} \Lambda_1 a = \Lambda_2$  for some  $a \in Z(g) \cap \text{GSp}$ . For any

torsion element  $g \in \text{GSp}$  and  $\mathbb{Z}_p$ -order  $\Lambda \subset Z(g)$ , put

$c_p(g, R, \Lambda) =$  the number of elements of  $M(g, \Lambda)$ ,

where  $M(g, \Lambda) = (Z(g) \cap G) \backslash M(g, R, \Lambda) / R^\times$  and

$$M(g, R, \Lambda) = \{ x \in \text{GSp}; x^{-1} g x \in R^\times, Z(g) \cap x R x^{-1} \sim \Lambda \}.$$

In the following sentences, we always denote by  $f(x)$

the principal polynomial of the elements in conjugacy classes treated there.

Proposition 2.1.    The total contribution of  $\pm 1 \in K(p)$  to  $\dim S_k(K(p))$  is given by:

$$(p^2+1)(2k-2)(2k-3)(2k-4)/2^9 3^3 5.$$

Proof    Obvious, because  $[Sp(2, Z) : B(p)] = (p^2+1)(p+1)^2$  and  $[K(p) : B(p)] = (p+1)^2$ .    q.e.d.

Proposition 2.2.    The representatives of  $K(p)/\{\pm 1\}$ -conjugacy classes with  $f(x) = (x-1)^2(x+1)^2$  are given by:

$$\delta_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \text{ and } \delta_2 = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & -1 & -1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}.$$

The contribution  $t(\delta_1)$ ,  $t(\delta_2)$  of each conjugacy class to  $\dim S_k(K(p))$  for  $k \geq 5$  is given by:

$$t(\delta_1) = (-1)^k(2k-2)(2k-4)/2^8 3^2,$$

$$t(\delta_2) = \begin{cases} (-1)^k(2k-2)(2k-4)/2^7 3, & \text{if } p \neq 2, \\ (-1)^k(2k-2)(2k-4)/2^9, & \text{if } p = 2. \end{cases}$$

Next, we treat the case where  $f(x) = (x-1)^2 g(x)$  and  $g(x)$  is an irreducible quadratic polynomial. Put  $F = Q[x]/g(x)$ . We identify the algebra  $M_2(Q_p) \times M_2(Q_p)$  with the algebra

$$\left\{ \begin{pmatrix} a & 0 & b & 0 \\ 0 & x & 0 & y \\ c & 0 & d & 0 \\ 0 & z & 0 & w \end{pmatrix} ; \quad a, b, c, d, x, y, z, w \in Q_p \right\}.$$

Put  $g = \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \omega \right)$ , where  $\omega = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ ,  $\begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$ , or  $\begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}$   
 for  $f(x) = (x-1)^2(x^2+1)$ ,  $(x-1)^2(x^2+x+1)$ , or  $(x-1)^2(x^2-x+1)$   
 , respectively.

Proposition 2.3.                      Notations being as above,

(i) if  $\left(\frac{F}{p}\right) = 1$ , then

$$c_p(g, R, \Lambda) = \begin{cases} 2 \dots & \text{if } \Lambda \sim M_2(\mathbb{Z}_p) \oplus \mathbb{Z}_p^2 \\ 0 \dots & \text{otherwise,} \end{cases}$$

(ii) if  $\left(\frac{F}{p}\right) = -1$ , or  $p = 3$  and  $f(x) = (x-1)^2(x^2-x+1)$ , then

$$c_p(g, R, \Lambda) = \begin{cases} 2 \dots & \text{if } \Lambda \sim M_2(\mathbb{Z}_p) \oplus \mathbb{Z}_p = \Lambda_1, \\ 0 \dots & \text{otherwise,} \end{cases}$$

and  $M(g, \Lambda_1) = \left\{ 1_4, \begin{pmatrix} 0 & p & 0 & 0 \\ p & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \right\}$  ,

(iii) if  $p = 3$  and  $f(x) = (x-1)^2(x^2+x+1)$ , then

$$c_p(g, R, \Lambda) = \begin{cases} 2 \dots & \text{if } \Lambda \sim \Lambda_1 = M_2(\mathbb{Z}_p) \oplus \mathbb{Z}_p^2 \\ 1 \dots & \text{if } \Lambda \sim \Lambda_2 \\ 0 \dots & \text{otherwise,} \end{cases}$$

where  $\Lambda_2 = \left\{ \left( \begin{pmatrix} a & 3b \\ c & d \end{pmatrix}, x+y\omega \right) ; a, b, c, d, x, y \in \mathbb{Z}_3, \right.$   
 $\left. x+y \equiv d \pmod{3} \right\}$  ,

$[\Lambda_1 \cap \text{GSp} : \Lambda_2 \cap \text{GSp}] = 6$ , and

$M(\Lambda_1)$  is as in (ii) ,  $M(\Lambda_2) = \left\{ \begin{pmatrix} -3 & 3 & -1 & 0 \\ 3 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix} \right\}$  .

(iv) if  $(\frac{F}{p}) = 0$  and  $p = 2$ ,

$$c_p(g, R, \Lambda) = \begin{cases} 2 \dots & \text{if } \Lambda \sim \Lambda_1 = M_2(\mathbb{Z}_p) \oplus \mathbb{Z}_p[\omega], \\ 1 \dots & \text{if } \Lambda \sim \Lambda_2, \\ 0 \dots & \text{otherwise,} \end{cases}$$

where  $\Lambda_2 = \left\{ \left( \begin{pmatrix} a & 2b \\ c & d \end{pmatrix}; x+y\omega \right); a, b, c, d, x, y \in \mathbb{Z}_2 \right. \\ \left. x-y \equiv d \pmod{2} \right\}$

$[\Lambda_1 \cap \text{GSp} : \Lambda_2 \cap \text{GSp}] = 3$ , and

$M(\Lambda_1)$  is as in (ii),  $M(\Lambda_2) = \left\{ \begin{pmatrix} -2 & 2 & 1 & 0 \\ 2 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix} \right\}$ .

Next, we treat the case where  $f(x) = g(x)^2$  and  $g(x)$  is an irreducible quadratic polynomial. First, we treat the case where  $\mathbb{Z}_0(g)$  is split. (As for the notation  $\mathbb{Z}_0(g)$ , see [7](I)). Put  $F = Q[x]/g(x)$ .

Proposition 2.4. Assumptions being as above,

(i) if  $(\frac{F}{p}) = -1$ , then  $c_p = 0$  for any  $\Lambda$ ,

(ii) if  $(\frac{F}{p}) = 1$ , take  $g = \begin{pmatrix} a1_2 & 0 \\ 0 & b1_2 \end{pmatrix}$ , where  $a, b \in \mathbb{Q}_p$  and

$g(x) = (x-a)(x-b)$ , then

$$c_p(g, R, \Lambda) = \begin{cases} 1 \dots & \text{if } \Lambda \sim \Lambda_1 = M_2(\mathbb{Z}_p) \oplus \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^{-1} M_2(\mathbb{Z}_p) \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}, \\ 0 \dots & \text{otherwise,} \end{cases}$$

$[\text{GSp} \cap \text{GL}_2(\mathbb{Z}_p)^2 : \Lambda_1 \cap \text{GSp}] = p+1$ , where we embed

$M_2(\mathbb{Q}_p)^2$  in  $M_4(\mathbb{Q}_p)$  diagonally.



(iii) if  $(\frac{F}{p}) = 0$  and  $p = 2$ , take  $g = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 1 & 0 & 0 & -1 \end{pmatrix}$ , then

$$c_p(g, R, \Lambda) = \begin{cases} 1 \dots & \text{if } \Lambda \sim \Lambda_1 = xRx^{-1} \cap Z(g), \\ 1 \dots & \text{if } \Lambda \sim \Lambda_2 = yRy^{-1} \cap Z(g), \\ 0 \dots & \text{otherwise,} \end{cases}$$

where  $x = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ -1 & 1 & -1 & -1 \\ 1 & 0 & 0 & -1 \end{pmatrix}$  and  $y = \begin{pmatrix} 4 & 0 & 1 & 2 \\ -4 & 4 & 1 & 0 \\ -4 & 4 & -3 & -4 \\ 4 & 0 & 1 & -2 \end{pmatrix}$ ,

$$d_2(\Lambda_1) = 3, e_2(\Lambda_1) = 2, d_2(\Lambda_2) = 6, e_2(\Lambda_2) = 2,$$

(iv) if  $(\frac{F}{p}) = 0$  and  $p = 3$ , take  $g = \begin{pmatrix} 1 & 1 & -1 & 0 \\ -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 1 \end{pmatrix}$ , then

$$c_p(g, R, \Lambda) = \begin{cases} 1 \dots & \text{if } \Lambda \sim \Lambda_1 = xRx^{-1} \cap Z(g), \\ 1 \dots & \text{if } \Lambda \sim \Lambda_2 = yRy^{-1} \cap Z(g), \\ 0 \dots & \text{otherwise,} \end{cases}$$

where  $x = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{pmatrix}$  and  $y = \begin{pmatrix} 3 & 0 & 0 & 1 \\ 3 & 3 & 1 & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ ,

$$d_3(\Lambda_1) = 1, e_3(\Lambda_1) = 1, d_3(\Lambda_2) = 8, e_3(\Lambda_2) = 2,$$

where  $d_p(\Lambda)$  and  $e_p(\Lambda)$  are as in [7]

Next, we treat the case where  $Z_0(g)$  is division.

Then,  $(\frac{F}{p}) \neq 1$  by definition of  $Z_0(g)$ .

Proposition 2.5.

(i) If  $(\frac{F}{p}) = -1$ , take  $g = \left( \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \omega \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}^{-1}, \omega \right)$ ,

where  $\omega$  are as in Prop.2.3, and  $g$  are regarded as  
elements of GSp as in Prop.2.3. Then,

$$c_p(g, R, \Lambda) = \begin{cases} 1 \dots \text{if } \Lambda \sim \Lambda_1 = M_2(Z_p[\omega]), \\ 0 \dots \text{otherwise,} \end{cases}$$

and  $d_p(\Lambda_1) = e_p(\Lambda_1) = 1,$

(ii) if  $(\frac{F}{p}) = 0$  and  $p = 2$ , take  $g = \begin{pmatrix} 0 & -1 \\ 1_2 & 0 \end{pmatrix}$ , then

$$c_p(g, R, \Lambda) = \begin{cases} 1 \dots \text{if } \Lambda \sim \Lambda_1 = xRx^{-1} \cap Z(g), \\ 1 \dots \text{if } \Lambda \sim \Lambda_2 = yRy^{-1} \cap Z(g), \\ 0 \dots \text{otherwise,} \end{cases}$$

where  $x = \begin{pmatrix} p1_2 & 1_2 \\ 0 & 1_2 \end{pmatrix}$ ,  $y = \begin{pmatrix} 2 & 0 & 0 & 1 \\ -2 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ ,

$d_2(\Lambda_1) = 6$ ,  $e_2(\Lambda_1) = 2$ ,  $d_2(\Lambda_2) = 1$ , and  $e_2(\Lambda_2) = 2.$

(iii) if  $(\frac{F}{p}) = 0$  and  $p = 3$ , take  $g = \pm \begin{pmatrix} 0 & 1_2 \\ -1_2 & 1_2 \end{pmatrix}$ , then

$$c_p(g, R, \Lambda) = \begin{cases} 1 \dots \text{if } \Lambda \sim \Lambda_1 = xRx^{-1} \cap Z(g), \\ 0 \dots \text{otherwise,} \end{cases}$$

where  $x = \begin{pmatrix} p & 0 & 0 & 0 \\ 0 & p & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ , and  $d_3(\Lambda_1) = 4$ ,  $e_3(\Lambda_1) = 2.$

Next, we treat the regular elements  $g \in K(p)$ . When  $Z[g]$  is the maximal order of  $Q[g]$ , it is fairly easy to classify global conjugacy classes. We sketch it here. Let

$\zeta \in Sp(2, Z)$  be an element whose principal polynomial is  $f(x) = (x^2+1)(x^2 \pm x+1)$ ,  $x^4 \pm x^3 + x^2 \pm x+1$ ,  $x^4+1$ , or  $x^4-x^2+1$ . (It exists and we fix it.) When  $f(x) = (x^2+1)(x^2 \pm x+1)$

more explicitly, put  $\zeta = \begin{pmatrix} -1 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$ .

Put  $J = \begin{pmatrix} 0 & 1_2 \\ -1_2 & 0 \end{pmatrix}$ . Assume that  $g^{-1}\zeta g \in K(p)$  for some

$g \in GL_4(Q)$ . Then,  $\zeta(gJ^t gJ^{-1}) = (gJ^t gJ^{-1})\zeta$ , and  $gJ^t gJ^{-1} \in Q(\zeta)$ . The map  $Q(\zeta) \ni h \rightarrow J^t h J^{-1} \in Q(\zeta)$  is

the complex conjugation on  $Q(\zeta)$ , and  $gJ^t gJ^{-1}$  is invariant by this map. So,  $gJ^t gJ^{-1} \in Q(\zeta + \zeta^{-1})$ . Put

$\delta = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & p & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ . Then,  $\delta^{-1} g^{-1} \zeta g \delta \in M_4(Z)$ . Now,

the class number of  $Q(\zeta)$  is one. So, by virtue of Chevalley [2],  $a g \delta \in GL_4(Z)$  for some  $a \in Q(\zeta)$ .

Lemma 2.6. Let  $f(x)$  be one of the above polynomials.

Then, the set of  $K(p)$ -conjugacy classes with principal polynomial  $f(x)$  is bijective to the set

$$\{ \alpha / p ; \alpha \in Z[\zeta + \zeta^{-1}], N(\alpha) = \pm p \} / N_{Q(\zeta)/Q(\zeta + \zeta^{-1})}(Z[\zeta]^{\times})$$

The map is given by:

$$g^{-1}\zeta g, (g \delta \in GL_4(Z)) \longrightarrow gJ^t gJ^{-1}.$$

Proof The injectivity is obvious. The surjectivity is proved by case by case process. q.e.d.

Proposition 2.7. The numbers of  $K(p)$ -conjugacy classes of above types are given as follows:

$$(x^2+1)(x^2+x+1) \dots 8$$

$$x^4+1 \dots \begin{cases} 0 \dots \text{if } \left(\frac{F}{p}\right) = -1, \\ 4 \dots \text{if } \left(\frac{F}{p}\right) = 0, \\ 8 \dots \text{if } \left(\frac{F}{p}\right) = 1, \end{cases}$$

$$x^4+x^3+x^2+x+1, \text{ and } \dots \text{ same as in } x^4+1, \\ x^4-x^3+x^2-x+1$$

$$x^4-x^2+1 \dots \begin{cases} 0 \dots \text{if } \left(\frac{F}{p}\right) = -1, \\ 2 \dots \text{if } \left(\frac{F}{p}\right) = 0, \\ 4 \dots \text{if } \left(\frac{F}{p}\right) = 1, \end{cases}$$

where  $F = Q(\zeta + \zeta^{-1})$ .

Next, we treat the case where  $f(x) = (x^2+x+1)(x^2-x+1)$ .

In this case,  $Z[x]/f(x)$  is not the maximal order, and we give the local data instead of giving global conjugacy classes. Put  $F = Q[x]/(x^2+x+1)$ . Put

$$g = \begin{pmatrix} a & 0 & 0 & 0 \\ 0 & -a & 0 & 0 \\ 0 & 0 & b & 0 \\ 0 & 0 & 0 & -b \end{pmatrix}, \text{ when } \left(\frac{F}{p}\right) = 1, \text{ where } f(x) = (x^2-a^2)(x^2-b^2), \\ a, b \in Q_p,$$

$$g_1 = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & -1 & 0 \\ 0 & -1 & 0 & 1 \end{pmatrix}, \quad g_2 = \begin{pmatrix} 0 & 0 & -1/p & 0 \\ 0 & 0 & 0 & 1 \\ p & 0 & -1 & 0 \\ 0 & -1 & 0 & 1 \end{pmatrix}, \quad \text{when } \left(\frac{F}{p}\right) = -1,$$

and

$$g_1 = \begin{pmatrix} 0 & 0 & -1/p & 0 \\ 0 & 0 & 0 & 1 \\ p & 0 & -1 & 0 \\ 0 & -1 & 0 & 1 \end{pmatrix}, \quad g_2 = \begin{pmatrix} 0 & 0 & 1/p & 0 \\ 0 & 0 & 0 & 0 \\ -p & 0 & -1 & 0 \\ 0 & -1 & 0 & 1 \end{pmatrix}, \quad \text{when } p = 3.$$

Proposition 2.8.

(i) If  $\left(\frac{F}{p}\right) = 1$ , then

$$c_p(g, R, \Lambda) = \begin{cases} 2 \dots & \text{if } \Lambda \sim Z_p^4 \\ 0 \dots & \text{otherwise,} \end{cases}$$

where  $Z_p^4$  is embedded diagonally in  $M_4(Z_p)$ , and

$$M(g, Z_p^4) = \left\{ 1_4, \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \right\},$$

(ii) If  $\left(\frac{F}{p}\right) = -1$ , then,

$c_p(g_1, R, \Lambda) = 0$  for any  $\Lambda$ , and

$$c_p(g_2, R, \Lambda) = \begin{cases} 2 \dots & \text{if } \Lambda \sim o_p, \\ 0 \dots & \text{otherwise,} \end{cases}$$

where  $o_p$  is the maximal order of  $F_p = Q_p(g_2)$ , and

$$M(g_2, o_p) = \left\{ 1_4, \begin{pmatrix} 0 & 1 & 0 & 0 \\ p & 0 & 0 & 0 \\ 0 & 0 & 0 & p \\ 0 & 0 & 1 & 0 \end{pmatrix} \right\}.$$

(iii) If  $\left(\frac{F}{p}\right) = 0$  ( $p = 3$ ), then

$$c_p(g_i, R, \Lambda) = \begin{cases} 2 \dots & \text{if } \Lambda \sim o_p, \\ 0 \dots & \text{otherwise,} \end{cases} \quad \text{for } i = 1, 2,$$

where  $o_p$  and  $M(g_i, o_p)$  are the same as in (ii).

Now, denote by  $H_1$  the total contribution to  $\dim S_k(K(p))$ , of those semi-simple conjugacy classes whose principal polynomials are of the form  $f_1(\pm x)$ , where the polynomials  $f_1(x)$  are defined as in [7] (I) p.590. We can give  $H_1$  explicitly as a corollary to the above results by using [7] and Hashimoto [6].

Theorem 2      Assume that  $k \geq 5$ , then  $H_1$  and  $H_2$  have been given in Prop.2.1, 2.2, and  $H_i (i \geq 3)$  are given as follows:

$$H_3 = \begin{cases} [k-2, -k+1, -k+2, k-1 ; 4] / 2^4 3, \dots & \text{if } p \neq 2, \\ 5 [k-2, -k+1, -k+2, k-1 ; 4] / 2^5 3, \dots & \text{if } p = 2, \end{cases}$$

$$H_4 = \begin{cases} [2k-3, -k+1, -k+2 ; 3] / 2^2 3^3, \dots & \text{if } p \neq 3, \\ 5 [2k-3, -k+1, -k+2 ; 3] / 2^2 3^3, \dots & \text{if } p = 3, \end{cases}$$

$$H_5 = [-1, -k+1, -k+2, 1, k-1, k-2 ; 6] / 2^2 3^2,$$

$$H_6 = \begin{cases} \frac{5(2k-3)(p+1)}{2^7 3} + \frac{(-1)^k (p+1)}{2^7} \dots & \text{if } p \equiv 1 \pmod{4}, \\ \frac{(2k-3)(p-1)}{2^7} + \frac{5(-1)^k (p-1)}{2^7 3} \dots & \text{if } p \equiv 3 \pmod{4}, \\ \frac{3(2k-3)}{2^7} + \frac{7(-1)^k}{2^7 3} \dots & \text{if } p = 2, \end{cases}$$

$$H_7 = \begin{cases} \frac{(2k-3)(p+1)}{2 \cdot 3^3} + \frac{(p+1)}{2^2 3^3} [0, -1, 1; 3] \dots \text{if } p \equiv 1 \pmod{3}, \\ \frac{(2k-3)(p-1)}{2^2 3^3} + \frac{(p-1)}{2 \cdot 3^3} [0, -1, 1; 3] \dots \text{if } p \equiv 2 \pmod{3}, \\ \frac{5(2k-3)}{2^2 3^3} + \frac{1}{3^3} [0, -1, 1; 3] \dots \text{if } p = 3, \end{cases}$$

$$H_8 = [1, 0, 0, -1, -1, -1, -1, 0, 0, 1, 1, 1; 12] / 2 \cdot 3,$$

$$H_9 = \begin{cases} 2 [1, 0, 0, -1, 0, 0] / 3^2 \dots \text{if } p \neq 2, \\ [1, 0, 0, -1, 0, 0] / 2 \cdot 3^2 \dots \text{if } p = 2, \end{cases}$$

$$H_{10} = (1 + (\frac{5}{p})) [1, 0, 0, -1, 0; 5] / 5,$$

$$H_{11} = (1 + (\frac{2}{p})) [1, 0, 0, -1; 4] / 2^3, \text{ and}$$

$$H_{12} = \begin{cases} [0, 1, -1; 3] / 2 \cdot 3 \dots \text{if } p \equiv 1 \pmod{12}, \\ (-1)^k / 2 \cdot 3 \dots \text{if } p \equiv 11 \pmod{12}, \\ (-1)^k / 2^2 \cdot 3 \dots \text{if } p = 2, 3, \\ 0 \dots \text{if } p \equiv 5, 7 \pmod{12}, \end{cases}$$

where  $(\frac{*}{p})$  is the Legendre symbol, and

$t = [t_0, t_1, \dots, t_{q-1}; q]$  means that

$t = t_j$  if  $k \equiv j \pmod{q}$ .

§3. Conjugacy classes of non semi-simple types

In this section, we shall give the representatives of non semi-simple  $K(p)$ -conjugacy classes which have non-zero contribution to  $\dim S_k(K(p))$ , and give their contribution to  $\dim S_k(K(p))$  ( $k \geq 5$ ).

Put

$$P_0 = \left\{ \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \in Sp(2, Q) \right\} \quad \text{and}$$

$$P_1 = \left\{ \begin{pmatrix} * & 0 & * & * \\ * & * & * & * \\ * & 0 & * & * \\ 0 & 0 & 0 & * \end{pmatrix} \in Sp(2, Q) \right\} .$$

Lemma 3.1.      Assume that  $g \in Sp(2, Q)$  is not semi-simple.  
Then, some  $Sp(2, Q)$ -conjugate of  $g$  is contained in  $P_0$  or  $P_1$ .

As for the proof, see Borel-Tits [1]. Next two lemmata are easy and the proof will be omitted.

Lemma 3.2.      The Satake compactification of  $K(p) \backslash Sp(2, R)$  has the unique zero-dimensional cusp and two one-dimensional cusps, that is

$$Sp(2, Q) = K(p)P_0$$

$$= K(p)P_1 \cup K(p) \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} K(p).$$



Lemma 3.3.     Assume that  $g \in K(p)$  is not semi-simple.  
Then, some  $K(p)$ -conjugate of  $g$  is contained in  $P_0, P_1,$   
or  $P_1^!$ , where

$$P_1^! = \left\{ \begin{pmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & 0 \\ 0 & * & * & * \end{pmatrix} \in Sp(2, Q) \right\} .$$

By this Lemma, we can assume that  $g \in P_0, P_1,$  or  $P_1^!$ .  
 Then, by case by case direct calculations, we can  
 give a complete list of  $K(p)$ -conjugacy classes which  
 are not semi simple and which have contribution to  
 $\dim S_k(K(p))$ . The proofs are lengthy but routine,  
 and will be omitted here.

Theorem 3     The representatives of  $K(p)$ -conjugacy  
classes which are of elliptic/parahoric,  $\delta$ -parabolic,  
parabolic, or paraelliptic (in the sense of Hashimoto[6]),  
are given in the following list, together with their  
contribution to  $\dim S_k(K(p)) (k \geq 5)$ . The contribution to  
 $\dim S_k(K(p))$ , of each set of conjugacy classes below,  
are denoted by  $I_i$ .

(I) elliptic/parabolic

(1)  $f(x) = (x-1)(x^2-x+1)$  and  $(x+1)^2(x^2+x+1)$ ,

$$\begin{aligned} & \pm \begin{pmatrix} 0 & 0 & 1/p & 0 \\ 0 & 1 & 0 & n \\ -p & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} & \pm \begin{pmatrix} 1 & 0 & 1/p & 0 \\ 0 & 1 & 0 & n \\ p & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\ & \pm \begin{pmatrix} 0 & 0 & n/p & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \end{pmatrix} & \pm \begin{pmatrix} 1 & 0 & n/p & 0 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \quad (n \in \mathbb{Z}, n \neq 0) \end{aligned}$$

The total contribution of the above conjugacy classes to  
 $\dim S_k(K(p))$  is given by:

$I_1 = [0, 1, 1, 0, -1, -1 ; 6] / 6,$

(2)  $f(x) = (x-1)^2(x^2+x+1)$  and  $(x+1)^2(x^2-x+1)$

(i) 
$$\begin{aligned} & \pm \begin{pmatrix} 0 & 0 & -1/p & 0 \\ 0 & 1 & 0 & n \\ p & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} & \pm \begin{pmatrix} -1 & 0 & 1/p & 0 \\ 0 & 1 & 0 & n \\ p & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\ & \pm \begin{pmatrix} 1 & 0 & n/p & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 \end{pmatrix} & \pm \begin{pmatrix} 1 & 0 & n/p & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} \end{aligned}$$

$(n \in \mathbb{Z}, n \neq 0)$

$: I_2 = [-2, 1, 1 ; 3] / 2 \cdot 3^2,$

$$(ii) \quad \pm \begin{pmatrix} 1 & 0 & n/p & 1 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & -1 \end{pmatrix} \quad \pm \begin{pmatrix} 1 & 0 & -n/p & 1 \\ 0 & -1 & -1 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & -1 & 0 \end{pmatrix}$$

$$\pm \begin{pmatrix} 0 & 0 & -1/p & 0 \\ 0 & 1 & 1 & n \\ p & 0 & -1 & p \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \pm \begin{pmatrix} -1 & 0 & 1/p & -1 \\ 0 & 1 & 1 & -n \\ -p & 0 & 0 & -p \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

( $n \in \mathbb{Z}$ ,  $n \neq 0$  if  $p \neq 3$  and  $n \neq -1$  if  $p = 3$ ) :

$$I_3 = \begin{cases} [-2, 1, 1 ; 3] / 3^2 & \dots \text{ if } p = 3, \\ 2 [-1, 1, 0 ; 3] / 3^2 & \dots \text{ if } p \equiv 1 \pmod{3}, \\ 2 [-1, 0, 1 ; 3] / 3^2 & \dots \text{ if } p \equiv 2 \pmod{3}, \end{cases}$$

(3)  $f(x) = (x-1)^2(x^2+1)$  and  $(x+1)^2(x^2+1)$ ,

$$\pm \begin{pmatrix} 0 & 0 & 1/p & 0 \\ 0 & 1 & 0 & n \\ -p & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \pm \begin{pmatrix} 0 & 0 & -1/p & 0 \\ 0 & 1 & 0 & n \\ p & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\pm \begin{pmatrix} 1 & 0 & n/p & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} \quad \pm \begin{pmatrix} 1 & 0 & n/p & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

( $n \in \mathbb{Z}$ ,  $n \neq 0$ ),

$$\pm \begin{pmatrix} 0 & 0 & 1/p & 0 \\ 0 & 1 & 1 & n \\ -p & 0 & 0 & -p \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \pm \begin{pmatrix} 0 & 0 & -1/p & 0 \\ 0 & 1 & 1 & -n \\ p & 0 & 0 & p \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\pm \begin{pmatrix} 1 & 0 & n/p & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & -1 & 0 \end{pmatrix} \quad \pm \begin{pmatrix} 1 & 0 & -n/p & 1 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \end{pmatrix}$$

( $n \in \mathbb{Z}$ ,  $n \neq 0$  if  $p \neq 2$  and  $n \neq 1$  if  $p = 2$ ):

$$I_4 = [-1, 1, 1, -1 ; 4] / 2^2,$$

(II)  $\delta$  -parabolic :  $f(x) = (x-1)^2(x+1)^2$

(i)  $\pm \begin{pmatrix} 1 & 0 & n/p & 0 \\ 0 & -1 & 0 & m \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \pm \begin{pmatrix} 1 & 0 & n/p & -1 \\ 0 & -1 & 1 & m \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$

$(n, m \in \mathbb{Z}, n \neq 0, m \neq 0) :$

$I_5 = (-1)^k/2^3,$

(ii)  $\begin{pmatrix} 1 & 1 & n/p & m \\ 0 & -1 & m & -2m \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & -1 \end{pmatrix} \begin{pmatrix} 1 & 1 & n/p & m-1 \\ 0 & -1 & m & -2m+1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & -1 \end{pmatrix}$

$\begin{pmatrix} -1 & 0 & 2m/p & m \\ -p & 1 & m-1 & n \\ 0 & 0 & -1 & -p \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} -1 & 0 & (2m-1)/p & m \\ -p & 1 & m-1 & n \\ 0 & 0 & -1 & -p \\ 0 & 0 & 0 & 1 \end{pmatrix}$

$(m, n \in \mathbb{Z},$  and  $(2n+pm, -2m), (4n+p(2m-1), -2m+1),$   
 $(2m, 2n-pm),$  or  $(2m-1, 4n-p(2m-1)),$  is not  
equal to  $(0, 0),$  respectively.):

$I_6 = (-1)^k(2 - (-\frac{1}{p}))/2^4,$

(iii)  $\pm \begin{pmatrix} 1 & 0 & & S \\ 0 & -1 & & \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix},$  where  $S = \begin{pmatrix} n/p & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & n \end{pmatrix},$   
 $\begin{pmatrix} n/p & -1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ 1 & n \end{pmatrix}$

$(n \in \mathbb{Z}, n \neq 0)$

$I_7 = -(-1)^k(2k-3)/2^3.$

(III) parahoric :  $f(x) = (x-1)^4$  and  $(x+1)^4$

$$(1) \pm \begin{pmatrix} 1_2 & S \\ 0 & 1_2 \end{pmatrix} ; S = \begin{pmatrix} 0 & 0 \\ 0 & n \end{pmatrix}, \quad \begin{pmatrix} n/p & 0 \\ 0 & 0 \end{pmatrix} \quad n \in \mathbb{Z}, n \neq 0,$$

$$I_8 = -p(2k-3)/2^4 \cdot 3^2.$$

Next, put  $L = \left\{ \begin{pmatrix} s_1 & s_{12} \\ s_{12} & s_2 \end{pmatrix} \quad s_1 \in p^{-1}\mathbb{Z}, s_{12}, s_2 \in \mathbb{Z} \right\}$

and for  $S_1, S_2 \in L$ , denote  $S_1 \sim S_2$  when  $S_1 = US_2^t U$  for  
some  $U \in \Gamma_0(p) \cup \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \Gamma_0(p)$ .

$$(2) \pm \begin{pmatrix} 1_2 & S \\ 0 & 1_2 \end{pmatrix} ; S \in \{ S \in L, \det S \in (Q^x)^2 \} / \sim ,$$

$$I_9 = -1 / 2^3 3$$

$$(3) \pm \begin{pmatrix} 1_2 & S \\ 0 & 1_2 \end{pmatrix} ; S \in \{ S \in L, S \text{ definite } \} / \sim ,$$

$$I_{10} = (p+1)/2^3 3,$$

$$(4) \pm \begin{pmatrix} 1_2 & S \\ 0 & 1_2 \end{pmatrix} ; S \in \{ S \in L, S \text{ indefinite, } \det S \in (Q^x)^2 \} / \sim$$

(the contribution to the dimension is zero),

(IV) paraelliptic

Put

$$g(d) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ d & 1 & 0 & 0 \\ 0 & 0 & 1 & -d \\ 0 & 0 & 0 & 1 \end{pmatrix}, \text{ where } d \text{ is some integer.}$$

(1)  $f(x) = (x^2+1)^2 :$

(i) If  $\left(\frac{-1}{p}\right) = -1$ , there exists none in  $K(p)$ ,

(ii) if  $\left(\frac{-1}{p}\right) = 1$ , then

$$g(d)^{-1} \begin{pmatrix} 0 & -1 & S & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix} g(d), \quad S = \begin{pmatrix} 0 & -n \\ n & 0 \end{pmatrix}, \begin{pmatrix} 1 & n \\ -n & 1 \end{pmatrix}$$

$(n \in \mathbb{Z}, n \neq 0),$

$$\begin{pmatrix} 0 & -n \\ n+1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & n+1 \\ -n & 1 \end{pmatrix}$$

$(n \in \mathbb{Z}),$

where d runs through a set of the representatives in  $\mathbb{Z}$  of the solutions of  $d^2+1 \equiv 0 \pmod{p}$ , and

(iii) if  $p = 2$ , then

$$g(1)^{-1} \begin{pmatrix} 0 & -1 & S & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix} g(1), \quad S = \begin{pmatrix} 0 & -n \\ n & 0 \end{pmatrix}, \begin{pmatrix} 2^{-1} & 2^{-1}-n \\ 2^{-1}+n & -2^{-1} \end{pmatrix}$$

$(n \in \mathbb{Z}, n \neq 0),$

$$\begin{pmatrix} 0 & -n \\ n+1 & 0 \end{pmatrix}, \begin{pmatrix} 2^{-1} & -2^{-1}-n \\ 2^{-1}+n & -2^{-1} \end{pmatrix}$$

$(n \in \mathbb{Z})$

$$I_{11} = -\frac{1}{8} \left(1 + \left(\frac{-1}{p}\right)\right),$$

(2)  $f(x) = (x^2+x+1)^2$  and  $(x^2-x+1)^2$  :

(i) if  $(\frac{-3}{p}) = -1$ , then, there exists none in  $K(p)$ ,

(ii) if  $(\frac{-3}{p}) = 1$ , then,

$$\pm g(d)^{-1} \begin{pmatrix} 0 & -1 & & S \\ 1 & -1 & & \\ 0 & 0 & -1 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix} g(d), \quad S = \begin{pmatrix} -n & -2n \\ n & -n \end{pmatrix},$$

( $n \in \mathbb{Z}, n \neq 0$ )

$$\begin{pmatrix} -n & -2n \\ n+1 & -n \end{pmatrix}, \begin{pmatrix} -n & -2n \\ n+2 & -n \end{pmatrix},$$

( $n \in \mathbb{Z}$ )

where d runs through a set of the representatives of the solutions of  $x^2+x+1 \equiv 0 \pmod{p}$ , and

(iii) if  $p = 3$ , then, besides the above conjugacy classes in (ii)(here, we put  $d = 1$ ), there exists following conjugacy classes:

$$\begin{pmatrix} 1 & -1 & & B \\ 3 & -2 & & \\ 0 & 0 & -2 & -3 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \quad \text{where } B = \begin{pmatrix} -m-2e-h & -3m-6e-h \\ 0 & -3m-6e+h \end{pmatrix},$$

$e = \pm 1/3$ ,  $h = 0, \pm 1$ , and m is any integer such that  $3m+6e+h \neq 0$  :

$I_{12} = - \frac{1}{6} (1 + (\frac{-3}{p})) .$

§4. Proof of Theorem 1

In this section, we prove Theorem 1. First, we get

Theorem 4 For any integer  $k \geq 5$  and any prime integer  $p$ , we have

$$\dim S_k(K(p)) = \sum_{i=1}^{12} H_i + \sum_{i=1}^{12} I_i ,$$

where  $H_i$  or  $I_i$  is given in Th.2 or Th.3, respectively.

Numerical examples of  $\dim S_k(K(p))$

$p \backslash k$	5	6	7	8	9	10	11	12	13	14	15	16
2	0	0	0	1	0	1	1	2	0	2	1	4
3	0	1	0	1	1	2	1	4	1	4	3	6
5	1	1	1	2	2	4	4	6	5	9	8	13
7	1	2	2	4	4	7	7	11	11	16	16	24
11	2	3	3	6	7	12	14	20	22	32	36	48
13	3	5	7	10	13	19	23	31	37	48	56	72



By virtue of [7] and Igusa [12], our Theorem 1 is a corollary to Theorem 4. But it is interesting to see the details of contribution of each conjugacy classes. We denote by  $J_i$  the contribution to

$$\dim S_k(K(p)) - 2 \dim S_k(\text{Sp}(2, Z)) - \dim M_{k-3}(U_2),$$

of those semi-simple conjugacy classes whose principal polynomials are of the form  $f_i(\pm x)$  ( $i = 1, \dots, 12$ ). (As for the notations  $f_i(x)$ , see [7] p.590, e.g.,

$$f_6(x) = (x^2+1)^2, f_7(x) = (x^2+x+1)^2, \text{ and } f_{12}(x) = x^4-x^2+1.)$$

We get the following result.

Proposition 4.1.      The numbers  $J_i$  ( $i = 1, \dots, 12$ ) are given as follows:

$$J_i = 0 \quad \text{if } i \neq 6, 7, 12, \quad \text{and}$$

$$J_6 = \frac{1}{2^4} \left(1 - \left(\frac{-1}{p}\right)\right) + \frac{(p-1)}{2^4 3} (-1)^k - \frac{k}{2^3 3} \left(1 - \left(\frac{-1}{p}\right)\right),$$

$$J_7 = \frac{1}{2^2 3} \left(1 - \left(\frac{-3}{p}\right)\right) + \frac{(p-1)}{2^2 3^2} [0, -1, 1; 3] - \frac{k}{2 \cdot 3^2} \left(1 - \left(\frac{-3}{p}\right)\right)$$

$$J_{12} = \frac{1}{2^2 3} \left(1 - \left(\frac{-3}{p}\right)\right) (-1)^k + \frac{1}{2^2 3} \left(1 - \left(\frac{-1}{p}\right)\right) [0, -1, 1; 3].$$

Proof. The contribution to  $\dim M_{k-3}(U_2)$  has been given in [7],  $\dim S_k(\text{Sp}(2, Z))$  in Hashimoto [6], and  $\dim S_k(K(p))$  in Theorem 2 of this paper.      q.e.d.

Remark This result is rather mysterious. Those elements with the principal polynomials  $f_i(x)$  ( $i=8, \dots, 12$ ) are regular elements. Among those, as stated above, only  $J_{12}$  is exceptionally non-zero. I do not know the intrinsic reason of this.

Next, we shall give the contribution to  $\dim S_k(K(p)) - 2 \dim S_k(\text{Sp}(2, Z))$ , of non-semi-simple conjugacy classes. (Note that there is no such contribution to  $M_{k-3}(U_2)$ .) More precisely, take a set  $\{\sigma\} \subset \text{Sp}(2, R)$  of non semi-simple elements, and denote by  $K(\{\sigma\})$  the contribution to  $\dim S_k(K(p)) - 2 \dim S_k(\text{Sp}(2, Z))$ , of those  $K(p)$ -conjugacy classes whose elements are  $\text{Sp}(2, R)$ -conjugates of one of  $\{\sigma\}$ . Put

$$\hat{S}(\pm 1, \pm 1) = \begin{pmatrix} 1 & 0 & \pm 1 & 0 \\ 0 & 1 & 0 & \pm 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad a = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$b = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \hat{\beta}(\theta, \lambda) = \begin{pmatrix} \cos \theta & 0 & \sin \theta & 0 \\ 0 & 1 & 0 & \lambda \\ -\sin \theta & 0 & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$\hat{\gamma}(\theta, \lambda) = \begin{pmatrix} \cos \theta & \sin \theta & \lambda \cos \theta & \lambda \sin \theta \\ -\sin \theta & \cos \theta & -\lambda \sin \theta & \lambda \cos \theta \\ 0 & 0 & \cos \theta & \sin \theta \\ 0 & 0 & -\sin \theta & \cos \theta \end{pmatrix}.$$

Proposition 4.2.      For  $k \geq 5$ , we have

$$K(\pm \hat{\beta}(\frac{2\pi}{3}, \pm 1)) = \frac{1}{3^2} (1 - (\frac{-3}{p})) [0, -1, 1; 3] ,$$

$$K(\hat{\delta}(\pm 1, \pm 1)) = \frac{(-1)^k}{2^4} (1 - (\frac{-1}{p})),$$

$$K(\pm a) = - \frac{p-1}{2^4 3^2} (2k-3),$$

$$K(\pm b) = \frac{p-1}{2^3 3} ,$$

$$K(\hat{\gamma}(\frac{\pi}{2}, \pm 1)) = \frac{1}{2^3} (1 - (\frac{-1}{p})),$$

$$K(\pm \hat{\gamma}(\frac{2\pi}{3}, \pm 1)) = \frac{1}{2 \cdot 3} (1 - (\frac{-3}{p})),$$

and  $K(\gamma) = 0$  for any other  $\gamma \in \text{Sp}(2, \mathbb{R})$  which is not  
 $\text{Sp}(2, \mathbb{R})$ -conjugate to one of the above.

Proof is obvious by virtue of Th.4 and Hashimoto[6] Th.6.2.

Now, denote six non zero values in Prop.4.2. by

$$K_i (i=1, \dots, 6), \text{ that is, } K_1 = K(\pm \hat{\beta}(\frac{2\pi}{3}, \pm 1)),$$

and so on. Then, for  $k \geq 5$ , we have,

$$\begin{aligned}
 & \dim S_k(K(p)) - 2 \dim S_k(\text{Sp}(2, Z)) - \dim M_{k-3}(U_2) \\
 &= J_6 + J_7 + J_{12} + \sum_{i=1}^6 K_i \\
 &= - \left\{ \frac{p-1}{12} + \frac{1}{4} \left( 1 - \left( \frac{-1}{p} \right) \right) + \frac{1}{3} \left( 1 - \left( \frac{-3}{p} \right) \right) \right\} \\
 & \quad \times \left\{ \frac{k}{6} - \frac{1}{3} [0, -1, 1; 3] - \frac{1}{4} (3 + (-1)^k) \right\} \\
 &= -\dim A_2(\Gamma_0(p)) \times \dim S_{2k-2}(\text{SL}_2(Z)).
 \end{aligned}$$

So, we obtain Theorem 1.

Remark We get also the following interesting result.

Put  $\Gamma_0(p) = B(p) \cup B(p)w_1B(p)$ ,  $\Gamma'_0(p) = B(p) \cup B(p)w_2B(p)$ ,

and  $\Gamma''_0(p) = B(p) \cup B(p)w_0B(p)$ . When  $p = 2$ , the

dimensions of cusp forms belonging to these groups

are easily calculated by using Igusa [14] (II) (cf. [11]).

We get the following equality for  $k \geq 3$ :

$$\begin{aligned}
 & \dim S_k(B(2)) - \dim S_k(\Gamma_0(2)) - \dim S_k(\Gamma'_0(2)) - \dim S_k(\Gamma''_0(2)) \\
 &+ \dim S_k(K(2)) + 2 \dim S_k(\text{Sp}(2, Z)) \\
 &= \dim M_{k-3}(U_0) - \dim M_{k-3}(U_1) - \dim M_{k-3}(U_2) + \begin{cases} 1 \dots & \text{if } k = 3, \\ 0 \dots & \text{if } k \neq 3, \end{cases}
 \end{aligned}$$

where the discriminant of  $B$  is two. This supports

the conjecture in [9]. This relation is extended in [8]

for all  $p$ .

References

- [1] A.Borel and J.Tits, Groupes reductifs, Publ.Math. IHES 27(1965)55-150.
- [2] C.Chevalley, Sur certains idéaux d'une algèbre simple, Abh.Math.Sem.Univ.Hambrug, (1934)83-105.
- [3] M.Eichler, Über die darstellbarkeit von Modulformen durch Theta Reihen, J.Reine Angew.Math.195(1956) 159-171.
- [4] M.Eichler, Quadratische Formen und Modulformen, Acta arithm.4(1958)217-239.
- [5] K.Hashimoto, On Brandt matrices associated with the positive definite quaternion hermitian forms, J.Fac.Sci.Univ.Tokyo Sec.IA 27(1980)227-245.
- [6] K.Hashimoto, The dimension of the space of cusp forms of Siegel upper half plane of degree two (I) J.Fac.Sci.Univ.Tokyo Sec.IA 30(1983)403-488; (II) to appear in Math.Ann.
- [7] K.Hashimoto and T.Ibukiyama, On class numbers of positive definite binary quaternion hermitian forms, (I) J.Fac.Sci.Univ.Tokyo Sec.IA 27(1980)549-601; (II) *ibid.*28(1982)695-699; (III) *ibid.*30(1983)393-401.
- [8] K.Hashimoto and T.Ibukiyama, On relations of dimensions of automorphic forms of  $Sp(2, R)$  and its compact twist  $Sp(2)$  (II) to appear.
- [9] T.Ibukiyama, On symplectic Euler factors of genus two, J.Fac.Sci.Univ.Tokyo Sect.IA Math.30(1984). and Proc.Japan Acad.57 Ser.A no.5(1981)271-275.

- [10] T.Ibukiyama, On automorphic forms of  $Sp(2, R)$  and its compact form  $Sp(2)$ , to appear in Sémi.Delange-Fisou-Poitou 1982.
- [11] T.Ibukiyama, On the graded rings of Siegel modular forms of genus two belonging to certain level two congruence subgroups, preprint.
- [12] J.Igusa, On Siegel modular forms of genus two, Amer.J.Math.84(1962)175-200, (II) ibid.86(1964)392-412.
- [13] Y.Ihara, On certain arithmetical Dirichlet series, J.Math.Soc.Japan 16(1964)214-225.
- [14] H.Jacquet and R.P.Langlands, Automorphic forms on  $GL(2)$ , Lecture Notes in Math.260, Springer(1972).
- [15] R.P.Langlands, Problems in the theory of automorphic forms, Lecture Notes in Math.170, Springer(1970)18-61.
- [16] R.P.Langlands, Stable conjugacy: Definitions and Lemmas, Canad.J.Math.31(1979)700-725.
- [17] H.Shimizu, On zeta functions of quaternion algebras, Ann.Math.81(1965)166-193.
- [18] J.Tits, Reductive groups over local fields, Proc.Symp.Pure Math.XXXIII part 1(1979)29-69.

Department of Mathematics  
College of General Education  
Kyushu University  
Ropponmatsu, Fukuoka  
810 Japan

and

Max-Planck-Institut für Mathematik  
Gottfried-Claren Str.26  
5300 Bonn 3, BRD