

GLOBAL BOUNDS FOR THE BETTI NUMBERS  
OF REGULAR FIBERS OF DIFFERENTIABLE  
MAPPINGS

Y. Yomdin

83 33

Sonderforschungsbereich  
Theoretische Mathematik  
Berlingstr. 4  
D-5300 Bonn 1

Max-Planck-Institut  
für Mathematik  
Gottfried-Claren-Straße 26  
D-5300 Bonn 3

MPI/SFB 83-33

GLOBAL BOUNDS FOR THE BETTI NUMBERS OF REGULAR  
FIBERS OF DIFFERENTIABLE MAPPINGS

Y. Yomdin

INTRODUCTION

It is well known, that the Betti numbers of any fiber  $p^{-1}(\xi)$  of a polynomial mapping  $p : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , are bounded by some constants, depending only on  $n$ ,  $m$  and the degree of  $p$  ( see e.g. [7] ).

Now let  $f$  be a  $k$  times differentiable mapping of a bounded domain, with all the derivatives of order  $k$ , bounded by a constant  $M_k$ . We can think  $M_k$  as the measure of a deviation of  $f$  from a polynomial mapping of degree  $k-1$ ; as far as the deviation in a  $C^j$ -norm is concerned,  $j \leq k-1$ , the Taylor formula gives the precise expression for it.

The important general phenomenon is that also in much more delicate questions, concerning the topology and the geometry of the mapping  $f$ , its "deviation" from the "polynomial behavior" can be bounded in terms of  $M_k$ .

In [11] this fact was established for the structure of critical points and values of  $f$ , and in [12] for some geometric properties of its fibers.

The aim of the present paper is to extend in the same

spirit to  $k$ -smooth mappings the property of polynomial ones, given above: the boundness of the Betti numbers of the fibers.

Clearly, it is impossible to bound the Betti numbers of each fiber: any closed set can be the set of zeroes of a  $C^\infty$  - smooth function. So the proper way to generalize the above property of polynomials is the following:

First, we prove for any  $f$  the existence of fibers with the Betti numbers, bounded by constants, depending only on  $M_k$  ( and, of course, on  $k$  and on the dimensions and the size of the domain and the image of  $f$  ).

Secondly, we estimate, in the same terms, the integral complexity of the fibers of  $f$  . In particular, we answer the question, concerning the conditions of integrability of the Banach indicatrix of a differentiable mapping, which was open for a long time ( see [1],[2],[9] ).

All the inequalities below have the following form: they consist of a term, corresponding to the case of polynomials, and of a "correction term", containing the factor  $M_k$  . Thus, for  $M_k = 0$ , i.e. for  $f$  a polynomial of degree  $k-1$ , we obtain, up to constants, the usual bounds.

Another important remark concerns the existence results below: in many cases we prove the existence of at least one value  $\xi$  in the image of  $f$  , for which the Betti numbers of the fiber  $f^{-1}(\xi)$  are bounded by suitable constants. Although we do not touch in this paper the question of the explicit finding of such values, we should mention that the corresponding results can be brought to a rather effective form: for instance, we can prove that in any regular net

with a sufficiently small ( explicitly given ) step, there are points  $\xi$  with the required properties.

The author would like to thank the Max-Planck-Institut für Mathematik, where this paper was written, for its support and kind hospitality.

## §1. CONNECTION BETWEEN TOPOLOGY OF FIBERS AND GEOMETRY OF CRITICAL VALUES

Although all the results below remain valid, with minor modifications, for any compact manifold, we shall consider only the mappings, defined on a closed ball  $B_r^n$  of radius  $r$  in  $R^n$ . In this case all the constants involved can be given explicitly.

We say that the mapping  $f : B_r^n \rightarrow R^m$  is  $q$ -smooth, where  $q = p + \alpha$ ,  $p \geq 1$  - an integer,  $0 < \alpha \leq 1$ , if  $f$  is  $p$  times continuously differentiable on  $B_r^n$ , and the  $p$ -th derivative  $d^p f$  satisfy on  $B_r^n$  the Hölder condition:

$$(1) \quad \|d^p f(x) - d^p f(y)\| \leq L \|x - y\|^\alpha,$$

with some constant  $L$ .

$$\text{Let } M_i(f) = \max_{y \in B_r^n} \|d^i f(y)\|, \quad i = 0, 1, \dots, p,$$

$M_q(f) = \text{infimum of } L \text{ in (1), and let } R_j(f) = M_j(f) r^j, \\ j = 0, 1, \dots, p, q. \text{ ( All the Euclidean spaces } R^s \text{ and the spaces of their linear and multilinear mappings are considered with the usual Euclidean norm ).}$

We always assume below that  $n \geq m$ . Let  $\Sigma(f)$  be the set of critical points of  $f$ , i.e. of points  $x \in B_R^n$ , where  $\text{rank } df(x) < m$ , or, if  $x$  belongs to the boundary  $S_R^{n-1}$  of  $B_R^n$ ,  $\text{rank } d(f/S_R^{n-1}) < m$ . Let

$\Delta(f) = f(\Sigma(f)) \subset R^m$  be the set of critical values of  $f$ .

For  $\xi \in R^m$  we denote by  $Y_\xi$  the fiber  $f^{-1}(\xi)$  of  $f$  over  $\xi$ . If  $\xi$  is a regular value of  $f$ , i.e.  $\xi \notin \Delta(f)$ ,  $Y_\xi$  is a compact  $n-m$ -dimensional manifold. We denote by  $b_i(Y_\xi)$ ,  $i = 0, \dots, n-m$ , the  $i$ -th Betti number of  $Y_\xi$ .

Let  $\rho(\xi) = d(\xi, \Delta(f))$  be the distance from  $\xi$  to  $\Delta(f)$ .

**Theorem 1.1.** Let  $f : B_R^n \rightarrow R^m$  be a  $q$ -smooth mapping,  $q = p + \alpha$ . Then for any regular value  $\xi \in R^m$  of  $f$ , and  $i = 0, \dots, n-m$ ,

$$b_i(Y_\xi) \leq \begin{cases} B_i, & \rho(\xi) \geq R_q(f) \\ B_i (R_q(f)/\rho(\xi))^{n/q}, & \rho(\xi) \leq R_q(f) \end{cases}$$

where the constants  $B_i$ ,  $i = 0, \dots, n-m$ , depend only on  $n, m$  and  $p$ .

**Proof.** Below  $K_j$  denote the constants, depending only on  $n, m, p$ . We also omit sometimes the index  $f$  in the notations of  $\Delta(f)$ ,  $M_i(f)$  and  $R_i(f)$ .

Denote by  $B$  an open ball of radius  $\rho(\xi)$ , centered at the given regular value  $\xi \in R^m \setminus \Delta(f)$ . All the points  $\xi' \in B$  are the regular values both of  $f$  and of the restriction  $f/S_R^{n-1}$ . Hence  $f : N \rightarrow B$ , where  $N = f^{-1}(B)$ ,

is a trivial fibration, and, in particular, we can find a retraction  $\pi : N \rightarrow Y_\xi$ ,  $\pi|_{Y_\xi} = \text{Id}$ .

We shall construct a semialgebraic set  $S \subset N$ , containing  $Y_\xi$ , such that the Betti numbers of  $S$  satisfy inequalities of theorem 1.1. The existence of a retraction  $\pi : S \rightarrow Y_\xi$  then shows that the Betti numbers of  $Y_\xi$  do not exceed those of  $S$ .

For a given  $\delta > 0$  let  $I_{k_1 \dots k_n}^\delta$  be the cube  $\{x = (x_1, \dots, x_n) \in \mathbb{R}^n / k_j \delta \leq x_j \leq (k_j + 1)\delta, j = 1, \dots, n\}$ ,  $k_j \in \mathbb{Z}$ . Let  $I_\beta^\delta$ ,  $\beta = 1, \dots, K(\delta)$ , be those of the cubes  $I_{k_1 \dots k_n}^\delta$ , which intersect  $B_r^n$ . Clearly, for  $\delta \leq r$ ,  $K(\delta) \leq K_1 (r/\delta)^n$ .

For each  $\beta = 1, \dots, K(\delta)$ , take some point  $x_\beta \in I_\beta^\delta \cap B_r^n$ , and let  $P_\beta^\delta$  be the Taylor polynomial of degree  $p$  of  $f$  at  $x_\beta$ . By Taylor formula we have for each  $x \in I_\beta^\delta$ :  $\|f(x) - P_\beta^\delta(x)\| \leq K_2 M_q \delta^q$ .

Now take  $\delta = \min(r, (\rho(\xi)/4K_2 M_q)^{1/q})$  and let

$$S_\beta = \{x \in I_\beta^\delta \cap B_r^n / \|P_\beta^\delta(x) - \xi\| \leq \frac{1}{2}\rho(\xi)\}, \quad S = \bigcup_{1 \leq \beta \leq K(\delta)} S_\beta.$$

$S$  is a semialgebraic set and we have  $Y_\xi \subset S \subset N$ .

Indeed, by the choice of  $\delta$ ,  $\|f(x) - P_\beta^\delta(x)\| \leq \frac{1}{4}\rho(\xi)$  for

$x \in I_\beta^\delta$ . Hence, if  $x \in Y_\xi \cap I_\beta^\delta$ ,  $\|P_\beta^\delta(x) - \xi\| =$

$$\|P_\beta^\delta(x) - f(x)\| \leq \frac{1}{4}\rho(\xi), \quad \text{i.e. } x \in S_\beta \subset S.$$

Conversly, for  $x \in S_\beta$ ,  $\|f(x) - \xi\| \leq \|f(x) - P_\beta^\delta(x)\| + \|P_\beta^\delta(x) - \xi\| \leq \frac{1}{4}\rho(\xi) + \frac{1}{2}\rho(\xi) < \rho(\xi)$ . i.e.  $x \in N$ .

It remains to estimate the Betti numbers of  $S$ . Each  $S_\beta$  is defined by polynomial inequalities, whose number depend only on  $n$ , and whose degrees do not exceed  $2p$ .

The same is true for any nonempty intersection of  $S_\beta$  (which occurs only if the corresponding cubes  $I_\beta^\delta$  are adjoint). Hence for each  $i = 0, \dots, n-m$ ,

$$b_i(S_{\beta_1} \cap \dots \cap S_{\beta_j}) \leq B'_i, \text{ where the constants } B'_i$$

depend only on  $n, m$  and  $p$  (Some explicit estimates of  $B'_i$  one can find by methods of [6],[7],[8] or [10]).

Using the Mayer - Vietoris sequence, we obtain immediately, that  $b_i(S) \leq B'_i K_3 K(\delta) \leq B'_i K_3 K_1 (r/\delta)^n = B'_i K_3 K_1 (4K_2)^{n/q} (M_q r^q / \rho(\xi))^{n/q} = B'_i (R_q / \rho(\xi))^{n/q}$ .

These computations are valid for  $\rho(\xi) \leq R_q \leq 4K_2 R_q$ , since in this case  $r \geq (\rho(\xi) / 4K_2 M_q)^{1/q}$ , and we take  $\delta$  equal to the last number. But for  $\rho(\xi) > R_q$  we can restrict our consideration to the ball of radius  $R_q$  at  $\xi$ . Theorem 1.1 is proved.

Easy examples show that the bound of theorem 1.1 is sharp, up to constants.

## §2. EXISTENCE OF FIBERS WITH SIMPLE TOPOLOGY

In this section we combine the result of theorem 1.1 with the information on the geometry of critical values of  $f$ ,

obtained in [11] .

For a  $q$  - smooth  $f : B_r^n \rightarrow R^m$  define  $R_{1q}(f)$  as follows:  $R_{1q}(f) = 2(V_m A (R_1(f) + R_q(f))^{m-1} R_q(f))^{1/m}$  , where  $V_m$  is the volume of the unit ball in  $R^m$  , and  $A = A(n,m,p)$  , depending only on  $n, m, p$  , is the maximum of the constants  $\bar{A}_i(n,m,p)$  ,  $i = 0, \dots, m$  , defined in theorem 1.1, [11] .

Denote  $n-m+1$  by  $s$  . Below we assume that the smoothness  $q$  of  $f$  is greater than  $s$  , and hence, by the Sard theorem, almost all values of  $f$  are regular.

**Theorem 2.1.** Let  $f : B_r^n \rightarrow R^m$  be a  $q$  - smooth mapping,  $q > s$  . Then in any set  $G \subset R^m$  there is a regular value  $\xi$  of  $f$  , such that for  $i = 0, \dots, n-m$ ,

$$b_i(Y_\xi) \leq \begin{cases} B_i & , & m(G) \geq R_{1q}^m(f) \\ B_i (R_{1q}^m(f)/m(G))^{n/q-s} & , & m(G) \leq R_{1q}^m(f) \end{cases} ,$$

where  $m(G)$  denotes the Lebesgue measure of  $G$  .

**Proof.** Let  $G \subset R^m$  with  $m(G) = \eta > 0$  be given. According to theorem 1.1, it is sufficient to find a point  $\xi \in G$  , which is "far away" from  $\Delta(f)$ .

We shall use theorem 1.1, [11], which gives an upper bound for the  $\epsilon$  - entropy of  $\Delta(f)$  ( see [2] ), or, which is the same, for the minimal number  $M(\epsilon, \Delta(f))$  of balls of a given radius  $\epsilon > 0$  , covering  $\Delta(f)$ . The following form of this bound, which can be deduced easily from the original general one, is appropriate for our case:



For any  $\epsilon \leq R_q(f)$ ,

$$(2) \quad M(\epsilon, \Delta(f)) \leq A (1/\epsilon)^{m-1} (R_q(f)/\epsilon)^{s/q} (R_1(f) + R_q(f))$$

Now let  $\epsilon > 0$ ,  $\epsilon \leq R_q$ , be fixed. Cover  $\Delta$  by  $M(\epsilon, \Delta)$  balls of radius  $\epsilon$ , and let  $\Omega_\epsilon$  be the union of open balls of radius  $2\epsilon$ , centered at the same points.  $\Omega_\epsilon$  contains an  $\epsilon$ -neighborhood of  $\Delta$ , and hence for any  $\xi \in R^m \setminus \Omega_\epsilon$ ,  $d(\xi, \Delta) \geq \epsilon$  and by theorem 1.1,

$$b_1(Y_\xi) \leq B_1 (R_q/\epsilon)^{n/q}.$$

Denote by  $C_1(t)$  the set of points  $\xi \in R^m$ , for which  $b_1(Y_\xi) > t$ . We obtain  $C_1(B_1 (R_q/\epsilon)^{n/q}) \subset \Omega_\epsilon$ ,

for  $\epsilon \leq R_q$ , or  $C_1(t) \subset \Omega_{\epsilon(t)}$ , where  $\epsilon(t) = R_q (B_1/t)^{q/n}$ ,  $t \geq B_1$ .

By (2) for the measure of  $\Omega_\epsilon$  we have:

$$m(\Omega_\epsilon) \leq V_m \epsilon^m M(\epsilon, \Delta) \leq V_m A (\epsilon/R_q)^{1-s/q} (R_1+R_q)^{m-1} R_q, \text{ or}$$

$$(3) \quad m(\Omega_\epsilon) \leq R_{1q}^m (\epsilon/R_q)^{1-s/q}.$$

Substituting here the value of  $\epsilon(t)$  as above, we obtain the following:

Proposition 2.2.

$$m(C_1(t)) \leq \begin{cases} V_m R_1^m, & 0 \leq t < B_1 \\ R_{1q}^m (B_1/t)^{\frac{q-s}{n}}, & t \geq B_1 \end{cases}$$

The first inequality here means simply, that  $b_1(Y) > 0$

only for  $\xi \in f(B_r^n)$ , and  $f(B_r^n)$  clearly is contained in a ball of radius  $R_1$ .

Now if  $m(C_1(t)) < \eta = m(G)$ , then  $G$  contains some points  $\xi \notin C_1(t)$ , i.e. with  $b_1(Y_\xi) \leq t$ . It remains to note that by proposition 2.2,  $m(C_1(t)) < \eta$  is satisfied for  $t = B_1$ , if  $\eta > R_{1q}^m$ , and for any

$t > B_1 (R_{1q}^m / \eta)^{\frac{n}{q-s}}$ , for  $\eta \leq R_{1q}^m$ . Theorem 2.1 is proved.

Notice that the use of the  $\epsilon$ -entropy of critical values instead of the Lebesgue or the Hausdorff measure, and, respectively, the use of the stronger theorem 1.1 [11] instead of the Sard theorem, is the crucial point here: no bounds on the measure of  $\Delta(f)$  allow to find points "far away" from this set.

The fiber  $Y_\xi$  in theorem 2.1 can occur to be empty, for instance, if all the points of  $B_r^n$  are critical for  $f$ . Now we consider situations, where nonempty fibers with simple topological structure can be found.

Corollary 2.3. Let  $f : B_r^n \rightarrow R^m$  be  $q$ -smooth,  $q > s$ , and let  $m(f(B_r^n)) = \eta > 0$ . Then there exists a nonempty fiber  $Y_\xi$  of  $f$  with

$$b_1(Y_\xi) \leq \begin{cases} B_1, & \eta \geq R_{1q}^m(f) \\ B_1 (R_{1q}^m(f) / \eta)^{\frac{n}{q-s}}, & \eta \leq R_{1q}^m(f) \end{cases} .$$

Especially simple form these inequalities have in the case  $m = 1$ :

Corollary 2.4. Let  $f : B_R^n \rightarrow \mathbb{R}$  be a  $q$ -smooth function,  $q > n$ . Then in any set  $G \subset \mathbb{R}$  with  $m(G) > 0$ , there is a point  $c$  with

$$b_i(Y_c) \leq \begin{cases} B_1, & m(G) \geq 4AR_q(f) \\ B_1 (4AR_q(f)/m(G))^{\frac{n}{q-n}}, & m(G) \leq 4AR_q(f) \end{cases}$$

In particular, for  $a = \min f$ ,  $b = \max f$ , there is  $c$ ,  $a < c < b$ , such that

$$b_i(Y_c) \leq \begin{cases} B_1, & b-a \geq 4AR_q(f) \\ B_1 (4AR_q(f)/(b-a))^{\frac{n}{q-n}}, & b-a \leq 4AR_q(f) \end{cases}$$

Let us formulate separately one important special case:

Corollary 2.5. Let  $f : B_R^n \rightarrow \mathbb{R}$  be a  $q$ -smooth function,  $q > n$ , and let  $\max f - \min f \geq 4AR_q(f)$ . Then there exists  $c$ ,  $\min f < c < \max f$ , such that  $b_i(Y_c) \leq B_1$  where the constant  $B_1$  depends only on  $n$  and  $p$ .

This corollary can be interpreted as the appearance of a "near-polynomiality" effect: if  $f$  is sufficiently close to a polynomial, in the sense that  $R_q(f)$  is sufficiently small with respect to  $\max f - \min f$ , then the Betti numbers of at least one nonempty fiber of  $f$  satisfy exactly the same kind of inequalities as the Betti numbers of the polynomial fibers.

It is interesting to compare this fact with the result of [12], which indicates another appearance of the same effect: if for a  $q$ -smooth  $f$ ,  $\max f - \min f \geq$

$q!2^q R_q(f)$ , then any fiber  $Y_c$  of  $f$  is similar to the fibers of a polynomial of degree  $p$  in the following sense:  $Y_c$  is contained in a countable union of compact smooth hypersurfaces in  $R^n$ , "many" straight lines cross  $Y_c$  in at most  $p$  points, and the  $n-1$  - volume  $v(Y_c)$  is bounded by  $Kr^{n-1}$ , where  $K$  depends only on  $n$  and  $p$ . However, easy examples show, that the Betti numbers of some fibers of  $f$  can be infinite.

The inequality of theorem 2.1 is rather precise. In example 1, §6, VI, [2], for any  $n$  and  $q > n$  the function  $f : B_1^n \rightarrow R$  is built with the following properties:

- i.  $f$  is  $q$  - smooth.
- ii. For any  $\eta > 0$  there is an interval  $I_\eta \subset R$  of length  $\eta$ , such that for any  $c \in I_\eta$ ,  $b_i(Y_c) \geq K(1/\eta)^{n/q}$ ,  $i = 0, \dots, n-1$ .

Hence the degree of  $1/m(G)$  in the bounds for  $b_i$  cannot be smaller than  $n/q$ . Our value  $\frac{n}{q-s}$  is "asymptotically" sharp, for  $q \rightarrow \infty$ .

Theorem 2.1 implies also the following fact: if there is at least one point  $x \in B_r^n$ , where the rank of  $df(x)$  is maximal (equal to  $m$ ), then the Betti numbers of some nonempty fibers of  $f$  can be effectively bounded. As usual in our "quantitative" approach, we must not simply assume the nondegeneracy of the differential of  $f$ , but to measure the degree of this nondegeneracy.

Let for a linear mapping  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ ,  $\omega(L)$  be the the minimal semiaxis of the ellipsoid  $L(B_1^n) \subset \mathbb{R}^m$ . For a smooth  $f : B_r^n \rightarrow \mathbb{R}^m$  define  $\gamma(f)$  as  $r \max_{x \in B_r^n} \omega(df(x))$ .

We also denote by  $R_{12q}(f)$  the constant  $\sqrt{20}(1/V_m)^{1/2m} (R_{1q}(f)R_2(f))$

To simplify the expressions below, we assume, that  $\gamma(f) \leq R_2(f)$ .

**Theorem 2.6.** Let  $f : B_r^n \rightarrow \mathbb{R}^m$  be a  $q$ -smooth mapping,  $q > s$ , with  $0 < \gamma(f) \leq R_2(f)$ . Then there exists a nonempty fiber  $Y_\xi$  of  $f$  with

$$b_i(Y_\xi) \leq \begin{cases} B_i, & \gamma(f) \geq R_{12q}(f) \\ B_i (R_{12q}(f)/\gamma(f))^{\frac{2mn}{q-s}}, & \gamma(f) \leq R_{12q}(f) \end{cases}$$

**Proof.** Fix some  $x \in B_r^n$  with  $\omega = \omega(df(x))$  maximal. Now let  $P$  be the  $m$ -dimensional plane through  $x$ , for which  $\omega(df(x)/P) = \omega$ .

Easy estimates, repeating the proof of the inverse function theorem, show that the ball of radius  $\omega/3M_2$  in  $P$  (or the part of this ball, containing in  $B_r^n$ ) is mapped by  $f$  diffeomorphically, and its image contains the ball of radius  $\omega^2/20M_2 = \gamma(f)^2/20R_2(f)$ . Hence  $m(f(B_r^n)) \geq V_m [\gamma(f)^2/20R_2(f)]^m$ . Substituting this value in the inequality of corollary 2.3, we obtain the required result.

Studying in more detail the structure of  $f$  in the case when  $\text{rank } df < m$  everywhere, one can prove the existence of a nonempty fiber of  $f$  with the Betti numbers, bounded

by the constants, depending only on  $R_q(f)$  and the diameter of the image  $f(B_r^n)$ , for any sufficiently smooth mapping  $f : B_r^n \rightarrow R^m$ , with no assumptions of nondegeneracy. This proof requires considerations, somewhat different from the ones used in this paper, and it will appear separately.

### §3. INTEGRAL COMPLEXITY OF THE FIBERS

In this section we give the bounds for the integrals of  $b_i(Y_\xi)$ , when  $\xi$  runs over  $R^m$ .

**Theorem 3.1.** Let  $f : B_r^n \rightarrow R^m$  be a  $q$ -smooth mapping,  $q > s$ , and let  $\nu > 0$  be given. Then for  $i = 0, \dots, n-m$ ,

$$\int_{R^m} b_i^{\nu}(Y_\xi) d\xi \leq B_i^{\nu} [V_m R_1^m(f) + R_{1q}^m(f) \int_1^{\infty} (1/t)^{\frac{q-s}{\nu}} dt] .$$

**Proof.** By the Fubini theorem,  $\int_{R^m} b_i^{\nu}(Y_\xi) d\xi = \int_0^{\infty} m(C_i(t^{1/\nu})) dt$ ,

and by proposition 2.2, the last integral is bounded by

$$\int_0^{B_i^{\nu}} V_m R_1^m dt + \int_{B_i^{\nu}}^{\infty} R_{1q}^m (B_i/t^{1/\nu})^{\frac{q-s}{\nu}} dt = B_i^{\nu} V_m R_1^m + B_i^{\nu} R_{1q}^m \int_1^{\infty} (1/t')^{\frac{q-s}{\nu}} dt' .$$

Theorem is proved.

Theorem 3.1, in particular, answers the following question, which sometimes is called the question of integrability of the Banach indicatrix: for given  $n \geq m$  and  $\nu > 0$  to find  $q(n,m,\nu)$  such that for any  $q$ -smooth mapping  $f : R^n \rightarrow R^m$  with compact support,  $q > q(n,m,\nu)$ ,

$$\int_{R^m} b_0^u(Y_\xi) d\xi < \infty \quad (\text{and, in particular, to prove the}$$

existence of such a  $q(n,m,u)$  ).

Some special cases have been settled: the case  $m = u = 1$ ,  $n$  arbitrary - in [9], the case  $m = 1$ ,  $n$  and  $u$  arbitrary - in [2], the cases  $u = 1$ ,  $n \geq m$  arbitrary and  $n = m$ ,  $u$  arbitrary - in [1] .

Theorem 3.1 implies immediately the following:

Corollary 3.2. For  $f : B_r^n \rightarrow R^m$  - a  $q$ -smooth mapping,  $q > s$ , and for a given  $u$  ,  $0 \leq u < \frac{q-s}{n}$  ,

$$\int_{R^m} b_i^u(Y_\xi) d\xi \leq B_i^u [V_m R_1^m(f) + R_{1q}^m(f) \frac{nu}{q-nu-s}] < \infty .$$

In particular,  $q(n,m,u) \leq un + s = (u+1)n - m + 1$  .

Examples of [2] show, that  $q(n,m,u) \geq un$ , so our bound for  $q(n,m,u)$  is sharp asymptotically, for  $u \rightarrow \infty$  .

#### §4. VOLUME OF THE FIBERS

In this section, using the results of §3 , we study the distribution of the volume of regular fibers of  $f$ . Here it is convenient first to obtain the integral bounds, and then to deduce the existence of fibers with "small" volume.

Let for  $\xi$  a regular value of  $f : B_r^n \rightarrow R^m$  ,  $v(Y_\xi)$  denote the  $n-m$  - dimensional volume of the compact  $n-m$  - dimensional submanifold  $Y_\xi$  in  $R^n$ .

**Theorem 4.1.** Let  $f : B_r^n \rightarrow R^m$  be a  $q$ -smooth mapping,  $u \geq 1$ . Assume that  $q > mu + 1$ . Then

$$\int_{R^m} [v(Y_\xi)]^u d\xi \leq B_0^u C^u r^{(n-m)u} [V_m R_1^m(f) + R_{1q}^m(f) \frac{mu}{q-mu-1}] < \infty,$$

where the constant  $C$  depend only on  $n$  and  $m$ .

**Proof.** By the standard integral-geometric formula,

$$v(Y_\xi) = \int_{G_n^m} b_0(Y_\xi \cap L) dL, \text{ where } G_n^m \text{ is the space}$$

of all the  $m$ -dimensional planes in  $R^n$  with the standard measure  $dL$ .  $b_0(Y_\xi \cap L)$  here for almost all  $L$  is simply the number of points in  $Y_\xi \cap L$ .

The integration above runs, in fact, only over the set  $H \subset G_n^m$  of planes  $L$ , intersecting the ball  $B_r^n$ , and the measure of  $H$  in  $G_n^m$  is equal to  $Cr^{n-m}$ , where  $C$  depends only on  $n$  and  $m$ .

$$\text{Hence } \int_{R^m} [v(Y_\xi)]^u d\xi = \int_{R^m} d\xi \left[ \int_H b_0(Y_\xi \cap L) dL \right]^u.$$

By the Hölder inequality,

$$\int_H b_0(Y_\xi \cap L) dL \leq \left( \int_H [b_0(Y_\xi \cap L)]^u dL \right)^{1/u} \left( \int_H 1 dL \right)^{1/u'},$$

where  $u' = \frac{u}{u-1}$ . Hence

$$\left[ \int_H b_0(Y_\xi \cap L) dL \right]^u \leq C^{u-1} r^{(n-m)(u-1)} \int_H [b_0(Y_\xi \cap L)]^u dL,$$



and by the Fubini theorem

$$\int_{R^m} [v(Y_\xi)]^u d\xi \leq C^{u-1} r^{(n-m)(u-1)} \int_H dL \int_{R^m} [b_0(Y_\xi \cap L)]^u d\xi$$

Now since  $L \cap B_r^n$  is the ball of radius  $\leq r$  in  $L \cong R^m$  and since all the derivatives of the restriction  $f/L$  do not exceed those of  $f$ , we have by corollary 3.2:

$$\int_{R^m} [b_0(Y_\xi \cap L)]^u d\xi \leq B_0^u [V_m R_1^m + R_1^m \frac{mu}{q-mu-1}] , \text{ and}$$

$$\int_{R^m} [v(Y_\xi)]^u d\xi \leq C^u r^{(n-m)u} B_0^u [V_m R_1^m + R_1^m \frac{mu}{q-mu-1}] .$$

Theorem 4.1 is proved.

The question of integrability of  $v(Y_\xi)^u$  was also studied for a long time: for  $n = 2$ ,  $m = 1$  it was settled in [4], and in a general case in [5]. However, our estimate of maximal  $u$ , for which the integral  $\int_{R^m} [v(Y_\xi)]^u d\xi$  converges, namely,  $u < \frac{q-1}{m}$ , is very close to the best possible,  $u \leq q/m$ , and is approximately twice better, than the Merkov's one [5]:  $u < q/2m + 1$ .

Using the integral bound of theorem 4.1, we can obtain the existence of regular fibers with the "small" volume:

**Theorem 4.2.** Let  $f : B_r^n \rightarrow R^m$  be a  $q$ -smooth mapping,  $q > m+1$ . Then for any  $\beta < \frac{q-1}{m}$ , there is a constant  $K$ , depending only on  $R_1(f)$ ,  $R_q(f)$ ,  $\beta$ ,  $n$ ,  $m$  and  $p$ , such

that in any  $G \subset \mathbb{R}^m$  there is  $\xi$  with

$$v(Y_\xi) \leq K(1/m(G))^{1/\beta}.$$

Proof. It follows immediately from the inequality of theorem 4.1, if we put

$$K = C r^{n-m} B_0 [V_m R_1^m + R_{1q}^m \frac{m\beta}{q-m\beta-1}]^{1/\beta}.$$

The results of this section include the situations, where the smoothness  $q$  of the mapping  $f$  is smaller than  $s = n-m+1$ . In these cases all the values of  $f$  may be critical and, respectively, all the fibers  $Y_\xi$  of  $f$  may not be the regular  $n-m$ -dimensional manifolds. Here we must understand  $v(Y_\xi)$  as the  $n-m$ -dimensional Hausdorff measure.

## §5. SOME INEQUALITIES BETWEEN THE DERIVATIVES OF $f$

In this section we show that all the constants in the inequalities above can be expressed in terms of the only two parameters of the mapping  $f : B_r^n \rightarrow \mathbb{R}^m$ : the remainder term  $R_q(f)$  and the diameter  $R_0(f)$  of the image  $f(B_r^n) \subset \mathbb{R}^m$ .

Proposition 5.1. There are constants  $N_j$ ,  $j = 1, \dots, p$ , depending only on  $n$ ,  $m$  and  $p$ , such that for any  $q = p+\alpha$ -smooth mapping  $f : B_r^n \rightarrow \mathbb{R}^m$ ,

$$R_j(f) \leq N_j (R_0(f) + R_q(f)), \quad j = 1, \dots, p.$$

Proof. For any polynomial mapping  $h : B_r^n \rightarrow \mathbb{R}^m$  of degree

$p$  the following Markov inequality is satisfied ( see e.g. [3] ):

$$(*) \quad R_j(h) \leq N_j^! R_0(h) \quad , \quad j = 1, \dots, p .$$

Now let  $h$  be the Taylor polynomial of  $f$  at the center of  $B_r^n$  . The Taylor formula shows that

$$(**) \quad R_j(h) - N_j^! R_q(f) \leq R_j(f) \leq R_j(h) + N_j^! R_q(f) \quad , \\ j = 1, \dots, p .$$

Combining  $(*)$  and  $(**)$ , we obtain the required inequalities.

**Corollary 5.2.** There is a constant  $D$ , depending only on  $n, m, p$ , such that for any  $q = p + \alpha$  - smooth mapping  $f : B_r^n \rightarrow R^m$

a) If  $R_0(f) \geq R_q(f)$  , then

$$R_{1q}(f) \leq D [R_0^{m-1}(f) R_q(f)]^{1/m} \quad ,$$

$$R_{12q}(f) \leq D [R_0^{2m-1}(f) R_q(f)]^{1/2m} \quad .$$

b) If  $R_0(f) \leq R_q(f)$  , then

$$R_{1q}(f) \leq D R_q(f) \quad ,$$

$$R_{12q}(f) \leq D R_q(f) \quad .$$

## REFERENCES

1. S.A.Gulevič, On the necessary condition of integrability of a Banach indicatrix of a differentiable mapping, *Izvestia AN SSSR, Ser. Mat.*, 47, 5 (1983), 1114 - 1134 .
2. L.D.Ivanov, *Variacii mnozestv i funkzii*, Nauka, 1975.
3. O.D.Kellogg, On bounded polynomials in several variables, *Math.Zeitschrift*, 27 (1928), 55 - 64 .
4. E.M.Landis, On the length of level curves of a function of two variables, *Dokl. AN SSSR*, 79, 3 (1951), 393 - 395 .
5. A.B.Merkov, On some properties of smooth mappings, *Vestnik mosk. Univ. Ser. Mat.* 2 (1979), 63 - 69 .
6. J.Milnor, On the Betti numbers of real varieties, *Proc. Amer.Math.Soc.* 15 (1964), 275 - 280 .
7. I.G.Petrovskii, O.A.Oleinik, On the topology of real algebraic surfaces, *Izvestia AN SSSR, Ser. Mat.*, 13 (1949), 389 -402.
8. R.Thom, Sur l'homologie des varietes algebriques reelles, In: *Differential and combinatorial topology. A symposium in honour of Marston Morse*, Princeton Univ. Press, 1965, 255 - 265 .
9. A.G.Vitushkin, *O mnogomernyh variaciah*, Gostehisdat, 1955 .
10. H.Whitney, Elementary structure of real algebraic varieties, *Ann,Math.*, 66 (1957), 545 - 556 .
11. Y.Yomdin, The geometry of critical and near-critical values of differentiable mappings, *Math.Ann.*, 264 (1983), 495 - 515 .

12. Y.Yomdin, The set of zeroes of an "almost polynomial" functions, to appear in the Proc.Amer.Math.Soc.

Max-Planck-Institut für Mathematik, Gottfried-Claren-Strasse 26  
D-5300 Bonn 3, Federal Republic of Germany

and

Department of Mathematics, Ben Gurion University of the Negev,  
Beer-Sheva 84120, Israel .