

REIDEMEISTER SPECTRUM FOR METABELIAN GROUPS OF THE FORM $Q^n \rtimes \mathbb{Z}$ AND $\mathbb{Z}[1/p]^n \rtimes \mathbb{Z}$, p PRIME.

ALEXANDER FEL'SHTYN AND DACIBERG L. GONÇALVES

ABSTRACT. In this note we study the Reidemeister spectrum for metabelian groups of the form $Q^n \rtimes \mathbb{Z}$ and $\mathbb{Z}[1/p]^n \rtimes \mathbb{Z}$. Particular attention is given to the R_∞ property of a subfamily of these groups.

1. INTRODUCTION

Let $\phi : G \rightarrow G$ be an automorphism of a group G . A class of equivalence $x \sim gx\phi(g^{-1})$ is called the *Reidemeister class* (or the ϕ -conjugacy class or the *twisted conjugacy class of ϕ*). The number of Reidemeister classes, denoted by $R(\phi)$, is called the *Reidemeister number* of ϕ . The interest in twisted conjugacy relations has its origins, in particular, in the Nielsen-Reidemeister fixed point theory (see [23, 4]), in Selberg theory (see [28, 1]), and Algebraic Geometry (see [20]). A current important problem of the field concerns obtaining a twisted analogue of the Burnside-Frobenius theorem [7, 4, 11, 12, 31, 10, 9], that is to show the coincidence of the Reidemeister number of ϕ and the number of fixed points of the induced homeomorphism of an appropriate dual object. One step in this process is to describe the class of groups G , such that $R(\phi) = \infty$ for any automorphism $\phi : G \rightarrow G$.

The work of discovering which groups belong to the mentioned class of groups was begun by Fel'shtyn and Hill in [7]. Later, it was shown by various authors that the following groups belong to this class: (1) non-elementary Gromov hyperbolic groups [5, 26] (2) Baumslag-Solitar groups $BS(m, n) = \langle a, b | ba^m b^{-1} = a^n \rangle$ except for $BS(1, 1)$ [6], (3) generalized Baumslag-Solitar groups, that is, finitely generated groups which act on a tree with all edge and vertex stabilizers infinite cyclic [25], (4) lamplighter groups $\mathbb{Z}_n \wr \mathbb{Z}$ iff

Date: August 21, 2009.

1991 Mathematics Subject Classification. 20E45;37C25; 55M20.

Key words and phrases. Reidemeister spectrum, twisted conjugacy classes, metabelian groups, divisible groups, R_∞ property.

$2|n$ or $3|n$ [19], (5) the solvable generalization Γ of $BS(1, n)$ given by the short exact sequence $1 \rightarrow \mathbb{Z} \left[\frac{1}{n} \right] \rightarrow \Gamma \rightarrow \mathbb{Z}^k \rightarrow 1$ as well as any group quasi-isometric to Γ [30], groups which are quasi-isometric to $BS(1, n)$ [29] (while this property is not a quasi-isometry invariant), (6) the R. Thompson group F [2], (7) saturated weakly branch groups (including the Grigorchuk group and the Gupta-Sidki group) [13], (8) mapping class groups, symplectic groups and braids groups [14], (9) relatively hyperbolic groups [15], (10) some classes of finitely generated free nilpotent groups [18, 27] and some classes of finitely generated free solvable groups [24], (11) some classes of crystallographic groups [3].

The paper [30] suggests a terminology for this property, which we would like to follow. Namely, a group G has *property* R_∞ (or is an R_∞ group) if all of its automorphisms ϕ have $R(\phi) = \infty$.

For the immediate consequences of the R_∞ property for the topological fixed point theory see, e.g., [29].

Following [24], we define the *Reidemeister spectrum of a group* G , denoted by $Spec(G)$, as the set of natural numbers k such that there is an automorphism $\phi \in Aut(G)$ with $R(\phi) = k$ (k can be infinite). In terms of the spectrum, the R_∞ -property of the group G simply means that $Spec(G)$ contains only one element which is the infinity.

It is easy to see that 1) $Spec(\mathbb{Z}) = \{2\} \cup \{\infty\}$, 2) for $n \geq 2$ the spectrum is full, i.e. $Spec(\mathbb{Z}^n) = \mathbb{N} \cup \{\infty\}$. Let $N = N_{rl}$ be the free nilpotent group of rank r and class l . Then for N_{22} (also known as discrete Heisenberg group) $Spec(N_{22}) = 2\mathbb{N} \cup \{\infty\}$ [21, 8, 24]. It is also known that $Spec(N_{23}) = \{2k^2 | k \in \mathbb{N}\} \cup \{\infty\}$ [24] and $Spec(N_{32}) = \{2n-1 | n \in \mathbb{N}\} \cup \{4n | n \in \mathbb{N}\} \cup \{\infty\}$ [24].

Let $X = L(m, q_1, \dots, q_r)$ be a generalized lens space and $f : X \rightarrow X$ a continuous map of degree d , where $|d| \neq 1$. Let $f_* : \pi_1(X) \rightarrow \pi_1(X)$ be induced homomorphism on the fundamental group $\pi_1(X) = \mathbb{Z}/m\mathbb{Z}$.

In 1943 Franz [16] has observed that $N(f) = R(f) = R(f_*) = \#Coker(1 - f_*) = \#(\mathbb{Z}/m\mathbb{Z})/(1 - d)(\mathbb{Z}/m\mathbb{Z}) = (1 - d, m)$, where $N(f)$ and $R(f)$ are Nielsen and Reidemeister numbers of the map f and where $(1 - d, m)$ denote the gcd of $1 - d$ and m . This gives a strong arithmetical restriction on the Reidemeister spectrum for endomorphisms of the group $\mathbb{Z}/m\mathbb{Z}$. We

observe that the knowledge of the Reidemeister spectrum of a group can be quite useful for fixed point theory.

It is interesting open problem to investigate $Spec(X) = \{N(f) | f \in C^0(X)\}$ - the *Nielsen spectrum of the space X* .

In this paper we study the Reidemeister spectrum and R_∞ -property for a subfamily of the family of the metabelian groups of the form $\mathbb{Q}^n \rtimes \mathbb{Z}$ and $\mathbb{Z}[1/p]^n \rtimes \mathbb{Z}$ for p a prime.

Acknowledgement. We would like to thank the Max-Planck-Institut für Mathematik (MPIM) in Bonn for its kind support and hospitality while the greater part of this work was completed.

2. PRELIMINARIES

In this section we show that for certain short exact sequences the kernel is characteristic. Then we compute $Aut(\mathbb{Q})$, $Aut(\mathbb{Z}[1/p])$ and the Reidemeister spectrum of the groups \mathbb{Q} and $\mathbb{Z}[1/p]$, where \mathbb{Q} denote the rational numbers and p a prime.

Let us consider a short exact sequence of the form $1 \rightarrow \mathbb{K} \rightarrow G \rightarrow \mathbb{Z} \rightarrow 1$, which of course splits.

Lemma 2.1. *Suppose the group \mathbb{K} has the property that for every $x \in \mathbb{K}$ there is a natural number $s > 1$ such that x is divisible by s (i.e. there is $y \in \mathbb{K}$ such that $y^s = x$). Then \mathbb{K} is characteristic in G .*

Proof. Let $\phi : G \rightarrow G$ be an automorphism. We know that $G \approx K \rtimes_\theta \mathbb{Z}$. Let $x \in \mathbb{K}$ and $\phi(x) = (z, r)$. We are going to show that $r = 0$. It follows from the definition of the operation on the semi-direct product that if s divides x then s also divides $\phi(x) = (z, r)$. Again by the definition of the operation on the semi-direct product s divides r . From the hypothesis it follows that there is an infinite sequence of positive integers numbers such that for each integer s of the sequence, r is divisible by this integer. Therefore, r has an infinite number of divisors and must be zero. \square

For the rational numbers we have:

Lemma 2.2. *The group of automorphisms of \mathbb{Q} is isomorphic to the multiplicative group of the rationals different from zero denoted by \mathbb{Q}^* .*

Proof. First, let us observe that an automorphism $\phi : \mathbb{Q} \rightarrow \mathbb{Q}$ is determined if we know the value of it at 1. For this let us consider an arbitrary element $p/q \in \mathbb{Q}$. Then $\phi(p/q) = p\phi(1/q)$ and it suffices to determine $\phi(1/q)$. Since \mathbb{Q} is torsion free follows that the divisibility in \mathbb{Q} is unique then $\phi(1/q)$ is uniquely determined. Conversely a multiplication by any rational number different from zero provides an automorphism, since multiplication by the inverse number provides the inverse homomorphism and the result follows. \square

Now we consider the group $\mathbb{Z}[1/p]$ for p a prime. We have a following similar result.

Lemma 2.3. *The group $\text{Aut}(\mathbb{Z}[1/p])$ is isomorphic to the multiplicative group of the elements of $\mathbb{Z}[1/p]$ generated by $\{\pm p\}$. This group is isomorphic $\mathbb{Z} + \mathbb{Z}_2$.*

Proof. The first part is similar to the proof of the previous Lemma. Namely, a homomorphism ϕ is determined by the value of the homomorphism at 1, and it is multiplication by this number. In order to have ϕ an automorphism then we need that $\phi(1)$ is invertible. Let $r/s \in \mathbb{Z}[1/p]$ where r/s is written in the reduced form. If $r/s \in \mathbb{Z}[1/p]$ then we have either $r = 1$ and $s = p^t$ or $s = 1$ and $r = p^t$ and the result follows. \square

Now we determine the Reidemeister spectrum of Q and $\mathbb{Z}[1/p]$. We need a Lemma which is going to be used also for the computation of the Reidemeister spectrum of other groups. Given $x \in \mathbb{Z}[1/p]$ it can be written uniquely in the form $\pm q/p^n$ where $n \in \mathbb{Z}$, $q \in \mathbb{N}$ (the natural numbers) and p, q relatively prime. Denote this number q by $v_P(x)$. Here P is the set of all primes relatively prime with p . This number will be also used later.

Lemma 2.4. *Let $\psi : A^n \rightarrow A^n$ be a homomorphism where A is either \mathbb{Q} or $\mathbb{Z}[1/p]$. Then ψ is invertible if and only if the determinant of the matrix of ψ is invertible. Furthermore, for $A = \mathbb{Q}$ the cardinality of the cokernel of ψ is infinite if $\det(\psi) = 0$, and it is 1 if $\det(\psi) \neq 0$. For $A = \mathbb{Z}[1/p]$ the cardinality of the cokernel of ψ is the natural number $v_P(x)$, defined above, for $x = \det(\psi)$, if $x \neq 0$. If $\det(\psi) = 0$ then the cardinality of the cokernel is infinite.*

Proof. That ψ is invertible if and only if $\det(\psi)$ is invertible is a classical fact since A is a commutative ring. Therefore if $A = \mathbb{Q}$ then $\det(\psi) \neq 0$ implies ψ invertible and we have the cardinality of the cokernel 1. If $\det(\psi) = 0$ then the result follows promptly from the fact that \mathbb{Q} is an infinite field.

Let $A = \mathbb{Z}[1/p]$ and we consider first the case where the homomorphism is injective. The matrix M of ψ have entries in A . Multiplication by p on A^n is an isomorphism as well multiplication by p^l for any l . So the cokernel of ψ and the cokernel of the composite of ψ with multiplication of p^l are isomorphic. For a sufficient large l we can assume that the matrix of the composite has entries of the form xp^i for x an integer (possibly negative) relatively prime with p and $i \geq 0$. Observe that the determinant of the new matrix M_1 is $p^l \det(\psi)$ and v_P of the two determinants are the same. Now consider the homomorphism $\psi_1 : \mathbb{Z}^n \rightarrow \mathbb{Z}^n$ defined by M_1 and $\psi_2 : A^n \rightarrow A^n$ the homomorphism defined by the same matrix.

Then we have the following commutative diagram:

$$\begin{array}{ccccccccc}
 1 & \longrightarrow & \mathbb{Z}^n & \xrightarrow{\psi_1} & \mathbb{Z}^n & \xrightarrow{\pi_1} & \text{coker}(\psi_1) & \longrightarrow & 1 \\
 & & \downarrow \iota & & \downarrow \iota & & \downarrow & & \\
 1 & \longrightarrow & \mathbb{Z}[1/p]^n & \xrightarrow{\psi_2} & \mathbb{Z}[1/p]^n & \xrightarrow{\pi_2} & \text{coker}(\psi_2) & \longrightarrow & 1
 \end{array}$$

where $\text{coker}(\psi_2)$ has no p torsion, as result of the p -divisibility of the group $\mathbb{Z}[1/p]$. After we take the tensor product with $\mathbb{Z}[1/p]$ over \mathbb{Z} the two first vertical homomorphisms becomes the identity. So we have an isomorphism between the tensor product of the cokernels. The first cokernel, since it is a finite abelian group is the direct product of two finite groups where the first has order a power of p and the other has order relatively prime to p . After take the tensor product we obtain only the finite subgroup of order relatively prime to p which is simultaneously the order of the coker of ψ_2 and $v_P(\det(\psi_1))$. But $v_P(\det(\psi_1)) = v_P(\det(\psi_2)) = v_P(\det(\psi))$ and the result follows.

For the second part if the homomorphism is not injective we have similar sequence without assume that the first horizontal homomorphisms are injective. Then the rest of the proof is similar to the proof above. Because the $\text{coker}(\psi_1)$ is infinite we obtain that $\text{coker}(\psi_2)$ is also infinite. The case when $A = \mathbb{Q}$ is simpler and we leave to the reader. \square

Proposition 2.5. a) The Reidemeister spectrum of \mathbb{Q} is

$$\text{Spec}(\mathbb{Q}) = \{1\} \cup \{\infty\}.$$

b) The Reidemeister spectrum of $\mathbb{Z}[1/p]$ for p an odd prime is $\text{Spec}(\mathbb{Z}[1/p]) = \{p^l + 1, p^{l+1} - 1, l \geq 0\} \cup \{\infty\}$

$$\text{and } \text{Spec}(\mathbb{Z}[1/2]) = \{2^l + 1, 2^l - 1, l \geq 1\} \cup \{\infty\}$$

Proof. Part a)- Given an automorphism $\phi : \mathbb{Q} \rightarrow \mathbb{Q}$ we know that it is multiplication by a rational number r . If $r = 1$ then the cokernel of multiplication of $r - 1 = 0$ is \mathbb{Q} so we obtain Reidemeister number infinite. Otherwise we have multiplication by $r - 1 \neq 0$ which is surjective. Therefore the Reidemeister number is 1 and the result follows.

Let us consider $\mathbb{Z}[1/2]$ and $P = \mathbb{P} - \{2\}$. An automorphism is multiplication by a rational number r such that $v_P(r)$ is 1. So we have $r = \pm 2^l$ $l \in \mathbb{Z}$. If $r = 1$ then we obtain that the Reidemeister number is infinite, and for $r = -1$ we obtain multiplication by -2 so we obtain Reidemeister number 1. So let $l \neq 0$. Then the numbers $2^l \pm 1$ are always odd and the result follows. The case p an odd prime is similar and simpler. We leave to the reader. \square

3. THE SEMI-DIRECT PRODUCT $\mathbb{Q}^n \rtimes \mathbb{Z}$, $\mathbb{Z}[1/p]^n \rtimes \mathbb{Z}$, $n \leq 2$

We begin by recall some basic facts. Given any automorphism ϕ of one of the groups $\mathbb{Q}^n \rtimes \mathbb{Z}$, $\mathbb{Z}[1/p]^n \rtimes \mathbb{Z}$, we know from section 2 that the subgroup either \mathbb{Q}^n or $\mathbb{Z}[1/p]^n$ is characteristic. So we obtain a homomorphism of short exact sequence. Whenever the induced homomorphism $\bar{\phi}$ on the quotient is the identity then by well known facts, see [17] implies that the Reidemeister number is infinite. Also from [17] in the case where $\bar{\phi} : \mathbb{Z} \rightarrow \mathbb{Z}$ is multiplication by -1(the only other possibility) then the Reidemeister number is computed as the sum of the Reidemeister number of ϕ' and the Reidemeister number of $\theta(1) \circ \phi'$. We will use the above procedure for the calculation which follows.

3.1. The case $n = 1$. In this subsection we have an action $\theta : \mathbb{Z} \rightarrow A$ where A is either \mathbb{Q} or $\mathbb{Z}[1/p]$. The homomorphism θ is completely determined by $\theta(1)$ which, in turn is determined by its values at $1 \in A$. So we identify $\theta(1)$ with its value at $1 \in A$.

Proposition 3.1. a) The Reidemeister spectrum of $\mathbb{Q} \rtimes_{\theta} \mathbb{Z}$ is $Spec(\mathbb{Q} \rtimes_{\theta} \mathbb{Z}) = \{\infty\}$ if $\theta(1)$ is a non zero rational number different from ± 1 and $Spec(\mathbb{Q} \rtimes \mathbb{Z}) = \{2\} \cup \{\infty\}$ otherwise.

b) For p an odd prime the Reidemeister spectrum of $\mathbb{Z}[1/p] \rtimes_{\theta} \mathbb{Z}$ is $Spec(\mathbb{Z}[1/p] \rtimes_{\theta} \mathbb{Z}) = \{\infty\}$ if $\theta(1) \in \mathbb{Z}[1/p]$ is a non zero invertible element different from ± 1 . If $r = 1$ then $Spec(\mathbb{Z}[1/p] \rtimes_{\theta} \mathbb{Z}) = \{p^l + 1, p^{l+1} - 1, l \geq 0\} \cup \{\infty\}$. If $r = -1$ then $Spec(\mathbb{Z}[1/p] \rtimes_{\theta} \mathbb{Z}) = \{2p^{l+1}, l \geq 0\} \cup \{\infty\}$ otherwise.

c) The Reidemeister spectrum of $\mathbb{Z}[1/2] \rtimes_{\theta} \mathbb{Z}$ is $Spec(\mathbb{Z}[1/2] \rtimes_{\theta} \mathbb{Z}) = \{\infty\}$ if $\theta(1) \in \mathbb{Z}[1/2]$ is a non zero invertible element different from ± 1 . If $\theta(1) = 1$ then $Spec(\mathbb{Z}[1/2] \rtimes_{\theta} \mathbb{Z}) = \{2(2^l + 1), 2(2^l - 1), l \geq 1\} \cup \{\infty\}$ and if $\theta(1) = -1$ then $Spec(\mathbb{Z}[1/2] \rtimes_{\theta} \mathbb{Z}) = \{2^{l+1}, l \geq 1\} \cup \{\infty\}$.

Proof. Let $\phi : \mathbb{Q} \rtimes_{\theta} \mathbb{Z}$ be a automorphism. From the discussion in the begin of the section we know that ∞ belongs to the spectrum and we will look at automorphisms such that $\bar{\phi}$ is multiplication by -1 . We compute the automorphisms $\phi' : \mathbb{Q} \rightarrow \mathbb{Q}$ which arises as restriction of such automorphisms. In order to an automorphism ϕ' be the restriction of an automorphism of the big group we must have the relation $\phi' \circ \theta(1) = \theta(-1) \circ \phi'$. This implies that $kr = r^{-1}k$ where $\phi'(1) = k$ (so different from 0) and $\theta(1) = r$. This implies that $r^2 = 1$. So if $r \neq \pm 1$ then there is no such automorphism and the Reidemeister spectrum of the group is $\{\infty\}$. If $r = \pm 1$ then k can assume any non zero value. If $r = 1$ then the two automorphisms on the fibers are multiplication by k and each one has Reidemeister number 1 if $k \neq 1$. If $r = -1$ then the two automorphisms on the fibers are multiplication by $k, -k$ respectively. So for $k \neq \pm 1$ the Reidemeister number is 2 and part a) follows.

Part b) The infinite certainly belongs to the spectrum because the Reidemeister number of the identity is infinite. The element $\theta(1)$ is invertible since it is an isomorphism. As in case a) we have that if $\theta(1) \neq \pm 1$ there is no such automorphism and follows that the group has spectrum $\{\infty\}$. Again as in part a) for $\theta(1) = \pm 1$ we have ϕ' an arbitrary automorphism of $\mathbb{Z}[1/p]$. For $r = 1$ the two automorphisms are the same and we obtain as Reidemeister number $2(p^l \pm 1)$ $l > 0$ and 4. If $r = -1$ then we have to look at $k - 1$ and

$k + 1$ for k an invertible, so of the form ϵp^t . If $t = 0$ then we get Reidemeister infinite. If $t \neq 0$ then we get as Reidemeister number $2p^l$. So the result follows.

Part c). The proof is similar to the proof of case b) where the only difference is because $2^0 + 1$ is not a Reidemeister number of a homomorphism for case c) but $p^0 + 1$ it is for case b). This justifies why the formulas are slight different. \square

3.2. The case $n = 2$. First we analyze the case where the kernel is \mathbb{Q}^2 . Let $G = \mathbb{Q}^2 \rtimes \mathbb{Z}$ and for an automorphism $\phi : G \rightarrow G$ let ϕ' be the restriction of ϕ to \mathbb{Q}^2 . If M is the matrix of ϕ' then we call $\det(\phi')$ the determinant of M .

Proposition 3.2. The Reidemeister spectrum of $\mathbb{Q}^2 \rtimes_{\theta} \mathbb{Z}$ is either $\{\infty\}$ or $\{2\} \cup \{\infty\}$.

The $\text{Spec}(\mathbb{Q}^2 \rtimes_{\theta} \mathbb{Z}) = \{\infty\}$ if there is no an invertible matrix N over \mathbb{Q} such that $N = MNM$ and $\det(Id - N)$ and $\det(Id - MN)$ are non zero. Here $\theta(1) = M$. Otherwise the Reidemeister spectrum is $\{2\} \cup \{\infty\}$.

Proof. From the considerations on the begin of this section and basic fact about Reidemeister number for homomorphisms of \mathbb{Q}^2 the result follows. \square

Examples-Now we show a family of groups which have Reidemeister spectrum $\{\infty\}$ and also a family of groups which have spectrum $\{2\} \cup \{\infty\}$.

Example 1- Let $\theta(1) = M$ be the automorphism given by

$$\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}$$

$r^2 \neq 1$, $rs \neq 0$ and 1 . There is no automorphism such that the induced homomorphism on the quotient is multiplication by -1 . This can be proven by showing that the only solution for the matrix N which satisfies $N = MNM$ is N the trivial matrix.

Example 2- Let $\theta(1)$ be the automorphism given by

$$\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}$$

$r^2 \neq 1$, $rs = 1$. By direct calculation we can find all matrices N and they are of the form

$$\begin{pmatrix} 0 & b \\ c & 0 \end{pmatrix}$$

for arbitrary b, c . The matrix MN is

$$\begin{pmatrix} 0 & rb \\ cs & 0 \end{pmatrix}$$

Then $\det(Id - N) = \det(Id - MN) = 1 - bc$ and whenever $1 - bc \neq 0$ we obtain an example where the Reidemeister number is 2. Therefore one example of a group where the spectrum is $\{2\} \cup \{\infty\}$.

Example 3- Let $\theta(1)$ be the automorphism given by

$$\begin{pmatrix} 0 & u \\ v & 0 \end{pmatrix}.$$

Let us consider two cases. Suppose that $|uv| \neq 1$. Then follows that there is no invertible N and follows that the Reidemeister number is infinite.

For the second case let $u = v = 1$ Then by direct calculation we can find all matrices N which are of the form

$$\begin{pmatrix} a & b \\ b & a \end{pmatrix}.$$

and the matrix MN is

$$\begin{pmatrix} b & a \\ a & b \end{pmatrix}.$$

The first matrix has determinant $a^2 - b^2$. The determinant of $Id - N$ and $Id - MN$ are respectively $1 - 2a + a^2 - b^2$ and $1 - 2b + b^2 - a^2$. There are plenty of rational values of a and b such that these 3 values are different from zero and consequently we obtain for each such values one example of a group which admits an automorphism which has Reidemeister number 2.

Now we consider the case $\mathbb{Z}[1/p]^2 \rtimes_{\theta} \mathbb{Z}$ for p a prime. We will compute the Reidemeister spectrum of $\mathbb{Z}[1/p]^2 \rtimes_{\theta} \mathbb{Z}$ for two families of action θ .

The Proposition 3.2 holds partially in this case.

Proposition 3.3. The Reidemeister spectrum of $\mathbb{Z}[1/p]^2 \rtimes_{\theta} \mathbb{Z}$ is $\{\infty\}$ if and only if there is no an invertible matrix N over $\mathbb{Z}[1/p]$ such that $N = MNM$ and $\det(Id - N)$ and $\det(Id - MN)$ are non zero. Here $\theta(1) = M$.

Proof. The proof is similat to the proof of Proposition 3.2. \square

The Reidemeister spectrum for the groups where there exist a solution N is not very simple in general but we can answer for two large families of actions θ . For example if $\theta(1)$ is the identity then the spectrum obtained by multiply by 2 the numbers(including infinite) which belong to the spectrum of the first factor. So the first step is to compute the Reidemeister spectrum of $\mathbb{Z}[1/p]^2$.

Proposition 3.4. The Reidemeister spectrum of $\mathbb{Z}[1/p]^2$ is $Spec(\mathbb{Z}[1/p]^2) = \{n|n \in \mathbb{N} \text{ and } (n, p) = 1\} \cup \{\infty\}$ where (n, p) denote the gcd of n and p .

Proof. Any matrix $N \in Gl(2, \mathbb{Z})$ can be regarded as an automorphism of $\mathbb{Z}[1/p]^2$. It is well known that the Reidemeister spectrum of $\mathbb{Z} + \mathbb{Z}$ is \mathbb{N} . So the Reidemeister spectrum of $\mathbb{Z}[1/p]^2$ contains $v_p(n)$ for every natural number by Lemma 2.4. So it contains all positive numbers relatively prime to p . But again by Lemma 2.4 an element of the spectrum has to be a positive integer relatively prime to p . So the result follows. \square

An immediate consequence of the Proposition above is that $Spec(\mathbb{Z}[1/p]^2 \times \mathbb{Z}) = \{2n|(n, p) = 1 \ n \in \mathbb{N}\} \cup \{\infty\}$ where (n, p) denote the gcd of n and p .

Let us start with the group $Spec(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z})$.

Proposition 3.5. Let $\theta(1)$ be of the form:

$$\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}$$

Then we have the following cases:

a) If $r = s = \pm 1$ then $Spec(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{2n|(n, 2) = 1 \ n \in \mathbb{N}\}$ where $(n, 2)$ denote the gcd of n and 2.

b) If $r = -s = \pm 1$ then the Reidemeister spectrum of the group $\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}$ is $Spec(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{2^{l+1}, 2^k(2^l \pm 1), \ l \geq 1, k \geq 2\} \cup \{\infty\}$.

c) If $rs = 1$ and $|r| \neq 1$ then $Spec(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{2(2^k + 1) \ k \geq 0, 2(2^k - 1) \ k \geq 1\} \cup \{\infty\}$

d) If either r or s does not have module equal to one, and $rs \neq 1$ then there is no automorphism of the group such that the induced on the quotient is multiplication by -1 and follows that $\text{Spec}(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{\infty\}$.

Proof. Let N be the matrix

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

The equation $N = MNM$ corresponds to the system $a = ar^2$, $b = brs$, $c = crs$ and $d = ds^2$. Suppose that $|r| \neq 1$ and $rs \neq 1$. This implies that $a = b = c = 0$ and so the system has no solution for a matrix N invertible. Similarly if we assume $|s| \neq 1$. So part d) follows.

For the part a) we have that all invertible matrices N are solutions. So the result follows from the Proposition 3.4 above.

For the part b) we have that the matrix N is diagonal. By straightforward calculation we can assume that the elements are of the form $\epsilon_1 2^i$ for $i \geq 0$. A direct calculation shows the values for the spectrum.

For the part c) we have that the matrix M is of the form

$$\begin{pmatrix} \epsilon 2^{\ell} & 0 \\ 0 & \epsilon 2^{-\ell} \end{pmatrix}.$$

and N of the form

$$\begin{pmatrix} 0 & \delta_1 2^u \\ \delta_2 2^v & 0 \end{pmatrix}.$$

The product MN is given by

$$\begin{pmatrix} 0 & \epsilon \delta_1 2^{u+\ell} \\ \epsilon \delta_2 2^{v-\ell} & 0 \end{pmatrix}.$$

Follows the matrices of $Id - N$ and $Id - MN$:

$$\begin{pmatrix} 1 & -\delta_1 2^u \\ -\delta_2 2^v & 0 \end{pmatrix}.$$

and

$$\begin{pmatrix} 1 & -\epsilon \delta_1 2^{u+\ell} \\ -\epsilon \delta_2 2^{v-\ell} & 1 \end{pmatrix}$$

respectively. The result follows by straightforward calculation.

□

Proposition 3.6. Let $\theta(1)$ be of the form:

$$\begin{pmatrix} 0 & u \\ v & 0 \end{pmatrix}$$

Then we have the following cases:

a) If $uv = 1$ then the Reidemeister spectrum of the group $\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}$ is $\text{Spec}(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{2^k(2^l \pm 1) \mid k \geq 2, l \geq 1\} \cup \{\infty\}$.

b) If $uv = -1$ then the Reidemeister spectrum of the group $\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}$ is $\text{Spec}(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{2(2^{2m} - 1) \mid m > 0\} \cup \{\infty\}$.

c) If $u^2v^2 \neq 1$ then there is no automorphism of the group such that the induced on the quotient is multiplication by -1 and follows that $\text{Spec}(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{\infty\}$.

Proof. In this case the entries of the matrix N

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

must satisfy the equations $uvd = a, u^2c = b, bv^2 = c, auv = d$. It follows that $a = (uv)^2a, d = (uv)^2d, b = (uv)^2b, c = (uv)^2c$. So part c) follows promptly from these equations.

Let us consider the case a). In this case the system of equations provide $a = d$. So the determinant of N becomes $a^2 - bc = a^2 - b^2v^2 = (a+bv)(a-bv)$ which is an invertible element, so $(a+bv)$ and $(a-bv)$ are also invertible. Since $uv = 1$ follows that $u = \delta 2^{-t}$ and $v = \delta 2^t$ for some integer t and $\delta \in \{1, -1\}$.

The matrix N is of the form

$$\begin{pmatrix} a & b \\ bv^2 & a \end{pmatrix}$$

and the matrix MN is

$$\begin{pmatrix} bv & a/v \\ av & bv \end{pmatrix}.$$

Our task is to compute v_P of $\det(Id - N)$ and $\det(Id - MN)$. We have $\det(Id - N) = 1 - 2a + a^2 - b^2v^2$ and $\det(Id - MN) = 1 - 2bv + b^2v^2 - a^2$ so $-\det(Id - MN) = -1 + 2bv + a^2 - b^2v^2$. We compute v_P of $-\det(Id - MN)$.

Since $a + bv$ and $a - bv$ are invertible they are of the form $a + bv = \epsilon_1 2^{l_1}$ and $a - bv = \epsilon_2 2^{l_2}$. Therefore $a = \epsilon_1 2^{l_1-1} + \epsilon_2 2^{l_2-1}$ and $b = \delta(\epsilon_1 2^{l_1-t-1} - \epsilon_2 2^{l_2-t-1})$. Also $\det(N) = a^2 - b^2v^2 = \epsilon_1 \epsilon_2 2^{l_1+l_2}$.

So we obtain $\det(Id - N) = \epsilon_1 \epsilon_2 2^{l_1+l_2} - \epsilon_1 2^{l_1} - \epsilon_2 2^{l_2} + 1$ and $-\det(Id - MN) = \epsilon_1 \epsilon_2 2^{l_1+l_2} + \epsilon_1 2^{l_1} - \epsilon_2 2^{l_2} - 1$. It is not difficult to see that the rational numbers, up to multiplication by a power of 2 (positive or negative) are the integers $2^{|l_1|+|l_2|} - \epsilon_2 2^{|l_1|} - \epsilon_1 2^{|l_2|} + 1$ and $2^{|l_1|+|l_2|} + \epsilon_2 2^{|l_1|} - \epsilon_1 2^{|l_2|} - 1$, respectively. Whenever these positive integers are odd they are the Reidemeister number of the correspondent matrices, which happens for $|l_1|, |l_2| > 0$. Then in this case the Reidemeister number is the sum equals to $2^{|l_2|+1}(2^{|l_1|} - \epsilon_1)$. In order to find the complete spectrum we have to analyze the particular cases. Suppose that $l_1 = l_2 = 0$. Then in this case we have four possibilities for the pair ϵ_1, ϵ_2 . By straightforward calculation for each case either $\det(Id - N)$ or $\det(Id - MN)$ is zero (if not both) and we obtain Reidemeister infinite. In details, for $(\epsilon_1, \epsilon_2) = (1, 1)$ then $(\det(Id - N), -\det(Id - MN)) = (0, 0)$, for $(\epsilon_1, \epsilon_2) = (1, -1)$ then $(\det(Id - N), -\det(Id - MN)) = (0, 0)$, for $(\epsilon_1, \epsilon_2) = (-1, 1)$ then $(\det(Id - N), -\det(Id - MN)) = (0, -4)$ for $(\epsilon_1, \epsilon_2) = (-1, -1)$ then $(\det(Id - N), -\det(Id - MN)) = (4, 0)$

Now let $l_2 = 0$ and $l_1 \neq 0$. By direct inspection for $\epsilon_2 = 1$ we obtain $\det(N) = 0$ and for $\epsilon_2 = -1$ we obtain $\det(MN) = 0$, hence we obtain Reidemeister infinite.

Finally let $l_1 = 0$ and $l_2 \neq 0$. If $\epsilon_1 = 1$ then $\det(Id - N) = \det(Id - MN) = 0$ and Reidemeister is infinite. If $\epsilon_1 = -1$ we get $\det(Id - N) = -2(\epsilon_2 2^{|l_2|} - 1)$ and $\det(Id - MN) = -2(\epsilon_2 2^{|l_2|} + 1)$. After we compute v_P of these numbers and add them up we obtain the Reidemeister number $2^{|l_2|+1}$ for $|l_2| \geq 1$ or 2^k $k \geq 2$. But these numbers were obtained already in previous cases and the result follows.

Let us consider the case b). Some of the calculations are similar and in this case we do not give all details. In this case the system of equations provide $a = -d$. So the determinant of N becomes $-a^2 - bc = -a^2 - b^2v^2 =$

which is an invertible element. Since $uv = -1$ follows that $u = -\delta 2^{-t}$ and $v = \delta 2^t$ for some integer t and $\delta \in \{1, -1\}$.

The matrix N is of the form

$$\begin{pmatrix} a & b \\ bv^2 & -a \end{pmatrix}$$

and the matrix MN is

$$\begin{pmatrix} -bv & a/v \\ av & bv \end{pmatrix}.$$

Our task is to commute v_P of $\det(Id - N)$ and $\det(Id - MN)$. We have $\det(Id - N) = 1 - a^2 - b^2v^2$ and $\det(Id - MN) = 1 - b^2v^2 - a^2$. So we compute v_P of $\det(Id - N)$.

Let $a = r2^m$ and $bv = s2^n$ where r, s are odd numbers (possible negative) and m, n integers. Because $a^2 + (bv)^2$ is invertible then we obtain that $r^2 + s^2$ is necessarily 1 and we obtain as possible solutions $a = \epsilon_1 2^m$ and $bv = 0$ (or $b = 0$ since $v \neq 0$) $a = 0$ and $bv = \epsilon_2 2^n$.

For the case $b = 0$ we obtain $\det(Id - N) = 1 - a^2 = 1 - 2^{2m}$. For $m = 0$ we obtain Reidemeister infinite otherwise we obtain the total Reidemeister number $2(2^{2m} - 1)$ for $m > 0$. For $a = 0$ we obtain $\det(Id - N) = 1 - b^2v^2 = 1 - 2^{2n}$. Then we obtain the same numbers as above and the result follows. \square

The example studied by Jabara in [22] is included in part a) above. Moreover, part a) above computes the Reidemeister spectrum such example.

For an arbitrary prime $p \neq 2$ we will have similar results.

Proposition 3.7. Let $\theta(1)$ be of the form:

$$\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}$$

Then we have the following cases:

- a) If $r = s = \pm 1$ then $\text{Spec}(\mathbb{Z}[1/p]^2 \rtimes_{\theta} \mathbb{Z}) = \{2n | n \in \mathbb{N}, (n, p) = 1\} \cup \{\infty\}$ where (n, p) denote the gcd of n and p .
- b) If $r = -s = \pm 1$ then the Reidemeister spectrum of the group $\mathbb{Z}[1/p]^2 \rtimes_{\theta} \mathbb{Z}$ is $\text{Spec}(\mathbb{Z}[1/p]^2 \rtimes_{\theta} \mathbb{Z}) = \{2p^l(p^k \pm 1), 4p^l | l, k > 0\} \cup \{\infty\}$
- c) If $rs = 1$ and $|r| \neq 1$ then $\text{Spec}(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{2(p^l \pm 1), 4 | l > 0\} \cup \{\infty\}$

d) If either r or s does not have module equal to one, and $rs \neq 1$ then there is no automorphism of the group such that the induced on the quotient is multiplication by -1 and follows that $\text{Spec}(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{\infty\}$.

Proof. As in Proposition 3.5 we have the system $a = ar^2$, $b = brs$, $c = crs$ and $d = ds^2$. Part a) and d) follows as in Proposition 3.5.

For the part b) from the equations $a = ar^2$, $b = brs$, $c = crs$ and $d = ds^2$ follows that the matrix N is diagonal. Let $a = \epsilon_1 p^{l_1}$ and $d = \epsilon_2 p^{l_2}$ since these elements are invertible. The $\det(Id - N) = 1 - a - d + ad = \epsilon_1 \epsilon_2 p^{l_1 + l_2} - \epsilon_1 p^{l_1} - \epsilon_2 p^{l_2} + 1$ and $-\det(Id - MN) = ad + a - d - 1 = \epsilon_1 \epsilon_2 p^{l_1 + l_2} + \epsilon_1 p^{l_1} - \epsilon_2 p^{l_2} - 1$.

Without loss of generality let us assume that $l_1, l_2 \geq 0$. First let $l_1 = l_2 = 0$. Then one of the two determinants is zero and the Reidemeister number is infinite. Now let $l_1 = 0$ and $l_2 \neq 0$. We have $\det(Id - N) = \epsilon_1 \epsilon_2 p^{l_2} - \epsilon_1 - \epsilon_2 p^{l_2} + 1$ and $-\det(Id - MN) = \epsilon_1 \epsilon_2 p^{l_2} + \epsilon_1 - \epsilon_2 p^{l_2} - 1$. If $\epsilon_1 = 1$ then $\det = 0$ in both cases and we have Reidemeister infinite. If $\epsilon_1 = -1$ then we obtain $\det(Id - N) = -2\epsilon_2 p^{l_2} + 2$ and $-\det(Id - MN) = -2\epsilon_2 p^{l_2} - 2$. Both numbers are not divisible by p and the Reidemeister number is the module of $4\epsilon_2 p^{l_2}$, $l_2 > 0$. Now let $l_2 = 0$ and $l_1 \neq 0$. We have $\det(Id - N) = \epsilon_1 \epsilon_2 p^{l_1} - \epsilon_2 - \epsilon_1 p^{l_1} + 1$ and $-\det(Id - MN) = \epsilon_1 \epsilon_2 p^{l_1} + \epsilon_1 p^{l_1} - \epsilon_2 - 1$. If $\epsilon_2 = 1$ then $\det(Id - N) = 0$ and we have Reidemeister infinite. If $\epsilon_2 = -1$ then $\det(Id - MN) = 0$ and we have Reidemeister infinite. Finally if $l_1, l_2 > 0$ then the two numbers $\epsilon_1 \epsilon_2 p^{l_1 + l_2} - \epsilon_1 p^{l_1} - \epsilon_2 p^{l_2} + 1$ and $\epsilon_1 \epsilon_2 p^{l_1 + l_2} + \epsilon_1 p^{l_1} - \epsilon_2 p^{l_2} - 1$ are not divisible by p and the Reidemeister number is the module of $2\epsilon_1 \epsilon_2 p^{l_1 + l_2} - 2\epsilon_2 p^{l_2} = 2\epsilon_2 p^{l_2}(\epsilon_1 p^{l_1} - 1)$. So the Reidemeister numbers are of the form $2p^l(p^k \pm 1)$, $l, k > 0$, and the result follows.

For the part c) from the equations $a = ar^2$, $b = brs$, $c = crs$ and $d = ds^2$ follows $a = d = 0$. So the matrix N is of the form

$$\begin{pmatrix} 0 & b \\ c & 0 \end{pmatrix}.$$

and MN is

$$\begin{pmatrix} 0 & rb \\ sc & 0 \end{pmatrix}.$$

By direct calculation, using that $rs = 1$, follows that $\det(Id - N) = \det(Id - MN) = 1 - bc$. Since bc is invertible then we can write $b = \epsilon_1 p^{l_1}$ and $c = \epsilon_2 p^{l_2}$ and follows that $\det(Id - N) = 1 - \epsilon_1 \epsilon_2 p^{l_1 + l_2}$. If $l_1 + l_2 = 0$ then we obtain for the determinant 0 or 2. So we obtain for Reidemeister number infinite and 4. If $l_1 + l_2 \neq 0$ then we obtain as Reidemeister numbers the numbers of the form $2(p^l \pm 1), l > 0$ and the result follows. \square

Proposition 3.8. Let $\theta(1)$ be of the form:

$$\begin{pmatrix} 0 & u \\ v & 0 \end{pmatrix}$$

Then we have the following cases:

a) If $uv = 1$ then the Reidemeister spectrum of the group $\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}$ is $\text{Spec}(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{2p^l(p^l \pm 1), l > 0\} \cup \{\infty\}$.

b) If $uv = -1$ and then the Reidemeister spectrum of the group $\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}$ is $\text{Spec}(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{2(p^l \pm 1), l > 0\} \cup \{\infty\}$.

c) If $u^2 v^2 \neq 1$ then there is no automorphism of the group such that the induced on the quotient is multiplication by -1 and follows that $\text{Spec}(\mathbb{Z}[1/2]^2 \rtimes_{\theta} \mathbb{Z}) = \{\infty\}$.

Proof. The proof follows the same steps as the proof of Proposition 3.6 and it is simpler. The matrix N

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

must satisfy the equations $uvd = a, u^2 c = b, bv^2 = c, auv = d$. It follows that $a = (uv)^2 a, d = (uv)^2 d, b = (uv)^2 b, c = (uv)^2 c$. So part c) follows promptly from these equations.

Let us consider the case a). In this case the system of equations provide $a = d$. So the determinant of N becomes $a^2 - bc = a^2 - b^2 v^2 = (a + bv)(a - bv)$ which is an invertible element, so $(a + bv)$ and $(a - bv)$ are also invertible. Since $uv = 1$ follows that $u = \delta p^{-t}$ and $v = \delta p^t$ for some integer t and $\delta \in \{1, -1\}$.

The matrix N is of the form

$$\begin{pmatrix} a & b \\ bv^2 & a \end{pmatrix}$$

and the matrix MN is

$$\begin{pmatrix} bv & a/v \\ av & bv \end{pmatrix}.$$

Our task is to compute v_P of $\det(Id - N)$ and $\det(Id - MN)$. We have $\det(Id - N) = 1 - 2a + a^2 - b^2v^2$ and $\det(Id - MN) = 1 - 2bv + b^2v^2 - a^2$ so $-\det(Id - MN) = -1 + 2bv + a^2 - b^2v^2$. We compute v_P of $-\det(Id - MN)$ which is the same as v_P of $\det(Id - MN)$.

Since $a + bv$ and $a - bv$ are invertible they are of the form $a + bv = \epsilon_1 p^{l_1}$ and $a - bv = \epsilon_2 p^{l_2}$. Therefore $2a = \epsilon_1 p^{l_1} + \epsilon_2 p^{l_2}$ and $2bv = \epsilon_1 p^{l_1} - \epsilon_2 p^{l_2}$. In order to have $\epsilon_1 p^{l_1} + \epsilon_2 p^{l_2}$ divisible by 2 we need to have $l_1 = l_2 = l$ and either $\epsilon_1 = \epsilon_2$ or $\epsilon_1 = -\epsilon_2$. In the first case we have $a = \epsilon_1 p^l$ and $2bv = 0$ so $b = 0$. In the latter case we have $a = 0$ and $bv = \epsilon_1 p^l$. Now we compute the Reidemeister for each of these two cases. Let $a = \epsilon_1 p^l$ and $b = 0$. Then $\det(Id - N) = 1 - 2a + a^2$ and $-\det(Id - MN) = a^2 - 1$. If $l = 0$ then $\det(Id - MN) = 0$ and we get Reidemeister infinite. If $l \neq 0$ then the determinants are not divisible by p and we get as Reidemister number the module of $2a^2 - 2a = 2a(a - 2) = 2\epsilon p^l(\epsilon p^l - 1)$. So the Reidemeister numbers are of the form $2p^l(p^l \pm 1)$, $l \neq 0$. So the result follows.

Let us consider the case b). Some of the calculations are similar to the corresponding case of the Proposition 3.6 and in this case we do not give the details. The system of equations provide $a = -d$. So the determinant of N becomes $-a^2 - bc = -a^2 - b^2v^2$ which is an invertible element. Since $uv = -1$ follows that $u = -\delta 2^{-t}$ and $v = \delta 2^t$ for some integer t and $\delta \in \{1, -1\}$.

The matrix N is of the form

$$\begin{pmatrix} a & b \\ bv^2 & -a \end{pmatrix}$$

and the matrix MN is

$$\begin{pmatrix} -bv & a/v \\ av & bv \end{pmatrix}.$$

Our task is to commute v_P of $\det(Id - N)$ and $\det(Id - MN)$. We have $\det(Id - N) = 1 - a^2 - b^2v^2$ and $\det(Id - MN) = 1 - b^2v^2 - a^2$. So we compute v_P of $\det(Id - N)$.

Let $a = rp^m$ and $bv = sp^n$ where r, s are relatively prime with p . Because $a^2 + (bv)^2$ is invertible then we obtain that $r^2 + s^2$ is necessarily 1 and we obtain as possible solution $a = \epsilon_1 p^m$ and $bv = 0$ (or $b = 0$ since $v \neq 0$), or $a = 0$ and $bv = \epsilon_2 p^n$.

For the case $b = 0$ we obtain $\det(Id - N) = 1 - a^2 = 1 - p^{2m}$. For $m = 0$ we obtain Reidemeister infinite otherwise we obtain the total Reidemeister number $2(p^{2m} - 1)m > 0$. For $a = 0$ we obtain $\det(Id - N) = 1 - b^2v^2 = 1 - p^{2n}$. Then we obtain the same numbers as above and the result follows. \square

4. FINAL COMMENTS ABOUT $Q^n \rtimes Z$, $\mathbb{Z}[1/p]^n \rtimes Z$, $n > 2$

There are some of the above results that extend easily to the groups $Q^n \rtimes Z$, $\mathbb{Z}[1/p]^n \rtimes Z$, $n > 2$. One of the results refer to the group $Q^n \rtimes Z$. The abelian group \mathbb{Q}^n has the property that it does not have a subgroup of finite index. Then an immediate consequence of this fact is that the Reidemeister number of any homomorphism is either 1 or infinite. So the following result holds.

Proposition 4.1. The Reidemeister spectrum of $\mathbb{Q}^n \rtimes_{\theta} \mathbb{Z}$ is either $\{\infty\}$ or $\{2\} \cup \{\infty\}$.

The $Spec(\mathbb{Q}^n \rtimes_{\theta} \mathbb{Z}) = \{\infty\}$ if there is no an invertible matrix N over \mathbb{Q} such that $N = MNM$ and $\det(Id - N)$ and $\det(Id - MN)$ are non zero. Here $\theta(1) = M$. Otherwise the Reidemeister spectrum is $\{2\} \cup \{\infty\}$.

Proof. (sketch) The only possible automorphism which can have Reidemeister finite is one such that the induced homomorphism on the quotient \mathbb{Z} is multiplication by -1 . From the procedure described in the begin of section 3 we have to compute the Reidemeister of the homomorphism given by N and MN . But a homomorphism of \mathbb{Q}^n is either surjective or has cokernel infinite. So the sum of the Reidemeister of the two homomorphisms is either infinite or 2. The case where both homomorphisms have Reidemeister 1 corresponds to say that $\det(Id - N)$ and $\det(Id - MN)$ are non zero and the result follows. \square

It is easy to construct examples which illustrated both situation.

Now let p be an arbitrary prime. It is not difficult to construct action $\theta(1)$ which has diagonal matrix such that there is no automorphism of the group such that the induced automorphisms on the quotient \mathbb{Z} is $-id$. This gives the examples of groups with the R_∞ property.

The complete calculation of the Reidemeister spectrum for all such groups is not clear for the moment.

REFERENCES

1. J. Arthur and L. Clozel, *Simple algebras, base change, and the advanced theory of the trace formula*, Princeton University Press, Princeton, NJ, 1989. MR90m:22041
2. C. Bleak, A. Fel'shtyn, and D. L. Gonçalves, Twisted conjugacy classes in R. Thompson's group F, *Pacific Journal of Mathematics*, no.1, v.238(2008), 1-6.
3. K. Dekimpe, P. Penninckx, The finiteness of the Reidemeister number of morphisms between almost-crystallographic groups, preprint, 2009.
4. A. Fel'shtyn, *Dynamical zeta functions, Nielsen theory and Reidemeister torsion*, Mem. Amer. Math. Soc. **147** (2000), no. 699, xii+146. MR2001a:37031
5. A. Fel'shtyn, *The Reidemeister number of any automorphism of a Gromov hyperbolic group is infinite*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **279** (2001), no. 6 (Geom. i Topol.), 229–240, 250. MR2002e:20081
6. A. Fel'shtyn and D. Gonçalves, *Reidemeister numbers of any automorphism of Baumslag-Solitar group, is infinite*, Geometry and Dynamics of Groups and Spaces, Progress in Mathematics, v. 265(2008), 286-306, Birkhauser. .
7. A. Fel'shtyn and R. Hill, *The Reidemeister zeta function with applications to Nielsen theory and a connection with Reidemeister torsion*, K-Theory **8** (1994), no. 4, 367–393. MR95h:57025
8. A. Fel'shtyn, F. Indukaev, and E. Troitsky, Twisted Burnside theorem for two- step torsion free nilpotent group. In: *C*-algebras and elliptic theory. II* Birkhäuser Trends in Math. series, 2008, 87–101.
9. A. Fel'shtyn and E. Troitsky, *Geometry of Reidemeister classes and twisted Burnside theorem*, Journal of K-theory **1**(2008), 1-40.
10. A. Fel'shtyn and E. Troitsky, Twisted Burnside - Frobenius theory for discrete groups, *J. Reine Angew. Math.* **613**(2007), 193-210.
11. A. Fel'shtyn, E. Troitsky, *A twisted Burnside theorem for countable groups and Reidemeister numbers*, Noncommutative Geometry and Number Theory (C. Consani and M. Marcolli, eds.), Vieweg, Braunschweig, 2006, pp. 141–154.
12. A. Fel'shtyn, E. Troitsky, and A. Vershik, *Twisted Burnside theorem for type II_1 groups: an example*, Math. Res. Lett. **13** (2006), no. 5, 719–728.

13. A. Fel'shtyn, Y. Leonov, E. Troitsky, Reidemeister numbers of saturated weakly branch groups. *Geometria Dedicata*, v.134(2008), 61-73.
14. Alexander Fel'shtyn and Daciberg L. Gonçalves, Twisted conjugacy classes in Symplectic groups, Mapping class groups and Braid groups(including an Appendix written with Francois Dahmani).E-print arxiv: math.GR/0708.2628.
15. Alexander Fel'shtyn, New directions in Nielsen-Reidemeister theory, E-print arXiv: math.GR/0712.2601.
16. W. Franz, Abbildungsklassen und fixpunktklassen dreidimensionaler Linsenräume. *J. Reine Angew. Math.* 185 (1943), 65–77.
17. Gonçalves D. L., The coincidence Reidemeister classes on nilmanifolds and nilpotent fibrations, *Top. and its Appl.* **83** (1998), 169–186.
18. D. Gonçalves and P. Wong , Twisted conjugacy classes in nilpotent groups, E-print arXiv: math.GR/0706.3425 .
19. D. Gonçalves and P. Wong, *Twisted conjugacy classes in wreath products*, International Journal of Algebra and Computation **16** No. 5 (2006), 875–886.
20. A. Grothendieck, *Formules de Nielsen-Wecken et de Lefschetz en géométrie algébrique*, Séminaire de Géométrie Algébrique du Bois-Marie 1965-66. SGA 5, Lecture Notes in Math., vol. 569, Springer-Verlag, Berlin, 1977, pp. 407–441.
21. F. K. Indukaev, *Twisted Burnside theory for the discrete Heisenberg group and wreath products of some groups*, (Russian) Vestn. Moscow Univ., Ser. 1, matem., mechan., 6 (2007), 9-17.
22. E. Jabara, *Automorphisms with finite Reidemeister number in residually finite groups*, *J. Algebra* **320** No. 10 (2008), 3671–3679.
23. B. Jiang, *Lectures on Nielsen fixed point theory*, Contemp. Math., vol. 14, Amer. Math. Soc., Providence, RI, 1983.
24. E. G. Kukina, V. Roma'nikov, On the Reidemeister spectrum and the R_∞ property for some free nilpotent groups, E-print arxiv:math.GR/0903.3455
25. G. Levitt, *On the automorphism group of generalised Baumslag-Solitar groups*, E-print arxiv:math.GR/0511083, 2005.
26. G. Levitt and M. Lustig, *Most automorphisms of a hyperbolic group have very simple dynamics.*, *Ann. Scient. Éc. Norm. Sup.* **33** (2000), 507–517.
27. V. Roma'nikov, Twisted conjugacy classes in nilpotent groups, E-print arxiv:math.GR/0903.3455
28. Salahoddin Shokranian, *The Selberg-Arthur trace formula*, Lecture Notes in Mathematics, vol. 1503, Springer-Verlag, Berlin, 1992, Based on lectures by James Arthur. MR1176101 (93j:11029)
29. Jennifer Taback and Peter Wong, *A note on twisted conjugacy and generalized Baumslag-Solitar groups*, E-print arXiv:math.GR/0606284, 2006.
30. Jennifer Taback and Peter Wong, *Twisted conjugacy and quasi-isometry invariance for generalized solvable Baumslag-Solitar groups*, E-print arxiv:math.GR/0601271, 2006.

31. E. Troitsky, *Noncommutative Riesz theorem and weak Burnside type theorem on twisted conjugacy*, *Funct. Anal. Pril.* **40** (2006), no. 2, 44–54, In Russian, English translation: *Funct. Anal. Appl.* **40** (2006), No. 2, 117–125 (Preprint 86 (2004), Max-Planck-Institut für Mathematik, math.OA/0606191).

INSTYTUT MATEMATYKI, UNIWERSYTET SZCZECIŃSKI, UL. WIELKOPOLSKA 15, 70-451 SZCZECIN, POLAND AND BOISE STATE UNIVERSITY, 1910 UNIVERSITY DRIVE, BOISE, IDAHO, 83725-155, USA

E-mail address: `felshtyn@diamond.boisestate.edu`, `felshtyn@mpim-bonn.mpg.de`

DEPT. DE MATEMÁTICA - IME - USP, CAIXA POSTAL 66.281 - CEP 05311-970, SÃO PAULO - SP, BRASIL

E-mail address: `dlgoncal@ime.usp.br`