

A CHARACTERIZATION OF COMPLEX MANIFOLDS
BIHOLOMORPHIC TO A CIRCULAR DOMAIN

by

Giorgio Patrizio

Sonderforschungsbereich 40
Theoretische Mathematik
Berlingstr. 4
D-5300 Bonn 1

Max-Planck-Institut
für Mathematik
Gottfried-Claren-Str.26
D-5300 Bonn 3

MPI/SFB 84-29

INTRODUCTION

W. Stoll [S], developing earlier ideas of Griffiths and King [GK], introduced in Value Distribution Theory the concept of parabolic exhaustion. This notion has recently found important applications in the classification of domains in \mathbb{C}^m and in general of complex manifolds. A strictly parabolic exhaustion on a non compact complex manifold M of dimension m is a C^∞ exhaustion function $\tau: M \rightarrow [0, \Delta^2)$, $0 < \Delta = \sup \sqrt{\tau} \leq \infty$, such that $dd^c \tau > 0$ on M and $dd^c \log \tau \geq 0$, $(dd^c \log \tau)^m \equiv 0$ on $M - \tau^{-1}(0)$. Using the information that τ yields and most noteworthy the existence of a Monge-Ampère foliation associated to τ (cfr. [BK]), Stoll in [SPM] shows that there exists a biholomorphic map $h: B(\Delta) \rightarrow M$, where $B(\Delta) = \{Z \in \mathbb{C}^m \mid \|Z\| < \Delta\}$, such that $\tau \circ h(Z) = \|Z\|^2$ (cfr. also [BU] and [W1] for other proofs). The properties of the exhaustion τ on $\tau^{-1}(0)$ play a decisive role in the classification. In fact, if we relax the differentiability assumptions on $\tau^{-1}(0)$, even assuming that this set reduces to a single point (which is always the case if τ is strictly parabolic), the range of possibilities for the underlying manifold is much wider. If $G \subset \subset \mathbb{C}^m$ is a smooth, strictly pseudoconvex, complete circular domain and σ is its Minkowski functional squared, then σ is strictly parabolic on $G - \{0\}$ but it is smooth on the origin if and only if G is biholomorphic to the ball. On the other hand the space of biholomorphic classes of such domains is infinite dimensional (cfr. [PW]). In this class of examples the Monge-Ampère foliation associated to the exhaustion is always holomorphic. This is also quite an exceptional

occurrence. Let $D \subset \subset \mathbb{C}^m$ be a smooth, strictly convex domain and $p \in D$ be any point. There exists an exhaustion $\tau_p: \bar{D} \rightarrow [0,1]$ such that $\tau_p^{-1}(0) = \{p\}$, $\tau|_{\partial D} \equiv 1$ and that τ_p is strictly parabolic on $D - \{p\}$ (cfr. [L1] and [P1]). For a generic strictly convex domain the Monge-Ampère foliation associated to τ_p is not holomorphic (cfr. [P1]).

In this paper we study complex manifolds M admitting an exhaustion $\tau: M \rightarrow [0, \Delta^2)$ which is strictly parabolic on $M - \tau^{-1}(0)$ and has a prescribed behavior on $\tau^{-1}(0)$. We require that there exists $p \in \tau^{-1}(0)$ such that, if $P: \tilde{M} \rightarrow M$ is the blow up of M at p , then $\tau \circ P \in C^\infty(\tilde{M})$ and such that, with respect to coordinates centered at p , we have $C \|z\|^2 \leq \tau(z) \leq K \|z\|^2$ for some positive constants C, K (cfr. Section 1 for detailed definitions). It is easy to see that this definition encompasses the above mentioned examples. Under these assumptions we are able to give an accurate description of the Monge-Ampère foliation associated to the exhaustion (cfr. Theorem 3.4 below). This is used to define, in analogy with our previous work [P1], a "model" (G, σ) for (M, τ) where G is a circular domain and σ , up to a factor, is its Minkowski functional squared. A homeomorphism $h: G \rightarrow M$ is constructed such that $\tau \circ h = \sigma$. We show that h is biholomorphic if and only if the Monge-Ampère foliation associated to τ is holomorphic. This, because of a theorem of Burns [Bu], is always true if $\Delta = \sup \sqrt{\tau} = \infty$. In this case M is biholomorphic to \mathbb{C}^m and τ pulls back to the Minkowski

functional squared of some circular domain (cfr. Theorem 4.4 for a precise statement).

The methods used here are a mixing of Stoll's technique ([SPM]) and of ideas developed in the special case of strictly convex domains ([P1]). The simplifications to Stoll's original proof due to Burns [Bu] and Wong [W1] cannot be used here because they are based on the consideration of the exponential map of the metric $dd^c \tau$ at the center point $\tau^{-1}(0)$ whereas in our situation the metric is not defined on $\tau^{-1}(0)$. It is interesting to note that we can define a "generalized" exponential map at the center which, however, is only homeomorphic. This map is smooth at the origin if and only if it is biholomorphic (which is the case if and only if τ is strictly parabolic on all M). Complex manifolds carrying an exhaustion which is strictly parabolic outside its zero set and whose associated Monge-Ampère foliation is holomorphic, have been classified also by Wong [W2]. His approach is based on the study of the differential geometry of the level sets of the exhaustion. His results do not overlap with ours but are clearly related.

Acknowledgements. I would like to thank here W. Stoll for sharing with me his ideas on this problem and I. Lieb for helpful discussions and remarks. I also gladly acknowledge the hospitality and the financial support of the Max-Planck-Institut für Mathematik and the SFB 40 Universität Bonn.

0. NOTATIONS

Let M be a complex manifold and let $T(M)$, $T^C(M)$ denote respectively the tangent bundle and the complexified tangent bundle of M . Then $T(M)$ is a subbundle of $T^C(M)$ and we have the splitting $T^C(M) = T^+(M) \oplus T^-(M)$ where $T^+(M), T^-(M)$ denote respectively the holomorphic and the antiholomorphic tangent bundle of M . The projections $\pi_+ : T^C(M) \rightarrow T^+(M)$ and $\pi_- : T^C(M) \rightarrow T^-(M)$ restrict to R -bundle isomorphisms $\pi_+ : T(M) \rightarrow T^+(M)$ and $\pi_- : T(M) \rightarrow T^-(M)$. To the complex structure of M is canonically associated an almost complex structure $J : T^C(M) \rightarrow T^C(M)$ which is a bundle isomorphism such that $J^2 = -\text{Id}$ and such that $J|_{T^+(M)}$ is the multiplication by i and $J|_{T^-(M)}$ is the multiplication by $-i$. By an hermitian metric k on M we intend here a C^∞ function $k : T^+(M) \oplus T^+(M) \rightarrow \mathbb{C}$ such that for all $p \in M$ its restriction $k_p : T_p^+(M) \oplus T_p^+(M) \rightarrow \mathbb{C}$ is a positive definite hermitian form. The canonical form associated to k is a positive $(1,1)$ form ω defined, for all $p \in M$ and $X, Y \in T_p^+(M)$, by $\omega_p(X, \bar{Y}) = (i/2\pi)k_p(X, Y)$. The metric k is Kähler if and only if $d\omega = 0$. Also to k is associated a Riemannian metric g defined for all $p \in M$ and $X, Y \in T(M)$, by $g_p(X, Y) = \text{Re } k_p(\pi_+(X), \pi_+(Y))$.

Given an exhaustion function $\tau : M \rightarrow [0, R^2)$, we shall use the following notations:

$$\begin{aligned}
 M[0] &= \{x \in M \mid \tau(x) = 0\} \\
 M(r) &= \{x \in M \mid \tau(x) < r^2\} \\
 (0.1) \quad M[r] &= \{x \in M \mid \tau(x) \leq r^2\} \\
 M\langle r \rangle &= \{x \in M \mid \tau(x) = r^2\} = M[r] - M(r) \\
 M_* &= \{x \in M \mid \tau(x) \neq 0\} = M - M[0] .
 \end{aligned}$$

In particular $\mathbb{E}(r)$, $\mathbb{E}[r]$ will always denote respectively the open and the closed disk of radius r in \mathbb{E} . For $m > 1$ we will denote the ball of radius r in \mathbb{E} by $\mathbb{B}(r)$ and write $\mathbb{B} = \mathbb{B}(1)$. The unit sphere in \mathbb{R}^n will be denoted by S^{n-1} or simply by S if the dimension is clear from the context. We will use the differential operators $d = \partial + \bar{\partial}$, $d^c = (i/4\pi)(\bar{\partial} - \partial)$ and $dd^c = (i/2\pi)\partial\bar{\partial}$. Finally, throughout the paper, upper indices will denote components and lower indices derivatives. Moreover, whenever possible, Einstein summation convention will be used. Also we shall use the Kroneker symbol $\delta_{\mu\nu}$.

1. DEFINITION AND PRELIMINARIES

Let M be a non compact complex manifold of dimension m and $\tau: M \rightarrow [0, \Delta^2)$, $0 < \Delta = \sup \sqrt{\tau} \leq \infty$, be an exhaustion function. We say the pair (M, τ) is a manifold of circular type if the following assumptions are satisfied:

(1.1) $\tau \in C^0(M) \cap C^\infty(M_*)$.

(1.2) $dd^c \tau > 0$ and $dd^c \log \tau \geq 0$ on M_* .

(1.3) $(dd^c \log \tau)^m = 0$ on M_* .

(1.4) There exists $p \in M[0]$ such that :

(i) with respect to local coordinates centered at p ,

if $\| \cdot \|$ denotes the euclidean norm, we have

$$C \|Z\|^2 \leq \tau(Z) \leq \|Z\|^2 \quad \text{for some positive constants}$$

$C, K;$

(ii) if $P: \tilde{M} \rightarrow M$ is the blow up of M at p , then

$$\tau \circ P \in C^\infty(\tilde{M}) .$$

In what follows we shall only use the following consequence of (1.4)

(1.4') There exists $p \in M[0]$ and a coordinate neighbourhood U_p , centered at p , such that:

(i) there exists positive constants C, K such that

$$C \|Z\|^2 \leq \tau(Z) \leq K \|Z\|^2 \quad \text{for all } Z \in U_p;$$

(ii) there exists $\varepsilon_0 > 0$ so that $tZ \in U_p$ if $|t| < \varepsilon_0$

and $\|Z\| < 2$ and such that the map $h: (-\varepsilon_0, \varepsilon_0) \times B(2) - \{0\} \rightarrow$

defined by $h(t, Z) = \tau(tZ)$ is of class C^∞ .

As it will turn out from Theorem 3.4, (1.4') is in fact equivalent to (1.4) if (1.1), (1.2), (1.3) are fulfilled. We also say that $\Delta = \sup \sqrt{\tau}$ is the radius of (M, τ) .

It is useful to bare in mind some concrete example from the very beginning. In fact, the basic motivation for this paper was to try to generalize Example 1 according to the lines suggested by Example 2 and 3.

Example 1. Stoll defines in [SPM] a strictly parabolic manifold to be a pair (M, τ) where M is a complex manifold of dimension m and $\tau: M \rightarrow (0, \Delta^2)$, $0 < \Delta = \sup \sqrt{\tau} \leq \infty$, is a C^∞ strictly plurisubharmonic exhaustion such that $(dd^c \log \tau)^m = 0$ on M_* . He proves that, up to exhaustion-preserving biholomorphic maps, the only such pairs are the balls of radius Δ in \mathbb{C}^m equipped with the exhaustion $\tau_0 = \| \cdot \|^2$ (simplified proofs of this theorem are given by Burns [Bu] and Wong [W 1]). Clearly, strictly parabolic manifolds are manifolds of circular type. This can be also easily checked directly without referring to the classification theorem (cfr. [SPM], Proposition 2.2).

Example 2. Let G be a strictly pseudoconvex, bounded, smooth, complete circular domain in \mathbb{C}^m . The Minkowski functional $\rho: \mathbb{C}^m \rightarrow \mathbb{R}_+$ associated to G is defined by

$$(1.5) \quad \rho(z) = \begin{cases} 0 & \text{if } z = 0 \\ \inf \{1/t \mid t > 0 \text{ and } tz \notin G\} & \text{if } z \neq 0. \end{cases}$$

It is not hard to check that (G, ρ^2) is a manifold of circular type with radius 1 (cfr. [BT 2] and [PW]).

Example 3. Let D be a smooth, bounded, strictly convex domain in \mathbb{C}^m . According to [L 1], for any $p \in D$ there exists a surjective C^∞ map $F: \mathbb{C}[1] \times S^{2m-1} \rightarrow \bar{D}$, which we called stationary representation of D at p in [P 1], such that for all $b \in S^{2m-1}$ $F(\cdot, b): \mathbb{C}(1) \rightarrow D$ is a proper holomorphic embedding with $F(0, b) = p$, $F'(0, b) = \|F'(0, b)\| b$ and such that, if K denotes the Kobayashi metric of D , then $K(p, b) = (\|F'(0, b)\|)^{-1}$. Also if $\lambda \in \mathbb{C}$ with $|\lambda| = 1$ and $(z, b) \in \mathbb{C}[1] \times S^{2m-1}$, then $F(\lambda z, b) = F(z, \lambda b)$. An exhaustion $\tau: \bar{D} \rightarrow [0, 1]$, called the Lempert exhaustion of D at p , is well defined by $\tau(F(z, b)) = |z|^2$. Using results of Lempert [L 1] one can show that (D, τ) is a manifold of circular type with radius 1. Properties of the Lempert exhaustions and applications to the classification of strictly convex were studied in [P 1].

In the remainder of this section we give some preliminar results on manifold of circular type. First we have the following.

Theorem 1.1. Let (M, τ) be a manifold of circular type. Then the set $M[0]$ consists exactly of one point which we call the center of M .

Proof. Since no use of the smoothness of τ on $M[0]$ is made, a word by word repetition of the proof of Theorem 2.5 of [SPM] shows that $M[0]$ is connected. On the other hand, since τ satisfies (1.4), then $M[0]$ has an isolated point and therefore the claim follows.

q.e.d.

Remark One can show that given any smooth, bounded, strictly convex domain $D \subset \mathbb{C}^m$ and any $p \in D$, the only exhaustion σ of \bar{D} such that (D, σ) is a manifold of circular type with center $\{p\}$ and radius 1 is the Lempert exhaustion at p . A proof of this fact based on the Bedford-Taylor minimum principle for the Monge-Ampère operator (cfr. [BT 1]) is in [L 2]. Likewise given a strictly pseudoconvex, bounded, smooth, complete circular domain $G \subset \mathbb{C}^m$ then the unique exhaustion τ of \bar{G} such that (G, σ) is a manifold of circular type with center at the origin and radius 1 is $\sigma = \rho^2$ where ρ is the Minkowski functional of G . The proof of this fact, which goes along the same lines of the case of strictly convex domains, can be found in [P 2].

Let (M, τ) be a manifold of circular type. We shall now collect some known facts which will be needed in this paper (cfr. for proofs [Bu] or [SPM] or [W 1]). A simple calculation shows that (1.2) and (1.3) imply that on M_* , with respect to any choice of local coordinates, the following fundamental formula holds

$$(1.6) \quad \tau_{\bar{\nu}} \tau^{\bar{\nu}\mu} \tau_{\mu} = \tau$$

where we set $(\tau^{\bar{\nu}\mu}) = (\tau_{\mu\bar{\nu}})^{-1}$. A Kähler metric k on M_* is defined by the form $dd^c \tau > 0$. A unique vector field X of type $(1,0)$, called the complex gradient of τ , exists which is dual to $\bar{\partial}\tau$ with respect to k i.e. such that:

$$(1.7) \quad \bar{\partial}\tau(\bar{W}) = k(X, W)$$

for all vector field W of type $(1,0)$. In local coordinates we

have the following expression for X :

$$(1.8) \quad X = X^\mu \frac{\partial}{\partial z^\mu} = \tau^{\bar{\nu}\mu} \tau_{\bar{\nu}} \frac{\partial}{\partial z^\mu} .$$

Because of (1.6) we have also

$$(1.9) \quad \bar{\partial}\tau(X) = k(X,X) = \tau .$$

From (1.2) and (1.3) it follows that the form $dd^c \log \tau$ has rank $m - 1$ at every point of M_* . Then a distribution \mathcal{A} of rank $m - 1$ is defined by

$$(1.10) \quad \mathcal{A} = \{W \in T^+(M_*) \mid dd^c \log \tau(W, \bar{V}) = 0 \text{ for all } V \in T^+(M_*)\} .$$

The distribution \mathcal{A} is integrable and the associated foliation of M_* in Riemann surfaces is called the Monge-Ampère foliation associated to τ . This foliation, first introduced by Bedford and Kalka [BK], has been focus of many studies recently (see for instance [B],[PW],[SPM],[W 1]). Here we shall just recall few well known properties. The distribution \mathcal{A} is a trivial subbundle of $T^+(M_*)$ spanned by the vector field X i.e. for every $p \in M_*$ we have

$$(1.11) \quad \mathcal{A}_p = \mathbb{C}X_p .$$

As a consequence we get that the Monge-Ampère foliation is holomorphic - i.e, \mathcal{A} is a holomorphic subbundle of $T^+(M_*)$ - if and only if X is holomorphic.

A further remarkable property of the Monge-Ampère foliation is that its leaves are totally geodesic submanifold of M_* with respect the Kähler metric k . In fact if g is the Riemannian metric associated to k and we define a vector field Y on M_* by

$$(1.12) \quad Y = \frac{1}{\sqrt{\tau}} (X + \bar{X}),$$

then we have

$$(1.13) \quad Y = \text{grad}_g \sqrt{\tau} \quad \text{and} \quad g(Y, Y) = 1 .$$

Every integral curve of Y lies on a leaf of the Monge-Ampère foliation and it is a geodesic for the metric g .

2. THE GRADIENT FLOW

In this section we shall give a precise description of the flow of the vector field Y defined in (1.12). We shall use the same method used by Stoll in [SPM] for the case of strictly parabolic manifolds. We have to recall first the concept of the double at a point of a differentiable manifold. Let N be a differentiable manifold with $\dim_{\mathbb{R}} N = n$. Let $L \subset N$ be a closed submanifold with $\dim_{\mathbb{R}} L = n - 1$. Let $h: N \rightarrow \mathbb{R}$ be a function of class C^∞ with $h|_L = 0$ and $a \in L$. We say that h vanishes of order k at a if there exists a neighbourhood U of a and a function $g: U \rightarrow \mathbb{R}$ with $g^{-1}(0) = L \cap U$ and $dg(x) \neq 0$ if $x \in U$, such that on a neighborhood $U_1 \subset U$ of a we have $h = g^k h_1$, where $h_1: U_1 \rightarrow \mathbb{R}$ is a function of class C^∞ with $h_1(a) \neq 0$. Clearly h need not vanish of any order at a . If M is another differentiable manifold, with $\dim_{\mathbb{R}} M = m$, and $f: N \rightarrow M$ is a differentiable map with $f|_L = b \in M$, we say that f branches of order s at $a \in L$ if, given a local coordinates around $a \in N$ and $b \in M$, one has that

$$D = \det \left(\frac{\partial f^i}{\partial x^j} \right)_{i,j=1,\dots,m}$$

vanishes of order $s + m - 1$ at $a \in L$. This definition does not depend on the choice of local coordinates and one easily sees that $s \geq 0$. Let $0 \in M$ be any point. We say that (N, r) is the double of M at 0 (or connected sum of M and M at 0) (cfr. Milnor [M] and Stoll [SPM]) if:

- (i) N is a differentiable manifold with $\dim_{\mathbb{R}} N = \dim_{\mathbb{R}} M = n$.
- (ii) The map $r: N \rightarrow M$ is proper, surjective and of class C^∞ .
- (iii) The inverse image $r^{-1}(0) = S_0$ is a compact submanifold of N diffeomorphic to S^{m-1} .
- (iv) The map r branches of order 0 at every point of S_0 .
- (v) There exists two open sets $N_1, N_2 \subset N$ such that $N = N_1 \cup N_2 \cup S_0$ is a disjoint union and $r_j = r|_{N_j}: N_j \rightarrow M - \{0\}$ is a diffeomorphism.

In a neighbourhood of S_0 the double of M can be described as follows (cfr. [SPM], pg. 113). Let U be a coordinate neighbourhood of 0 in M and let $B(\epsilon) \subset U$ be a ball of radius ϵ centered at 0 . Let $V = r^{-1}(B(\epsilon))$. If ϵ is small enough, there exists a diffeomorphism $f: V \rightarrow (-\epsilon, \epsilon) \times S^{m-1}$ such that $f(S_0) = \{0\} \times S^{m-1}$ and $r(x) = r_0(f(x))$ for all $x \in V$ where r_0 is defined by $r_0(t, b) = tb$. When working near S_0 , we shall identify V with $(-\epsilon, \epsilon) \times S^{m-1}$ so that $f = \text{Id}$ and $r = r_0$ on V .

Let now (M, τ) be a manifold of circular type of dimension m and radius Δ . To describe the flow of the vector field Y defined in (1.12) we will use the following strategy. First we pull back Y to a vector field \tilde{Y} on the double (N, r) of M at the center $0 \in M$. Then we integrate \tilde{Y} on N and finally we push down the resulting flow. Of course one has to make precise this strategy and carry it out correctly. To this aim is devoted this section.

Let U_0 be the coordinate neighbourhood of the center 0 of M with the properties listed in (1.4). Also the notations of Section 1 will be assumed. Furthermore here we write $S = S^{2m-1}$. Define $U = \{Z \in U_0 \mid Z = tb \text{ with } t \in (-\epsilon, \epsilon) \text{ and } b \in S\}$. If (N, r) is the double of M at 0 then set $V = r^{-1}(U)$. We can choose ϵ small enough so that we can identify $V = (-\epsilon, \epsilon) \times S$ and so that $r(t, b) = tb$. We also set $V_1 = N_1 \cap V = (-\epsilon, 0) \times S$ and $V_2 = N_2 \cap V = (0, \epsilon) \times S$.

LEMMA 2.1. There exists a function $G: V - \{0\} \rightarrow \mathbb{R}_+$ of class C^∞ and constants $C_2 > C_1 > 0$, such that for all $Z \in U - \{0\}$ we have $\tau(Z) = G(Z) \|Z\|^2$ and $C_1 < G(Z) < C_2$. Furthermore there exists $c: V \rightarrow \mathbb{R}$, $H: S \rightarrow \mathbb{R}_+$, $R: V \rightarrow \mathbb{R}$ of class C^∞ such that $c^2(t, b) = \tau(r(t, b))$ for all $(t, b) \in V$ with $c|_{V_1} < 0$, $c|_{V_2} > 0$ and such that

$$(2.1) \quad c(t, b) = \sqrt{H(b)} + t^2 R(t, b) .$$

PROOF The fact that a function G on $U - \{0\}$ exists with the above properties is an immediate consequence of (1.4') part (i). Let $h: (-\epsilon_0, \epsilon_0) \times \mathbb{B}(2) - \{0\} \rightarrow \mathbb{R}_+$ be defined as in (1.4') part (ii). Then its restriction $h: V \rightarrow \mathbb{R}_+$ is of class C^∞ . There exists $G: V \rightarrow \mathbb{R}_+$ of class C^∞ and such that $\tilde{G} > 0$ on V so that $h(t, b) = \tilde{G}(t, b)t^2$. Then we have $\dot{h}(0, b) = 0$ and $\ddot{h}(0, b) > 0$ for all $b \in S$ where we have set $\dot{h} = \frac{\partial h}{\partial t}$. Define $H: S \rightarrow \mathbb{R}_+$ by

$$(2.2) \quad H(b) = \frac{1}{2} \tilde{h}(0,b) .$$

Then there exists $T:V \rightarrow \mathbb{R}$ of class C^∞ such that if $(t,b) \in V$, then $0 \leq h(t,b) = t^2 H(b) + t^3 T(t,b)$. In particular we have that $H(b) + t T(t,b) > 0$ for all $(t,b) \in V$. The function $L:V \rightarrow \mathbb{R}$ defined by $L(t,b) = (H(b) - t T(t,b))^{1/2} - (H(b))^{1/2}$ is of class C^∞ and $L(0,b) = 0$ for all $b \in S$. Thus there exists $R:V \rightarrow \mathbb{R}$ of class C^∞ such that $L(t,b) = t R(t,b)$. But then $c:V \rightarrow \mathbb{R}$ given by

$$c(t,b) = t(H(b))^{1/2} + t^2 R(t,b) = t(H(b) + t T(t,b))^{1/2}$$

is the required function.

q.e.d.

LEMMA 2.2 Let $W = (-\epsilon_0, \epsilon_0) \times (B(2) - \{0\})$. There exists a map $A:W \rightarrow \mathbb{C}^m$ of class C^∞ so that if X is the vector field defined in (1.7), $(t,Z) \in W$ and $t \neq 0$, then

$$(2.3) \quad X(t,Z) = tZ + t^2 A(t,Z) .$$

PROOF Let $(t,Z) \in W$ and $t \neq 0$, then because of Lemma 2.1 we have for $\mu, \nu = 1, \dots, m$:

$$\begin{aligned} \tau(tZ) &= t^2 G(tZ) \|Z\|^2 \\ \tau_{\bar{\nu}}(tZ) &= t G_{\bar{\nu}}(tZ) z^\nu + t^2 G_{\bar{\nu}}(tZ) \|Z\|^2 \\ \tau_{\mu\bar{\nu}}(tZ) &= G_{\mu\bar{\nu}}(tZ) \delta_{\mu\nu} + t G_{\mu\bar{\nu}}(tZ) z^\nu + t G_{\bar{\nu}}(tZ) \bar{z}^\mu \\ &\quad + t^2 G_{\mu\bar{\nu}}(tZ) \|Z\|^2 \end{aligned}$$

Because of (1.4') there exists $\tilde{G}:W \rightarrow (0,\infty)$ of class C^∞ such that $h(t,Z) = \tilde{G}(t,Z)\|tZ\|^2$ and, in fact, $\tilde{G}(t,Z) = G(tZ)$ if $(t,Z) \in W$ and $t \neq 0$. Functions $\hat{G}^\mu:W \rightarrow \mathbb{C}$, $\hat{G}^\nu:W \rightarrow \mathbb{C}$ and $\hat{G}^{\mu\bar{\nu}}:W \rightarrow \mathbb{C}$ of class C^∞ are defined by

$$\hat{G}^\mu(t,Z) = \begin{cases} (1/t)\tilde{G}_\mu(t,Z) = G_\mu(tZ) & \text{if } t \neq 0 \\ \dot{\tilde{G}}_\mu(t,Z) & \text{if } t = 0 \end{cases}$$

$$\hat{G}^\nu(t,Z) = \begin{cases} (1/t)\tilde{G}_\nu(t,Z) = G_\nu(tZ) & \text{if } t \neq 0 \\ \dot{\tilde{G}}_\nu(t,Z) & \text{if } t = 0 \end{cases}$$

$$\hat{G}^{\mu\bar{\nu}}(t,Z) = \begin{cases} (1/t)G_{\mu\bar{\nu}}^\mu(t,Z) = G_{\mu\bar{\nu}}(tZ) & \text{if } t \neq 0 \\ \dot{G}_{\mu\bar{\nu}}^\mu(t,Z) & \text{if } t = 0 \end{cases}$$

where the dot denotes the derivative with respect to t .

Then we have for $(t,Z) \in W$ and $t \neq 0$

$$(2.4) \quad \begin{aligned} \tau_{\bar{\nu}}(tZ) &= t\tilde{G}(t,Z)z^\nu + t^2\hat{G}^{\bar{\nu}}(t,Z)\|Z\|^2 \\ \tau_{\mu\bar{\nu}}(tZ) &= \tilde{G}(t,Z)\delta_{\mu\nu} + tB^{\mu\bar{\nu}}(t,Z) \end{aligned}$$

where $B^{\mu\bar{\nu}}:W \rightarrow \mathbb{C}$ are functions of class C^∞ defined by

$$B^{\mu\bar{\nu}}(t,Z) = \hat{G}^\mu(t,Z)z^\nu + \hat{G}^\nu(t,Z)\bar{z}^\mu + t\hat{G}^{\mu\bar{\nu}}(t,Z)\|Z\|^2.$$

There exist functions $R^{\mu\bar{\nu}}:W \rightarrow \mathbb{C}$ of class C^∞ so that if

$(t, Z) \in W$ and $t \neq 0$ we have

$$\tau^{\bar{\nu}\mu}(tZ) = \frac{1}{\tilde{G}(t, Z)} \delta_{\mu\nu} + t R^{\bar{\nu}\mu}(t, Z)$$

where $(\tau^{\bar{\nu}\mu}) = (\tau_{\mu\bar{\nu}})^{-1}$. But recalling (1.8), we get at once (2.2) where $A:W \rightarrow \mathbb{C}^m$ is the map of class C^∞ defined by

$$A^\mu(t, Z) = \sum_{\nu=1}^m \frac{\|Z\|^2}{\tilde{G}(t, Z)} \delta_{\mu\nu} \hat{G}^{\bar{\nu}}(t, Z) + \tilde{G}(t, Z) R^{\mu\bar{\nu}}(t, Z) Z^\nu + t \|Z\|^2 R^{\mu\bar{\nu}}(t, Z) \hat{G}^{\bar{\nu}}(t, Z).$$

q.e.d.

In order to study vector fields on the double (N, r) of M we make the following identifications. Let $(t, b) \in V$. We identify $T_{(t,b)}(V) = \mathbb{R} \oplus T_b(S)$ where $T_b(S)$ is the tangent space of S at b . Also if $H_b(S)$ is the holomorphic tangent space to S we identify $T_{(t,b)}(V) = \mathbb{R} \oplus i\mathbb{R}b \oplus H_b(S)$. As seen above we can consider V as a neighbourhood of $S_0 \subset N$ and set $r(t, b) = tb$ if $(t, b) \in V$. Also if $p \in V$ then we identify $T_p(U) = T_p^+(U) = T_p^-(U) = \mathbb{C}^m$ so that the projection π_+ , π_- are the identity, the almost complex structure J is the multiplication by i and the conjugation is obtained by conjugation of the coordinates. We denote the hermitian product in \mathbb{C}^m by $(|)$ so that the real scalar product $\langle | \rangle$ is given by $\langle Z | W \rangle = \text{Re}(Z | W)$. Then we have the following formula:

$$(2.5) \quad dr^{-1}(t, b)(Z) = (\text{Re}(Z | b), \frac{\text{Im}(Z | b)}{t} | b + \frac{Z}{t} - \frac{(Z | b)}{t}) \in \mathbb{R} \oplus i\mathbb{R}b \oplus H_b(S)$$

Let J be the almost complex structure on M . On $N - S_0$ a complex structure is defined by pulling back by r the complex structure of M . The associated almost complex structure is denoted by \tilde{J} and we have $\tilde{J} = dr^{-1} \circ J \circ dr$. Thus if $(t,b) \in V - S_0$ and $(u, ivb + w) \in \mathbb{R} \oplus i\mathbb{R} \oplus H_b(S)$

$$(2.6) \quad \tilde{J}_{t,b}(u, ivb + w) = (-tv, \frac{u}{t}ib + iw) .$$

On $N - S_0$ we can define vector fields of class C^∞ \tilde{X} and \tilde{Y} by lifting via r the vector fields X, Y . Also define $\tilde{\tau}: N \rightarrow \mathbb{R}_+$ by $\tilde{\tau} = \tau \circ r$.

LEMMA 2.3 The function $\tilde{\tau}$ is of class C^∞ on N . There exists $c: N \rightarrow \mathbb{R}$ of class C^∞ such that $c^2 = \tilde{\tau}$, $c|_{N_1} < 0$ and $c|_{N_2} > 0$. Furthermore \tilde{X} , $\tilde{J}\tilde{X}$ and \tilde{Y} extend to vector fields of class C^∞ on N .

PROOF. We need only to check the claim on V . It is an immediate consequence of Lemma 2.1 that $\tilde{\tau}$ is of class C^∞ and that $c: N \rightarrow \mathbb{R}$ exists with the required properties. Using

(2.3) and (2.5) we have for $(t,b) \in V - S_0$:

$$(2.7) \quad \begin{aligned} \tilde{X}(t,b) &= dr^{-1}(t,b)(X(r(t,b))) \\ &= t(F_1(t,b), F_2(t,b)ib + F_3(t,b)) \end{aligned}$$

where $F_1: V \rightarrow \mathbb{R}$, $F_2: V \rightarrow \mathbb{R}$, $F_3: V \rightarrow H_b(S)$ are of class C^∞

and defined by

$$(2.8) \quad \begin{cases} F_1(t,b) = 1 + t \operatorname{Re}(A(t,b) | b) \\ F_2(t,b) = \operatorname{Im}(A(t,b) | b) \\ F_3(t,b) = A(t,b) - (A(t,b) | b) . \end{cases}$$

But then \tilde{X} extends of class C^∞ on all V and thus on N .
From (2.6) and (2.7) we have for $(t,b) \in V - S_0$

$$(2.9) \quad \tilde{J}\tilde{X}(t,b) = (-t^2 F_2(t,b), F_1(t,b)ib + itF_2(t,b))$$

and thus also $\tilde{J}\tilde{X}$ extends of class C^∞ on all N .
Finally, since if $(t,b) \in V - S_0$ we have

$$\tilde{Y}(t,b) = \frac{1}{c(t,b)} \left(\tilde{X}(t,b) + \overline{\tilde{X}(t,b)} \right) ,$$

\tilde{Y} extends of class C^∞ on N because of (2.1) and (2.7)

q.e.d.

REMARK Using (2.8) and (2.9) we also have the following useful formula for $b \in S$:

$$(2.10) \quad \tilde{J}\tilde{X}(0,b) = (0, ib) .$$

PROPOSITION 2.4 There exists a number $\eta > 0$ and a map

$\Psi: (-\eta, \Delta) \times S \rightarrow N$ of class C^∞ such that

- (i) $\Psi(0, b) = (0, b) \in S_0 = \{0\} \times S$ for all $b \in S$;
- (ii) $\dot{\Psi}(t, b) = \tilde{Y}(\Psi(t, b))$ for all $(t, b) \in (-\eta, \Delta) \times S$;
- (iii) The map $\Psi: [0, \Delta] \rightarrow S_0 \cup N_2$ is a diffeomorphism;
- (iv) $c(\Psi(t, b)) = t$ for all $(t, b) \in [0, \Delta] \times S$;
- (v) If $t \geq 0$, define $\Psi_t: S \rightarrow N$ by $\Psi_t(b) = \Psi(t, b)$ and $N_t = \{c = t\}$. Then $\Psi_t: S \rightarrow N_t$ is a diffeomorphism.

PROOF Since $c: N \rightarrow \mathbb{R}$ is a function of class C^∞ such that $c^{-1}([0, \Delta])$ is non empty, non compact and connected, $c^{-1}([0, t])$ is compact for all $t \in [0, \Delta]$, $S_0 = c^{-1}(\{0\}) \neq \emptyset$ and \tilde{Y} is a vector field of class C^∞ on N , according to Theorem 4.1 of [SPM] the map Ψ with the properties (i) - (v) exists provided $dc(p, \tilde{Y}(p)) = 1$ for all $p \in c^{-1}([0, \Delta]) = S_0 \cup N_2$. To show this observe that

$$\begin{aligned} dr(p, \tilde{Y}(p)) &= -Y(r(p)) \quad \text{on } N_1 \\ dr(p, \tilde{Y}(p)) &= Y(r(p)) \quad \text{on } N_2 \quad . \end{aligned}$$

But then on N_1 using (1.13) we have

$$\begin{aligned} dc(p, \tilde{Y}(p)) &= -d\sqrt{r}(r(p), dr(p, \tilde{Y}(p))) \\ &= -d\sqrt{r}(r(p), -Y(r(p))) = 1 \end{aligned}$$

In the same way one obtains $dc(p, \tilde{Y}(p)) = 1$ on N_2 and thus by continuity the claim is true on N .

q.e.d.

LEMMA 2.5 Let $r_0 > 0$ small enough so that $\Psi((-r_0, r_0) \times S) \subset V$ and let $\Psi_1: (-r_0, r_0) \times S \rightarrow \mathbb{R}$, $\Psi_2: (-r_0, r_0) \times S \rightarrow S$ such that $\Psi = (\Psi_1, \Psi_2)$. There exist $\Psi_3: (-r_0, r_0) \times S \rightarrow \mathbb{R}$ and $\Psi_4: (-r_0, r_0) \times S \rightarrow \mathbb{C}^m$ of class C^∞ such that

$$(2.11) \quad \Psi_1(t, b) = \frac{1}{\sqrt{H(b)}} t + t^2 \Psi_3(t, b) ,$$

$$(2.12) \quad \Psi_2(t, b) = b + t \Psi_4(t, b) .$$

PROOF Because of (i) of Theorem 2.4 there exist $\Psi_4: (-r_0, r_0) \times S \rightarrow \mathbb{C}^m$ and $\Psi_5: (-r_0, r_0) \times S \rightarrow \mathbb{R}$ of class C^∞ such that (2.12) holds and $\Psi_1(t, b) = t \Psi_5(t, b)$. If π_+ is the isomorphism defined in Section 0, then

$$\begin{aligned} \lim_{t \rightarrow 0} \pi_+(\tilde{Y}(t, b)) &= \lim_{t \rightarrow 0} \frac{1}{c(t, b)} \tilde{X}(t, b) \\ &= \frac{1}{\sqrt{H(b)}} (1, F_2(0, b)ib + F_3(0, b)) \end{aligned}$$

where we have used (2.1), (2.7) and (2.8). Thus because of (ii) of Theorem 2.4 we have $\Psi_5(0, b) = \dot{\Psi}_1(0, b) = (H(b))^{-1/2}$. But then there exists $\Psi_3: (-r_0, r_0) \times S \rightarrow \mathbb{R}$ of class C^∞ such that $\Psi_5(t, b) = (H(b))^{-1/2} + t \Psi_3(t, b)$ and thus the proof is completed.

q.e.d.

We have now all the ingredients to describe in detail the flow on M of the vector field Y defined in (1.12). Here we identify $T_0(M) = \mathbb{C}^m$.

THEOREM 2.6 There exists a map of class C^∞

$$(2.13) \quad \varphi: [0, \Delta) \times S \rightarrow M$$

with the following properties

- (i) $\varphi(0, b) = 0$ for all $b \in S$;
- (ii) $\dot{\varphi}(0, b) = (H(b))^{-1/2} b$ for all $b \in S$;
- (iii) $\dot{\varphi}(t, b) = Y(\varphi(t, b))$ for all $(t, b) \in (0, \Delta) \times S$;
- (iv) $\varphi: (0, \Delta) \times S \rightarrow M_*$ is a diffeomorphism;
- (v) $\tau(\varphi(t, b)) = t^2$ for all $(t, b) \in [0, \Delta) \times S$;
- (vi) the map $\varphi_t: S \rightarrow M\langle t \rangle$ defined by $\varphi_t(b) = \varphi(t, b)$ is a diffeomorphism;
- (vii) $t\pi_+(\dot{\varphi}(t, b)) = X(\varphi(t, b))$ for all $(t, b) \in [0, \Delta) \times S$;
- (viii) The curve $\varphi(\square, b): (0, \Delta) \times S \rightarrow M_*$ is geodesic with respect to the Kähler metric defined by $dd^c \tau > 0$ for all $b \in S$.

PROOF If $r: N \rightarrow M$ is the projection from the double, we define $\varphi = r \circ \Psi: [0, \Delta) \rightarrow M$. Then (i), (iii), (iv), (v), (vi) and (vii) are immediate from Proposition 2.4. Also, using the usual identification of the tangent space of M at the center 0 with \mathbb{C}^m , with respect to local coordinates around 0 we have recalling (2.11) and (2.12):

$$\begin{aligned} \varphi(t, b) &= r(\Psi(t, b)) \\ &= r(\Psi_1(t, b), \Psi_2(t, b)) \\ &= t(H(b))^{-1/2} b + t^2 T(t, b) \end{aligned}$$

where T is of class C^∞ and thus (ii) follows. Finally, (viii) is a consequence of the well known fact recalled in Section 1 that the integral curves of Y are geodesics.

q.e.d.

COROLLARY 2.7 The map $E: \mathcal{B}(\Delta) \rightarrow M$ defined by

$$(2.14) \quad E(Z) = \begin{cases} 0 & \text{if } Z=0 \\ \varphi(\|Z\|, Z/\|Z\|) & \text{if } Z \neq 0 \end{cases}$$

is a homeomorphism and its restriction $E: \mathcal{B}(\Delta) - \{0\} \rightarrow M_*$ is a diffeomorphism.

PROOF. The map $E: \mathcal{B}(\Delta) - \{0\} \rightarrow M_*$ is clearly a diffeomorphism. Also $\tau(E(Z)) = \|Z\|^2$ and hence $\lim_{Z \rightarrow 0} E(Z) = 0$ and $\lim_{p \rightarrow 0} E^{-1}(p) = 0$ so that the proof is complete.

q.e.d.

REMARK The metric k defined on M_* by $dd^c \tau$ is not defined on the center 0 of M . Thus one cannot define in the usual way an exponential map at 0 . The map E defined in (2.14) is a generalized exponential map at the "singular" point 0 with respect to the metric k . The existence of an exponential map at a singular point of a Riemannian manifold has been investigated by Stone [St] in terms of curvature conditions in a neighbourhood of the singularity. Here it follows from the behavior of the exhaustion which defines the metric. It would be interesting to

understand the relationship between Stone's hypothesis and ours in this particular case. Notice also that the map E is differentiable at 0 if and only if the metric extends over 0 and this is the case if M is strictly parabolic (in which case E is biholomorphic). This will be evident from the results of the next section.

3. THE LEAF PARAMETRIZATION MAP

In order to give a parametrization of the leaves of the Monge-Ampère foliation induced by τ we must consider also the flow of the vector field Z defined on M_* by

$$(3.1) \quad Z = i(X - \bar{X}) = J(X + \bar{X}) .$$

where X is as usual the complex gradient of τ . Using (1.9) we have

$$d\tau(Z) = i\partial\tau(X) - i\bar{\partial}\tau(\bar{X}) = i\tau - i\tau = 0$$

and therefore any integral curve of Z is contained in some level set of τ which is compact. But this implies that Z is complete i.e. there exists a unique map $\sigma : \mathbb{R} \times M_* \rightarrow M_*$ such that

$$(3.2) \quad \sigma(\square, p) : \mathbb{R} \rightarrow M_* \text{ is an integral curve of } Z \text{ such that} \\ \sigma(0, p) = p.$$

$$(3.3) \quad \text{For all } y \in \mathbb{R} \text{ the map } \sigma(y, \square) : M_* \rightarrow M_* \text{ is a diffeomorphism.}$$

$$(3.4) \quad \text{For all } y_1, y_2 \in \mathbb{R} \text{ and } p \in M_* \text{ we have} \\ \sigma(y_1 + y_2, p) = (y_1, \sigma(y_2, p)).$$

Define a map $\alpha : [0, \Delta) \times M_* \rightarrow M$ by

$$(3.5) \quad \alpha(t, p) = \varphi(t, \varphi^{-1}(\frac{p}{\sqrt{\tau(p)}})).$$

Clearly α is continuous and of class C^∞ on $(0, \Delta) \times M_*$.

LEMMA 3.1 For all $t \in \mathbb{R}$, $s \in (0, \Delta)$ and $p \in M_*$ we have

$$(3.6) \quad \sigma(t, \alpha(s, p)) = \alpha(s, \sigma(t, p))$$

PROOF. Since α is such that for all $p \in M_*$ $\alpha(\square, p) : (0, \Delta) \rightarrow M_*$ is the unique integral curve of Y such that $\alpha(\sqrt{\tau(p)}, p) = p$,

(3.6) is equivalent to $[Y, Z] = 0$ on M_* . Define $F = X + \bar{X}$ so that $Y = (\tau)^{-1/2} F$ and $Z = JF$. Then

$$[Y, Z] = \frac{1}{\sqrt{\tau}} [F, JF] - JF \left(\frac{1}{\sqrt{\tau}} \right) F.$$

Since $JF((\tau)^{-1/2}) = 0$ follows easily from (1.9) to show $[Y, Z]$ we have only to prove $[F, JF] = 0$. If $[F, JF] = H^\mu \frac{\partial}{\partial z^\mu} + \bar{H}^\mu \frac{\partial}{\partial \bar{z}^\mu}$ then one computes $H^\mu = 2i \bar{X}^\nu X^\mu_{\bar{\nu}}$. On the other hand

$$\begin{aligned} \bar{X}^\nu X^\mu_{\bar{\nu}} &= \tau_\alpha \tau^{\bar{\nu}\alpha} \left(\tau_{\bar{\beta}} \tau^{\bar{\beta}\mu} \right)_{\bar{\nu}} \\ &= \tau_\alpha \tau^{\bar{\nu}\alpha} \tau_{\bar{\beta}\bar{\nu}} \tau^{\bar{\beta}\mu} + \tau_\alpha \tau^{\bar{\nu}\alpha} \tau_{\bar{\beta}} \tau^{\bar{\beta}\mu}_{\bar{\nu}}. \end{aligned}$$

Differentiating the equality $\tau_{\sigma\bar{\gamma}} \tau^{\bar{\nu}\sigma} = \delta_{\gamma\nu}$ and (1.6)

we obtain the following equalities:

$$\tau_{\sigma\bar{\gamma}} \tau^{\bar{\nu}\sigma}_{\bar{\beta}} + \tau_{\bar{\gamma}\bar{\beta}\sigma} \tau^{\bar{\nu}\sigma} = 0$$

$$\tau_{\bar{\beta}\bar{\nu}} \tau^{\bar{\nu}\alpha} \tau_{\alpha} + \tau_{\bar{\nu}\bar{\beta}} \tau^{\bar{\nu}\alpha} \tau_{\alpha} = 0 .$$

But then we have

$$\begin{aligned} \tau_{\alpha} \tau^{\bar{\nu}\alpha} \tau_{\bar{\beta}\bar{\nu}} \tau^{\bar{\beta}\mu} &= -\tau_{\alpha} \tau^{\bar{\nu}\alpha} \tau_{\bar{\nu}} \tau^{\bar{\beta}\mu} \\ &= -\tau_{\bar{\nu}} \tau^{\bar{\gamma}\alpha} \tau_{\sigma\bar{\gamma}} \tau^{\bar{\nu}\sigma} \tau_{\alpha} \tau^{\bar{\beta}\mu} \\ &= \tau_{\bar{\nu}} \tau^{\bar{\gamma}\alpha} \tau_{\sigma\bar{\gamma}\bar{\beta}} \tau^{\bar{\nu}\sigma} \tau_{\alpha} \tau^{\bar{\beta}\mu} \end{aligned}$$

and

$$\begin{aligned} \tau_{\bar{\beta}} \tau^{\bar{\beta}\mu} \tau_{\alpha} \tau^{\bar{\nu}\alpha} &= \tau_{\bar{\beta}} \tau^{\bar{\gamma}\mu} \tau_{\sigma\bar{\gamma}\bar{\nu}} \tau^{\bar{\beta}\sigma} \tau_{\alpha} \tau^{\bar{\nu}\alpha} \\ &= -\tau_{\bar{\beta}} \tau^{\bar{\gamma}\mu} \tau_{\sigma\bar{\gamma}\bar{\nu}} \tau^{\bar{\beta}\sigma} \tau_{\alpha} \tau^{\bar{\nu}\alpha} \\ &= -\tau_{\bar{\nu}} \tau^{\bar{\beta}\mu} \tau_{\sigma\bar{\beta}\bar{\gamma}} \tau^{\bar{\nu}\sigma} \tau_{\alpha} \tau^{\bar{\gamma}\alpha} \end{aligned}$$

where in the last line we exchange the indices in the following way : $\beta \rightarrow \nu, \gamma \rightarrow \beta, \nu \rightarrow \gamma$. Our computation shows that $\bar{X}^{\nu} X^{\mu} = 0$ and thus $[F, JF] = 0$.

q.e.d.

We need now some notations. For $R \in (0, \infty]$ define

$$(3.7) \quad H(R) = \begin{cases} \{z \in \mathbb{Z} \mid \operatorname{Re} z < \log R\} & \text{if } R < \infty \\ \mathbb{Z} & \text{if } R = \infty. \end{cases}$$

Also fix $x_0 \in (0, \Delta)$. Write $q = \varphi_{x_0}$. A one parameter group of diffeomorphisms $\zeta: \mathbb{R} \times S \rightarrow S$ is defined by

$$(3.8) \quad \zeta(y, b) = q^{-1}(\sigma(y, q(b))) .$$

Finally we define a surjective map $W: H(\Delta) \times S \rightarrow M_*$ of class C^∞ by

$$(3.9) \quad W(x+iy, b) = \sigma(y, \varphi(e^x, b))$$

From (3.5), (3.6) and (3.8) one has at once

$$\begin{aligned} W(x+iy, b) &= \sigma(y, \alpha(e^x, q(b))) \\ (3.10) \quad &= \alpha(e^x, \sigma(y, q(b))) \\ &= \varphi(e^x, \zeta(y, b)) . \end{aligned}$$

LEMMA 3.2. For all $b \in S$ the map $W(\square, b): H(\Delta) \rightarrow M_*$ is holomorphic and $W'(z, b) = X(W(z, b))$. Furthermore if $x + iy, s + it \in H(\Delta)$ and $b, c \in S$, then $W(x + iy, b) = W(s + it, c)$ if and only if $x = s$ and $c = \zeta(y - t, b)$.

PROOF. If we set $p = W(x+iy, b)$, then by (3.10):

$$\begin{aligned} \frac{\partial}{\partial x} W(x+iy, b) &= \dot{\varphi}(e^x, \zeta(y, b)) e^x = \sqrt{\tau(p)} Y(p) = X(p) + \overline{X(p)} \\ \frac{\partial}{\partial x} W(x+iy, b) &= \dot{\sigma}(y, \varphi(e^x, b)) = Z(p) = J(X(p) + \overline{X(p)}) . \end{aligned}$$

But then $\frac{\partial W}{\partial x} + J \frac{\partial W}{\partial y} = \frac{\partial W}{\partial x} = \frac{\partial W}{\partial x} + JJ \frac{\partial W}{\partial x} = 0$ and $W(\square, b)$ is thus holomorphic. Moreover $W'(z, b) = \pi_+ \left(\frac{\partial W(z, b)}{\partial x} \right) = X(W(z, b))$. Finally if $W(x+iy, b) = W(s+it, c)$ then $\varphi(e^x, \zeta(y, b)) = \varphi(e^s, \zeta(t, b))$ and therefore $x = s$ and $\zeta(y, b) = \zeta(t, c)$ which implies $c = \zeta(y-t, b)$ since ζ is a one parameter group. In the other direction the implication is immediate.

q.e.d.

We need now to compute explicitly $\zeta(y, b)$ for $y \in \mathbb{R}$ and $b \in S$. To this end we shall again use the notations introduced in Section 2. Let (N, r) be the double of M at the center 0 , $V = (-\epsilon, \epsilon) \times S \subset N$ and let $U = r(V)$. If Ψ is the map defined in Proposition 2.4, let $r_0 > 0$ be such that $\Psi([0, r_0] \times S) \subset V$. Then, since $\varphi = r \circ \Psi$, we have also $\varphi([0, r_0] \times S) \subset U$. It is not restrictive to assume that, if $x_0 \in (0, \Delta)$ is the number used in the definition (3.8) of ζ , then $x_0 \in (0, r_0)$.

PROPOSITION 3.3 For all $y \in \mathbb{R}$ and $b \in S$ we have

$$(3.11) \quad \zeta(y, b) = e^{1y} b .$$

PROOF. Fix $b \in S$. For $t \in (0, r_0)$ and $y \in \mathbb{R}$ define $\gamma(t, y) = r^{-1}(\varphi(t, \zeta(y, b)))$. Then γ is of class C^∞ on $(0, r_0) \times \mathbb{R}$ and we have

$$\gamma(t, y) = \Psi(t, \zeta(y, b)) = r^{-1}(W(\log t + iy, b)) .$$

Using (2.11) and (2.12) we have

$$\begin{aligned} \gamma(t,y) &= (\Psi_1(t,\zeta(y,b)) , \Psi_2(t,\zeta(y,b))) \\ &= \left(\frac{t}{\sqrt{H(\zeta(y,b))}} + t^2\Psi_3(t,\zeta(y,b)), \zeta(y,b) + t\Psi_4(t,\zeta(y,b)) \right) \end{aligned}$$

Thus

$$\begin{aligned} \gamma_y(t,y) &= \left(t \frac{\partial}{\partial y} \left(\frac{1}{\sqrt{H(\zeta(y,b))}} + t\Psi_3(t,\zeta(y,b)) \right), \zeta_y(y,b) \right. \\ &\quad \left. + t \frac{\partial}{\partial y} \Psi_4(t,\zeta(y,b)) \right) \end{aligned}$$

The limit of $\gamma_y(t,y)$ as $t \rightarrow 0$ exists and

$$\gamma_y(0,y) = \lim_{t \rightarrow 0} \gamma_y(t,y) = (0, \zeta_y(y,b)) .$$

On the other hand since

$$W_y(\log t + iy, b) = iW'(\log t + iy, b) = iX(W(\log t + iy, b))$$

we have:

$$\begin{aligned} \gamma_y(t,b) &= dr^{-1}(W(\log t + iy, b), W_y(\log t + iy, b)) \\ &= dr^{-1}(\varphi(t,\zeta(y,b)), iX(\varphi(t,\zeta(y,b)))) \\ &= \tilde{J}\tilde{X}(\Psi(t,\zeta(y,b))) \end{aligned}$$

Thus using (2.10) we have:

$$\begin{aligned}\gamma_y(0,y) &= \lim_{t \rightarrow 0} \gamma_y(t,y) \\ &= \tilde{J}\tilde{X}(\Psi_1(0,\zeta(y,b)), \Psi_2(0,\zeta(y,b))) \\ &= \tilde{J}\tilde{X}(0,\zeta(y,b)) \\ &= i\zeta(y,b)\end{aligned}$$

Comparing the two expressions that we have obtained for $\gamma_y(0,y)$ we get

$$\zeta_y(y,b) = i\zeta(y,b).$$

Integrating this equation and imposing the initial condition $\zeta(0,b) = b$ we conclude that (3.11) holds.

q.e.d.

We can now state the main result of this section. As usual we shall identify the tangent space to M at the center 0 with \mathbb{C}^m and we shall denote by $\| \cdot \|$ the euclidean norm on it.

THEOREM 3.4. There exists uniquely a surjective map, called the leaf parametrization map,

$$(3.12) \quad F: \mathbb{C}(\Delta) \times S \rightarrow M$$

of class C^∞ and with the following properties:

$$(i) \quad F(0,b) = 0 \quad \text{for all } b \in S;$$

- (ii) $\tau(F(z,b)) = |z|^2$ for all $b \in S$ and $z \in \mathbb{C}(\Delta)$;
- (iii) $F(t,b) = \varphi(t,b)$ for all $b \in S$ and $t \in [0,\Delta)$;
- (iv) $F(z,\lambda b) = F(\lambda z,b)$ for all $z \in \mathbb{C}(\Delta)$, $b \in S$ and $\lambda \in \mathbb{C}$ with $|\lambda| = 1$;
- (v) $F(\square,b) : \mathbb{C}(\Delta) \rightarrow M$ is holomorphic, proper and injective;
- (vi) $X(F(z,b)) = zF'(z,b)$;
- (vii) $F'(0,b) = \|F'(0,b)\| b = \frac{b}{\sqrt{H(b)}}$.

PROOF Since $\exp: \mathbb{H}(\Delta) \rightarrow \mathbb{C}(\Delta) - \{0\}$ is the universal cover and, because of Lemma 3.2 and Proposition 3.3, we have $W(z_1,b) = W(z_2,b)$ if and only if $z_2 = z_1 + 2k\pi i$, a unique map $F: \mathbb{C}(\Delta) - \{0\} \rightarrow M_*$ of class C^∞ is defined by $F(e^z,b) = W(z,b)$. Also

$F(\square,b): \mathbb{C}(\Delta) - \{0\} \rightarrow M_*$ is holomorphic and injective. Define $F(0,b) = 0$ for all $b \in S$. Then $\tau(F(0,b)) = 0$. If $z \in \mathbb{C}(\Delta) - \{0\}$, then $z = e^{x+iy}$ with $x + iy \in \mathbb{H}(\Delta)$. Thus $\tau(F(z,b)) = \tau(W(x + iy,b)) = e^{2x} = |z|^2$ and (ii) is verified. Clearly $F: \mathbb{C}(\Delta) \times S \rightarrow M$ so defined is continuous and, by Riemann extension theorem, $F(\square,b): \mathbb{C}(\Delta) \rightarrow M$ is holomorphic for all $b \in S$. Let $U \subset M$ be a coordinate neighbourhood centered at $0 \in M$. With no loss of generality we can identify it with an open subset of \mathbb{C}^m . Let $t_0 \in (0,\Delta)$ be such that $F(\mathbb{C}(t_0) \times S) \subset U$. Let $s \in (0,t_0)$ then if $z \in \mathbb{C}(s)$ and $b \in S$ we have

$$F(z,b) = \frac{1}{2\pi i} \int_{|w|=s} \frac{F(w,b)}{w-z} dw$$

and hence F is of class C^∞ on $\mathbb{C}(s) \times S$ and therefore on $\mathbb{C}(\Delta) \times S$. If $K \subset M$ is compact then $K \subset M[r]$ for some $r \in (0,\Delta)$.

Then $F^{-1}(K) \subset \mathbb{C}[r] \times S$ and therefore F is proper and thus so is $F(\square, b)$ for all $b \in S$. Also (iii) follows by construction and F is surjective because so is φ . As a consequence of the definition of F , Lemma 3.2 and Proposition 3.3 one obtains (iv) immediately. Again from Lemma 3.2 one gets (vi). Finally (vii) is a consequence of (iii) and of (ii) of Theorem 2.6.

q.e.d.

REMARK If $E: \mathbb{B}(\Delta) \rightarrow M$ is the map defined in (2.14), because of (iii) and (iv) of Theorem 3.4 we have that $E(zb) = F(z, b)$ for all $(z, b) \in \mathbb{C}(\Delta) \times S$. In other words the restriction of E on any disk through the origin is holomorphic. This proves the claim made earlier that E is biholomorphic if and only if it is of class C^∞ at the origin.

EXAMPLES Before moving on it may clarify the matter to write down the leaf parametrization map for some explicit example. Let G be a strictly pseudoconvex, bounded, smooth, complete circular domain in \mathbb{C}^m and let ρ its Minkowski functional. If $\sigma = \rho^2$ one computes easily that the complex gradient vector field X of σ is given by $X(z) = z^\mu \frac{\partial}{\partial z^\mu}$. It follows then that the map $F: \mathbb{C}(1) \times S \rightarrow G$, relative to the manifold of circular type (G, σ) , is defined by $F(z, b) = z(\rho(b))^{-1}b$. Let $D \subset \mathbb{C}^m$ be a bounded, smooth, strictly convex domain, $p \in D$ and τ be the Lempert exhaustion at p . From the results of [P 1] it follows that the map F for (D, τ) is just the stationary parametrization map which defines τ . However, it is not

hard to check directly that $F(\square, b) : \mathbb{E}(1) \rightarrow D$ is a stationary map in the sense of [L 1] for all $b \in S$ and thus that the map F is indeed the stationary parametrization of D .

4. THE CIRCULAR REPRESENTATION

Let as before (M, τ) be a manifold of circular type and Δ be its radius. We shall now define an "approximation" (G, σ) of (M, τ) where G is a circular domain and σ is, up to a factor, its Minkowski functional squared. Let $F: \mathbb{C}(\Delta) \rightarrow S \times M$ be the leaf parametrization map of (M, τ) . As usual we identify freely $T_0(M)$ with \mathbb{C}^m . Define $\sigma: \mathbb{C}^m \rightarrow \mathbb{R}_+$ by $\sigma(0) = 0$ and

$$(4.1) \quad \sigma(Z) = \left(\frac{\|Z\|}{\|F'(0, Z/\|Z\|)\|} \right)^2 = H(Z/\|Z\|) \|Z\|^2$$

if $Z \neq 0$ where H is defined in (2.7). One easily sees that $\sigma \in C^0(\mathbb{C}^m) \cap C^\infty(\mathbb{C}^m - \{0\})$. Also from the definition of σ and (iv) of Theorem 3.4 one has for all $Z \in \mathbb{C}^m$ and $\lambda \in \mathbb{C}$:

$$(4.2) \quad \sigma(\lambda Z) = |\lambda|^2 \sigma(Z) .$$

For $r \in (0, \infty]$ we define also complete circular domains $G(r) \subset \mathbb{C}^m$ by

$$(4.3) \quad G(r) = \begin{cases} \{Z \in \mathbb{C}^m \mid \sigma(Z) < r^2\} & \text{if } r < \infty \\ \mathbb{C}^m & \text{if } r = \infty \end{cases} .$$

Clearly the Minkowski functional of $G(r)$ is given by $\sqrt{\sigma}/r$.

PROPOSITION 4.1. For all $r \in (0, \infty]$ the pair $(G(r), \sigma)$ is a manifold of circular type with radius r .

PROOF Since (1.1) and (1.4) hold by construction only (1.2) and (1.3) must be shown. But if $dd^C_\sigma > 0$ on $G(r) - \{0\}$ using (4.2) the other statements are obtained by simple calculations. We shall now show $dd^C_\sigma > 0$ on $G(r) - \{0\}$. Since $\sigma_{\mu\bar{\nu}}(tZ) = \sigma_{\mu\bar{\nu}}(Z)$ because of (4.2), it will be enough to show that for some $\varepsilon_0 > 0$ one has $\sigma_{\mu\bar{\nu}}(Z)v^\mu \bar{v}^\nu > 0$ for all $Z \neq 0$ with $\|Z\| < \varepsilon_0$ and $v \in \mathbb{C}^m$ with $\|v\| = 1$. In a small enough neighbourhood U of $0 \in M$ we can choose coordinates so that for some $\varepsilon, \varepsilon_0 > 0$ if $0 < t < \varepsilon$ and $Z \in \mathbb{C}^m - \{0\}$ with $\|Z\| < \varepsilon_0$, then $tZ \in U$ and we have

$$\sigma(tZ) = \tau(tZ) - t^3 L(t, Z)$$

where L is a function of class C^∞ on $(-\varepsilon, \varepsilon) \times (\mathbb{B}(\varepsilon_0) - \{0\})$ (cfr. Lemma 2.1). Furthermore, recalling (2.4), there are functions \tilde{G} and $B^{\mu\bar{\nu}}$ of class C^∞ defined on $(-\varepsilon, \varepsilon) \times (\mathbb{B}(\varepsilon_0) - \{0\})$ such that

$$\tau_{\mu\bar{\nu}}(tZ) = \tilde{G}(t, Z) \sigma_{\mu\bar{\nu}} + t B^{\mu\bar{\nu}}(t, Z) .$$

Also there is a constant $C > 0$ such that $\tilde{G}(t, Z) > C$ for all $(t, Z) \in (-\varepsilon, \varepsilon) \times (\mathbb{B}(\varepsilon_0) - \{0\})$. Fix $Z \in \mathbb{B}(\varepsilon_0) - \{0\}$ and define

$$M = \max_{\substack{|t| \leq \varepsilon/2 \\ v \in S}} \left\| \sum_{\mu\nu} B^{\mu\bar{\nu}}(t, Z) v^\mu \bar{v}^\nu \right\|$$

$$N = \max_{\substack{|t| \leq \varepsilon/2 \\ v \in S}} \left\| \sum_{\mu\nu} L_{\mu\bar{\nu}}(t, Z) v^\mu \bar{v}^\nu \right\|$$

Then for $0 < t < \min \{ \epsilon/2, C/(M + N) \}$ we have

$$\begin{aligned} \sigma_{\mu\bar{\nu}}(Z) V^{\mu}\bar{V}^{\nu} &= \sigma_{\mu\bar{\nu}}(tZ) V^{\mu}\bar{V}^{\nu} \\ &= \tau_{\mu\bar{\nu}}(tZ) V^{\mu}\bar{V}^{\nu} - tL_{\mu\bar{\nu}}(tZ) V^{\mu}\bar{V}^{\nu} \\ &\geq C - t(M + N) > 0 \end{aligned}$$

q.e.d.

Define $\rho : \mathbb{C}^m \rightarrow \mathbb{R}_+$ by

$$(4.4) \quad \rho = \sqrt{\sigma} .$$

Then a map $h:G(\Delta) \rightarrow M$, called the circular representation of (M, τ) , is defined by $h(0) = 0$ and

$$(4.5) \quad h(Z) = F(\rho(Z), Z/\|Z\|),$$

where F is the leaf parametrization map of (M, τ) defined in (3.12), if $Z \neq 0$.

PROPOSITION 4.2 The circular representation has the following properties:

- (i) $h:G(\Delta) \rightarrow M$ is a homeomorphism with $\tau \circ h = \sigma$;
- (ii) $h:G(\Delta) - \{0\} \rightarrow M_*$ is a diffeomorphism;
- (iii) If L is any complex line through the origin in \mathbb{C}^m , then $h|_{G(\Delta) \cap L}$ is a holomorphic map;

(iv) $h:G(\Delta) \rightarrow M$ is a biholomorphic map if and only if it is of class C^∞ at the origin.

PROOF It is clear by construction that $\tau \circ h = \sigma$. Also if we define $Q:G(\Delta) \rightarrow B(\Delta)$ by

$$Q(Z) = \frac{\rho(Z)}{\|Z\|} Z$$

then Q is a homeomorphism and $Q:G(\Delta) - \{0\} \rightarrow B(\Delta) - \{0\}$ is a diffeomorphism. Also if $E:B(\Delta) \rightarrow M$ is the map defined in (2.14) then $h=Q \circ E$. Thus (i) and (ii) follow from Corollary 2.7. Also since $\rho(\lambda Z) \leq |\lambda| \rho(Z)$ for all $\lambda \in \mathbb{C}$ and $Z \in \mathbb{C}^m$, we have $Q(\lambda Z) = \lambda Q(Z)$ for all $\lambda \in \mathbb{C}$ and $Z \in G(\Delta)$ and thus (iii) follows from the fact remarked at the end of Section 3 that E is holomorphic on each disk through the origin. Finally, since a smooth function on a complete circular domain which is holomorphic on each disk through the origin is holomorphic (cfr. [P 1], Lemma 8.1 for a detailed proof), (iii) implies (iv).

q.e.d.

REMARK One can show in fact that the map h is differentiable at the origin (not necessarily of class C^1 !) and that $dh(0) = Id$. The proof is the same as in the case of strictly convex domains (cfr [P 1], Theorem 6.2).

PROPOSITION 4.3 The circular representation $h:G(\Delta) \rightarrow M$ is biholomorphic if and only if the Monge-Ampère foliation associated to τ is holomorphic.

PROOF By the definition of h and (vi) of Theorem 3.4, the disks through the origin in $G(\Delta)$ are mapped by h into the leaves of the Monge-Ampère foliation induced by τ . Thus, if h is biholomorphic, the Monge-Ampère foliation on M is holomorphic. Assume now that the Monge-Ampère foliation is holomorphic. As recalled in Section 1 this is equivalent to the fact that the complex gradient X of τ is holomorphic on M_* . Thus X extends holomorphically on all M with $X(0) = 0$. Moreover, if we regard X as a vector function with values in \mathbb{C}^m , we have, for instance from (2.3), that $dX(0) = \text{Id}$. There exist a neighbourhood U of $0 \in M$ and a holomorphic map $Q:U \rightarrow \mathbb{C}^m$ such that on U

$$(4.6) \quad X = \text{Id} + Q .$$

Define a map $A:\mathbb{C}(1) \times G(\Delta) \rightarrow M$ by $A(z,Z) = h(zZ)$. Then from (4.5) and (iv) of Theorem 3.4, for $Z \neq 0$, one has $A(z,Z) = F(z\rho(Z), Z/\|Z\|)$ so that from (vi) of Theorem 3.4 we have

$$(4.7) \quad zA'(z,Z) = X(A(z,Z)) .$$

For every $j \geq 1$, there exists $A_j:G(\Delta) \rightarrow \mathbb{C}$ such that

$$A(z,Z) = \sum_{j=1}^{\infty} A_j(Z) z^j .$$

Thus from (4.6) and (4.7) we obtain

$$\begin{aligned}
 Q(A(z,Z)) &= zA'(z,Z) - A(z,Z) \\
 (4.8) \quad &= \sum_{j=2}^{\infty} (j-1)A_j(Z)z^j
 \end{aligned}$$

Define $P(z,Z) = \sum_{j=1}^{\infty} A_j(Z)z^{j-1}$. If $Q = \sum_{k=2}^{\infty} Q_k$ is the development of Q in homogeneous vector polynomials of degree k , then there exist vector polynomials Q_{kr} such that

$$Q_k(P(z,Z)) = \sum_{r=0}^{\infty} Q_{kr}(A_1(Z), \dots, A_{r+1}(Z))z^r.$$

Thus

$$\begin{aligned}
 (4.9) \quad Q(A(z,Z)) &= \sum_{k=2}^{\infty} Q_k(P(z,Z))z^k \\
 &= \sum_{k=2}^{\infty} \sum_{r=0}^{\infty} Q_{kr}(A_1(Z), \dots, A_{r+1}(Z))z^{r+k} \\
 &= \sum_{j=2}^{\infty} \left(\sum_{j=k+r} Q_{kr}(A_1(Z), \dots, A_{r+1}(Z)) \right) z^j
 \end{aligned}$$

Comparing (4.8) and (4.9), we get

$$(j-1)A_j(Z) = \sum_{j=k+r} Q_{kr}(A_1(Z), \dots, A_{r+1}(Z))$$

Since $k \geq 2$, we have $r+1 \leq j-1$ and hence there exists a vector polynomial R_j for all $j \geq 2$ such that

$$(4.10) \quad (j-1)A_j(Z) = R_j(A_1(Z), \dots, A_{j-1}(Z)).$$

Since $A_1 = dX(0) = \text{Id}$ is holomorphic, using (4.10) it follows by induction on j that A_j is holomorphic for all $j \geq 1$. On the other hand there exists a number $\varepsilon > 0$ such that $B(2\varepsilon) \subset G(\Delta)$, and $\sum_{j=1}^{\infty} A_j(Z) z^j$ converges uniformly if $\|Z\| < 2\varepsilon$ and $|z| < 1/2$. If $\|Z\| < \varepsilon$ then we have

$$h(Z) = A(1/2, 2Z) = \sum_{j=1}^{\infty} A_j(2Z) (1/2)^j$$

where the series converges uniformly. Therefore h is holomorphic on a neighbourhood of the origin and hence of class C^∞ there. By Proposition 4.2 it follows that in fact h is biholomorphic.

q.e.d

We can now state our main result.

THEOREM 4.4. Let (M, τ) be a manifold of circular type of radius Δ . Then

- (i) If $\Delta = \infty$ there exists a biholomorphic map $h: \mathbb{C}^m \rightarrow M$ such that $(\tau \circ h)^{1/2}$ is the Minkowski functional of some circular domain $G \subset \mathbb{C}^m$.
- (ii) If $\Delta < \infty$ and the Monge-Ampère foliation associated to τ is holomorphic, then there exists a circular domain $G(\Delta) \subset \mathbb{C}^m$ and a biholomorphic map $h: G(\Delta) \rightarrow M$ such that

$\Delta^{-1}(\tau \circ h)^{1/2}$ is the Minkowski functional of $G(\Delta)$.

PROOF Proposition 4.2 and 4.3 yield immediately (ii). To prove (i) we have only to recall that according to a theorem of Burns [Bu], when $\Delta = \infty$ the Monge-Ampère foliation associated to τ is always holomorphic.

q.e.d.

REFERENCES

- [BK] Bedford, E., Kalka, M., Foliations and complex Monge-Ampère equations, Comm. Pure Appl. Math. 30 (1977), 510-538.
- [BT 1] Bedford, E., Taylor, B.A., The Dirichlet problem for a complex Monge-Ampère equation, Invent Math. 37 (1976), 1-44.
- [BT 2] Bedford, E., Taylor, B.A., Variational properties of the complex Monge-Ampère equation, II Intrinsic norms., Amer. J. of Math., 101 (1979), 1131-1166.
- [Bu] Burns, D., Curvature of the Monge-Ampère foliations and parabolic manifolds, Ann. of Math. 115 (1982), 349-373.
- [GK] Griffiths, P.-King J., Nevanlinna theory and holomorphic mappings between algebraic varieties, Acta Math., 130 (1973), 145-220.
- [L 1] Lempert, L., La métrique de Kobayashi et la représentation des domaines sur la boule, Bull. Soc. Mat. France, 109 (1981), 427-474.
- [L 2] Lempert, L., Solving the Degenerate Complex Monge-Ampère Equation with one Concentrated Singularity, Math. Ann., Ann., 263 (1983), 515-532.
- [M] Milnor, J. Sommes de variétés différentiables et structure différentiables des sphères, Bull. Soc. Math. France, 87 (1959), 439-444.
- [P 1] Patrizio, G., Parabolic exhaustions for strictly convex domains, to appear in Manuscripta Math.
- [P 2] Patrizio, G., Holomorphic maps between circular domains, Preprint.

- [PW] Patrizio, G., Wong. P.M., Stability of the Monge-Ampère foliation, Math. Ann., 263 (1983), 13-29.
- [S] Stoll, W., Value distribution on parabolic spaces, Lecture Notes in Math. 600, Springer-Verlag, Berlin-Heidelberg-New York, 1977.
- [SPM] Stoll, W., The characterization of strictly parabolic manifolds, Ann. Scuola Norm. Sup. Pisa, 7 (1980), 81-154.
- [St] Stone, D.A., The exponential map at an isolated singular point, Memoirs of the A.M.S., 256 (1982).
- [W 1] Wong, P.M., Geometry of the Homogeneous Complex Monge-Ampère Equation, Invent. Math., 67 (1982), 261-274.
- [W 2] Wong, P.M., On Umbilical Hypersurfaces and Uniformization of Circular Domains, in Complex Analysis of Several Variables, Y.T. Siu ed., Symp. Pure Math. 41, A.M.S., Providence, 1984.