

BILLIARDS ON ALMOST INTEGRABLE

POLYHEDRAL SURFACES

Eugene Gutkin

Columbia University

New York

USA

Max-Planck-Institut

für Mathematik

Gottfried-Claren-Straße 26

D - 5300 Bonn 3 BRD

SFB / MPI 83 - 26

Introduction Geodesic flows on Euclidean Polyhedra is an old subject that goes as far back as 1906 (see [10] and a related paper [9]). An example of such flow is the motion of a billiard ball inside a polygon. If the angles of the polygon are rational multiples of π , the direction of any geodesic takes only a finite number of values as time varies. Fixing these values one obtains invariant "surfaces" of the billiard flow and the induced flow on the typical invariant surface is minimal ([13]). It is not known whether the flow is typically ergodic (with respect to the invariant Lebesgue measure). The general expectation is that the answer is yes. This question is related to the question whether the typical interval exchange transformation is ergodic where the answer is positive ([5], [12]).

For arbitrary polyhedral surfaces the condition that vertex angles are π -rational does not insure the existence of invariant surfaces for the geodesic flow. One needs an extra condition that a certain holonomy group is finite. I call surfaces satisfying this condition almost integrable because the geodesic flow which is a Hamiltonian system with two degrees of freedom has an additional integral of motion which "almost commutes" with the Hamiltonian.

In section 1 I associate with any almost integrable surface S a Riemann surface R in a purely geometric way. The genus of R is determined by the vertex angles of S (formulas (8) - (11)). The main tool is the developing map of the universal covering of S onto the complex plane. The typical invariant surface of the geodesic flow of S (which I call the billiard flow) is isomorphic to R . Topology of the foliation of the phase space of the billiard flow into invariant surfaces is discussed in [4] .

In section 2 I show that the billiard flow of S is equivalent to a family b_θ^t $0 \leq \theta < 2\pi$ of flows on R (b_θ^t is the billiard flow in direction θ). They are pulled back by the developing map from the linear flows l_θ^t on \mathbb{C} . At the end of the section I extend to b_θ^t the minimality results of [13] and [3].

In section 3 I consider a class of almost integrable surfaces S given essentially by condition that the full holonomy group of S is discrete. For these surfaces Theorem 3 shows the billiard flow b_θ^t is minimal if and only if it is uniquely ergodic (which is false for general almost integrable billiards). Moreover in this case b_θ^t extends the linear flow l_θ^t on a certain torus intrinsically defined by S , thus b_θ^t is minimal if and only if θ is an irrational direction. If θ is irrational the discrete spectrum of b_θ^t coincides with the spectrum of l_θ^t , i.e. b_θ^t is weakly mixing modulo l_θ^t . It is reasonable to expect that for almost integrable polyhedral surfaces outside of this class the billiard flow b_θ^t is typically weakly mixing.

Some results of the paper generalize to billiards on higher dimensional polyhedra which will be discussed elsewhere. It is worth mentioning that almost integrable billiards (both classical and quantum) are of interests to physicists (cf. [8], [1]). Finally I would like to thank A. Katok and W. Veech for useful consultations.

1. Developing map.

By Euclidean polygon I mean a closed bounded polygon P in \mathbb{C} such that its interior $P \setminus \partial P$ is connected. If P has more than one connected component we say that P has obstacles. If an obstacle has two adjacent sides with angle 2π between them we say that P has a slit (see fig. 1).

A Euclidean polyhedron S (of dimension 2) is a collection of Euclidean polygons with some sides identified by isometries. These polygons, their sides and vertices are the faces, edges and vertices of S . A Euclidean polyhedron has a natural topology.

Definition 1. A polyhedral surface S is a connected Euclidean polyhedron homeomorphic to a topological surface.

If a polyhedron S has vertices with an infinite number of adjacent faces then S can not be a polyhedral surface. We say that S is a polyhedral surface with vertices at infinity if S with those vertices punctured becomes a topological surface. Henceforth polyhedron will mean a polyhedral surface possibly with vertices at infinity. A polyhedral surface is closed if it has no boundary, $\partial S = \emptyset$. If $\partial S \neq \emptyset$ two copies of S glued along the boundary make a closed polyhedral surface dS called the doubling of S .

There is a canonical complex structure on any oriented polyhedral surface. Assume first that $\partial S = \emptyset$. Every face and every edge of S define a coordinate patch in an obvious way. Let A be a vertex of S and let P_1, \dots, P_n be the adjacent faces in an orientation preserving order. Let $\alpha_1, \dots, \alpha_n$ be their respective angles. The sum $\alpha = \alpha_1 + \dots + \alpha_n$ is called the angle of A . Cut $U = P_1 \cup \dots \cup P_n$ along an edge b and unfold it on \mathbb{C} so that A goes into 0 and b goes into the positive real axis. Let z be the complex coordinate in \mathbb{C} . Then $u = z^{2\pi/\alpha}$ is a well defined coordinate

in U . It is straightforward to check that the transition functions of the covering are complex analytic and that the complex structure thus defined does not depend on the choices made. If

$\partial S \neq \emptyset$ the imbedding $S \subset dS$ defines the complex structure on S . Thus any polyhedral surface is a Riemann surface.

Polyhedra S_1, S_2 are isomorphic if there is an invertible mapping $f: S_1 \rightarrow S_2$ which maps faces isometrically onto faces. Given a polyhedron S one can always draw new edges on the faces of S . This operation does not change S essentially.

Definition 2. Polyhedra S_1, S_2 are called equivalent if they can be made isomorphic by adding new edges.

A group G of automorphisms of a polyhedron S acts properly discontinuously if for any face $P \subset S$ there is only a finite number of $g \in G$ such that $gP \cap P \neq \emptyset$. The quotient S/G is naturally a polyhedron.

Definition 3. A mapping $f: S_1 \rightarrow S_2$ of polyhedra is a covering if for any $x \in S_1$ there is S_1' equivalent to S_1 , a subpolyhedron R of S_1' containing x and a group G acting properly discontinuously on R such that S_2 is equivalent to R/G and $f|_R$ coincides with the natural projection $R \rightarrow R/G$.

If G acts on S properly discontinuously the covering $S \rightarrow S/G$ is the regular covering with the group G of deck transformations. The reader should be aware of the fact that coverings of polyhedra are usually branched. If $\partial S \neq \emptyset$ the natural involution of dS defines a regular covering $dS \rightarrow S$ with the group $\mathbb{Z}/2$ of deck transformations branched at $\partial S \subset dS$.

Let S be a closed polyhedron and let x_0 be an interior point of a face $P_0 \subset S$. Consider the set of continuous loops on S starting at x_0 and avoiding vertices. The set of homotopy classes of these loops endowed with the usual composition becomes a group

$\pi_f(S)$ called the full fundamental group of S . If \dot{S} is the topological surface obtained by puncturing S at the vertices then $\pi_f(S) = \pi_1(\dot{S})$. Let $\partial S \neq \emptyset$. Imbed S into dS and denote by S_0, S_1 the image and the mirror image of S respectively. Let $x_1 \in S_1$ be the mirror image of $x_0 \in S_0$. To define $\pi_f(S)$ we start from the set of piecewise smooth loops transversal to ∂S . Any such loop γ has a unique lifting $\tilde{\gamma}$ on dS if we agree that $\tilde{\gamma}$ passes from S_0 to S_1 or vice versa each time as γ bounces off ∂S . We say that γ_1 and γ_2 are equivalent if $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ are homotopic with fixed ends. The set of equivalence classes with the usual operation is the group $\pi_f(S)$. There is an obvious exact sequence

$$1 \rightarrow \pi_f(dS) \rightarrow \pi_f(S) \rightarrow \mathbb{Z}/2 \rightarrow 1$$

A natural class of coverings of S is associated with the subgroups of $\pi_f(S)$. Let $\partial S = \emptyset$ and let $H < \pi_f(S)$ be a subgroup. The imbedding $\dot{S} \subset S$ identifies H with a subgroup of $\pi_1(\dot{S})$. Let \dot{S}^H be the unbranched covering of \dot{S} corresponding to H . Filling in the punctures we obtain a closed polyhedron S^H and the covering $p_H: S^H \rightarrow S$. If $\partial S \neq \emptyset$ we define S^H for any subgroup $H < \pi_f(dS)$ to be $(dS)^H$ with the projection $p_H: (dS)^H \rightarrow dS \rightarrow S$. The coverings $p_H: S^H \rightarrow S$ are branched at the vertices and above ∂S .

Proposition 1. Let S be a closed polyhedron and let H, G be subgroups of $\pi_f(S)$.

1. The inclusion $H < G$ holds if and only if there is a covering $q: S^H \rightarrow S^G$ such that the diagram below commutes

$$\begin{array}{ccc} S^H & \xrightarrow{q} & S^G \\ p_H \downarrow & \text{id} & \downarrow p_G \\ S & \longrightarrow & S \end{array} \quad (1)$$

2. For any subgroup $H < \mathcal{T}_f(S)$ we have $\mathcal{T}_f(S^H) = H$. If H is normal then $p_H: S^H \rightarrow S$ is a regular covering with the group $\mathcal{T}_f(S)/H$ of deck transformations.

The proof as well as the generalization to the case $\partial S \neq \emptyset$ are straightforward and are left to the reader.

Definition 4. The covering of S corresponding to the trivial subgroup of $\mathcal{T}_f(S)$ is called the universal covering and is denoted by \tilde{S} .

Definition 5. Let S be a closed polyhedron. A mapping $\varphi: S \rightarrow \mathbb{C}$ is called a developing map if it is an isometry on every face of S and if for any edge b there is a neighbourhood U of the interior of b such that φ is an isometry on U .

A closed polyhedron S is called developable if there exists a developing map $\varphi: S \rightarrow \mathbb{C}$.

Proposition 2. 1. For any closed polyhedron S the following are equivalent:

- a) S is developable
- b) there exists an isometry $\varphi_0: P_0 \rightarrow \mathbb{C}$ of a face P_0 which continues to a developing map $\varphi: S \rightarrow \mathbb{C}$
- c) any isometry $\varphi_0: P_0 \rightarrow \mathbb{C}$ uniquely extends to a developing map $\varphi: S \rightarrow \mathbb{C}$.

2. Let $O(\mathbb{C})$ denote the group of isometries of \mathbb{C} . Then for any two developing maps φ, ψ of S there is a unique $g \in O(\mathbb{C})$ such that $\psi = g \circ \varphi$.

The proof is obvious and is left to the reader. For general polyhedra S the universal covering \tilde{S} is the minimal developable covering of S .

Definition 6. A polyhedral surface S is called rational if any vertex angle of S is π times a rational number.

Let S be a closed rational polyhedral surface. Let A_i with

angles $2\pi m_i/n_i$ (m_i and n_i are coprime) $i \in I$ be the vertices of S . Denote by $g_i \in \mathcal{T}_f(S)$ the equivalence class of a simple loop around A_i . Let $\hat{\mathcal{T}}(S)$ be the minimal normal subgroup of $\mathcal{T}_f(S)$ containing $g_i^{n_i}, i \in I$.

Definition 7. The corresponding covering \hat{S} of S is called the universal rational covering. If $\partial S \neq \emptyset$ we set $\hat{S} = (dS)^\wedge$.

For any polyhedron S we denote by $\mathcal{T}_1(S)$ the fundamental group of the underlying topological space. We say that S is simply connected if $\mathcal{T}_1(S) = 1$.

Theorem 1. Let S be a compact connected rational polyhedral surface. Let $\alpha_i = \pi m_i/n_i$ $i = 1, \dots, M$ and $\alpha_j = 2\pi m_j/n_j$ $j = M + 1, \dots, N$ be the angles of boundary, respectively interior vertices of S .

1. \hat{S} is a closed connected simply connected noncompact developable polyhedral surface.

2. \hat{S} with its canonical complex structure is isomorphic to \mathbb{C} if and only if all the numerators $m_i = 1$ $i = 1, \dots, N$. Otherwise \hat{S} is isomorphic to the hyperbolic plane \mathbb{H} . The isomorphism $f: \hat{S} \rightarrow \mathbb{H}$ ($\hat{S} \rightarrow \mathbb{C}$) carries the group $\mathcal{T}_r(S) = \mathcal{T}_f(S) / \hat{\mathcal{T}}(S)$ of deck transformations of the covering $\hat{S} \rightarrow S$ into a discrete group G of isometries of \mathbb{H} (respectively \mathbb{C}). If S is oriented the projection $p: \hat{S} \rightarrow S$ is a covering of Riemann surfaces.

3. For any developing map $\varphi: \hat{S} \rightarrow \mathbb{C}$ there is a homomorphism $h: \mathcal{T}_r(S) \rightarrow O(\mathbb{C})$ such that φ is h -equivariant i.e. for any $g \in \mathcal{T}_r(S)$, $\varphi \circ g = h(g) \circ \varphi$. The developing map φ is a holomorphic branched covering. The branching locus is contained in the set of vertices of \hat{S} . The branching number at a vertex $\hat{A} \in \hat{S}$ with angle $2\pi m$ is m .

Proof. By Proposition 1, $\mathcal{T}_f(\tilde{S}) = 1$ therefore $\mathcal{T}_1(\tilde{S}) = 1$. The regular covering $q: \tilde{S} \rightarrow \hat{S}$ induces a homomorphism

$q_* : \hat{\mathcal{T}}(S) \rightarrow \mathcal{T}_1(\hat{S}) \rightarrow 1$. The group $\hat{\mathcal{T}}(S)$ is generated by conjugates of g_1^n (taking dS if necessary we can assume without loss of generality that $\partial S = \emptyset$). Each g_1 fixes some vertex at infinity of \tilde{S} , thus $\hat{\mathcal{T}}(S)$ is generated by transformations with fixed points. Those generate the kernel of q_* , thus $\mathcal{T}_1(\hat{S}) = 1$. Let \hat{A} be a vertex of \hat{S} above A with angle $2\pi m/n$. Going around A once and lifting to \hat{S} we rotate \hat{S} about \hat{A} by $2\pi m/n$. Repeating it n times we make a circle around \hat{A} , thus \hat{A} has a finite number of adjacent faces and its angle is $2\pi m$.

The covering \tilde{S} is developable almost by definition. Choose a reference face P_0 and an imbedding $\varphi_0 : P_0 \rightarrow \mathbb{C}$. Continue φ_0 analytically along a path γ going from P_0 to some face P . The path γ defines a face $\tilde{P} \subset \tilde{S}$ above P . If γ' is another path homotopic to γ we continue φ_0 analytically along the homotopy from γ to γ' and conclude that the imbedding of \tilde{P} depends only on the homotopy class of γ . Thus φ_0 uniquely continues to a developing map $\varphi : \tilde{S} \rightarrow \mathbb{C}$. Any automorphism g of \tilde{S} defines another developing map $\psi = \varphi \circ g$. By proposition 2, there is an isometry h of \mathbb{C} such that $\psi = h \circ \varphi$. Thus $h = h(g)$ is a homomorphism of $\mathcal{T}_f(S)$ into $O(\mathbb{C})$ such that φ is h -equivariant. Let $\hat{A} \in \tilde{S}$ be a vertex above $A \in S$ with angle $2\pi m/n$. Let $g \in \mathcal{T}_f(S)$ correspond to a simple loop around A . Then $h(g)$ is the rotation of \mathbb{C} by $2\pi m/n$ around $\varphi(\hat{A})$. Thus $h(g^n) = 1$ therefore $\hat{\mathcal{T}}(S) \subset \text{Ker } h$. Therefore $\varphi : \tilde{S} \rightarrow \mathbb{C}$ is invariant with respect to $\hat{\mathcal{T}}(S)$ and it descends uniquely to a developing map of S . Since S is developable it is not compact. By now we have proved 1) and 3).

Since the Riemann surface \hat{S} is simply connected and noncompact, by Riemann mapping theorem \hat{S} is either \mathbb{C} or \mathbb{H} . Denote both by D and let $f : \hat{S} \rightarrow D$ be an isomorphism. Any $g \in \mathcal{T}_r(S)$ is an automorphism of \hat{S} therefore it preserves (reverses) the canonical

complex structure of \hat{S} if g preserves (reverses) the orientation. Thus f induces an isomorphism of $\mathcal{T}_r(S)$ onto a discrete subgroup G of the group $O(D)$ of isometries of D and identifies S with the quotient D/G . If S is oriented then $S = D/G$ as a Riemann surface.

It remains to prove that $D = \mathbb{C}$ if and only if $m_i = 1$ $i = 1, \dots, N$. Doubling S if necessary we assume that $\partial S = \emptyset$. Any polyhedral surface S defines an orbifold ([11] ch. 13). The orbifold S has an Euler number $\chi_0(S)$ which is given by ([11])

$$\chi_0(S) = \chi(S) - \sum_{i=1}^N (1 - 1/n_i) \quad (2)$$

where $\chi(S)$ is the Euler characteristic of S . For any closed compact polyhedral surface S with vertex angles α_i $i = 1, \dots, N$ we have

$$\chi(S) = \sum_{i=1}^N (1 - \alpha_i/2\pi) \quad (3)$$

The proof of (3) is an elementary computation and is left to the reader. If S is rational with vertex angles $2\pi m_i/n_i$, (2) and (3) yield

$$\chi_0(S) = \sum_{i=1}^N (1 - m_i)/n_i \quad (4)$$

Thus $\chi_0(S) \leq 0$ and the equality takes place if and only if $m_i = 1$ for all i . Let G_0 be a subgroup of G of a finite index n which acts freely on D . Then $D/G_0 = R \rightarrow S$ is a covering with n sheets and $\chi(R) = n \chi_0(R)$. For a covering $R \rightarrow S$ of orbifolds with n sheets we have ([11], 13.3.4)

$$\chi_0(R) = n \chi_0(S) \quad (5)$$

If $D = \mathbb{C}$ then $\chi(R) = 0$ which is equivalent to $m_i = 1$ $i = 1, \dots, N$

Definition 8. A rational polyhedral surface is called flat (respectively hyperbolic) if the corresponding orbifold is isomorphic to \mathbb{C}/G (resp. \mathbb{H}/G) for some discrete group G of isometries.

The following Corollary has been established in the course of proof of Theorem 1 (compare with [11] , 13.3.6).

Corollary 1. A rational polyhedral surface S is flat if and only if the numerators of all vertex angles of S are equal to 1. Otherwise S is hyperbolic.

Corollary 2. Let S be a polyhedral surface with boundary vertex angles $\pi m_i/n_i$ $i = 1, \dots, M$ and interior vertex angles $2\pi m_i/n_i$ $i = M+1, \dots, N$. Then the Euler number of the orbifold modeled on S is

$$\chi_0(S) = (1/2) \sum_{i=1}^M (1-m_i)/n_i + \sum_{i=M+1}^N (1-m_i)/n_i$$

Proof. If $\partial S = \emptyset$ (6) becomes (4). If $\partial S \neq \emptyset$ (6) follows from (4) for dS and $\chi_0(dS) = 2\chi_0(S)$ by (5).

From now on we identify \hat{S} with D ($D = \mathbb{C}$ or \mathbb{H}) and the group $\mathcal{T}_\Gamma(S)$ of the deck transformations of \hat{S} with $G \subset O(D)$. We choose a developing map $\varphi: D \rightarrow \mathbb{C}$ and let $\Gamma \subset O(\mathbb{C})$ be the image of G under the homomorphism $h: G \rightarrow O(\mathbb{C})$. Choose an origin in \mathbb{C} , let C be the unit circle around it and let $O(\mathbb{C})$ be the group of isometries of \mathbb{C} . Then $O(\mathbb{C})$ contains $O(C)$, the normal subgroup \mathbb{C} of translations and $O(\mathbb{C}) = O(C) \cdot \mathbb{C}$ is the semidirect product. Denote by $\bar{h}: G \rightarrow O(C)$ the composition of h and the homomorphism $O(\mathbb{C}) \rightarrow O(C)$. Let $\bar{\Gamma} = \bar{h}(G) \subset O(C)$. *no face*

— For reasons that will become clear in Section 2, h (resp. \bar{h}) is called the holonomy (resp. restricted holonomy) homomorphism. The group Γ (resp. $\bar{\Gamma}$) is called the holonomy (resp. restricted holonomy) group of S .

Denote by $\mathbb{Z}/n \subset O(\mathbb{C})$ the group of rotations of order n and by $D_n \subset O(\mathbb{C})$ the group generated by reflections in two axes meeting

at the angle π/n . These groups exhaust all finite subgroups of $O(\mathbb{C})$.

Proposition 3. Let S be a compact rational polyhedral surface and let n be the least common multiple of denominators of the vertex angles of S . Then $\overline{\Gamma}$ contains \mathbb{Z}/n . If $\partial S \neq \emptyset$ or if S is not orientable then $\overline{\Gamma}$ contains D_n .

Proof. Let $\partial S = \emptyset$ and let A_i $i = 1, \dots, N$ be the vertices of S with angles $2\pi m_i/n_i$. Let $g_i \in G$ be the simple loops around A_i . Then $h(g_i)$ is the rotation of \mathbb{C} around $\varphi(A_i)$ by $2\pi m_i/n_i$. Thus $h(g_i)$ is a primitive rotation of order n_i . Together they generate \mathbb{Z}/n . Let S be nonorientable and let S' be the orientable 2-sheeted covering of S . Then G is generated by $G(S')$ and an element r such that $r^2 \in G(S')$. Then $h(r)$ reverses orientation so $\overline{h}(r)$ is a reflection. Thus $\overline{\Gamma}$ contains \mathbb{Z}/n and a reflection, so $D_n \subset \overline{\Gamma}$. If $\partial S \neq \emptyset$ then G is generated by $G(\partial S)$ and an element r such that $r^2 = 1$. Then $h(r)$ is a reflection therefore $D_n \subset \overline{\Gamma}$.

Definition 9. A rational polyhedral surface S is called almost integrable if the restricted holonomy group $\overline{\Gamma}$ is finite.

In order to state Proposition 4 we need the notion of the developing map along a path. Let $\gamma(t)$ $0 \leq t \leq 1$ be a piecewise differentiable path on a polyhedron S going through faces P_1, \dots, P_n . Any isometry $\varphi_1: P_1 \rightarrow \mathbb{C}$ uniquely analytically continues to an immersion $\varphi: \bigcup_{i=1}^n P_i \rightarrow \mathbb{C}$ which is an isometry on each P_i . The mapping φ is the developing map along γ .

Proposition 4. 1. A polygon P is almost integrable if and only if the angle between any two sides of P is π -rational.

2. Let a rational polyhedral surface S be homeomorphic to the sphere with n holes, $n = 0, 1, \dots$. Let Q_1, \dots, Q_n be the connected components of ∂S . Choose a side a_i of Q_i and a path γ_i from a_i to a_{i+1} $i = 1, \dots, n-1$. Then S is almost integrable if

and only if developing S along γ_i we map a_i, a_{i+1} into intervals a'_i, a'_{i+1} respectively with a \mathcal{N} -rational angle between them.

3. Let a rational polyhedral surface S be homeomorphic to the projective plane \mathbb{P}^2 . Then S is almost integrable.

Proof. The group G is generated by simple loops g_i around vertices, $\mathcal{N}_1(S)$ and the flipping r if $\partial S \neq \emptyset$. Proving Proposition 2 we have shown that $\bar{h}(g_i) \quad i = 1, \dots, N$ and $\bar{h}(r)$ generate a finite group $\bar{\Gamma}' \subset \bar{\Gamma}$. In case 1, $\bar{h}(\mathcal{N}_1(S)) = 1$ and in case 2 $\bar{h}(\mathcal{N}_1(S))$ is generated by rotations of finite order, thus $\bar{\Gamma}$ is finite. In case 3 let r be the generator of $\mathcal{N}_1(S)$. Then $\bar{h}(g_i)$ generate \mathbb{Z}/n and $\bar{h}(r)$ is a reflection, thus $\bar{\Gamma} = D_n$.

Corollary (of the proof). 1. Let S be a polygon with \mathcal{N} -rational angles between the sides and let n be the least common multiple of denominators. Then $\bar{\Gamma} = D_n$.

2. Let S be a rational polyhedral surface homeomorphic to S^2 (resp. \mathbb{P}^2 or the disc) and let n be the least common multiple of denominators of the vertex angles. Then $\bar{\Gamma} = \mathbb{Z}/n$ (resp. D_n).

Let S be an almost integrable polyhedral surface. Denote by G_0 the kernel of $\bar{h}: G \rightarrow O(C)$ and by R the quotient D/G_0 . Then $R = S^{G_0}$ is the regular covering of S with the group $\bar{\Gamma}$ of deck transformations. In this notation we have

Theorem 2. 1. Let n be the least common multiple of denominators of the vertex angles of S . Then $\bar{\Gamma} = \mathbb{Z}/n'$ if S is orientable and $\partial S = \emptyset$, otherwise $\bar{\Gamma} = D_{n'}$, where n' is divisible by n .

2. R is a compact Riemann surface without boundary and $G_0 = \mathcal{N}_1(R)$. The vertex angles of R are multiples of 2π . The Euler characteristic $\chi(R)$ is equal to $n'\chi_0(S)$ (resp. $2n\chi_0(S)$) if S is closed orientable (resp. otherwise) where $\chi_0(S)$ is the Euler number of S given by (6).

3. The group $\Gamma_0 = h(G_0)$ is a finitely generated group of parallel translations. The developing map $\varphi: D \rightarrow \mathfrak{C}$ yields a commutative diagramm

$$\begin{array}{ccc} D & \xrightarrow{\varphi} & \mathfrak{C} \\ G_0 \downarrow & & \downarrow \Gamma_0 \\ R & & \end{array} \quad (7)$$

where one corner is left empty because in general Γ_0 is not discrete.

Proof. By Proposition 4, the group $\bar{\Gamma}$ is cyclic if S is closed orientable and $\bar{\Gamma}$ is dihedral otherwise. By Proposition 3, $\bar{\Gamma}$ contains Z/n , D_n in the first, resp. second case.

2. If $g \in G$ has fixed points then g has a fixed vertex A . Thus $h(g)$ fixes $\varphi(A)$ therefore $\bar{h}(g) \neq 1$. Analogously if g reverses orientation then $h(g)$ does and $\bar{h}(g) \neq 1$. Thus G_0 acts freely by conformal automorphisms of D inducing a complex structure on $R = D/G_0$ and $\pi_1(R) = G_0$. Besides $\chi(R) = \chi_0(R) = |\bar{\Gamma}| \chi_0(S)$ by (5). The vertex angles of \hat{S} are multiples of 2π and G_0 acts freely, thus vertex angles of S/G_0 are the same.

3. By definition $\bar{h}(G_0) = 1$ so $\bar{\Gamma}_0 \subset \mathfrak{C}$. If p is the genus of R then G_0 and therefore Γ_0 have $2p$ generators. The rest is obvious.

Riemann surface R is called the canonical covering of an almost integrable surface S . In cases of particular interest we can calculate $\bar{\Gamma}$ and the genus $g(R)$ from the angles of S .

Proposition 5. 1. Let S be a polygon and let n be the least common multiple of the denominators of angles between sides of S . Then $\bar{\Gamma} = D_n$. Let π_{m_i/n_i} $i = 1, \dots, M$ be the vertex angles. Then

$$g(R) = 1 + (n/2) \sum_{i=1}^M (m_i - 1)/n_i \quad (8)$$

2. Let S be homeomorphic to the disc and let π_{m_i/n_i} $i = 1, \dots, (2\pi_{m_i/n_i}$ $i = M+1, \dots, N)$ be the boundary (interiour) vertex angles of S . Let n be the least common multiple of n_i . Then $\bar{\Gamma} = D_n$

and

$$g(R) = 1 + n \left[(1/2) \sum_{i=1}^M (m_i - 1)/n_i + \sum_{i=M+1}^N (m_i - 1)/n_i \right] \quad (9)$$

3. Let S be homeomorphic to the sphere (resp. projective plane), let $2\pi m_i/n_i$ $i = 1, \dots, N$ be the vertex angles and let n be the least common multiple of n_i . Then $\bar{r} = 2/n$ (resp. $\bar{r} = D_i$) and

$$g(R) = 1 + (n/2) \sum_{i=1}^N (m_i - 1)/n_i \quad (10)$$

respectively

$$g(R) = 1 + n \sum_{i=1}^N (m_i - 1)/n_i \quad (11)$$

Proof. The groups \bar{r} were calculated in Proposition 4. Using that $2-2g(R) = \chi(R) = |\bar{r}| \chi_0(S)$ and formula (6) we obtain (8) -

2. Invariant surfaces of the billiard flow

If a polygon P is imbedded in \mathbb{C} we define the unit tangent bundle $T(P)$ to be the set of tangent vectors in \mathbb{C} of length one with base points in P and looking into P . The unit tangent bundle $T(S)$ of a polyhedral surface S is made from $T(P_i)$ where P_i runs over the the faces of S , with obvious identifications.

The set $T(S)$ is the phase space of the geodesic flow on S which is modeled on the movement of the billiard ball on S . The ball goes straight inside each face. It bounces off boundary edges in an obvious way. Let the ball come to an edge b between two faces P and Q , at an interior point of b . We imbed $P \cup Q$ in \mathbb{C} and let the ball cross straight from P to Q . We agree to stop the ball at the vertices. We will see later that the trajectory has a natural continuation through a boundary (interior) vertex if and only if its angle is π/n ($2\pi/n$).

Definition 10. The flow on $T(S)$ defined above is the billiard flow of S .

If $x \in S$ is not a vertex, the fiber $T(S)_x$ is isomorphic to C and the measure on $T(S)$ which is locally the product of Lebesgue measures on \mathbb{C} and C is preserved by the flow (cf. [7]). For any $x \in S$ the set of directions which will bring the ball from x to a vertex is countable therefore the set of $v \in T(S)$ which generate finite lifetime trajectories has measure 0.

Let $x, y \in S$ be not vertices and let γ be a piecewise differentiable curve on S going from x to y and avoiding vertices. Developing S on \mathbb{C} along γ we obtain an isomorphism $T(S)|_\gamma = \gamma \times C$ and an isometry $T_\gamma: T(S)_x \rightarrow T(S)_y$ which depends on γ . It is called the parallel translation along γ . Moreover T_γ depends only on the homotopy class $[\gamma] \in \pi_1(S)$.

From now on S is assumed to be almost integrable. Choose a base point $x_0 \in P_0$ and an imbedding $\varphi_0: P_0 \rightarrow \mathbb{C}$. For any $x \in S$ a choice of curve γ from x to x_0 gives an isometry $T_\gamma: T(S)_x \rightarrow C$. Varying γ changes T_γ by the action of \bar{F} on C which yields a mapping $\bar{\Theta}: T(S) \rightarrow C/\bar{F}$. Let now R be the canonical covering of S and identify P_0 with a face of R . Since the restricted holonomy group of R is trivial the construction above yields a mapping $\Theta: T(R) \rightarrow C$. The action of \bar{F} on R obviously lifts to $T(R)$ and we have $T(S) = T(R)/\bar{F}$. The projection $p_*: T(R) \rightarrow T(S)$ of unit tangent bundles commutes with the billiard flows on $T(R)$ and $T(S)$, so the billiard on S is the quotient of the billiard on R .

Proposition 6. 1. The mapping Θ ($\bar{\Theta}$) uniquely extends to a continuous mapping $\Theta: T(R) \rightarrow C$ ($\bar{\Theta}: T(S) \rightarrow C/\bar{F}$) invariant under the geodesic flow and such that the following diagram commutes

$$\begin{array}{ccc} T(R) & \xrightarrow{\Theta} & C \\ p_* \downarrow & & \downarrow \\ T(S) & \xrightarrow{\bar{\Theta}} & C/\bar{F} \end{array} \quad (12)$$

2. For $\theta \in C$ ($\bar{\theta} \in C/\bar{F}$) denote by $R_\theta \subset T(R)$ ($R_{\bar{\theta}} \subset T(S)$) the

set given by the equation $\Theta(v) = \theta$ ($\bar{\Theta}(v) = \bar{\theta}$) and by b_θ^t ($b_{\bar{\theta}}^t$) the flow induced on R_θ ($R_{\bar{\theta}}$) by the billiard flow on $T(R)$ ($T(S)$). Let $d: C \rightarrow C/\bar{F}$ be the projection and say that $\bar{\theta} \in C/\bar{F}$ is regular if any $\theta \in d^{-1}(\bar{\theta})$ has a trivial isotropy subgroup. Then

i) for any $\theta \in C$ the projection $q: T(R) \rightarrow R$ induces a mapping $q_*: R_\theta \rightarrow R$ which is one to one everywhere except over vertices of R with angles $2\pi m > 2\pi$ where it is m to one.

ii) for any regular $\bar{\theta} \in C/\bar{F}$ and any $\theta \in d^{-1}(\bar{\theta})$ the projection p_* induces an isomorphism of R_θ onto $R_{\bar{\theta}}$ which commutes with b_θ^t and $b_{\bar{\theta}}^t$ on R_θ and $R_{\bar{\theta}}$ respectively. Let $\bar{\theta} \in C/\bar{F}$ be not regular and let $\bar{\Gamma}_\theta$ be the isotropy subgroup of $\theta \in d^{-1}(\bar{\theta})$. Then p_* acts on R_θ and $R_{\bar{\theta}} = R_\theta/\bar{\Gamma}_\theta$ with the flows b_θ^t and $b_{\bar{\theta}}^t$ respectively.

Proof. First of all we can define the billiard flow and the parallel translations on $T(\hat{S})$ where everything commutes with the projection $q_*: T(\hat{S}) \rightarrow T(R)$. Parallel translations on $T(\hat{S})$ defined by $\hat{\Theta}: T(\hat{S}) \rightarrow C$. Parallel translations and the geodesic flow on $T(\hat{S})$ are induced by the developing map $\psi: \hat{S} \rightarrow \mathbb{C}$, thus $\hat{\Theta}$ is induced by the composition of $\psi_*: T(\hat{S}) \rightarrow T(\mathbb{C}) = \mathbb{C} \times \mathbb{C}$ and the projection $\mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C}$. Therefore $\hat{\Theta}$ is defined everywhere including the vertices, and commutes with the geodesic flow. All the mappings are compatible with the action of relevant groups, so we have the following commutative diagram

$$\begin{array}{ccccc}
 T(S) & \xrightarrow{G_0} & T(R) & \xrightarrow{\bar{F}} & T(S) \\
 \hat{\Theta} \downarrow & & \Theta \downarrow & & \downarrow \bar{\Theta} \\
 C & \xrightarrow{id} & C & \longrightarrow & C/\bar{F}
 \end{array} \tag{13}$$

which implies (12).

Fixing some $\theta \in C$ we have the constant vector field X_θ on $T(R)$ which is a cross-section of $T(R) \rightarrow R$. The set $\hat{S}_\theta \subset T(\hat{S})$ is the pullback of X_θ by ψ_* . The branching properties of ψ imply

(see Theorem 1) that the projection $\hat{S}_\theta \rightarrow \hat{S}$ is one to one everywhere except above the vertices with angle $2\pi m$ where it is m to 1. Factoring out by the action of G_0 we obtain the same property for the projection $R_\theta \rightarrow R$. Commutative diagram (12) implies that for any $\theta \in C/\bar{\Gamma}$ the action of $\bar{\Gamma}$ on $T(R)$ permutes R_{θ_i} for $\theta_i \in d^{-1}(\bar{\theta})$ according to the action of $\bar{\Gamma}$ on $d^{-1}(\bar{\theta})$. The assertion 2, ii) follows.

Proposition 6 shows that for regular $\bar{\theta} \in C/\bar{\Gamma}$ ($C/\bar{\Gamma}$ has two irregular points if $\bar{\Gamma}$ is dihedral and none otherwise) the billiard flow on the invariant surface $R_{\bar{\theta}} \subset T(S)$ is equivalent to that on $R_\theta \subset T(R)$ for any $\theta \in d^{-1}(\bar{\theta})$. The projection $q_\theta : R_\theta \rightarrow R$ being essentially one to one, we transfer by it the flows b_θ^t from R_θ to R , denote them by the same symbol and call them the billiard flows on R . It is clear from the construction that the flows b_θ^t have singularities at the vertices of R with angles $2\pi m > 2\pi$.

Proposition 7. 1. Let $\theta \in C$ be arbitrary. Let l_θ^t be the linear flow on C in direction θ , i.e. $l_\theta^t z = z + te^{i\theta}$. Let $\varphi_*^{-1} l_\theta^t$ be the pullback of l_θ^t on \hat{S} by the developing map $\varphi : \hat{S} \rightarrow C$. Then the flow $\varphi_*^{-1} l_\theta^t$ is invariant under the action of G_0 and the induced flow on $R = D/G_0$ is b_θ^t .

2. Each vertex of R with angle $2\pi m > 2\pi$ is a singular point of b_θ^t with uniformly spaced m incoming and m outgoing separatrices. Other points of R are nonsingular. The flows b_θ^t are obtained from anyone of them by rotation.

3. The billiard flow on $T(R)$ is isomorphic to $R \times C$ with the flow b_θ^t on $R \times \theta$.

Proof. From the proof of proposition 6 we see that the geodesic flow on \hat{S} is induced by the developing map $\varphi : \hat{S} \rightarrow C$. Proposition 7 follows easily from that, the information about the branching of φ (Theorem 1) and the fact that $h(G_0) = \Gamma_0$ is a group

of translations, thus it leaves l_θ^t invariant.

Recall that a flow b^t on a manifold M smooth everywhere except at a finite number of multisaddle singular points is called minimal (quasiminimal in [13]) if every infinite semitrajectory of b^t is dense in M .

Definition 11. A direction θ is called rational with respect to a set $X \subset \mathbb{C}$ (with respect to a group Γ of translations) if there exists a straight line in direction θ which meets X in two points (which contains points z_1, z_2 such that the vector $z_1 - z_2 \in \Gamma$).

Proposition 8. Let S, R, Γ_0 and b_θ^t be as before. Denote by $V \subset \mathbb{C}$ the image of the branching locus of $\varphi: D \rightarrow \mathbb{C}$. Then the flow b_θ^t has a periodic trajectory (resp. is not minimal) only if θ is rational with respect to Γ_0 (resp. θ is rational with respect to V).

Proof. Let $\gamma(t)$ be a periodic trajectory of b_θ^t with period T . Then the lifting $\hat{\gamma}(t)$ on D has the property $\hat{\gamma}(t+T) = g \hat{\gamma}(t)$ for some $g \in G_0$. The image $\varphi_* \hat{\gamma}(t)$ is the straight line in direction θ , thus the equivariance of φ implies that θ is rational with respect to Γ_0 . Let b_θ^t have a nondense infinite semitrajectory. Arguing as in [3] or as in [13] (they used Theorem 9 of [6]) we conclude that b_θ^t has a trajectory γ going from one singular point to another. Thus both ends of $\hat{\gamma}$ are vertices of \hat{S} and $\varphi(\hat{\gamma})$ is a straight interval with both ends in V .

Corollary. The flow b_θ^t is not minimal for at most a countable set of θ 's.

Proof. The set of branching points of $\varphi: S \rightarrow \mathbb{C}$ is at most countable.

3. Ergodicity and spectrum

We keep notation of the previous sections. We will assume

that the reader is familiar with the basic notions of ergodic theory (cf. [7]). The flow b_θ^t preserves the Lebesgue measure on \mathbb{R} and let U_θ^t be the corresponding group of unitary operators on $L_2(\mathbb{R})$. For any discrete group L of translations the linear flow l_θ^t define a flow on the torus $T = \mathbb{C}/L$ denoted by the same symbol. When L is fixed we call $\theta \in \mathbb{C}$ rational if it is rational with respect to L . It is well known that the flow l_θ^t on the torus is ergodic if and only if it is minimal if and only if θ is an irrational direction. Moreover for irrational θ the flow is uniquely ergodic and for rational θ it is periodic. The spectrum of l_θ^t is discrete. We will generalize these facts to a certain class of billiards.

Denote by $V_0 \subset V$ the set of fixed points of rotations in Γ . Any two points $x, y \in \mathbb{C}$ define a translation vector $y-x$.

Definition 12. A point $z \in \mathbb{C}$ is called rational if there exist $z_0 \in V_0$ and $a_1, a_2 \in \Gamma_0$ such that $z-z_0 = r_1 a_1 + r_2 a_2$ for some rational numbers r_1, r_2 .

Theorem 3. Let S be an almost integrable polyhedral surface such that the monodromy group $\Gamma \subset O(\mathbb{C})$ is discrete and such that all points of V are rational. Then

1. The following are equivalent:
 - i) direction θ is irrational
 - ii) flow b_θ^t is minimal
 - iii) flow b_θ^t is ergodic with respect to the Lebesgue measure
 - iv) b_θ^t is uniquely ergodic
2. Let l_θ^t be the linear flow on $T = \mathbb{C}/\Gamma_0$. For every irrational direction θ the discrete spectrum of b_θ^t coincides with the spectrum of l_θ^t . Continuous spectrum of b_θ^t is empty if and only if S is flat.

3. For every rational θ the flow b_θ^t is periodic.

Proof. By Theorem 2, if Γ_0 is discrete we can complete the com-

mutative diagramm (7)

$$\begin{array}{ccc}
 D & \xrightarrow{\psi} & C \\
 G_0 \downarrow & & \downarrow \Gamma_0 \\
 R & \xrightarrow{p} & T
 \end{array} \tag{14}$$

and obtain a branched covering $p: R \rightarrow T$ which projects b_θ^t on 1_θ^t .

It is well known ([2]) that there are 17 types of discrete groups $\Gamma < O(\mathbb{C})$. The set V_0 of centers of rotations together with the axes of reflexions and sliding reflexions of Γ form a pattern of lines and points in \mathbb{C} . Choose two generators a, b of Γ_0 and take some $0 \in V_0$ for origin in \mathbb{C} . An obvious choice of coordinates x, y identifies a with $(1,0)$, b with $(0,1)$ and Γ_0 with the integer lattice. It is easy to verify case by case that every $A \in V_0$ has rational coordinates and that every axis of (sliding) reflection has two points with rational coordinates. By assumption all points of V have rational coordinates. Observe that all points of $V \setminus V_0$ come from (boundary) vertices of S with angle $(\pi m > \pi)$ $2\pi m > 2\pi$ and that V is invariant under the action of Γ_0 . There is a finite number of orbits of V under the action of Γ_0 , therefore coordinates of $z \in V$ have a common denominator, say N . Let Γ'_0 be the group generated by a/N and b/N , denote by x, y the new coordinates and by $Q = \{0 \leq x, y \leq 1\}$ the fundamental parallelogram of Γ'_0 . Now the points of V have integer coordinates and the whole plane is the union of $Q_{m,n} = \{m-1 \leq x \leq m, n-1 \leq y \leq n\}$. Denote the torus \mathbb{C}/Γ'_0 by T' and let $p': T \rightarrow T'$ and $p_1 = p'p: R \rightarrow T_1$ be the natural projections. The covering p_1 is branched only over $(0,0) \in T_1$, therefore R can be represented as a union of Q_i $i \in I$ such that $p_1: Q_i \rightarrow Q$ is 1 to 1 for all i . Choose a fundamental domain $\tilde{R} \in D$ for R , then $\tilde{R} = \bigcup_I Q_i$. Since the composition of $\psi: D \rightarrow \mathbb{C}$ and the projection $\mathbb{C} \rightarrow T_1$ maps each Q_i onto T_1 , ψ must map Q_i on some Q_π for all $i \in I$. Thus we have represented R by the union of a finite

number of $Q_{m,n}$ where each $Q_{m,n}$ may be taken with some multiplicity. Let us call this object a polygon in the lattice Γ'_0 and denote it by R' . Identifications on the boundary of R' are made by elements of Γ_0 .

This construction represents the billiard flow b_θ^t on R by the linear flow in the direction θ on R' . When the ball reaches a boundary edge of R' it gets transferred by some $g \in \Gamma_0$ to another edge and then it keeps rolling in the same direction. We think of Q_i $i \in I$ as parallelograms on the plane and let X be the union of their bases. Thus $X = [0,1) \times I$. The flow b_θ^t that started on the base of some Q_i reaches after a certain lapse of time T_θ the tops of two parallelograms of R' which are identified with the bases of say Q_j and $Q_{j'}$ (see fig. 2). Every point of the base of Q_i comes to its destination translated by the rotation number α of the flow l_θ^t . Besides the $[0,1-\alpha)$ part of Q_i comes to Q_j and the $[1-\alpha, 1)$ part of Q_i comes to $Q_{j'}$. Denote by β_α the rotation of $[0,1)$ by α . The previous remarks can be summarized as follows. There exists a function $w(x)$ on $[0,1)$ with values in the permutation group Σ_I of I symbols, constant on $[0,1-\alpha)$ and $[1-\alpha, 1)$, so that the first return to X map τ_α is $\tau_\alpha(x, i) = (x+\alpha, w(x)i)$ for $x \in [0,1)$, $i \in I$. Thus τ_α is the extension of rotation β_α with the squewing function $w(x)$. The flow b_θ^t is the suspension of τ_α with the constant time of return function. To show 1) we use an unpublished result of W. Veech.

Theorem (W. Veech). Any extension τ_α of an irrational rotation β_α with the squewing function constant on $[0,1-\alpha)$ and $[1-\alpha, 1)$ is ergodic if and only if it is minimal. The ergodic components of τ_α are in 1 to 1 correspondence with the orbits on I of the group $W \subset \Sigma_I$ generated by $w([0,1-\alpha))$ and $w([1-\alpha, 1))$.

Ergodicity or minimality of τ_α is equivalent to the ergodi-

city or minimality of b_θ^t . If the direction θ is irrational then b_θ^t is minimal by Proposition 8 and α is irrational. Thus by Veech's Theorem, τ_α is ergodic. For finite extensions of irrational rotations ergodicity is equivalent to unique ergodicity which proves 1). If θ is rational then l_θ^t is periodic and b_θ^t being a finite extension of it is too.

Pulling back by $p: R \rightarrow T$ we imbed $L_2(T)$ isometrically into $L_2(R)$ as functions constant on the fibers. We want to show that $L_2(R) \ominus L_2(T)$ does not contain eigenfunctions of b_θ^t . Every non-trivial eigenfunction of b_θ^t comes from an eigenfunction $f(x, i)$ of τ_α with an eigenvalue λ . Then $g(x, i, j) = f(x, i)\bar{f}(x, j)$ is a fixed function of the transformation τ'_α of $[0, 1) \times I \times I$ given by $(x, i, j) \rightarrow (x + \alpha, w(x)i, w(x)j)$. Thus by Veech's Theorem, $g(x, i, j) = h(i, j)$ where $h(wi, wj) = h(i, j)$ for any $w \in W$. Since α is irrational, W acts transitively on I and let W_0 be the isotropy subgroup of 1. Set $h(i) = h(1, i)$, then h is W_0 -invariant, $|h| = 1$ and

$$f(x, i) = h(i)f(x, 1) \tag{15}$$

Applying τ_α to both sides of (15) we get

$$f(x + \alpha, w(x)i) = h(w(x)i)f(x + \alpha, 1) = \lambda f(x, i) = \lambda h(i)f(x, 1).$$

Thus for each generator $w \in W$ there is an interval $J \subset [0, 1)$ such that for all $x \in J$

$$h(wi)h(i)^{-1} = \lambda f(x, 1)f(x + \alpha, 1)^{-1} \tag{16}$$

Since the right hand side of (16) does not depend on i , there is $c(w) \in \mathbb{C}$ such that

$$h(wi) = c(w)h(i) \tag{17}$$

for all $i \in I$. It follows from (17) that $w \rightarrow c(w)$ is a character of W trivial on W_0 . Its kernel W_1 is a normal subgroup containing W_0 . Let $m = [W:W_1]$ and let $I_1 \subset I$ be the W_1 -orbit of 1. Then the

character c induces an isomorphism $c: W/W_1 \rightarrow Z/m$, the group W_1 has m orbits I_1, \dots, I_m of the same magnitude $|I|/m$ and Z/m permutes them cyclically. Therefore if we represent elements of I by pairs (i, j) $1 \leq i \leq m, 1 \leq j \leq |I|/m$ we have $\tau_\alpha(x, i, j) = (x + \alpha, c(w(x))i, w_1(x, i)j)$. The transformation $\sigma_\alpha: (x, i) \rightarrow (x + \alpha, c(w(x))i)$ is a m -point extension of ρ_α and τ_α is a $|I|/m$ -point extension of σ_α . The eigenfunction f of τ_α is the pullback of an eigenfunction \tilde{f} of σ_α and σ_α is equivalent to the rotation by α of $[0, m)$.

Going back to the flows we conclude that there is a torus \tilde{T} , a branched covering $\tilde{p}: R \rightarrow \tilde{T}$ with $|I|/m$ sheets and a covering $q: \tilde{T} \rightarrow T_1$ (unbranched) with m sheets such that $p_1 = q\tilde{p}$ and the diagram below commutes

$$\begin{array}{ccccc}
 R & \xrightarrow{\tilde{p}} & \tilde{T} & \xrightarrow{q} & T_1 \\
 b_\theta^t \downarrow & & \downarrow l_\theta^t & & \downarrow l_\theta^t \\
 R & \xrightarrow{p} & T & \xrightarrow{q} & T_1
 \end{array} \tag{18}$$

Thus we know that any eigenfunction of b_θ^t is the pullback from some torus \tilde{T} . Let $\tilde{T} = \mathbb{C}/\tilde{\Gamma}_0$. Since R is the polygon \tilde{R} with sides identified by some g_i , the group $\tilde{\Gamma}_0$ must contain all g_i . But g_i generate Γ_0 , thus $\Gamma_0 \subset \tilde{\Gamma}_0$ which implies that the covering $\tilde{p}: R \rightarrow \tilde{T}$ factors through p and some $\tilde{q}: T \rightarrow \tilde{T}$ where all the mappings commute with the relevant flows. This argument shows that T is the maximal torus covered by R . Therefore we can pull an eigenfunction back from \tilde{T} to R in two steps: first from \tilde{T} to T and then from T to R . Continuous spectrum of b_θ^t is empty if and only if $L_2(R) = L_2(T)$ which means that $p: R \rightarrow T$ is an isomorphism.

Corollary 1. Let P be a polygon with \mathcal{N} -rational angles between the sides. Assume that the obstacles of P have no slits. If the group Γ generated by reflections in the sides of P is discrete then the billiard flow b_θ^t is uniquely ergodic for any irrational

direction θ .

Proof. Since P has no vertices with angles $\pi m > \pi$. we have $V = V_0$. Points of V_0 are rational with respect to Γ by definition, therefore assumptions of Theorem 3 are satisfied.

Corollary 2. Let P be a polygon with π -rational angles between sides such that the group Γ generated by reflections in the sides of P is discrete. Let A be a vertex of a slit and continue the line of the slit until it crosses a side b of P . Let B be the point of intersection. If the vector $B-A$ is rational with respect to Γ (for every slit in P) then b_θ^t is uniquely ergodic for any irrational direction θ .

Proof. The slit and the side b define axes of reflections in Γ . It was mentioned in the proof of Theorem 3 that those are rational lines. The intersection of rational lines is a rational point. Since B is rational and $B-A$ is by assumption, A is rational and we are in the range of Theorem 3.

Corollary 3. Let S be a rational polyhedral surface homeomorphic to the sphere or the disc. Assume that S has no boundary (interior) vertices with angles πm ($2\pi m$), $m > 1$. Let the holonomy group Γ of S be discrete. Then for any irrational direction θ the billiard flow b_θ^t is uniquely ergodic.

Proof. By Proposition 4, 2), S is almost integrable. Apply Theorem 3.


Corollary 4. Let S be any Platonic solid except the dodecahedron. Then the billiard flow on S in any irrational direction is uniquely ergodic.

Proof. Developing S on the plane one sees that Γ is not discrete only for the dodecahedron.

REFERENCES


1. B. Eckhardt, J. Ford, F. Vivaldi. Analytically solvable dynamical systems which are not integrable. Preprint 1983.
2. E. S. Fedorov. Zap. Imp. S. Petersburg Mineral. OBSC. 28 (1971), 345 - 390 (in Russian).
3. R. H. Fox, R. B. Kershner. Concerning the transitive properties of geodesics on a rational polyhedron. Duke Math. J. 2 (1936), 147 - 150.
4. E. Gutkin. Geodesic flows on rational polyhedra. Preprint 1982.
5. H. Masur. Interval exchange transformations and measured foliations. Ann. Math. 115 (1982), 169 - 200.
6. A. Mayer. Trajectories on the closed orientable surfaces. Mat. Sbornik 12 (1943), 71 - 84 (in Russian).
7. Y. Sinai. Introduction into ergodic theory. Princeton University Press: Princeton 1976.
8. P. T. Richens, M. V. Berry. Pseudointegrable systems in classical and quantum mechanics. Physica 2D (1981), 495 - 512.
9. C. Rodenberg. Geodatische Linien auf Polyederflächen. Rend. Circ. Mat. Palermo 23 (1907), 107 - 125.
10. P. Stäckel. Geodatische Linien auf Polyederflächen. Rend. Circ. Mat. Palermo 22 (1906), 141 - 151.
11. W. Thurston. The geometry and topology of three-manifolds. Mimeographed notes: Princeton 1980.
12. W. A. Veech. Gauss measures for transformations on the space of interval exchange maps. Ann. Math. 115 (1982), 201 - 242.
13. A. N. Zemlyakov, A. B. Katok. Topological transitivity of billiards in polygons. Matem. Zametki 18:2 (1975), 291 - 300 (in Russian, English transl. in Mat. Notes 18:2 (1976), 760 - 764.

fig. 1



The shaded regions are obstacles. Obstacle 2 has a slit and obstacle 3 is just a slit.

fig. 2



Boundary edges are identified by parallel translation.
The shaded area does not belong to R' , it comes from an obstacle.