


Max-Planck-Institut für Mathematik Bonn

On symplectic vortex equations over a compact orbifold
Riemann surface

by

Hironori Sakai



On symplectic vortex equations over a compact orbifold Riemann surface

Hironori Sakai

Max-Planck-Institut für Mathematik
Vivatsgasse 7
53111 Bonn
Germany

On symplectic vortex equations over a compact orbifold Riemann surface

Hironori Sakai

Abstract

Making use of theory of differentiable stacks, we study symplectic vortex equations over a compact orbifold Riemann surface. We discuss the category of representable morphisms from a compact orbifold Riemann surface to a quotient stack. After that we define symplectic vortex equations over a compact orbifold Riemann surface. We also discuss the moduli space of solutions to the equations for linear actions of the circle group on the complex plane.

1 Introduction

The symplectic vortex equations (SVE for short) are introduced by Salamon and Mundet independently. Let G be a compact Lie group and \mathfrak{g} its Lie algebra. We choose an invariant inner product on \mathfrak{g} to identify \mathfrak{g} with its dual space \mathfrak{g}^\vee . Suppose that G acts on a symplectic manifold M in a Hamiltonian fashion with a moment map $\mu : M \rightarrow \mathfrak{g}$. Let $P \rightarrow \Sigma$ be a principal G -bundle over a compact Riemann surface Σ . The SVE (over Σ) are defined by the PDEs

$$\bar{\partial}_A u = 0, \quad *F_A + \mu(u) = 0$$

for G -equivariant map $u : P \rightarrow M$ and a connection A on P . Here $\bar{\partial}_A$ is the anti-holomorphic part of the covariant derivative d_A , F_A is the curvature of A , and $*$ is the Hodge $*$ -operator. (We also have to fix a volume form on Σ and an almost complex structure on M compatible with the symplectic structure and the G -action.) Under some analytic assumptions, we can define Hamiltonian invariants as integrations over the moduli space of solutions to the SVE. The invariants are called symplectic vortex invariants.

The most important feature of the SVE is that we can obtain the equation of pseudo-holomorphic curve from Σ to the Marsden–Weinstein quotient $M//G$ as a certain limit of the SVE. This limit is called the adiabatic limit, and this is the most important idea for an identity between Gromov–Witten invariants (with fixed marked points) of the smooth quotient $M//G$ and symplectic vortex invariants for the Hamiltonian G -space M [8].

It is conjectured that the identity extends to Gromov–Witten invariants of the orbifold Marsden–Weinstein quotient $M//G$ (after a suitable extension of theory of SVE) [15, §12.7].

Gromov–Witten invariants of closed symplectic orbifold are defined as integration over a certain compactification of the moduli space of representable pseudo-holomorphic maps from orbifold Riemann surfaces [4]. Therefore it is natural to think about an extension of the SVE such that its adiabatic limit is the equation of representable pseudo-holomorphic maps. But if we put everything into orbifold settings (e.g. orbifold, etc.) for SVE, then the theory becomes extremely complicated.

To solve this problem, we make use of theory of differentiable stacks. Roughly speaking, the symplectic vortex equations over a Riemann surface Σ can be thought as “differential equations” of maps from Σ to the quotient stack $[M/G]$: a (representable) morphism of stacks $\varphi : \Sigma \rightarrow [M/G]$ corresponds to the 2-cartesian square

$$\begin{array}{ccc} P & \xrightarrow{u} & M \\ \pi^P \downarrow & & \downarrow \pi^M \\ \Sigma & \xrightarrow{\varphi} & [M/G], \end{array}$$

where u is a G -equivariant map and $\pi^P : P \rightarrow \Sigma$ is a principal G -bundle over Σ .

In this paper, we see that we can replace Σ with a stack corresponding to compact orbifold Riemann surface in the above diagram. One of the advantages of the replacement is that the upper part of the above diagram still belongs to the category of smooth manifolds. Finally we find that we can use the same PDEs for an extension of SVE after several observations on differentiable stacks.

This paper is organised as follows. In §2, we review briefly theory of differentiable stacks. In §3, we discuss the category of representable morphisms from an orbifold Riemann surface to a quotient stack. We show that the category is equivalent to a certain category similar to the category of principal G -bundles. In §4, we fix notations for an orbifold Riemann surface by using statements developed in §3, and we consider (representable) pseudo-holomorphic maps from a compact orbifold Riemann surface Σ to a (orbifold) Marsden–Weinstein quotient. After that we define SVE over Σ and discuss the moduli space of solutions to the SVE for the case when the circle group S^1 acts on the complex plane \mathbb{C} .

Notations We fix the following notations through this paper.

We denote by ψ any right action map: if we have a right G -space M , then ψ is defined by

$$\psi : M \times G \rightarrow M; (m, g) \mapsto \psi(m, g) = \psi_g(m) = m \cdot g = mg.$$

Let \mathfrak{g} be the Lie algebra of a Lie group G . For a G -space M and $\xi \in \mathfrak{g}$, we define the infinitesimal action $\xi^M \in \text{Vect}(M)$ by

$$\xi^M(m) = \begin{cases} \left. \frac{d}{dt} \right|_{t=0} m \cdot \exp(t\xi) & \text{(if } M \text{ is a right } G\text{-space),} \\ \left. \frac{d}{dt} \right|_{t=0} \exp(-t\xi) \cdot m & \text{(if } M \text{ is a left } G\text{-space)} \end{cases} \quad (m \in M).$$

Here $\text{Vect}(M)$ is the space of vector fields on M . The above convention is the same as Bott-Tu [3] and opposite to Cieliebak–Gaio–Salamon [5]. Set

$$\mathfrak{g}^M(m) = \{\zeta^M(m) \in T_m M \mid \zeta \in \mathfrak{g}\} \text{ and } \mathfrak{g}^M = \{\zeta^M(m) \in TM \mid \zeta \in \mathfrak{g}, m \in M\}.$$

If the G -action is locally free, then \mathfrak{g}^M is a subbundle of the tangent bundle TM .

For a (2-)fibred product $\mathfrak{X}_1 \times_{F_1} \times_{F_2} \mathfrak{X}_2$, the i -th ($i = 1, 2$) projection is denoted by pr_i .

2 Differentiable stacks

We review theory of differentiable stacks and geometric objects (e.g. differential forms) on stacks. First we recall the bicategory of Lie groupoids and bibundles. Secondly we introduce the 2-category of stacks over the smooth category and geometric objects on Deligne–Mumford stacks. Finally we describe an orbifold Riemann surface as a Deligne–Mumford stack.

2.1 Lie groupoids and bibundles

For details of Lie groupoids and bibundles, see Lerman [12], Metzler [16], Moerdijk [20] and Moerdijk–Mrčun [17].

A **groupoid** is a category whose arrows are all invertible. We write $X_1 \rightrightarrows X_0$ for a groupoid whose class of objects and class of arrows are X_0 and X_1 respectively. The source map and the target map are always denoted by src and tgt respectively: for $a \in \text{Hom}(x, y)$, $\text{src}(a) = x$ and $\text{tgt}(a) = y$. We also write $a : x \rightarrow y$ for such an arrow.

Definition 2.1. A groupoid $X_1 \rightrightarrows X_0$ is called a **Lie groupoid** if

- (i) both X_0 and X_1 are smooth manifolds,
- (ii) both the source and the target map are surjective submersions, and
- (iii) the composition map

$$X_1 \times_{\text{src}} \times_{\text{tgt}} X_1 \mapsto X_1; (a, b) \mapsto ab,$$

the unit map $x \mapsto 1_x$ ($x \in X_0$) and the inverse map $a \mapsto a^{-1}$ ($a \in X_1$) are all smooth.

Let $X_1 \rightrightarrows X_0$ be a Lie groupoid. For an object $x \in X_0$, the submanifold

$$(X_1 \rightrightarrows X_0)_x = \{a \in X_1 \mid \text{src}(a) = x = \text{tgt}(a)\}$$

of X_1 is a Lie group. The Lie group is called the **stabiliser group** at x .

Two objects $x, y \in X_0$ are said to lie in the same orbit ($x \sim y$) if there is an arrow $a : x \rightarrow y$. The quotient space of X_0 with respect to the equivalent relation \sim is called the **underlying space** of $X_1 \rightrightarrows X_0$ and denoted by X_0/X_1 . If $x \sim y$, then the stabiliser group at x is naturally isomorphic to the stabiliser group at y .

Example 2.2. Let G be a Lie group and M a right G -space. The **action groupoid** $M \rtimes G$ of the G -action on M is a Lie groupoid $M \times G \rightrightarrows M$ whose structure maps are given as follows: $\text{src}(m, g) = mg$, $\text{tgt}(m, g) = m$, $(m, g)(mg, h) = (m, gh)$, $1_m = (m, 1)$ and $(m, g)^{-1} = (mg, g^{-1})$ for $g, h \in G$ and $m \in M$. The stabiliser group at $m \in M$ is the ordinary stabiliser group at m and the underlying space of the action groupoid $M \rtimes G$ is the quotient space M/G .

Definition 2.3.

- A Lie groupoid $X_1 \rightrightarrows X_0$ is said to be **proper** if $(\text{src}, \text{tgt}) : X_1 \rightarrow X_0 \times X_0$ is a proper map.
- A Lie groupoid is called an **étale groupoid** if the source and the target maps are étale (locally diffeomorphic).

Next we recall bibundles. Regarding a bibundle as an arrow between Lie groupoids, we obtain the bicategory of Lie groupoids and bibundles.

Definition 2.4. A **right action** of a Lie groupoid $Y_1 \rightrightarrows Y_0$ on a manifold P consists of two maps $\alpha : P \rightarrow Y_0$ and

$$\mu : P_{\alpha} \times_{\text{tgt}} Y_1 \rightarrow P; (p, a) \mapsto pa$$

satisfying that $\alpha(pa) = \text{src}(a)$, $p1_{\alpha(p)} = p$ and $p(ab) = (pa)b$ for any $(a, b) \in Y_{1, \text{src}} \times_{\text{tgt}} Y_1$ and $p \in P$ with $\alpha(p) = \text{tgt}(a)$. The map α is called the **anchor map**.

A left action of a Lie groupoid can be defined in a similar way.

Definition 2.5. Let $Y_1 \rightrightarrows Y_0$ be a Lie groupoid. A **principal** $(Y_1 \rightrightarrows Y_0)$ -**bundle** over a manifold B is a smooth map $\pi : P \rightarrow B$ with a right action of $Y_1 \rightrightarrows Y_0$ on P satisfying the following conditions.

- The map π is $(Y_1 \rightrightarrows Y_0)$ -invariant, i.e. $\pi(pa) = \pi(p)$ for any $(p, a) \in P_{\alpha} \times_{\text{tgt}} Y_1$.
- The map π is a surjective submersion.
- The map

$$P_{\alpha} \times_{\text{tgt}} Y_1 \rightarrow P_{\pi} \times_{\pi} P; (p, a) \mapsto (p, pa)$$

is a diffeomorphism. Here α is the anchor map for the action.

Definition 2.6. Let $X_1 \rightrightarrows X_0$ and $Y_1 \rightrightarrows Y_0$ be two Lie groupoids. A **bibundle** (or Hilssum-Skandalis morphism) $f = (\alpha_L, P, \alpha_R)$ is a manifold P equipped with

- a left action of $X_1 \rightrightarrows X_0$ with an anchor map α_L and
- a right action of $Y_1 \rightrightarrows Y_0$ with an anchor map α_R

satisfying following conditions.

- The map $\alpha_L : P \rightarrow X_0$ is $(Y_1 \rightrightarrows Y_0)$ -invariant and $\alpha_R : P \rightarrow Y_0$ is $(X_1 \rightrightarrows X_0)$ -invariant.
- The actions of $X_1 \rightrightarrows X_0$ and $Y_1 \rightrightarrows Y_0$ are compatible, i.e. $(ap)b = a(pb)$ for any $a \in X_1$, $p \in P$ and $b \in Y_1$ with $\text{src}(a) = \alpha_L(p)$ and $\alpha_R(p) = \text{tgt}(b)$.

(iii) The map $\alpha_L : P \rightarrow X_0$ is a principal $(Y_1 \rightrightarrows Y_0)$ -bundle over X_0 .

Given two bibundles $f = (\alpha_L, P, \alpha_R)$ and $f' = (\alpha'_L, P', \alpha'_R)$, a **2-isomorphism** from f to f' is a diffeomorphism $\varphi : P \rightarrow P'$ which commutes with both $(X_1 \rightrightarrows X_0)$ - and $(Y_1 \rightrightarrows Y_0)$ -actions.

The composition of bibundles is not strictly associative, but associative up to a (canonical) 2-isomorphism. Therefore Lie groupoids and bibundles do not form a category, but form a bicategory. We denote by \mathfrak{Bi} the bicategory of Lie groupoids and bibundles.

Any bibundle induces a continuous map between underlying spaces and a group homomorphism between stabiliser groups. Suppose that we have a bibundle $f = (\alpha_L, P, \alpha_R)$ from $X_1 \rightrightarrows X_0$ to $Y_1 \rightrightarrows Y_0$. For any $p, p' \in P$ and $a \in X_1$ satisfying $\text{src } a = \alpha_R(p)$ and $\text{tgt } a = \alpha_R(p')$, there is a unique $b \in Y_1$ such that $ap = pb$, $\text{tgt } b = \alpha_R(p')$ and $\text{src } b = \alpha_R(p)$. This fact guarantees that the map

$$F^f : X_0/X_1 \rightarrow Y_0/Y_1; [\alpha_R(p)] \mapsto [\alpha_L(p)] \quad (p \in P)$$

is a well-defined continuous map and the map

$$(X_1 \rightrightarrows X_0)_{\alpha_R(p)} \rightarrow (Y_1 \rightrightarrows Y_0)_{\alpha_L(p)}; a \mapsto b$$

is a group homomorphism. (Two isomorphic bibundles induces the same continuous map and the group homomorphism.) If two groupoids $X_1 \rightrightarrows X_0$ and $Y_1 \rightrightarrows Y_0$ are equivalent in the bicategory \mathfrak{Bi} , then the underlying spaces X_0/X_1 and Y_0/Y_1 are homeomorphic and the group structures of stabiliser groups at the points corresponding under the homeomorphism are isomorphic. Therefore topological structures on the underlying space and stabiliser groups are invariants of the category \mathfrak{Bi} .

2.2 Differentiable stacks

For details of stacks, see Behrend–Xu [2], Heinloth [9], Metzler [16] and Lerman [12].

Roughly speaking, a stack is a category \mathfrak{X} equipped with a functor from \mathfrak{X} to a base site satisfying some conditions including a so-called “gluing condition”. Here a **site** is a category equipped with a Grothendieck topology. Theory of stacks can be discussed for several base sites, but we only use the category \mathfrak{Diff} of smooth manifolds and smooth maps as a base site in this paper.

We define a Grothendieck topology on \mathfrak{Diff} as follows. For $U \in \mathfrak{Diff}$, a family $\{f_i : U_i \rightarrow U\}_i$ of smooth maps to U is called a **covering family** of U if $f_i : U_i \rightarrow U$ is a local diffeomorphism for each i and the total map $\bigsqcup U_i \rightarrow U$ is surjective. The function K assigning to each object $U \in \mathfrak{Diff}$ the collection $K(U)$ of covering families define a basis of Grothendieck topology (cf Metzler [16, Definition 5]). The basis of Grothendieck topology generates a Grothendieck topology on the category \mathfrak{Diff} .

Remark 2.7. For general definition of Grothendieck topology see Metzler [16] and Vistoli [25]. Note that a basis for a Grothendieck topology (pretopology) is called a Grothendieck topology in Vistoli [25, Definition 2.24]. We follow Behrend–Xu [2] for definition of covering families in $\mathfrak{D}\text{iff}$ and the same definition is used in Lerman [12, Remark 2.17] and Lerman–Malkin [13]. Metzler uses a slightly different definition for covering family, but the Grothendieck topology is the same as our Grothendieck topology.

Definition 2.8. A **category fibred in groupoids (over $\mathfrak{D}\text{iff}$)** is a category \mathfrak{X} equipped with a functor $F_{\mathfrak{X}} : \mathfrak{X} \rightarrow \mathfrak{D}\text{iff}$ satisfying the following conditions.

- (i) For any $f : V \rightarrow U$ in $\mathfrak{D}\text{iff}$ and any $x \in \mathfrak{X}$ with $F_{\mathfrak{X}}(x) = U$, there is an arrow $a : y \rightarrow x$ in \mathfrak{X} such that $F_{\mathfrak{X}}(a) = f$.
- (ii) Suppose we have two arrows $a_1 : y_1 \rightarrow x$ and $a_2 : y_2 \rightarrow x$. For any smooth map $f : F_{\mathfrak{X}}(y_1) \rightarrow F_{\mathfrak{X}}(y_2)$ with $F_{\mathfrak{X}}(a_2) \circ f = F_{\mathfrak{X}}(a_1)$ there is a unique arrow $b : y_1 \rightarrow y_2$ such that $a_2 b = a_1$ and $F_{\mathfrak{X}}(b) = f$.

The functor $F_{\mathfrak{X}} : \mathfrak{X} \rightarrow \mathfrak{D}\text{iff}$ is called the **base functor of the category fibred in groupoids**.

Omitting the base functor $F_{\mathfrak{X}}$, we often say that \mathfrak{X} is a category fibred in groupoids. Moreover $F_{\mathfrak{X}}$ is described as “ $\underline{\quad}$ ”. Namely $\underline{x} = F_{\mathfrak{X}}(x)$ for $x \in \mathfrak{X}$ and $\underline{a} = F_{\mathfrak{X}}(a)$ for an arrow a in \mathfrak{X} .

The collection of all categories fibred in groupoids form a 2-category:

Definition 2.9. A **morphism of categories fibred in groupoids** from \mathfrak{X} to \mathfrak{Y} is a functor $\varphi : \mathfrak{X} \rightarrow \mathfrak{Y}$ which commutes with the base functors.

For two morphisms of categories fibred in groupoids $\varphi_1, \varphi_2 : \mathfrak{X} \rightarrow \mathfrak{Y}$, a **2-morphism of categories fibred in groupoids** is a natural transformation $\alpha : \varphi_1 \rightarrow \varphi_2$ such that the horizontal composition $F_{\mathfrak{Y}} * \alpha$ is the identity transformation of $F_{\mathfrak{X}}$.

The **fibre of \mathfrak{X} over $U \in \mathfrak{D}\text{iff}$** is the groupoid \mathfrak{X}_U with objects $\{x \in \mathfrak{X} \mid \underline{x} = U\}$ and arrows $\{\text{arrow } a \text{ in } \mathfrak{X} \mid \underline{a} = \text{id}_U\}$.

Given a category fibred in groupoids \mathfrak{X} , for every object $x \in \mathfrak{X}$ and every $f : V \rightarrow \underline{x}$ in $\mathfrak{D}\text{iff}$ we choose an arrow $a : y \rightarrow x$ in \mathfrak{X} such that $\underline{a} = f$ (cf Definition 2.8). The object y is called the **pullback of the object x via f** and denoted by f^*x .

Let $a : x \rightarrow y$ be an arrow in a fibre \mathfrak{X}_U and $f : V \rightarrow U$ a smooth map. Then we have two arrows $b_x : f^*x \rightarrow x$ and $b_y : f^*y \rightarrow y$ which we have chosen. Then by Definition 2.8, there is a unique arrow $\tilde{a} : f^*x \rightarrow f^*y$ such that \tilde{a} belongs to \mathfrak{X}_V . The arrow \tilde{a} is called the **pullback of the arrow a via f** and denoted by f^*a .

We can assign the descent category $\mathfrak{X}_{\{U_i \rightarrow U\}}$ to each covering family $\{f_i : U_i \rightarrow U\}$ and there is a natural functor from the fibre \mathfrak{X}_U to the descent category $\mathfrak{X}_{\{U_i \rightarrow U\}}$. The category fibred in groupoids is called a **stack (over $\mathfrak{D}\text{iff}$)** if for any covering family the natural functor is an equivalence of categories.

The collection of all stacks (over $\mathfrak{D}iff$) form a 2-subcategory $\mathfrak{StD}iff$ of the 2-category of categories fibred in groupoids. We regard two stacks as the same stack if they are equivalent.

Remark 2.10. Some authors say two stacks which are equivalent are isomorphic. But we follow the ordinary terminology of theory of 2-categories.

Let G be a Lie group. For a right G -space M , we can define a stack $[M/G]$ as follows. An object of $[M/G]$ is a pair (π, ε) of a principal G -bundle $\pi : P \rightarrow U$ over a manifold U and a G -equivariant map $\varepsilon : P \rightarrow M$. The object (π, ε) is often denoted by $U \xleftarrow{\pi} P \xrightarrow{\varepsilon} M$. An arrow from $V \xleftarrow{\pi^Q} Q \xrightarrow{\varepsilon^Q} M$ to $U \xleftarrow{\pi^P} P \xrightarrow{\varepsilon^P} M$ is a pair (f, \tilde{f}) of smooth map $f : V \rightarrow U$ and a G -equivariant map $\tilde{f} : Q \rightarrow P$ which makes the following diagram commutative:

$$\begin{array}{ccccc} V & \xleftarrow{\pi^Q} & Q & \xrightarrow{\varepsilon^Q} & M \\ f \downarrow & & \tilde{f} \downarrow & & \nearrow \\ U & \xleftarrow{\pi^P} & P & \xrightarrow{\varepsilon^P} & M \end{array}$$

The category $[M/G]$ is a category fibred in groupoid with the base functor

$$[M/G] \rightarrow \mathfrak{D}iff; \begin{cases} (U \xleftarrow{\pi} P \xrightarrow{\varepsilon} M) \mapsto U & \text{for objects,} \\ (f, \tilde{f}) \mapsto f & \text{for arrows.} \end{cases}$$

For any object $U \xleftarrow{\pi} P \xrightarrow{\varepsilon} M$ over U and any smooth map $f : U \rightarrow V$, the diagram $V \xleftarrow{\text{pr}_1} V_f \times_{\pi} P \xrightarrow{\varepsilon \circ \text{pr}_2} M$ gives an object of over V and we have an arrow in $[M/G]$.

$$\begin{array}{ccccc} V & \xleftarrow{\text{pr}_1} & V_f \times_{\pi} P & \xrightarrow{\varepsilon \circ \text{pr}_2} & M \\ f \downarrow & & \text{pr}_2 \downarrow & & \nearrow \\ U & \xleftarrow{\pi} & P & \xrightarrow{\varepsilon} & M \end{array}$$

We choose the object $V \xleftarrow{\text{pr}_1} V_f \times_{\pi} P \xrightarrow{\varepsilon \circ \text{pr}_2} M$ as the pullback of $U \xleftarrow{\pi} P \xrightarrow{\varepsilon} M$ via f . We can show that the category fibred in groupoids $[M/G]$ is a stack. The stack is called the **quotient stack** associated to the G -space.

An arbitrary manifold M can be considered as a quotient stack $[M/\{1\}]$: An object is a smooth map f whose target is M and an arrow from $g : V \rightarrow M$ to $f : U \rightarrow M$ is a smooth map $a : V \rightarrow U$ with $f \circ a = g$. Thanks to the 2-Yoneda embedding, the category $\mathfrak{D}iff$ can be embedded into the 2-category of stacks $\mathfrak{StD}iff$. We identify every manifold M with the stack $[M/\{1\}]$. A stack \mathfrak{X} is said to be **representable** if there is a manifold equivalent to the stack \mathfrak{X} .

In $\mathfrak{StD}iff$ we can always take a 2-fibre product.

Definition 2.11. Let $\varphi_1 : \mathfrak{Y}_1 \rightarrow \mathfrak{X}$ and $\varphi_2 : \mathfrak{Y}_2 \rightarrow \mathfrak{X}$ be two morphisms of stacks. The **(2-)fibre product** $\mathfrak{Y}_1 \times_{\varphi_1, \varphi_2} \mathfrak{Y}_2$ is a stack defined as follows. The class of objects is

$$\{(y_1, y_2, a) \mid y_1 \in \mathfrak{Y}_1, y_2 \in \mathfrak{Y}_2, \underline{y} = \underline{y}' = U, a : \varphi(y) \rightarrow \varphi'(y_2) \text{ in } \mathfrak{X}_U\}.$$

An arrow from (y_1, y_2, a) to (y'_1, y'_2, a') is a pair of arrows $(b_1 : y_1 \rightarrow y'_1, b_2 : y_2 \rightarrow y'_2)$ satisfying $\underline{b_1} = \underline{b_2}$ and $a' \circ \varphi_1(b_1) = \varphi_2(b_2) \circ a$. The base functor $\mathfrak{Y}_1 \times_{\varphi_1 \times \varphi_2} \mathfrak{Y}_2 \rightarrow \mathfrak{D}iff$ is given by $(\underline{y_1}, \underline{y_2}, \underline{a}) = \underline{y_1}$ for objects and $(\underline{b_1}, \underline{b_2}) = \underline{b_1}$ for arrows. In particular $\mathfrak{Y}_1 \times \mathfrak{Y}_2$ is defined as the 2-fibre product $\mathfrak{Y}_1 \times_{\varphi_1 \times \varphi_2} \mathfrak{Y}_2$ for $\varphi_1 : \mathfrak{Y}_1 \rightarrow \text{pt}$ and $\varphi_2 : \mathfrak{Y}_2 \rightarrow \text{pt}$.

Definition 2.12. A morphism of stacks $\varphi : \mathfrak{Y} \rightarrow \mathfrak{X}$ is said to be **representable** if for any morphism of stack F from a manifold U to \mathfrak{X} the fibre product $U_F \times_{\varphi} \mathfrak{Y}$ is representable.

Definition 2.13. A representable morphism φ from a manifold X to a stack \mathfrak{X} is called an **atlas** (resp. **étale atlas**) of \mathfrak{X} if for any morphism F from a manifold U to the stack \mathfrak{X} the projection map from the representable stack $U_F \times_{\varphi} X$ to U is a surjective submersion (resp. surjective local diffeomorphism).

Let $\varphi : X_0 \rightarrow \mathfrak{X}$ be an atlas of a stack \mathfrak{X} . Choosing a manifold X_1 which is equivalent to the representable stack $X_0 \times_{\varphi} X_0$, we obtain a 2-cartesian diagram:

$$\begin{array}{ccc} X_1 & \xrightarrow{s} & X_0 \\ t \downarrow & & \downarrow \varphi \\ X_0 & \xrightarrow{\varphi} & \mathfrak{X}. \end{array}$$

We can see that $X_1 \rightrightarrows X_0$ is a Lie groupoid whose source map and target map are s and t , respectively. The Lie groupoid $X_1 \rightrightarrows X_0$ is called the **presentation** of \mathfrak{X} associated to the atlas.

Two presentations of the same stack associated to two atlases are equivalent in $\mathfrak{B}i$, therefore any invariant of the category $\mathfrak{B}i$ (e.g. the underlying space, the stabiliser groups, etc.) is also an invariant of differentiable stacks. See Lerman [12] for more details.

Definition 2.14. A stack \mathfrak{X} is called a **differentiable stack** (resp **Deligne–Mumford stack**) if \mathfrak{X} has an atlas (resp étale atlas) and the presentation associated to the atlas is proper.

Definition 2.15. Let $X_1 \rightrightarrows X_0$ be the presentation of \mathfrak{X} . Suppose that both X_1 and X_0 have constant dimensions. We define the **dimension** of \mathfrak{X} by $2 \dim X_0 - \dim X_1$.

Definition 2.16. Let $X_1 \rightrightarrows X_0$ be the presentation of \mathfrak{X} associated to an atlas.

- A differentiable stack \mathfrak{X} is said to be **compact** if so is X_0/X_1 .
- A differentiable stack \mathfrak{X} is said to be **connected** if so is X_0/X_1 .

Definition 2.17. A stack \mathfrak{X} is said to be **type-R** if

- (i) the stack \mathfrak{X} is a compact and connected Deligne–Mumford stack, and
- (ii) a generic stabiliser group of \mathfrak{X} is trivial.

Example 2.18. Let G be a compact Lie group and M a (right) G -space. The quotient stack $[M/G]$ always has an atlas $\pi^M : M \rightarrow [M/G]$ called the **natural projection**. For objects,

$\pi^M(m : U \rightarrow M) = (U \xleftarrow{\text{pr}_1} U \times G \xrightarrow{\varepsilon} M)$, where $\varepsilon(u, g) = m(u)g$. Therefore the quotient stack is differentiable. The action groupoid $M \rtimes G$ is a presentation of the atlas. If the G -action is locally free, then the quotient stack is Deligne–Mumford. (This does not mean that the natural projection is an étale atlas.)

2.3 Differential forms and vector fields over stacks

We describe geometric objects (e.g. differential forms, vector fields, etc) on Deligne–Mumford stacks using sheaves over the stack. The general definition of a sheaves over stacks can be found in Behrend–Xu [2] and Metzler [16]. Kashiwara and Schapira [11] explain a general theory of sheaves over sites. Note that a Grothendieck topology on a stack can be induced by the Grothendieck topology on the category $\mathfrak{D}\text{iff}$.

Differential forms over a differentiable stack [1, 2] We define the sheaf $\Omega_{\mathfrak{X}}^k$ of differential k -forms on a differentiable stack \mathfrak{X} as follows. For an object $x \in \mathfrak{X}$ over U , $\Omega_{\mathfrak{X}}^k$ is defined by the space $\Omega^k(U)$ of k -forms on U . For an arrow $a : x \rightarrow y$ in \mathfrak{X} with $\underline{a} : U \rightarrow V$, we assign the pullback map $\underline{a}^* : \Omega^k(V) \rightarrow \Omega^k(U)$ to $\Omega_{\mathfrak{X}}^k(y) \rightarrow \Omega_{\mathfrak{X}}^k(x)$. This presheaf $\Omega_{\mathfrak{X}}^k : \mathfrak{X} \rightarrow (\mathbb{R}\text{-Vect})$ satisfies the conditions to be a sheaf over \mathfrak{X} . Since the exterior derivative d commutes with pullbacks, d makes the abelian sheaves $\Omega_{\mathfrak{X}}^*$ a complex. The complex $\Omega_{\mathfrak{X}}^*$ is called the (big) de Rham complex of \mathfrak{X} . The de Rham cohomology of \mathfrak{X} is defined as the hypercohomology of \mathfrak{X} with values in $\Omega_{\mathfrak{X}}^*$.

Using the presentation $X_1 \rightrightarrows X_0$ associated to an atlas of \mathfrak{X} , we can calculate the de Rham cohomology $H_{\text{DR}}^*(\mathfrak{X})$ more explicitly. The set of (global) k -forms on \mathfrak{X} is given by

$$\Omega^k(\mathfrak{X}) = \{\eta \in \Omega^k(X_0) \mid \text{src}^* \eta = \text{tgt}^* \eta\}.$$

The de Rham cohomology is isomorphic to the cohomology of the complex $(\Omega^k(\mathfrak{X}), d)$.

Remark 2.19. In general, the set of global sections of a sheaf \mathcal{F} over a stack \mathfrak{X} is defined by applying the global section functor Γ to \mathcal{F} . If \mathfrak{X} is differentiable, the set is canonically isomorphic to the equaliser of the two restriction maps $\mathcal{F}(X_1) \rightrightarrows \mathcal{F}(X_0)$, where $X_1 \rightrightarrows X_0$ is a presentation associated to an atlas $X_0 \rightarrow \mathfrak{X}$.

Remark 2.20. For any vector space V , we can define the sheaf of differential k -forms on a differentiable stack \mathfrak{X} in a similar way.

Vector fields over a Deligne–Mumford stack [13] If we have a Deligne–Mumford stack \mathfrak{X} , then we can define the tangent sheaf $\mathcal{T}_{\mathfrak{X}}$ as follows. Let $Z_1 \rightrightarrows Z_0$ be the presentation of \mathfrak{X}

associated to the étale atlas $p : Z_0 \rightarrow \mathfrak{X}$. For $x \in \mathfrak{X}_U$ we have the following cartesian diagrams:

$$\begin{array}{ccccc} V_1 & \rightrightarrows & V_0 & \xrightarrow{q} & U \\ \downarrow f_1 & & \downarrow f_0 & & \downarrow x \\ Z_1 & \rightrightarrows & Z_0 & \xrightarrow{p} & \mathfrak{X} \end{array}$$

Note that $V_1 \rightrightarrows V_0$ is an étale groupoid and a presentation of U . Since $Z_1 \rightrightarrows Z_0$ is étale, $\text{src}^* TZ_0 = \text{tgt}^* TZ_0 = TZ_1$. According to descent theory, there is a vector bundle $E \rightarrow U$ (which is unique up to isomorphisms) such that the pullback q^*E is isomorphic to $f_0^* TZ_0$. Then define $\mathcal{T}_{\mathfrak{X}}(x)$ as the set of global sections of the bundle $E \rightarrow U$. For an arrow $a : x \rightarrow y$, $\mathcal{T}_{\mathfrak{X}}(a) : \mathcal{T}_{\mathfrak{X}}(y) \rightarrow \mathcal{T}_{\mathfrak{X}}(x)$ is naturally defined. Moreover the tangent sheaf $\mathcal{T}_{\mathfrak{X}}$ is independent of the choice of the étale atlas. The set of global sections of the tangent sheaf is denoted by $\text{Vect}(\mathfrak{X})$ and an element of $\text{Vect}(\mathfrak{X})$ is called a vector field on \mathfrak{X} .

Proposition 2.21 (Lerman–Malkin [13]). *Let $X_1 \rightrightarrows X_0$ be a presentation of an atlas $p : X_0 \rightarrow \mathfrak{X}$ of a Deligne–Mumford stack \mathfrak{X} .*

- (i) *Let A be the pullback bundle of $\ker(d \text{src}) \rightarrow X_1$ along the unit map $X_0 \rightarrow X_1$. Then the map $d \text{tgt} : A \rightarrow TX_0$ is an injective bundle map. (Thus we regard A as a subbundle of TX_0 .)*
- (ii) *The (small) sheaf over X_0 induced by the tangent sheaf $\mathcal{T}_{\mathfrak{X}}$ is the sheaf of sections of the bundle TX_0/A .*
- (iii) *The (small) sheaf over X_1 induced by the tangent sheaf $\mathcal{T}_{\mathfrak{X}}$ is the sheaf of sections of the bundle $TX_1/(\ker d \text{src} + \ker d \text{tgt})$.*
- (iv) *The set of vector fields $\text{Vect}(\mathfrak{X})$ is explicitly given by the following quotient vector space:*

$$\mathcal{V}/\{(v_1, v_0) \in \mathcal{V} \mid v_1 \in \ker(d \text{src}) + \ker(d \text{tgt})\}.$$

Here $\mathcal{V} = \{(v_1, v_0) \in \text{Vect}(X_1) \times \text{Vect}(X_0) \mid d \text{src} \circ v_1 = v_0 \circ \text{src}, d \text{tgt} \circ v_1 = v_0 \circ \text{tgt}\}$ and $\text{Vect}(X_i)$ is the space of the (ordinary) vector fields on X_i ($i = 0, 1$). In particular, if the atlas $p : X_0 \rightarrow \mathfrak{X}$ is étale, then $\text{Vect}(\mathfrak{X})$ is isomorphic to \mathcal{V} .

Symplectic forms [13] Let $X_1 \rightrightarrows X_0$ be a presentation of a Deligne–Mumford stack \mathfrak{X} associated to an atlas $p : X_0 \rightarrow \mathfrak{X}$. For a vector field $(v_1, v_0) \in \text{Vect}(\mathfrak{X})$, we can define the interior product

$$\iota(v_1, v_0) : \Omega^k(\mathfrak{X}) \rightarrow \Omega^{k-1}(\mathfrak{X}); \eta \mapsto \iota(v_0)\eta.$$

A 2-form $\omega \in \Omega^2(\mathfrak{X})$ on \mathfrak{X} is said to be nondegenerate if the map

$$\text{Vect}(\mathfrak{X}) \rightarrow \Omega^1(\mathfrak{X}); (v_1, v_0) \mapsto \iota(v_1, v_0)\omega$$

is a linear isomorphism. This is equivalent to the condition that $\ker \omega = A$. A closed nondegenerate 2-form ω on \mathfrak{X} is called a symplectic form on \mathfrak{X} and a symplectic Deligne–Mumford stack is a pair (\mathfrak{X}, ω) of a Deligne–Mumford stack \mathfrak{X} and a symplectic form ω on \mathfrak{X} .

Orientations and almost complex structures Let \mathfrak{X} be a differentiable stack of dimension n . The stack \mathfrak{X} is said to be orientable if $\Omega^n(\mathfrak{X})$ is isomorphic to $\mathcal{O}(\mathfrak{X})$ as an $\mathcal{O}(\mathfrak{X})$ -module. For an orientable stack \mathfrak{X} we define

$$\Omega^n(\mathfrak{X})_{\text{gen}} = \{\omega \in \Omega^n(\mathfrak{X}) \mid \mathcal{O}(\mathfrak{X})\omega = \Omega^n(\mathfrak{X})\}.$$

and we denote by $\mathcal{O}(\mathfrak{X})_+$ the space of (global) positive functions. An orientation of \mathfrak{X} is a choice of an element of the quotient set $\Omega^n(\mathfrak{X})_{\text{gen}}/\mathcal{O}(\mathfrak{X})_+$. For a Deligne–Mumford stack \mathfrak{X} , an orientation is nothing but a pair of orientations of Z_0 and Z_1 compatible with the source and target maps.

Let \mathfrak{X} be a Deligne–Mumford stack. An almost complex structure is an endomorphism $J : \mathcal{T}_{\mathfrak{X}} \rightarrow \mathcal{T}_{\mathfrak{X}}$ of $\mathcal{O}_{\mathfrak{X}}$ -modules satisfying $J^2 = -\text{id}_{\mathcal{T}_{\mathfrak{X}}}$. For an atlas $p : X_0 \rightarrow \mathfrak{X}$, an almost complex structure J on \mathfrak{X} induces a complex structure of the vector bundle TX_0/A compatible with the source and the target map. In particular, for an étale atlas $\zeta : Z_0 \rightarrow \mathfrak{X}$, an almost complex structure J induces almost complex structures on Z_0 and Z_1 which are compatible with the source and target maps. Here $Z_1 \rightrightarrows Z_0$ is a presentation associated to the atlas $\zeta : Z_0 \rightarrow \mathfrak{X}$. In particular, an almost complex structure induces a canonical orientation on \mathfrak{X} .

Integral over a Deligne–Mumford stack [1] A partition of unity of a proper étale groupoid $Z_1 \rightrightarrows Z_0$ is a smooth function ρ on Z_0 such that $\text{src}^* \rho$ has proper support with respect to $\text{tgt} : Z_1 \rightarrow Z_0$ and $\text{tgt}_! \text{src}^* \rho = 1$. Not all proper étale groupoid has a partition of unity, but for any proper étale groupoid $Z_1 \rightrightarrows Z_0$, we can find a proper étale groupoid $Z'_1 \rightrightarrows Z'_0$ which is equivalent to $Z_1 \rightrightarrows Z_0$ and has a partition of unity.

Let \mathfrak{X} be an oriented and compact Deligne–Mumford stack with an étale atlas $\zeta : Z_0 \rightarrow \mathfrak{X}$ and $Z_1 \rightrightarrows Z_0$ a presentation associated to the atlas. If the presentation $Z_1 \rightrightarrows Z_0$ has a partition of unity $\rho : Z_0 \rightarrow \mathbb{R}$, then we can define an integral

$$\int_{\mathfrak{X}} : \Omega^n(\mathfrak{X}) \rightarrow \mathbb{R}; \quad \int_{\mathfrak{X}} \omega = \int_{Z_0} \rho \omega.$$

Here $n = \dim \mathfrak{X} = \dim Z_0$. This gives rise to a linear map $H^n(\mathfrak{X}) \rightarrow \mathbb{R}$ as usual. Moreover the Poincaré duality holds: The pairing

$$H_{\text{DR}}^p(\mathfrak{X}) \otimes H_{\text{DR}}^{n-p}(\mathfrak{X}) \rightarrow \mathbb{R}; \quad ([\omega], [\eta]) \mapsto \int_{\mathfrak{X}} \omega \wedge \eta$$

is nondegenerate.

2.4 An orbifold Riemann surface as a stack

In this paper we consider an orbifold Riemann surface as a Deligne–Mumford stack. We describe how to construct the stack in this subsection. We basically follow terminologies of Moerdijk–Prunk [18, 19] for orbifolds.

A compact orbifold Riemann surface is a quadruple $\Sigma = (\Sigma, j, \mathbf{z}, \mathbf{m})$ of a closed Riemann surface Σ , a complex structure j on Σ , a k -tuple $\mathbf{z} = (z_1, \dots, z_k)$ of distinct points on Σ and a k -tuple $\mathbf{m} = (m_1, \dots, m_k)$ of positive integers. We define the **multiplicity** m of $p \in \Sigma$ as follows:

$$m = \begin{cases} m_i & \text{if } p = z_i \text{ for some } i, \\ 1 & \text{else.} \end{cases}$$

A point $p \in \Sigma$ is said to be **smooth** if its multiplicity is 1.

First we recall an orbifold atlas for the compact orbifold Riemann surface Σ . For a point p of multiplicity m , we choose a complex coordinate system $w_p : V_p \rightarrow \mathbb{C}$ centred at p so that $V_p \setminus \{p\}$ consists only of smooth points. We also choose an open disc $U_p \subset \mathbb{C}$ centred at the origin and a holomorphic map $\pi_p : U_p \rightarrow \Sigma$ satisfying $\pi_p(0) = p$, $\pi_p(U_p) = V_p$ and $(w_p \circ \pi_p)(z) = z^m$. Here z is the standard coordinate function on \mathbb{C} . Then the group

$$G_p = \left\{ e^{\frac{2\pi i k}{m}} \in \mathbb{C} \mid k = 0, \dots, m-1 \right\} \cong \mathbb{Z}/m\mathbb{Z}$$

acts on U_p as multiplication and the map π_p induces a homeomorphism from U_p/G_p to V_p . Then (U_p, G_p, π_p) is an orbifold chart for $\pi_p(U_p)$ and the family $\mathcal{U} = \{(U_p, G_p, \pi_p)\}_{p \in \Sigma}$ is an orbifold atlas on Σ .

We consider the tangent bundle $T\Sigma$. This is a complex orbibundle over Σ . Choosing a metric compatible with the complex structure, we obtain the S^1 -bundle P consisting of unit tangent vectors. It is easy to see that the total space of the bundle is a smooth manifold. Moreover the circle group S^1 acts locally freely on P as complex multiplication (on the right).

Since the orbifold Σ can be obtained as the quotient of P with respect to the S^1 -action, we regard the quotient stack $\mathfrak{X} = [P/S^1]$ as the orbifold Riemann surface Σ . By the construction of \mathfrak{X} , the stack \mathfrak{X} is type-R.

By Proposition 2.21, the sheaf over P induced by the tangent sheaf is the sheaf of sections of $N = TP/\mathfrak{g}^P$. Here \mathfrak{g} is the Lie algebra of S^1 . The bundle N is an S^1 -equivariant vector bundle, and $N/S^1 \rightarrow P/S^1$ is naturally identified with the tangent bundle $T\Sigma$. Therefore the vector bundle N is equipped with a complex structure compatible with the S^1 -action. The complex structure gives the almost complex structure on \mathfrak{X} which will be denoted by j as well.

3 The category of representable morphisms

In this section, we discuss the category of representable morphisms of stacks from a (type-R) stack to a quotient stack. The category of (representable) morphisms from a manifold to a quotient stack is a prototype. Let M be a G -space. By the 2-Yoneda lemma and the definition of quotient stacks, any (representable) morphism of stacks φ from a manifold U to the quotient

stack $[M/G]$ corresponds to the 2-cartesian square

$$\begin{array}{ccc} P & \xrightarrow{u} & M \\ \pi^P \downarrow & & \downarrow \pi^M \\ U & \xrightarrow{\varphi} & [M/G]. \end{array}$$

Here $\pi^P : P \rightarrow U$ is a principal G -bundle and $u : P \rightarrow M$ is a G -equivariant map. To replace U with a (type-R) stack \mathfrak{X} , we discuss a certain category $\mathfrak{P}^G(\mathfrak{X})$ which is similar to the category of principal G -bundles. After that we show that the group of automorphisms of the category for the type-R base \mathfrak{X} is the same as the ordinary gauge group action. Finally we discuss the category of representable morphisms from \mathfrak{X} to the quotient stack $[M/G]$.

3.1 The category $\mathfrak{P}^G(\mathfrak{X})$

For a stack \mathfrak{X} and a compact Lie group G , we construct a category $\mathfrak{P}^G(\mathfrak{X})$ as follows.

Definition 3.1. An object of $\mathfrak{P}^G(\mathfrak{X})$ is a pair of a morphism $\pi^P : P \rightarrow \mathfrak{X}$ from a right G -space P to \mathfrak{X} and a 2-morphisms of stacks $\sigma : \pi^P \circ \text{pr}_1 \rightarrow \pi^P \circ \psi$

$$\begin{array}{ccc} P \times G & \xrightarrow{\psi} & P \\ \text{pr}_1 \downarrow & & \downarrow \pi^P \\ P & \xrightarrow{\pi^P} & \mathfrak{X}. \end{array} \quad \begin{array}{c} \nearrow \sigma \\ \searrow \end{array}$$

satisfying the following conditions.

- (i) The morphism $\pi^P : P \rightarrow \mathfrak{X}$ is an atlas.
- (ii) The morphism of stacks

$$P \times G \rightarrow P \times_{\pi^P} P; \quad \begin{cases} (p, g) \rightarrow (p, pg, \sigma(p, g)) & \text{for objects,} \\ f \rightarrow (f, f) & \text{for arrows} \end{cases}$$

is an equivalence. (Therefore the above square is 2-cartesian.)

- (iii) For any $p \in P, g, h \in G$ with $\underline{p} = \underline{g} = \underline{h}$, the following identity holds.

$$\sigma(p, gh) = \sigma(pg, h)\sigma(p, g). \quad (1)$$

Remark 3.2. If $(\pi : P \rightarrow \mathfrak{X}, \sigma)$ is an object of $\mathfrak{P}^G(\mathfrak{X})$, the stack \mathfrak{X} is equivalent to the quotient stack $[P/G]$. On the other hand, for any quotient stack $[P/G]$ we can naturally obtain an object $(\pi^P : P \rightarrow [P/G], \sigma^P)$ of $\mathfrak{P}^G([P/G])$. (For $(p, g) \in (P \times G)_U$, $\sigma^P(p, g) : U \times G \rightarrow U \times G$ is defined by $\sigma^P(p, g)(u, a) = (u, g(u)^{-1}a)$.) The point of the definition of $\mathfrak{P}^G(\mathfrak{X})$ is that we can stick with a fixed base stack \mathfrak{X} .

Definition 3.3. An arrow between two objects $(\pi : P \rightarrow \mathfrak{X}, \sigma)$ and $(\pi' : P' \rightarrow \mathfrak{X}, \sigma')$ of $\mathfrak{P}^G(\mathfrak{X})$ is a pair of a G -equivariant diffeomorphism $\varphi : P \rightarrow P'$ and a 2-morphism of stacks $\tau : \pi \rightarrow \pi' \circ \varphi$ such that for any $(p, g) \in P \times G$ the identity $\tau(pg)\sigma(p, g) = \sigma'(\varphi(p), g)\tau(p)$ hold.

$$\begin{array}{ccc} P & \xrightarrow{\varphi} & P' \\ & \searrow \pi & \swarrow \pi' \\ & \mathfrak{X} & \end{array} \quad \begin{array}{c} \xrightarrow{\tau} \\ \end{array}$$

For two arrows $(\varphi, \tau) : (\pi, \sigma) \rightarrow (\pi', \sigma')$ and $(\varphi', \tau') : (\pi', \sigma') \rightarrow (\pi'', \sigma'')$ in $\mathfrak{P}^G(\mathfrak{X})$, the composition $(\varphi', \tau')(\varphi, \tau)$ is defined by $(\varphi' \circ \varphi, (\tau' \circ \text{id}_\varphi) \bullet \tau)$, where \circ is the horizontal composition and \bullet is the vertical composition. The identity arrow of $(\pi : P \rightarrow \mathfrak{X}, \sigma)$ is $(\text{id}_P, \text{id}_\pi)$. Then $\mathfrak{P}^G(\mathfrak{X})$ form a groupoid.

Remark 3.4. For a manifold X , $\mathfrak{P}^G(X)$ is the category of principal G -bundles over X .

Remark 3.5. A G -stack is defined by Romagny [22] as a generalisation of G -spaces. If we regard a stack \mathfrak{X} as a trivial G -stack, then any object (π, σ) of $\mathfrak{P}^G(\mathfrak{X})$ gives a morphisms of G -stack. Moreover for any arrow $(\varphi, \tau) : (\pi, \sigma) \rightarrow (\pi', \sigma')$ in $\mathfrak{P}^G(\mathfrak{X})$, the 2-morphism of stacks τ gives a 2-morphism of G -stacks $\pi \rightarrow \pi' \circ \varphi$.

Let $(\pi^P : P \rightarrow \mathfrak{X}, \sigma)$ be an object of $\mathfrak{P}^G(\mathfrak{X})$. For any representable morphism $F : \mathfrak{Y} \rightarrow \mathfrak{X}$, we can construct an object of $\mathfrak{P}^G(\mathfrak{Y})$ by taking a pullback. Now we have a manifold Q and the 2-cartesian square

$$\begin{array}{ccc} Q & \xrightarrow{\varepsilon} & P \\ \pi \downarrow & & \downarrow \pi^P \\ \mathfrak{Y} & \xrightarrow{F} & \mathfrak{X} \end{array} \quad \begin{array}{c} \nearrow \alpha \\ \end{array} \quad (2)$$

Define $\varepsilon' : Q \times G \rightarrow P$ by $\varepsilon'(q, g) = \varepsilon(q)g$. Then the map

$$\text{Ob}(Q \times G) \rightarrow \text{Ar}(\mathfrak{X}); (q, g) \mapsto \sigma(\varepsilon(q), g)\alpha(q)$$

gives a 2-morphism of stacks $F \circ (\pi \circ \text{pr}_1) \rightarrow \pi^P \circ \varepsilon'$. Since the above diagram is 2-cartesian, there is a unique smooth map $\psi' : Q \times G \rightarrow Q$ and a 2-morphism of stacks $\sigma' : \pi \circ \text{pr}_1 \rightarrow \pi \circ \psi'$ such that $\varepsilon \circ \psi' = \varepsilon'$ and

$$\alpha(\psi'(q, g))F(\sigma'(q, g)) = \sigma(\varepsilon(q), g)\alpha(q) \quad (3)$$

for any $(q, g) \in Q \times G$.

Proposition 3.6. *The maps $\psi' : Q \times G \rightarrow Q$ defines a right G -action on Q .*

The following lemma is a direct conclusion of the universality of the 2-cartesian square (2).

Lemma 3.7. Let $\tilde{\pi} : R \rightarrow \mathfrak{Y}$ and $\tilde{\varepsilon} : R \rightarrow P$ be morphisms of stacks. Suppose we have two pairs (f_1, β_1) and (f_2, β_2) of a smooth map $f_i : R \rightarrow Q$ with $\tilde{\varepsilon} = \varepsilon \circ f_i$ and a 2-morphism of stacks $\beta_i : \tilde{\pi} \rightarrow \pi \circ f_i$ ($i = 1, 2$):

$$\begin{array}{ccccc}
 & & R & & \\
 & \swarrow \tilde{\pi} & \downarrow f_i & \searrow \tilde{\varepsilon} & \\
 \mathfrak{Y} & \xleftarrow{\pi} & Q & \xrightarrow{\varepsilon} & P.
 \end{array}$$

If $\alpha(f_1(r))F(\beta_1(r)) = \alpha(f_2(r))F(\beta_2(r))$ for any $r \in R$, then $f_1 = f_2$ and $\beta_1 = \beta_2$.

Proof of Proposition 3.6. First we show that $\psi'(q, 1) = q$ holds for any $q \in Q$. Define $s : Q \rightarrow Q \times G$ by $s(q) = (q, 1)$. Let $(f_1, \beta_1) = (\psi \circ s, \sigma' \circ \text{id}_s)$ and $(f_2, \beta_2) = (\text{id}_Q, \text{id}_\pi)$, then we obtain the following 2-commutative diagram ($i = 1, 2$):

$$\begin{array}{ccccc}
 & & Q & & \\
 & \swarrow \pi & \downarrow f_i & \searrow \varepsilon & \\
 \mathfrak{Y} & \xleftarrow{\pi} & Q & \xrightarrow{\varepsilon} & P.
 \end{array}$$

Then the identity (3) implies $\alpha(f_1(q))F(\beta_1(q)) = \alpha(f_2(q))F(\beta_2(q))$ for any $q \in Q$. Thus Lemma 3.7 implies $f_1 = f_2$ i.e. $\psi'(q, 1) = q$.

Next we show that $\psi'(\psi'(q, g), h) = \psi'(q, gh)$ holds for any $(q, g, h) \in Q \times G \times G$. Define $s_1 : Q \times G \times G \rightarrow Q \times G$ by $s_1(q, g, h) = (q, gh)$ and $s_2 : Q \times G \times G \rightarrow Q \times G$ by $s_2(q, g, h) = (\psi'(q, g), h)$. Let $(f_1, \beta_1) = (\psi' \circ s_1, \sigma' \circ \text{id}_{s_1})$ and $(f_2, \beta_2) = (\psi' \circ s_2, (\sigma' \circ \text{id}_{s_2}) \bullet (\sigma' \circ \text{pr}_{12}))$, then we obtain the following 2-commutative diagram ($i = 1, 2$):

$$\begin{array}{ccccc}
 & & Q \times G \times G & & \\
 & \swarrow \pi \circ \text{pr}_1 & \downarrow f_i & \searrow (q, g, h) \mapsto \varepsilon(q)gh & \\
 \mathfrak{Y} & \xleftarrow{\pi} & Q & \xrightarrow{\varepsilon} & P.
 \end{array}$$

Then the identities (1) and (3) imply $\alpha(f_1(q))F(\beta_1(q)) = \alpha(f_2(q))F(\beta_2(q))$ for any $q \in Q$. Thus Lemma 3.7 implies $f_1 = f_2$ i.e. $\psi'(q, gh) = \psi'(\psi'(q, g), h)$. \square

From now on, $\psi'(q, g)$ is denoted by qg . Then we can write the identity (3) in the form

$$\alpha(qg)F(\sigma'(q, g)) = \sigma(\varepsilon(q), g)\alpha(q) \quad (4)$$

for any $(q, g) \in Q \times G$. The identity $\varepsilon \circ \psi' = \varepsilon'$ implies that the smooth map $\varepsilon : Q \rightarrow P$ is G -equivariant. Note that we also obtain the following identity

$$\sigma'(q, gh) = \sigma'(qg, h)\sigma'(q, g) \quad (5)$$

for any $(q, g, h) \in Q \times G \times G$ in the proof of the above proposition.

Proposition 3.8. *The pair $(\pi : Q \rightarrow \mathfrak{Y}, \sigma')$ is an object of $\mathfrak{P}^G(\mathfrak{Y})$.*

Proof. It is sufficient to see that the morphism of stacks

$$\Phi : Q \times G \rightarrow Q_{\pi \times \pi} Q; \begin{cases} (q, g) \rightarrow (q, qg, \sigma'(q, g)) & \text{for objects,} \\ a \rightarrow (a, a) & \text{for arrows} \end{cases}$$

is an equivalence (over each manifold U). First we show that the morphism Φ is fully faithful over U . Let $(q_1, g_1), (q_2, g_2) \in Q \times G$ be objects over U . Suppose that we have an arrow $(q_1, q_1g_1, \sigma'(q_1, g_1)) \rightarrow (q_2, q_2g_2, \sigma'(q_2, g_2))$ in $(Q_{\pi \times \pi} Q)_U$. Since such an arrow is an identity arrow, we have $(q_1, q_1g_1, \sigma'(q_1, g_1)) = (q_2, q_2g_2, \sigma'(q_2, g_2))$. Moreover this implies that $(\varepsilon(q_1), \varepsilon(q_1)g_1, \sigma(\varepsilon(q_1), g_1)) = (\varepsilon(q_2), \varepsilon(q_2)g_2, \sigma(\varepsilon(q_2), g_2)))$ in $(P_{\pi^P} \times_{\pi^P} P)_U$. Therefore $g_1 = g_2$ and the morphism Φ is fully faithful over U .

Next we show that the morphism Φ is essentially surjective over U . Let (q_1, q_2, a) be an object of $(Q_{\pi \times \pi} Q)_U$. Since $(\varepsilon(q_1), \varepsilon(q_2), \alpha(q_2)F(a)\alpha(q_1)^{-1})$ is an object of $(P_{\pi^P} \times_{\pi^P} P)_U$, there is $(p, g) \in (P \times G)_U$ such that $(\varepsilon(q_1), \varepsilon(q_2), \alpha(q_2)F(a)\alpha(q_1)^{-1}) = (p, pg, \sigma(p, g))$. Now the pair $(\sigma'(q_1, g)a^{-1}, 1)$ gives an arrow from $(\pi(q_2), \varepsilon(q_2), \alpha(q_2))$ to $(\pi(q_1g), \varepsilon(q_1g), \alpha(q_1g))$. In fact,

$$\begin{aligned} \alpha(q_1g)F(\sigma'(q_1, g)a^{-1}) &= \alpha(q_1g)F(\sigma'(q_1, g))F(a)^{-1} \\ &= \sigma(p, g)\alpha(q_1)F(a)^{-1} && (\because (4).) \\ &= \alpha(q_2). && (\because \sigma(p, g) = \alpha(q_2)F(a)\alpha(q_1)^{-1}.) \end{aligned}$$

Since the diagram (2) is 2-cartesian, we can conclude that $q_2 = q_1g$ and $a = \sigma'(q_1, g)$, i.e. $(q_1, q_2, a) = (q_1, q_1g, \sigma'(q_1, g))$. Therefore the morphism Φ is essentially surjective over U . \square

3.2 Automorphisms in $\mathfrak{P}^G(\mathfrak{X})$

Let \mathfrak{X} be a differentiable stack. For an object $(\pi : P \rightarrow \mathfrak{X}, \sigma)$ of $\mathfrak{P}^G(\mathfrak{X})$, we consider the group of automorphisms $\text{Aut}(\pi, \sigma)$ of (π, σ) in $\mathfrak{P}^G(\mathfrak{X})$. Let $(\varphi, \tau) \in \text{Aut}(\pi, \sigma)$. Since τ is a 2-morphism of stacks $\pi \circ \text{id}_P \rightarrow \pi \circ \varphi$, we have a unique smooth map $\gamma_{(\pi, \sigma)} = \gamma : P \rightarrow G$ making the following diagram commutative.

$$\begin{array}{ccccc} P & & \xrightarrow{\varphi} & & P \\ & \searrow^{(\text{id}_P, \gamma)} & & & \downarrow \pi \\ & P \times G & \xrightarrow{\psi} & & P \\ & \downarrow \text{pr}_1 & & \nearrow \sigma & \downarrow \pi \\ P & & \xrightarrow{\pi} & & \mathfrak{X}. \end{array}$$

Therefore $\varphi(p) = p\gamma(p)$ for any $p \in P$. Since φ is G -equivariant, $p\gamma(p)g = pg\gamma(pg)$ for any $p \in P$ and $g \in G$. We also have $\tau(p) = \sigma(p, \gamma(p))$ for $p \in P$.

Now we assume that the stack \mathfrak{X} is type-R (cf. Definition 2.17). Since $[P/G]$ is equivalent to \mathfrak{X} , the G -action of P must be locally free and effective. Therefore $\gamma(pg) = g^{-1}\gamma(p)g$ holds for any $g \in G$. Define $\mathcal{G}(P)$ by

$$\mathcal{G}(P) = \{\gamma \in \mathcal{C}^\infty(P, G) \mid \gamma(pg) = g^{-1}\gamma(p)g \text{ for all } p \in P \text{ and } g \in G\}. \quad (6)$$

The space $\mathcal{G}(P)$ inherits the group structure from G . It is easy to see that the above assignment $(\varphi, \tau) \mapsto \gamma_{(\varphi, \tau)}$ gives an injective group anti-homomorphism from $\text{Aut}(\pi, \sigma)$ to $\mathcal{G}(P)$. Suppose $\gamma \in \mathcal{G}(P)$. Putting $\varphi(p) = p\gamma(p)$ and $\tau(p) = \sigma(p, \gamma(p))$, we can easily check that $(\varphi, \tau) \in \text{Aut}(\pi, \sigma)$. Thus the group anti-homomorphism is surjective.

Proposition 3.9. *If \mathfrak{X} is type-R, then $\text{Aut}(\pi : P \rightarrow \mathfrak{X}, \sigma)$ is anti-isomorphic to the group $\mathcal{G}(P)$.*

3.3 The category of representable morphisms

Let M be a right G -space and \mathfrak{X} a differentiable stack. Making use of the category $\mathfrak{P}^G(\mathfrak{X})$ we construct a category which is equivalent to the category of representable morphisms from \mathfrak{X} to the quotient stack $[M/G]$.

Definition 3.10. We define a category $\mathfrak{R}^G(\mathfrak{X}, M)$ as follows.

- An object of $\mathfrak{R}^G(\mathfrak{X}, M)$ is a triple $(\pi, \sigma, \varepsilon)$ such that (π, σ) is an object of $\mathfrak{P}^G(\mathfrak{X})$ and ε is a G -equivariant map $P \rightarrow M$.
- An arrow $(\pi, \sigma, \varepsilon) \rightarrow (\pi', \sigma', \varepsilon')$ in $\mathfrak{R}^G(\mathfrak{X}, M)$ is an arrow $(\varphi, \tau) : (\pi, \sigma) \rightarrow (\pi', \sigma')$ in $\mathfrak{P}^G(\mathfrak{X})$ satisfying $\varepsilon' \circ \varphi = \varepsilon$.
- The composition of two arrows $(\varphi, \tau) : (\pi, \sigma, \varepsilon) \rightarrow (\pi', \sigma', \varepsilon')$ and $(\varphi', \tau') : (\pi', \sigma') \rightarrow (\pi'', \sigma'')$ is defined by the composition $(\varphi', \tau')(\varphi, \tau)$ in $\mathfrak{P}^G(\mathfrak{X})$.
- The identity arrow of $(\pi, \sigma, \varepsilon)$ is defined by the identity arrow $1_{(\pi, \sigma)}$ in $\mathfrak{P}^G(\mathfrak{X})$.

The category of representable morphisms from \mathfrak{X} to $[M/G]$ is denoted by $\text{Mor}^{\text{Rep}}(\mathfrak{X}, [M/G])$.

Theorem 3.11. *The category $\text{Mor}^{\text{Rep}}(\mathfrak{X}, [M/G])$ is equivalent to the category $\mathfrak{R}^G(\mathfrak{X}, M)$.*

Define a functor $\Psi : \text{Mor}^{\text{Rep}}(\mathfrak{X}, [M/G]) \rightarrow \mathfrak{R}^G(\mathfrak{X}, M)$ as follows. For each representable morphism $F : \mathfrak{X} \rightarrow [M/G]$, we can choose a manifold P^F , a morphism $\pi^F : P^F \rightarrow \mathfrak{X}$ and a smooth map $\varepsilon^F : P^F \rightarrow M$ so that the following square is 2-cartesian:

$$\begin{array}{ccc} P^F & \xrightarrow{\varepsilon^F} & M \\ \pi^F \downarrow & \nearrow \alpha^F & \downarrow \pi^M \\ \mathfrak{X} & \xrightarrow{F} & [M/G]. \end{array}$$

Propositions 3.6 and 3.8 say that there is a 2-morphism of stacks $\sigma^F : \text{pr}_1 \circ \pi^F \rightarrow \psi \circ \pi^F$ such that $(\pi^F : P^F \rightarrow \mathfrak{X}, \sigma^F)$ is an object of $\mathfrak{P}^G(\mathfrak{X})$ and ε^F is G -equivariant i.e. $(\pi^F, \sigma^F, \varepsilon^F)$ is an object

of $\mathfrak{R}^G(\mathfrak{X}, M)$. We define $\Psi(F)$ by $(\pi^F, \sigma^F, \varepsilon^F)$ (cf. Remark 3.2). Note that the 2-morphism of stacks σ^F satisfies

$$\alpha^F(pg)F(\sigma^F(p, g)) = \sigma^M(\varepsilon^F(p), g)\alpha^F(p) \quad (7)$$

for any $(p, g) \in P \times G$.

Let F_1 and F_2 be representable morphisms $\mathfrak{X} \rightarrow [M/G]$ and $\theta : F_1 \rightarrow F_2$ a 2-morphisms of stacks. Since we have a 2-morphism of stacks $\alpha^{F_1} \bullet (\theta^{-1} \circ \text{id}_{\pi^{F_1}}) : F_2 \circ \pi^{F_1} \rightarrow \varepsilon^{F_1} \circ \pi^M$, we obtain a unique smooth map $\varphi^\theta : P^{F_1} \rightarrow P^{F_2}$ satisfying $\varepsilon^{F_2} \circ \varphi = \varepsilon^{F_1}$ and a unique 2-morphism of stacks $\tau^\theta : \pi^{F_1} \rightarrow \pi^{F_2} \circ \varphi$:

$$\begin{array}{ccccc}
 P^{F_1} & & \xrightarrow{\varepsilon^{F_1}} & & M \\
 \downarrow \varphi^\theta & & \searrow & & \downarrow \varepsilon^{F_2} \\
 P^{F_2} & \xrightarrow{\varepsilon^{F_2}} & & & M \\
 \downarrow \tau^\theta & \downarrow \pi^{F_2} & & \nearrow \alpha^{F_2} & \downarrow \pi^M \\
 \mathfrak{X} & \xrightarrow{F_2} & & & [M/G].
 \end{array}$$

Here the 2-morphism of stacks τ^θ satisfies

$$\alpha^{F_2}(\varphi^\theta(p))F_2(\tau^\theta(p)) = \alpha^{F_1}(p)\theta(\pi^{F_1}(p))^{-1} \quad (8)$$

for any $p \in P^{F_1}$.

We see that $(\varphi^\theta, \tau^\theta)$ is an arrow $(\pi^{F_1}, \sigma^{F_1}, \varepsilon^{F_1}) \rightarrow (\pi^{F_2}, \sigma^{F_2}, \varepsilon^{F_2})$ in the category $\mathfrak{R}^G(\mathfrak{X}, M)$. Let $(f_1, \beta_1) = (\varphi^\theta \circ \psi, (\tau^\theta \circ \text{id}_\psi) \bullet \sigma^{F_1})$ and $(f_2, \beta_2) = (\psi \circ (\varphi^\theta \times \text{id}_G), (\sigma^{F_2} \circ \text{id}_{\varphi^\theta \times \text{id}_G}) \bullet (\tau^\theta \circ \text{id}_{\text{pr}_1}))$. Then we have the following 2-commutative diagrams ($i = 1, 2$):

$$\begin{array}{ccccc}
 & & P^{F_1} \times G & & \\
 & \swarrow \pi^{F_1} \circ \text{pr}_1 & \downarrow f_i & \searrow (p, g) \mapsto \varepsilon^{F_1}(p)g & \\
 \mathfrak{X} & \xleftarrow{\pi^{F_2}} & P^{F_2} & \xrightarrow{\varepsilon^{F_2}} & P.
 \end{array}$$

Then for $(p, g) \in P^{F_1} \times G$ we have

$$\begin{aligned}
 \alpha^{F_2}(f_1(p, g))F_2(\beta_1(p, g)) &= \alpha^{F_2}(\varphi^\theta(pg))F_2(\tau^\theta(pg))F_2(\sigma^{F_1}(p, g)) \\
 &= \alpha^{F_1}(pg)\theta(\pi^{F_1}(pg))^{-1}F_2(\sigma^{F_1}(p, g)) & (\because (8).) \\
 &= \alpha^{F_1}(pg)F_1(\sigma^{F_1}(p, g))\theta(\pi^{F_1}(p))^{-1} \\
 &= \sigma^M(\varepsilon^{F_1}(p), g)\alpha^{F_1}(p)\theta(\pi^{F_1}(p))^{-1}. & (\because (7).)
 \end{aligned}$$

On the other hand, we have

$$\begin{aligned}
 \alpha^{F_2}(f_2(p, g))F_2(\beta_2(p, g)) &= \alpha^{F_2}(\varphi^\theta(p)g)F_2(\sigma^{F_2}(\varphi^\theta(p), g))F_2(\tau^\theta(p)) \\
 &= \sigma^M(\varepsilon^{F_2}(\varphi^\theta(p)), g)\alpha^{F_2}(\varphi^\theta(p))F_2(\tau^\theta(p)) & (\because (7).) \\
 &= \sigma^M(\varepsilon^{F_1}(p), g)\alpha^{F_1}(p)\theta(\pi^{F_1}(p))^{-1}. & (\because (8).)
 \end{aligned}$$

Applying Lemma 3.7, we conclude that $f_1 = f_2$ and $\beta_1 = \beta_2$, i.e. for $(p, g) \in P^{F_1} \times G$ we have $\varphi^\theta(pg) = \varphi^\theta(p)g$ and $\tau^\theta(pg)\sigma^{F_1}(p, g) = \sigma^{F_2}(\varphi^\theta(p), g)\tau^\theta(p)$. Similarly we can see that φ^θ is a diffeomorphism. (In fact, $\varphi^{(\theta^{-1})}$ is the inverse of φ^θ .) Therefore $(\varphi^\theta, \tau^\theta)$ is an arrow $(\pi^{F_1}, \sigma^{F_1}, \varepsilon^{F_1}) \rightarrow (\pi^{F_2}, \sigma^{F_2}, \varepsilon^{F_2})$ in the category $\mathfrak{R}^G(\mathfrak{X}, M)$. We define $\Psi(\theta) = (\varphi^\theta, \tau^\theta)$.

Lemma 3.12. *The functor $\Psi : \text{Mor}^{\text{Rep}}(\mathfrak{X}, [M/G]) \rightarrow \mathfrak{R}^G(\mathfrak{X}, M)$ is faithful.*

Proof. Let θ and θ' be two 2-morphisms of stacks from F_1 to F_2 in $\text{Mor}^{\text{Rep}}(\mathfrak{X}, [M/G])$. Suppose that we have an arrow $(\varphi, \tau) : \Psi(\theta) \rightarrow \Psi(\theta')$ in $\mathfrak{R}^G(\mathfrak{X}, M)$. Because of Identity (8), we have

$$\theta(\pi^{F_1}(p)) = F_2(\tau(p))^{-1} \alpha^{F_2}(\varphi(p))^{-1} \alpha^{F_1}(p) = \theta'(\pi^{F_1}(p))$$

for any $p \in P$. Therefore it suffices to show that for any object $x \in \mathfrak{X}$ the arrow $\theta(x)$ can be calculated without θ . Let $x \in \mathfrak{X}_U$ and $\{U_i \rightarrow U\}$ be a covering family of U in $\mathfrak{D}\text{iff}$. Since $[M/G]$ is a stack, $\theta(x)$ can be uniquely determined by its pullbacks $f_i^*\theta(x)$. Therefore it suffices to show that $f_i^*\theta(x)$ can be calculated without θ .

Choosing a manifold P_x equivalent to $U_x \times_{\pi^{F_1}} P^{F_1}$, we have a principal G -bundle $\pi_x : P_x \rightarrow U$, a G -equivariant map $\varepsilon_x : P_x \rightarrow P^{F_1}$ and a 2-morphism of stacks $\beta_x : x \circ \pi_x \rightarrow \pi^{F_1} \circ \varepsilon_x$. We may assume that the pullback of the principal G -bundle $\pi_x : P_x \rightarrow U$ via f_i is trivial:

$$\begin{array}{ccccc} U_i \times G & \longrightarrow & P_x & \xrightarrow{\varepsilon_x} & P^{F_1} \\ \text{pr}_1 \downarrow & \nearrow s_i & \downarrow \pi_x & \searrow \beta_x & \downarrow \pi^{F_1} \\ U_i & \xrightarrow{f_i} & U & \xrightarrow{x} & \mathfrak{X}. \end{array}$$

Here $s_i : U_i \rightarrow P_x$ is a smooth map corresponding to a trivialisation. Using $s_i \in (P_x)_{U_i}$, we obtain the following arrows in $[M/G]_{U_i}$:

$$\begin{array}{ccccc} f_i^* F_1(x) & \longleftarrow & F_1(f_i^* x) & \xrightarrow{F_1(\beta_x(s_i))} & F_1(\pi^{F_1}(p_i)) \\ \downarrow f_i^* \theta(x) & & \downarrow \theta(f_i^* x) & & \downarrow \theta(\pi^{F_1}(p_i)) \\ f_i^* F_2(x) & \longleftarrow & F_2(f_i^* x) & \xrightarrow{F_2(\beta_x(s_i))} & F_2(\pi^{F_1}(p_i)). \end{array}$$

Because of Identity (8), $\theta(\pi^{F_1}(p_i))$ can be calculated without θ , and so is $f_i^*\theta(x)$. \square

Let F be a representable morphism from \mathfrak{X} to $[M/G]$. Making use of the atlas $P^F \times G$ of \mathfrak{X} , we can describe F in an explicit way:

Proposition 3.13. *Let $F \in \text{Mor}^{\text{Rep}}(\mathfrak{X}, [M/G])$. If we identify \mathfrak{X} with $[P^F/G]$, then the morphism of stacks F is isomorphic to the morphism of stacks \hat{F} defined by*

$$\hat{F} : [P^F/G] \rightarrow [M/G]; \quad \begin{cases} (\pi, \varepsilon) \mapsto (\pi, \varepsilon^F \circ \varepsilon) & \text{for objects,} \\ (f, \tilde{f}) \mapsto (f, \tilde{f}) & \text{for arrows.} \end{cases}$$

Remark 3.14. There is an embedding \mathfrak{B} of the bicategory \mathfrak{Bi} of Lie groupoids and bibundles into the 2-category $\mathfrak{St}\mathfrak{Diff}$ of stacks over \mathfrak{Diff} . For a Lie groupoid $X_1 \rightrightarrows X_0$, $\mathfrak{B}(X_1 \rightrightarrows X_0)$ is the category of principal $(X_1 \rightrightarrows X_0)$ -bundles. For a bibundle f_P from $X_1 \rightrightarrows X_0$ to $Y_1 \rightrightarrows Y_0$, the morphism of stacks $\mathfrak{B}(f_P) : \mathfrak{B}(X_1 \rightrightarrows X_0) \rightarrow \mathfrak{B}(Y_1 \rightrightarrows Y_0)$ is defined as follows. Any principal $(X_1 \rightrightarrows X_0)$ -bundle $\alpha : Q \rightarrow M$ naturally defines a bibundle f_Q from $M \rightrightarrows M$ to $X_1 \rightrightarrows X_0$. The composition $f_P \circ f_Q$ includes a principal $(Y_1 \rightrightarrows Y_0)$ -bundle, and it is the image of f_Q via $\mathfrak{B}(f_P)$. The image of the embedding \mathfrak{B} is the 2-category of differentiable stacks (without the condition to be proper). For details of the embedding \mathfrak{B} , see Lerman [12, §4].

Proof. A bibundle corresponding a morphism of stacks can be obtained by taking the following pullback [12]:

$$\begin{array}{ccccc}
P^F \times G & \xrightarrow{\psi} & P^F & \xrightarrow{\varepsilon^F} & M \\
\text{pr}_1 \downarrow & & \downarrow \sigma^F & & \downarrow \pi^M \\
P^F & \xrightarrow{\pi^F} & \mathfrak{X} & \xrightarrow{F} & [M/G].
\end{array}$$

The triple $(\text{pr}_1, P^F, \varepsilon^F \circ \psi)$ is a bibundle from $P^F \times G$ to $M \times G$ corresponding to $F : \mathfrak{X} \rightarrow [M/G]$. It is easy to see that the morphism of stacks $\mathfrak{B}(\text{pr}_1, P^F, \varepsilon^F \circ \psi) : [P^F/G] \rightarrow [M/G]$ is isomorphic to \hat{F} . \square

Lemma 3.15. *The functor $\Psi : \text{Mor}^{\text{Rep}}(\mathfrak{X}, [M/G]) \rightarrow \mathfrak{R}^G(\mathfrak{X}, M)$ is essentially surjective.*

Proof. Let $(\pi : P \rightarrow \mathfrak{X}, \sigma, \varepsilon)$ be an object of $\mathfrak{R}^G(\mathfrak{X}, M)$ and identify \mathfrak{X} with $[P/G]$. Consider the morphism of stacks defined by

$$F : [P/G] \rightarrow [M/G]; \begin{cases} (\pi', \varepsilon') \mapsto (\pi', \varepsilon \circ \varepsilon') & \text{for objects,} \\ (f, \tilde{f}) \mapsto (f, \tilde{f}) & \text{for arrows.} \end{cases}$$

It is easy to see that the following square is 2-cartesian:

$$\begin{array}{ccc}
P & \xrightarrow{\varepsilon} & M \\
\pi^P \downarrow & & \downarrow \pi^M \\
[P/G] & \xrightarrow{F} & [M/G].
\end{array}$$

(The identity transformation gives a 2-morphism of stack in the above square.) This also implies that F is representable. Then we have a unique smooth map $\varphi : P^F \rightarrow P$ and a 2-

morphism of stacks $\tau : \pi^P \rightarrow \varphi \circ \pi$ such that the following diagram is 2-commutative.

$$\begin{array}{ccccc}
 P^F & & & & \\
 \downarrow \varphi & \xrightarrow{\varepsilon^F} & & & \\
 P & \xrightarrow{\varepsilon} & M & & \\
 \downarrow \pi & & \downarrow \pi^P & & \\
 [P/G] & \xrightarrow{F} & [M/G] & &
 \end{array}$$

Modifying the above discussion for $\Psi(\theta)$ slightly, we can show that the pair (φ, τ) gives an arrow from $\Psi(F)$ to $(\pi, \sigma, \varepsilon)$ in $\mathfrak{R}^G(\mathfrak{X}, [M/G])$. \square

Lemma 3.16. *The functor $\Psi : \text{Mor}^{\text{Rep}}(\mathfrak{X}, [M/G]) \rightarrow \mathfrak{R}^G(\mathfrak{X}, M)$ is full.*

Proof. Let $F_1, F_2 \in \text{Mor}^{\text{Rep}}(\mathfrak{X}, [M/G])$. Suppose we have an arrow (φ, τ) from $\Psi(F_1)$ to $\Psi(F_2)$. Since φ is a G -equivariant diffeomorphism, it is obvious that the following morphism E is an equivalence:

$$E : [P^{F_1}/G] \rightarrow [P^{F_2}/G]; \begin{cases} (\pi, \varepsilon) \mapsto (\pi, \varphi \circ \varepsilon) & \text{for objects,} \\ (f, \tilde{f}) \mapsto (f, \tilde{f}) & \text{for arrows.} \end{cases}$$

We identify $[P^{F_2}/G]$ with $[P^{F_1}/G]$ via the above morphism. Then we can also identify F_1 and F_2 because of Proposition 3.13 and $\varepsilon^{F_2} \circ \varphi = \varepsilon^{F_1}$. Therefore the functor Ψ is full. \square

4 Symplectic vortex equation

In this section, we define symplectic vortex equations over a compact Riemann surface Σ , regarding Σ as a type-R stack \mathfrak{X} . First we fix notations for an orbifold Riemann surface. Secondly we see the definition of (representable) pseudo-holomorphic curve from \mathfrak{X} to a Marsden–Weinstein quotient (also known as a symplectic quotient). After that we define symplectic vortex equations and an energy functional. Finally we discuss the moduli space of solutions to the equations for a linear action of the circle group S^1 on the complex plane \mathbb{C} .

In this section, G is a compact and connected Lie group. The Lie algebra of G is denoted by \mathfrak{g} . We fix an invariant inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{g} to identify \mathfrak{g} with its dual \mathfrak{g}^\vee .

A Hamiltonian G -space is a symplectic manifold (M, ω) equipped with a (right) G -action and a G -equivariant map $\mu : M \rightarrow \mathfrak{g}$ satisfying

$$\mu(mg) = \text{Ad}(g^{-1})\mu(m) \quad \text{for any } m \in M \text{ and } g \in G \quad (9)$$

and

$$d\langle \mu, \xi \rangle = -\iota(\xi^M)\omega \quad \text{for any } \xi \in \mathfrak{g}. \quad (10)$$

Here Ad is the adjoint representation of G . Note that the G -action preserves ω since G is connected. This map μ is called a moment map for the G -action.

4.1 An orbifold Riemann surface, revisited

Let $\Sigma = (\Sigma, j, \mathbf{z}, \mathbf{m})$ be an orbifold Riemann surface and \mathfrak{X} the type-R stack corresponding to Σ (§2.4) and $(\pi : P \rightarrow \mathfrak{X}, \sigma)$ an object of $\mathfrak{P}^G(\mathfrak{X})$. Since \mathfrak{X} is type-R, the right G -action on P is locally free and effective (cf. §3.2). In particular the space of connections

$$\mathcal{A}(P) = \{A \in \Omega^1(P, \mathfrak{g}) \mid \iota(\xi^P)A = \xi \ (\forall \xi \in \mathfrak{g}) \text{ and } \psi_g^*A = \text{Ad}(g^{-1})A \ (\forall g \in G)\}$$

is nonempty.

The sheaf over P induced by the tangent sheaf $\mathcal{T}_{\mathfrak{X}}$ is the sheaf of sections of the bundle TP/\mathfrak{g}^P . Note that the bundle TP/\mathfrak{g}^P has a complex structure coming from the complex structure j on \mathfrak{X} , and we also denote by j the complex structure on TP/\mathfrak{g}^P . Since j is compatible with the source and target maps, the identity $(j[v])g = j[vg]$ holds for any $v \in TP$ and $g \in G$.

We will need local slices for integrations over \mathfrak{X} and §4.4. Let $p \in P$ and (U, φ, Γ) a triple of

- (i) a finite subgroup $\Gamma \subset G$, which acts on \mathbb{C} in an orthogonal linear way,
- (ii) a Γ -invariant open neighbourhood U of 0 in \mathbb{C} , and
- (iii) a smooth map $\varphi : U \rightarrow P$ satisfying $\varphi(0) = p$ and $\varphi(gu) = \varphi(u)g^{-1}$.

The triple is called a (complex) slice at $p \in P$ if the map

$$U \times_{\Gamma} G \rightarrow P; [u, a] \mapsto \varphi(u)a$$

is a G -equivariant diffeomorphism onto a G -invariant open neighbourhood of p . Here Γ acts on $U \times G$ by $(u, a) \cdot g = (g^{-1}u, g^{-1}a)$ ($u \in U, a \in G$ and $g \in \Gamma$).

We may choose a family $\{(U_i, \varphi_i, \Gamma_i)\}_i$ of slices so that $\{\varphi_i(U_i)G\}$ is an open cover of P . Let Z_0 be the disjoint union $\bigsqcup_i U_i$ and $\iota : Z_0 \rightarrow P$ is the map whose restriction to U_i is the inclusion map $\varphi_i : U_i \rightarrow P$. Let Z_1 be a fibred product of the diagram:

$$\begin{array}{ccccc} Z_1 & \xrightarrow{\iota_1} & P \times S^1 & \xrightarrow{\text{Pr}_2} & S^1 \\ (\text{tgt,src}) \downarrow & & \downarrow (\text{tgt,src}) & & \\ Z_0 \times Z_0 & \xrightarrow{\iota \times \iota} & P \times P & & \end{array}$$

Then we obtain a proper étale groupoid $Z_1 \rightrightarrows Z_0$ which is equivalent to $P \rtimes G$ in the bicategory \mathfrak{Bi} (cf. Lerman [12, Definition 2.25]). It is easy to see that the composition $\pi \circ \iota : Z_0 \rightarrow \mathfrak{X}$ is an atlas of \mathfrak{X} , and the groupoid $Z_1 \rightrightarrows Z_0$ is a presentation associated to the atlas. Note that the groupoid $Z_1 \rightrightarrows Z_0$ has a partition of unity [1].

The space of 2-forms on \mathfrak{X} is given by the space of basic 2-forms on P

$$\Omega^2([P/G]) = \{\eta \in \Omega^2(P) \mid \iota(\xi^P)\eta = 0 \ (\forall \xi \in \mathfrak{g}) \text{ and } \psi_g^*\eta = \eta \ (\forall g \in G)\}.$$

In terms of $Z_1 \rightrightarrows Z_0$, the space of 2-forms on \mathfrak{X} is the space

$$\Omega^2(Z_1 \rightrightarrows Z_0) = \{\theta \in \Omega^2(Z_0) \mid \text{src}^*\theta = \text{tgt}^*\theta\}$$

and an explicit correspondence between $\Omega^2([P/G])$ and $\Omega^2(Z_1 \rightrightarrows Z_0)$ is given by $\iota^* \eta = \theta$ for $\eta \in \Omega^2([P/G])$ and $\theta \in \Omega^2(Z_1 \rightrightarrows Z_0)$.

We denote by $\rho : Z_1 \rightarrow S^1$ the composition of $\text{pr}_2 \circ \iota_1$. The map ρ satisfies $\iota(\text{tgt } \alpha)\rho(\alpha) = \iota(\text{src } \alpha)$ for any $\alpha \in Z_1$. The map

$$\varepsilon : Z_0 \times S^1 \rightarrow P; (z, t) \mapsto \iota(z)t$$

is an atlas of P , and a presentation associated to the atlas $\varepsilon : Z_0 \times S^1 \rightarrow P$ is given by

$$Z_1 \times S^1 \rightrightarrows Z_0 \times S^1; \text{src}(\alpha, t) = (\text{src } \alpha, t), \text{tgt}(\alpha, t) = (\text{tgt } \alpha, \rho(\alpha)t).$$

Note that the map $\varepsilon : Z_0 \times S^1 \rightarrow P$ is S^1 -equivariant with respect to the right S^1 -action on $Z_0 \times S^1$ defined by $(z, t) \cdot t' = (z, tt')$.

4.2 Pseudo-holomorphic curves in a Marsden–Weinstein quotient

For details of (smooth) Marsden–Weinstein quotients, see McDuff–Salamon [14]. Cieliebak–Gao–Salamon [5] is also helpful for our setting.

Let (M, ω) be a Hamiltonian G -space with a moment map $\mu : M \rightarrow \mathfrak{g}$. Assume that $0 \in \mathfrak{g}$ is a regular value of μ . Then $\mu^{-1}(0)$ is a G -invariant submanifold of M and the G -action on $\mu^{-1}(0)$ is locally free. Therefore the quotient stack $M//G = [\mu^{-1}(0)/G]$ is Deligne–Mumford.

The quotient stack $M//G$ inherits a symplectic form from (M, ω) as follows. Let ω_μ be the restriction of ω to $\mu^{-1}(0)$. A direct calculation shows that the 2-form ω_μ is basic: $\omega_\mu \in \Omega^2(M//G)$. Since the kernel of the 2-form ω_μ at $m \in \mu^{-1}(0)$ coincides with $\mathfrak{g}^{\mu^{-1}(0)}(m)$, ω_μ is nondegenerate and therefore ω_μ gives a symplectic form on $M//G$. The symplectic Deligne–Mumford stack $M//G$ is called the Marsden–Weinstein quotient.

Choose an almost complex structure J compatible with the G -action and ω i.e. J is an almost complex structure on M such that $\psi_g \circ J = J \circ \psi_g$ for any $g \in G$ and the bilinear form

$$g(v_1, v_2) = \omega(v_1, Jv_2) \tag{11}$$

defines a Riemannian metric on M . Since the G -action on $\mu^{-1}(0)$ is locally free, $\mathfrak{g}^{\mu^{-1}(0)}$ is a trivial subbundle of $T\mu^{-1}(0)$. The sheaf induced by the tangent sheaf $\mathcal{T}_{M//G}$ is the sheaf of sections of the quotient bundle $T\mu^{-1}(0)/\mathfrak{g}^{\mu^{-1}(0)}$. The bundle $T\mu^{-1}(0)/\mathfrak{g}^{\mu^{-1}(0)}$ inherits a complex structure from TM and it gives rise to an almost complex structure on $\mathcal{T}_{M//G}$.

We consider a representable morphism from a compact orbifold Riemann surface $(\Sigma, j, \mathbf{z}, \mathbf{m})$ to the Marsden–Weinstein quotient $M//G$. We regard the compact orbifold Riemann surface as a type-R stack \mathfrak{X} as Section 2.4.

By Theorem 3.11, a representable morphism F from \mathfrak{X} to $M//G$ corresponds to an object of the category $\mathfrak{R}^G(\mathfrak{X}, \mu^{-1}(0))$, i.e. a pair of an object $(\pi : P \rightarrow \mathfrak{X}, \sigma)$ of $\mathfrak{P}^G(\mathfrak{X})$ and a G -equivariant

map $u : P \rightarrow \mu^{-1}(0)$:

$$\begin{array}{ccc}
 P & \xrightarrow{u} & \mu^{-1}(0) \\
 \pi \downarrow & \nearrow \alpha & \downarrow \pi^M \\
 \mathfrak{X} & \xrightarrow{F} & M//G.
 \end{array}$$

We will stick with the presentation $P \rtimes G$ to describe geometric concepts on \mathfrak{X} (cf. §4.1).

Definition 4.1. The representable morphism $F : \mathfrak{X} \rightarrow M//G$ is said to be **pseudo-holomorphic** if the map $du : TP/\mathfrak{g}^P \rightarrow T\mu^{-1}(0)/\mathfrak{g}^M$ is compatible with the complex structures:

$$J \circ du = du \circ j \text{ as a map } TP/\mathfrak{g}^P \rightarrow T\mu^{-1}(0)/\mathfrak{g}^{\mu^{-1}(0)}. \quad (12)$$

Remark 4.2. This definition is independent of the choice of the object (π, σ, u) of $\mathfrak{R}^G(\mathfrak{X}, \mu^{-1}(0))$ up to isomorphism.

The above condition (12) can be described more explicitly by using a connection. For a connection $A \in \mathcal{A}(P)$ the covariant derivative $d_A u \in \Omega^1(P, u^*TM)$ of u is defined by

$$d_A u : TP \rightarrow u^*TM; d_A u(v) = du(v) - A(v)^M(u(p)) \text{ for } v \in T_p P.$$

We denote by $\bar{d}_A u$ the complex antiholomorphic part of $d_A u$:

$$\bar{d}_A u = \frac{1}{2} (d_A u + J \circ d_A u \circ j) \in \Omega^1(P, u^*TM).$$

Here $d_A u \circ j : TP \rightarrow TM$ is defined as follows. For $v \in T_p P$ there is a tangent vector $\tilde{v} \in T_p P$ such that $[\tilde{v}] = j[v]$. Define $(d_A u \circ j)(v) = d_A u(\tilde{v})$. This definition is independent of the choice of \tilde{v} because the identity $d_A u(\xi^P(p)) = 0$ holds for any $\xi \in \mathfrak{g}$.

Proposition 4.3. A representable morphism $F : \mathfrak{X} \rightarrow M//G$ is pseudo-holomorphic if and only if there is a connection $A \in \mathcal{A}(P)$ satisfying that $\bar{d}_A u = 0$.

Proof. First we note that we have the orthogonal decomposition

$$T_{u(p)}M = N_{u(p)} \oplus \mathfrak{g}^M(u(p)) \oplus J\mathfrak{g}^M(u(p)). \quad (13)$$

Here $N_{u(p)}$ is the orthogonal complement of $\mathfrak{g}^M(u(p))$ in $T_{u(p)}\mu^{-1}(0)$. We can canonically identify the bundle $T\mu^{-1}(0)/\mathfrak{g}^{\mu^{-1}(0)}$ with $N \rightarrow \mu^{-1}(0)$. We denote by ν_1 and ν_2 the projections from $N_{u(p)} \oplus \mathfrak{g}^M(u(p))$ to the first and second component respectively. The condition (12) is equivalent to the identity $J\nu_1(du(p)v) = \nu_1(du(p)jv)$ for every $p \in P$ and $v \in T_p P$.

Suppose a representable morphism $F : \mathfrak{X} \rightarrow M//G$ to be pseudo-holomorphic. Define a 1-form $A \in \Omega^1(P, \mathfrak{g})$ by the composition $T_p P \rightarrow T_{u(p)}\mu^{-1}(0) \rightarrow \mathfrak{g}^{\mu^{-1}(0)}(u(p)) \cong \mathfrak{g}$. It is easy to see that $A \in \mathcal{A}(P)$ and $\bar{d}_A u = 0$.

Conversely suppose that we have a connection $A \in \mathcal{A}(P)$ satisfying $\bar{\partial}_A u = 0$ i.e.

$$J(d_A u)v = (d_A u)jv$$

for any $v \in TP$. Applying the orthogonal decomposition (13) to the above identity, we can easily see that the identity $Jv_1(du(p)v) = v_1(du(p)jv)$ holds for every $p \in P$ and $v \in T_p P$. \square

4.3 Symplectic vortex equations

The symplectic vortex equations are defined for the following data:

- A Hamiltonian G -space (M, ω) with a moment map $\mu : M \rightarrow \mathfrak{g}$.
- An almost complex structure J of M compatible with the G -action.
- A compact orbifold Riemann surface $(\Sigma, j, \mathbf{z}, \mathbf{m})$. We regard it as a stack \mathfrak{X} .
- A volume form $d\text{vol}_{\mathfrak{X}}$ of \mathfrak{X} .
- An object $(\pi^P : P \rightarrow \mathfrak{X}, \sigma)$ of the category $\mathfrak{P}^G(\mathfrak{X})$.

A volume form $d\text{vol}_{\mathfrak{X}}$ of \mathfrak{X} is a nowhere-vanishing 2-form on \mathfrak{X} . Using a G -invariant Riemannian metric on P , we can easily see that a volume form of \mathfrak{X} always exists. Moreover for any 2-form α on \mathfrak{X} there is a unique function $f \in \mathcal{O}(\mathfrak{X})$ such that $\alpha = f d\text{vol}_{\mathfrak{X}}$.

Since \mathfrak{X} is 2-dimensional, a choice of volume form $d\text{vol}_{\mathfrak{X}}$ gives an orientation on \mathfrak{X} . On the other hand, the complex structure j on \mathfrak{X} also gives an orientation on \mathfrak{X} . Thus we choose a volume form $d\text{vol}_{\mathfrak{X}}$ so that both orientations are the same. More explicitly, we assume that the following inequality holds:

$$d\text{vol}_{\mathfrak{X}}(v, \tilde{v}) > 0 \text{ for any nonzero } [v] \in T_p P / \mathfrak{g}^P. \quad (14)$$

Here \tilde{v} is a tangent vector at p satisfying $[\tilde{v}] = j[v]$, and $[\]$ is the projection map from TP to the quotient bundle TP/\mathfrak{g}^P . The left hand side of the inequality is independent of the choice of \tilde{v} since $d\text{vol}_{\mathfrak{X}}$ is basic.

Fix a volume form $d\text{vol}_{\mathfrak{X}}$ of \mathfrak{X} . We define the Hodge $*$ -operator by

$$* : \Omega^2(\mathfrak{X}) \rightarrow \Omega^0(\mathfrak{X}); * \alpha = *(f d\text{vol}_{\mathfrak{X}}) = f.$$

The Hodge $*$ -operator can naturally extend to a map $\Omega^2(\mathfrak{X}, \mathfrak{g}) \rightarrow \Omega^0(\mathfrak{X}, \mathfrak{g})$.

Definition 4.4. The **symplectic vortex equations** are the partial differential equations

$$\bar{\partial}_A u = 0, \quad *F_A + \mu(u) = 0 \quad (15)$$

for $(A, u) \in \mathcal{A}(P) \times \mathcal{C}_G^\infty(P, M)$. Here $\mathcal{C}_G^\infty(P, M)$ is the space of G -equivariant smooth maps from P to M . Note that the curvature F_A belongs to $\Omega^2(\mathfrak{X}, \mathfrak{g})$.

Remark 4.5. Assume that $0 \in \mathfrak{g}$ is a regular value of the moment map μ . If we replace $d\text{vol}_{\mathfrak{X}}$ with $\varepsilon^{-2} d\text{vol}_{\mathfrak{X}}$ for $\varepsilon > 0$, then the equations (15) become

$$\bar{\partial}_A u = 0, \quad *F_A + \varepsilon^{-2} \mu(u) = 0.$$

The limit of the above equations of ε , as ε approaches to 0, are nothing but the equations for pseudo-holomorphic morphisms from \mathfrak{X} to $M//G$ (cf. Proposition 4.3):

$$\bar{\partial}_A u = 0, \quad \mu(u) = 0.$$

We define the the **energy functional** (or the action functional) $E : \mathcal{A}(P) \times C_G^\infty(P, M) \rightarrow \mathbb{R}$ by

$$E(A, u) = \frac{1}{2} \int_{\mathfrak{X}} \left(|d_A u|^2 + |F_A|^2 + |\mu(u)|^2 \right) d\text{vol}_{\mathfrak{X}}. \quad (16)$$

Here we use the G -invariant inner product on \mathfrak{g} and the Riemannian metric on M defined as (11). In the above formula, $|d_A u|^2$ can be calculated as follows. First we note that the complex structure j on TP/\mathfrak{g}^P and the volume form $d\text{vol}_{\mathfrak{X}}$ define a metric on TP/\mathfrak{g}^P :

$$g_{\mathfrak{X}}([v], [w]) = d\text{vol}_{\mathfrak{X}}(v, \tilde{w}) \text{ for } v, w \in T_p P,$$

where \tilde{w} is a tangent vector at p satisfying $[\tilde{w}] = j[w]$, and the above definition is well-defined because $d\text{vol}_{\mathfrak{X}}$ is basic. Since TP/\mathfrak{g}^P is a vector bundle of rank 2, it is easy to see that $g_{\mathfrak{X}}$ form a metric on TP/\mathfrak{g}^P . Then define

$$|d_A u| = \|[v]\|^{-1} \sqrt{|d_A u(v)|^2 + |(d_A u \circ j)(v)|^2} \quad (17)$$

for nonzero $[v] \in TP/\mathfrak{g}^P$, where $\|[v]\| = \sqrt{g_{\mathfrak{X}}([v], [v])}$. The right hand side is independent of the choice of $[v]$.

Proposition 4.6. *For every $(A, u) \in \mathcal{A}(P) \times C_G^\infty(P, M)$, the identity*

$$E(A, u) = \int_{\mathfrak{X}} \left(|\bar{\partial}_A u|^2 + \frac{1}{2} |*F_A + \mu(u)|^2 \right) d\text{vol}_{\mathfrak{X}} + R(\omega, \mu, u) \quad (18)$$

holds. Here

$$R(\omega, \mu, u) = \int_{\mathfrak{X}} \left((d_A u)^* \omega - \langle \mu(u), F_A \rangle \right)$$

and $|\bar{\partial}_A u|$ is calculated in a similar way to $|d_A u|$.

Remark 4.7. We can see as follows that the value $R(\omega, \mu, u)$ is independent of the choice of $A \in \mathcal{A}(P)$. First we consider the Cartan model $\Omega_G^2(P)$ for the G -space P . Then $u^*(\omega - \mu)$ is a G -equivariant 2-form after we identify \mathfrak{g} with \mathfrak{g}^\vee . We can assign the Cartan operator

$$\Omega_G^2(P) \rightarrow \Omega^2([P/G]); \quad \eta \rightarrow \eta_A$$

to each connection $A \in \mathcal{A}(P)$. In terms of the Cartan operator,

$$R(\omega, \mu, u) = \int_{\mathfrak{X}} u^*(\omega - \mu)_A,$$

and this integration is independent of the choice of the connection A [6, Proposition 4.2].

Proof. We follow the notation in §4.1. Let $\chi \in C^\infty(Z_0)$ be a partition of unity. Then

$$E(u, A) = \frac{1}{2} \sum_i \int_{U_i} \chi_i \varphi_i^* \left(|d_A u|^2 + |F_A|^2 + |\mu(u)|^2 \right) \text{dvol}_{\mathfrak{X}},$$

where $\chi_i = \chi|_{U_i}$. Let $s + it$ be the standard complex coordinate on \mathbb{C} . Then $[d\varphi_i(\partial_t)] = j[d\varphi_i(\partial_s)]$ and $\varphi_i^* A = \Phi_i ds + \Psi_i dt$ for some $\Phi_i, \Psi_i \in C^\infty(U_i, \mathfrak{g})$. Put $u_i = u \circ \varphi_i$. Then the pullback of $d_A u, F_A$ and $\text{dvol}_{\mathfrak{X}}$ to U_i via φ_i are given by

$$\begin{aligned} \varphi_i^* d_A u &= \left(\frac{\partial u_i}{\partial s} + \Phi_i^M \right) ds + \left(\frac{\partial u_i}{\partial t} + \Psi_i^M \right) dt, \\ \varphi_i^* F_A &= \left(\frac{\partial \Psi_i}{\partial s} - \frac{\partial \Phi_i}{\partial t} + [\Phi_i, \Psi_i] \right) ds \wedge dt, \\ \varphi_i^* \text{dvol}_{\mathfrak{X}} &= \lambda_i^2 ds \wedge dt \end{aligned}$$

for some positive function λ_i on U_i . The rest of the proof is similar to Proposition 3.1 in Cieliebak–Gaio–Salamon [5]. \square

Corollary 4.8. *A pair $(A, u) \in \mathcal{A}(P) \times C_G^\infty(P, M)$ is a solution of the symplectic vortex equation (15) if and only if $E(A, u) = R(\omega, \mu, u)$.*

Since \mathfrak{X} is type-R, the group of automorphisms of $(\pi^P : P \rightarrow \mathfrak{X}, \sigma)$ in $\mathfrak{B}^G(\mathfrak{X})$ is anti-isomorphic to the group $\mathcal{G}(P)$ defined as (6). The group $\mathcal{G}(P)$ acts on the space $\mathcal{A}(P) \times C_G^\infty(P, M)$ by

$$\mathcal{A}(P) \times C_G^\infty(P, M) \curvearrowright \mathcal{G}(P); \quad \gamma^*(A, u) = (\gamma^{-1} d\gamma + \gamma^{-1} A \gamma, u \gamma).$$

Lemma 4.9. *The energy functional E is $\mathcal{G}(P)$ -invariant.*

This lemma can be shown by direct calculation and implies the following proposition.

Proposition 4.10. *The action of $\mathcal{G}(P)$ on $\mathcal{A}(P) \times C_G^\infty(P, M)$ preserves the space of solutions to the symplectic vortex equations (15).*

4.4 The case of $G = S^1$ and $M = \mathbb{C}$

In this subsection, we restrict ourselves to the cases when $G = S^1$ and $M = \mathbb{C}$, and we consider the moduli space of solutions to the symplectic vortex equations by comparing it with the moduli space of Kähler vortices [21].

First we fix notation for the circle group $G = S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$. The Lie algebra $\mathfrak{g} = \mathfrak{i}\mathbb{R}$ is equipped with an S^1 -invariant inner product: $\langle \xi_1, \xi_2 \rangle = -\xi_1 \xi_2$ for $\xi_1, \xi_2 \in \mathfrak{i}\mathbb{R}$. Fix a positive integer a we define a left S^1 -action on \mathbb{C} by

$$S^1 \curvearrowright \mathbb{C}; \quad t \cdot z = t^{-a} z.$$

(Therefore the right action is defined as $z \cdot t = t^{-1} \cdot z = t^a z$.) A moment map is given by

$$\mu : \mathbb{C} \mapsto i\mathbb{R}; z \mapsto \frac{i}{2}(a|z|^2 - \tau),$$

where τ is a positive real number. Then the Marsden–Weinstein quotient $\mathbb{C} // S^1$ is called a weighted projective space $\mathbb{P}(a)$ (cf. Sakai [23, Section 3.3]).

Let \mathfrak{X} be a compact orbifold Riemann surface $(\Sigma, j, \mathbf{z}, \mathbf{m})$. Choose an object $(\pi : P \rightarrow \mathfrak{X}, \sigma)$ of $\mathfrak{P}^G(\mathfrak{X})$. Then P is a 3-dimensional closed manifold equipped with a locally free S^1 -action, i.e. a Seifert fibred space [21, 24].

The symplectic vortex equations (15) are given by

$$\bar{\partial}_A u = 0, \quad *F_A + \frac{i}{2}(a|u|^2 - \tau) = 0, \quad (19)$$

where $A \in \mathcal{A}(P) \subset \Omega^1(P, i\mathbb{R})$ is a connection on P , and u belongs to

$$\mathcal{C}_{S^1}^\infty(P, \mathbb{C}) = \{u \in \mathcal{C}^\infty(P, \mathbb{C}) \mid u(pt) = t^a u(p) \text{ for all } p \in P \text{ and } t \in S^1\}.$$

The gauge group $\mathcal{G}(P) = \{\gamma \in \mathcal{C}^\infty(P, S^1) \mid \gamma(pt) = \gamma(p) \text{ for all } p \in P \text{ and } t \in S^1\}$ acts on the space of solutions to the equations (19) by $\gamma^*(A, u) = (\gamma^{-1}d\gamma + \gamma^{-1}A\gamma, \gamma^a u)$. We denote by $\mathcal{M}(P)$ the moduli space of the solutions to (19):

$$\mathcal{M}(P) = \{(A, u) \in \mathcal{A}(P) \times \mathcal{C}_{S^1}^\infty(P, \mathbb{C}) \mid (A, u) \text{ satisfies (19)}\} / \mathcal{G}(P).$$

To consider the moduli space of solutions of the symplectic vortex equations (19), we assume the following condition for \mathfrak{X} .

Assumption 4.11. Let $\mathbf{m} = (m_1, \dots, m_k)$. Then a is a common multiple of m_1, \dots, m_k .

The reason for the assumption is as follows. As Remark 4.5, a certain limit of the symplectic vortex equations is the equation for pseudo-holomorphic curve in the Marsden–Weinstein quotient $\mathbb{C} // S^1$:

$$\bar{\partial}_A u = 0, \quad |u|^2 = \frac{\tau}{a}.$$

It is natural to assume that the above equations have a solution. Let (A, u) be a solution. For $p \in P$ and a stabiliser $t \in S^1$ at p , the map $u : P \rightarrow \mathbb{C}$ satisfies $u(p) = t^a u(p)$. By the second equation we have $u(p) \neq 0$, therefore $t^a = 1$. This implies the assumption.

Remark 4.12. If $a = 1$, then the orbifold Riemann surface \mathfrak{X} must be non-singular.

To compare $\mathcal{M}(P)$ with the moduli space of Kähler vortices, we describe the moduli space $\mathcal{M}(P)$ by using slices of the S^1 -space P (cf. §4.1).

Taking pullbacks via the atlas ε of P , we can identify $\mathcal{C}^\infty(P, \mathbb{C})$ with the space

$$\mathcal{C}^\infty(Z_1 \times S^1 \rightrightarrows Z_0 \times S^1, \mathbb{C}) = \{\tilde{u} \in \mathcal{C}^\infty(Z_0 \times S^1, \mathbb{C}) \mid \text{src}^* \tilde{u} = \text{tgt}^* \tilde{u}\}.$$

By this fact, there is a one-to-one correspondence between $\mathcal{C}_{S^1}^\infty(P, \mathbb{C})$ and the space

$$\mathcal{C}^\infty(Z_0, \mathbb{C})_\rho = \{f \in \mathcal{C}^\infty(Z_0, \mathbb{C}) \mid \text{src}^* f = \rho^a \text{tgt}^* f\}.$$

The correspondence is explicitly given by $f = u \circ \iota$.

Moreover the pullback $\varepsilon^* : \Omega^1(P, \mathbb{i}\mathbb{R}) \rightarrow \Omega^1(Z_0 \times S^1, \mathbb{i}\mathbb{R})$ gives a linear isomorphism from the space $\mathcal{A}(P)$ of connections to the space $\mathcal{A}(Z_1 \times S^1 \rightrightarrows Z_0 \times S^1)$ defined by

$$\{B \in \Omega^1(Z_0 \times S^1, \mathbb{i}\mathbb{R}) \mid \text{src}^* B = \text{tgt}^* B, B(\zeta^{Z_0 \times S^1}) = \zeta \ (\forall \zeta \in \mathbb{i}\mathbb{R}) \text{ and } \psi_t^* B = B \ (\forall t \in S^1)\}.$$

Let B_{MC} be the Maurer–Cartan form for the product bundle $Z_0 \times S^1 \rightarrow Z_0$. Define

$$\Omega^1(Z_0, \mathbb{i}\mathbb{R})_\rho = \{\theta \in \Omega^1(Z_0, \mathbb{i}\mathbb{R}) \mid \text{src}^* \theta = \rho^{-1} d\rho + \text{tgt}^* \theta\}.$$

Then the pullback $\text{pr}_1^* : \Omega^1(Z_0, \mathbb{i}\mathbb{R}) \rightarrow \Omega^1(Z_0 \times S^1, \mathbb{i}\mathbb{R})$ gives rise to the linear isomorphism

$$\Omega^1(Z_0, \mathbb{i}\mathbb{R})_\rho \rightarrow \mathcal{A}(Z_1 \times S^1 \rightrightarrows Z_0 \times S^1); \theta \mapsto B_{\text{MC}} + \text{pr}_1^* \theta.$$

Therefore we obtain a one-to-one correspondence between $\mathcal{A}(P)$ and $\Omega^1(Z_0, \mathbb{i}\mathbb{R})_\rho$. The explicit correspondence is given by $\varepsilon^* A = B_{\text{MC}} + \text{pr}_1^* \theta$ (or $\iota^* A = \theta$) for $A \in \mathcal{A}(P)$ and $\theta \in \Omega^1(Z_0, \mathbb{i}\mathbb{R})_\rho$.

The gauge group $\mathcal{G}(P)$ can be identified with

$$\mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1) = \{g \in \mathcal{C}^\infty(Z_0, S^1) \mid \text{src}^* g = \text{tgt}^* g\}$$

via the identification $\gamma \circ \iota = g$ for $\gamma \in \mathcal{G}(P)$ and $g \in \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1)$ and the action of the gauge group $\mathcal{G}(P)$ on $\mathcal{A}(P) \times \mathcal{C}_{S^1}^\infty(P, \mathbb{C})$ can be identified with the following action:

$$\Omega^1(Z_0, \mathbb{i}\mathbb{R})_\rho \times \mathcal{C}^\infty(Z_0, \mathbb{C})_\rho \curvearrowright \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1); (\theta, f) \cdot g = (g^{-1} d g + \theta, g^a f).$$

Under the above identification, the equations (19) can be identified the following PDEs by taking pullbacks via $\iota : Z_0 \rightarrow P$:

$$\bar{\partial}_{-a\theta} f = 0, \quad *F_\theta + \frac{i}{2}(a|f|^2 - \tau) = 0 \tag{20}$$

for $(\theta, f) \in \Omega^1(Z_0, \mathbb{i}\mathbb{R})_\rho \times \mathcal{C}^\infty(Z_0, \mathbb{C})_\rho$. Here, for the $*$ -operator on Z_0 , the volume form on Z_0 is the pullback $\iota^* \text{dvol}_x$. Therefore $\mathcal{M}(P)$ is identified with

$$\mathcal{M}(Z_1 \rightrightarrows Z_0) = \{(\theta, f) \in \Omega^1(Z_0, \mathbb{i}\mathbb{R})_\rho \times \mathcal{C}^\infty(Z_0, \mathbb{C})_\rho \mid (\theta, f) \text{ satisfies (20)}\} / \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1).$$

Next we describe the moduli space of Kähler vortices in terms of $Z_1 \rightrightarrows Z_0$. The S^1 -action on $P \times \mathbb{C}$ by $(p, x) \cdot t = (pt, t^a x)$ is proper and locally free, and therefore the action induces an orbifold structure on the quotient space $P \times_{S^1} \mathbb{C}$. Moreover the quotient map $P \times_{S^1} \mathbb{C} \rightarrow P/S^1 = \Sigma$ gives rise to a Hermitian orbifold line bundle. The Hermitian orbifold line bundle can be described as the following proper étale groupoid

$$Z_1 \times \mathbb{C} \rightrightarrows Z_0 \times \mathbb{C}; \begin{cases} \text{src}(\alpha, x) = (\text{src } \alpha, x), \\ \text{tgt}(\alpha, x) = (\text{tgt } \alpha, \rho(\alpha)^a x). \end{cases}$$

The space of orbisections is naturally identified with $\mathcal{C}^\infty(Z_0, \mathbb{C})_\rho$, and the space of the compatible orbifold connections is given by

$$\Omega^1(Z_0, \mathbb{R})_a = \{B \in \Omega^1(Z_0, \mathbb{R}) \mid \text{src}^* B = -a\rho^{-1}d\rho + \text{tgt}^* B\}.$$

The gauge group is nothing but $\mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1)$ and the (right) action of the gauge group on $\Omega^1(Z_0, \mathbb{R})_a \times \mathcal{C}^\infty(Z_0, \mathbb{C})_\rho$ is defined by

$$\Omega^1(Z_0, \mathbb{R})_a \times \mathcal{C}^\infty(Z_0, \mathbb{C})_\rho \curvearrowright \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1); (B, f) \cdot g = (g^{-1}dg + B, g^{-1}f).$$

The equations of Kähler vortices over the orbifold Riemann surface $(\Sigma, j, \mathbf{z}, \mathbf{m})$ are the PDEs

$$\bar{\partial}_B f = 0, \quad *F_B + \frac{i}{2}(|f|^2 - a\tau) = 0 \quad (21)$$

for $(B, f) \in \Omega^1(Z_0, \mathbb{R})_a \times \mathcal{C}^\infty(Z_0, \mathbb{C})_\rho$.

Remark 4.13. The above equations are obtained after we replace F_{K_Σ} with $-ai\tau \text{dvol}_\Sigma$ in the equations of Kähler vortices [21, Section 5.5]. This replacement does not produce any problems on the following discussion (Proposition 4.16) in this paper.

Therefore the moduli space of Kähler vortices is defined by

$$\mathcal{M}^K = \{(B, f) \in \Omega^1(Z_0, \mathbb{R})_a \times \mathcal{C}^\infty(Z_0, \mathbb{C})_\rho \mid (B, f) \text{ satisfies (21)}\} / \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1).$$

Mrowka, Ozsváth and Yu show that the moduli space of Kähler vortices has a structure of complex manifold and the complex dimension is equal to the integer called the background degree. To calculate it, we recall briefly Seifert invariants. (The details of Seifert invariants can be found in Mrowka–Ozsváth–Yu [21, Section 2] and Furuta–Steer [7].)

Orbifold line bundles over an orbifold Riemann surface can be classified by pairs of integers called Seifert invariants:

$$\mathbf{b} = (b, \beta_1, \dots, \beta_k). \quad (22)$$

Here k is the number of singular points of the base orbifold, β_i is an integer with $0 \leq \beta_i < m_i$, and b is the integer which is called the background degree of the orbifold line bundle. The integer β_i is defined by local structure of the orbifold line bundle around the i -th singular point. The background degree b is the degree of the de-singularisation of the orbifold line bundle. For an orbifold line bundle $E \rightarrow \Sigma$ with the Seifert invariant (22), the degree of the line bundle is calculated by the formula

$$\text{deg}(E) = b + \frac{\beta_1}{m_1} + \dots + \frac{\beta_k}{m_k}. \quad (23)$$

Define d as the real number $\frac{i}{2\pi} \int_{[P/S^1]} F_A$ which is independent of the choice of $A \in \mathcal{A}(P)$.

Lemma 4.14. *The Seifert invariant of the orbifold line bundle $P \times_{S^1} \mathbb{C} \rightarrow \Sigma$ is $\mathbf{b} = (ad, 0, \dots, 0)$.*

Proof. Note that the triple $(\widetilde{U}_\alpha \times \mathbb{C}, \varphi_\alpha \times \text{id}_{\mathbb{C}}, \Gamma_\alpha)$ is a slice for the S^1 -action on $P \times \mathbb{C}$. Here the right action of Γ_α on $U_\alpha \times \mathbb{C}$ is given by

$$(z, x) \cdot t = (zt, t^a x) = (zt, x).$$

The last identity follows Assumption 4.11, and therefore $\beta_1 = \dots = \beta_k = 0$ in the description of Seifert invariant (22).

Because of the identity (23), the background degree is equal to the degree of the orbifold line bundle $P \times_{S^1} \mathbb{C} \rightarrow \Sigma$. Direct calculation shows that the degree is equal to ad . \square

Remark 4.15. The above lemma implies that ad is an integer under Assumption 4.11.

Applying Theorem 5 of Mrowka–Ozsváth–Yu [21], we obtain the following statement.

Proposition 4.16. *The moduli space \mathcal{M}^K of Kähler vortices is empty if*

$$d > \frac{\tau \text{Vol}(\Sigma)}{4\pi},$$

and it is naturally diffeomorphic to the ad -fold symmetric product of Σ , $\text{Sym}^{ad}(\Sigma)$ if

$$d < \frac{\tau \text{Vol}(\Sigma)}{4\pi}. \quad (24)$$

Let $\widetilde{\mathcal{M}}(Z_1 \rightrightarrows Z_0)$ and $\widetilde{\mathcal{M}}^K$ be the spaces of solutions to (20) and (21), respectively. By definition $\mathcal{M}(Z_1 \rightrightarrows Z_0) = \widetilde{\mathcal{M}}(Z_1 \rightrightarrows Z_0)/\mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1)$ and $\mathcal{M}^K = \widetilde{\mathcal{M}}^K/\mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1)$ hold. Now we compare $\mathcal{M}(Z_1 \rightrightarrows Z_0)$ with \mathcal{M}^K . The following linear isomorphism

$$\Psi : \Omega^1(Z_0, i\mathbb{R})_\rho \times \mathcal{C}^\infty(Z_0, \mathbb{C})_\rho \rightarrow \Omega^1(Z_0, i\mathbb{R})_a \times \mathcal{C}^\infty(Z_0, \mathbb{C})_\rho; (\theta, f) \mapsto (-a\theta, af)$$

gives a one-to-one correspondence between $\widetilde{\mathcal{M}}(Z_1 \rightrightarrows Z_0)$ and $\widetilde{\mathcal{M}}^K$. Moreover the map Ψ satisfies

$$\Psi((\theta, f) \cdot g) = \Psi(\theta, f) \cdot g^{-a} \quad (25)$$

for $\theta \in \Omega^1(Z_0, i\mathbb{R})_\rho$, $f \in \mathcal{C}^\infty(Z_0, \mathbb{C})_\rho$ and $g \in \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1)$. Therefore Ψ induces a well-defined surjective map $\overline{\Psi} : \mathcal{M}(Z_1 \rightrightarrows Z_0) \rightarrow \mathcal{M}^K$.

In general, the map $\overline{\Psi}$ is not injective. Define a covering map $\sigma : S^1 \rightarrow S^1$ by $\sigma(t) = t^a$ and a map $\tilde{\sigma} : \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1) \rightarrow \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1)$ by $\tilde{\sigma}(g) = \sigma \circ g$. Then Ψ induces a bijection between $\mathcal{M}(Z_1 \rightrightarrows Z_0)$ and $\widetilde{\mathcal{M}}^K/\text{image } \tilde{\sigma}$, and each fibre of the natural projection $\widetilde{\mathcal{M}}^K/\text{image } \tilde{\sigma} \rightarrow \widetilde{\mathcal{M}}^K/\mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1) = \mathcal{M}^K$ can be identified with $\text{coker } \tilde{\sigma}$.

$$\begin{array}{ccc} \mathcal{M}(Z_1 \rightrightarrows Z_0) & \xleftrightarrow{1:1} & \widetilde{\mathcal{M}}^K/\text{image } \tilde{\sigma} \\ & \searrow \overline{\Psi} & \downarrow \\ & & \mathcal{M}^K. \end{array}$$

Remark 4.17. If the inequality (24) holds, then the action of $\mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1)$ on $\widetilde{\mathcal{M}}^K$ is free.

In general, the cokernel of $\tilde{\sigma}$ is complicated, but it is trivial if the underlying space Σ of \mathfrak{X} is simply-connected.

Lemma 4.18. *If the underlying space Σ is simply-connected (i.e. of genus zero), then for any $g \in \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1)$ there is $g_0 \in \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1)$ such that $g = g_0^a$.*

Proof. Let $\pi : Z_0 \rightarrow Z_0/Z_1 = \Sigma$ be the quotient map. Any $g \in \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1)$ induces a continuous map $h : \Sigma \rightarrow S^1$ such that $g = h \circ \pi$. Since Σ is simply-connected, h has a lift: a continuous map $\tilde{h} : \Sigma \rightarrow S^1$ satisfying $\sigma \circ \tilde{h} = h$.

Since $\sigma \circ (\tilde{h} \circ \pi) = h \circ \pi = g$, the continuous map $\tilde{h} \circ \pi : Z_0 \rightarrow S^1$ is a lift of $g : Z_0 \rightarrow S^1$, hence the composition $\tilde{h} \circ \pi$ is smooth. Since $\tilde{h} \circ \pi \circ \text{src} = \tilde{h} \circ \pi \circ \text{tgt}$, $\tilde{h} \circ \pi \in \mathcal{C}^\infty(Z_1 \rightrightarrows Z_0, S^1)$. The map $g_0 = \tilde{h} \circ \pi$ is what we want. \square

Identifying $\mathcal{M}(P)$ with $\mathcal{M}(Z_1 \rightrightarrows Z_0)$, we obtain the following statement.

Theorem 4.19. *If Σ is simply-connected, then the moduli space $\mathcal{M}(P)$ is empty if*

$$d > \frac{\tau \text{Vol}(\Sigma)}{4\pi},$$

and it has a smooth structure which is diffeomorphic to the complex projective space $\mathbb{C}P^{ad}$ if

$$d < \frac{\tau \text{Vol}(\Sigma)}{4\pi}.$$

Acknowledgments The author would like to thank the Max-Planck-Institut für Mathematik in Bonn for providing financial support and excellent environment during my stay. He would also like to thank Andreas Ott and Nuno Miguel Romão for valuable comments and discussions.

References

- [1] Kai Behrend. Cohomology of stacks. In *Intersection theory and moduli*, ICTP Lect. Notes, XIX, pages 249–294 (electronic). Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004.
- [2] Kai Behrend and Ping Xu. Differentiable stacks and gerbes. *J. Symplectic Geom.*, 9(3):285–341, 2011.
- [3] Raoul Bott and Loring W. Tu. Equivariant characteristic classes in the Cartan model. In *Geometry, analysis and applications (Varanasi, 2000)*, pages 3–20. World Sci. Publ., River Edge, NJ, 2001.

- [4] Weimin Chen and Yongbin Ruan. Orbifold Gromov-Witten theory. In *Orbifolds in mathematics and physics (Madison, WI, 2001)*, volume 310 of *Contemp. Math.*, pages 25–85. Amer. Math. Soc., Providence, RI, 2002.
- [5] Kai Cieliebak, Ana Rita Gaio, and Dietmar A. Salamon. J -holomorphic curves, moment maps, and invariants of Hamiltonian group actions. *Internat. Math. Res. Notices*, (16):831–882, 2000. For historical comments, see arXiv:math/9909122v1.
- [6] Kai Cieliebak, Ignasi Mundet i Riera, and Dietmar A. Salamon. Equivariant moduli problems, branched manifolds, and the Euler class. *Topology*, 42(3):641–700, 2003.
- [7] Mikio Furuta and Brian Steer. Seifert fibred homology 3-spheres and the Yang-Mills equations on Riemann surfaces with marked points. *Adv. Math.*, 96(1):38–102, 1992.
- [8] A. Rita Gaio and Dietmar A. Salamon. Gromov-Witten invariants of symplectic quotients and adiabatic limits. *J. Symplectic Geom.*, 3(1):55–159, 2005.
- [9] Jochen Heinloth. Notes on differentiable stacks. In *Mathematisches Institut, Georg-August-Universität Göttingen: Seminars Winter Term 2004/2005*, pages 1–32. Universitätsdrucke Göttingen, Göttingen, 2005.
- [10] I. Mundet i Riera. *Yang-Mills theory for symplectic fibrations*. PhD thesis, Madrid, 1999.
- [11] Masaki Kashiwara and Pierre Schapira. *Categories and sheaves*, volume 332 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2006.
- [12] Eugene Lerman. Orbifolds as stacks? *Enseign. Math. (2)*, 56(3-4):315–363, 2010.
- [13] Eugene Lerman and Anton Malkin. Hamiltonian group actions on symplectic Deligne–Mumford stacks and toric orbifolds. *Adv. Math.*, 229:984–1000, 2012.
- [14] Dusa McDuff and Dietmar Salamon. *Introduction to symplectic topology*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1998.
- [15] Dusa McDuff and Dietmar Salamon. *J-holomorphic curves and symplectic topology*, volume 52 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2004.
- [16] David Metzler. Topological and Smooth Stacks. Preprint, arXiv:math/0306176v1, 2003.
- [17] I. Moerdijk and J. Mrčun. *Introduction to foliations and Lie groupoids*, volume 91 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2003.
- [18] I. Moerdijk and D. A. Pronk. Orbifolds, sheaves and groupoids. *K-Theory*, 12(1):3–21, 1997.

- [19] I. Moerdijk and D. A. Pronk. Simplicial cohomology of orbifolds. *Indag. Math. (N.S.)*, 10(2):269–293, 1999.
- [20] Ieke Moerdijk. Orbifolds as groupoids: an introduction. In *Orbifolds in mathematics and physics (Madison, WI, 2001)*, volume 310 of *Contemp. Math.*, pages 205–222. Amer. Math. Soc., Providence, RI, 2002.
- [21] Tomasz Mrowka, Peter Ozsváth, and Baozhen Yu. Seiberg-Witten monopoles on Seifert fibered spaces. *Comm. Anal. Geom.*, 5(4):685–791, 1997.
- [22] Matthieu Romagny. Group actions on stacks and applications. *Michigan Math. J.*, 53(1):209–236, 2005.
- [23] Hironori Sakai. The symplectic Deligne-Mumford stack associated to a stacky polytope. *Results Math.*, 2012. <http://dx.doi.org/10.1007/s00025-012-0240-3>.
- [24] Peter Scott. The geometries of 3-manifolds. *Bull. London Math. Soc.*, 15(5):401–487, 1983.
- [25] Angelo Vistoli. Notes on Grothendieck topologies, fibered categories and descent theory. Available at <http://homepage.sns.it/vistoli/papers.html>, 2008.

Max-Planck-Institut für Mathematik
Vivatsgasse 7
53111 Bonn
GERMANY
E-mail: sakai@blueskyproject.net