

On the Diophantine Approximations of logarithms in cyclotomic fields.

L.A. Gutnik

*To 100th birthday of
Professor A.O.Gelfond.*

Acknowledgments. I express my deepest thanks to Professors
B.Z.Moroz, I.I. Piatetski-Shapiro,
A.G.Aleksandrov, P.Bundshuh and S.G.Gindikin for help and support.

Let $T \in \mathbb{R}$, $\{\Delta, m, n\} \in \mathbb{N}$, $\Delta \geq 2$, $K_m = \mathbb{Q}[\exp(2\pi i/m)]$ is a cyclotomic field, \mathbb{Z}_{K_m} is the ring of all the integers in K_m , $\Lambda(n)$ is the Mangold's function, $\epsilon^2 = \epsilon$. Let $\Lambda_0(m) = 0$, if m is odd and $\Lambda_0(m) = \Lambda(m/2)$, if m is even. Let $\omega_1(m) = (m-1)/2$, if m is odd, $\omega_1(m) = m/2 - 2$, if $m \equiv 2 \pmod{4}$ and $\omega_1(m) = m/2 - 1$, if $m \equiv 0 \pmod{4}$. Let

$$(1) \quad w_\Delta(T) = \sqrt{\frac{\sqrt{(\Delta^2(3-T^2)+1)^2 + 16\Delta^4 T^2} + \Delta^2(3-T^2) + 1}{2}},$$

$$(2) \quad V_\Delta^* = (\Delta + 1) + \log((\Delta - 1)^{(\Delta-1)/2}(\Delta + 1)^{(\Delta+1)/2}\Delta^{-\Delta}) +$$

$$\frac{\pi}{2} \sum_{\mu=0}^1 (1 - 2\mu) \sum_{\kappa=1}^{[(d-1)/2]+\mu} \cot\left(\frac{\pi\kappa}{d-1+2\mu}\right),$$

$$(3) \quad V_\Delta(m) = V^* + (\Delta + 1)\Lambda_0(m)/\phi(m),$$

$$(4) \quad l_\Delta(\epsilon, T) = -\log(4(\Delta + 1)^{\Delta+1}(1 - 1/\Delta)^{(\Delta-1)}) +$$

$$\begin{aligned} & \frac{1}{2} \log \left((2\Delta + (-1)^\epsilon w_\Delta(T) + (\Delta + 1))^2 + T^2 \Delta^2 \left(1 + \frac{(-1)^\epsilon 2\Delta}{w_\Delta(T)} \right)^2 \right) + \\ & \frac{1}{2} \log \left((2\Delta + (-1)^\epsilon w_\Delta(T) - (\Delta + 1))^2 + T^2 \Delta^2 \left(1 + \frac{(-1)^\epsilon 2\Delta}{w_\Delta(T)} \right)^2 \right) + \\ & \frac{(\Delta - 1)}{2} \log \left((2\Delta + (-1)^\epsilon w_\Delta(T))^2 + T^2 \Delta^2 \left(1 + \frac{(-1)^\epsilon 2\Delta}{w_\Delta(T)} \right)^2 \right), \end{aligned}$$

$$(5) \quad g_{\Delta, \epsilon}(m) = (-1)^\epsilon (l_\Delta(\epsilon, \tan(\pi\omega_1(m)/m) + V_\Delta(m))),$$

$$(6) \quad h_\Delta(m) = -V_\Delta(m) - l_\Delta(1, \tan(\pi/m)),$$

where $m \neq 2$, $k = 0, 1$. Let

$$\beta(\Delta, m) = g_{d,0}(m)/h_\Delta(m), \quad \alpha(\Delta, m) = \beta(\Delta, m) - 1 + g_{\Delta,1}(m)/h_\Delta(m).$$

Theorem. *Let $m \in \mathbb{N} \setminus \{1, 2, 6\}$ $\Delta \in \{5, 7\}$. Then*

$$(7) \quad h_\Delta(m) > 0$$

and for each $\varepsilon > 0$ there exists $C_{\Delta, m}(\varepsilon) > 0$ such that

$$(8) \quad \max_{\sigma \in \text{Gal}(K/\mathbb{Q})} (|q^\sigma \log((2 + \exp(2\pi i/m))^\sigma) - p^\sigma|) \geq C_{\Delta, m}(\varepsilon) \left(\max_{\sigma \in \text{Gal}(K_m/\mathbb{Q})} (|q^\sigma|)^{-\alpha(\Delta, m) - \varepsilon} \right),$$

where $p \in \mathbb{Z}_{K_m}$ and $q \in \mathbb{Z}_{K_m} \setminus \{0_{K_m}\}$; moreover, for any $q \in \mathbb{Z}_{K_m} \setminus \{0_{K_m}\}$ and any $\varepsilon > 0$ there exists $C_{\Delta, m}^*(q, \varepsilon) > 0$ such that

$$(9) \quad b^{\beta(\Delta, m) + \varepsilon} \max_{\sigma \in \text{Gal}(K/\mathbb{Q})} (|q^\sigma b \log((2 + \exp(2\pi i/m))^\sigma) - p^\sigma|) \geq C_{\Delta, m}^*(q, \varepsilon),$$

where $p \in \mathbb{Z}_{K_m}$, $b \in \mathbb{N}$.

For the proof I use the same method, as in [37] – [67]. I work on the Riemann surface \mathfrak{F} of the function $\text{Log}(z)$ and identify it with the direct product of the multiplicative group $\mathbb{R}_+^* = \{r \in \mathbb{R} : r > 0\}$ of all the positive real numbers with the operation \times , not to be written down explicitly as usual, and the additive group \mathbb{R} of all the real numbers, so that

$$z_1 z_2 = (r_1 r_2, \phi_1 + \phi_2)$$

for any two points $z_1 = (r_1, \phi_1)$ and $z_2 = (r_2, \phi_2)$ on \mathfrak{F} . I will illustrate the appearing situations on the half plain (ϕ, r) , where $r > 0$.

For each $z = (r, \phi) \in \mathfrak{F}$, let

$$\theta_0(z) = r \exp i\phi, \quad \text{Log}(z) = \ln(r) + i\phi, \quad \eta_\alpha^*(z) = (r, \phi - \alpha),$$

where $\alpha \in \mathbb{R}$. Clearly, $\text{Log}(z_1 z_2) = \text{Log}(z_1) + \text{Log}(z_2)$ for any $z_1 \in \mathfrak{F}$ $z_2 \in \mathfrak{F}$. Let $\rho(z_1, z_2) = |\text{Log}(z_1) - \text{Log}(z_2)|$, where $z_1 \in \mathfrak{F}$ and $z_2 \in \mathfrak{F}$; clearly, (\mathfrak{F}, ρ) is a metric space. Clearly, $\rho(z z_1, z z_2) = \rho(z_1, z_2)$ for any z_1, z_2 and z in \mathfrak{F} . Clearly, $\theta_0(z) = \exp(\text{Log}(z))$ for any $z \in \mathfrak{F}$. Clearly, for any $\alpha \in \mathbb{R}$ the map $z \rightarrow \eta_\alpha^*(z)$ is the bijection of \mathfrak{F} onto \mathfrak{F} and

$$\theta_0((\eta_\alpha^*)^m(z)) = \exp(-im\alpha)\theta_0(z)$$

for each $z = (r, \phi) \in \mathfrak{F}$, $\alpha \in \mathbb{R}$ and $m \in \mathbb{Z}$. Clearly, the group \mathfrak{F} may be considered as \mathbb{C} -linear space, if for any $z \in \mathfrak{F}$ and any $s \in \mathbb{C}$ we let

$$z^s = (|\exp(s\text{Log}(z))|, \Im(s\text{Log}(z))).$$

Let us fix a domain D in \mathfrak{F} . Let $f(z) = f^\wedge(r, \phi)$ for a complex-valued function $f(z)$ on D , It is well known that $f(z)$ is holomorphic in D if the complex-valued function $f^\wedge(r, \phi)$ of two real variables r and ϕ has continuous partial derivatives in D , and the Cauchy-Riemann conditions

$$(10) \quad r((\partial/\partial r)f^\wedge)(r, \phi) = -i((\partial/\partial \phi)f^\wedge)(r, \phi) :=$$

$$(\delta f)(z) := \theta_0(z)((\partial/\partial z)f)(z)$$

are satisfied for every point $z = (r, \phi) \in D$. The equalities (10) determine a differentiations $\frac{\partial}{\partial z}$ and $\delta = \theta_0(z)\frac{\partial}{\partial z}$ on the ring of all the holomorphic in the domain D functions. In particular, the function $\text{Log}(z)$ is holomorphic on \mathfrak{F} and we have the equalities

$$((\partial/\partial z)\text{Log})(z) = \theta_0(z^{-1}), \quad (\delta \text{Log})(z) = 1.$$

For the proof I use the functions of C.S.Mejer. Let $\Delta \in \mathbb{N} + 1$, $\delta_0 = 1/\Delta$,

$$\gamma_1 = (1 - \delta_0)/(1 + \delta_0), \quad d_l = \Delta + (-1)^l, \quad l = 1, 2.$$

To introduce the first of my auxiliary function $f_1(z, \nu)$, I use the auxiliary set

$$\Omega_0 = \{z \in \mathfrak{F} : |z| \leq 1\}.$$

I prove that, for each $\nu \in \mathbb{N}$, the function $f_1(z, \nu)$ belongs to the ring $\mathbb{Q}[\theta_0(z)]$; therefore using the principle of analytic continuation we may regard it as being defined in \mathfrak{F} . For $\nu \in \mathbb{N}$, let

$$(11) \quad f_1(z, \nu) = -(-1)^{\nu(\Delta+1)} G_{2,2}^{(1,1)} \left(z \left| \begin{matrix} -\nu d_1, 1 + \nu d_2 \\ 0, \nu \end{matrix} \right. \right) \\ = -(-1)^{\nu(\Delta+1)} \frac{1}{2\pi i} \int_{L_1} g_{2,2}^{(1,1)}(s) ds,$$

where

$$g_{2,2}^{(1,1)}(s) = \theta_0(z^s)\Gamma(-s)\Gamma(1 + d_1\nu + s)/(\Gamma(1 - \nu + s)\Gamma(1 + d_2\nu - s))$$

and the curve L_1 passes from $+\infty$ to $+\infty$ encircling the set $\mathbb{N} - 1$ in the negative direction, but not including any point of the set $-\mathbb{N}$. So, for the parameters of the Meyer's functions we have

$$p = q = 2, m = n = 1, a_1 = -\nu d_1, a_2 = 1 + \nu d_2, b_1 = 0, b_2 = \nu,$$

$$\Delta^* = \left(\sum_{k=1}^q b_k \right) - \sum_{j=1}^p a_j = -\nu - 1 < -1,$$

and, since we take $|z| \leq 1$, convergence conditions of the integral in (11) hold. To compute the function $f_1(z, \nu)$, we use the following formula

$$(12) \quad G = (-1)^k \sum_{s \in S_k} Res(g; s),$$

where $k = 1$, G denotes the integral (11) with $L = L_k$, g denotes the integrand of the integral (11), S_k denotes the set of all the unremovable singularities of g encircled by L_k , and $Res(g; s)$ denotes the residue of the function g at the point s . Then we obtain the equality

$$f_1(z, \nu) = (\nu d_1)! / (\nu \Delta)! z^\nu (-1)^{\nu \Delta} \sum_{k=0}^{\nu \Delta} (-\theta_0(z))^k \binom{\nu \Delta}{k} \binom{\nu \Delta + k}{\nu d_1}.$$

Therefore, as it has been already remarked, using the principle of analytic continuation we may regard it as being defined in \mathfrak{F} . Let

$$\Omega_1 = \{z \in \mathfrak{F} : |z| \geq 1\}.$$

Now, let me introduce my second auxiliary function defined for $z \in \Omega_1$. For $\nu \in \mathbb{N}$, let

$$(13) \quad f_2(z, \nu) = -(-1)^{\nu(\Delta+1)} G_{2,2}^{(2,1)} \left(z \left| \begin{matrix} -\nu d_1, & 1 + \nu d_2 \\ 0, & \nu \end{matrix} \right. \right) = -(-1)^{\nu(\Delta+1)} \frac{1}{2\pi i} \int_{L_2} g_{2,2}^{(2,1)}(s) ds,$$

where

$$g_{2,2}^{(2,1)}(s) = \theta_0((\eta_\pi(z))^s)\Gamma(-s)\Gamma(\nu - s)\Gamma(1 + d_1\nu + s)/\Gamma(1 + d_2\nu - s).$$

and the curve L_2 passes from $-\infty$ to $-\infty$ encircling the set $-\mathbb{N}$ in the positive direction, but not including any point of the set $\mathbb{N} - 1$. So, for the parameters of the Meyer's functions we have

$$p = q = m = 2, n = 1, a_1 = -\nu d_1, a_2 = 1 + \nu d_2, b_1 = 0, b_2 = \nu,$$

$$\Delta^* = \left(\sum_{k=1}^q b_k \right) - \sum_{j=1}^p a_j = -nu - 1 < -1,$$

and, since we take $|z| \geq 1$, convergence conditions of the integral in (13) hold. To compute the function $f_2(z, \nu)$, we use the formula (12) where $k = 2$, G denotes the integral in (13) with $L = L_k$, g denotes the integrand of the integral in (13), S_k denotes the set of all the unremovable singularities of g encircled by L_k , and $Res(g; s)$ denotes the residue of the function g at the point s . Then we obtain the equality

$$(14) \quad f_2(z, \nu)(\nu\Delta)!/(\nu d_1)! = (-1)^\nu \sum_{t=\nu+1}^{\infty} R_0(t; \nu)\theta_0(z^{-t+\nu}),$$

where

$$R_0(t; \nu) = (\nu\Delta)!/(\nu d_1)! \left(\prod_{\kappa=\nu+1}^{\nu\Delta} (t - \kappa) \right) \prod_{\kappa=0}^{\nu\Delta} (t + \kappa)^{-1}.$$

Let further

$$(15) \quad f_k^*(z, \nu) = f_k(z, \nu)(\nu\Delta)!/(\nu d_1!),$$

where $k = 1, 2$. Expanding the function $R_0(t; \nu)$ into partial fractions, we obtain the equality

$$R_0(t; \nu) = \sum_{k=0}^{\nu\Delta} \alpha_{\nu, k}^*/(t + k)$$

with

$$(16) \quad \alpha_{\nu, k}^* = (-1)^{\nu+\nu\Delta+k} \binom{\nu\Delta}{k} \binom{\nu\Delta + k}{\nu\Delta - \nu},$$

where $k = 0, \dots, \nu\Delta$. It follows from (13), (14), (15) and (16) that

$$(17) \quad \begin{aligned} f_2^*(z, \nu) &= (-\theta_0(z))^\nu \sum_{t=1+\nu}^{+\infty} (\theta_0(z))^{-t} R_0(t; \nu) = \\ &= (-\theta_0(z))^\nu \sum_{t=1+\nu}^{+\infty} (\theta_0(z))^{-t-k+k} \sum_{k=0}^{\nu\Delta} \alpha_{\nu, k}^*/(t + k) \\ &= (-\theta_0(z))^\nu \sum_{t=1+\nu}^{+\infty} ((\theta_0(z))^{-t-k}/(t + k)) \sum_{k=0}^{\nu\Delta} \alpha_{\nu, k}^*(\theta_0(z))^k = \\ &= (-\theta_0(z))^\nu \sum_{k=0}^{\nu\Delta} \alpha_{\nu, k}^*(\theta_0(z))^k \sum_{\tau=1+\nu+k}^{+\infty} ((\theta_0(z))^{-\tau}/\tau) = \\ &= \alpha^*(z; \nu)(-\log(1 - 1/\theta_0(z))) - \phi^*(z; \nu), \end{aligned}$$

where $\log(\zeta)$ is a branch of $Log(\zeta)$ with $|\arg(\zeta)| < \pi$,

$$(18) \quad \alpha^*(z; \nu) = (-\theta_0(z))^\nu \sum_{k=0}^{\nu\Delta} \alpha_{\nu, k}^*(\theta_0(z))^k = f_1^*(z; \nu),$$

$$\begin{aligned}
(19) \quad \phi^*(z; \nu) &= (-\theta_0(z))^\nu \sum_{k=0}^{\nu\Delta} \alpha_{\nu,k}^* (\theta_0(z))^k \sum_{\tau=1}^{\nu+k} ((\theta_0(z))^{-\tau}/\tau) = \\
&(-\theta_0(z))^\nu \sum_{\tau=1}^{\nu} ((\theta_0(z))^{-\tau} \alpha^*(z; \nu)/\tau + \\
&(-\theta_0(z))^\nu \sum_{k=0}^{\nu\Delta} \alpha_{\nu,k}^* (\theta_0(z))^k \sum_{\tau=1+\nu}^{\nu+k} ((\theta_0(z))^{-\tau}/\tau)).
\end{aligned}$$

The change of order of summation by passage to (17) is possible, because the series in the second sum in (17) is convergent, if $|z| \geq 1$ and $\theta_0(z) \neq 1$. Since

$$\deg_t \left(\prod_{\kappa=\nu+1}^{\nu\Delta} (t - \kappa) \right) - \deg_t \left(\prod_{\kappa=0}^{\nu\Delta} (t + \kappa) \right) = -\nu - 1,$$

it follows that

$$\alpha^*(1; \nu) = \text{Res}(R_0(t; \nu); t = \infty) = 0$$

So in the domain $D_0 = \{z \in \mathfrak{F} : |z| > 1\}$ the function $f_2^*(z, \nu)$ coincides with the function

$$(20) \quad f_0^*(z, \nu) = \alpha^*(z; \nu)(-\log(1 - 1/\theta_0(z))) - \phi^*(z; \nu),$$

The form (20) may be used for various applications. Especially it is pleasant, when both $1/\theta_0(z)$ and $\alpha^*(z; \nu)$ for some z is integer algebraic number. The following Lemma corresponds to this remark.

Lemma 1. *Let $m \in \mathbb{N}$, $m > 2$, $m \neq 2p^\alpha$, where p run over the all the prime numbers and α run over \mathbb{N} . Then $1 + \exp(2\pi i/m)$ belongs to the group of the units of the field K_m . If $m = 2p^\alpha$, where p is a prime number and $\alpha \in \mathbb{N}$, then the ideal $\mathfrak{l} = (1 + \exp(2\pi i/m))$ is a prime ideal in the field K_m , and $\mathfrak{l}^{\phi(m)} = (p)$.*

Proof. Let polynomial $\Phi_m(z)$ is irreducible over \mathbb{Q} , has the leading coefficient equal to one and $\Phi_m(\exp(2\pi i/m)) = 0$. Let $\Lambda(n)$, as usual, denotes the Mangold's function. Since (see, for example, [27], end of the chapter 3)

$$\Phi_m(z) = \prod_{d|m} (z^{m/d} - 1)^{\mu(d)},$$

it follows that

$$\Phi_m(-1) = (-2)^{\left(\sum_{d|m} \mu(d)\right)} = 1,$$

if $m \in 1 + 2\mathbb{N}$,

$$\Phi_m(z) = \prod_{d|(m/2)} (((z)^{m/(2d)} - 1)^{\mu(2)} ((-z)^{m/d} - 1)/((-z) - 1))^{\mu(d)},$$

$$\begin{aligned}
\Phi_m(-1) &= \lim_{z \rightarrow -1} \prod_{d|(m/2)} (((-z)^{m/d} - 1)/((-z) - 1))^{\mu(d)} \times \\
&(-2)^{\mu(2)} \left(\sum_{d|(m/2)} \mu(d) \right) =
\end{aligned}$$

$$\exp\left(\sum_{d|(m/2)} \ln(m/(2d))\mu(2d)\right) = \exp(\Lambda(m/2)),$$

if $m \in 2(1 + 2\mathbb{N})$,

$$\Phi_m(z) = \prod_{d|(m/2)} (((-z)^{m/d} - 1)/((-z) - 1))^{\mu(d)},$$

and

$$\Phi_m(-1) = \lim_{z \rightarrow -1} \prod_{d|(m/2)} (((-z)^{m/d} - 1)/((-z) - 1))^{\mu(d)} =$$

$$\exp\left(\sum_{d|m/2} \ln(m/(2d))\mu(d)\right) = \exp(\Lambda(m/2)),$$

if $m \in 4\mathbb{N}$. If $m = 2p^\alpha$ with $\alpha \in \mathbb{N}$, then $\Phi_m(-1) = \exp(\Lambda(m/2)) = p$, and ideals $\mathfrak{l}_k = (1 + \exp(2\pi ik/m))$, where $(k, m) = 1$, divide each other and in the standard equality $efg = n$ (see, [27], chapter 3, section 10) we have

$$e = n = \phi(m), \quad f = g = 1. \blacksquare$$

In connection with the above remark and with the Lemma 1, the following case is interesting for us:

$$(21) \quad \begin{aligned} \theta_0(z) &= (-\rho)(1 + \exp(-i\beta)) = -(\rho \exp(i\beta/2))/(2\cos(\beta/2)) = \\ &= -(\rho \exp(i\psi))(2\cos(\psi)) = -(1 + i \tan(\psi))/2 \end{aligned}$$

with $\rho > 2/3$, $|\beta| < \pi$ and $-\pi/2 < \psi = \beta/2 < \pi/2$; then

$$\Re(1 - 1/\theta_0(z)) = \Re(2 + \exp(i\beta)/\rho) > 1/2,$$

and we have no problems with $\log(1 - 1/\theta_0(z))$. Of course, according to the Lemma 1, the case $\rho = 1$ is interesting especially. So, we will take further

$$(22) \quad z = (\rho/(2\cos(\psi)), \psi - \pi) = (\rho/(-2\cos(\theta)), \theta),$$

where $\rho > 2/3$, $|\psi| < \pi/2$ and $-3\pi/2 < \theta = \psi - \pi < -\pi/2$; clearly, the function (20) is analytic in the domain

$$\begin{aligned} D_1 &= \{z = (\rho(2\cos(\psi))^{-1}, \psi - \pi) : \rho > 2/3, -\pi/2 < \psi < \pi/2\} = \\ &= \{z = ((-2\rho\cos(\theta))^{-1}, \theta) : \rho > 2/3, -3\pi/2 < \theta < -\pi/2\}. \end{aligned}$$

Let

$$(23) \quad D_2(\delta_0) = \{z \in \mathfrak{F} : |z| > 1 + \delta_0/2\}, \quad D_3 = D_2(\delta_0) \cup D_1.$$

So, the function $f_2^*(z, \nu)$ coincides with the function (20) in $D_2(\delta_0) \subset D_0$. Since $D_2(\delta_0) \cap D_1 \neq \emptyset$, it follows that the join $D_3 = D_2(\delta_0) \cup D_1$ of the domains $D_2(\delta_0)$ and D_1 is a domain in \mathfrak{F} and the function (20) is analytic in this domain.

The conditions, which imply the equality

$$(24) \quad (-1)^{m+p-n} \exp(-i\alpha)\theta_0(z) \times \\ \left(\left(\prod_{j=1}^p (\delta + 1 - a_j) \right) (G \circ \eta_\alpha^*) \right) (z) = \left(\left(\prod_{k=1}^q (\delta - b_k) \right) (G \circ \eta_\alpha^*) \right) (z)$$

hold in our case for the Mejer's function

$$G = G_{p,q}^{(m,n)} \left(z \left| \begin{array}{c} a_1, \dots, a_p \\ b_1, \dots, b_q \end{array} \right. \right).$$

We have $p = q = 2$, $m = n = 1$, $\alpha = 0$ for the function $f_1(z, \nu)$ and the equation (24) takes the form

$$\theta_0(z)((\delta + 1 + d_1\nu)(\delta - d_2\nu)f_1)(z, \nu) = (\delta(\delta - \nu)f_1)(z, \nu)$$

We have $p = q = m = 2$, $n = 1$, $\alpha = \pi$ for the function $f_2(z, \nu)$ and the equation (24) takes the form

$$\theta_0(z)((\delta + 1 + d_1\nu)(\delta - d_2\nu)f_2)(z, \nu) = (\delta(\delta - \nu)f_2)(z, \nu).$$

We see that both the functions $f(z, \nu) = f_k^*(z, \nu)$, where $k = 1, 2$ satisfy to the same differential equation

$$(25) \quad \theta_0(z)(\delta + 1 + d_1\nu)(\delta - d_2\nu)f(z, \nu) = (\delta(\delta - \nu)f)(z, \nu).$$

in the domain D_0 . According to the general properties of the Mejer's functions we have the equality

$$(26) \quad \left(\prod_{\kappa=1}^{\Delta-1} (\nu(\Delta - 1) + \kappa) \right) \prod_{\kappa=1}^{d_2} (\delta - d_2\nu - \kappa) f_k^*(z, \nu + 1) = \\ \left(\prod_{\kappa=1}^{\Delta} (\nu\Delta + \kappa) \right) (\delta - \nu) \prod_{\kappa=1}^{d_1} (\delta + d_1\nu + \kappa) f_k^*(z, \nu),$$

where $k = 1, 2$ and $z \in D_0$. Since $f_0^*(z, \nu)$ and polynomial $f_1^*(z, \nu)$ are analytic in the domain $D_0 \cup D_1$, and $f_0^*(z, \nu)$ coincides with $f_2^*(z, \nu)$, it follows that the equations (25) and (26) hold in $D_0 \cup D_1$ for $k = 0, 1$.

Let

$$(27) \quad D^\vee(w, \eta) = (\eta + 1)(\eta + \gamma_1) - 2(1 + \gamma_1)w\eta,$$

$$(28) \quad D^\wedge(z, \eta) = D^\vee(\theta_0(z), \eta),$$

where, in view of (21),

$$(29) \quad w = \theta_0(z) = -r \exp(i\psi), \quad r = 1/(2\cos(\psi)), \quad |\psi| < \pi/2.$$

In view of (29), the polynomial (27) coincides with the polynomial (1) in [59]. Let

$$(30) \quad h^\sim(\eta) = (\eta - 1)(1 - \delta_0)^{-d_1}(\eta + 1)2^{-2} \eta^{d_1}.$$

As in [51], we consider ν^{-1} as an independent variable taking its values in the field \mathbb{C} including 0. Let F be a bounded closed subset of \mathfrak{F} (in particular, this compact F may be an one-point set). Let $\mathfrak{H}_0(F)$ be the subring of all those functions in $\mathbb{Q}(w)$, which are well defined for every $w \in \theta_0(F)$. For $\varepsilon \in (0, 1)$, let $\mathfrak{H}(F, \varepsilon)$ be the subring of all those functions in $\mathbb{Q}(w, \nu^{-1})$, which are well defined for every (w, ν^{-1}) with $w \in \theta_0(F)$, $|\nu^{-1}| \leq \varepsilon_0$.

Lemma 2. *Let F be a closed bounded subset of $D_0 \cup D_1$ (in particular, F may be an one-point set). Let further for any $z \in F$ the polynomial (28) has only simple roots and on the set of all the roots η of the polynomial $D^\wedge(z, \eta)$ the map*

$$(31) \quad \eta \rightarrow h^\sim(\eta)$$

is injective. Then there is $\varepsilon \in (0, 1)$ such that, for any $z \in F, \nu \in \mathbb{N} + [1/\varepsilon]$, the functions $f_0^(z, \nu), f_1^*(z, \nu) = \alpha^*(z; \nu)$ and $\phi^*(z; \nu)$ are solutions of the difference equation*

$$(32) \quad x(z, \nu + 2) + \sum_{j=0}^1 q_j^*(z, \nu^{-1})x(z, \nu + j) = 0,$$

moreover,

$$(33) \quad q_j^*(z, \nu^{-1}) \in \mathfrak{H}(F, \varepsilon)$$

for $j = 0, 1$, and trinomial

$$(34) \quad w^2 + \sum_{j=0}^1 q_j^*(z, 0)w^j$$

coincides with

$$(35) \quad \prod_{k=0}^1 (w - h(\eta_k)),$$

if

$$\prod_{k=0}^1 (w - \eta_k),$$

coincides with $D^\vee(w, \eta)$ from (27).

Proof. Proof may be found in [51]. ■

This Lemma shows the importance of the properties of the roots of the polynomial (27). In correspondence with (22) and with notations in [59], let

$$(36) \quad \rho > 2/3, r = \rho/(2 \cos(\psi)), t = \cos(\psi), |\psi| < \pi/2.$$

Let $u = r^2, \delta_0 \leq 1/2 < 2/3 < \rho$. Then

$$(37) \quad 2\delta_0 \leq 2/5 < 2/3 < \rho < 2\sqrt{u} = 2r.$$

Clearly,

$$(\partial/\partial\psi)r = (\rho \sin(\psi))/(2 \cos^2(\psi)) = -2\rho(\sin(\psi) - 1) - 2\rho/(\sin(\psi) + 1),$$

$(\partial/\partial\psi)^2 r = (2\rho \cos(\psi))/(\sin(\psi) - 1)^2 + (2\rho \cos(\psi))/(\sin(\psi) + 1)^2 > 0$,
if $|\psi| < \pi/2$ In view of (3.1.10) in [52],

$$(38) \quad |D_0(r, \psi, \delta_0)|^2 = r^4 + r^2 + (\delta_0/2)^4 + \\ 2r^2(\delta_0/2)^2(2t^2 - 1) + 2r(r^2 + (\delta_0/2)^2)t = \\ u^2 + u + (\delta_0/2)^4 + (\delta_0/2)^2(\rho^2 - 2u) + \rho(u + (\delta_0/2)^2) = \\ u^2 + u(\rho + 1 - (\delta_0)^2/2) + (\delta_0/2)^2(\rho^2 + \rho + (\delta_0/2)^2),$$

$$(39) \quad |R_0(r, \psi, \delta_0)|^2 = |D_0(r, \psi, \delta_0)| = \\ \sqrt{u^2 + u(\rho + 1 - (\delta_0)^2/2) + (\delta_0/2)^2(\rho^2 + \rho + (\delta_0/2)^2)}.$$

In view of (3.1.41) - (3.1.43) in [52] and (39),

$$(40) \quad p_1 = 8(|R_0^*(r, \psi, \delta_0)|^2 + |R_0(r, \psi, \delta_0)|^2)/(1 + \delta_0)^2 = \\ 8(r^2 + rt + 1/4 + |D_0(r, \psi, \delta_0)|)/(1 + \delta_0)^2 = 8(1 + \delta_0)^{-2} \times \\ \left(u + \rho/2 + 1/4 + \sqrt{u^2 + u(\rho + 1 - (\delta_0)^2/2) + (\delta_0/2)^2(\rho^2 + \rho + (\delta_0/2)^2)} \right),$$

$$(41) \quad p_2 = (8(|R_1^*(r, \psi, \delta_0)|^2 + |R_0(r, \psi, \delta_0)|^2))/(1 + \delta_0)^2 = \\ 8(r^2 - r\delta_0 t + (\delta_0)^2/4 + |D_0(r, \psi, \delta_0)|)/(1 + \delta_0)^2 = \\ 8(u - \delta_0\rho/2 + (\delta_0)^2/4)/(1 + \delta_0)^2 + \\ 8(1 + \delta_0)^{-2} \sqrt{u^2 + u(\rho + 1 - (\delta_0)^2/2) + (\delta_0/2)^2(\rho^2 + \rho + (\delta_0/2)^2)} = \\ 8(1 + \delta_0)^{-2} u(2 + (\rho + 1 - \delta_0\rho)/(2u) + O(1/u^2)),$$

$$(42) \quad q_1(r, \psi, \delta_0) = ((1 - \delta_0)/(1 + \delta_0))^2, \quad q_2(r, \psi, \delta_0) = \\ (4r/(1 + \delta_0))^2 = (16u)/(1 + \delta_0)^2.$$

In view of (91) in [59], (36) and (37),

$$(43) \quad s = s_0(r, \psi) = |r \exp(i\psi) + 1|/2 = \sqrt{(r^2 + 1 + 2r \cos(\psi))/4} = \\ \sqrt{(u + 1 + \rho)/4} \in (\max(|r - 1|/2, \delta_0/4), (r + 1)/2]$$

and

$$t = \cos(\psi) = (4s^2 - r^2 - 1)/(2r).$$

In view of (3.1.68) in [52], (3.1.70) - (3.1.71) in [52] and (39),

$$|R_{-1}^*(r, \psi, \delta_0)|^2 = r^2 + (2 + \delta_0)^2/4 + r(2 + \delta_0) \cos(\psi) = \\ u + (2 + \delta_0)^2/4 + \rho(2 + \delta_0)/2,$$

$$(44) \quad p_0 = 8(|R_{-1}^*(r, \psi, \delta_0)|^2 + |R_0(r, \psi, \delta_0)|^2)/(1 + \delta_0)^2 =$$

$$\frac{8(u + (2 + \delta_0)^2/4 + \rho(2 + \delta_0)/2)/(1 + \delta_0)^2 + 8(1 + \delta_0)^{-2}\sqrt{u^2 + u(\rho + 1 - (\delta_0)^2/2) + (\delta_0/2)^2(\rho^2 + \rho + (\delta_0/2)^2)}}{8(1 + \delta_0)^{-2}\sqrt{u^2 + u(\rho + 1 - (\delta_0)^2/2) + (\delta_0/2)^2(\rho^2 + \rho + (\delta_0/2)^2)}},$$

$$(45) \quad q_0(r, \psi, \delta_0)(1 + \delta_0)^2/16 = (r^2 + 1 + 2r\cos(\psi)) = (u + 1 + \rho).$$

According to Lemma 4.4 in [59], (23) and (37),

$$(46) \quad |\eta_1^\wedge(r, \psi, \delta_0) + \epsilon| < |\eta_0^\wedge(r, \psi, \delta_0) + \epsilon|,$$

if $\epsilon^2 = \epsilon$ and $z \in D_3$. Therefore, according to (40), (42) and (46),

$$(47) \quad (-1)^k(\partial/\partial u)|\eta_k^\wedge(r, \psi, \delta_0)| > 0,$$

where $\frac{1}{3} < \rho/2 < \sqrt{u} = r$, $k^2 = k$. According to a) and c) of the Lemma 4.6 in [59], and in view of (23) and (43),

$$(48) \quad |\eta_1^\wedge(r, \psi, \delta_0) - 1| < |\eta_0^\wedge(r, \psi, \delta_0) - 1|,$$

if $z \in D_3$. In view of (38),

$$(49) \quad |D_0(r, \psi, \delta_0)|^2 = u^2 + u(\rho + 1 - (\delta_0)^2/2) + (\delta_0/2)^2(\rho^2 + \rho + (\delta_0/2)^2) = (u + (\rho + 1)/2 - (\delta_0)^2/4)^2 + (\delta_0/2)^2(\rho^2 + \rho + (\delta_0/2)^2) - (((\rho + 1)/2)^2 - (\rho + 1)(\delta_0)^2/4 + (\delta_0/2)^4) = (u + (\rho + 1)/2 - (\delta_0)^2/4)^2 + (\delta_0/2)^2(\rho^2 + 2\rho + 1) - (\rho + 1)^2/4 = (u + (\rho + 1)/2 - (\delta_0)^2/4)^2 - (\rho + 1)^2(1 - (\delta_0)^2)/4.$$

Consequently,

$$(50) \quad |D_0(r, \psi, \delta_0)| = u + \frac{\rho + 1}{2} - \frac{(\delta_0)^2}{4} + O(1/u),$$

where $u \geq 1/4$. Since $u \geq 1/4 > (\delta_0)^2/4$, it follows that

$$u + (\rho + 1)/2 - (\delta_0)^2/4 > \sqrt{1 - (\delta_0)^2}(\rho + 1)/2.$$

If $\rho = 1, u = 1/4$ then in view of (49),

$$|D_0(r, \psi, \delta_0)|^2 = (5/4 - (\delta_0)^2/4)^2 - (1 - (\delta_0)^2) = (\tau - 5/4)^2 + 4\tau - 1,$$

where $0 < \tau = \frac{(\delta_0)^2}{4} < \frac{1}{100}$; moreover, in this case

$$(\partial/\partial \tau)|D_0(r, \psi, \delta_0)|^2 = 2\tau - 5/2 + 4 > 0;$$

therefore if $\delta_0 \leq 1/5$, then

$$|D_0(r, \psi, \delta_0)|^2 \Big|_{u=1/4, \rho=1} \leq (1, 24)^2 - 0, 96 = 0, 5776$$

and

$$|D_0(r, \psi, \delta_0)|^2 \Big|_{u=1/4, \rho=1} \leq 0, 76.$$

In view of (49),

$$1 < (\partial/\partial u)|D_0(r, \psi, \delta_0)| = \sqrt{\frac{(u + (\rho + 1)/2 - (\delta_0)^2/4)^2}{(u + (\rho + 1)/2 - \frac{(\delta_0)^2}{4})^2 - (\rho + 1)^2(1 - (\delta_0)^2)/4}} = 1 + O(1/u^2),$$

in view of (40), (41) and (44),

$$(51) \quad (\partial/\partial u)p_\epsilon = 8(2 + O(1/u^2))/(1 + \delta_0)^2,$$

where $\epsilon^3 = \epsilon$, and $(\partial/\partial u)|D_0(r, \psi, \delta_0)|$ decreases with increasing u ; consequently,

$$(\partial/\partial u)^2|D_0(r, \psi, \delta_0)| < 0,$$

if $u \geq 1/4$. In view of (40), (41) and (44),

$$(52) \quad (\partial/\partial u)^2 p_\epsilon = (\partial/\partial u)^2|D_0(r, \psi, \delta_0)| < 0,$$

where $u \geq 1/4$, $0 < \delta_0 < 2/3 < \rho$, $\epsilon^3 = \epsilon$. In view of (41), (42), (49) and (50), if $\rho = 1$, $u > 1/4$, $0 < \delta_0 \leq 1/5$, then

$$(53) \quad \begin{aligned} & q_2((\partial/\partial u)p_2)/(\partial/\partial u)q_2 - p_2/2 = \\ & 8u(1 + (u + 1 - (\delta_0)^2/4))/|D_0(r, \psi, \delta_0)|/(1 + \delta_0)^2 - \\ & 4(u - \delta_0/2 + (\delta_0)^2/4 + |D_0(r, \psi, \delta_0)|)/(1 + \delta_0)^2 = \\ & 4(u + \delta_0/2 - (\delta_0)^2/4)/(1 + \delta_0)^2 + \\ & 4((1 + \delta_0)^2|D_0(r, \psi, \delta_0)|)^{-1}(2u^2 + u(2 - (\delta_0)^2/2) - \\ & 4((1 + \delta_0)^2|D_0(r, \psi, \delta_0)|)^{-1}(u^2 + u(2 - (\delta_0)^2/2) + (\delta_0/2)^2(2 + (\delta_0/2)^2)) = \\ & 4(u + \delta_0/2 - (\delta_0)^2/4)/(1 + \delta_0)^2 + \\ & 4(1 + \delta_0)^2|D_0(r, \psi, \delta_0)|^{-1}(u^2 - (\delta_0/2)^2(2 + (\delta_0/2)^2)) > 0, \\ & q_2((\partial/\partial u)p_2)/(\partial/\partial u)q_2 - p_2 = \frac{8}{u}(1 + (u + 1 - (\delta_0)^2/4))/|D_0(r, \psi, \delta_0)|/(1 + \delta_0)^2 - \\ & 8(u - \delta_0/2 + (\delta_0)^2/4 + |D_0(r, \psi, \delta_0)|)/(1 + \delta_0)^2 = \\ & 8u(2 + O(1/u^2))/(1 + \delta_0)^2 - \\ & 8(u - \frac{\delta_0}{2} + \frac{(\delta_0)^2}{4} + u + 1 - \frac{(\delta_0)^2}{4} + O(1/u))/(1 + \delta_0)^2 = \\ & -8(1 - \frac{\delta_0}{2} + O(1/u))/(1 + \delta_0)^2. \end{aligned}$$

In view of (44), (45), (53), (49), (51), (50), if $\rho = 1$, $u > 1/4$, $0 < \delta_0 \leq 1/5$, then

$$(u + 1)(\partial/\partial u)p_0 - p_0/2 > 8(2u + 2)/(1 + \delta_0)^2 - 4(u + (2 + \delta_0)^2/4 + (2 + \delta_0)/2 + u + 1 - (\delta_0)^2/4)/(1 + \delta_0)^2 =$$

$$\frac{8}{7}1/2 + u - (3\delta_0)/4)/(1 + \delta_0)^2 > 0,$$

$$(54) \quad q_0((\partial/\partial u)p_0)/(\partial/\partial u)q_0 - p_0/2 = (u + 2)(\partial/\partial u)p_0 - p_0/2 > \\ (u + 1)(\partial/\partial u)p_0 - p_0/2 > 0,$$

$$(55) \quad q_0(\partial/\partial u)p_0)/(\partial/\partial u)q_0 - p_0 = 8(u + 2)(2 + O(1/u^2))/(1 + \delta_0)^2 - \\ 8(u + (2 + \delta_0)^2/4 + (2 + \delta_0)/2 + u + 1 - (\delta_0)^2/4)/(1 + \delta_0)^2 = \\ 8(4 + O(1/u))/(1 + \delta_0)^2 - (2 + \delta_0)^2/4 - (2 + \delta_0)/2 - 1 + (\delta_0)^2/4 + O(\frac{1}{7}u) = \\ 8(1 - (3/2)\delta_0 + O(1/u))/(1 + \delta_0)^2,$$

where $u > 1/4$. In view of (45), (54) and (52),

$$(\partial/\partial u)((q_0(\partial/\partial u)p_0)/(\partial/\partial u)q_0 - p_0)(\partial/\partial u)p_0 + (\partial/\partial u)q_0 = \\ (\partial/\partial u)((u + 2)(\partial/\partial u)p_0 - p_0)(\partial/\partial u)p_0 = \\ ((\partial/\partial u)p_0)^2 + ((u + 2)(\partial/\partial u)^2p_0 - (\partial/\partial u)p_0)(\partial/\partial u)p_0 + \\ ((u + 2)(\partial/\partial u)p_0 - p_0)(\partial/\partial u)^2p_0 = \\ ((u + 2)(\partial/\partial u)^2p_0)(\partial/\partial u)p_0 + ((u + 2)(\partial/\partial u)p_0 - p_0)(\partial/\partial u)^2p_0 = \\ (2(u + 2)(\partial/\partial u)p_0 - p_0)(\partial/\partial u)^2p_0 < 0.$$

Therefore, according to (55), (51) and (45),

$$(56) \quad \inf\{((u + 2)(\partial/\partial u)p_0 - p_0)(\partial/\partial u)p_0 + (\partial/\partial u)q_0 : u \geq 1/4\} = \\ \lim_{u \rightarrow +\infty} ((u(\partial/\partial u)p_0 - p_0)(\partial/\partial u)p_0 + (\partial/\partial u)q_0) = \\ 128(1 - (3/2)\delta_0)/(1 + \delta_0)^4 + 16/(1 + \delta_0)^2 > 0.$$

According to the Lemma 4.17 in [59] and in view of (53), (54), (56),

$$(57) \quad (\partial/\partial u)|\eta_0(r, \psi, \delta_0) + \epsilon|^2 > 0,$$

where $\epsilon^2 = 1$, $u > 1/4$,

$$(58) \quad (\partial/\partial u)|\eta_1(r, \psi, \delta_0) - 1|^2 < 0,$$

where $u > 1/4$. The following Lemma describes the behavior of the value $h^\sim(\eta_k(r, \psi, \delta_0))$ with $k^2 = k$ and h^s in (30).

Lemma 3. *If $\Delta \geq 5$, then*

$$(59) \quad (\partial/\partial u)(|h^\sim(\eta_0(r, \psi, \delta_0))|) > 0,$$

$$(\partial/\partial u)(|h^\sim(\eta_1(r, \psi, \delta_0))|) < 0,$$

where $u \in (1/4, +\infty)$.

Proof. The inequality (59) directly follows from (46), (57) and (30). So, we must prove the inequality (30) Clearly, if $\beta < 1$, $u > 1/4$ then

$$(\partial/\partial u)(u^{3/4} + (3/4)\beta u^{-1/4}) > 0,$$

We take

$$\beta = (4/3)(\delta_0/2)^2(2 + (\delta_0)^2)/4/(2 - (\delta_0)^2/2).$$

Then, clearly, $\beta < (\delta_0)^2 = 1/(\Delta)^2 < 1$. Therefore, in view of (40) and (49), if $\rho = 1$, then

$$p_1 u^{-1/4} = 8(1 + \delta_0)^{-2} \times \\ (u^{3/4} + (3/4)u^{-1/4} + \sqrt{u^{3/2} + u^{1/4}(\rho + 1 - (\delta_0)^2/2)(u^{3/4} + (3/4)\beta u^{-1/4})})$$

increases together with increasing $u \in (1/4, +\infty)$, and, in view of (42),

$$(60) \quad |\eta_0(r, \psi, \delta_0)|^2 u^{-1/4} = p_1 u^{-1/4}/2 + \sqrt{(p_1 u^{-1/4}/2)^2 - q_1 u^{-1/2}}$$

increases together with increasing $u \in (1/4, +\infty)$.

In view of (47), (42), (60), (57) and (58), if $\Delta \geq 5$, then

$$|\eta_1(r, \psi, \delta_0)|^{2(\Delta-1)} |(\eta_1(r, \psi, \delta_0))^2 - 1|^2 = \\ |\eta_1(r, \psi, \delta_0)|^{2(\Delta-5)} \frac{(q_1)^4}{(|\eta_0(r, \psi, \delta_0)|^2 u^{-1/4})^4} \times \\ \frac{16}{(1 + \delta_0)^2} |\eta_0(r, \psi, \delta_0) + 1|^{-2} |\eta_1(r, \psi, \delta_0) - 1|^2$$

decreases together with increasing $u \in (1/4, +\infty)$. ■

Let D is bounded domain in \mathbb{C} or \mathfrak{F} . and D^* is closure of D . Let

$$(61) \quad a_0^\sim(z), \dots, a_n^\sim(z)$$

are the functions continuous on D^* and analytic in D . Let $a_n^\sim(z) = 1$ for any $z \in D^*$. Let

$$(62) \quad T(z, \lambda) = \sum_{i=0}^n a_i^\sim(z) \lambda^k.$$

Let $s \in \mathbb{N}$, $n_i \in \mathbb{N} - 1$, where $i = 1, \dots, s$ and $\sum_{i=1}^s n_i = n$. We say that polynomial $T(z, \lambda)$ has (n_1, \dots, n_s) -disjoint system of roots on D^* , if for any $z \in D^*$ the set of all the roots λ of the polynomial $T(z, \lambda)$ splits in s classes $\mathfrak{K}_1(z), \dots, \mathfrak{K}_s(z)$ with following properties:

a) the sum of the multiplicities of the roots of the class \mathfrak{K}_i is equal to n_i for $i = 1, \dots, s$;

b) if $i \in [1, s] \cap \mathbb{N}$, $j \in (i, s] \cap \mathbb{N}$ and $n_i n_j \neq 0$, then the absolute value of each roots of the class $\mathfrak{K}_i(z)$ is greater than absolute value of the each roots of the class $\mathfrak{K}_j(z)$.

If the polynomial (62) has (n_1, \dots, n_s) -disjoint system of roots on D^* , then for each $i = 1, \dots, s$ we denote by $\rho_{i,0}^*(z)$ and $\rho_{i,1}^*(z)$ respectively the maximal and minimal absolute value of the roots of the class $\mathfrak{K}_i(z)$.

Let D is bounded domain in \mathfrak{F} such that $D^* \in D_3$. Let

$$(63) \quad F^\wedge(z, \eta) = \prod_{i=1}^2 (\theta_0(z) - h(\eta_{i-1}(r, \psi, \delta_0))),$$

$$n = s = 2, n_1 = n_2 = 1, \mathfrak{K}_i(z) = \{h(\eta_{i-1}(r, \psi, \delta_0))\},$$

$$\rho_{i,0} = \rho_{i,1} = |h(\eta_{i-1}(r, \psi, \delta_0))|,$$

where $i = 1, 2$.

Lemma 4. *The polynomial $F^\wedge(z, \eta)$ in (63) has (1, 1)-disjoint system of roots on D^* .*

Proof. The assertion of the Lemma follows from (46) and (48). ■

Corollary. *The map (31) is injective for every $z \in D^*$; all the conditions of the Lemma 2 are fulfilled for the functions $f_0^*(z, \nu)$ from (20), $\alpha^*(z, \nu)$ from (18) and $\phi^*(z, \nu)$ from (19) in every $z \in D^*$; therefore for every $z \in D^*$ these functions are solutions of the difference equation of Poincaré type (32), and the polynomial (35) coincides with characteristical polynomial of this equation. ■*

Let for each $\nu \in \mathbb{N} - 1$ are given continuous on D^* functions

$$(64) \quad a_0(z; \nu), \dots, a_n(z, \nu),$$

which are analytic in D .

Let $a_n(z; \nu) = 1$ for any $z \in D^*$ and any $\nu \in \mathbb{N} - 1$. Let for any $i = 1, \dots, n - 1$ the sequence of functions $a_i(z; \nu)$ converges to $a_i^\sim(z)$ uniformly on D^* , when $\nu \rightarrow \infty$. Let us consider now the difference equation

$$(65) \quad a_0(z; \nu)y(\nu + 0) + \dots + a_n(z; \nu)y(\nu + n) = 0,$$

i.e. we consider a difference equation of the Poincaré type, coefficients (64) of this equation are continuous on D^* and analytic in D , and they uniformly converge to limit functions (61), when $\nu \rightarrow \infty$.

Lemma 5. *Let polynomial (62) has (n_1, \dots, n_s) -disjoint system of roots on D^* . Let $y(z, \nu)$ is a solution of the equation (65), and this solution is continuous on D^* and analytic on D . Let further $i \in [1, s] \cap \mathbb{Z}$. Let us consider the set of all the $z \in D$, for which the following inequality holds*

$$(66) \quad \limsup_{\nu \in \mathbb{N}, \nu \rightarrow \infty} |y(z, \nu)|^{1/\nu} < \rho_{i,1}(z);$$

if this set has a limit point in D , then the inequality (66) holds in D^* .

Proof. The proof may be found in [31] (Theorem 1 and its Corollary). ■

Lemma 6. *Let D is bounded domain in \mathfrak{F} such that $D^* \in D_3$. Then*

$$(67) \quad \limsup_{\nu \in \mathbb{N}, \nu \rightarrow \infty} |f_0^*(z, \nu)|^{1/\nu} < \rho_{1,1}(z) = |h^\sim(\eta_0(r, \psi, \delta_0))|$$

for any $z \in D^*$.

Proof. In view of (23), expanding the domain D , if necessary, we can suppose that $\{(r, \phi) : r \in [2, 3], \phi = 0\} \in D$. Making use the same arguments, as in [55], Lemma 4.2.1, we see that the inequality (67) holds for

any point $z = (r, \phi) \in \{r \in [2, 3], \phi = 0\}$. According to the Lemma 5, the inequality (67) holds for any $z \in D^*$. ■

For each prime $p \in \mathbb{N}$ let v_p denotes the p -adic valuation on \mathbb{Q} .

Lemma 7. *Let $p \in \mathbb{N} + 2$ is a prime number,*

$$d \in \mathbb{N} - 1, r \in \mathbb{N} - 1, r < p.$$

Then

$$v_p((dp + r)! / ((-p)^d d! r!) - 1) \geq 1.$$

Lemma 8. *Let $p \in \mathbb{N} + 2$ is a prime number, $d \in \mathbb{N} - 1, d_1 \in \mathbb{N} - 1,$*

$$(68) \quad r \in [0, p - 1] \cap \mathbb{N}, r_1 \in [0, p - 1] \cap \mathbb{N}, d_1 p + r_1 \leq dp + r.$$

Then

$$(69) \quad v_p \left(\binom{dp + r}{d_1 p + r_1} \right) = v_p \left(\binom{d}{d_1} \right),$$

if $r_1 \leq r,$

$$(70) \quad v_p \left(\binom{dp + r}{d_1 p + r_1} \left(\binom{d}{d_1} \binom{r}{r_1} \right)^{-1} - 1 \right) \geq 1,$$

if $r_1 \leq r,$

$$(71) \quad v_p \left(\binom{dp + r}{d_1 p + r_1} \right) = 1 + v_p \left((d - d_1) \binom{d}{d_1} \right),$$

if $r < r_1,$

$$(72) \quad v_p \left((-1)^{r_1 - r - 1} \binom{dp + r}{d_1 p + r_1} \binom{r_1}{r} (r_1 - r) \left(p \binom{d}{d_1} (d - d_1) \right)^{-1} - 1 \right) \geq 1,$$

Proof. Clearly, $d_1 \leq d$. If $r_1 \leq r$, then let $r_2 = r - r_1, d_2 = d - d_1$. On the other hand, if $r_1 > r$, then, in view of (68), $d \geq d_1 + 1$; therefore in this case we let

$$(73) \quad r_2 = p + r - r_1, d_2 = d - d_1 - 1.$$

Then $d = d_1 + d_2, r = r_1 + r_2,$

$$\binom{dp + r}{d_1 p + r_1} = (dp + r)! / ((d_1 p + r_1)! (d_2 p + r_2)!)^{-1}.$$

Accordingg to the Lemma 7,

$$(74) \quad v_p \left(\binom{dp + r}{d_1 p + r_1} (-p)^{-d + d_1 + d_2} d_1! r_1! d_2! r_2! / (d! r!) - 1 \right) \geq 1,$$

$$(75) \quad v_p \left(\binom{dp + r}{d_1 p + r_1} \right) = d - d_1 - d_2 + v_p(d! r! / (d_1! r_1! d_2! r_2!)).$$

The equality (69) and the inequality (13) directly follow from (74) and (75).
If

the inequality $r < r_1$ holds, then in view of (73) – (75),

$$r_2! \prod_{j=1}^{r_1-r-1} (p+r-r_1+j) = (p-1)!, \quad v_p(r_2!(r_1-r-1)!(-1)^{r_1-r}-1) \geq 1,$$

and (72) holds.

Corollary 1. *Let $p \in \mathbb{N}$ is a prime number,*

$$d \in \mathbb{N} - 1, r \in \mathbb{N} - 1, d_1 \in \mathbb{N} - 1, d_2 \in \mathbb{N} - 1, r_1 \in \mathbb{N} - 1, r_2 \in \mathbb{N} - 1,$$

$$\max(r_1, r_2) < p.$$

Then

$$p^{-d}(dp+r)! \in (-1)^d d!r! + p\mathbb{Z},$$

$$\binom{(d_1+d_2)p+r_1+r_2}{d_1p+r_1} \in \binom{d_1+d_2}{d_1} \binom{r_1+r_2}{r_1} + p\mathbb{Z}.$$

Proof. This is direct corollary of the Lemma 7 and Lemma 8. See also Lemma 9 in [54]. ■

Corollary 2. *Let $p \in \mathbb{N} + 2$ is a prime number,*

$$d \in \mathbb{N}, r_1 \in \mathbb{N}, r_1 < p, d_1 \in \mathbb{N} - 1, d_1 < d.$$

Then

$$(76) \quad v_p \left(\binom{dp}{d_1p+r_1} \left(d \binom{d-1}{d_1} \binom{p}{r_1} \right)^{-1} + 1 \right) \geq 1$$

Proof. Since,

$$d \binom{d-1}{d_1} = (d-d_1) \binom{d}{d_1}, \quad v_p \left(\binom{p}{r_1} r_1/p - (-1)^{r_1} \right) \geq 1,$$

the equality (76) directly follows from (72). ■

Corollary 3. *Let $p \in \mathbb{N} + 2$ is a prime number,*

$$d \in \mathbb{N}, r_1 \in \mathbb{N}, r_1 < p, d^\sim \in \mathbb{N} - 1, d^\sim < d.$$

Then

$$\binom{dp}{d_1p+r_1} \in d \binom{d-1}{d^\sim} \binom{p}{r_1} + p^2\mathbb{Z}.$$

Proof. This is a corollary of the Corollary 2. See also Lemma 10 in [54]. ■

Let let p be prime in $(2, +\infty)$, let K be a finite extension of \mathbb{Q} let \mathfrak{p} be a prime ideal in \mathbb{Z}_K and $p \in \mathfrak{p}$, let f be the degree of \mathfrak{p} , let $(p) = \mathfrak{p}^e \mathfrak{b}$, with entire ideal \mathfrak{b} not contained in \mathfrak{p} , let $v_{\mathfrak{p}}$ be additive \mathfrak{p} -valuation, which prolongs v_p ; so, if π is a \mathfrak{p} -prime number, then $v_{\mathfrak{p}}(\pi) = 1/e$. If f is the degree of the ideal \mathfrak{p} then

$$(77) \quad v_{\mathfrak{p}} \left(w^{p^\beta} - w \right) \geq 1,$$

where $\beta \in \mathbb{N}f$, $w \in K$ and

$$v_{\mathfrak{p}}(w) \geq 0.$$

In view of (77), (18), and (16),

$$v_{\mathfrak{p}}(\alpha^*(z; p^\beta l) - \alpha^*(z; l)) > 1/e,$$

if $\beta \in \mathbb{N}f$, $\theta_0(z) \in K$ and $v_{\mathfrak{p}}(\theta_0(z)) \geq 0$. In view of (19),

$$(78) \quad \begin{aligned} \phi^*(z; \nu) &= (-\theta_0(z))^\nu \sum_{k=0}^{\nu\Delta} \alpha_{\nu,k}^*(\theta_0(z))^k \sum_{\tau=1}^{\nu+k} ((\theta_0(z))^{-\tau}/\tau) = \\ &(-\theta_0(z))^\nu \sum_{\tau=1}^{\nu} ((\theta_0(z))^{-\tau} \alpha^*(z; \nu)/\tau + \\ &(-\theta_0(z))^\nu \sum_{k=0}^{\nu\Delta} \alpha_{\nu,k}^*(\theta_0(z))^k \sum_{\tau=1+\nu}^{\nu+k} ((\theta_0(z))^{-\tau}/\tau) = \\ &(-1)^\nu \sum_{\tau=1}^{\nu(\Delta+1)} \frac{1}{\tau} \sum_{k=\max(0, \tau-\nu)}^{\nu\Delta} \alpha_{\nu,k}^*(\theta_0(z))^{\nu-\tau+k}; \end{aligned}$$

therefore, if $\nu = p^\beta l$, $f = 1$, $\beta \in \mathbb{N}f$, $p > l(\Delta + 1)$, $\theta_0(z) \in K$ and $v_{\mathfrak{p}}(\theta_0(z)) \geq 0$, then, according to the Lemma 2,

$$(79) \quad 1 - \beta \leq$$

$$v_{\mathfrak{p}} \left(\phi^*(z; \nu) - \sum_{\substack{\eta \in [1, \Delta+1] \cap \mathbb{Z} \\ k \in [p^\beta(\eta-l), p^\beta l \Delta] \cap \mathbb{Z} \\ k \geq 0, v_{\mathfrak{p}}(k) > 0}} \frac{(-1)^{p^l}}{p^{\beta\eta}} (\theta_0(z))^{p^\beta(l-\eta)+k} \alpha_{\nu,k}^* \right),$$

$$(80) \quad 1/e - \beta \leq$$

$$v_{\mathfrak{p}} \left(\phi^*(z; \nu) - \sum_{\substack{\eta \in [1, \Delta+1] \cap \mathbb{Z} \\ k \in [p^{\beta-1}(\eta-l), p^{\beta-1} l \Delta] \cap \mathbb{Z} \\ k \geq 0}} \frac{(-1)^{p^l} p}{p^{\beta\eta}} (\theta_0(z))^{p^{\beta-1}(l-\eta)+k} \alpha_{\nu/p,k}^* \right).$$

We make the pass (79) \rightarrow (80) β times and obtain the inequality

$$(81) \quad 1/e - \beta \leq$$

$$v_{\mathfrak{p}}(\phi^*(z; p^\beta l) - p^{-\beta} \phi^*(z; l)),$$

where

$$\{l, \beta\} \subset \mathbb{N}, p > l(\Delta + 1), p \in \mathfrak{p}$$

and \mathfrak{p} is ideal of the first degree.

Lemma 9. *If $m \in \mathbb{N} + 1$, $K = \mathbb{Q}[\exp(2\pi i/m)]$,*

$$\alpha^*(z; l_1) \phi^*(z; l_2) \neq 0$$

for some $z \in K \setminus \{0\}$, $l_1 \in \mathbb{N}$, $l_2 \in \mathbb{N}$, then for any $l \in \mathbb{N}$ the sequences

$$(82) \quad \alpha^*(z; \nu), \phi^*(z; \nu),$$

where $\nu \in l + \mathbb{N}$ form a linear independent system over K .

Proof. There exists $d^* \in \mathbb{N}$ such that

$$d^*z \in \mathbb{Z}_K, d^*z\alpha^*(z; l_1) \in \mathbb{Z}_K, d^*z\phi^*(z; l_2) \in \mathbb{Z}_K.$$

Let a prime $p \in \mathbb{N}m + 1$ satisfies to the inequality

$$p > |Nm_{K/\mathbb{Q}}(d^*z\alpha^*(z; \nu))| + |Nm_{K/\mathbb{Q}}(d^*z\phi^*(z; \nu))| + \\ |Nm_{K/\mathbb{Q}}(d^*z)| + |Nm_{K/\mathbb{Q}}(d^*) + (\Delta + 1)(l_1 + l_2).$$

Let \mathfrak{p} is a prime ideal containing p . Then

$$v_{\mathfrak{p}}(\alpha^*(z; l_1)) = v_{\mathfrak{p}}(\phi^*(z; l_2)) = 0,$$

and, in view of (81),

$$v_{\mathfrak{p}}(\phi^*(z; p^\beta l_2)) = -\beta,$$

but

$$v_{\mathfrak{p}}(\alpha^*(z; p^\beta l_1)) = 0.$$

with $\beta \in \mathbb{N}$ ■.

Let $m \in \mathbb{N}$, $k \in \mathbb{Z}$, $2 \leq 2|k| < m$, and let m and k have no common divisor with exception ± 1 . Let further $K_m = \mathbb{Q}[\exp(2\pi i/m)]$ is a cyclotomic field, \mathbb{Z}_{K_m} is the ring of all the integers of the field K_m .

Lemma 10. Let $\Delta \in \{5, 7\}$. In corresponde with (21), (22) and (23), let $z = (1/(2 \cos(k\pi/m)), k\pi/m - \pi)$, where $|k| < m/2$, $(|k|, m) = 1$.

Then for each $l \in \mathbb{N}$ the two sequences (82) form a linear independent system over \mathbb{C} .

Proof. We check the fulfilment of the conditions of the Lemma 9.

Let $\mathfrak{M} = \mathbb{N} \setminus \{1, 2, 6\}$ and $\mathfrak{M}_0 = \{m \in \mathfrak{M}: \Lambda_0(m) = 0\}$. According to the condition of the Lemma, $\theta_0(z) = -1/(1 + \exp(2i\pi/m))$ with $m \in \mathfrak{M}$. If $m \in \mathfrak{M}$ and $\phi(m) > \Delta$, then, in view of (18) and (16), $\alpha^*(z; 1) \neq 0$, because the numbers $(1 + \exp(2i\pi/m))^k$, where $k = 0, \dots, \phi(m) - 1$, form a basis of the field K_m . Let $\Delta = p \in 2\mathbb{N} + 1$, where p is a prime, \mathfrak{p} is a prime ideal containing p , and, as before, let $(p) = \mathfrak{b}p^e$, $1_{K_m} \in \mathfrak{b} + \mathfrak{p}$. Then

$$(83) \quad \binom{2p-1}{p} \binom{p}{p-1} \equiv p \pmod{p^2}, v_{\mathfrak{p}} \left(\binom{p+k}{1+k} \binom{p}{k} \right) = 2,$$

where $k = 1, \dots, p-2$,

$$(84) \quad \binom{p}{1} \binom{p}{0} = p, \binom{2p}{p+1} \binom{p}{p} \equiv 2p \pmod{p^2}.$$

If $m \in \mathfrak{M}$ and $(m, p) = 1$, or, if $m \in \mathfrak{M}_0$, then, according to the Lemma 1,

$$(85) \quad (1 + \exp(2i\pi/m), p) = (1)$$

and, according to the Lemmata 7 and 8,

$$(86) \quad \alpha^*(z; 1)/(p\theta_0(z)) \equiv 1 + (\theta_0(z))^{p-1} - 2(\theta_0(z))^p \equiv \\ 1 + (\exp(2i\pi/m) + 3)/(1 + \exp(2pi\pi/m)) \equiv \\ (\exp(2ip\pi/m) + \exp(2i\pi/m) + 4)/(1 + \exp(2pi\pi/m)) \pmod{p}.$$

If $m = q^\alpha$ with $\alpha \in \mathbb{N}$ and prime q and there exists l in $\{0, \dots, \phi(m) - 1\}$ such that $p \equiv l \pmod{m}$, then

$$(87) \quad \exp(2ip\pi/m) + \exp(2i\pi/m) + 4 \not\equiv 0 \pmod{p}.$$

If $m = 2q^\alpha$ with odd prime q and $\alpha \in \mathbb{N}$, and there exist l in $\{0, \dots, \phi(m/2) - 1\}$ such that $p \equiv 2l \pmod{m/2}$, then (87) holds.

If $p = 5$, then $\{3, 4, 5, 8, 10, 12\} = \{m \in \mathfrak{M}: \phi(\mathfrak{m}) \leq \mathfrak{p}\}$.

If $m = 3, 4, 5, 8, 10$ then, clearly, (87) holds.

If $m = 12$, then $1, \exp(i\pi/2), \exp(2i\pi/3), \exp(i\pi/6)$, form a entire basis of K_{12} , $\exp(5i\pi/6) = \exp(i\pi/2) - \exp(i\pi/6)$, and (87) holds.

If $p = 7$ then $\{3, 4, 5, 7, 8, 9, 10, 12, 14, 18\} = \{m \in \mathfrak{M}: \phi(\mathfrak{m}) \leq \mathfrak{p}\}$.

If $m = 3, 4, 5, 7, 9, 14$, then, clearly, (87) holds.

If $m = 8$, then $\exp(7i\pi/4) = -\exp(3i\pi/4)$ and (87) holds.

If $m = 12$, then $1, \exp(i\pi/2), \exp(2i\pi/3), \exp(i\pi/6)$, form a entire basis of K_{12} , $\exp(7i\pi/6) = -\exp(i\pi/6)$, and (87) holds.

If $m = 18$, then

$$\exp(7i\pi/9) = -\exp(-2i\pi/9) = \exp(4i\pi/9) + \exp(10i\pi/9),$$

and (87) holds.

The coefficient at $(\theta_0(z))^0$ in the expression (19) of $\phi^*(z; \nu)$ is equal to

$$\sum_{k=0}^{\nu\Delta} (-1)^k \alpha_{\nu,k}^* / (\nu + k)$$

and, if $\Delta = p, \nu = 1$, then in view (83) – (84), the value of v_p on this coefficient is equal to 0. Therefore, if $m \in \mathfrak{M}$ and $\phi(m) > p = \Delta$, then $\phi^*(z; 1) \neq 0$.

If $m \in \mathfrak{M} \setminus \mathfrak{M}_0$, and $m \equiv 0 \pmod{p}$ then $m = 2p^\alpha$, where $\alpha \in \mathbb{N}$. According to the Lemma 1, $\mathfrak{p} = (1 + \exp(2i\pi/m))$ is a prime ideal in K_m , and, furthermore, $\mathfrak{p}^{\phi(m)} = (p)$. Let $v_{\mathfrak{p}}$ is the \mathfrak{p} -adic valuation, which prolongs the valuation v_p . Clearly, $v_{\mathfrak{p}}(1 + \exp(2i\pi/m)) = 1/\phi(m)$, $v_{\mathfrak{p}}(\theta_0(z)) = -1/\phi(m)$ In view of (19) with $\nu = 1$, for the summands of the sum

$$\sum_{k=1}^{\nu\Delta} \alpha_{\nu,k}^* (\theta_0(z))^{1+k} \sum_{\tau=2}^{1+k} ((\theta_0(z))^{-\tau}/\tau)$$

we have the inequality

$$v_{\mathfrak{p}}((\theta_0(z))^{\Delta+k-y\alpha_{\nu,k}^*/y}) \geq -(k-1)/\phi(m) + 2 - v_{\mathfrak{p}}(\tau) \geq -(p-3)/\phi(m) + 2,$$

if $k = 1, \dots, p-2$, because in this case $\tau \in [2, p-1]$,

$$v_p((\theta_0(z))^{\Delta+k-y\alpha_{\nu,k}^*/y}) \geq -(k)/\phi(m) + 1 - v_p(\tau) \geq -(p-1)/\phi(m),$$

where $k \in \{q-1, q\}$, and the equality reaches only for $k = \tau = p$; on the other hand, $v_{\mathfrak{p}}(\alpha^*(z; 1)) \geq 1 - (p+1)/\phi(m) \geq -2/(p-1) \geq -2/\phi(m)$. So, if $p \geq 5$, then $v_{\mathfrak{p}}(\phi^*(z; 1)) = -(p-1)/\phi(m)$. If $m \in \mathfrak{M} \setminus \mathfrak{M}_0$, then $m = 2q^\alpha$, with prime q , according to the Lemma 1, $\mathfrak{l} = (1 + \exp(2i\pi/m))$ is a prime ideal in K_m , and $\mathfrak{l}^{\phi(m)} = (q)$. Therefore in this case $v_{\mathfrak{p}}(\theta_0(z)) = 0$ If $m \in \mathfrak{M}_0$, then, according to the Lemma 1, $v_{\mathfrak{p}}(\theta_0(z)) = 0$. According to (19), in both last cases,

$$v_{\mathfrak{p}}(\phi^*(z; 1)) + \alpha_{\nu, p-1}/p + \theta_0(z)\alpha_{\nu, p}/p \geq 1.$$

In view of (83), (84),

$$\begin{aligned} v_{\mathfrak{p}}(\alpha_{\nu, p-1}/p + \theta_0(z)\alpha_{\nu, p}/p) = \\ v_{\mathfrak{p}}(\exp(2i\pi/m) - 1)/(\exp(2i\pi/m) + 1). \end{aligned}$$

If $p = 5$ and $m \in \{3, 4, 5, 7, 8, 9, 10\}$ then, clearly,

$$(88) \quad v_{\mathfrak{p}}(\exp(2i\pi/m) - 1) \leq 1/4.$$

If $p = 5$ and $m = 12$, then $Nm_{K_{12}}(\exp(i\pi/6) - 1) = 3$ and (88) holds.

If $p = 7$, and $m \in \{3, 4, 5, 7, 8, 9, 10, 12, 14, 18\}$, then

$$v_{\mathfrak{p}}(\exp(2i\pi/m) - 1) \leq 1/6.$$

■

Lemma 11. *Let are fulfilled all the conditions of the Lemma 10. Then*

$$(89) \quad \limsup_{\nu \in \mathbb{N}, \nu \rightarrow \infty} \left(|f_0^*(z, \nu)|^{1/\nu} = \rho_{2,1}(z) \Big|_{\theta_0(z) = -1/(1+\exp(2ik\pi/m))} \right) = |h^\sim(\eta_1(1/(2 \cos(k\pi i/m)), k\pi i/m, \delta_0))|,$$

where $h^\sim(\eta)$ is defined in (30).

Proof. According to the Lemma 2, (20) and Lemma 10, $f_0^*(z, \nu)$ is a nonzero solution of the Poincaré type difference equation (32). According to the Perron's theorem and Lemma 5, the equality (89) holds. ■

Let K/\mathbb{Q} be the finite extension of the field \mathbb{Q} ,

$$[K : \mathbb{Q}] = d.$$

Let the field K has r_1 real places and r_2 complex places. Each such place is the monomorphism of the field K in the field \mathbb{R} , if a place is real, or in the field \mathbb{C} , if a place is not real; we will denote these monomorphisms respectively by $\sigma_1, \dots, \sigma_{r_1+r_2}$. Then $d = r_1 + 2r_2$. Let \mathfrak{B} be the fixed integer basis

$$\omega_1, \dots, \omega_d$$

of the field K over \mathbb{Q} . Clearly, K is an algebra over \mathbb{Q} . With extension of the ground field from \mathbb{Q} to \mathbb{R} appears an isomorphism of the algebra $\mathfrak{K} = K \otimes \mathbb{R}$ onto direct sum

$$\underbrace{\mathbb{R} \oplus \dots \oplus \mathbb{R}}_{r_1 \text{ times}} \oplus \underbrace{\mathbb{C} \oplus \dots \oplus \mathbb{C}}_{r_2 \text{ times}}$$

of r_1 copies of the field \mathbb{R} and r_2 copies of the field \mathbb{C} . We identify by means of this isomorphism the algebra \mathfrak{K} with the specified direct sum. We denote below by π_j , where $j = 1, \dots, r_1 + r_2$, the projection of \mathfrak{K} onto its j -th direct summand and also the extension of this projection onto all kinds of matrices which have all the elements in \mathfrak{K} . So, $\pi_j(\mathfrak{K}) = \mathbb{R}$ for $j = 1, \dots, r_1$ and $\pi_j(\mathfrak{K}) = \mathbb{C}$ for $j = r_1 + 1, \dots, r_1 + r_2$. Further by $\mathbf{i}_{\mathfrak{K}}$ we denote the embedding of \mathbb{R} in \mathfrak{K} in diagonal way and also the extension of this embedding onto all kinds of the real matrices. So, \mathbb{R} is imbedded by means of $\mathbf{i}_{\mathfrak{K}}$ in \mathfrak{K} in diagonal way. Each element $Z \in \mathfrak{K}$ has a unique representation in the form:

$$Z = \begin{pmatrix} z_1 \\ \vdots \\ z_{r_1+r_2} \\ \hline z_{r_1+1} \\ \vdots \\ z_{r_1+r_2} \end{pmatrix},$$

with $z_j = \pi_j(Z) \in \mathbb{R}$ for any $j = 1, \dots, r_1$ and with $z_j = \pi_j(Z) \in \mathbb{C}$ for any $j = r_1 + 1, \dots, r_1 + r_2$. Further by $Tr_{\mathfrak{K}}(Z)$ we denote the sum

$$\begin{aligned} \sum_{j=1}^{r_1} z_j + \sum_{j=r_1+1}^{r_1+r_2} 2\Re(z_j) = \\ \sum_{j=1}^{r_1} \pi_j(Z) + \sum_{j=r_1+1}^{r_1+r_2} 2\Re(\pi_j(Z)), \end{aligned}$$

and by $q_{\infty}^{(\mathfrak{K})}(Z)$ we denote the value

$$\begin{aligned} \max(|z_1|, \dots, |z_{r_1+r_2}|) = \\ \max(|\pi_1(Z)|, \dots, |\pi_{r_1+r_2}(Z)|). \end{aligned}$$

Clearly,

$$\begin{aligned} q_{\infty}^{(\mathfrak{K})}(Z_1 Z_2) &\leq q_{\infty}^{(\mathfrak{K})}(Z_1) q_{\infty}^{(\mathfrak{K})}(Z_2), \\ q_{\infty}^{(\mathfrak{K})}(Z_1 + Z_2) &\leq q_{\infty}^{(\mathfrak{K})}(Z_1) + q_{\infty}^{(\mathfrak{K})}(Z_2), \\ q_{\infty}^{(\mathfrak{K})}(\mathbf{i}_{\mathfrak{K}}(\lambda)Z) &= |\lambda| q_{\infty}^{(\mathfrak{K})}(Z) \end{aligned}$$

for any $Z_1 \in \mathfrak{K}$, $Z_2 \in \mathfrak{K}$, $Z \in \mathfrak{K}$ and $\lambda \in \mathbb{R}$. The natural extension of the norm $q_{\infty}^{(\mathfrak{K})}$ on the set of all the matrices, which have all the elements in \mathfrak{K} (i.e. the maximum of the norm $q_{\infty}^{(\mathfrak{K})}$ of all the elements of the matrix) also will be denoted by $q_{\infty}^{(\mathfrak{K})}$. If

$$Z = \begin{pmatrix} z_1 \\ \vdots \\ z_d \end{pmatrix} \in K,$$

then

$$z_j = \sigma_j(Z),$$

where $j = 1, \dots, r_1 + r_2$,

$$z_{r_1+r_2+j} = \overline{\sigma_{r_1+j}(Z)},$$

where $j = 1, \dots, r_2$. In particular,

$$\omega_k = \begin{pmatrix} \sigma_1(\omega_k) \\ \vdots \\ \sigma_{r_1+r_2}(\omega_k) \\ \hline \sigma_{r_1+1}(\omega_k) \\ \vdots \\ \sigma_{r_1+r_2}(\omega_k) \end{pmatrix},$$

As usually, the ring of all the integer elements of the field K will be denoted by \mathbb{Z}_K . The ring \mathbb{Z}_K is embedded in the ring \mathfrak{K} as discrete lattice. Moreover, if $Z \in \mathbb{Z}_K \setminus \{0\}$, then

$$\left(\prod_{i=1}^{r_1} |\sigma_j(Z)| \right) \prod_{i=1}^{r_2} |\sigma_{r_1+i}(Z)|^2 = |Nm_{K/\mathbb{Q}}(Z)| \in \mathbb{N}$$

and therefore $q_\infty^{(\mathfrak{K})}(Z) \geq 1$. for any $Z \in \mathbb{Z}_K \setminus \{0\}$. The elements of \mathbb{Z}_K we name below by K -integers. For each $Z \in \mathfrak{K}$ let

$$\|Z\|_K = \inf_{W \in \mathbb{Z}_K} \{q_\infty^{(\mathfrak{K})}(Z - W)\}.$$

Let $\{m, n\} \subset \mathbb{N}$,

$$a_{i,k} \in \mathfrak{K}$$

for $i = 1, \dots, m$, $k = 1, \dots, n$,

$$\alpha_j^\wedge(\nu) \in \mathbb{Z}_K,$$

where $j = 1, \dots, m + n$ and $\nu \in \mathbb{N}$. Let there are $\gamma_0, r_1^\wedge \geq 1, \dots, r_m^\wedge \geq 1$ such that

$$q_\infty^{(\mathfrak{K})}(\alpha_i(\nu)) < \gamma_0 (r_i^\wedge)^\nu$$

where $i = 1, \dots, m$ and $\nu \in \mathbb{N}$. Let

$$y_k(\nu) = -\alpha_{m+k}^\wedge(\nu) + \sum_{i=1}^m a_{i,k} \alpha_i^\wedge(\nu)$$

where $k = 1, \dots, n$ and $\nu \in \mathbb{N}$. If $X = \begin{pmatrix} Z_1 \\ \vdots \\ Z_n \end{pmatrix} \in \mathfrak{K}^n$, then let

$$y^\wedge(X) = y^\wedge(X, \nu) = \sum_{k=1}^n y_k^\wedge(\nu) Z_k$$

for $\nu \in \mathbb{N}$, let

$$\phi_i(X) = \sum_{k=1}^n a_{i,k} Z_k$$

for $i = 1, \dots, m$, and let

$$\alpha_0^\wedge(X, \nu) = \sum_{k=1}^n \alpha_{m+k}^\wedge(\nu) Z_k$$

for $\nu \in \mathbb{N}$. Clearly,

$$y^\wedge(X, \nu) = -\alpha_0^\wedge(X, \nu) + \sum_{i=1}^m \alpha_i^\wedge(\nu) \phi_i(X)$$

for $X \in \mathfrak{K}^n$ and $\nu \in \mathbb{N}$,

$$\alpha_0^\wedge(X, \nu) \in \mathbb{Z}_{\mathbb{K}}$$

for $X \in (\mathbb{Z}_K)^n$ and $\nu \in \mathbb{N}$.

Lemma 12. Let $\{l, n\} \subset \mathbb{N}$, $\gamma_1 > 0$, $\gamma_2 > \frac{1}{2}$, $R_1 \geq R_2 > 1$,

$$\alpha_i = (\log(r_i^\wedge R_1 / R_2)) / \log(R_2),$$

where $i = 1, \dots, m$, let $X \in (\mathbb{Z}_K)^n \setminus \{(0)\}$,

$$\gamma_3 = \gamma_1 (R_1)^{(-\log(2\gamma_2 R_2)) / \log(R_2)}, \gamma_4 = \gamma_3 \left(\sum_{i=1}^m \gamma_0 (r_i^\wedge)^{(\log(2\gamma_2)) / \log(R_2) + l} \right)^{-1}$$

and let for each $\nu \in \mathbb{N} - 1$ hold the inequalities

$$\gamma_1 (R_1)^{-\nu} q_\infty^{(\mathfrak{K})}(X) \leq \sup\{q_\infty^{(\mathfrak{K})}(y^\wedge(X, \kappa)) : \kappa = \nu, \dots, \nu + l - 1\},$$

$$q_\infty^{(\mathfrak{K})}(y^\wedge(X, \nu)) \leq \gamma_2 (R_2)^{-\nu} q_\infty^{(\mathfrak{K})}(X)$$

Then

$$\sup\{\|\phi_i(X)\|_K (q_\infty^{(\mathfrak{K})}(X))^{\alpha_i} : i = 1, \dots, m\} \geq \gamma_4.$$

Proof. Proof may be found in [56], Theorem 2.3.1. ■

Corollary. Let $a \in \mathfrak{K}$,

$$(90) \quad \alpha_1^\wedge(\nu) \in \mathbb{Z}_K, \alpha_2^\wedge(\nu) \in \mathbb{Z}_K, y(\nu) = -\alpha_2^\wedge(\nu) + a\alpha_1^\wedge(\nu)$$

where $\nu \in \mathbb{N}$. Let there are $\gamma_0, r_1^\wedge \geq 1$ such that

$$q_\infty^{(\mathfrak{K})}(\alpha_1(\nu)) < \gamma_0 (r_1^\wedge)^\nu,$$

where $\nu \in \mathbb{N}$. Let $l \in \mathbb{N}$, $\gamma_1 > 0$, $\gamma_2 > \frac{1}{2}$, $R_1 \geq R_2 > 1$,

$$\alpha_1 = (\log(r_1^\wedge R_1 / R_2)) / \log(R_2), \gamma_3 = \gamma_1 (R_1)^{(-\log(2\gamma_2 R_2)) / \log(R_2)},$$

$$\gamma_4 = \gamma_3 (\gamma_0 (r_1^\wedge)^{(\log(2\gamma_2)) / \log(R_2) + l})^{-1},$$

$X \in \mathbb{Z}_K$ and let for each $\nu \in \mathbb{N} - 1$ hold the inequalities

$$\gamma_1 (R_1)^{-\nu} q_\infty^{(\mathfrak{K})}(X) \leq \sup\{q_\infty^{(\mathfrak{K})}(y_1(\kappa)X) : \kappa = \nu, \dots, \nu + l - 1\},$$

$$q_{\infty}^{(\mathfrak{R})}(y(\nu)X) \leq \gamma_2(R_2)^{-\nu} q_{\infty}^{(\mathfrak{R})}(X)$$

Then

$$(91) \quad \|aX\|_K(q_{\infty}^{(\mathfrak{R})}(X))^{\alpha} \geq \gamma_4.$$

Proof. This Corrolary is the Lemma 12 for $m = n = 1$. ■

Let $B \in \mathbb{N}$, $D^*(B) = \inf\{q \in \mathbb{N} : d/\kappa \in \mathbb{N}, \kappa \in \mathbb{N}, \kappa \leq B\}$. It is known that

$$D^*(B) = \exp(B + O(B/\log(B))).$$

Let $d_0^*(\Delta, \nu) = D^*(\nu(\Delta + 1))$. Then

$$(92) \quad d_0^*(\Delta, \nu) = \exp(\nu(\Delta + 1) + O(\nu/\log(\nu))),$$

when $\nu \rightarrow \infty$.

Probably G.V. Chudnovsky was the first man, who discovered, that the numbers (16) have a great common divisor; Hata ([17]) in details studied this effect. Therefore I name the mentioned common divisor by Chudnovsky-Hata's multiplier and denote it by $d_1^*(\Delta, \nu)$. According to the Hata's results,

$$(93) \quad \log(d_1^*(\Delta, \nu)) = (1 + o(1))\nu \times$$

$$\sum_{\mu=0}^1 \left(\frac{\Delta + (-1)^{\mu}}{2} \log \left(\frac{\Delta}{\Delta + (-1)^{\mu}} \right) + (-1)^{\mu} \frac{\pi}{2} \sum_{\kappa=1}^{\lfloor \frac{\Delta + (-1)^{\mu}}{2} \rfloor} \cot \left(\frac{\pi \kappa}{\Delta + (-1)^{\mu}} \right) \right).$$

In view of (92),

$$(94) \quad d_0^*(5, \nu) = \exp(6\nu(\Delta + 1) + O(\nu/\log(\nu))), d_0^*(7, \nu) = \exp(8\nu(8) + O(\nu/\log(\nu))).$$

In view of (94)

$$(95) \quad \log(d_1^*(5, \nu)) = (1 + o(1))\nu \times (-3 \log(1.2) + 2 \log(0.8) + (\pi/2)(\cot(\pi/6) + \cot(\pi/3) + \cot(\pi/4))) = (1 + o(1))\nu \times 1.956124\dots,$$

$$(96) \quad \log(d_1^*(7, \nu)) = (1 + o(1))\nu \times (4 \log(7/8) + 3 \log(7/6)) + (1 + o(1))(\pi/2)\nu \times (-\cot(\pi/6) - \cot(\pi/3) + \cot(\pi/8) + \cot(3\pi/8) + \cot(\pi/4)) = (1 + o(1))\nu(4 \log(7/8) + 3 \log(7/6) + \pi(-2/\sqrt{3} + 2/\sqrt{2} + 1/2)) = (1 + o(1))\nu \times 2.314407\dots,$$

when $\nu \rightarrow \infty$.

In view of (18) and (19),

$$\alpha^*(z; \nu)d_0^*(\nu)/d_1^*(\nu) \in \mathbb{Z}[z],$$

$$\phi^*(z; \nu)d_0^*(\nu)/d_1^*(\nu) \in \mathbb{Z}[z].$$

Let

$$(97) \quad U_{\Delta}(m, \nu) = d_0^*(\nu)/d_1^*(\nu), \quad \Lambda_0(m) = 0,$$

if $m \neq 2p^\alpha$, where p run over the all the prime numbers and α run over \mathbb{N} and let

$$(98) \quad U_{\Delta}(m, \nu) = \frac{d_0^*(\nu)}{d_1^*(\nu)} p^{[(\Delta+1)\nu/\phi(m)]+1}, \quad \Lambda_0(m) = \Lambda(m/2),$$

if $m = 2p^\alpha$, where p is a prime number and $\alpha \in \mathbb{N}$. In view of the (18), (19) and Lemma 1,

$$(99) \quad \alpha^*(z; \nu) \Big|_{z=\left(\frac{1}{2\cos\left(\frac{k\pi i}{m}\right)}, \frac{k\pi i}{m} - \pi\right)} U_{\Delta}(m, \nu) \in \mathbb{Z}_{\mathbb{Q}[\exp(2i\pi/m)]},$$

$$(100) \quad \phi^*(z; \nu) \Big|_{z=\left(\frac{1}{2\cos\left(\frac{k\pi i}{m}\right)}, \frac{k\pi i}{m} - \pi\right)} U_{\Delta}(m, \nu) \in \mathbb{Z}_{\mathbb{Q}[\exp(2i\pi/m)]},$$

where $(k, m) = 1$. In view of (98), (97), (93), (92), (2) and (3)

$$(101) \quad \frac{d_0^*(\nu)}{d_1^*(\nu)} = \nu(1 + o(1))V_{\Delta}^* \log(U_{\Delta}(m, \nu)) = \nu(1 + o(1))V_{\Delta}(m),$$

when $\nu \rightarrow \infty$.

The polynomial (28) take the form

$$D^{\wedge}(z, \eta) = (\eta + 1) \left(\eta + \frac{\Delta - 1}{\Delta + 1} \right) + \frac{2\Delta \exp(i\psi)\eta}{(\Delta + 1) \cos(\psi)} =$$

$$((\Delta + 1)\eta^2 + 2\Delta(2 + iT)\eta + (\Delta - 1))/(\Delta + 1),$$

where $\psi \in (-\pi/2, \pi/2)$ and $T = \tan(\psi)$; its roots are equal to

$$(102) \quad -(2\Delta + \Delta iT + R)/(\Delta + 1),$$

where $R^2 = \Delta^2(3 - T^2) + 1 + 4\Delta^2 iT$. In view of (1), Then

$$R \in \{\pm (w_{\Delta}(T) + i2\Delta^2 iT/w_{\Delta}(T))\}.$$

In view of (102) and (46),

$$\eta_j^{\wedge}(r, \psi, \delta_0) =$$

$$-\frac{2\Delta + \Delta iT + (-1)^j (w_{\Delta}(T) + i2\Delta^2 iT/w_{\Delta}(T))}{\Delta + 1} =$$

$$-\frac{2\Delta + (-1)^j w_{\Delta}(T) + iT\Delta (1 + (-1)^j 2\Delta/w_{\Delta}(T))}{\Delta + 1},$$

where $j = 0, 1$,

$$|\eta_j^{\wedge}(r, \psi, \delta_0) + k|^2 =$$

$$\frac{(2\Delta + (-1)^j w_\Delta(T) - k(\Delta + 1))^2 + T^2 \Delta^2 (1 + (-1)^j 2\Delta/w_\Delta(T))^2}{(\Delta + 1)^2},$$

where $j = 0, 1$; $k = 0, 1, -1$. Therefore, in view of (30) and (4)

$$(103) \quad \ln |h^\sim(\eta_j^\wedge(r, \psi, \delta_0))| =$$

$$(\eta_j(r, \psi, \delta_0) - 1)(1 - \delta_0)^{-d_1}(\eta_j(r, \psi, \delta_0) + 1)2^{-2}\eta_j(r, \psi, \delta_0)^{d_1} =$$

$$-\log(4(\Delta + 1)^{\Delta+1}(1 - 1/\Delta)^{\Delta-1}) +$$

$$\frac{1}{2} \log \left((2\Delta + (-1)^j w_\Delta(T) + (\Delta + 1))^2 + T^2 \Delta^2 \left(1 + \frac{(-1)^j 2\Delta}{w_\Delta(T)} \right)^2 \right) +$$

$$\frac{1}{2} \log \left((2\Delta + (-1)^j w_\Delta(T) - (\Delta + 1))^2 + T^2 \Delta^2 \left(1 + \frac{(-1)^j 2\Delta}{w_\Delta(T)} \right)^2 \right) +$$

$$\frac{(\Delta - 1)}{2} \log \left((2\Delta + (-1)^j w_\Delta(T))^2 + T^2 \Delta^2 \left(1 + \frac{(-1)^j 2\Delta}{w_\Delta(T)} \right)^2 \right) =$$

$$l_\Delta(j, T),$$

where $j = 0, 1$. Clearly,

$$w_\Delta(0) = \sqrt{3\Delta^2 + 1},$$

$$\eta_j^\wedge(1/2, 0, \delta_0) = -\frac{2\Delta + (-1)^j \sqrt{3\Delta^2 + 1}}{\Delta + 1},$$

where $j = 0, 1$,

$$|\eta_j^\wedge(1/2, 0, \delta_0) + k| = \left| \frac{2\Delta + (-1)^j \sqrt{3\Delta^2 + 1} - k(\Delta + 1)}{\Delta + 1} \right|,$$

where $j = 0, 1$; $k = 0, 1, -1$. Therefore

$$(104) \quad l_\Delta(\epsilon, 0) = (\log |h^\sim(\eta_\epsilon^\wedge(1/2, 0, \delta_0))|) =$$

$$\log(|(\eta_\epsilon(1/2, 0, \delta_0) - 1)(1 - \delta_0)^{-d_1}(\eta_\epsilon(1/2, 0, \delta_0) + 1)2^{-2}\eta_\epsilon(1/2, 0, \delta_0)^{d_1}|) =$$

$$-\log(4(\Delta + 1)^{\Delta+1}(1 - 1/\Delta)^{\Delta-1}) +$$

$$\log(|2\Delta + (-1)^\epsilon \sqrt{3\Delta^2 + 1} - (\Delta + 1)|) +$$

$$\log(|2\Delta + (-1)^\epsilon \sqrt{3\Delta^2 + 1} + (\Delta + 1)|) +$$

$$(\Delta - 1) \log(|2\Delta + (-1)^\epsilon \sqrt{3\Delta^2 + 1}|).$$

Consequently

$$l_5(1, 0) = -\log(4) - 6 \log 6 - 4 \log(0.8) +$$

$$\log(\sqrt{76} - 4) + \log(16 - \sqrt{76}) + 4 \log(10 - \sqrt{76})$$

I made computations below "by hands" using calculator of the firm "CASIO."

$$\log 4 = 1,386294361\dots; 6 \log(6) = 10,7505682\dots;$$

$$\begin{aligned}
4 \log(0.8) &= -0,892574205\dots; \\
\sqrt{76} &= 8,717797887\dots; \sqrt{76} - 4 = 4,717797887\dots; \\
16 - \sqrt{76} &= 7,282202113\dots; 10 - \sqrt{76} = 1,282202113\dots; \\
\log(\sqrt{76} - 4) &= 1.551342141\dots; \log(16 - \sqrt{76}) = 1.985433305\dots; \\
\log(10 - \sqrt{76}) &= 0.248579\dots; 4 \log(10 - \sqrt{76}) = 0,994316001\dots;
\end{aligned}$$

$$(105) \quad l_5(1, 0) = -6.713196909\dots;$$

$$\begin{aligned}
l_7(1, 0) &= -\log(4) - 8 \log(8) - 6 \log(6) + 6 \log(7) + \\
&\log(\sqrt{148} - 6) + \log(22 - \sqrt{148}) + 6 \log(14 - \sqrt{148}); \\
8 \log 8 &= 16,63553233\dots; 6 \log 6 = 10,75055682\dots; 6 \log 7 = 11,67546089\dots; \\
\sqrt{148} &= 12,16552506\dots; \sqrt{148} - 6 = 6,16552506\dots \\
22 - \sqrt{148} &= 9,83474939\dots; 14 - \sqrt{148} = 1,83474939\dots; \\
\log(\sqrt{148} - 6) &= 1,818973301; \log(22 - \sqrt{148}) = 2,285894063\dots; \\
\log(14 - \sqrt{148}) &= 0,606758304\dots; 6 \log(14 - \sqrt{148}) = 3,640549824\dots;
\end{aligned}$$

$$(106) \quad l_7(1, 0) = -9,35150543\dots$$

In view of (2), (92), (93), (95), (96) and (101),

$$(107) \quad V_5^* = 6 - 1.956124\dots = 4,04387\dots; V_7^* = 8 - 2.314407 = 5,685593.$$

In view (105) – (107),

$$(108) \quad -V_5^* - l_5(1, 0) > 0, -V_7^* - l_7(1, 0) > 0.$$

So, the key inequalities (108) are checked "by hands". In view of (103), (108) and Lemma 3,

$$-V_5^* - l_5(1, \tan(\pi/m)) > 0, -V_7^* - l_7(1, \tan(\pi/m)) > 0,$$

where $m > 2$. Since $(\log(p))/(p^{\alpha-1}(p-1))$ decreases together with increasing of $p \in (3, +\infty)$ with fixed $\alpha \geq 1$, or increasing of $\alpha \in (1, +\infty)$ with fixed $p \geq 2$ (or, of course, increasing both $\alpha \in (1, +\infty)$ and $p \in (3, +\infty)$), and

$$\lim_{p \rightarrow \infty} ((\log(p))/(p^{\alpha-1}(p-1))) = 0,$$

where $\alpha \geq 1$,

$$\lim_{\alpha \rightarrow \infty} ((\log(p))/(p^{\alpha-1}(p-1))) = 0,$$

where $p \geq 2$, it follows that the inequality (7) holds for all the sufficient big integers m . Computations on computer of class "Pentium" show that the inequality (7) holds for $m = 3$, $m = 4$, $m = 5$ and $m = 2 \times 5$; therefore

inequality (7) holds for all the $m > 2 \times 3$. Let $\varepsilon_0 = h_\Delta(m)/2$, with $h_\Delta(m)$ defined in (6). In view of (7), $\varepsilon_0 > 0$. We take now $K = K_m = \mathbb{Q}[\exp(2\pi i/m)]$. Let further $\{\sigma_1, \dots, \sigma_{\phi(m)}\} = \text{Gal}(K/\mathbb{Q})$. For each $j = 1, \dots, \phi(m)$ there exists $k_j \in (-m/2, m/2) \cap \mathbb{Z}$ such that

$$(|k_j|, m) = 1, \sigma_j \left(\exp \left(\frac{2\pi i}{m} \right) \right) = \exp \left(\frac{2\pi i k_j}{m} \right).$$

Let a be the element of \mathfrak{K} , such that

$$\pi_j(a) = \log(2 + \sigma_j(\exp(2\pi i/m))) = \log(2 + \exp(2\pi i k_j/m)),$$

where $j = 1, \dots, \phi(m)$; we suppose that $k_1 = 1$. In view of (99) and (100), let $\alpha_1^\vee(\nu)$, $\alpha_1^\wedge(\nu)$, $\alpha_2^\vee(\nu)$, $\alpha_2^\wedge(\nu)$, are elements in \mathfrak{K} such that

$$\pi_j(\alpha_1^\vee(\nu)) = \alpha^*(z; \nu) \Big|_{z = \left(\frac{1}{2 \cos(\frac{k_j \pi i}{m})}, \frac{k_j \pi i}{m} - \pi \right)},$$

$$\pi_j(\alpha_2^\vee(\nu)) = \phi^*(z; \nu) \Big|_{z = \left(\frac{1}{2 \cos(\frac{k_j \pi i}{m})}, \frac{k_j \pi i}{m} - \pi \right)},$$

$$(109) \quad \pi_j(\alpha_1^\wedge(\nu)) = \alpha^*(z; \nu) \Big|_{z = \left(\frac{1}{2 \cos(\frac{k_j \pi i}{m})}, \frac{k_j \pi i}{m} - \pi \right)} U_\Delta(m, \nu),$$

$$(110) \quad \pi_j(\alpha_2^\wedge(\nu)) = \phi^*(z; \nu) \Big|_{z = \left(\frac{1}{2 \cos(\frac{k_j \pi i}{m})}, \frac{k_j \pi i}{m} - \pi \right)} U_\Delta(m, \nu),$$

where $j = 1, \dots, \phi(m)$. Then $\alpha_k^\wedge(\nu) \in \mathbb{Z}_K$ for $k = 1, 2$.

$$(111) \quad y^\vee(\nu) = -\alpha_2^\vee(\nu) + a\alpha_1^\vee(\nu),$$

and let $y(\nu)$ is defined by means the equality (90). According to the Corrolary of the Lemma 4, to the Theorem 4 in [58] (or Theorem 7 in [66]), to the Lemma 8, to (103), there exist $m_1^* \in \mathbb{N}$ having the following property:

for any $\varepsilon \in (0, \varepsilon_0)$ there exist $\gamma_0(\varepsilon) > 0$, $\gamma_1(\varepsilon) > 0$, and $\gamma_2(\varepsilon) > 0$ such that

$$(112) \quad |\pi_j(\alpha_k^\vee(\nu))| \leq \gamma_0(\varepsilon) \exp((l_\Delta(\tan((k_j \pi i)/m), 0) + \varepsilon/3)\nu),$$

where $k = 1, 2$, $j = 1, \dots, \phi(m)$ and $\nu \in \mathbb{N} - 1 + m_1^*$,

$$(113) \quad \gamma_1(\varepsilon) \exp((l_\Delta(\tan((k_j \pi i)/m), 1) - \varepsilon/3)\nu) \leq \max(|\pi_j(y^\vee(\nu))|, |\pi_j(y^\vee(\nu + 1))|) \leq \gamma_2(\varepsilon) \exp((l_\Delta(\tan((k_j \pi i)/m), 1) + \varepsilon/3)\nu),$$

where $j = 1, \dots, \phi(m)$ and $\nu \in \mathbb{N} - 1 + m_1^*$.

Let $\omega_1(m) = (m-1)/2$, if m is odd, $\omega_1(m) = m/2 - 2$, if $m \equiv 2 \pmod{4}$ and $\omega(m) = m/2 - 1$, if $m \equiv 0 \pmod{4}$. Then

$$\omega_1(m) = \sup\{k \in \mathbb{N}: k_j < m/2, (k, m) = 1\}.$$

According to the Lemma 3 and (103),

$$(114) \quad l_\Delta(\tan((k_j\pi i)/m), 0) \leq l_\Delta(\tan((\omega_1(m)\pi i)/m), 0),$$

$$(115) \quad l_\Delta(\tan((\omega_1(m)\pi i)/m), 1) \leq l_\Delta(\tan((k_j\pi i)/m), 1) \leq l_\Delta(\tan((\pi i)/m), 1)$$

where $j = 1, \dots, \phi(m)$. In view of (112) – (115),

$$(116) \quad |\pi_j(\alpha_k^\vee(\nu))| \leq \gamma_0(\varepsilon) \exp((l_\Delta(\tan((\omega_1(\nu)\pi i)/m), 0) + \varepsilon/3)\nu),$$

where $k = 1, 2, j = 1, \dots, \phi(m)$ and $\nu \in \mathbb{N} - 1 + m_1^*$,

$$(117) \quad \gamma_1(\varepsilon) \exp((l_\Delta(\tan((\omega_1(m)\pi i)/m), 1) - \varepsilon/3)\nu) \leq \max(|\pi_j(y^\vee(\nu))|, |\pi_j(y^\vee(\nu+1))|) \leq \gamma_2(\varepsilon) \exp((l_\Delta(\tan((\pi i)/m), 1) + \varepsilon/3)\nu),$$

where $j = 1, \dots, \phi(m)$ and $\nu \in \mathbb{N} - 1 + m_1^*$. In view of (101), there exists $m_2^* \in \mathbb{N} - 1 + m_1^*$, such that

$$(118) \quad \exp(V_\Delta(m) - \varepsilon/3)\nu \leq U_\Delta(m, \nu) \leq \exp(V_\Delta(m) - \varepsilon/3)\nu$$

where $\nu \in \mathbb{N} - 1 + m_2^*$.

In view of (115) – (118), (109) – (111), (6), (5),

$$(119) \quad |\pi_j(\alpha_k(\nu))| \leq \gamma_0(\varepsilon) \exp((g_{\Delta,0}(m) + 2\varepsilon/3)\nu),$$

where $k = 1, 2, j = 1, \dots, \phi(m)$ and $\nu \in \mathbb{N} - 1 + m_2^*$,

$$(120) \quad \gamma_1(\varepsilon) \exp((-g_{\Delta,1}(m) - 2\varepsilon/3)\nu) \leq \max(|\pi_j(y^\vee(\nu))|, |\pi_j(y^\vee(\nu+1))|) \leq \gamma_2(\varepsilon) \exp((-h_\Delta(m) + 2\varepsilon/3)\nu),$$

where $j = 1, \dots, \phi(m)$ and $\nu \in \mathbb{N} - 1 + m_2^*$.

Let $X \in \mathbb{Z}_{K_m} \setminus \{0\}$. Then, in view of (119) and (120),

$$(121) \quad |\pi_j(X\alpha_k(\nu))| \leq \gamma_0(\varepsilon) \exp((g_{\Delta,0}(m) + 2\varepsilon/3)\nu) |\pi_j(X)| \leq \gamma_0(\varepsilon) \exp((g_{\Delta,0}(m) + 2\varepsilon/3)\nu) q_\infty^{(\mathbb{R})}(X),$$

where $k = 1, 2, j = 1, \dots, \phi(m)$ and $\nu \in \mathbb{N} - 1 + m_2^*$,

$$(122) \quad \gamma_1(\varepsilon) \exp((-g_{\Delta,1}(m) - 2\varepsilon/3)\nu) |\pi_j(X)| \leq$$

$$\begin{aligned} & \max(|\pi_j(Xy^\vee(\nu))|, |\pi_j(Xy^\vee(\nu+1))|) \leq \\ & \max(q_\infty^{(\mathfrak{R})}(Xy^\vee(\nu)), q_\infty^{(\mathfrak{R})}(Xy^\vee(\nu+1))), \end{aligned}$$

where $j = 1, \dots, \phi(m)$ and $\nu \in \mathbb{N} - 1 + m_2^*$,

$$(123) \quad \begin{aligned} & \max(|\pi_j(Xy^\vee(\nu))|, |\pi_j(Xy^\vee(\nu+1))|) \leq \\ & \gamma_2(\varepsilon) \exp((-h_\Delta(m) + 2\varepsilon/3)\nu) |\pi_j(X)| \leq \\ & \gamma_2(\varepsilon) \exp((-h_\Delta(m) + 2\varepsilon/3)\nu) q_\infty^{(\mathfrak{R})}(X), \end{aligned}$$

where $j = 1, \dots, \phi(m)$ and $\nu \in \mathbb{N} - 1 + m_2^*$.

In view of (121)

$$(124) \quad \begin{aligned} & q_\infty^{(\mathfrak{R})}(X\alpha_k(\nu)) \leq \\ & \gamma_0(\varepsilon) \exp((g_{\Delta,0}(m) + 2\varepsilon/3)\nu) q_\infty^{(\mathfrak{R})}(X), \end{aligned}$$

where $k = 1, 2$, and $\nu \in \mathbb{N} - 1 + m_2^*$. In view of (123),

$$(125) \quad \begin{aligned} & \max(q_\infty^{(\mathfrak{R})}(Xy^\vee(\nu)), q_\infty^{(\mathfrak{R})}(Xy^\vee(\nu+1))) = \\ & \sup(\{|\pi_j(Xy^\vee(\nu+\epsilon))|, : \epsilon \in \{0, 1\}, j = 1, \dots, \phi(m)\}) \leq \\ & \gamma_2(\varepsilon) \exp((-h_\Delta(m) + 2\varepsilon/3)\nu) q_\infty^{(\mathfrak{R})}(X), \end{aligned}$$

where $\nu \in \mathbb{N} - 1 + m_2^*$.

Taking in account (124), (125) and (122), we see that all the conditions of the Corollary of the Lemma 12 are fulfilled for

$$\begin{aligned} & \varepsilon \in (0, \varepsilon_0), \gamma_0(\varepsilon), \gamma_1(\varepsilon), \gamma_2(\varepsilon), y = y(\nu), \alpha_1(\nu), \alpha_2(\nu), \\ & r_1 = r_1(\varepsilon) = \exp(g_{\Delta,0}(m) + 2\varepsilon/3), \\ & R_1 = R_1(\varepsilon) = \exp(g_{\Delta,1}(m) + 2\varepsilon/3), \\ & R_2 = R_2(\varepsilon) \exp(h_\Delta(m) - 2\varepsilon/3), \end{aligned}$$

and this proves the part of our Theorem connected with the inequality (8).

Let again $X \in \mathbb{Z}_{K_m} \setminus \{0\}$ and let

$$q_{min}^{(\mathfrak{R})}(X) = \inf(|\{\pi_j(X)\}| : j = 1, \dots, \phi(m))$$

Clearly, $q_{min}^{(\mathfrak{R})}(X) > 0$ According to the Theorem 4 in [58], or to the Theorem 7 in [66], there exist $m_1^* \in \mathbb{N}$ having the following property: for any $\varepsilon \in (0, \varepsilon_0)$ there exist $\gamma_0^*(X, \varepsilon) > 0$, $\gamma_1^*(X, \varepsilon) > 0$, and $\gamma_2^*(X, \varepsilon) > 0$ such that

$$|\pi_j(\alpha_k^\vee(\nu))| \leq \gamma_0^*(\varepsilon) \exp((l_\Delta(\tan((\omega_m \pi i)/m), 0) + \varepsilon/3)\nu),$$

where $k = 1, 2, j = 1, \dots, \phi(m)$ and $\nu \in \mathbb{N} - 1 + m_1^*$,

$$\begin{aligned} & \gamma_1^*(X\varepsilon) \exp((l_\Delta(\tan((\pi i)/m), 1) - \varepsilon/3)\nu) \leq \\ & \max(|\pi_j(y^\vee(\nu))|, |\pi_j(y^\vee(\nu+1))|) \leq \\ & \gamma_2(\varepsilon) \exp((l_\Delta(\tan((\pi i)/m), 1) + \varepsilon/3)\nu), \end{aligned}$$

where $j = 1, \dots, \phi(m)$ and $\nu \in \mathbb{N} - 1 + m_1^*$. Repeating the previous considerations, we see that all the conditions of the Corollary of the Lemma 12 are fulfilled for $\varepsilon \in (0, \varepsilon_0)$,

$$\gamma_0 = \gamma_0^*(X, \varepsilon), \gamma_1 = \gamma_1^*(X, \varepsilon), \gamma_2 = \gamma_2^*(X, \varepsilon),$$

$$y = y(\nu), \alpha_1(\nu), \alpha_2(\nu), r_1 = r_1(\varepsilon) = \exp(g_{\Delta,0}(m) + 2\varepsilon/3),$$

and

$$R_1 = R_2 = R_2(\varepsilon) \exp(h_{\Delta}(m) - 2\varepsilon/3),$$

and this proves the part of our Theorem connected with the inequality (9).

■

Below are values of β and α computed for $\Delta \in \{5, 7\}$ and some $m \in \mathbb{N}$.

$$\begin{aligned} (m; \Delta; \beta; \alpha) &= (3; 5; 3, 111228\dots; 3, 111228\dots), \\ (m; \Delta; \beta; \alpha) &= (3; 7; 3, 073525\dots; 3, 073525\dots), \\ (m; \Delta; \beta; \alpha) &= (4; 5; 11, 458947\dots; 11, 458947\dots), \\ (m; \Delta; \beta; \alpha) &= (4; 7; 10, 551730\dots; 10, 551730\dots), \\ (m; \Delta; \beta; \alpha) &= (5; 5; 4, 826751\dots; 5, 607961\dots), \\ (m; \Delta; \beta; \alpha) &= (5; 7; 4, 837858\dots; 5, 684622\dots), \\ (m; \Delta; \beta; \alpha) &= (7; 5; 5, 701485\dots; 6, 977258\dots), \\ (m; \Delta; \beta; \alpha) &= (7; 7; 5, 724804\dots; 7, 114963\dots), \\ (m; \Delta; \beta; \alpha) &= (8; 5; 8, 337857\dots; 9, 436901\dots), \\ (m; \Delta; \beta; \alpha) &= (8; 7; 8, 253047\dots; 9, 433260\dots), \\ (m; \Delta; \beta; \alpha) &= (9; 5; 6, 312056\dots; 7, 960502\dots), \\ (m; \Delta; \beta; \alpha) &= (9; 7; 6, 335274\dots; 8, 134962\dots), \\ (m; \Delta; \beta; \alpha) &= (10; 5; 43, 546644\dots; 46, 230614\dots), \\ (m; \Delta; \beta; \alpha) &= (10; 7; 35, 648681\dots; 38, 043440\dots), \\ (m; \Delta; \beta; \alpha) &= (11; 5; 6, 786990\dots; 8, 735234\dots), \\ (m; \Delta; \beta; \alpha) &= (11; 7; 6, 806087\dots; 8, 934922\dots), \\ (m; \Delta; \beta; \alpha) &= (12; 5; 5, 638541\dots; 6, 813222\dots), \\ (m; \Delta; \beta; \alpha) &= (12; 7; 5, 696732\dots; 6, 983870\dots), \\ (m; \Delta; \beta; \alpha) &= (13; 5; 7, 177155\dots; 9, 376030\dots), \\ (m; \Delta; \beta; \alpha) &= (13; 7; 7, 190814\dots; 9, 594580\dots), \\ (m; \Delta; \beta; \alpha) &= (14; 5; 19, 659885\dots; 21, 835056\dots), \\ (m; \Delta; \beta; \alpha) &= (14; 7; 18, 447228\dots; 20, 668254\dots), \\ (m; \Delta; \beta; \alpha) &= (15; 5; 7, 508714\dots; 9, 922761\dots), \\ (m; \Delta; \beta; \alpha) &= (15; 7; 7, 516606\dots; 10, 156245\dots), \\ (m; \Delta; \beta; \alpha) &= (16; 5; 7, 951153\dots; 9, 876454\dots), \end{aligned}$$

$$\begin{aligned}
(m; \Delta; \beta; \alpha) &= (16; 7, 7, 945763\dots; 10, 039605\dots), \\
(m; \Delta; \beta; \alpha) &= (17; 5; 7, 797153\dots; 10, 399610\dots), \\
(m; \Delta; \beta; \alpha) &= (17; 7, 7, 799343\dots; 10, 645404\dots), \\
(m; \Delta; \beta; \alpha) &= (18; 5, 9, 486110\dots; 10, 955534\dots), \\
(m; \Delta; \beta; \alpha) &= (18; 7, 9, 406368\dots; 10, 989150\dots), \\
(m; \Delta; \beta; \alpha) &= (19; 5; 8, 052478\dots; 10, 822446\dots), \\
(m; \Delta; \beta; \alpha) &= (19; 7; 8, 049182\dots; 11, 078690\dots), \\
(m; \Delta; \beta; \alpha) &= (20; 5; 6, 696241\dots; 8, 559091\dots), \\
(m; \Delta; \beta; \alpha) &= (20; 7; 6, 733979\dots; 8, 774063\dots), \\
(m; \Delta; \beta; \alpha) &= (21; 5; 8, 281548\dots; 11, 202268\dots), \\
(m; \Delta; \beta; \alpha) &= (21; 7; 8, 273039\dots; 11, 467583\dots), \\
(m; \Delta; \beta; \alpha) &= (22; 5; 13, 134623\dots; 15, 504916\dots), \\
(m; \Delta; \beta; \alpha) &= (22; 7; 12, 815391\dots; 15, 331975\dots), \\
(m; \Delta; \beta; \alpha) &= (23; 5; 8, 489281\dots; 11, 547024\dots), \\
(m; \Delta; \beta; \alpha) &= (23; 7; 8, 475843\dots; 11, 820351\dots), \\
(m; \Delta; \beta; \alpha) &= (24; 5; 7, 088338\dots; 9, 210037\dots), \\
(m; \Delta; \beta; \alpha) &= (24; 7; 7, 116679\dots; 8, 439782\dots), \\
(m; \Delta; \beta; \alpha) &= (25; 5; 8, 679328\dots; 11, 862643\dots), \\
(m; \Delta; \beta; \alpha) &= (25; 7; 8, 661235\dots; 12, 143143\dots), \\
(m; \Delta; \beta; \alpha) &= (26; 5; 12, 172520\dots; 14, 674949\dots), \\
(m; \Delta; \beta; \alpha) &= (26; 7; 11, 944943\dots; 14, 618461\dots), \\
&\dots \\
(m; \Delta; \beta; \alpha) &= (32; 5; 8, 654733\dots; 11, 466214\dots), \\
(m; \Delta; \beta; \alpha) &= (32; 7; 8, 637697\dots; 11, 705492\dots), \\
(m; \Delta; \beta; \alpha) &= (33; 5; 9, 310125\dots; 12, 911341\dots), \\
(m; \Delta; \beta; \alpha) &= (33; 5; 9, 275806\dots; 13, 214792\dots),
\end{aligned}$$

References.

- [1] R.Apéry, Interpolation des fractions continues
et irrationalite de certaines constantes,
Bulletin de la section des sciences du C.T.H., 1981, No 3, 37 – 53;
- [2] F.Beukers, A note on the irrationality of $\zeta(2)$ and $\zeta(3)$,
Bull. London Math. Soc., 1979, 11, 268 – 272;
- [3] A.van der Porten, A proof that Euler missed...Apéry's proof of the irrationality of $\zeta(3)$,
Math Intellegencer, 1979, 1, 195 – 203;
- [4] W. Maier, Potenzreihen irrationalen Grenzwertes,
J.reine angew. Math., 156, 1927, 93 – 148;
- [5] E.M. Nikišin, On irrationality of the values of the functions $F(x,s)$ (in Russian),
Mat.Sb. 109 (1979), 410 – 417;
English transl. in Math. USSR Sb. 37 (1980), 381 – 388;

- [6] G.V. Chudnovsky, Pade approximations to the generalized hyper-geometric functions
I,J.Math.Pures Appl., 58, 1979, 445 – 476;
- [7] —————, Transcendental numbers, Number Theory, Carbondale,
Lecture Notes in Math, Springer-Verlag, 1979, 751, 45 – 69;
- [8] —————, Approximations rationnelles des logarithmes de nombres rationnelles
C.R.Acad.Sc. Paris, Série A, 1979, 288, 607 – 609;
- [9] —————, Formules d’Hermite pour les approximants de Padé de logarithmes
et de fonctions binômes, et mesures d’irrationalité,
C.R.Acad.Sc. Paris, Série A, 1979, t.288, 965 – 967;
- [10] —————, Un système explicite d’approximants de Padé
pour les fonctions hypérgéométriques généralisées,
avec applications à l’arithmétique,
C.R.Acad.Sc. Paris, Série A, 1979, t.288, 1001 – 1004;
- [11] —————, Recurrences defining Rational Approximations
to the irrational numbers, Proceedings
of the Japan Academie, Ser. A, 1982, 58, 129 – 133;
- [12] —————, On the method of Thue-Siegel,
Annals of Mathematics, 117 (1983), 325 – 382;
- [13] K.Alladi and M. Robinson, Legendre polynomials and irrationality,
J. Reine Angew.Math., 1980, 318, 137 – 155;
- [14] A. Dubitskas, An approximation of logarithms of some numbers,
Diophantine approximations II, Moscow, 1986, 20 – 34;
- [15] —————, On approximation of $\pi/\sqrt{3}$ by rational fractions,
Vestnik MGU, series 1, 1987, 6, 73 – 76;
- [16] S.Eckmann, Über die lineare Unabhängigkeit der Werte gewisser Reihen,
Results in Mathematics, 11, 1987, 7 – 43;
- [17] M.Hata, Legendre type polynomials and irrationality measures,
J. Reine Angew. Math., 1990, 407, 99 – 125;
- [18] A.O. Gelfond, Transcendental and algebraic numbers (in Russian),
GIT-TL, Moscow, 1952;
- [19] H.Bateman and A.Erdélyi, Higher transcendental functions, 1953,
New-York – Toronto – London, Mc. Grow-Hill Book Company, Inc.;
- [20] O.Perron, Über die Poincaresche Differenzgleichung,
Journal für die reine und angewandte mathematik,
1910, 137, 6 – 64;
- [21] A.O.Gelfond, Differenzenrechnung (in Russian), 1967, Nauka, Moscow.
- [22] A.O.Gelfond and I.M.Kubenskaya, On the theorem of Perron
in the theory of difference equations (in Russian),
IAN USSR, math. ser., 1953, 17, 2, 83 – 86.
- [23] M.A.Evgrafov, New proof of the theorem of Perron
(in Russian), IAN USSR, math. ser., 1953, 17, 2, 77 – 82;
- [24] G.A.Frejman, On theorems of Poincaré and Perron
(in Russian), UMN, 1957, 12, 3 (75), 243 – 245;
- [25] N.E.Nörlund, Differenzenrechnung, Berlin, Springer Verlag, 1924;
- [26] I.M.Vinogradov, Foundations of the Number Theory, (in Russian), 1952, GIT-TL;
- [27] —————, H.WEYL, Algebraic theory of numbers, 1940,
Russian translation by L.I.Kopejkina,
- [28] L.A.Gutnik, On the decomposition of the difference operators of Poincaré type
(in Russian), VINITI, Moscow, 1992, 2468 – 92, 1 – 55;
- [29] —————, On the decomposition of the difference operators
of Poincaré type in Banach algebras
(in Russian), VINITI, Moscow, 1992, 3443 – 92, 1 – 36;
- [30] —————, On the difference equations of Poincaré type
(in Russian), VINITI, Moscow 1993, 443 – B93, 1 – 41;
- [31] —————, On the difference equations of Poincaré type in normed algebras
(in Russian), VINITI, Moscow, 1994, 668 – B94, 1 – 44;
- [32] —————, On the decomposition of the difference equations of Poincaré type
(in Russian), VINITI, Moscow, 1997, 2062 – B97, 1 – 41;
- [33] —————, The difference equations of Poincaré type

- with characteristic polynomial having roots equal to zero
(in Russian), VINITI, Moscow, 1997, 2418 – 97, 1 – 20;
- [34] —————, On the behavior of solutions
of difference equations of Poincaré type
(in Russian), VINITI, Moscow, 1997, 3384 – B97, 1 - 41;
- [35] —————, On the variability of solutions of difference equations of Poincaré type
(in Russian), VINITI, Moscow, 1999, 361 – B99, 1 – 9;
- [36] —————, To the question of the variability of solutions
of difference equations of Poincaré type (in Russian),
VINITI, Moscow, 2000, 2416 – B00, 1 – 22;
- [37] —————, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$,
Max-Plank-Institut für Mathematik,
Bonn, Preprint Series, 2000, 3, 1 – 13;
- [38] —————, On the Irrationality of Some Quantities Containing $\zeta(3)$ (in Russian),
Uspekhi Mat. Nauk, 1979, 34, 3(207), 190;
- [39] —————, On the Irrationality of Some Quantities Containing $\zeta(3)$,
Eleven papers translated from the Russian,
American Mathematical Society, 1988, 140, 45 - 56;
- [40] —————, Linear independence over \mathbb{Q} of dilogarithms at rational points
(in Russian), UMN, 37 (1982), 179-180;
english transl. in Russ. Math. surveys 37 (1982), 176-177;
- [41] —————, On a measure of the irrationality of dilogarithms at rational points
(in Russian), VINITI, 1984, 4345-84, 1 – 74;
- [42] —————, To the question of the smallness of some linear forms
(in Russian), VINITI, 1993, 2413-B93, 1 – 94;
- [43] —————, About linear forms,
whose coefficients are logarithms
of algebraic numbers (in Russian),
VINITI, 1995, 135-B95, 1 – 149;
- [44] —————, About systems of vectors, whose coordinates
are linear combinations of logarithms of algebraic numbers
with algebraic coefficients (in Russian),
VINITI, 1994, 3122-B94, 1 – 158;
- [45] —————, On the linear forms, whose
coefficients are \mathbb{A} - linear combinations
of logarithms of \mathbb{A} - numbers,
VINITI, 1996, 1617-B96, pp. 1 – 23.
- [46] —————, On systems of linear forms, whose
coefficients are \mathbb{A} - linear combinations
of logarithms of \mathbb{A} - numbers,
VINITI, 1996, 2663-B96, pp. 1 – 18.
- [47] —————, About linear forms, whose coefficients
are \mathbb{Q} -proportional to the number $\log 2$, and the values
of $\zeta(s)$ for integer s (in Russian),
VINITI, 1996, 3258-B96, 1 – 70;
- [48] —————, The lower estimate for some linear forms,
coefficients of which are proportional to the values
of $\zeta(s)$ for integer s (in Russian),
VINITI, 1997, 3072-B97, 1 – 77;
- [49] —————, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2000, 3, 1 – 13;
- [50] —————, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$
(the detailed version, part 1), Max-Plank-Institut für Mathematik,
Bonn, Preprint Series, 2001, 15, 1 – 20;
- [51] —————, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$
(the detailed version, part 2), Max-Plank-Institut für Mathematik,
Bonn, Preprint Series, 2001, 104, 1 – 36;
- [52] —————, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$

- (the detailed version, part 3), Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2002, 57, 1 – 33;
- [53] —————, On the rank over \mathbb{Q} of some real matrices (in Russian), VINITI, 1984, 5736-84; 1 – 29;
- [54] —————, On the rank over \mathbb{Q} of some real matrices, Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2002, 27, 1 – 32;
- [55] —————, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$ (the detailed version, part 4), Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2002, 142, 1 – 27;
- [56] —————, On the dimension of some linear spaces over finite extension of \mathbb{Q} (part 2), Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2002, 107, 1 – 37;
- [57] —————, On the dimension of some linear spaces over \mathbb{Q} (part 3), Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2003, 16, 1 – 45.
- [58] —————, On the difference equation of Poincaré type (Part 1). Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2003, 52, 1 – 44.
- [59] —————, On the dimension of some linear spaces over \mathbb{Q} , (part 4) Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2003, 73, 1 – 38.
- [60] —————, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$ (the detailed version, part 5), Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2003, 83, 1 – 13.
- [61] —————, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$ (the detailed version, part 6), Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2003, 99, 1 – 33.
- [62] —————, On the difference equation of Poincaré type (Part 2). Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2003, 107, 1 – 25.
- [63] —————, On the asymptotic behavior of solutions of difference equation (in English). Chebyshevskij sbornik, 2003, v.4, issue 2, 142 – 153.
- [64] —————, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$, Bonner Mathematische Schriften Nr. 360, Bonn, 2003, 360.
- [65] —————, On the dimension of some linear spaces over \mathbb{Q} , (part 5) Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2004, 46, 1 – 42.
- [66] —————, On the difference equation of Poincaré type (Part 3). Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2004, 9, 1 – 33.
- [67] —————, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$ (the detailed version, part 7), Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2004, 88, 1 – 27.

E-mail: gutnik@gutnik.mccme.ru