

# ON CERTAIN 5-MANIFOLDS WITH FUNDAMENTAL GROUP OF ORDER 2

IAN HAMBLETON AND YANG SU

ABSTRACT. In this paper, an explicit classification result for certain 5-manifolds with fundamental group  $\mathbb{Z}/2$  is obtained. These manifolds include total spaces of circle bundles over simply-connected 4-manifolds.

## 1. INTRODUCTION

The classification of manifolds with certain properties is a central topic of topology, and in dimensions  $\geq 5$  methods from handlebody theory and surgery have been successfully applied to a number of cases. One of the first examples was the complete classification of simply-connected 5-manifolds by Smale [16] and Barden [1] in 1960's. This result has been very useful for studying the existence of other geometric structures on 5-manifolds, such as the existence of Riemannian metrics with given curvature properties. We consider this as a model and motivation for studying the classification of non-simply connected 5-manifolds.

An orientable 5-manifold  $M$  is said to be of *fibred type* if  $\pi_2(M)$  is a trivial  $\mathbb{Z}[\pi_1(M)]$ -module. In this paper, we will be concerned with closed, orientable fibred type 5-manifolds  $M^5$  with  $\pi_1(M) \cong \mathbb{Z}/2$ , and torsion free  $H_2(M; \mathbb{Z})$ . A classification of these manifolds in the smooth (or PL) and the topological category is given in Section 3. We give a simple set of invariants, namely the rank of  $H_2(M; \mathbb{Z})$  and the  $\text{Pin}^\dagger$ -bordism ( $\text{TopPin}^\dagger$ -bordism) class of the characteristic submanifold, which determine the diffeomorphism (homeomorphism) types. In the smooth case, the main result of the classification is:

**Theorem 3.1.** *Two smooth, closed, orientable fibred type 5-manifolds  $M$  and  $M'$  with fundamental group  $\mathbb{Z}/2$  and torsion free second homology group are diffeomorphic if and only if they have the same  $w_2$ -type,  $\text{rank } H_2(M) = \text{rank } H_2(M')$ , and  $[P] = [P'] \in \Omega_4^{\text{Pin}^\dagger} / \pm$ , where  $P$  and  $P'$  are characteristic submanifolds and  $\dagger = c, -, +$  for  $w_2$ -types I, II, III respectively.*

The homeomorphism classification is given in Theorem 3.4. We also determine all the relation among these invariants (Theorem 3.5), and give a list of standard forms for these manifolds (Theorem 3.6, Theorem 3.10).

One motivation for this classification problem comes from the study of circle bundles  $M^5$  over simply-connected 4-manifolds, since their total spaces are of fibred type. Duan-Liang [5] gave an explicit geometric description of  $M^5$  for simply-connected total spaces,

---

*Date:* June 24, 2009.

Research partially supported by NSERC Research Grant A4000. The first author would like to thank the Max Planck Institut für Mathematik in Bonn for its hospitality and support.

making essential use of the results of Smale and Barden. As an application of our results, in Section 4 we give an explicit geometric description when the total spaces have fundamental group  $\mathbb{Z}/2$ .

**Theorem 4.5** (type II). *Let  $X$  be a closed, simply-connected, topological spin 4-manifold,  $\xi: S^1 \hookrightarrow M^5 \rightarrow X$  be a circle bundle over  $X$  with  $c_1(\xi) = 2$ -primitive. Then we have*

(1) *if  $KS(X) = 0$ , then  $M$  is smoothable and  $M$  is diffeomorphic to*

$$(S^2 \times \mathbb{R}P^3) \#_{S^1} ((\#_k S^2 \times S^2) \times S^1);$$

(2) *if  $KS(X) = 1$ , then  $M$  is non-smoothable and  $M$  is homeomorphic to*

$$*(S^2 \times \mathbb{R}P^3) \#_{S^1} ((\#_k S^2 \times S^2) \times S^1).$$

Where  $k = \text{rank } H_2(X)/2 - 1$ .

In the statement,  $*(S^2 \times \mathbb{R}P^3)$  denotes a non-smoothable manifold homotopy equivalent to  $S^2 \times \mathbb{R}P^3$ . The corresponding results for the other  $w_2$ -types are given in Theorem 4.7 and Theorem 4.8.

Classification results can also be useful in studying the existence problem for geometric structures on fibered type 5-manifolds. For example, a closed, orientable 5-manifold with  $\pi_1 = \mathbb{Z}/2$ , such that  $w_2$  vanishes on homology, admits a contact structure by the work of Geiges and Thomas [6]. They showed that all such manifolds can be obtained by surgery on 2-dimensional links from exactly one of ten model manifolds. The topology of such manifolds of fibered type are described explicitly for the first time by our results, and we note that all the manifolds in the list in Theorem 3.6 satisfy the necessary condition  $W_3 = 0$  for the existence of contact structures. It may be possible to obtain similar information for fibered type 5-manifolds which admit Sasakian or Einstein metrics by using the work of Boyer and Galicki [2].

The surgery exact sequence of Wall [18] provides a way to classify manifolds within a given (simple) homotopy type. However, in the application to concrete problems, one often faces homotopy theoretical difficulties. In our situation, the setting of the problems is appropriate for the application of the modified surgery methods developed by Kreck [11]. The proofs in Section 5 and Section 6 are based on this theory.

In dimension 5, the smooth category and the PL category are equivalent. By convention,  $M$  stands for either a smooth or a topological manifold when not specified.

**Acknowledgement.** The second author would like to thank the James Stewart Centre for Mathematics at McMaster and the Institute for Mathematical Sciences of the National University of Singapore for research visits in Fall, 2008.

## 2. PRELIMINARIES

Let  $M$  be a closed, orientable 5-manifold with  $\pi_1(M) \cong \mathbb{Z}/2$  and universal cover  $\widetilde{M}$ . The manifold  $M$  is said to be of *type II* if  $w_2(M) = 0$ , of *type III* if  $w_2(M) \neq 0$  and  $w_2(\widetilde{M}) = 0$ , of *type I* if  $w_2(\widetilde{M}) \neq 0$ . By the universal coefficient theorem, there is an exact sequence

$$0 \rightarrow \text{Ext}(H_1(M), \mathbb{Z}/2) \rightarrow H^2(M; \mathbb{Z}/2) \rightarrow \text{Hom}(H_2(M), \mathbb{Z}/2) \rightarrow 0.$$

**Lemma 2.1.**  $M$  is of type II  $\Leftrightarrow w_2(M) = 0$ ; of type III  $\Leftrightarrow w_2(M) \neq 0$  and  $w_2(M) \in \text{Ext}(H_1(M), \mathbb{Z}/2)$ ; of type I  $\Leftrightarrow$  otherwise.

*Proof.* There is a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ext}(H_1(M), \mathbb{Z}/2) & \longrightarrow & H^2(M; \mathbb{Z}/2) & \longrightarrow & \text{Hom}(H_2(M), \mathbb{Z}/2) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \text{Ext}(H_1(\widetilde{M}), \mathbb{Z}/2) & \longrightarrow & H^2(\widetilde{M}; \mathbb{Z}/2) & \longrightarrow & \text{Hom}(H_2(\widetilde{M}), \mathbb{Z}/2) \longrightarrow 0. \end{array}$$

Let  $p: \widetilde{M} \rightarrow M$  be the covering map, then  $T\widetilde{M} = p^*TM$  and  $w_2(\widetilde{M}) = p^*w_2(M)$ . By the exact sequence  $\pi_2(M) \rightarrow H_2(M) \rightarrow H_2(\mathbb{Z}/2) \rightarrow 0$  and the fact  $H_2(\mathbb{Z}/2) = 0$  (cf. [3]), it is seen that the map  $H_2(\widetilde{M}) \rightarrow H_2(M)$  is surjective, therefore the last vertical map in the diagram  $\text{Hom}(H_2(M), \mathbb{Z}/2) \rightarrow \text{Hom}(H_2(\widetilde{M}), \mathbb{Z}/2)$  is a monomorphism. Thus  $w_2(\widetilde{M}) = 0$  if and only if  $w_2(M) \in \text{Ext}(H_1(M), \mathbb{Z}/2)$ .  $\square$

**Remark 2.2.** By this lemma, the type II and type III manifolds are manifolds having second Stiefel-Whitney class equal to zero on homology, which are studied in [6].

Recall that for a manifold  $M^n$  with fundamental group  $\mathbb{Z}/2$ , a *characteristic submanifold*  $P^{n-1} \subset M$  is defined in the following way (see [6, §5]): there is a decomposition  $\widetilde{M} = A \cup T A$  such that  $\partial A = \partial T A = \widetilde{P}$ , where  $T$  is the deck-transformation. Then  $P := \widetilde{P}/T$  is called the characteristic submanifold of  $M$ . For example, if  $M = \mathbb{R}P^n$ , then  $P = \mathbb{R}P^{n-1}$ . In general, let  $f: M \rightarrow \mathbb{R}P^N$  ( $N$  large) be the classifying map of the universal cover, transverse to  $\mathbb{R}P^{N-1}$ , then  $P$  can be taken as  $f^{-1}(\mathbb{R}P^{n-1})$ . By equivariant surgery we may assume that  $\pi_1(P) \cong \mathbb{Z}/2$  and that the inclusion  $i: P \subset M$  induces an isomorphism on  $\pi_1$ . The above construction also holds in the topological category by topological transversality [9].

Recall that there are central extensions of  $O(n)$  by  $\mathbb{Z}/2$  (see [10, §1] and [7, §2]):

$$1 \rightarrow \mathbb{Z}/2 \rightarrow \text{Pin}^\pm(n) \rightarrow O(n) \rightarrow 1,$$

and central extension

$$1 \rightarrow U(1) \rightarrow \text{Pin}^c(n) \rightarrow O(n) \rightarrow 1.$$

Let  $\dagger \in \{c, +, -\}$ . A  $\text{Pin}^\dagger$ -structure on a vector bundle  $\xi$  over a space  $X$  is the fiber homotopy class of lifts of the classifying map  $c_E: X \rightarrow BO$ .

**Lemma 2.3.** [7, Lemma 1]

(1) A vector bundle  $E$  over  $X$  admits a  $\text{Pin}^\dagger$ -structure if and only if

$$\begin{aligned} \beta(w_2(E)) &= 0 & \text{for } \dagger = c, \\ w_2(E) &= 0 & \text{for } \dagger = +, \\ w_2(E) &= w_1(E)^2 & \text{for } \dagger = -, \end{aligned}$$

where  $\beta: H^2(X; \mathbb{Z}/2) \rightarrow H^3(X; \mathbb{Z})$  is the Bockstein operator induced from the exact coefficient sequence  $\mathbb{Z} \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}/2$ .

- (2)  $\text{Pin}^\pm$ -structures are in bijection with  $H^1(X; \mathbb{Z}/2)$  and  $\text{Pin}^c$ -structures are in bijection with  $H^2(X; \mathbb{Z})$ .

In the smooth category, the division of the manifolds under consideration into 3 types corresponds to different  $\text{Pin}^\dagger$ -structures on their characteristic submanifold, compare [6, Lemma 9] for  $\dagger = \pm$ .

**Lemma 2.4.** *Let  $M$  be a smooth, orientable 5-manifold with  $\pi_1(M) \cong \mathbb{Z}/2$  and  $H_2(M; \mathbb{Z})$  torsion free. Let  $P \subset M$  be the characteristic submanifold (with  $\pi_1(P) \cong \mathbb{Z}/2$ ). Then  $TP$  admits a  $\text{Pin}^\dagger$ -structure, where*

$$\dagger = \begin{cases} c & \text{if } M \text{ is of type I} \\ - & \text{if } M \text{ is of type II} \\ + & \text{if } M \text{ is of type III} \end{cases}$$

*Proof.* Let  $i: P \subset M$  be the inclusion and  $\nu$  be the normal bundle of this inclusion. Then  $TP \oplus \nu = i^*TM$ , hence by the product formula of the Stiefel-Whitney classes we have  $w_2(P) + w_1(P) \cdot w_1(\nu) = i^*w_2(M)$ . Because  $P$  is nonorientable,  $\nu$  is the nonorientable line bundle. Now  $H^1(P; \mathbb{Z}/2) \cong \mathbb{Z}/2$ , we see that  $w_1(P) = w_1(\nu)$ , therefore  $w_2(P) + w_1(P)^2 = i^*w_2(M)$ .

If  $M$  is of type II,  $w_2(M) = 0$ , then  $w_2(P) + w_1(P)^2 = 0$ , and  $P$  admits a  $\text{Pin}^-$ -structure. If  $M$  is of type III,  $w_2(M)$  is the nonzero element in  $\text{Ext}(H_1(M), \mathbb{Z}/2) \cong \mathbb{Z}/2$ . Then  $i^*w_2(M) = w_1(P)^2$  because of the isomorphism

$$i^*: \text{Ext}(H_1(M), \mathbb{Z}/2) \rightarrow \text{Ext}(H_1(P), \mathbb{Z}/2)$$

and the fact that the nonzero element in  $\text{Ext}(H_1(P), \mathbb{Z}/2)$  is  $w_1(P)^2$ .

In general, since  $H^3(M; \mathbb{Z}) \cong H_2(M; \mathbb{Z})$  is torsion free, the Bockstein homomorphism  $\beta: H^2(M; \mathbb{Z}/2) \rightarrow H^3(M; \mathbb{Z})$  is trivial. Therefore  $\beta i^*w_2(M) = i^*\beta w_2(M) = 0$ . In the classifying space, the Bockstein homomorphism  $\beta: H^2(\mathbb{R}P^\infty; \mathbb{Z}/2) \rightarrow H^3(\mathbb{R}P^\infty; \mathbb{Z})$  is trivial, since  $\beta$  equals to the Steenrod square  $Sq^1$ , which is trivial. Now  $w_1(P)$  is the pullback of the nonzero element in  $H^1(\mathbb{R}P^\infty; \mathbb{Z}/2)$ , and therefore  $w_1(P)^2$  is the pullback of the nonzero element in  $H^2(\mathbb{R}P^\infty; \mathbb{Z}/2)$ . Thus we have  $\beta(w_1(P)^2) = 0$  by naturality. Therefore  $\beta w_2(P) = 0$  and  $P$  admits a  $\text{Pin}^c$ -structure.  $\square$

### 3. MAIN RESULTS

For a given  $M^5$ , different characteristic submanifolds are bordant in the corresponding  $\text{Pin}^\dagger$ -bordism group, where a bordism is obtained from a homotopy between the relevant classifying maps. A concrete  $\text{Pin}^\dagger$ -structure is given by a  $\text{Spin}$ -structure on a stable vector bundle related to  $\nu M$ , which will be explained in Section 5. The two different  $\text{Pin}^\dagger$ -structures on the characteristic submanifold correspond to a pair of elements inverse to each other in  $\Omega_4^{\text{Pin}^\dagger}$ .

**Theorem 3.1.** *Two smooth, closed, orientable fibered type 5-manifolds  $M$  and  $M'$  with fundamental group  $\mathbb{Z}/2$  and torsion free second homology group are diffeomorphic if and only if they have the same  $w_2$ -type,  $\text{rank } H_2(M) = \text{rank } H_2(M')$ , and  $[P] = [P'] \in \Omega_4^{\text{Pin}^\dagger} / \pm$ , where  $P$  and  $P'$  are the characteristic submanifolds and  $\dagger = c, -, +$  for types I, II, III respectively.*

**Remark 3.2.** We will see in a moment that  $\Omega_4^{\text{Pin}^-} = 0$ . Therefore for the type II manifolds  $\text{rank } H_2(M)$  is the only diffeomorphism invariant.

There are topological versions of the central extensions mentioned above and we have groups  $\text{TopPin}^\dagger(n)$ ,  $\dagger \in \{c, +, -\}$ . Lemma 2 holds in the topological case. For the preliminaries on  $\text{TopPin}^\dagger(n)$  we refer to [10] and [7]. Therefore we have corresponding results in the topological category.

**Lemma 3.3.** *Let  $M$  be a topological, orientable 5-manifold with  $\pi_1(M) \cong \mathbb{Z}/2$  and  $H_2(M; \mathbb{Z})$  torsion free. Let  $P \subset M$  be a characteristic submanifold (with  $\pi_1(P) \cong \mathbb{Z}/2$ ). Then  $TP$  admits a  $\text{TopPin}^\dagger$ -structure, where*

$$\dagger = \begin{cases} c & \text{if } M \text{ is of type I} \\ - & \text{if } M \text{ is of type II} \\ + & \text{if } M \text{ is of type III} \end{cases}$$

**Theorem 3.4.** *Two topological, closed, orientable fibered type 5-manifolds  $M$  and  $M'$  with fundamental group  $\mathbb{Z}/2$  and torsion free second homology group are homeomorphic if and only if they have the same  $w_2$ -type,  $\text{rank } H_2(M) = \text{rank } H_2(M')$  and  $[P] = [P'] \in \Omega_4^{\text{TopPin}^\dagger} / \pm$ , where  $P$  and  $P'$  are characteristic submanifolds and  $\dagger = c, -, +$  for type I, II, III respectively.*

The groups  $\Omega_4^{\text{Pin}^\pm}$  and  $\Omega_4^{\text{TopPin}^\pm}$  are computed in [10].  $\Omega_4^{\text{TopPin}^c}$  is computed in [7, p.654]. (Note that the rôle of  $\text{Pin}^+$  and  $\text{Pin}^-$  in [7] are reversed since in that paper the authors consider normal structures whereas here we use the convention in [10], looking at the tangential Gauss-map.) In a similar way we will compute  $\Omega_4^{\text{Pin}^c}$  below. We list the values of these groups:

$\dagger$	$\Omega_4^{\text{Pin}^\dagger}$	invariants	generators
$c$	$\mathbb{Z}/8 \oplus \mathbb{Z}/2$	$(\text{arf}, w_2^2)$	$\mathbb{RP}^4, \mathbb{CP}^2$
$+$	$\mathbb{Z}/16$	?	$\mathbb{RP}^4$
$-$	$0$	$-$	$-$
$\dagger$	$\Omega_4^{\text{TopPin}^\dagger}$	invariants	generators
$c$	$\mathbb{Z}/2 \oplus \mathbb{Z}/8 \oplus \mathbb{Z}/2$	$(KS, \text{arf}, w_2^2)$	$E_8, \mathbb{RP}^4, \mathbb{CP}^2$
$+$	$\mathbb{Z}/2 \oplus \mathbb{Z}/8$	$(KS, \text{arf})$	$E_8, \mathbb{RP}^4$
$-$	$\mathbb{Z}/2$	$KS$	$E_8$

Computation of  $\Omega_4^{\text{Pin}^c}$ : the extension

$$1 \rightarrow \text{Pin}^- \rightarrow \text{Pin}^c \rightarrow U(1) \rightarrow 1$$

induces Gysin-sequence (compare [7, p.654])

$$\dots \rightarrow \Omega_4^{\text{Pin}^-} \rightarrow \Omega_4^{\text{Pin}^c} \xrightarrow{\cap c} \Omega_2^{\text{Pin}^-}(BU(1)) \rightarrow \Omega_3^{\text{Pin}^-} \rightarrow \dots$$

Since  $\Omega_4^{\text{Pin}^-} = \Omega_3^{\text{Pin}^-} = 0$  (see [10]), we have an isomorphism

$$\Omega_4^{\text{Pin}^c} \xrightarrow{\cap c} \Omega_2^{\text{Pin}^-}(BU(1))$$

and the latter group is the same as  $\Omega_2^{\text{TopPin}^-}(BU(1))$ , which is computed in [7].

The invariants in Theorem 3.1 are subject to certain relations. Denote  $r = \text{rank } H_2(M)$ ,  $q = [P] \in \Omega_4^{\text{Pin}^+}/\pm = \{0, 1, \dots, 8\}$  and  $(q, s) = [P] \in \Omega_4^{\text{Pin}^c}/\pm = \{0, 1, \dots, 4\} \times \{0, 1\}$ . As an application of the semi-characteristic class [13], we have

**Theorem 3.5.** *Let  $M$  be a smooth, orientable 5-manifold with  $\pi_1(M) \cong \mathbb{Z}/2$  and torsion free  $H_2(M)$ , having the invariants as above. Then these invariants subject to the following relations*

<i>type</i>	<i>relation</i>
I	$q + s + r \equiv 1 \pmod{2}$
II	$r \equiv 1 \pmod{2}$
III	$q + r \equiv 1 \pmod{2}$

Now we give a list of all the manifolds under consideration, realizing the possible invariants. We need some preliminaries.

By a computation of the surgery exact sequence, it is shown in [18] that in the smooth (or PL) category, there are 4 distinct diffeomorphism types of manifolds which are homotopy equivalent to  $\mathbb{R}P^5$ , these are called fake  $\mathbb{R}P^5$ . An explicit construction using links of singularities (Brieskorn spheres) can be found in [6]. Following the notations there, we denote these fake  $\mathbb{R}P^5$  by  $X^5(q)$ ,  $q = 1, 3, 5, 7$ , with  $X^5(1) = \mathbb{R}P^5$ . These manifolds fall into the class of manifolds under consideration. They are of type III and the  $\text{Pin}^+$ -bordism class of the corresponding characteristic submanifold is  $q \in \Omega_4^{\text{Pin}^+}/\pm = \{0, 1, \dots, 8\}$ , see [6]. In our list of standard forms these fake projective spaces will serve as building blocks under the operation  $\sharp_{S^1}$ —“connected-sum along  $S^1$ ”, which we explain now, compare [7].

Let  $M_i$  ( $i = 1, 2$ ) be oriented 5-manifolds with fundamental group  $\mathbb{Z}/2$  or  $\mathbb{Z}$ , and at least one of the fundamental groups is  $\mathbb{Z}/2$ . Denote the trivial oriented 4-dimensional real disc bundle over  $S^1$  by  $E$ . Choose embeddings of  $E$  into  $M_1$  and  $M_2$ , representing a generator of  $\pi_1(M_i)$ , such that the first embedding preserves the orientation and the second reverses it. Then we define

$$M_1 \sharp_{S^1} M_2 := (M_1 - E) \cup_{\partial} (M_2 - E).$$

Note that if one of the 5-manifolds admits an orientation reversing automorphism, then the construction doesn't depend on the orientations, and this is the case for the building blocks in the list below, namely,  $S^2 \times \mathbb{R}P^3$ ,  $S^2 \times S^2 \times S^1$ ,  $X^5(q)$  and  $\mathbb{C}P^2 \times S^1$  admit orientation reversing automorphisms. (The fact that  $X^5(q)$  admits orientation reversing automorphisms follows from that  $\mathbb{R}P^5$  admits orientation reversing automorphisms and that the action of  $\text{Aut}(\mathbb{R}P^5)$  on the structure set  $\mathcal{S}(\mathbb{R}P^5)$  is trivial.)

The Seifert-van Kampen theorem implies that  $\pi_1(M_1 \sharp_{S^1} M_2) \cong \mathbb{Z}/2$ . The Mayer-Vietoris exact sequence implies that  $H_2(M_1 \sharp_{S^1} M_2)$  is torsion free, and hence  $M_1 \sharp_{S^1} M_2$  is of fibered type. The homology rank  $H_2(M_1 \sharp_{S^1} M_2) = \text{rank } H_2(M_1) + \text{rank } H_2(M_2) + 1$  if

both fundamental groups are  $\mathbb{Z}/2$ , and  $\text{rank } H_2(M_1 \#_{S^1} M_2) = \text{rank } H_2(M_1) + \text{rank } H_2(M_2)$  if one of the fundamental groups is  $\mathbb{Z}$ .

Since  $\pi_1 SO(4) \cong \mathbb{Z}/2$ , there are actually two possibilities to form  $M_1 \#_{S^1} M_2$ . However, from the classification result, it turns out that this ambiguity happens only when we construct  $X^5(q) \#_{S^1} X^5(q')$ . This does depend on the framings, and therefore  $X^5(q) \#_{S^1} X^5(q')$  represents two manifolds. Note that the characteristic submanifold of  $M_1 \#_{S^1} M_2$  is  $P_1 \#_{S^1} P_2$  (see [7, p.651] for the definition of  $\#_{S^1}$  for nonorientable 4-manifolds with fundamental group  $\mathbb{Z}/2$ ). Therefore if we fix  $\text{Pin}^+$ -structures on each of the characteristic submanifolds, then  $X^5(q) \#_{S^1} X^5(q')$  is well-defined.

This construction allows us to construct manifolds with a given bordism class of characteristic submanifold. Note that  $P_1 \#_{S^1} P_2$  corresponds to the addition in the bordism group  $\Omega_4^{\text{Pin}^+}$ . Now for  $q = 0, 2, 4, 6, 8$ , choose  $l, l' \in \{1, 3, 5, 7\}$  and appropriate  $\text{Pin}^+$ -structures on the characteristic submanifolds of  $X^5(l)$  and  $X^5(l')$ , we can form a manifold  $X^5(l) \#_{S^1} X^5(l')$  such that the characteristic submanifold  $[P] = q \in \Omega_4^{\text{Pin}^+}/\pm$ . We denote this manifold also by  $X^5(q)$ . For example, we can form  $X^5(0) = X^5(1) \#_{S^1} X^5(1)$  and  $X^5(2) = X^5(1) \#_{S^1} X^5(1)$  with different glueing maps.

With these notations, the list of standard forms of the manifolds under consideration is given as follows:

**Theorem 3.6.** *Every closed smooth orientable fibered type 5-manifold with fundamental group  $\mathbb{Z}/2$  and second homology group  $\mathbb{Z}^r$  is diffeomorphic to exactly one of the following standard forms:*

$$\text{type I : } X^5(q) \#_{S^1} (S^2 \times \mathbb{RP}^3) \#_{S^1} ((\#_k S^2 \times S^2) \times S^1), \quad r = 2k + (5 + (-1)^q)/2, \quad q \in \{0, \dots, 4\};$$

$$X^5(q) \#_{S^1} (\mathbb{CP}^2 \times S^1) \#_{S^1} ((\#_k S^2 \times S^2) \times S^1), \quad r = 2k + (3 + (-1)^q)/2, \quad q \in \{0, \dots, 4\};$$

$$\text{type II : } (S^2 \times \mathbb{RP}^3) \#_{S^1} ((\#_k S^2 \times S^2) \times S^1), \quad r = 2k + 1;$$

$$\text{type III : } X^5(q) \#_{S^1} ((\#_k S^2 \times S^2) \times S^1), \quad r = 2k + (1 + (-1)^q)/2, \quad q \in \{0, \dots, 8\}.$$

Where  $\#_k S^2 \times S^2$  is the connected sum of  $k$  copies of  $S^2 \times S^2$ .

**Remark 3.7.** There can be other descriptions of the manifolds in the list. For example, we have a (more symmetric) description of the type II standard forms

$$\underbrace{(S^2 \times \mathbb{RP}^3) \#_{S^1} \cdots \#_{S^1} (S^2 \times \mathbb{RP}^3)}_{k \text{ times}}.$$

**Remark 3.8.** Note that the universal covers of the manifolds under consideration have torsion free second homology, therefore, according to the results of Smale and Barden, are diffeomorphic to  $\#_r(S^2 \times S^3)$  or  $B\#_{r-1}(S^2 \times S^3)$ , where  $B$  is the nontrivial  $S^3$ -bundle over  $S^2$ . From this point of view, Theorem 3.6 gives the classification of orientation preserving free involutions on  $\#_r(S^2 \times S^3)$  and  $B\#_{r-1}(S^2 \times S^3)$ , which act trivially on  $H_2$ . For example, consider the orientation preserving free involution on  $S^2 \times S^3$  given by  $(x, y) \mapsto (r(x), -y)$ , where  $r: S^2 \rightarrow S^2$  is the reflection along a line and  $-: S^3 \rightarrow S^3$  is the

antipodal map. Then the quotient space is actually the sphere bundle of the nontrivial orientable  $\mathbb{R}P^3$ -bundle over  $\mathbb{R}P^3$ . From Theorem 3.1 it is easy to see that this is just  $X^5(0)$ .

**Remark 3.9.** The above list may be of use in the study of geometric structures on these manifolds. Geiges and Thomas [6] show that the type II and type III manifolds admit contact structures. On the other hand, a necessary condition for the existence of contact structures on  $M^{2n+1}$  is the reduction of the structure group of  $TM$  to  $U(n)$ , hence the vanishing of integral Stiefel-Whitney classes  $W_{2i+1}(M)$ . It is easy to see that the type I manifolds satisfy this necessary condition. These manifolds also satisfy the necessary conditions on the cup length and Betti numbers in [2] for the existence of Sasakian structures. Therefore it would be interesting to study these geometric structures on these manifolds.

To give a list of standard forms of the manifolds under consideration in the topological case, we need a topological 5-manifold which is homotopy equivalent to  $S^2 \times \mathbb{R}P^3$  and whose characteristic submanifold represents the nontrivial element in  $\Omega_4^{\text{TopPin}^-} = \mathbb{Z}/2$ . Note that by Theorem 3.4, if such manifolds exist, then the homeomorphism type is unique. Following the notation in [7], we denote this manifold by  $*(S^2 \times \mathbb{R}P^3)$ . We now give the construction of  $*(S^2 \times \mathbb{R}P^3)$ .

Let  $W = S^2 \times \mathbb{R}P^3 \#_{S^1} E_8 \times S^1$ , so that  $\pi_1(W) = \mathbb{Z}/2$  and the characteristic submanifold of  $W$  is  $S^2 \times \mathbb{R}P^2 \# E_8$ . Let  $h: W \rightarrow S^2 \times \mathbb{R}P^3$  be a degree 1 normal map which extends the degree 1 normal map  $f: S^2 \times \mathbb{R}P^2 \# E_8 \rightarrow S^2 \times \mathbb{R}P^2$ . Then by doing codimension 1 surgery on  $h$  we obtain a  $W'$  with characteristic submanifold  $P = *(S^2 \times \mathbb{R}P^2)$  and a degree 1 normal map  $h': W' \rightarrow S^2 \times \mathbb{R}P^3$  extending a homotopy equivalence  $f': *(S^2 \times \mathbb{R}P^2) \rightarrow S^2 \times \mathbb{R}P^2$  (cf. [7] for the construction of  $*(S^2 \times \mathbb{R}P^2)$ ). The  $\pi$ - $\pi$  theorem allows us to do further surgeries on the complement of a tubular neighbourhood of  $P$  to obtain a homotopy equivalence.

In the topological category there are four fake  $\mathbb{R}P^5$ 's. Two of them are smoothable. We denote these manifolds by  $X^5(p, q)$  ( $p \in \{0, 1\}$ ,  $q \in \{1, 3\}$ ) such that the characteristic submanifold of  $X^5(p, q)$  is  $(p, q) \in \Omega_4^{\text{TopPin}^+} / \pm = \{0, 1\} \times \{0, 1, 2, 3, 4\}$ . Similar to the smooth case, we can also construct  $X^5(p, q)$  ( $p \in \{0, 1\}$ ,  $q \in \{0, 2, 4\}$ ) by circle connected sum of fake  $\mathbb{R}P^5$ . (Note that the Kirby-Siebenmann invariant is additive under the connected sum operation [15]).

**Theorem 3.10.** *Every closed topological orientable fibered type 5-manifold with fundamental group  $\mathbb{Z}/2$  and second homology group  $\mathbb{Z}^r$  is homeomorphic to exactly one of the following standard forms:*

$$\begin{aligned} \text{type I : } X^5(p, q) \#_{S^1} (S^2 \times \mathbb{R}P^3) \#_{S^1} ((\#k S^2 \times S^2) \times S^1), \\ r = 2k + (5 + (-1)^q)/2, q \in \{0, \dots, 4\}, p = 0, 1; \end{aligned}$$

$$\begin{aligned} X^5(p, q) \#_{S^1} (\mathbb{C}P^2 \times S^1) \#_{S^1} ((\#k S^2 \times S^2) \times S^1), \\ r = 2k + (3 + (-1)^q)/2, q \in \{0, \dots, 4\}, p = 0, 1; \end{aligned}$$

$$\text{type II : } (S^2 \times \mathbb{R}P^3) \#_{S^1} ((\#k S^2 \times S^2) \times S^1), r = 2k + 1;$$



$$*(S^2 \times \mathbb{R}P^3) \#_{S^1} ((\#k S^2 \times S^2) \times S^1), r = 2k + 1;$$

$$\text{type III} : X^5(p, q) \#_{S^1} ((\#k S^2 \times S^2) \times S^1), r = 2k + (1 + (-1)^q)/2, q \in \{0, \dots, 4\}, p = 0, 1.$$

From the above list, we can also give a homotopy classification.

**Theorem 3.11.** *The homotopy type of  $M^5$  is determined by its  $w_2$ -type,  $\text{rank } H_2(M)$ , and in the type I case the number  $\langle w_2(M)^2 \cup t + t^5, [M] \rangle \in \mathbb{Z}/2$ , where  $t \in H^1(M; \mathbb{Z}/2)$  is the nonzero element.*

#### 4. CIRCLE BUNDLES OVER 1-CONNECTED 4-MANIFOLDS

Let  $X^4$  be a simply-connected 4-manifold, smooth or topological. Let  $\xi$  be a complex line bundle over  $X$ , with first Chern class  $c_1(\xi) \in H^2(X; \mathbb{Z})$ . Choose a Riemannian metric on  $\xi$ , then the total space of the corresponding circle bundle is a 5-manifold  $M$ . The homotopy long exact sequence of the fiber bundle shows that  $\pi_1(M) \cong \mathbb{Z}/m$  if  $c_1(\xi)$  is an  $m$  multiple of a primitive element.

In [5], a classification of  $M$  in terms of the topological invariants of  $X$  and  $c_1(\xi)$  is obtained for  $m = 1$ , using the classification theorem of Smale and Barden. It is also known that  $H_2(M)$  is torsion free of rank  $H_2(X) - 1$  and that  $M$  is of fibered type. In this Section, we will apply the results in last section to the  $m = 2$  case, give classification of  $M$  in terms of the topological invariants of  $X$  and  $c_1(\xi)$ . We will also identify  $M$  in the list of standard forms in Theorem 3.6 and Theorem 3.10.

**§4A. Invariants of  $M$ .** In this subsection we collect the basic algebraic-topological invariants of  $M$ .

**Proposition 4.1.** *Let  $M^5$  be a circle bundle over a simply-connected 4-manifold  $X$ , with first Chern class  $c_1(\xi) = 2 \cdot \text{primitive}$ , then*

- (1)  $\pi_1(M) \cong \mathbb{Z}/2$
- (2)  $H_2(M) \cong \mathbb{Z}^r$  where  $r = \text{rank } H_2(X) - 1$ .
- (3) the  $\pi_1(M)$ -action on  $H_2(\widetilde{M})$  is trivial.
- (4) the type of  $M^5$  is given by

type I	type II	type III
$w_2(X) \neq 0$		
$w_2(X) \not\equiv c_1(\widetilde{\xi}) \pmod{2}$	$w_2(X) = 0$	$w_2(X) \equiv c_1(\widetilde{\xi}) \pmod{2}$

*Proof.* First of all, the homotopy long exact sequence

$$\pi_1(S^1) \rightarrow \pi_1(M) \rightarrow \pi_1(X)$$

implies that  $\pi_1(M)$  is a cyclic group. The Gysin sequence

$$0 \rightarrow H_2(M) \rightarrow H_2(X) \xrightarrow{\cap c_1} H_0(X) \rightarrow H_1(M) \rightarrow 0$$

shows that  $H_2(M)$  is torsion free of rank equal to  $\text{rank } H_2(X) - 1$  and  $H_1(M) \cong \mathbb{Z}/2$  since  $c_1(\xi) = 2 \cdot \text{primitive}$ . Note that the universal cover  $\widetilde{M}$  is a circle bundle over  $X$ , denoted

by  $\tilde{\xi}$ , with first Chern class  $c_1(\tilde{\xi}) = \frac{1}{2}c_1(\xi)$ . The  $\pi_1(M)$ -action on  $\tilde{M}$  is the antipodal map on each fiber, thus the commutative diagram

$$\begin{array}{ccc} H_2(\tilde{M}) & \xrightarrow{T_*} & H_2(\tilde{M}) \\ & \searrow p_* & \downarrow p_* \\ & & H_2(X) \end{array}$$

shows that the action on  $H_2(\tilde{M})$  is trivial. For the Stiefel-Whitney class, if  $X$  is smooth, we have  $TM \oplus \mathbb{R} = p^*(TX \oplus \xi)$  (where  $p$  is the projection map), this implies  $w_2(M) = p^*w_2(X)$ . In general,  $X - pt$  admits a smooth structure, then the same argument holds, see [5, Lemma 3].  $\square$

#### §4B. Smoothings of $M$ .

**Proposition 4.2.** *Let  $\xi: S^1 \hookrightarrow M^5 \rightarrow X$  be a nontrivial circle bundle over a closed, simply-connected, topological 4-manifold. If  $c_1(\xi)$  is an odd multiple of a primitive element, then  $M$  is smoothable; if  $c_1(\xi)$  is an even multiple of a primitive element, then  $M$  admits a smooth structure if and only if  $KS(X) = 0$ .*

*Proof.* Let  $M^5$  be a topological 5-manifold, then by [9], the obstruction for smoothing  $M$  lies in  $H^4(M; \pi_3(Top/O)) = H^4(M; \pi_3(Top/PL)) = H^4(M; \mathbb{Z}/2) \cong H_1(M; \mathbb{Z}/2)$ . The latter group is trivial if  $c_1(\xi)$  is an odd multiple of a primitive element. On the other hand, we have  $TM \oplus \mathbb{R} = \pi^*(TX \oplus \xi)$ , where  $\pi$  is the projection map. Therefore the obstruction for smoothing  $M$  is  $\pi^*KS(X)$ . It is seen from the Gysin sequence that  $\pi^*: H^4(X; \mathbb{Z}/2) \rightarrow H^4(M; \mathbb{Z}/2)$  is injective if  $c_1(\xi)$  is an even multiple of a primitive element. Therefore  $M$  admits a smooth structure if and only if  $KS(X) = 0$ .  $\square$

Now we give a geometric description of the characteristic submanifold of a circle bundle over simply-connected  $X^4$ .

**Lemma 4.3.** *Let  $\xi: S^1 \hookrightarrow M^5 \rightarrow X$  be a circle bundle,  $\pi_1(M) \cong \mathbb{Z}/2$ . Let  $F \subset X$  be an embedded surface dual to  $c_1(\tilde{\xi})$ ,  $N(F)$  be a tubular neighborhood of  $F$  in  $X$ ,  $S^1 \hookrightarrow B \rightarrow F$  be the restriction of  $\xi$  on  $F$ . Then there is a double cover map  $\partial N(F) \rightarrow B$  and the characteristic submanifold of  $M$  is  $P^4 = (X - \mathring{N}(F)) \cup_{\partial} B$ .*

In other words, the characteristic submanifold  $P$  is obtained by removing a tubular neighborhood of an embedded surface dual to  $c_1(\tilde{\xi})$  and then identifying antipodal points on each fiber.

*Proof.* Since  $c_1(\xi) = 2$ -primitive, the circle bundle is the pull-back of the circle bundle over  $\mathbb{C}P^2$  with first Chern class = 2-primitive:

$$\begin{array}{ccc} S^1 & \xrightarrow{=} & S^1 \\ \downarrow & & \downarrow \\ M^5 & \xrightarrow{f} & \mathbb{R}P^5 \\ \downarrow & & \downarrow \\ X & \xrightarrow{g} & \mathbb{C}P^2 \end{array} \quad \begin{array}{l} \supset \mathbb{R}P^4 = \mathbb{R}P^3 \cup D^4 \\ \\ = \mathbb{C}P^1 \cup D^4 \end{array}$$

Now  $P = f^{-1}(\mathbb{R}P^4) = f^{-1}(D^4 \cup_{S^3} \mathbb{R}P^3)$ . Let  $F = g^{-1}(\mathbb{C}P^1)$  be the transverse preimage of  $\mathbb{C}P^1$ , then the normal bundle  $\nu$  of  $F$  in  $X$  is the pullback of the Hopf bundle, and the restriction of  $\xi$  on  $F$  is  $\nu \otimes \nu$ , therefore there is a double cover  $\partial N(F) \rightarrow B$ . It is easy to see that  $P^4 = (X - \mathring{N}(F)) \cup_{\partial} B$ .  $\square$

**Lemma 4.4.** *Let  $P$  be as above. Then  $KS(P) = KS(X)$ .*

*Proof.* We identify  $N(F)$  with the normal 2-disk bundle, let  $V$  be the associated  $\mathbb{R}P^2$ -bundle obtained by identifying antipodal points on  $\partial N(F)$ . Then by the construction,

$$P = X \cup_{N(F) \times \{0\}} N(F) \times I \cup_{N(F) \times \{1\}} V.$$

Therefore  $P$  is bordant to  $X \sqcup V$ . It was shown by Hsu [8] and Lashof-Taylor [12] that the Kirby-Siebenmann invariant is a bordism invariant, thus  $KS(P) = KS(X) + KS(V) = KS(X)$  since  $V$  is smooth.  $\square$

**§4C. Classification.** Now we can give a classification of circle bundles over 1-connected 4-manifolds, identify them with the standard forms in Theorem 3.6 and Theorem 3.10, in terms of the topology of  $X$  and  $\xi$ .

For the type II manifolds it is an immediate consequence of Theorem 3.1 and Theorem 3.4.

**Theorem 4.5** (type II). *Let  $X$  be a closed, simply-connected, topological spin 4-manifold,  $\xi: S^1 \hookrightarrow M^5 \rightarrow X$  be a circle bundle over  $X$  with  $c_1(\xi) = 2$ -primitive. Then we have*

(1) *if  $KS(X) = 0$ , then  $M$  is smoothable and  $M$  is diffeomorphic to*

$$(S^2 \times \mathbb{R}P^3) \#_{S^1} ((\#_k S^2 \times S^2) \times S^1);$$

(2) *if  $KS(X) = 1$ , then  $M$  is non-smoothable and  $M$  is homeomorphic to*

$$*(S^2 \times \mathbb{R}P^3) \#_{S^1} ((\#_k S^2 \times S^2) \times S^1).$$

Where  $k = \text{rank } H_2(X)/2 - 1$ .

**Remark 4.6.** Note that for a spin 4-manifold  $X$ ,  $\text{rank } H_2(X)$  is even, and thus  $k$  is an integer.

For smooth manifolds of type III, we do not know a good invariant detecting the bordism group  $\Omega_4^{\text{Pin}^+}$ . Therefore we could only determine the diffeomorphism type up to an ambiguity of order 2. This is based on the following exact sequence (see [10, §5])

$$0 \rightarrow \mathbb{Z}/2 \rightarrow \Omega_4^{\text{Pin}^+} \xrightarrow{\cap w_1^2} \Omega_2^{\text{Pin}^-} \rightarrow 0,$$

where  $\cap w_1^2$  is the operation of taking a submanifold dual to  $w_1^2$ . The generators of  $\Omega_2^{\text{Pin}^-}$  is  $\pm\mathbb{R}P^2$  and  $\cap w_1^2$  maps  $\pm\mathbb{R}P^4$  to  $\pm\mathbb{R}P^2$ . The image of  $[P]$  in  $\Omega_2^{\text{Pin}^-}$  can be determined from the data of the circle bundle.

In the topological case, we have an epimorphism (see [10, §9])

$$\Omega_4^{\text{TopPin}^+} \rightarrow \Omega_2^{\text{TopPin}^-} \cong \mathbb{Z}/8,$$

which is an isomorphism on the subgroup generated by  $\mathbb{R}P^4$ . By Lemma 4.4, we have  $KS(P) = KS(X)$ . Therefore by Theorem 3.4, we have a complete topological classification.

**Theorem 4.7** (type III). *Let  $X$  be a closed, simply-connected topological 4-manifold,  $\xi: S^1 \hookrightarrow M^5 \rightarrow X$  be a circle bundle over  $X$  with  $c_1(\xi) = 2 \cdot \text{primitive}$ , and  $w_2(X) \equiv c_1(\tilde{\xi}) \pmod{2}$ . Then we have*

- (1) *if  $X$  is smooth, then the diffeomorphism type of  $M$  (with the induced smooth structure) is determined up to an ambiguity of order 2 by  $\text{rank } H_2(X)$  and  $\langle c_1(\tilde{\xi})^2, [X] \rangle \in (\mathbb{Z}/8)/\pm = \{0, 1, 2, 3, 4\}$ .*
- (2)  *$M$  is homeomorphic to  $X^5(p, q) \#_{S^1} (\#_k S^2 \times S^2) \times S^1$ , where  $q = \langle c_1(\tilde{\xi})^2, [X] \rangle \in (\mathbb{Z}/8)/\pm = \{0, 1, 2, 3, 4\}$ ,  $k = (\text{rank } H_2(X) - (3 + (-1)^q)/2)$ ,  $p = KS(X)$ .*

*Proof.* We only need to prove (1), the proof of (2) is similar. We see from the proof of Lemma 4.3 that  $P = f^{-1}(\mathbb{R}P^4)$ , where  $f: P \rightarrow \mathbb{R}P^4$  induces an isomorphism on  $\pi_1$ . If the mod 2 degree of  $f$  is 1, then the submanifold dual to  $w_1(P)$  is  $f^{-1}(\mathbb{R}P^3)$ , and the submanifold  $V$  dual to  $w_1(P)^2$  is  $f^{-1}(\mathbb{R}P^2)$ . Now we have the following commutative diagram

$$\begin{array}{ccc} S^1 & \xrightarrow{=} & S^1 \\ \downarrow & & \downarrow \\ \partial N(F) & \xrightarrow{f} & \mathbb{R}P^3 & \supset \mathbb{R}P^2 = D^2 \cup S^1 \\ \downarrow & & \downarrow \\ F & \xrightarrow{g} & \mathbb{C}P^1 & = D^2 \cup pt \end{array}$$

Let  $d = \deg g = \langle c_1(\tilde{\xi})^2, [X] \rangle$  and  $D = g^{-1}(pt) = \{p_1, \dots, p_d\}$ , it is seen that  $V = f^{-1}(\mathbb{R}P^2) = (F - D) \cup_{\partial} d \cdot S^1$  (where the gluing map is of degree 2) and  $[V] = d \cdot [\mathbb{R}P^2] \in \Omega_2^{\text{Pin}^-}$ . If the mod 2 degree of  $f$  is zero, then we consider the circle bundle over  $X \# \mathbb{C}P^2$  with first Chern class  $(c_1(\xi), 2)$ . The corresponding map has nonzero mod 2 degree, the image of the corresponding characteristic submanifold in  $\Omega_2^{\text{Pin}^-}$  equals to that of the original one plus 1. Finally  $\langle (c_1(\tilde{\xi}), 1)^2, [X \# \mathbb{C}P^2] \rangle = \langle c_1(\tilde{\xi})^2, [X] \rangle + 1$ . This proves the theorem.  $\square$

For the manifolds of type I, we have

**Theorem 4.8** (type I). *Let  $X$  be a closed, simply-connected non-spin topological 4-manifold,  $\xi: S^1 \hookrightarrow M^5 \rightarrow X$  be a circle bundle over  $X$  with  $c_1(\xi) = 2 \cdot \text{primitive}$ , and  $w_2(X) \not\equiv c_1(\tilde{\xi}) \pmod{2}$ . We have*

- (1) if  $KS(X) = 0$ , then  $M$  is smoothable and
- if  $\langle w_2(X)^2, [X] \rangle \equiv \langle c_1(\tilde{\xi})^2, [X] \rangle \pmod{2}$ , then  $M$  is diffeomorphic to  $X^5(q) \#_{S^1}(S^2 \times \mathbb{R}P^3) \#_{S^1}((\#_k S^2 \times S^2) \times S^1)$ ,  
where  $q = \langle c_1(\tilde{\xi})^2, [X] \rangle \in (\mathbb{Z}/8)/\pm = \{0, 1, 2, 3, 4\}$  and  $k = \frac{1}{2}(\text{rank } H_2(X) - \frac{1}{2}(7 + (-1)^q))$ ;
  - if  $\langle w_2(X)^2, [X] \rangle \not\equiv \langle c_1(\tilde{\xi})^2, [X] \rangle \pmod{2}$ , then  $M$  is diffeomorphic to  $X^5(q) \#_{S^1}(\mathbb{C}P^2 \times S^1) \#_{S^1}((\#_k S^2 \times S^2) \times S^1)$ ,  
where  $q = \langle c_1(\tilde{\xi})^2, [X] \rangle \in (\mathbb{Z}/8)/\pm = \{0, 1, 2, 3, 4\}$  and  $k = \frac{1}{2}(\text{rank } H_2(X) - \frac{1}{2}(5 + (-1)^q))$ .
- (2) if  $KS(X) = 1$ , then  $M$  is non-smoothable and
- if  $\langle w_2(X)^2, [X] \rangle \equiv \langle c_1(\tilde{\xi})^2, [X] \rangle \pmod{2}$ , then  $M$  is homeomorphic to  $X^5(1, q) \#_{S^1}(S^2 \times \mathbb{R}P^3) \#_{S^1}((\#_k S^2 \times S^2) \times S^1)$ ,  
where  $q = \langle c_1(\tilde{\xi})^2, [X] \rangle \in (\mathbb{Z}/8)/\pm = \{0, 1, 2, 3, 4\}$  and  $k = \frac{1}{2}(\text{rank } H_2(X) - \frac{1}{2}(7 + (-1)^q))$ ;
  - if  $\langle w_2(X)^2, [X] \rangle \not\equiv \langle c_1(\tilde{\xi})^2, [X] \rangle \pmod{2}$ , then  $M$  is homeomorphic to  $X^5(1, q) \#_{S^1}(\mathbb{C}P^2 \times S^1) \#_{S^1}((\#_k S^2 \times S^2) \times S^1)$ ,  
where  $q = \langle c_1(\tilde{\xi})^2, [X] \rangle \in (\mathbb{Z}/8)/\pm = \{0, 1, 2, 3, 4\}$  and  $k = \frac{1}{2}(\text{rank } H_2(X) - \frac{1}{2}(5 + (-1)^q))$ .

**Remark 4.9.** Note that for 4-manifold  $X$ ,  $\langle w_2(X)^2, [X] \rangle \equiv \text{rank } H_2(X) \pmod{2}$ . This ensures that  $k$  is an integer.

*Proof.* We only need to prove (1), the proof of (2) is similar. Recall that we have  $\Omega_4^{\text{Pin}^c} \cong \mathbb{Z}/8 \oplus \mathbb{Z}/2$ , with generators  $\mathbb{R}P^4$  and  $\mathbb{C}P^2$ . Thus the  $q$ -component is determined as in the type III case. The  $s$ -component of  $P$  is determined by the bordism number  $\langle w_2(P)^2, [P] \rangle \in \mathbb{Z}/2$ . (Here we use the notations given before Theorem 3.5.) Since  $KS(X) = 0$ , there exists an integer  $m$  such that  $X_0 = X \# m(S^2 \times S^2)$  is smooth. Note that if we do the same construction on  $X_0$  we get  $P_0 = P \# m(S^2 \times S^2)$ , and  $\langle w_2(P_0)^2, [P_0] \rangle = \langle w_2(P)^2, [P] \rangle$ . Therefore, to compute the  $s$ -component, we may assume that  $X$  is smooth. Recall that  $P = (X - \hat{N}(F)) \cup_{\partial} B$ , it is seen that the bordism class of  $P$  is determined by the bordism class of the pair  $(X, F)$ , which can be viewed as a singular manifold  $(X, f) \in \Omega_4(BU(1)) \cong \Omega_4 \oplus H_4(BU(1))$ . We have two homomorphisms

$$\Omega_4(BU(1)) \rightarrow \mathbb{Z}/2, \quad [X, F] \mapsto \langle w_2(P)^2, [P] \rangle$$

and

$$\Omega_4(BU(1)) \rightarrow \mathbb{Z}/2, \quad [X, c_1(\tilde{\xi})] \mapsto \langle w_2(X)^2 + c_1(\tilde{\xi})^2, [X] \rangle.$$

By a check on the generators  $(\mathbb{C}P^2 \sharp (S^2 \times S^2), c_1(\tilde{\xi}) = (1, 0, 1))$  and  $(\mathbb{C}P^2 \sharp (S^2 \times S^2), c_1(\tilde{\xi}) = (0, 0, 1))$ , we see that  $s = \langle w_2(P)^2, [P] \rangle = \langle w_2(X)^2 + c_1(\tilde{\xi})^2, [X] \rangle \pmod{2}$ . The two cases correspond to the values  $s = 0$  and  $s = 1$ . For the proof of (2), the only change is that  $\Omega_4^{\text{Top}} \cong \mathbb{Z} \oplus \mathbb{Z}/2$  with generators  $\mathbb{C}P^2$  and  $*\mathbb{C}P^2 \sharp \overline{\mathbb{C}P^2}$  [8].  $\square$

## 5. BORDISM AND SURGERY

The main tool used in the solution of the classification problem is the modified surgery developed in [11]. We first recall some basic definitions and facts. Let  $p: B \rightarrow BO$  be a fibration,  $\bar{\nu}: M^5 \rightarrow B$  be a lift of the normal Gauss map  $\nu: M \rightarrow BO$  ( $\bar{\nu}$  is called a normal  $B$ -structure of  $M$ ).  $\bar{\nu}$  is called a normal 2-smoothing if it is a 3-equivalence. Manifolds with normal  $B$ -structures form a bordism theory. Suppose  $(M_i^5, \bar{\nu}_i)$  ( $i = 1, 2$ ) are two normal 2-smoothings in  $B$ ,  $(W^6, \bar{\nu})$  is a  $B$ -bordism between  $(M_1^5, \bar{\nu}_1)$  and  $(M_2^5, \bar{\nu}_2)$ . Then  $W$  is bordant rel. boundary to an  $s$ -cobordism (implying that  $M_1$  and  $M_2$  are diffeomorphic) if and only if an obstruction  $\theta(W, \bar{\nu}) \in L_6(\pi_1(B))$  is zero [11, p.730].

The obstruction group  $L_6(\pi_1)$  is related to the ordinary Wall's  $L$ -group in the following exact sequence

$$0 \rightarrow L_6^s(\pi_1) \rightarrow L_6(\pi_1) \rightarrow \text{Wh}(\pi_1),$$

where  $L_6^s(\pi_1)$  is the Wall's  $L$ -group of  $\pi_1$  and  $\text{Wh}(\pi_1)$  is the Whitehead group. If  $\pi_1 = \mathbb{Z}/2$ , the map  $L_6^s(\pi_1) \rightarrow L_6(\pi_1)$  is an isomorphism since  $\text{Wh}(\mathbb{Z}/2) = 0$  ([14]). The elements in  $L_6^s(\mathbb{Z}/2)$  are detected by the Kervaire-Arf invariant. If  $\theta(W, \bar{\nu})$  is nonzero in  $L_6^s(\pi_1)$ , then one can do surgery on  $(W, \bar{\nu})$  such that the result manifold  $(W', \bar{\nu}')$  has trivial surgery obstruction [18]. Therefore we have proved the following

**Proposition 5.1.** *Two smooth 5-manifolds  $M_1$  and  $M_2$  with fundamental group  $\mathbb{Z}/2$  are diffeomorphic if they have bordant normal 2-smoothings in some fibration  $B$ .*

The fibration  $B$  is called the normal 2-type of  $M$  if  $p$  is 3-coconnected. This is an invariant of  $M$ . Because of this proposition, the solution to the classification problem consists of two steps: first determine the normal 2-types  $B$  for the 5-manifolds under consideration and then determine invariants of the corresponding bordism groups  $\Omega_5^{(B,p)}$ .

**§5A. Normal 2-types.** Let  $M^5$  be a fibered type 5-manifold. The universal coefficient theorem implies that  $H_2(\widetilde{M}) \otimes_{\mathbb{Z}[\pi_1]} \mathbb{Z} \rightarrow H_2(M)$  is an isomorphism. Since the  $\pi_1(M)$ -action on  $H_2(\widetilde{M})$  is trivial, we have  $H_2(\widetilde{M}) \otimes_{\mathbb{Z}[\pi_1]} \mathbb{Z} = H_2(\widetilde{M})$ , therefore  $H_2(\widetilde{M}) \rightarrow H_2(M)$  is an isomorphism, also is the second Hurewicz map  $\pi_2(M) \rightarrow H_2(M)$ . Now suppose  $\pi_1(M) \cong \mathbb{Z}/2$  and  $H_2(M) \cong \mathbb{Z}^r$ .

We start with the description of the normal 2-types for type II manifolds. It is the simplest situation and illuminates the ideas.

Type II: consider the fibration

$$p: B = \mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r \times B\text{Spin} \rightarrow BO,$$

where  $p: B \rightarrow BO$  is trivial on the first two factors and on  $B\text{Spin}$  it is the canonical projection from  $B\text{Spin}$  onto  $BO$ . A lift  $\bar{\nu} \rightarrow B$  is given as follows: the map to  $\mathbb{R}P^\infty$  is the classifying map of the fundamental group; choose a basis  $\{u_1, \dots, u_r\}$  of the free part of  $H^2(M) \cong \mathbb{Z}^r \oplus \mathbb{Z}/2$ , by realizing each element  $u_i$  by a map to  $\mathbb{C}P^\infty$  we get a map to

$(\mathbb{C}P^\infty)^r$ ; a Spin-structure on  $\nu M$  gives rise to a map to  $B\text{Spin}$ . It's easy to see that  $(B, p)$  is the normal 2-type of type II manifolds and that  $\bar{\nu}$  induces an isomorphism on  $\pi_1$  and  $H_2$ . Since the second Hurewicz maps  $\pi_2(M) \rightarrow H_2(M)$  and  $\pi_2((\mathbb{C}P^\infty)^r) \rightarrow H_2((\mathbb{C}P^\infty)^r)$  are isomorphisms,  $\bar{\nu}$  is a normal 2-smoothing.

Type III: let  $\eta$  be the canonical real line bundle over  $\mathbb{R}P^\infty$ ,  $2\eta = \eta \oplus \eta$ . Consider the fibration

$$p: B = \mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r \times B\text{Spin} \xrightarrow{p_1 \times p_2} BO \times BO \xrightarrow{\oplus} BO,$$

where  $p_1: \mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r \rightarrow BO$  is the classifying map of  $\pi^*(2\eta)$ , (where  $\pi: \mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r \rightarrow \mathbb{R}P^\infty$  is the projection map,)  $p_2: B\text{Spin} \rightarrow BO$  is the canonical projection and  $\oplus: BO \times BO \rightarrow BO$  is the  $H$ -space structure on  $BO$  induced by the Whitney sum of vector bundles. A lift  $\bar{\nu} \rightarrow B$  is given as follows: the map to  $\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r$  is the same as in type II. Since  $w_2(2\eta) = w_1(\eta)^2$  is the nonzero element in  $\text{Ext}(H_1(\mathbb{R}P^\infty), \mathbb{Z}/2)$  and  $w_2(M)$  is the nonzero element in  $\text{Ext}(H_1(M), \mathbb{Z}/2)$ , we have  $w_2(\bar{\nu}^*2\eta) = w_2(\nu M)$ . This implies that  $\nu M - \bar{\nu}^*2\eta$  admits a Spin-structure. Such a structure induces a map to  $B\text{Spin}$ . Then  $\bar{\nu}$  is a lift of  $\nu$ . It is easy to see that  $(B, p)$  is the normal 2-type of type III manifolds and  $\bar{\nu}$  is a normal 2-smoothing.

Type I: let  $\gamma$  be the canonical complex line bundle over  $\mathbb{C}P^\infty$ . Consider the fibration

$$p: B = \mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r \times B\text{Spin} \xrightarrow{p_1 \times p_2} BO \times BO \xrightarrow{\oplus} BO,$$

where  $p_1: \mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r \rightarrow BO$  is the classifying map of  $\pi^*\gamma$ ,  $\pi: \mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r \rightarrow \mathbb{C}P^\infty$  is the projection map to the first  $\mathbb{C}P^\infty$ . A lift  $\bar{\nu} \rightarrow B$  is given as follows: since the Bockstein homomorphism  $\beta: H^2(M; \mathbb{Z}/2) \rightarrow H^3(M; \mathbb{Z})$  is trivial,  $w_2(M)$  is the mod 2 reduction of an integral cohomology class. Since  $w_2(M)$  is not contained in  $\text{Ext}(H_1(M), \mathbb{Z}/2)$ , this integral cohomology class can be taken as a primitive one, say,  $u_1$  and we extend it to a basis  $\{u_1, \dots, u_r\}$ . Then the map to  $\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r$  is the same as above. Now  $\nu M - \bar{\nu}^*\gamma$  admits a Spin-structure, this gives rise to a map  $M \rightarrow B\text{Spin}$ . Then  $\bar{\nu}$  is a lift of  $\nu$ . It is easy to see that  $(B, p)$  is the normal 2-type of type I manifolds and  $\bar{\nu}$  is a normal 2-smoothing.


**§5B. Computation of the bordism groups.** As in the last subsection, we start with the type II manifolds, which is the simplest case.

Type II: recall that the normal 2-type is

$$p: B = \mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r \times B\text{Spin} \rightarrow BO,$$

where  $p: B \rightarrow BO$  is trivial on the first two factors and is the canonical projection from  $B\text{Spin}$  onto  $BO$ . Therefore the bordism group  $\Omega_5^{(B,p)}$  is the Spin-bordism group  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r)$ . To compute this bordism group, we apply the Atiyah-Hirzebruch spectral sequence. The  $E^2$ -terms are  $E_{p,q}^2 = H_p(\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r; \Omega_q^{\text{Spin}})$ .

To illuminate the situation, we first consider the group  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times \mathbb{C}P^\infty)$ . The relevant terms and differentials in the spectral sequence are depicted as follows:



The  $E^2$ -terms are:

- $E_{1,4}^2 = H_1(\mathbb{R}P^\infty \times \mathbb{C}P^\infty) \cong \mathbb{Z}/2$ ,
- $E_{2,2}^2 = H_2(\mathbb{R}P^\infty \times \mathbb{C}P^\infty; \mathbb{Z}/2) \cong \mathbb{Z}/2 \oplus \mathbb{Z}/2$ ,
- $E_{3,1}^2 = E_{3,2}^2 = H_3(\mathbb{R}P^\infty \times \mathbb{C}P^\infty; \mathbb{Z}/2) \cong \mathbb{Z}/2 \oplus \mathbb{Z}/2$ ,
- $E_{4,1}^2 = E_{4,2}^2 = H_4(\mathbb{R}P^\infty \times \mathbb{C}P^\infty; \mathbb{Z}/2) \cong (\mathbb{Z}/2)^3$ ,
- $E_{5,0}^2 = H_5(\mathbb{R}P^\infty \times \mathbb{C}P^\infty) \cong (\mathbb{Z}/2)^3$ ,
- $E_{5,1}^2 = H_5(\mathbb{R}P^\infty \times \mathbb{C}P^\infty; \mathbb{Z}/2) \cong (\mathbb{Z}/2)^3$ ,
- $E_{6,0}^2 = H_6(\mathbb{R}P^\infty \times \mathbb{C}P^\infty) \cong \mathbb{Z}/2$ .

The differentials  $d_2$  are dual to the Steenrod square  $Sq^2$ . On the  $E^3$ -page, by comparison with the Atiyah-Hirzebruch spectral sequence of  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty)$ , the differential  $d_3: E_{4,2}^3 \rightarrow E_{1,4}^3$  is nontrivial: it is computed in [10] that  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty)$  is zero (see Remark 5.5 below), therefore the  $E_{1,4}^3$ -term must be killed. Therefore on the  $E^\infty$ -page, on the line  $p+q=5$ , the nontrivial terms are  $E_{5,0}^\infty = H_3(\mathbb{R}P^\infty) \otimes H_2(\mathbb{C}P^\infty) \cong \mathbb{Z}/2$  and  $E_{4,1}^\infty = H_2(\mathbb{R}P^\infty; \mathbb{Z}/2) \otimes H_2(\mathbb{C}P^\infty; \mathbb{Z}/2) \cong \mathbb{Z}/2$ .

To state the result of our calculation, let  $\alpha \in H^1(\mathbb{R}P^\infty; \mathbb{Z}/2)$ ,  $\beta \in H^2(\mathbb{C}P^\infty; \mathbb{Z}/2)$  denote the nonzero elements, and let  $\tau: \mathbb{C}P^\infty \rightarrow \mathbb{C}P^\infty$  be the involution on  $\mathbb{C}P^\infty$  with  $\tau_* = -1$  on  $H_2(\mathbb{C}P^\infty)$ .

**Lemma 5.2.** *The short exact sequence*

$$0 \rightarrow \mathbb{Z}/2 \rightarrow \Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times \mathbb{C}P^\infty) \rightarrow \mathbb{Z}/2 \rightarrow 0$$

is nonsplit, and  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times \mathbb{C}P^\infty) \cong \mathbb{Z}/4$ . A generating bordism class  $[X^5, f]$  is detected by the invariant  $\langle \alpha^3 \cup \beta, f_*[X] \rangle \in \mathbb{Z}/2$ . Furthermore, we have the relation  $\langle \alpha \cup \beta^2, f_*[X] \rangle = 0$ , and  $[X, (\text{id}_{\mathbb{R}P^\infty} \times \tau) \circ f] = -[X, f]$ .

*Proof.* There is a product map

$$\varphi: \Omega_3^{\text{Spin}}(\mathbb{R}P^\infty) \otimes \Omega_2^{\text{Spin}}(\mathbb{C}P^\infty) \rightarrow \Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times \mathbb{C}P^\infty),$$



induced by the product of manifolds. There is a corresponding product map on the Atiyah-Hirzebruch spectral sequences

$$\Phi: E_{p,q}^r(1) \otimes E_{s,t}^r(2) \rightarrow E_{p+s,q+t}^r(3),$$

where on the  $E^\infty$ -page  $\Phi$  is compatible with the filtrations on the bordism groups and on the  $E^2$ -page it is just the cross product map (see [17, p. 352]). It is easy to see that the Atiyah-Hirzebruch spectral sequence of  $\Omega_3^{\text{Spin}}(\mathbb{R}P^\infty)$  collapses on the line  $p+q=3$ , the Atiyah-Hirzebruch spectral sequence of  $\Omega_2^{\text{Spin}}(\mathbb{C}P^\infty)$  collapses on the line  $p+q=2$ , and  $\Phi$  is surjective on  $E_{5,0}^\infty(3)$  and  $E_{4,1}^\infty(3)$ . Therefore  $\varphi$  is surjective. Now

$$\Omega_3^{\text{Spin}}(\mathbb{R}P^\infty) \cong \Omega_2^{\text{Pin}^-} \cong \mathbb{Z}/8,$$

generated by  $[\mathbb{R}P^3, \text{inclusion}]$  (see [10]) and  $\Omega_2^{\text{Spin}}(\mathbb{C}P^\infty) \cong \Omega_2^{\text{Spin}} \oplus H_2(\mathbb{C}P^\infty)$ . The group  $\Omega_2^{\text{Spin}}$  is generated by  $T^2$  with the Lie group spin structure. The product  $\varphi(\mathbb{R}P^3, T^2) = 0$ , since the map  $\mathbb{R}P^3 \times T^2 \rightarrow \mathbb{R}P^\infty \times \mathbb{C}P^\infty$  factors through  $\mathbb{R}P^\infty$  and  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty) = 0$ . Therefore we have a surjection

$$\mathbb{Z}/8 \otimes \mathbb{Z} \rightarrow \Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times \mathbb{C}P^\infty).$$

This shows that  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times \mathbb{C}P^\infty) \cong \mathbb{Z}/4$  and  $[X, (\text{id}_{\mathbb{R}P^\infty} \times \tau) \circ f] = -[X, f]$ . The relation comes from the fact that the dual of  $d_2$  maps  $\alpha\beta$  to  $\alpha\beta^2$ .  $\square$

In general, in the Atiyah-Hirzebruch spectral sequence for  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r)$ , the nontrivial terms for the line  $p+q=5$  on the  $E^\infty$ -page are

$$E_{5,0}^\infty = \oplus_i H_3(\mathbb{R}P^\infty) \otimes H_2(\mathbb{C}P_i^\infty) \oplus \oplus_{i \neq j} H_1(\mathbb{R}P^\infty) \otimes H_2(\mathbb{C}P_i^\infty) \otimes H_2(\mathbb{C}P_j^\infty) \cong (\mathbb{Z}/2)^{r+r(r-1)/2}$$

and

$$E_{4,1}^\infty = \oplus_i H_2(\mathbb{R}P^\infty; \mathbb{Z}/2) \otimes H_2(\mathbb{C}P_i^\infty; \mathbb{Z}/2) \cong (\mathbb{Z}/2)^r.$$

Using the same argument as in Lemma 5.2, we have the following

**Proposition 5.3** (type II). *The bordism group  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r)$  is isomorphic to  $(\mathbb{Z}/4)^r \oplus (\mathbb{Z}/2)^{r(r-1)/2}$ . Let  $\alpha \in H^1(\mathbb{R}P^\infty; \mathbb{Z}/2)$ ,  $\beta_i \in H^2(\mathbb{C}P_i^\infty; \mathbb{Z}/2)$  be the nonzero elements, then*

- (1) *the  $\mathbb{Z}/2$ -factors are determined by the invariants  $\langle \alpha \cup \beta_i \cup \beta_j, f_*[X] \rangle \in \mathbb{Z}/2$ , with  $i, j = 1, \dots, r$ , and  $i > j$ ,*
- (2) *the bordism classes  $\{\pm 1\} \subset \mathbb{Z}/4$  are detected by the invariants  $\langle \alpha^3 \cup \beta_i, f_*[X] \rangle \in \mathbb{Z}/2$ , with  $i = 1, \dots, r$ , and*
- (3) *the bordism classes  $\{\pm 1\} \subset \mathbb{Z}/4$  are interchanged if we compose  $f$  with the involution  $\tau_i$  on  $\mathbb{C}P_i^\infty$ .*
- (4) *Furthermore we have the relation  $\langle \alpha \cup \beta_i^2, f_*[X] \rangle = 0$  for all  $i$ .*

Type III: the normal 2-type is

$$p: B = \mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r \times B\text{Spin} \rightarrow BO,$$

where the map on  $\mathbb{R}P^\infty$  is the classifying map of the vector bundle  $2\eta$ . Therefore the bordism group  $\Omega_5^{(B,p)}$  is the twisted Spin-bordism group

$$\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r; \pi^*2\eta) = \tilde{\Omega}_7^{\text{Spin}}(\text{Th}(\pi^*2\eta)).$$

In the Atiyah-Hirzebruch spectral sequence, the  $E^2$ -terms are

$$E_{p,q}^2 = H_p(\mathrm{Th}(\pi^*2\eta); \Omega_q^{\mathrm{Spin}}).$$

Since  $2\eta$  is orientable, we may apply the Thom isomorphism and after a degree shift  $p \mapsto p - 2$  we have  $E_{p,q}^2 = H_p(\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r; \Omega_q^{\mathrm{Spin}})$ . Therefore the  $E^2$ -terms are the same as in the type II case, where the differentials  $d_2$  are dual to  $Sq^2 + w_2(2\eta)$ . The differential  $d_3: E_{4,2}^3 \rightarrow E_{1,4}^3$  is determined by the comparison with the Atiyah-Hirzebruch spectral sequence of  $\Omega_5^{\mathrm{Spin}}(\mathbb{R}P^\infty; 2\eta)$  as follows:

The following construction is given in [6, §5]. Let  $[X, f] \in \Omega_5^{\mathrm{Spin}}(\mathbb{R}P^\infty; 2\eta)$  be a singular manifold, where by doing 1-surgeries we can assume that  $f$  is an isomorphism on  $\pi_1$ . If  $P$  be the characteristic submanifold of  $X$ , then  $P$  admits a  $\mathrm{Pin}^+$ -structure, which is induced from the  $\mathrm{Spin}$ -structure on  $\nu X \oplus f^*(2\eta)$ . The bordism class of  $P$  depends only on the bordism class of  $[X, f]$ , and therefore we have a homomorphism  $\Omega_5^{\mathrm{Spin}}(\mathbb{R}P^\infty; 2\eta) \rightarrow \Omega_4^{\mathrm{Pin}^+}$ .

**Lemma 5.4.** *The homomorphism  $\Omega_5^{\mathrm{Spin}}(\mathbb{R}P^\infty; 2\eta) \rightarrow \Omega_4^{\mathrm{Pin}^+}$  is an isomorphism.*

*Proof.* The injectivity is ensured by [6, Lemma 10]. The generator of  $\Omega_4^{\mathrm{Pin}^+}$  is  $\mathbb{R}P^4$ , which is the image of  $(\mathbb{R}P^5, i)$ . This proves the surjectivity.  $\square$

**Remark 5.5.** Similarly it can be shown that the homomorphism  $\Omega_5^{\mathrm{Spin}}(\mathbb{R}P^\infty) \rightarrow \Omega_4^{\mathrm{Pin}^-}$  is also an isomorphism, where both groups are 0.

Now we know  $\Omega_5^{\mathrm{Spin}}(\mathbb{R}P^\infty; 2\eta) \cong \Omega_4^{\mathrm{Pin}^+} \cong \mathbb{Z}/16$ , and therefore the differential  $d_3: E_{4,2}^3 \rightarrow E_{1,4}^3$  must be trivial. Therefore we have

**Proposition 5.6** (type III). *There is a short exact sequence*

$$0 \rightarrow G \rightarrow \Omega_5^{\mathrm{Spin}}(\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r; \pi^*(2\eta)) \rightarrow \Omega_4^{\mathrm{Pin}^+} \rightarrow 0,$$

where  $G \cong (\mathbb{Z}/4)^r \oplus (\mathbb{Z}/2)^{r(r-1)/2}$ . Then

- (1) the  $\mathbb{Z}/2$ -factors are determined by the invariants  $\langle \alpha \cup \beta_i \cup \beta_j, f_*[X] \rangle \in \mathbb{Z}/2$ , with  $i, j = 1, \dots, r$ , and  $i > j$ ,
- (2) the bordism classes  $\{\pm 1\} \subset \mathbb{Z}/4$  are detected by the invariants  $\langle \alpha^3 \cup \beta_i, f_*[X] \rangle \in \mathbb{Z}/2$ , with  $i = 1, \dots, r$ , and
- (3) the bordism classes  $\{\pm 1\} \subset \mathbb{Z}/4$  are interchanged if we compose  $f$  with the involution  $\tau_i$  on  $\mathbb{C}P_i^\infty$ .
- (4) Furthermore we have  $\langle \alpha \cup \beta_i^2, f_*[X] \rangle = \langle \alpha^3 \cup \beta_i, f_*[X] \rangle$  for all  $i$ .

Type I: recall that the normal 2-type is

$$p: B = \mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r \times B\mathrm{Spin} \rightarrow BO,$$

where the map  $p$  on the first  $\mathbb{C}P^\infty$  is the classifying map of the vector bundle  $\gamma$ . Therefore the bordism group  $\Omega_5^{(B,p)}$  is the twisted  $\mathrm{Spin}$ -bordism group

$$\Omega_5^{\mathrm{Spin}}(\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r; \pi^*\gamma) = \tilde{\Omega}_7^{\mathrm{Spin}}(\mathrm{Th}(\pi^*\gamma)).$$

As in the type III case we apply the Thom isomorphism and the  $E^2$ -terms in the Atiyah-Hirzebruch spectral sequence are  $E_{p,q}^2 = H_p(\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r; \Omega_q^{\mathrm{Spin}})$ , where the differentials  $d_2$  are dual to  $Sq^2 + w_2(\gamma)$ .

As in the type III case, we have a homomorphism  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times \mathbb{C}P^\infty; \pi^*\gamma) \rightarrow \Omega_4^{\text{Pin}^c}$ , mapping  $[X, f]$  to its characteristic submanifold.

**Lemma 5.7.** *The homomorphism  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times \mathbb{C}P^\infty; \pi^*\gamma) \rightarrow \Omega_4^{\text{Pin}^c}$  is an isomorphism.*

*Proof.* The proof of the injectivity is analogous to the argument in [6, Lemma 10]: suppose that  $P$  and  $P'$  are bordant in  $\Omega_4^{\text{Pin}^c}$ , then  $\widetilde{P}$  and  $\widetilde{P}'$  are bordant in  $\Omega_4^{\text{Spin}^c}$ . Let  $V^5$  be such a bordism, then  $A \cup_{\widetilde{P}} V \cup_{\widetilde{P}'} A'$  together with the reference map is an element in  $\Omega_5^{\text{Spin}}(\mathbb{C}P^\infty; \gamma)$ , where the reference maps on each part to  $\mathbb{C}P^\infty$  are induced from the  $\text{Spin}^c$ -structures. An easy computation with the Atiyah-Hirzebruch spectral sequence shows that  $\Omega_5^{\text{Spin}}(\mathbb{C}P^\infty; \gamma) = 0$ , and what follows are exactly the same as in the proof of [6, Lemma 10]. The Atiyah-Hirzebruch spectral sequence shows that the order of  $\Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times \mathbb{C}P^\infty; \pi^*\gamma)$  is 16, and we have shown that  $\Omega_4^{\text{Pin}^c} \cong \mathbb{Z}/8 \oplus \mathbb{Z}/2$ . This proves the surjectivity.  $\square$

**Proposition 5.8** (type I). *There is a short exact sequence*

$$0 \rightarrow G \rightarrow \Omega_5^{\text{Spin}}(\mathbb{R}P^\infty \times (\mathbb{C}P^\infty)^r; \gamma) \rightarrow \Omega_4^{\text{Pin}^c} \rightarrow 0,$$

where  $G \cong (\mathbb{Z}/4)^{r-1} \oplus (\mathbb{Z}/2)^{r(r-1)/2}$ . Then

- (1) the  $\mathbb{Z}/2$ -factors are determined by the invariants  $\langle \alpha \cup \beta_i \cup \beta_j, f_*[X] \rangle \in \mathbb{Z}/2$ , with  $i, j = 1, \dots, r$ , and  $i > j$ ,
- (2) the bordism classes  $\{\pm 1\} \subset \mathbb{Z}/4$  are detected by the invariants  $\langle \alpha^3 \cup \beta_i, f_*[X] \rangle \in \mathbb{Z}/2$ , with  $i = 1, \dots, r$ , and
- (3) the bordism classes  $\{\pm 1\} \subset \mathbb{Z}/4$  are interchanged if we compose  $f$  with the involution  $\tau_i$  on  $\mathbb{C}P_i^\infty$ .
- (4) We have  $\langle \alpha^5 + \alpha^3 \cup \beta_1, f_*[X] \rangle = 0$  and  $\langle \alpha \cup \beta_i^2, f_*[X] \rangle = \langle \alpha \cup \beta_1 \cup \beta_i, f_*[X] \rangle$  for all  $i$ .

## 6. PROOFS OF THE MAIN RESULTS

Now that the bordism groups are determined, we can prove the main results of the paper. First we need a lemma for the classification theorem.

**Lemma 6.1.** *Let  $M^5$  be a fibered type manifold with  $\pi_1(M) \cong \mathbb{Z}/2$  and  $H_2(M) \cong \mathbb{Z}^r$ . Let  $t \in H^1(M; \mathbb{Z}/2)$  be the nonzero element, and let  $\{t^2, x_1, \dots, x_r\}$  be a basis of  $H^2(M; \mathbb{Z}/2)$ . Then  $\{t^3, tx_1, \dots, tx_r\}$  is a basis of  $H^3(M; \mathbb{Z}/2)$ .*

*Proof.* Consider the Leray-Serre cohomology spectral sequence for the fibration  $\widetilde{M} \rightarrow M \rightarrow \mathbb{R}P^\infty$  with  $\mathbb{Z}/2$ -coefficient. Note that  $\dim H^2(M; \mathbb{Z}/2) = r+1$  and  $\dim H^2(\widetilde{M}; \mathbb{Z}/2) = r$ . This implies that the differential

$$d_2: E_2^{0,2} = H^2(\widetilde{M}; \mathbb{Z}/2) \rightarrow E_3^{3,0} = H^3(\mathbb{R}P^\infty; \mathbb{Z}/2)$$

must be trivial. Therefore, the elements  $t^3, tx_1, \dots, tx_r$  all survive to form a basis of  $H^3(M; \mathbb{Z}/2)$ .  $\square$

The proof of Theorem 3.1.

Type II: let  $\bar{\nu}: M^5 \rightarrow B$  be a normal 2-smoothing in  $B$ . Let  $t = \bar{\nu}^*\alpha$ ,  $x_i = \bar{\nu}^*\beta_i$ . Then  $\{t^2, x_1, \dots, x_r\}$  is a basis of  $H^2(M; \mathbb{Z}/2)$ . Consider the symmetric bilinear form

$$\lambda: H^2(M; \mathbb{Z}/2) \times H^2(M; \mathbb{Z}/2) \xrightarrow{\cup} H^4(M; \mathbb{Z}/2) \xrightarrow{\cup t} H^5(M; \mathbb{Z}/2) \cong \mathbb{Z}/2.$$

By Poincaré duality and Lemma 6.1,  $\lambda$  is nondegenerate. Since  $\lambda(t^2, t^2) = t^5 = 0$  and  $\lambda(x_i, x_i) = tx_i^2 = 0$  (the relation in Proposition 5.3), we can find a symplectic basis of  $\lambda$ ,  $\{t^2, u_1, \dots, u_r\}$ , such that  $\lambda(t^2, u_1) = t^3u_1 = 1$ ,  $\lambda(t^2, u_i) = t^3u_i = 0$  for  $i > 1$ , and  $\lambda(u_i, u_j) = tu_iu_j = 0$  or 1. Make a basis change by letting  $u'_i = u_i + u_1$  for  $i > 1$ , then  $t^3u'_i = 1$  for all  $i$  and  $tu'_iu'_j = tu_iu_j$ .  $\{u'_1, \dots, u'_r\}$  lift to a basis of the free part of  $H^2(M)$ , which gives a normal 2-smoothing. By Proposition 5.3, the bordism class of this normal 2-smoothing is independent on  $M$ . Therefore all these manifolds are  $B$ -bordant under appropriate normal 2-smoothings. Proposition 5.1 implies that they are diffeomorphic.

Type III: first note that by the relation in Proposition 5.6, for all  $v \in H^2(M; \mathbb{Z}/2)$ ,  $\lambda(t^2, v) = \lambda(v, v)$ . There are two different cases:

- (1)  $\lambda(t^2, t^2) = 0$ : then there exists a  $u_1$  such that  $\lambda(t^2, u_1) = t^3u_1 = 1$ . On the orthogonal complement of  $\text{span}(t^2, u_1)$ ,  $\lambda(v, v) = 0$ , thus there exists a symplectic basis  $\{u_2, \dots, u_r\}$ . Then the argument is the same as in the previous case.
- (2)  $\lambda(t^2, t^2) = 1$ : if  $\lambda(t^2, x_i) = 1$ , then let  $x'_i = x_i + t^2$  and otherwise let  $x'_i = x_i$ . Then  $\lambda(t^2, x'_i) = \lambda(x'_i, x'_i) = 0$ .  $\{x'_1, \dots, x'_r\}$  is a basis of the orthogonal complement of  $\text{span}(t^2)$ . There exists a symplectic basis on the complement,  $\{u_1, \dots, u_r\}$ , such that  $t^2u_i = 0$  and  $tu_iu_j = 0$  or 1. Let  $u'_i = u_i + t^2$ , then  $t^3u'_i = 1$  for all  $i$  and  $tu'_iu'_j = tu_iu_j + 1$ . The remaining argument is the same as in the previous case.

Therefore all the manifolds with the same  $[P] \in \Omega_4^{\text{Pin}^+} / \pm$  are  $B$ -bordant under appropriate normal 2-smoothings. Proposition 5.1 implies that they are diffeomorphic.

Type I: by the relation in Proposition 5.8,  $\lambda(x_1, x_i) = \lambda(x_i, x_i)$  and  $\lambda(t^2, t^2) = \lambda(t^2, x_1)$ . We have four cases:

- (1)  $\lambda(t^2, t^2) = 1$  and  $\lambda(x_1, x_1) = 0$ : then  $\lambda$  is nondegenerate on  $\text{span}(t^2, x_1)$ , and for all  $v \notin \text{span}(t^2, x_1)$ ,  $\lambda(v, v) = \lambda(v, x_1) = 0$ . Especially on the orthogonal complement of  $\text{span}(t^2, x_1)$  there exists a symplectic basis  $\{u_2, \dots, u_r\}$ .
- (2)  $\lambda(t^2, t^2) = 0$  and  $\lambda(x_1, x_1) = 0$ : then exists a  $u_2$  such that  $\lambda(x_1, u_2) = 1$  and  $\lambda(t^2, u_2) = 0$ .  $\lambda$  is nondegenerate on  $\text{span}(x_1, u_2)$ . On the orthogonal complement we have  $\lambda(v, v) = \lambda(v, x_1) = 0$ . Therefore there is a symplectic basis  $\{t^2, u_3, \dots, u_r\}$  such that  $\lambda(t^2, u_3) = 1$ ,  $\lambda(t^2, u_i) = 0$  for  $i > 3$  and  $\lambda(u_i, u_j) = 0$  or 1.
- (3)  $\lambda(t^2, t^2) = 1$  and  $\lambda(x_1, x_1) = 1$ : let  $e_0 = t^2 + x_1$ , then  $\lambda(e_0, x_1) = 0$ . On the orthogonal complement of  $\text{span}(x_1)$ , there exists a symplectic basis  $\{e_0, u_2, \dots, u_r\}$  such that  $\lambda(e_0, u_2) = \lambda(t^2, u_2) = 1$ ,  $\lambda(e_0, u_i) = \lambda(t^2, u_i) = 0$  for  $i > 2$  and  $\lambda(u_i, u_j) = 0$  or 1.
- (4)  $\lambda(t^2, t^2) = 0$  and  $\lambda(x_1, x_1) = 1$ : then on the orthogonal complement of  $\text{span}(x_1)$  there is a symplectic basis  $\{t^2, u_2, \dots, u_r\}$  such that  $\lambda(t^2, u_2) = 1$ ,  $\lambda(t^2, u_i) = 0$  for  $i > 2$  and  $\lambda(u_i, u_j) = 0$  or 1.

In all these situations we may find appropriate normal 2-smoothings such that the manifolds having equal  $[P] \in \Omega_4^{\text{Pin}^c}/\pm$  are  $B$ -bordant. Proposition 5.1 implies that they are diffeomorphic.  $\square$

Although the relations among the invariants are essentially seen in the previous proof, we would like to give a more conceptual one here.

*The proof of Theorem 3.5.* We will use the semi-characteristic class defined by R. Lee in [13]. We work with  $\mathbb{Q}$ -coefficient, in this case, the semi-characteristic class of an odd dimensional manifold with a free  $\mathbb{Z}/2$ -action is a homomorphism

$$\chi_{1/2}: \Omega_5(\mathbb{Z}/2) \rightarrow L^5(\mathbb{Q}[\mathbb{Z}/2]) \cong \mathbb{Z}/2,$$

where  $\Omega_5(\mathbb{Z}/2)$  is the bordism group of closed smooth oriented manifolds with an orientation-preserving free  $\mathbb{Z}/2$ -action, and  $L^5(\mathbb{Q}[\mathbb{Z}/2])$  is the symmetric  $L$ -group of the rational group ring  $\mathbb{Q}[\mathbb{Z}/2]$ . We refer to [13] and [4] for details.

Let  $M^5$  be an oriented smooth 5-manifold with fundamental group  $\mathbb{Z}/2$ , then the semi-characteristic class  $\chi_{1/2}(\widetilde{M}; \mathbb{Q}) \in \mathbb{Z}/2$  is defined. There is a characteristic class formula [4, Theorem C]

$$\chi_{1/2}(\widetilde{M}; \mathbb{Q}) = \langle w_4(M) \cup f^*(\alpha), [M] \rangle,$$

where  $f: M \rightarrow \mathbb{RP}^\infty$  is the classifying map of the covering and  $\alpha \in H^1(\mathbb{RP}^\infty; \mathbb{Z}/2)$  is the nonzero element. On the other hand,  $\chi_{1/2}(\widetilde{M}; \mathbb{Q})$  is identified with (see [4, p.57])

$$\begin{aligned} \hat{\chi}_{1/2}(\widetilde{M}; \mathbb{Q}) &:= \dim_{\mathbb{Q}} H_0(\widetilde{M}; \mathbb{Q}) + \dim_{\mathbb{Q}} H_1(\widetilde{M}; \mathbb{Q}) + \dim_{\mathbb{Q}} H_2(\widetilde{M}; \mathbb{Q}) \pmod{2} \\ &\equiv 1 + r \pmod{2}. \end{aligned}$$

Type II: the Wu classes of  $M$  are  $v_1 = 0$  and  $v_2 = 0$  since  $w_1(M) = w_2(M) = 0$ . Therefore  $w_4(M) = Sq^2v_2 = 0$ . This means  $r$  is odd.

Type III: the Wu classes of  $M$  are  $v_1 = 0$  and  $v_2 = w_2(M) = t^2$ . Therefore  $w_4(M) = Sq^2v_2 = t^4$  and  $\langle w_4(M) \cup f^*(\alpha), [M] \rangle = \langle \alpha^5, \bar{\nu}_*[M] \rangle$ . By the Atiyah-Hirzebruch spectral sequence, there is a nonsplit exact sequence

$$0 \rightarrow \mathbb{Z}/8 \rightarrow \Omega_5^{\text{Spin}}(\mathbb{RP}^\infty; 2\eta) \rightarrow H_5(\mathbb{RP}^\infty) \rightarrow 0,$$

and therefore  $\bar{\nu}_*[M] \equiv q \pmod{2}$ . This implies  $r + q$  is odd.

Type I: the Wu classes of  $M$  are  $v_1 = 0$  and  $v_2 = w_2(M) = \bar{\nu}^*w_2(\gamma)$ . Therefore  $w_4(M) = Sq^2v_2 = \bar{\nu}^*w_2(\gamma)^2$  and  $\langle w_4(M) \cup f^*(\alpha), [M] \rangle = \langle \alpha \cup \beta^2, \bar{\nu}_*[M] \rangle$ . Check on the generators of  $\Omega_5^{\text{Spin}}(\mathbb{RP}^\infty \times \mathbb{CP}^\infty; \gamma)$ ,  $\mathbb{RP}^5 \#_{S^1}(S^2 \times \mathbb{RP}^3)$  with  $(q = 1, s = 0)$  and  $\mathbb{RP}^5 \#_{S^1}(\mathbb{CP}^2 \times S^1)$  with  $(q = 1, s = 1)$ , it is seen that  $\langle \alpha \cup \beta^2, \bar{\nu}_*[M] \rangle \equiv q + s \pmod{2}$ . This implies the relation  $q + s + r \equiv 1 \pmod{2}$ .  $\square$

*The proof of Theorem 3.6.* By the Van-Kampen theorem and the Mayer-Vietoris sequence it is easy to see that all the manifolds in the list are orientable, with fundamental group  $\mathbb{Z}/2$  and torsion free  $H_2$ , and the  $\pi_1$ -action on  $H_2$  is trivial. We only need to show that these manifolds have different invariants and realize all the possible invariants.

Type II:  $\text{rank } H_2((S^2 \times \mathbb{RP}^3) \#_{S^1}((\#_k S^2 \times S^2) \times S^1)) = 2k + 1$ .

Type III: the characteristic submanifold of  $X^5(q) \#_{S^1}((\#_k S^2 \times S^2) \times S^1)$  is just that of  $X^5(q)$ , which corresponds to  $q \in \Omega_4^{\text{Pin}^+}/\pm = \{0, \dots, 8\}$ .

Type I: similarly, the manifold  $X^5(q) \sharp_{S^1}(\mathbb{C}P^2 \times S^1) \sharp_{S^1}((\sharp_k S^2 \times S^2) \times S^1)$  has characteristic submanifold invariant  $(q, 1) \in \Omega_4^{\text{Pin}^c}/\pm$ .  $\square$

*The proof of Theorem 3.11.* Note that  $X^5(q)$  and  $X^5(p, q)$  are homotopy equivalent to  $\mathbb{R}P^5$  and the operation  $\sharp_{S^1}$  preserves homotopy equivalence. This proves the theorem for the type II and III cases. For type I manifolds, the  $s$ -component of the characteristic submanifold  $P$  is determined by  $\langle w_2(P)^2, [P] \rangle$ . Since  $w_2(P) = i^*(w_2(M) + t^2)$ ,  $\langle w_2(P)^2, [P] \rangle = \langle w_2(M)^2 \cup t + t^5, [M] \rangle$ , and this is a homotopy invariant.  $\square$

#### REFERENCES

- [1] D. Barden, *Simply-connected five-manifolds*, Ann. of Math. (2) **82** (1965), 365–385.
- [2] C. Boyer and K. Galicki, *Sasakian geometry* Oxford University Press, 2008.
- [3] K. S. Brown, *Cohomology of Groups*, Springer, 1982.
- [4] J. F. Davis and R. J. Milgram, *Semicharacteristic, bordism and free group actions*, Trans. Am. Math. Soc. **312** (1989), 55–83.
- [5] H. B. Duan and C. Liang, *Circle bundles over 4-manifolds*, Arch. Math. **85** (2005), 278–282.
- [6] H. Geiges and C. B. Thomas, *Contact topology and the structure of 5-manifolds*, Ann. Inst. Fourier **48** (1998), 1167–1188.
- [7] I. Hambleton, M. Kreck, and P. Teichner, *Nonorientable 4-manifolds with fundamental group of order 2*, Trans. Am. Math. Soc. **344** (1994), 649–665.
- [8] F. Hsu, *4-dimensional topological bordism*, Topology and its Application **26** (1987), 281–285.
- [9] R. C. Kirby and L. C. Siebenmann, *Foundational Essays on Topological Manifolds, Smoothings, and Triangulations*, Annals of Mathematics Studies **88**, Princeton, N. J., Princeton University Press and University of Tokyo Press, 1977.
- [10] R. C. Kirby and L. R. Taylor, *Pin-structures on low-dimensional manifolds*, Geometry of Low-Dimensional Manifolds. 2 (S. K. Donaldson and C. B. Thomas, eds.) London Math. Soc. Lecture Note Ser., vol. 151, Cambridge Univ. Press, London and New York, 1990.
- [11] M. Kreck, *Surgery and duality*, Ann. Math. (2) **149** (1999), 707–754.
- [12] R. Lashof and L. Taylor, *Smoothing theory and Freedman’s work on 4-manifolds*, Lecture Notes Math. 1051 (Springer, Berlin) 271–290.
- [13] R. Lee, *Semicharacteristic classes*, Topology **12** (1973), 189–199.
- [14] J. Milnor, *Whitehead torsion*, Bull. Am. Math. Soc. **72** (1966), 358–426.
- [15] F. Quinn, *Smooth structures on 4-manifolds*, Four-Manifold Theory (N. H. Durham, 1982), 473–479, Contemp. Math. 35, Amer. Math. Soc., Providence, RI, 1984.
- [16] S. Smale, *On the structure of 5-manifolds*, Ann. of Math. (2) **75** (1962), 38–46.
- [17] R. M. Switzer, *Algebraic Topology—Homology and Homotopy*, Springer, 2002.
- [18] C. T. C. Wall, *Surgery on Compact Manifolds*, Second edition, Edited and with a foreword by A. A. Ranicki, Mathematical Surveys and Monographs, 69, American Mathematical Society, Providence, RI, 1999.

DEPARTMENT OF MATHEMATICS & STATISTICS  
 MCMASTER UNIVERSITY  
 HAMILTON, ON L8S 4K1, CANADA  
 E-mail address: hambleton@mcmaster.ca

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE  
 CHINESE ACADEMY OF SCIENCES  
 BEIJING, 100190, CHINA  
 E-mail address: suyang@math.ac.cn