

# LOCAL INDEX THEOREM FOR PROJECTIVE FAMILIES

MOULAY-TAHAR BENAMEUR AND ALEXANDER GOROKHOVSKY

ABSTRACT. We give a superconnection proof of the Mathai-Melrose-Singer index theorem for the family of twisted Dirac operators [20, 21].

## CONTENTS

1. Introduction	2
2. Preliminaries	3
2.1. Cyclic homology	3
2.2. Notion of a gerbe	4
2.3. Connections on gerbes	5
2.4. Twisted cohomology	7
2.5. Twisted bundles	8
2.6. Horizontally twisted bundles	14
3. Projective families and the analytic index	15
3.1. Families of pseudodifferential operators	15
3.2. Projective families	16
3.3. Analytic index	17
3.4. Chern character of the index	18
4. Dirac operators and superconnections	20
4.1. Dirac operators	20
4.2. Superconnections and index	22
4.3. The local index theorem	25
References	26

---

1991 *Mathematics Subject Classification.* 19L47, 19M05, 19K56.

*Key words and phrases.* Key words: Gerbe, Twisted K-theory, Family index, Superconnection.  
The second author was partially supported by the NSF grant DMS-0900968.

## 1. INTRODUCTION

Let  $\pi : M \rightarrow B$  be a smooth fibration. Given a class  $\delta \in H^3(B, \mathbb{Z})$  Mathai, Melrose and Singer defined an algebra of twisted by this class (also called projective) families of pseudodifferential operators. In order to give this definition in [20] it is assumed that  $\delta$  is a torsion class. In [21] the corresponding assumption is that  $\delta = \alpha \cup \beta$ ,  $\alpha \in H^1(B, \mathbb{Z})$ ,  $\beta \in H^2(B, \mathbb{Z})$  and  $\pi^*\beta = 0$ .

There is a notion of ellipticity for such a twisted pseudodifferential family and for an elliptic operator one can then define its index as an element of the  $K$ -theory of the algebra of smoothing operators. Mathai, Melrose and Singer than prove in [20, 21] an index theorem for such elliptic families thus giving an extension of the Atiyah-Singer family index theorem [1] to the twisted case.

In this paper we give a superconnection proof of the cohomological version of this index theorem for a projective family of Dirac operators. We assume that we are given a gerbe  $\mathcal{L}$  on  $B$  with a class  $\delta = [\mathcal{L}] \in H^3(B, \mathbb{Z})$ . Our conditions on this class are apparently somewhat weaker than in [20, 21]; namely we assume only that  $\pi^*\delta$  is a torsion element in  $H^3(M, \mathbb{Z})$ . Assume that we are given a  $\mathbb{Z}_2$ -graded  $\mathcal{L}$ -twisted Clifford module  $\mathcal{E}$  on  $M$  as defined in the Section 4.1. We note that such Clifford modules always exist under our assumptions on the class of the gerbe  $\mathcal{L}$ . We define the algebra  $\Psi_{\mathcal{L}}(M|B; \mathcal{E})$  of projective families of pseudodifferential operators on  $\mathcal{E}$ . One can then define the index of (the positive part of) the twisted Dirac operator  $\text{ind } D^+ \in K_0(\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E}))$ . Here by  $\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E})$  we denote the algebra of smoothing  $\mathcal{L}$ -twisted pseudodifferential operators on  $\mathcal{E}$ .

In order to define the Chern character of the index we use cyclic homology and the map constructed in [22]. This Chern character takes values in the twisted cohomology of  $B$  defined as follows. Let  $\Omega \in \Omega^3(B)$  be a form representing the class  $[\mathcal{L}]$  and let  $u$  be a formal variable of degree  $-2$ . Twisted cohomology  $H_{\mathcal{L}}^{\bullet}(B)$  is then the homology of the complex  $\Omega^*(B)[u]$  with the differential  $d_{\Omega} = ud + u^2\Omega \wedge \cdot$ . Note that the form  $\Omega$  and thus the complex depend on the choice of connection on  $\mathcal{L}$ . Nevertheless the homologies of all the complexes thus obtained are canonically isomorphic. Following Mathai and Stevenson [22] one defines a morphism of complexes  $\Phi_{\nabla\mathcal{H}} : CC_{\bullet}^{-}(\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E})) \rightarrow (\Omega^*(B)[u], d_{\Omega})$ . Here  $CC_{\bullet}^{-}$  denotes the negative cyclic complex. By composing  $\Phi_{\nabla\mathcal{H}}$  (or rather the map induced by it on homology) with the Chern character  $\text{ch} : K_0(\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E})) \rightarrow HC_0^{-}(\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E}))$  we obtain the class  $\Phi_{\nabla\mathcal{H}}(\text{ch}(\text{ind } D^+)) \in H_{\mathcal{L}}^{\bullet}(B)$ . The main result of the paper is the proof of the following theorem expressing the Chern character of the index in terms of characteristic classes:

**Theorem 1.1.** *Let  $D$  be a projective family of Dirac operators on a horizontally  $\mathcal{L}$ -twisted Clifford module  $\mathcal{E}$  on a fibration  $\pi : M \rightarrow B$ . Then the following equality holds in  $H_{\mathcal{L}}^{\bullet}(B)$ :*

$$[\Phi_{\nabla\mathcal{H}}(\text{ch}(\text{ind } D^+))] = \left[ u^{-\frac{k}{2}} \int_{M|B} \widehat{A} \left( \frac{u}{2\pi i} R^{M|B} \right) \text{Ch}_{\mathcal{L}}(\mathcal{E}/\mathcal{S}) \right],$$

where  $k = \dim M - \dim B$  is the dimension of the fibers.

The characteristic classes appearing in the right hand side are defined in the Section 4.1.

Our proof uses the Bismut superconnection formalism [5]. We extend the notion of superconnection to the twisted context and show in the Theorem 4.6 that the Chern character

of superconnection adapted to  $D$  computes the Chern character of  $\text{ind } D^+$ . Then in the Theorem 4.8 using the results of [5] and [6] we compute the limit of the Chern character of the rescaled superconnection obtaining the expression in the right hand side of the index formula. Together these results imply the Theorem 1.1.

The paper is organized as follows. In the Section 2 we give a brief review of cyclic homology, gerbes and connections on them. We also discuss twisted bundles and their characteristic forms. In the Section 3 we give the definitions of the algebra of twisted families of pseudodifferential operators. Finally in the section 4 we define the projective family of Dirac operators on a twisted Clifford module and give the proofs of the main theorems of the paper.

This paper is a byproduct of a joint project with E. Leichtnam. The authors would like to thank him and M. Lesch, V. Mathai and R. Nest for helpful discussions. The authors worked on this paper while visiting CIRM, Luminy as well as Hausdorff institute and Max Planck Institute for Mathematics in Bonn.

## 2. PRELIMINARIES

**2.1. Cyclic homology.** The general reference for this material is the book [19].

Let  $A$  be a complex unital algebra. Set  $C_k(A) = A \otimes (A/\mathbb{C}1)^{\otimes k}$ . Let  $u$  be a formal variable of degree  $-2$ . The space of negative cyclic chains of degree  $l \in \mathbb{Z}$  is defined by

$$CC_l^-(A) = (C_\bullet(A)[[u]])_l = \prod_{-2n+k=l, n \geq 0} u^n C_k(A).$$

The boundary is given by  $b+uB$  where  $b$  and  $B$  are the Hochschild and Connes boundaries of the cyclic complex. The homology of this complex is denoted  $HC_\bullet^-(A)$ . When the algebra  $A$  is  $\mathbb{Z}_2$  graded they incorporate the relevant signs. If  $A$  is not necessarily unital denote by  $A^+$  its unitalisation and set  $CC_l^-(A) = CC_l^-(A^+)/CC_l^-(\mathbb{C})$ . If  $I$  is an ideal in a unital algebra  $A$  the relative cyclic complex is defined by  $CC_\bullet^-(A, I) = \text{Ker}(CC_\bullet^-(A) \rightarrow CC_\bullet^-(A/I))$ . One has a natural morphism of complexes  $\iota: CC_\bullet^-(I) \rightarrow CC_\bullet^-(A, I)$  induced by the homomorphism  $I^+ \rightarrow A$ .

Recall that for an algebra  $A$  we have Chern character in cyclic homology  $\text{ch}: K_0(A) \rightarrow HC_0^-(A)$ . It is defined by the following formula. Let  $P, Q \in M_n(A^+)$  be two idempotents in  $n \times n$  matrices of the algebra  $A^+$ , representing a class  $[P - Q] \in K_0(A)$ . Then

(2.1.1)

$$\text{Ch}([P - Q]) = \text{tr}(P - Q) + \sum_{n=1}^{\infty} (-u)^n \frac{(2n)!}{n!} \text{tr} \left( \left( P - \frac{1}{2} \right) \otimes P^{\otimes (2n)} - \left( Q - \frac{1}{2} \right) \otimes Q^{\otimes (2n)} \right)$$

We will use the notation  $\text{Ch}([P - Q])$  for the cyclic cycle defined above and  $\text{ch}([P - Q])$  for its class in cyclic homology  $HC_0^-(A)$ .

We will also need to use the entire cyclic complex. For our purpose the algebraic version from [11], IV.7.α Remark 7 b. will be sufficient. First recall that one has the periodic cyclic complex  $(CC_\bullet^{\text{per}}(A), b+uB)$  where  $CC_\bullet^{\text{per}}(A) = C_\bullet(A)[u^{-1}, u]$ . Assume we are given a periodic chain  $\alpha = \sum_{k \geq 0} \alpha_k u^k \in CC_m^{\text{per}}(A)$ ,  $\alpha_k \in C_{2k+m}(A)$ . Then  $\alpha$  is called entire if there exist a finite dimensional subspace  $V \subset A$ ,  $1 \in V$  and  $C > 0$  (depending on  $\alpha$ ) such

that  $\alpha_k \in V \otimes (V/\mathbb{C}1)^{\otimes k}$  and  $\|\alpha_k\| \leq C^k k!$ . Here the norms on  $V \otimes (V/\mathbb{C}1)^{\otimes k}$  are induced by an arbitrary norm on  $V$ . We denote the entire cyclic complex of  $A$  by  $CC_{\bullet}^{\text{entire}}(A)$ .

Note that the chain  $\text{Ch}([P - Q])$  defined in (2.1.1) is an element in  $CC_0^{\text{entire}}(A)$ .

**2.2. Notion of a gerbe.** We give here only the general overview, referring the reader to [8] and [17] for the details. We will describe the gerbes in terms of their descent data.

Let  $M$  be a smooth manifold. Given an open cover  $(U_\alpha)_{\alpha \in \Lambda}$  of  $M$ , we set as usual

$$U_{\alpha_1 \dots \alpha_k} = \bigcap_{1 \leq j \leq k} U_{\alpha_j}.$$

**Definition 2.1.** A descent datum for a gerbe  $\mathcal{L}$  on  $M$  is the collection  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  where  $(U_\alpha)_{\alpha \in \Lambda}$  is an open cover of  $M$ ,  $(\mathcal{L}_{\alpha\beta} \rightarrow U_{\alpha\beta})_{\alpha, \beta \in \Lambda}$  is a collection of line bundles and  $\mu_{\alpha\beta\gamma} : \mathcal{L}_{\alpha\beta} \otimes \mathcal{L}_{\beta\gamma} \rightarrow \mathcal{L}_{\alpha\gamma}$  are line bundle isomorphisms over each triple intersection  $U_{\alpha\beta\gamma}$  such that over each quadruple intersection  $U_{\alpha\beta\gamma\delta}$ , the following diagram commutes

$$\begin{array}{ccc} \mathcal{L}_{\alpha\beta} \otimes \mathcal{L}_{\beta\gamma} \otimes \mathcal{L}_{\gamma\delta} & \xrightarrow{\mu_{\alpha\beta\gamma} \otimes \text{id}} & \mathcal{L}_{\alpha\gamma} \otimes \mathcal{L}_{\gamma\delta} \\ \text{id} \otimes \mu_{\beta\gamma\delta} \downarrow & & \downarrow \mu_{\alpha\gamma\delta} \\ \mathcal{L}_{\alpha\beta} \otimes \mathcal{L}_{\beta\delta} & \xrightarrow{\mu_{\alpha\beta\delta}} & \mathcal{L}_{\alpha\delta} \end{array}$$

Notice that we don't assume in this definition that the open sets  $(U_\alpha)_{\alpha \in \Lambda}$  are contractible.

Given two descent data  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  and  $(U_\alpha, \mathcal{L}'_{\alpha\beta}, \mu'_{\alpha\beta\gamma})$  on the same open cover  $\{U_\alpha\}$  an isomorphism between them is given by line bundles  $S_\alpha$  on  $U_\alpha$  and isomorphisms of line bundles  $\lambda_{\alpha\beta} : S_\alpha^{-1} \otimes \mathcal{L}_{\alpha\beta} \otimes S_\beta \rightarrow \mathcal{L}'_{\alpha\beta}$  over  $U_{\alpha\beta}$  so that the diagram

$$\begin{array}{ccc} S_\alpha^{-1} \otimes \mathcal{L}_{\alpha\beta} \otimes S_\beta \otimes S_\beta^{-1} \otimes \mathcal{L}_{\beta\gamma} \otimes S_\gamma & \xrightarrow{\text{id} \otimes \mu_{\alpha\beta\gamma} \otimes \text{id}} & S_\alpha^{-1} \otimes \mathcal{L}_{\alpha\gamma} \otimes S_\gamma \\ \lambda_{\alpha\beta} \otimes \lambda_{\beta\gamma} \downarrow & & \downarrow \lambda_{\alpha\gamma} \\ \mathcal{L}'_{\alpha\beta} \otimes \mathcal{L}'_{\beta\gamma} & \xrightarrow{\mu'_{\alpha\beta\gamma}} & \mathcal{L}'_{\alpha\gamma} \end{array}$$

commutes.

Given two isomorphisms  $(S_\alpha, \lambda_{\alpha\beta})$  and  $(S'_\alpha, \lambda'_{\alpha\beta})$  between  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  and  $(U_\alpha, \mathcal{L}'_{\alpha\beta}, \mu'_{\alpha\beta\gamma})$ , a two-morphism between them is a collection of line bundle isomorphisms  $\nu_\alpha : S_\alpha \rightarrow S'_\alpha$  such that

$$\lambda'_{\alpha\beta} \circ (\nu_\alpha^{-1} \otimes \text{id} \otimes \nu_\beta) = \lambda_{\alpha\beta}$$

where we denote by  $\nu_\alpha^{-1}$  the isomorphism  $S_\alpha^{-1} \rightarrow (S'_\alpha)^{-1}$  induced by  $\nu_\alpha$ .

Let  $(V_i, \varrho)_{i \in I}$  be a refinement of the open cover of  $(U_\alpha)_{\alpha \in \Lambda}$  of  $M$ . So  $(V_i)_{i \in I}$  is an open cover of  $M$  and

$$\varrho : I \rightarrow \Lambda \quad \text{such that} \quad V_i \subset U_{\varrho(i)}.$$

Then restriction to the refinement of  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  is the descent datum  $\mathcal{L}' = (V_i, \mathcal{L}'_{ij}, \mu'_{ijk})$  given by:

$$\mathcal{L}'_{ij} := \mathcal{L}_{\varrho(i)\varrho(j)}|_{V_{ij}} \text{ and } \mu'_{ijk} := \mu_{\varrho(i)\varrho(j)\varrho(k)}|_{V_{ijk}}.$$

Similarly one defines restriction of the isomorphisms and 2-morphisms to a refinement. We do not distinguish between a descent datum, isomorphisms of descent data, etc., and their restrictions to a refinement. Thus for instance, the isomorphism between two descent data  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  and  $(U'_\alpha, \mathcal{L}'_{\alpha\beta}, \mu'_{\alpha\beta\gamma})$  is an isomorphism between their restrictions to some common refinement of  $\{U_\alpha\}$  and  $\{U'_\alpha\}$ .

An equivalence class of Dixmier-Douady gerbes on  $M$  is an equivalence class of the descent data on  $M$ . More precisely a gerbe is a maximal collection of descent data  $\mathcal{D}_i$ ,  $i \in A$  together with the isomorphisms  $s_{ij}: \mathcal{D}_j \rightarrow \mathcal{D}_i$  for each  $i, j \in A$  and 2-morphisms  $\nu_{ijk}: s_{ij}s_{jk} \rightarrow s_{ik}$  satisfying the natural associativity condition. We refer the reader to the book [8] for the details.

If the cover  $U_\alpha$  is good [7] all the bundles  $\mathcal{L}_{\alpha\beta}$  are trivializable. After choice of such trivialization the collection  $(\mu_{\alpha\beta\gamma})$  can be viewed as a Čech 2-cochain with coefficients in the sheaf  $\underline{\mathbb{C}}^*$  of smooth functions with values in the nonzero complex numbers  $\mathbb{C}^*$ . The compatibility condition over  $U_{\alpha\beta\gamma\delta}$  tells us that  $\mu$  is a 2-cocycle and hence defines a cohomology class  $[\mu] \in H^2(M; \underline{\mathbb{C}}^*) \cong H^3(M, \mathbb{Z})$ . This class is a well defined invariant of the gerbe called the Dixmier-Douady class. We denote this class by  $[\mathcal{L}]$ . Every class in  $H^3(M, \mathbb{Z})$  is a class of a gerbe defined by this class uniquely up to an isomorphism (see [8]).

Given a smooth map  $f: M' \rightarrow M$  between smooth manifolds  $M'$  and  $M$ , we can pull-back any descent datum for a gerbe on  $M$  to a descent datum on  $M'$ . The pull backs of isomorphic descent data are isomorphic and thus we obtain a well-defined pull-back of a gerbe. Clearly the Dixmier-Douady class of the pull-back is the pull-back of the Dixmier-Douady class.

An unitary descent datum is  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  together with a choice of metric on each  $\mathcal{L}_{\alpha\beta}$  such that each  $\mu_{\alpha\beta\gamma}$  is an isometry. A notion of unitary equivalence of two unitary descent data on the same open cover  $U_\alpha$  is obtained from the notion of equivalence above by requiring that each line bundle  $S_\alpha$  is Hermitian and each  $\lambda_{\alpha\beta}$  is an isometry. The definition of 2-morphisms is modified by requiring each  $\nu_\alpha$  to be an isometry. It is clear that the restriction of a unitary descent datum to a refinement is again unitary. Then a unitary gerbe is an equivalence class of unitary descent data in the sense described above.

### 2.3. Connections on gerbes.

**Lemma 2.2.** *Let  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  be a descent datum on  $M$ . There exists a collection  $(\nabla_{\alpha\beta})$  of connections on  $(\mathcal{L}_{\alpha\beta})$  such that for any  $(\alpha, \beta, \gamma) \in \Lambda^3$  with  $U_{\alpha\beta\gamma} \neq \emptyset$ :*

$$\mu_{\alpha\beta\gamma}^* \nabla_{\alpha\gamma} = \nabla_{\alpha\beta} \otimes \text{id} + \text{id} \otimes \nabla_{\beta\gamma}.$$

*If the descent datum is unitary each  $\nabla_{\alpha\beta}$  can be chosen Hermitian.*

*Proof.* Fix for any  $\alpha, \beta$  with  $U_{\alpha\beta} \neq \emptyset$  a connection  $\nabla'_{\alpha\beta}$  on  $\mathcal{L}_{\alpha\beta}$ . We set for  $U_{\alpha\beta\gamma} \neq \emptyset$ ,

$$A_{\alpha\beta\gamma} := \mu_{\alpha\beta\gamma}^* \nabla_{\alpha\gamma} - \nabla_{\alpha\beta} \otimes \text{id} - \text{id} \otimes \nabla_{\beta\gamma}.$$

Then using the identification

$$\text{End}(\mathcal{L}_{\alpha\beta} \otimes \mathcal{L}_{\beta\gamma}) \simeq U_{\alpha\beta\gamma} \times \mathbb{C};$$

we see that  $A_{\alpha\beta\gamma}$  is identified with a differential 1-form on the open set  $U_{\alpha\beta\gamma}$ . We also have for  $U_{\alpha\beta\gamma\delta} \neq \emptyset$

$$A_{\beta\gamma\delta} + A_{\alpha\beta\delta} = A_{\alpha\gamma\delta} + A_{\alpha\beta\gamma}.$$

Therefore, there exists  $A' = (A'_{\alpha\beta})$  such that

$$A_{\alpha\beta\gamma} = A'_{\beta\gamma} - A'_{\alpha\gamma} + A'_{\alpha\beta}.$$

The collection of connections  $(\nabla_{\alpha\beta} = \nabla'_{\alpha\beta} + A'_{\alpha\beta})$  is then a connection on the gerbe  $\mathcal{L}$ .  $\square$

**Lemma 2.3.** *Let  $(\nabla_{\alpha\beta})$  be as above, and denote by  $\omega_{\alpha\beta} = \nabla_{\alpha\beta}^2$  the curvatures of the connections  $\nabla_{\alpha\beta}$ . Then there exists a collection of differential 2-forms  $\omega_\alpha \in \Omega^2(U_\alpha)$  such that*

$$\omega_{\alpha\beta} = \omega_\alpha - \omega_\beta, \quad \text{for } U_{\alpha\beta} \neq \emptyset.$$

*Proof.* The collection  $\omega_{\alpha\beta}$  satisfies

$$\omega_{\alpha\gamma} = \omega_{\alpha\beta} + \omega_{\beta\gamma}.$$

This shows that  $(\omega_{\alpha\beta})$  is a Čech cocycle and hence by the acyclicity of the Čech complex of forms, there exist  $(\omega_\alpha)$  such that

$$\omega_{\alpha\beta} = \omega_\alpha - \omega_\beta. \quad \square$$

We will say that the collection  $(\nabla_{\alpha\beta}, \omega_\alpha)$  is a connection on the descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta})$ . Given  $\varepsilon_\alpha \in \Omega^1(U_\alpha)$  we can construct another connection  $(\nabla'_{\alpha\beta}, \omega'_\alpha)$  where  $\nabla'_{\alpha\beta} = \nabla_{\alpha\beta} + \varepsilon_\alpha - \varepsilon_\beta$  and  $\omega'_\alpha = \omega_\alpha + d\varepsilon_\alpha$ . We say that the connections  $(\nabla_{\alpha\beta}, \omega_\alpha)$  and  $(\nabla'_{\alpha\beta}, \omega'_\alpha)$  are equivalent.

Connection on descent datum yields a connection on its restriction to a refinement in an obvious manner. If a connection  $(\nabla'_{\alpha\beta}, \omega'_\alpha)$  on  $(U_\alpha, \mathcal{L}'_{\alpha\beta}, \mu'_{\alpha\beta})$  is given and  $s = (S_\alpha, \lambda_{\alpha\beta})$  is an isomorphism between descent data  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta})$  and  $(U_\alpha, \mathcal{L}'_{\alpha\beta}, \mu'_{\alpha\beta})$ , choice of connections  $\nabla_\alpha$  on each  $S_\alpha$  yields a connection  $s^*(\nabla'_{\alpha\beta}, \omega'_\alpha) = (\nabla_{\alpha\beta}, \omega_\alpha)$  on  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta})$  as follows. Let  $\pi_\alpha = \nabla_\alpha^2$  be the curvatures of these connections and let  $\nabla_\alpha^{-1}$  be the dual connection on  $S_\alpha^{-1}$ . Then we have (using the isomorphism  $\mathcal{L}_{\alpha\beta} = S_\alpha \otimes (S_\alpha^{-1} \otimes \mathcal{L}_{\alpha\beta} \otimes S_\beta) \otimes S_\beta^{-1}$ ):

$$(2.3.1) \quad \nabla_{\alpha\beta} = (\lambda_{\alpha\beta})^*(\nabla_\alpha \otimes \text{id} \otimes \text{id} + \text{id} \otimes \nabla'_{\alpha\beta} \otimes \text{id} + \text{id} \otimes \text{id} \otimes \nabla_\beta^{-1})$$

and

$$(2.3.2) \quad \omega_\alpha = \omega'_\alpha + \pi_\alpha.$$

The equivalence class of this connection is independent of the choice of  $\nabla_\alpha$ . If  $s'$  is another isomorphism between  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta})$  and  $(U_\alpha, \mathcal{L}'_{\alpha\beta}, \mu'_{\alpha\beta})$  such that there exists a 2-morphism between  $s$  and  $s'$ , the equivalence classes of connections induced by  $s$  and  $s'$  coincide. A connection on a gerbe  $\mathcal{L}$  is a choice of equivalence class of connections on each descent datum of  $\mathcal{L}$  in a manner compatible with isomorphism of descent data.

**Lemma 2.4.** *Let  $\mathcal{L}$  be a gerbe represented by a descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$ . Choose a connection on  $\mathcal{L}$  represented by a connection  $(\nabla_{\alpha\beta}, \omega_\alpha)$  on the descent datum.*

- (1) *There exists a well-defined closed form  $\Omega \in \Omega^3(M)$  – curvature 3-form of the connection – such that  $\Omega|_{U_\alpha} = \frac{d\omega_\alpha}{2\pi i}$ .*

- (2) Let  $(\nabla'_{\alpha\beta}, \omega'_\alpha)$  be another connection on  $\mathcal{L}$  and let  $\Omega'$  be the corresponding curvature 3-form. Then there exists a canonical  $\eta \in \Omega^2(M)/d\Omega^1(M)$  such that  $\Omega' = \Omega + d\eta$ .
- (3) Let  $(\nabla_{\alpha\beta}, \omega_\alpha)$ ,  $(\nabla'_{\alpha\beta}, \omega'_\alpha)$ ,  $(\nabla''_{\alpha\beta}, \omega''_\alpha)$  be 3 connections on  $\mathcal{L}$  with the corresponding curvature 3-forms  $\Omega$ ,  $\Omega'$ ,  $\Omega''$ . Let  $\eta, \eta', \eta'' \in \Omega^2(M)/d\Omega^1(M)$  be the canonical elements constructed above such that  $\Omega' - \Omega = d\eta$ ,  $\Omega'' - \Omega' = d\eta'$ ,  $\Omega'' - \Omega = d\eta''$ . Then  $\eta'' = \eta + \eta'$ .

The 3-form  $\Omega$  is a de Rham representative of the Dixmier-Douady class of the gerbe.

*Proof.* We use notations of the previous Lemma. For the first part notice that since  $d\omega_{\alpha\beta} = 0$  for any  $\alpha, \beta$  we see that

$$d\omega_\alpha|_{U_{\alpha\beta}} = d\omega_\beta|_{U_{\alpha\beta}}$$

which shows the existence of  $\Omega$ ; it is clearly closed. It clearly does not change if one restricts the descent datum to a refinement or replaces a connection on descent datum by an equivalent one. The equation (2.3.2) implies that if one replaces a descent datum by an isomorphic one with the induced connection the 3-curvature form does not change.

We proceed to the proof of the second part. Let  $(\nabla_{\alpha\beta}, \omega_\alpha)$  and  $(\nabla'_{\alpha\beta}, \omega'_\alpha)$  be two connections on the descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$ . Set  $\delta_{\alpha\beta} = \nabla'_{\alpha\beta} - \nabla_{\alpha\beta} \in \Omega^1(U_{\alpha\beta})$ . Choose  $\delta_\alpha \in \Omega^1(U_\alpha)$  such that  $\delta_{\alpha\beta} = \delta_\alpha - \delta_\beta$  on  $U_{\alpha\beta}$ . Then  $\omega'_{\alpha\beta} - \omega_{\alpha\beta} = d\delta_{\alpha\beta} = d\delta_\alpha - d\delta_\beta$ . Hence  $\omega'_\alpha - \omega_\alpha - d\delta_\alpha = \omega'_\beta - \omega_\beta - d\delta_\beta$  on  $U_{\alpha\beta}$ . Therefore there exists  $\eta \in \Omega^2(M)$  such that  $\eta|_{U_\alpha} = \frac{1}{2\pi i}(\omega'_\alpha - \omega_\alpha - d\delta_\alpha)$ . Then  $\Omega' - \Omega = d\eta$ . The formula for  $\eta$  given above depends on the choice of  $\delta_\alpha$ . If  $\bar{\delta}_\alpha$  is a different such choice, then  $\bar{\delta}_\alpha - \delta_\alpha = \epsilon|_{U_\alpha}$  for some  $\epsilon \in \Omega^1(M)$ . Then  $\bar{\eta} = \eta - d\epsilon$ . Hence the class of  $\eta \in \Omega^2(M)/d\Omega^1(M)$  does not depend on the choice made. It is easy to see that it does not depend on the choice of the particular descent datum used to represent the connections.

The verification of the third statement is straightforward and left to the reader.  $\square$

**2.4. Twisted cohomology.** For a smooth manifold  $M$  let  $\Omega \in \Omega^3(M)$  be a closed 3-form. Denote by  $u$  a formal variable of degree  $-2$ . The twisted de Rham complex is defined as the complex  $\Omega^*(M)[u]$  with the differential  $d_\Omega = ud + u^2\Omega \wedge \cdot$ . Note that if  $\Omega' = \Omega + d\eta$  is cohomologous to  $\Omega$  then the complexes  $(\Omega^*(M)[u], d_\Omega)$  and  $(\Omega^*(M)[u], d_{\Omega'})$  are isomorphic via the isomorphism

$$(2.4.1) \quad I_\eta: \xi \mapsto e^{-u\eta} \wedge \xi$$

**Lemma 2.5.** *The map induced by  $I_\eta$  on cohomology depends only on the class of  $\eta$  in  $\Omega^2(M)/(d\Omega^1(M))$ .*

*Proof.* Indeed, for  $\epsilon \in \Omega^1(M)$  define  $h_\epsilon: (\Omega^*(M)[u])_\bullet \rightarrow (\Omega^*(M)[u])_{\bullet-1}$  by

$$h_\epsilon = u\epsilon \left( \sum_k \frac{(u\eta)^k}{(k+1)!} \right) \wedge \cdot.$$

Then

$$I_{\eta+d\epsilon} - I_\eta = h_\epsilon \circ d_\Omega + d_{\Omega'} \circ h_\epsilon$$

and therefore the maps  $I_\eta$  and  $I_{\eta+d\epsilon}$  are chain homotopic.  $\square$

The identity  $d_{\Omega+\Omega'}(\xi \wedge \eta) = d_{\Omega}\xi \wedge \eta + (-1)^{|\xi|}\xi \wedge d_{\Omega'}\eta$  implies that the product of forms induces the product  $H_{\Omega}^{\bullet}(M) \otimes H_{\Omega'}^{\bullet}(M) \rightarrow H_{\Omega+\Omega'}^{\bullet}(M)$  and in particular endows  $H_{\Omega}^{\bullet}(M)$  with the structure of  $H^{\bullet}(M)[u]$ -module.

We will also need to consider the following situation. Let  $\pi: M \rightarrow B$  be an oriented fibration with compact fibers. Then we have an integration along the fibers map  $\int_{M|B}: \Omega^*(M) \rightarrow \Omega^{*-k}(B)$ ,  $k = \dim M - \dim B$ . Let  $\Omega \in \Omega^3(B)$  be a closed form. Then  $\int_{M|B} d\eta = d \int_{M|B} \eta$ ,  $\int_{M|B} \pi^* \Omega \wedge \eta = \Omega \wedge \int_{M|B} \eta$  and therefore

$$\int_{M|B} d_{\pi^*\Omega} \eta = d_{\Omega} \int_{M|B} \eta.$$

Hence we obtain a chain map  $\int_{M|B}: (\Omega^*(M), d_{\pi^*\Omega})_{\bullet} \rightarrow (\Omega^*(B), d_{\Omega})_{\bullet-k}$ .

Let  $\mathcal{L}$  be a gerbe. The choice of connection defines a closed 3-form  $\Omega$ . We can therefore consider the complex  $(\Omega^*(M)[u], d_{\Omega})$ . Different choice of connection leads to a different complex  $(\Omega^*(M)[u], d_{\Omega'})$ . Lemma 2.4 however implies that there exists however a canonical isomorphism  $I$  of the homologies of these complexes. We denote this homology  $H_{\mathcal{L}}^{\bullet}(B)$

## 2.5. Twisted bundles.

**Definition 2.6.** A descent datum for twisted vector bundle  $\mathcal{E}$  consists of a descent datum  $(U_{\alpha}, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  for a gerbe  $\mathcal{L}$  together with a collection  $(\mathcal{E}_{\alpha} \rightarrow U_{\alpha})_{\alpha \in \Lambda}$  of vector bundles and a collection of vector bundle isomorphisms  $\varphi_{\alpha\beta}: \mathcal{E}_{\alpha} \otimes \mathcal{L}_{\alpha\beta} \cong \mathcal{E}_{\beta}$  such that for every  $\alpha, \beta, \gamma$  the following diagram commutes

$$\begin{array}{ccc} \mathcal{E}_{\alpha} \otimes \mathcal{L}_{\alpha\beta} \otimes \mathcal{L}_{\beta\gamma} & \xrightarrow{\text{id} \otimes \mu_{\alpha\beta\gamma}} & \mathcal{E}_{\alpha} \otimes \mathcal{L}_{\alpha\gamma} \\ \varphi_{\alpha\beta} \otimes 1 \downarrow & & \downarrow \varphi_{\alpha\gamma} \\ \mathcal{E}_{\beta} \otimes \mathcal{L}_{\beta\gamma} & \xrightarrow{\varphi_{\beta\gamma}} & \mathcal{E}_{\gamma} \end{array}$$

Restriction of the descent datum for  $\mathcal{E}$  to a refinement is given by the restriction of the descent datum for  $\mathcal{L}$  together with restriction of the vector bundles  $\mathcal{E}_{\alpha}$ .

**Definition 2.7.** An (iso)morphism between two descent data  $(U_{\alpha}, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_{\alpha}, \varphi_{\alpha\beta})$  and  $(U'_{\alpha}, \mathcal{L}'_{\alpha\beta}, \mu'_{\alpha\beta\gamma}, \mathcal{E}'_{\alpha}, \varphi'_{\alpha\beta})$  is given by the collection  $(\rho_{\alpha}, S_{\alpha}, \lambda_{\alpha\beta})$  where  $(S_{\alpha}, \lambda_{\alpha\beta})$  is an isomorphism between  $(U_{\alpha}, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  and  $(U'_{\alpha}, \mathcal{L}'_{\alpha\beta}, \mu'_{\alpha\beta\gamma})$  and  $\rho_{\alpha}: \mathcal{E}_{\alpha} \otimes S_{\alpha} \rightarrow \mathcal{E}'_{\alpha}$  is a collection of (iso)morphisms such that the diagram

$$\begin{array}{ccc} \mathcal{E}_{\alpha} \otimes S_{\alpha} \otimes S_{\alpha}^{-1} \otimes \mathcal{L}_{\alpha\beta} \otimes S_{\beta} & \xrightarrow{\varphi_{\alpha\beta} \otimes \text{id}} & \mathcal{E}_{\beta} \otimes S_{\beta} \\ \rho_{\alpha} \otimes \lambda_{\alpha\beta} \downarrow & & \downarrow \rho_{\beta} \\ \mathcal{E}'_{\alpha} \otimes \mathcal{L}'_{\alpha\beta} & \xrightarrow{\varphi'_{\alpha\beta\gamma}} & \mathcal{E}'_{\beta} \end{array}$$

commutes.



An isomorphism between two descent data on two different covers is defined as an isomorphism between their restriction on a common refinement. A 2-morphism between two isomorphisms is the 2-morphisms between the corresponding isomorphisms of the gerbe descent data.

A twisted bundle is then defined as an equivalence class of descent data of twisted vector bundles. “Forgetting” the bundle data we obtain from the descent datum for a twisted vector bundle a descent datum for a gerbe, and the same applies to morphisms and 2-morphisms. We say that  $\mathcal{E}$  is an  $\mathcal{L}$ -twisted vector bundle if “forgetting” the bundle data one obtains equivalence class of the gerbe descent data defining  $\mathcal{L}$ .

Assume now that the gerbe  $\mathcal{L}$  is unitary. An Hermitian descent datum for  $\mathcal{E}$  consists of a unitary descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  for  $\mathcal{L}$  and a collection  $h_\alpha$  of metrics on  $\mathcal{E}_\alpha$  such that the maps  $\varphi_{\alpha\beta}$  are isometries. One obtains a notion of isomorphism of Hermitian descent data by requiring  $\rho_\alpha$  to be isometries. An Hermitian twisted bundle is then an equivalence class of Hermitian descent data.

Given a gerbe  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  on  $M$ , it is well known that (finite dimensional) twisted vector bundles exist if and only if the gerbe is torsion (see e.g. [20] and references therein).

**Lemma 2.8.** *Let  $\mathcal{L}$  be a gerbe on  $M$  and  $\mathcal{E}$  an  $\mathcal{L}$ -twisted bundle. Let  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  be a descent datum for  $\mathcal{L}$ . Then there exists a descent datum for  $\mathcal{E}$  isomorphic to the one of the form  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$ .*

*Proof.* Let  $(V_i, L_{ij}, m_{ijk}, E_i, r_{ij}), i, j, k \in I$  be a descent datum for  $\mathcal{E}$ . Without a loss of generality we may assume that  $\{V_i\}$  is refinement of  $\{U_\alpha\}$  given by the map  $\varrho : I \rightarrow \Lambda$  and that there exists an isomorphism  $(S_i, \lambda_{ij})$  between  $(V_i, L_{ij}, m_{ijk})$  and restriction of  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  to  $\{V_i\}$ .

We define the vector bundle  $\mathcal{E}_\alpha$  as follows. On every non empty open set  $U_\alpha \cap V_i$ , we set

$$\mathcal{E}_\alpha^{(i)} := E_i \otimes S_i \otimes \mathcal{L}_{\varrho(i)\alpha}.$$

Notice that  $V_i \cap U_\alpha \subset U_{\varrho(i)\alpha}$  so that this definition makes sense as a vector bundle over  $V_i \cap U_\alpha$ . Moreover, if  $(i, j) \in I^2$  is such that  $V_{ij} \cap U_\alpha \neq \emptyset$ , then we have a bundle isomorphism

$$\psi_\alpha^{(ij)} : \mathcal{E}_\alpha^{(j)}|_{V_{ij} \cap U_\alpha} \longrightarrow \mathcal{E}_\alpha^{(i)}|_{V_{ij} \cap U_\alpha}.$$

defined by the composition

$$\begin{aligned} E_j \otimes S_j \otimes \mathcal{L}_{\varrho(j)\alpha} &\xrightarrow{\varphi_{ij}^{-1} \otimes \text{id}} E_i \otimes L_{ij} \otimes S_j \otimes \mathcal{L}_{\varrho(j)\alpha} \longrightarrow E_i \otimes S_i \otimes S_i^{-1} \otimes L_{ij} \otimes S_j \otimes \mathcal{L}_{\varrho(j)\alpha} \\ &\xrightarrow{\text{id} \otimes \lambda_{ij} \otimes \text{id}} E_i \otimes S_i \otimes \mathcal{L}_{\varrho(i)\varrho(j)} \otimes \mathcal{L}_{\varrho(j)\alpha} \xrightarrow{\text{id} \otimes \mu_{\varrho(i)\varrho(j)\alpha}} E_i \otimes S_i \otimes \mathcal{L}_{\varrho(i)\alpha} \end{aligned}$$

It is easy to see that

$$\psi_\alpha^{(ik)} = \psi_\alpha^{(ij)} \circ \psi_\alpha^{(jk)}.$$

So, we can glue the bundles  $(\mathcal{E}_\alpha^{(i)})_i$  for  $V_i \cap U_\alpha \neq \emptyset$  together and form a vector bundle  $\mathcal{E}_\alpha$  over  $U_\alpha$ . More precisely

$$\mathcal{E}_\alpha := (\coprod_{V_i \cap U_\alpha \neq \emptyset} \mathcal{E}_\alpha^{(i)}) / \{\psi_\alpha^{(ij)}\}.$$

Next we introduce for  $i \in I$  with  $V_i \cap U_{\alpha\beta} \neq \emptyset$ ,

$$\varphi_{\alpha\beta}^{(i)} := \text{id} \otimes \mu_{\varrho(i)\alpha\beta} : \mathcal{E}_\alpha^{(i)} \otimes \mathcal{L}_{\alpha\beta} \rightarrow \mathcal{E}_\beta^{(i)}.$$

When  $V_{ij} \cap U_{\alpha\beta} \neq \emptyset$ , a straightforward inspection shows again that the following diagram commutes

$$\begin{array}{ccc} \mathcal{E}_\alpha^{(i)} \otimes \mathcal{L}_{\alpha\beta} & \xrightarrow{\varphi_{\alpha\beta}^{(i)}} & \mathcal{E}_\beta^{(i)} \\ \psi_\alpha^{(ji)} \otimes \text{id} \downarrow & & \downarrow \psi_\beta^{(ji)} \otimes \text{id} \\ \mathcal{E}_\alpha^{(j)} \otimes \mathcal{L}_{\alpha\beta} & \xrightarrow{\varphi_{\alpha\beta}^{(j)}} & \mathcal{E}_\beta^{(j)} \end{array}$$

Therefore, the isomorphisms  $\varphi_{\alpha\beta}^{(i)}$  induce an isomorphism

$$\varphi_{\alpha\beta} : \mathcal{E}_\alpha \otimes \mathcal{L}_{\alpha\beta} \rightarrow \mathcal{E}_\beta.$$

We leave it to the reader to show that  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  is a descent datum for a twisted vector bundle isomorphic to  $(V_i, L_{ij}, m_{ijk}, E_i, r_{ij})$ .  $\square$

We now fix a gerbe  $\mathcal{L}$  with a descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  on the smooth manifold  $M$  together with a twisted vector bundle  $\mathcal{E}$  represented by  $(\mathcal{E}_\alpha, \varphi_{\alpha\beta})$ . We denote by  $\mathcal{A}_\alpha$  the collection of bundles of algebras

$$\mathcal{A}_\alpha := \text{End}(\mathcal{E}_\alpha), \quad \alpha \in \Lambda.$$

For any  $U_{\alpha\beta} \neq \emptyset$ , we have a canonical vector bundle isomorphism over  $U_{\alpha\beta}$

$$\rho_{\alpha\beta} : \text{End}(\mathcal{E}_\alpha \otimes \mathcal{L}_{\alpha\beta}) \longrightarrow \mathcal{A}_\alpha,$$

extending the canonical isomorphism  $\text{End}(\mathcal{L}_{\alpha\beta}) \simeq U_{\alpha\beta} \times \mathbb{C}$ . Therefore, the bundle isomorphism  $\varphi_{\alpha\beta}$  together with the identification  $\rho_{\alpha\beta}$  induce the isomorphism of algebra bundles over  $U_{\alpha\beta}$  given by

$$\mathcal{A}_\beta \xrightarrow{\varphi_{\alpha\beta}^*} \text{End}(\mathcal{E}_\alpha \otimes \mathcal{L}_{\alpha\beta}) \xrightarrow{\rho_{\alpha\beta}^*} \mathcal{A}_\alpha.$$

We denote this isomorphism by  $\varphi_{\alpha\beta}^*$ . It is then easy to check that

$$\varphi_{\alpha\beta}^* \circ \varphi_{\beta\gamma}^* = \varphi_{\alpha\gamma}^*, \quad \text{over } U_{\alpha\beta\gamma}.$$

Therefore, the collection  $(\mathcal{A}_\alpha)$  defines a bundle  $\mathcal{A}$  of algebras over  $M$ , which we denote  $\text{End}(\mathcal{E})$ . We leave to the reader an easy check that the isomorphism class of  $\text{End}(\mathcal{E})$  depends only on the isomorphism class of  $\mathcal{E}$ .

Note that for every  $\alpha$  we have the trace  $\text{tr}_\alpha : \text{End}(\mathcal{E}_\alpha) \rightarrow C^\infty(U_\alpha)$ . For  $a \in \Gamma(U_{\alpha\beta}; \mathcal{A}_\beta)$   $\text{tr}_\alpha(\varphi_{\alpha\beta}^*(a)) = \text{tr}_\beta(a)$ . Therefore we obtain the trace  $\text{tr} : \text{End}(\mathcal{E}) \rightarrow C^\infty(M)$  defined by

$$\text{tr}(a)|_{U_\alpha} = \text{tr}_\alpha(a|_{U_\alpha}) \text{ for } a \in \text{End}(\mathcal{E})$$

If the bundle  $\mathcal{E}$  is  $\mathbb{Z}_2$ -graded then  $\text{End}(\mathcal{E})$  is a bundle of  $\mathbb{Z}_2$ -graded algebras with a supertrace  $\text{str} : \text{End}(\mathcal{E}) \rightarrow C^\infty(M)$ .

**Definition 2.9.** The bundle  $\mathcal{A} = \text{End}(\mathcal{E})$  is called the Azumaya bundle associated with the  $\mathcal{L}$ -twisted bundle  $\mathcal{E}$ .

If  $\mathcal{E}$  is an Hermitian twisted bundle, each of the bundles  $\text{End}(\mathcal{E}_\alpha)$  is a bundle of  $*$ -algebras, with the  $*$ -operation given by taking the adjoint endomorphism. This induces a structure of a bundle of  $*$ -algebras on  $\text{End}(\mathcal{E})$ .

Notice that, more generally, if  $\mathcal{E}$  and  $\mathcal{E}'$  are  $\mathcal{L}$ -twisted bundles, we have a well defined bundle  $\text{Hom}(\mathcal{E}, \mathcal{E}')$ , defined similarly by  $\text{Hom}(\mathcal{E}, \mathcal{E}')|_{U_\alpha} = \text{Hom}(\mathcal{E}|_{U_\alpha}, \mathcal{E}'|_{U_\alpha})$ .

Let  $\mathcal{L}$  be a gerbe on  $M$  and  $\mathcal{E}$  an  $\mathcal{L}$ -twisted bundle on  $M$ . Let  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  be a descent datum for  $\mathcal{E}$ .

**Definition 2.10.** A connection on  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  is a collection  $(\nabla_\alpha, \nabla_{\alpha\beta}, \omega_\alpha)$  where  $(\nabla_{\alpha\beta}, \omega_\alpha)$  is a connection on  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  and each  $\nabla_\alpha$  is a connection on  $\mathcal{E}_\alpha$  such that the identities

$$(2.5.1) \quad \varphi_{\alpha\beta}^* \nabla_\beta = \nabla_\alpha \otimes \text{id} + \text{id} \otimes \nabla_{\alpha\beta}$$

hold for  $U_{\alpha\beta} \neq \emptyset$ .

**Lemma 2.11.** *Let  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  be a descent datum for an  $\mathcal{L}$ -twisted bundle  $\mathcal{E}$ . Then every connection  $(\nabla_{\alpha\beta}, \omega_\alpha)$  on the descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  for  $\mathcal{L}$  can be extended to a connection for the descent datum for  $\mathcal{E}$ .*

*Proof.* We start with a collection  $(\nabla'_\alpha)$  of connections on  $(\mathcal{E}_\alpha)$ . Then we set for  $U_{\alpha\beta} \neq \emptyset$ :

$$A_{\alpha\beta} = \varphi_{\alpha\beta}^* \nabla'_\beta - \nabla'_\alpha \otimes \text{id} - \text{id} \otimes \nabla_{\alpha\beta}$$

So,  $A_{\alpha\beta}$  is differential 1-form on  $U_{\alpha\beta}$  with coefficients in the bundle  $\text{End}(\mathcal{E}_\alpha \otimes \mathcal{L}_{\alpha\beta})$ . This latter being canonically isomorphic to  $\text{End}(\mathcal{E}_\alpha)$ , we see that

$$A_{\alpha\beta} \in \Omega^1(U_{\alpha\beta}, \text{End}(\mathcal{E}_\alpha)).$$

Using the functoriality conditions on the isomorphisms  $(\varphi_{\alpha\beta})$ , it is then easy to check that

$$A_{\alpha\gamma} = A_{\alpha\beta} + \varphi_{\alpha\beta}^* A_{\beta\gamma},$$

where  $\varphi_{\alpha\beta}$  is viewed here as the isomorphism over  $U_{\alpha\beta}$  between  $\text{End}(\mathcal{E}_\beta)$  and  $\text{End}(\mathcal{E}_\alpha)$ . Since the sheaf of sections of the bundle  $\text{End}(\mathcal{E})$  is soft, there exists a collection  $A_\alpha \in \Omega^1(U_\alpha; \text{End}(\mathcal{E}_\alpha))$  such that

$$A_{\alpha\beta} = A_\alpha - \varphi_{\alpha\beta}^* A_\beta.$$

The collection  $\nabla_\alpha = \nabla'_\alpha + A_\alpha$  is then a connection on the  $\mathcal{L}$ -twisted vector bundle  $\mathcal{E}$  compatible with the curving  $\nabla_{\alpha\beta}$ .  $\square$

Given a connection  $(\nabla_\alpha, \nabla_{\alpha\beta}, \omega_\alpha)$  and a collection  $\epsilon_\alpha \in \Omega^1(U_\alpha)$  we can construct another connection  $(\nabla'_\alpha, \nabla'_{\alpha\beta}, \omega'_\alpha)$  where  $\nabla'_\alpha = \nabla_\alpha - \epsilon_\alpha$ ,  $\nabla'_{\alpha\beta} = \nabla_{\alpha\beta} + \epsilon_\alpha - \epsilon_\beta$  and  $\omega'_\alpha = \omega_\alpha + d\epsilon_\alpha$ . We will say that these connections are equivalent.

Given a morphism  $s = (\rho_\alpha, S_\alpha, \lambda_{\alpha\beta})$  between two descent data  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  and  $(U'_\alpha, \mathcal{L}'_{\alpha\beta}, \mu'_{\alpha\beta\gamma}, \mathcal{E}'_\alpha, \varphi'_{\alpha\beta})$  for  $\mathcal{E}$  and connection  $(\nabla'_\alpha, \nabla'_{\alpha\beta}, \omega'_\alpha)$  on the second datum, choice of connections on each  $S_\alpha$  allows one to define a pulled-back connection on the first. Namely we set  $\nabla_\alpha = \rho_\alpha^* \nabla'_\alpha \otimes \text{id} + \text{id} \otimes \nabla_{\alpha\beta}^{-1}$ , and the rest of the data is defined as in (2.3.1), (2.3.2). The equivalence class of the pull-back connection is independent of the choice.

A connection on a twisted bundle is a choice of equivalence class of connections on descent data of this twisted bundle compatible with the isomorphisms.

**Proposition 2.12.** *Let  $\mathcal{E}$  be an  $\mathcal{L}$ -twisted bundle with connection. Choose a descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  with a connection  $(\nabla_\alpha, \nabla_{\alpha\beta}, \omega_\alpha)$  representing  $\mathcal{E}$ . Then the collection  $(\theta_\alpha + \omega_\alpha)$ , where  $\theta_\alpha = \nabla_\alpha^2$  is the curvature of  $\nabla_\alpha$ , defines a global differential 2-form  $\theta$  on  $M$  with coefficients in the Azumaya bundle  $\mathcal{A} = \text{End}(\mathcal{E})$ . This form is independent of the choice of the representing descent datum.*

*Proof.* We have for any  $\alpha \in \Lambda$ ,  $\theta_\alpha \in \Omega^2(U_\alpha, \mathcal{A}_\alpha)$  where  $\mathcal{A}_\alpha = \text{End}(\mathcal{E}_\alpha)$  is the Azumaya bundle associated with the twisted bundle  $\mathcal{E}$ . The equation (2.5.1) implies that

$$\varphi_{\alpha\beta}^* \theta_\beta = \theta_\alpha + \omega_{\alpha\beta} \in \Omega^2(U_{\alpha\beta}, \text{End}(\mathcal{E}_\alpha)).$$

Therefore, the collection  $(\theta_\alpha + \omega_\alpha)$  of elements of  $\Omega^2(U_\alpha, \mathcal{A}_\alpha)$  satisfies the relations

$$\varphi_{\alpha\beta}^*(\theta_\beta + \omega_\beta) = \theta_\alpha + \omega_\alpha.$$

It is easy to see that the form  $\theta$  is independent of the equivalence class of the connection and is functorial with respect to the isomorphism of descent data.  $\square$

In the notations above let  $\nabla: \mathcal{A} \rightarrow \Omega^1(M, \mathcal{A})$  be connection defined for a fixed descent datum by

$$(2.5.2) \quad (\nabla\xi)|_{U_\alpha} = [\nabla_\alpha, \xi].$$

It is easy to see that  $\nabla$  is well defined and by derivation with respect to the product on  $\mathcal{A}$ .

Note that

$$(2.5.3) \quad \nabla^2 = [\theta, \cdot] \text{ and } \nabla\theta = 2\pi i\Omega$$

where  $\Omega$  is the 3-curvature form of the connection on  $\mathcal{L}$ , see Lemma 2.4.

**Proposition 2.13.** (1) *Let  $\mathcal{E}$  be an  $\mathcal{L}$ -twisted bundle and  $\nabla$  a connection on  $\mathcal{E}$ . Set  $\text{Ch}_{\mathcal{L}}(\nabla) = \text{tr} e^{-\frac{u\theta}{2\pi i}} \in \Omega^*(M)[u]$ . Then  $d_\Omega \text{Ch}_{\mathcal{L}}(\nabla) = 0$*   
(2) *The class of  $\text{Ch}_{\mathcal{L}}(\nabla)$  in  $H_\Omega(M)$  is independent of choice of connection. Namely assume we are given a different connection  $\nabla'$  on  $\mathcal{E}$  (and therefore on  $\mathcal{L}$ ) and let  $\Omega'$  be the associated 3-curvature form. Then  $I([\text{Ch}_{\mathcal{L}}(\nabla)]) = [\text{Ch}_{\mathcal{L}}(\nabla')]$ , where  $I$  is the canonical isomorphism of cohomology of  $(\Omega^*(M)[u], d_\Omega)$  with cohomology of  $(\Omega^*(M)[u], d_{\Omega'})$ .*

We denote the class of  $\text{Ch}_{\mathcal{L}}(\nabla)$  by  $\text{Ch}_{\mathcal{L}}(\mathcal{E})$ .

*Proof.* Since  $\Omega$  is central,  $\nabla(e^{-\frac{u\theta}{2\pi i}}) = -u\Omega e^{-\frac{u\theta}{2\pi i}}$ . Hence

$$d \text{tr} e^{-\frac{u\theta}{2\pi i}} = \text{tr} \nabla(e^{-\frac{u\theta}{2\pi i}}) = -u\Omega \wedge \text{Ch}_{\mathcal{L}}(\nabla)$$

and  $d_\Omega \text{Ch}_{\mathcal{L}}(\nabla) = 0$ , which proves the first statement.

Fix now a descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  for  $\mathcal{E}$  together with a connection  $\nabla = (\nabla_\alpha, \nabla_{\alpha\beta}, \omega_\alpha)$  and let  $\nabla' = (\nabla'_\alpha, \nabla'_{\alpha\beta}, \omega'_\alpha)$  be another connection on  $\mathcal{E}$ . Set  $\delta_{\alpha\beta} = \nabla'_{\alpha\beta} - \nabla_{\alpha\beta}$  and let  $\delta_\alpha$  be such that  $\delta_\alpha - \delta_\beta = \delta_{\alpha\beta}$ . Recall that the isomorphism  $I$  is induced by the map of complexes

$$\xi \longmapsto e^{-u\eta} \wedge \xi$$

where  $\eta \in \Omega^2(M)$  is defined by  $\eta|_{U_\alpha} = \frac{1}{2\pi i}(\omega'_\alpha - \omega_\alpha - d\delta_\alpha)$ .

Consider now the manifold  $\widetilde{M} = \mathbb{R} \times M$ . Denote by  $\pi: \widetilde{M} \rightarrow M$  the projection on the second factor and by  $t: \widetilde{M} \rightarrow \mathbb{R}$  the projection on the first factor. Consider on  $\widetilde{M}$  the gerbe  $\widetilde{\mathcal{L}} = \pi^*\mathcal{L}$  given by the descent datum  $(\pi^{-1}U_\alpha, \pi^*\mathcal{L}_{\alpha\beta}, \pi^*\mu_{\alpha\beta\gamma})$ . Then  $\pi^*\mathcal{E}$ ,  $\pi^*\phi_{\alpha\beta}$  describe a  $\widetilde{\mathcal{L}}$  twisted bundle  $\widetilde{\mathcal{E}}$ . Moreover,

$$\widetilde{\nabla}_{\alpha\beta} := (1-t)\pi^*\nabla_{\alpha\beta} + t\pi^*\nabla'_{\alpha\beta} \text{ and } \widetilde{\omega}_\alpha = (1-t)\pi^*\omega_\alpha + t\pi^*\omega'_\alpha + dt \wedge \delta_\alpha$$

define respectively, connective structure and curving of  $\widetilde{L}$ . Similarly,  $\widetilde{\nabla}_\alpha = (1-t)\pi^*\nabla_\alpha + t\pi^*\nabla'_\alpha$  defines a compatible connection on  $\widetilde{\mathcal{E}}$ . The corresponding 3-form is given by

$$\widetilde{\Omega} = (1-t)\pi^*\Omega + t\pi^*\Omega' + \frac{1}{2\pi i} dt \wedge (\omega'_\alpha - \omega_\alpha - d\delta_\alpha) = (1-t)\pi^*\Omega + t\pi^*\Omega' + dt \wedge \pi^*\eta.$$

Now, the map  $\xi \mapsto e^{ut\eta} \wedge \xi$  is an isomorphism of complexes  $(\Omega^*(\widetilde{M})[u], d_{\widetilde{\Omega}})$  and  $(\Omega^*(\widetilde{M})[u], d_{\pi^*\Omega})$ . Set  $\widetilde{\mathcal{A}} = \pi^*\mathcal{A}$  and let  $\widetilde{\theta} \in \Omega^2(\widetilde{M}, \widetilde{\mathcal{A}})$  be the form defined by  $\pi^*\theta_\alpha + \widetilde{\omega}_\alpha$ . By the result of the first part of the proposition the differential form  $\text{tr} e^{-\frac{u\theta}{2\pi i}}$  is a cocycle in the complex  $(\Omega^*(\widetilde{M})[u], d_{\widetilde{\Omega}})$ . Hence  $\text{tr} e^{ut\eta} \wedge e^{-\frac{u\theta}{2\pi i}}$  is a cocycle in the complex  $(\Omega^*(\widetilde{M})[u], d_{\pi^*\Omega})$ . This implies the relation

$$\text{tr} e^{u\eta} \wedge e^{-\frac{u\theta'}{2\pi i}} - \text{tr} e^{-\frac{u\theta}{2\pi i}} = d_\Omega \int_0^1 \iota_{\frac{\partial}{\partial t}} \text{tr} e^{ut\eta} \wedge e^{-\frac{u\theta}{2\pi i}} dt.$$

Therefore, we finally deduce that the differential forms  $\text{tr} e^{u\eta} \wedge e^{-\frac{u\theta'}{2\pi i}}$  and  $\text{tr} e^{-\frac{u\theta}{2\pi i}}$  are cohomologous in  $(\Omega^*(M)[u], d_\Omega)$ , and hence that the differential forms  $\text{tr} e^{-\frac{u\theta'}{2\pi i}}$  and  $\text{tr} e^{-u\eta} \wedge e^{-\frac{u\theta}{2\pi i}}$  are cohomologous in  $(\Omega^*(M)[u], d_{\Omega'})$ , which finishes the proof.  $\square$

We will need also a notion of superconnection on the twisted bundle. We now briefly indicate the modifications which need to be made to the notion of connection to obtain that of superconnection. Assume that we are given a gerbe  $\mathcal{L}$  and a  $\mathbb{Z}_2$ -graded  $\mathcal{L}$ -twisted vector bundle  $\mathcal{E} = \mathcal{E}^+ \oplus \mathcal{E}^-$ . Let  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  be a descent datum for  $\mathcal{E}$ .

**Definition 2.14.** A superconnection  $\mathbb{A}$  on the descent datum is  $\mathbb{A} = (\mathbb{A}_\alpha, \nabla_{\alpha\beta}, \omega_\alpha)$  where  $(\nabla_{\alpha\beta}, \omega_\alpha)$  is a connection on the descent datum for  $\mathcal{L}$  and each  $\mathbb{A}_\alpha$  is a superconnection on  $\mathcal{E}_\alpha$  satisfying the relations

$$\varphi_{\alpha\beta}^* \mathbb{A}_\beta = \mathbb{A}_\alpha \otimes \text{id} + \text{id} \otimes \nabla_{\alpha\beta}, \quad \text{for } U_{\alpha\beta} \neq \emptyset.$$

Each superconnection  $\mathbb{A}_\alpha$  can be written as  $\mathbb{A}_\alpha = \sum_{k \geq 0} \mathbb{A}_\alpha^{[k]}$  where  $\mathbb{A}_\alpha^{[k]} \in \Omega^k(U_\alpha; \text{End}(\mathcal{E}_\alpha)^-)$  for  $k$ -even,  $\mathbb{A}_\alpha^{[k]} \in \Omega^k(U_\alpha; \text{End}(\mathcal{E}_\alpha)^+)$  for  $k$ -odd,  $k \neq 1$ , and  $\mathbb{A}_\alpha^{[1]}$  is a grading preserving connection on  $\mathcal{E}_\alpha$ . It is easy to see that for each  $k \neq 1$  there exists a form  $\mathbb{A}^{[k]} \in \Omega^k(M; \text{End}(\mathcal{E}))$  such that  $\mathbb{A}^{[k]}|_{U_\alpha} = \mathbb{A}_\alpha^{[k]}$ . For  $k = 1$   $(\mathbb{A}_\alpha^{[1]}, \nabla_{\alpha\beta}, \omega_\alpha)$  defines a connection on the (descent datum of)  $\mathcal{E}$ .

Let now  $u^{1/2}$  be a formal variable of degree  $-1$  such that  $(u^{1/2})^2 = u$ . Define the rescaled superconnection

$$\mathbb{A}_{u^{-1}} := \sum u^{(k-1)/2} \mathbb{A}^{[k]}.$$

Let  $\Omega$  be as before the curvature 3-form of the connection on  $\mathcal{L}$ . Define the curvature of the rescaled superconnection  $\theta^{\mathbb{A}_{u^{-1}}}$  by

$$\theta^{\mathbb{A}_{u^{-1}}}|_{U_\alpha} = (\mathbb{A}_\alpha)_{u^{-1}}^2 + \omega_\alpha.$$

Then  $u\theta^{\mathbb{A}_{u^{-1}}} \in \Omega^{even}(M, \text{End}(\mathcal{E})^+)[u] + u^{1/2}\Omega^{odd}(M, \text{End}(\mathcal{E})^-)[u]$ . We therefore have the differential form  $\exp\left(-\frac{u\theta^{\mathbb{A}_{u^{-1}}}}{2\pi i}\right)$  which belongs to  $\Omega^{even}(M, \text{End}(\mathcal{E})^+)[u] + u^{1/2}\Omega^{odd}(M, \text{End}(\mathcal{E})^-)[u]$  and the differential form  $\text{str} \exp\left(-\frac{u\theta^{\mathbb{A}_{u^{-1}}}}{2\pi i}\right)$  which belongs to  $\Omega^{even}(M)[u]$ . The following is an analogue of the Proposition 2.13 for the superconnections with the essentially identical proof.

**Proposition 2.15.**

- (1) Set  $\text{Ch}_{\mathcal{L}}(\mathbb{A}) = \text{str} \exp\left(-\frac{u\theta^{\mathbb{A}_{u^{-1}}}}{2\pi i}\right)$ . Then  $d_\Omega \text{Ch}_{\mathcal{L}}(\mathbb{A}) = 0$
- (2) The class of  $\text{Ch}_{\mathcal{L}}(\mathbb{A})$  in  $H_\Omega(M)$  is independent of choice of superconnection. Specifically, assume we are given a different superconnection  $\mathbb{A}'$  on  $\mathcal{E}$  (and therefore a different connection on  $\mathcal{L}$ ) and let  $\Omega'$  be the associated 3-curvature form. Then  $I([\text{Ch}_{\mathcal{L}}(\mathbb{A})]) = [\text{Ch}_{\mathcal{L}}(\mathbb{A}')]$ , where  $I$  is the canonical isomorphism of cohomology of  $(\Omega^*(M)[u], d_\Omega)$  with cohomology of  $(\Omega^*(M)[u], d_{\Omega'})$ .

**2.6. Horizontally twisted bundles.** Let  $\pi : M \rightarrow B$  be a smooth fibration. Let  $\mathcal{L}$  be a gerbe on  $B$ .

**Definition 2.16.** A descent datum for a horizontally  $\mathcal{L}$ -twisted bundle  $\mathcal{E}$  on  $M$  consists of the descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  for  $\mathcal{L}$  together with a collection  $(\mathcal{E}_\alpha \rightarrow \pi^{-1}U_\alpha)_{\alpha \in \Lambda}$  of vector bundles and a collection of vector bundle isomorphisms  $\varphi_{\alpha\beta} : \mathcal{E}_\alpha \otimes \pi^*\mathcal{L}_{\alpha\beta} \cong \mathcal{E}_\beta$  so that

$$(\pi^{-1}U_\alpha, \pi^*\mathcal{L}_{\alpha\beta}, \pi^*\mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$$

is a descent datum for a twisted vector bundle on  $M$ .

An (iso)morphism between two such descent data

$$(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta}) \text{ and } (U'_\alpha, \mathcal{L}'_{\alpha\beta}, \mu'_{\alpha\beta\gamma}, \mathcal{E}'_\alpha, \varphi'_{\alpha\beta})$$

is given by the collection  $(\rho_\alpha, S_\alpha, \lambda_{\alpha\beta})$  where  $(S_\alpha, \lambda_{\alpha\beta})$  is an isomorphism between  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma})$  and  $(U'_\alpha, \mathcal{L}'_{\alpha\beta}, \mu'_{\alpha\beta\gamma})$  and  $\rho_\alpha : \mathcal{E}_\alpha \otimes \pi^*S_\alpha \rightarrow \mathcal{E}'_\alpha$  is such that  $(\rho_\alpha, \pi^*S_\alpha, \pi^*\lambda_{\alpha\beta})$  is an (iso)morphism between  $(\pi^{-1}U_\alpha, \pi^*\mathcal{L}_{\alpha\beta}, \pi^*\mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  and  $(\pi^{-1}U'_\alpha, \pi^*\mathcal{L}'_{\alpha\beta}, \pi^*\mu'_{\alpha\beta\gamma}, \mathcal{E}'_\alpha, \varphi'_{\alpha\beta})$ .

With these definitions one can now define a horizontally twisted bundle as an equivalence class of descent data. Let  $Tw(\pi^*\mathcal{L})$  denote the set of isomorphism classes of all  $\pi^*\mathcal{L}$ -twisted bundles on  $M$  and  $Tw_h(\mathcal{L})$  denote the set of isomorphism classes of all horizontally  $\mathcal{L}$ -twisted bundles. Then we have an obvious map  $Tw_h(\mathcal{L}) \rightarrow Tw(\pi^*\mathcal{L})$ . According to Lemma 2.8 this map is surjective. In particular  $Tw_h(\mathcal{L}) \neq \emptyset$  if and only if  $\pi^*[\mathcal{L}]$  is torsion in  $H^3(M, \mathbb{Z})$ . It is however not injective. Indeed, if  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  is a descent datum for a horizontally twisted bundle and  $S$  is a line bundle on  $M$  then  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha \otimes S|_{\pi^{-1}U_\alpha}, \varphi_{\alpha\beta} \otimes \text{id})$  is another such descent datum. These data define the same element of  $Tw(\pi^*\mathcal{L})$  but, unless  $S$  is a pull-back of a line bundle from  $B$ , different elements of  $Tw_h(\mathcal{L})$ .

A connection on the descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  is a collection  $(\nabla_\alpha, \nabla_{\alpha\beta}, \omega_\alpha)$  where  $(\nabla_{\alpha\beta}, \omega_\alpha)$  is a connection on the descent datum for  $\mathcal{L}$  and  $\nabla_\alpha$  is a connection on  $\mathcal{E}_\alpha$  such

that  $(\nabla_\alpha, \pi^*\nabla_{\alpha\beta}, \pi^*\omega_\alpha)$  is a connection on the descent datum  $(\pi^{-1}U_\alpha, \pi^*\mathcal{L}_{\alpha\beta}, \pi^*\mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$ . Connections  $(\nabla_\alpha, \nabla_{\alpha\beta}, \omega_\alpha)$  and  $(\nabla_\alpha - \pi^*\epsilon_\alpha, \nabla_{\alpha\beta} + \epsilon_\alpha - \epsilon_\beta, \omega_\alpha + d\epsilon_\alpha)$ ,  $\epsilon_\alpha \in \Omega^1(U_\alpha)$  are equivalent. With these definitions one can now define a notion of connection on the horizontally  $\mathcal{L}$ -twisted bundle. If  $\nabla$  is such a connection and  $\Omega$  is the curvature 3-form of the gerbe  $\mathcal{L}$ , one defines  $\text{Ch}_\mathcal{L}(\nabla)$  – a closed form in  $(\Omega^*(M), d_{\pi^*\Omega})_\bullet$ . The analogues of Propositions 2.12 and 2.13 hold in this context with the same proofs.

### 3. PROJECTIVE FAMILIES AND THE ANALYTIC INDEX

**3.1. Families of pseudodifferential operators.** Here we collect several facts about the (untwisted) families of pseudodifferential operators.

Let  $\pi: X \rightarrow Y$  be a smooth fibration and  $E$  a vector bundle on  $X$ .

We denote by  $\Psi_\mathcal{L}^m(X|Y; E)$  the space of classical fiberwise pseudodifferential operators of order  $\leq m$  on  $\pi$ , acting on the sections of the vector bundle  $E$ . As usual, we set

$$\Psi(X|Y; E) := \bigcup_{m \in \mathbb{Z}} \Psi^m(X|Y; E)$$

and

$$\Psi^{-\infty}(X|Y; E) := \bigcap_{m \in \mathbb{Z}} \Psi^m(X|Y; E).$$

Recall that composition endows each  $\Psi(X|Y; E)$  with the structure of a filtered algebra and that  $\Psi^{-\infty}(X|Y; E)$  is an ideal in this algebra.  $\Psi(X|Y; E)$  is also a module over  $C^\infty(Y)$  and the composition is  $C^\infty(Y)$ -linear.

We have the following elementary general result:

**Lemma 3.1.** *Assume  $\pi: X \rightarrow Y$  is a smooth fibration,  $E$  a vector bundle on  $X$ ,  $L$  a line bundle on  $Y$ . Define a map  $\chi_L: \Psi(X|Y; E) \rightarrow \Psi(X|Y; E \otimes \pi^*L)$  by*

$$\chi_L(D)(e \otimes \pi^*(l)) = D(e) \otimes \pi^*l$$

for  $D \in \Psi_\mathcal{L}(X|Y; \mathcal{E})$ ,  $e \in \Gamma_c(E)$ ,  $l \in \Gamma(L)$ . Then  $\chi_L$  is a well-defined isomorphism of algebras and  $C^\infty(Y)$ -modules.

$$\chi_{L_1 \otimes L_2} = \chi_{L_2} \circ \chi_{L_1}$$

If we have two vector bundles  $E, E'$  on  $X$  we denote by  $\Psi(X|Y; E, E')$  the set of fiberwise pseudodifferential operators  $\Gamma_c(E) \rightarrow \Gamma(E')$ . For  $L$  – line bundle on  $Y$  we again have the isomorphism of  $C^\infty(Y)$ -modules  $\chi_L: \Psi(X|Y; E, E') \rightarrow \Psi(X|Y; E \otimes \pi^*L, E' \otimes \pi^*L)$  defined by the same formula.

We have the vertical cotangent bundle  $T^*(X|Y) = T^*X / (\text{Ker } \pi_*)^\perp$ .  $\mathring{T}^*(X|Y)$  denotes (the total space of) this bundle with the zero section removed, and  $p: \mathring{T}^*(X|Y) \rightarrow X$  is the natural projection. Recall that for  $P \in \Psi^m(X|Y; E, E')$  the principal symbol  $\sigma_m(P)$  is an  $m$ -homogeneous smooth section over  $\mathring{T}^*(X|Y)$  of vector bundle  $p^* \text{Hom}(E, E')$ . Then identifying canonically isomorphic bundles  $\text{Hom}(E, E')$  and  $\text{Hom}(E \otimes \pi^*L, E' \otimes \pi^*L)$  we have  $\sigma_m(P) = \sigma_m(\chi_L(P))$ .

**3.2. Projective families.** Let  $\pi : M \rightarrow B$  be a smooth fibration with compact fibers. Let  $\mathcal{L}$  be a gerbe on  $B$  such that  $\pi^*[\mathcal{L}]$  is a torsion class in  $H^3(M, \mathbb{Z})$ . Let  $\mathcal{E}$  be a horizontally  $\mathcal{L}$ -twisted bundle on  $M$ , cf. Section 2.6. We fix a descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  for  $\mathcal{E}$ . For any  $(\alpha, \beta) \in \Lambda^2$  with  $U_{\alpha\beta} \neq \emptyset$  we have an isomorphism of filtered algebras, respecting the  $C^\infty(U_{\alpha\beta})$ -module structure:

$$(3.2.1) \quad \phi_{\alpha\beta} : \Psi(\pi^{-1}U_{\alpha\beta}|U_{\alpha\beta}; \mathcal{E}_\beta) \rightarrow \Psi(\pi^{-1}U_{\alpha\beta}|U_{\alpha\beta}; \mathcal{E}_\alpha),$$

It is defined as the composition

$$\Psi_{\mathcal{L}}(\pi^{-1}U_{\alpha\beta}|U_{\alpha\beta}; \mathcal{E}_\beta) \xrightarrow{\psi_{\alpha\beta}} \Psi_{\mathcal{L}}(\pi^{-1}U_{\alpha\beta}|U_{\alpha\beta}; \mathcal{E}_\beta \otimes \pi^*\mathcal{L}_{\alpha\beta}) \xrightarrow{\varphi_{\alpha\beta}} \Psi_{\mathcal{L}}(\pi^{-1}U_{\alpha\beta}|U_{\alpha\beta}; \mathcal{E}_\alpha)$$

where  $\psi_{\alpha\beta} = \chi_{\mathcal{L}_{\alpha\beta}}^{-1} = \chi_{\mathcal{L}_{\beta\alpha}}$ .

Recall (cf. [4]) that for every  $\alpha \in \Lambda$  we have an infinite dimensional bundle  $\pi_*\mathcal{E}_\alpha$  on  $U_\alpha$  defined by  $\Gamma(V, \pi_*\mathcal{E}_\alpha) = \Gamma(\pi^{-1}V, \mathcal{E}_\alpha)$ ,  $V \subset U_\alpha$ . Over  $U_{\alpha\beta}$  we have isomorphisms  $\pi_*\varphi_{\alpha\beta} : \pi_*\mathcal{E}_\alpha \otimes \mathcal{L}_{\alpha\beta} \rightarrow \pi_*\mathcal{E}_\beta$  defined by

$$\pi_*\varphi_{\alpha\beta}(\xi \otimes l) = \varphi(\xi \otimes \pi^*(l)).$$

Here  $\xi \in \Gamma(U_{\alpha\beta}, \pi_*\mathcal{E}_\alpha) = \Gamma(\pi^{-1}U_{\alpha\beta}, \mathcal{E}_\alpha)$ ,  $l \in \Gamma(U_{\alpha\beta}, \mathcal{L}_{\alpha\beta})$ .

Note that the isomorphisms  $\pi_*\varphi_{\alpha\beta} : \pi_*\mathcal{E}_\alpha \otimes \mathcal{L}_{\alpha\beta} \rightarrow \pi_*\mathcal{E}_\beta$  induce the isomorphisms

$$(\pi_*\varphi_{\alpha\beta})^* : \text{End}(\pi_*\mathcal{E}_\beta) \rightarrow \text{End}(\pi_*\mathcal{E}_\alpha \otimes \mathcal{L}_{\alpha\beta}) \cong \text{End}(\pi_*\mathcal{E}_\alpha)$$

over  $U_{\alpha\beta}$ . The restriction of this isomorphism to  $\Psi(\pi^{-1}U_{\alpha\beta}|U_{\alpha\beta}, \mathcal{E}_\beta) \subset \text{End}(\pi_*\mathcal{E}_\beta)$  coincides with the isomorphism  $\phi_{\alpha\beta} : \Psi(\pi^{-1}U_{\alpha\beta}|U_{\alpha\beta}, \mathcal{E}_\beta) \rightarrow \Psi(\pi^{-1}U_{\alpha\beta}|U_{\alpha\beta}, \mathcal{E}_\alpha)$ . Since the isomorphisms  $(\pi_*\varphi_{\alpha\beta})^*$  satisfy the natural cocycle identity we have the following:

**Lemma 3.2.** *The isomorphisms  $\phi_{\alpha\beta}$  satisfy*

$$\phi_{\alpha\beta} \circ \phi_{\beta\gamma} = \phi_{\alpha\gamma}$$

whenever  $U_{\alpha\beta\gamma} \neq \emptyset$ .

Recall that the isomorphisms  $\varphi_{\alpha\beta}$  induce the natural isomorphisms  $\varphi_{\alpha\beta}^* : \text{End}(\mathcal{E}_\beta) \rightarrow \text{End}(\mathcal{E}_\alpha)$ . Then

$$(3.2.2) \quad \sigma_m \circ \phi_{\alpha\beta} = p^* (\varphi_{\alpha\beta}^*) \circ \sigma_m$$

**Definition 3.3.** A fiberwise pseudodifferential operator  $P$  of order  $\leq m$  with coefficients in the horizontally  $\mathcal{L}$ -twisted vector bundle  $\mathcal{E}$  is a collection  $\{P_\alpha\}_{\alpha \in \Lambda}$ ,  $P_\alpha \in \Psi^m(\pi^{-1}U_\alpha|U_\alpha; \mathcal{E}_\alpha)$  such that

$$P_\alpha = \phi_{\alpha\beta}(P_\beta).$$

where  $\phi_{\alpha\beta}$  is defined in Equation 3.2.1. The space of fiberwise pseudodifferential operators of order  $\leq m$ , with coefficients in the  $\pi^*\mathcal{L}$ -twisted vector bundle  $\mathcal{E}$ , is denoted by  $\Psi_{\mathcal{L}}^m(M|B; \mathcal{E})$ .

Note that the equation (3.2.2) implies that if  $P = \{P_\alpha\} \in \Psi_{\mathcal{L}}^m(M|B; \mathcal{E})$  then the collection  $\sigma_m(P_\alpha)$  defines a section of the (untwisted) bundle  $p^*\text{End}(\mathcal{E})$ . We will call this section the principal symbol of  $P = \{P_\alpha\}$



*Remark 3.4.* We define in the same way the space  $\Psi_{\mathcal{L}}^m(M|B; \mathcal{E}, \mathcal{E}')$  of fiberwise pseudodifferential operators of order  $\leq m$ , from the horizontally  $\mathcal{L}$ -twisted vector bundle  $\mathcal{E}$  to the horizontally  $\mathcal{L}$ -twisted vector bundle  $\mathcal{E}'$ . In particular  $\Psi_{\mathcal{L}}^m(M|B; \mathcal{E}, \mathcal{E}) = \Psi_{\mathcal{L}}^m(M|B; \mathcal{E})$ . We also have a principal symbol map  $\sigma_m: \Psi_{\mathcal{L}}^m(M|B; \mathcal{E}, \mathcal{E}) \rightarrow p^* \text{Hom}(\mathcal{E}, \mathcal{E}')$ .

We set

$$\Psi_{\mathcal{L}}(M|B; \mathcal{E}) := \bigcup_{m \in \mathbb{Z}} \Psi_{\mathcal{L}}^m(M|B; \mathcal{E}) \text{ and } \Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E}) := \bigcap_{m \in \mathbb{Z}} \Psi_{\mathcal{L}}^m(M|B; \mathcal{E}).$$

Introduce now a composition in  $\Psi_{\mathcal{L}}(M|B; \mathcal{E})$  by

$$\{P_{\alpha}\} \circ \{Q_{\alpha}\} = \{P_{\alpha}Q_{\alpha}\}$$

Since  $\phi_{\alpha\beta}$  are algebra isomorphisms the right hand side of this equality defines an element in  $\Psi_{\mathcal{L}}(M|B; \mathcal{E})$ .

**Proposition 3.5.** *The composition of operators is  $C^{\infty}(B)$ -linear and endows  $\Psi_{\mathcal{L}}(M|B; \mathcal{E})$  with the structure of associative algebra;  $\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E})$  is an ideal in  $\Psi_{\mathcal{L}}(M|B; \mathcal{E})$ .*

We can now define the algebra of forms on  $B$  with values in  $\Psi_{\mathcal{L}}(M|B; \mathcal{E})$  by

$$\Omega^*(B, \Psi_{\mathcal{L}}(M|B; \mathcal{E})) = \Omega^*(B) \otimes_{C^{\infty}(B)} \Psi_{\mathcal{L}}(M|B; \mathcal{E}).$$

Recall that for every  $\alpha$  and  $V \subset U_{\alpha}$  we have a fiberwise trace  $\text{Tr}_{\alpha}: \Psi^{-\infty}(\pi^{-1}V|V; \mathcal{E}_{\alpha}) \rightarrow C^{\infty}(V)$ . It is easy to see that for  $P \in \Psi^{-\infty}(\pi^{-1}U_{\alpha\beta}|U_{\alpha\beta}; \mathcal{E}_{\beta})$

$$\text{Tr}_{\alpha} \phi_{\alpha\beta}(P) = \text{Tr}_{\beta}(P).$$

We therefore obtain a well defined map  $\text{Tr}: \Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E}) \rightarrow C^{\infty}(B)$  by setting  $\text{Tr}\{P_{\alpha}\}|_{U_{\alpha}} = \text{Tr}_{\alpha}(P_{\alpha})$ . This trace is a  $C^{\infty}(B)$ -module map satisfying  $\text{Tr}[A, B] = 0$ . It extends naturally to define a map  $\text{Tr}: \Omega^*(B, \Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E})) \rightarrow \Omega^*(B)$ .

If the bundle  $\mathcal{E}$  is  $\mathbb{Z}_2$  graded we have a similarly defined supertrace

$$\text{Str}: \Omega^*(B, \Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E})) \rightarrow \Omega^*(B).$$

Note that the definition of the  $\Psi_{\mathcal{L}}(M|B; \mathcal{E})$  depends on the descent datum for  $\mathcal{E}$ . It is straightforward however to see that an isomorphism of descent data defines canonically an isomorphism of the corresponding bundles of algebras.

### 3.3. Analytic index.

**Definition 3.6.** Let  $m \geq 0$  be fixed. A fiberwise pseudodifferential operator  $P \in \Psi_{\mathcal{L}}^m(M|B; \mathcal{E}, \mathcal{E}')$  is fiberwise elliptic if the principal symbol  $\sigma_m(P) \in \Gamma(p^* \text{Hom}(\mathcal{E}, \mathcal{E}'))$  is an isomorphism.

We say that  $Q \in \Psi_{\mathcal{L}}^{-m}(M|B; \mathcal{E}', \mathcal{E})$  is a parametrix of  $P \in \Psi_{\mathcal{L}}^m(M|B; \mathcal{E}, \mathcal{E}')$  if  $PQ - 1 \in \Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E}', \mathcal{E}')$  and  $QP - 1 \in \Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E}, \mathcal{E})$ .

**Lemma 3.7.** *Every elliptic  $P \in \Psi_{\mathcal{L}}^m(M|B; \mathcal{E}, \mathcal{E}')$  has a parametrix.*

*Proof.* Construct first a parametrix  $R_{\alpha}$  for  $P_{\alpha}$ . Let  $\rho_{\alpha}$  be a partition of unity subordinate to  $\{U_{\alpha}\}$ . Then define  $R'_{\alpha} = \sum_{\beta} \rho_{\beta} \phi_{\alpha\beta}(R_{\beta})$ . Then each  $R'_{\alpha}$  is a parametrix for  $P_{\alpha}$  and  $\phi_{\alpha\beta}(R'_{\beta}) = R'_{\alpha}$ .  $\square$

Let  $D \in \Psi_{\mathcal{L}}^m(M|B; \mathcal{E}, \mathcal{E}')$  be elliptic. Let  $F \in \Psi_{\mathcal{L}}^0(M|B; \mathcal{E}, \mathcal{E}')$  be such that  $\sigma_0(F)|_{S^*(M|B)} = \sigma_m(D)|_{S^*(M|B)}$ . Choose a parametrix  $R$  for  $F$ . Let  $U_D \in \Psi_{\mathcal{L}}^0(M|B; \mathcal{E} \oplus \mathcal{E}')$  be an invertible operator such that  $U_D - \begin{bmatrix} 0 & -R \\ F & 0 \end{bmatrix} \in \Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}')$ . An explicit construction of an example of such an operator is as follows. Let  $S_0 = 1 - RF$ ,  $S_1 = 1 - FR$ . Then set  $U_D = \begin{bmatrix} S_0 & -(1 + S_0)R \\ F & S_1 \end{bmatrix}$ . With such a choice the inverse is given by an explicit formula  $U_D^{-1} = \begin{bmatrix} S_0 & (1 + S_0)R \\ -F & S_1 \end{bmatrix}$ .

**Definition 3.8.** The index of  $D$  is the  $K$ -theory class of the algebra  $\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}')$  defined by

$$\text{ind}(D) = [P_D - Q] \in K_0(\Psi_{\mathcal{L}}^0(M|B, \mathcal{E} \oplus \mathcal{E}'), \Psi_{\mathcal{L}}^{-\infty}(M|B, \mathcal{E} \oplus \mathcal{E}')) \cong K_0(\Psi_{\mathcal{L}}^{-\infty}(M|B, \mathcal{E} \oplus \mathcal{E}')),$$

where  $P_D$  and  $Q$  are the idempotents given by  $P_D = U_D \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} U_D^{-1}$  and  $Q = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ .

We leave to the reader the standard  $K$ -theoretic proof that the index is well defined and is stable under the homotopies of  $D$  in the class of elliptic operators in  $\Psi_{\mathcal{L}}^m(M|B; \mathcal{E}, \mathcal{E}')$ . Assume that the horizontally  $\mathcal{L}$ -twisted bundles  $\mathcal{E}$  and  $\mathcal{E}'$  are hermitian (and in particular  $\mathcal{L}$  is unitary) and that fibers of  $\pi$  are equipped with smoothly varying volume forms. In this situation for a projective family  $D$  we can define an formally adjoint projective family  $D^*$  by forming the formal adjoints for each family  $D_\alpha$ .

**Lemma 3.9.** *Then, identifying  $K_0(\Psi_{\mathcal{L}}^{-\infty}(M|B, \mathcal{E} \oplus \mathcal{E}'))$  with  $K_0(\Psi_{\mathcal{L}}^{-\infty}(M|B, \mathcal{E}' \oplus \mathcal{E}))$  we have*

$$\text{ind } D^* = -\text{ind } D.$$

*Proof.* We have

$$\text{ind } D^* = \left[ U_{D^*} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} U_{D^*}^{-1} - \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \right] \in K_0(\Psi_{\mathcal{L}}^{-\infty}(M|B, \mathcal{E} \oplus \mathcal{E}')).$$

By deforming  $D$  we may assume that  $\sigma_m(D)|_{S^*(M|B)}$  is an isometry. In this case we may choose  $U_{D^*} = U_D^{-1}$ , and the statement follows.  $\square$

**3.4. Chern character of the index.** We continue in the notations of the previous section. We assume that we are given a horizontally  $\mathcal{L}$ -twisted bundle  $\mathcal{E}$  with a connection represented by a descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  with connection  $(\nabla_\alpha, \nabla_{\alpha\beta}, \omega_\alpha)$  see the Section 2.6. Following Mathai and Stevenson [22], we describe in this paragraph a morphism of complexes  $CC_\bullet^-(\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E})) \rightarrow (\Omega^*(M)[u], d_\Omega)_\bullet$ .

Recall the bundles  $\pi_*\mathcal{E}_\alpha$  and isomorphisms  $\pi_*\varphi_{\alpha\beta}$  defined in the Section 3.2. It is easy to see that  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \pi_*\mathcal{E}_\alpha, \pi_*\varphi_{\alpha\beta})$  is a descent datum for an infinite dimensional twisted bundle. We now proceed to define a connection on this descent datum.

Choose horizontal distribution i.e. a subbundle  $\mathcal{H} \subset TM$  such that  $TM = \mathcal{H} \oplus T(M|B)$ . This choice together with connections  $\nabla_\alpha^\mathcal{E}$  defines for each  $\alpha$  a connection  $\nabla_\alpha^\mathcal{H}$  as follows:

$$(\nabla_\alpha^\mathcal{H})_X \xi = (\nabla_\alpha^\mathcal{E})_{X^\mathcal{H}} \xi$$

where  $X^\mathcal{H}$  is the horizontal lift of  $X \in \Gamma(B, TB)$ .

**Lemma 3.10.**  $(\pi_*\varphi_{\alpha\beta})^*\nabla_{\beta}^{\mathcal{H}} = \nabla_{\alpha}^{\mathcal{H}} \otimes \text{id} + \text{id} \otimes \nabla_{\alpha\beta}$

The curvature of the connection  $\nabla_{\alpha}^{\mathcal{H}}$  is a 2-form  $\theta_{\alpha}^{\mathcal{H}}$  on  $U_{\alpha}$  with values in fiberwise differential operators given by

$$\theta_{\alpha}^{\mathcal{H}}(X, Y) = \theta_{\alpha}^{\mathcal{E}}(X^{\mathcal{H}}, Y^{\mathcal{H}}) + (\nabla_{\alpha}^{\mathcal{E}})_{T(X, Y)}.$$

where

$$(3.4.1) \quad T^{\mathcal{H}}(X, Y) = [X^{\mathcal{H}}, Y^{\mathcal{H}}] - [X, Y]^{\mathcal{H}}, \quad X, Y \in \Gamma(B, TB).$$

Each  $\nabla_{\alpha}^{\mathcal{H}}$  defines a filtration-preserving derivation  $\partial_{\alpha}^{\mathcal{H}}$  of the algebra of fiberwise pseudodifferential operators

$$\partial_{\alpha}^{\mathcal{H}}: \Psi(\pi^{-1}U_{\alpha}|U_{\alpha}, \mathcal{E}_{\alpha}) \rightarrow \Omega^1(U_{\alpha}, \Psi(\pi^{-1}U_{\alpha}|U_{\alpha}, \mathcal{E}_{\alpha})) \text{ defined by } \partial_{\alpha}^{\mathcal{H}}(D) = [\nabla_{\alpha}^{\mathcal{H}}, D].$$

If  $D \in \Psi(\pi^{-1}U_{\alpha\beta}|U_{\alpha\beta}, \mathcal{E}_{\beta})$  then the result of Lemma 3.10 implies that

$$\partial_{\alpha}^{\mathcal{H}}(\phi_{\alpha\beta}(D)) = \phi_{\alpha\beta}(\partial_{\beta}^{\mathcal{H}}(D)).$$

Therefore if  $\{D_{\alpha}\}$ ,  $D_{\alpha} \in \Psi_{\mathcal{L}}(\pi^{-1}U_{\alpha}|U_{\alpha}, \mathcal{E}_{\alpha})$ , defines an element in  $\Psi_{\mathcal{L}}^m(M|B, \mathcal{E})$  then  $\{\partial_{\alpha}^{\mathcal{H}}(D_{\alpha})\} \in \Omega^1(B, \Psi_{\mathcal{L}}^m(M|B, \mathcal{E}))$ . We therefore obtain a derivation

$$\partial^{\mathcal{H}}: \Psi_{\mathcal{L}}(M|B, \mathcal{E}) \rightarrow \Omega^1(B, \Psi_{\mathcal{L}}(M|B, \mathcal{E})),$$

which extends to a derivation of the algebra  $\Omega^*(B, \Psi_{\mathcal{L}}(M|B, \mathcal{E}))$ .

**Lemma 3.11.** *There exists  $\theta^{\mathcal{H}} \in \Omega^2(B, \Psi_{\mathcal{L}}^1(M|B, \mathcal{E}))$  such that*

$$\theta^{\mathcal{H}}|_{U_{\alpha}} = \theta_{\alpha}^{\mathcal{H}} + \pi^*\omega_{\alpha}$$

*Proof.* By Lemma 3.10,  $\phi_{\alpha\beta}^*\theta_{\beta}^{\mathcal{H}} = \theta_{\alpha}^{\mathcal{H}} + \pi^*(\omega_{\alpha} - \omega_{\beta})$ , and the statement follows as in Proposition 2.12.  $\square$

We have

$$(\partial^{\mathcal{H}})^2(D) = [\theta^{\mathcal{H}}, D] \text{ and } \partial^{\mathcal{H}}(\theta^{\mathcal{H}}) = 2\pi i(\pi^*\Omega).$$

where  $\Omega$  is the 3-curvature form of the connection on  $\mathcal{L}$ .

Following Mathai and Stevenson [22] one can construct the morphism of complexes

$$\Phi_{\nabla^{\mathcal{H}}}: CC_{\bullet}^{-}(\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}')) \rightarrow (\Omega^*(B)[u], d_{\Omega})_{\bullet}$$

as follows. (Note that in the nontwisted case similar morphism was constructed in [14, 25, 26].) Denote by

$$\Delta^k := \{(t_0, \dots, t_k) \in \mathbb{R}^k \mid 0 \leq t_i, \sum_{i=0}^k t_i = 1\}.$$

the standard  $k$ -simplex.

Define the maps  $\Phi_{\nabla^{\mathcal{H}}}^k: C_k(\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}')) \rightarrow \Omega^*(M)[u]$  by

$$\Phi_{\nabla^{\mathcal{H}}}^k(A_0, \dots, A_k) := \int_{\Delta^k} \text{Tr} \left( A_0 e^{-ut_0 \frac{\theta^{\mathcal{H}}}{2\pi i}} \partial^{\mathcal{H}}(A_1) e^{-ut_1 \frac{\theta^{\mathcal{H}}}{2\pi i}} \dots e^{-ut_{k-1} \frac{\theta^{\mathcal{H}}}{2\pi i}} \partial^{\mathcal{H}}(A_k) e^{-ut_k \frac{\theta^{\mathcal{H}}}{2\pi i}} \right) dt_1 \dots dt_k,$$

for  $A_0 \otimes \dots \otimes A_k \in C_k(\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}'))$ . Then let  $\Phi_{\nabla^{\mathcal{H}}} = \sum_{k=0}^{\infty} \Phi_{\nabla^{\mathcal{H}}}^k$ .

**Theorem 3.12.** [22] *The map  $\Phi_{\nabla^{\mathcal{H}}}: (CC_{\bullet}^{-}(\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E})), b + uB) \rightarrow (\Omega^*(B)[u], d_{\Omega})_{\bullet}$  is a morphism of complexes.*

This morphism depends on the choice of horizontal distribution  $\mathcal{H}$ . However the results of [22] show that a different choice of  $\mathcal{H}$  leads to a chain homotopic morphism.

Assume now that  $D \in \Psi_{\mathcal{L}}(M|B; \mathcal{E}, \mathcal{E}')$  is an twisted elliptic family. In the Definition 3.8 we defined  $\text{ind}(D) = [P_D - Q] \in K_0(\overline{\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E}, \mathcal{E}')} )$ . Here the idempotents  $P_D$  and  $Q$  belong to the algebra  $\overline{\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}')} -$  the unitalization of  $\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}')$  with the multiplier  $Q \in \Psi_{\mathcal{L}}^0(M|B; \mathcal{E} \oplus \mathcal{E}')$  adjoined. It follows that if we directly apply the formula (2.1.1) to  $[P_D - Q]$  we obtain a 0-chain  $\text{Ch}[P_D - Q]$  in the relative cyclic complex  $CC_{\bullet}^{-}(\overline{\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}')} , \Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}'))$ . We have the natural morphism  $\iota: CC_{\bullet}^{-}(\overline{\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}')} ) \rightarrow CC_{\bullet}^{-}(\overline{\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}')} , \Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}'))$ , and the following equality of homology classes:  $\iota(\text{ch}(\text{ind } D)) = [\text{Ch}[P_D - Q]]$ . It is straightforward to see that the map  $\Phi_{\nabla \mathcal{H}}$  extends to a morphism

$$\Phi_{\nabla \mathcal{H}}: CC_{\bullet}^{-}(\overline{\Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}')} , \Psi_{\mathcal{L}}^{-\infty}(M|B; \mathcal{E} \oplus \mathcal{E}')) \rightarrow (\Omega^*(B)[u], d_{\Omega})_{\bullet}$$

defined by the same formula, and that we have an equality  $\Phi_{\nabla \mathcal{H}} \circ \iota = \Phi_{\nabla \mathcal{H}}$ . It follows that

$$\Phi_{\nabla \mathcal{H}}(\text{ch}(\text{ind } D)) = [\Phi_{\nabla \mathcal{H}}(\text{Ch}[P_D - Q])].$$

#### 4. DIRAC OPERATORS AND SUPERCONNECTIONS

**4.1. Dirac operators.** The goal of this section is to give a superconnection proof of the family index theorem for projective families of Dirac operators. We assume that the fibers of the smooth fibration  $\pi: M \rightarrow B$  are even dimensional compact Riemannian manifolds.

We begin with the definition of a horizontally twisted Clifford module. Denote by  $C(M|B)$  the Clifford algebra of the fiberwise cotangent bundle  $T^*(M|B) = T^*M/(\ker \pi_*)^{\perp}$ . Let  $\mathcal{L}$  be a unitary gerbe on  $B$ .

##### Definition 4.1.

- A twisted Clifford module is a horizontally  $\mathcal{L}$ -twisted Hermitian  $\mathbb{Z}_2$ -graded vector bundle  $\mathcal{E} = \mathcal{E}^+ \oplus \mathcal{E}^-$  on  $M$  together with the homomorphism  $c: C(M|B) \rightarrow \text{End}(\mathcal{E})$  of bundles of unital  $\mathbb{Z}_2$ -graded  $*$ -algebras.
- A Clifford connection  $\nabla^{\mathcal{E}}$  on  $\mathcal{E}$  is an Hermitian connection such that  $\nabla^{\mathcal{E}}(c(a)) = c(\nabla^{M|B}(a))$ .

Clifford connections on Clifford modules always exist; the proof is analogous to the proof of existence of twisted connections in Lemma 2.11.

Choose horizontal distribution i.e. a subbundle  $\mathcal{H} \subset TM$  such that  $TM = \mathcal{H} \oplus T(M|B)$ . This choice together with the Riemannian metric on the fibers of  $\pi$  allows one to define a connection  $\nabla^{M|B}$  on the fiberwise tangent bundle  $T(M|B)$ , see [4] Section 10.1. We denote by  $R^{M|B}$  the curvature of this connection.

Set  $\text{End}_{C(M|B)}(\mathcal{E}) = \{A \in \text{End}(\mathcal{E}) \mid [A, c(a)] = 0 \text{ for every } a \in C(M|B)\}$ . Let  $\Gamma \in C(M|B)$  be the chirality operator defined locally by  $\Gamma = i^{k/2} e^1 \dots e^k$  where  $k = \dim M - \dim B$  and  $e^1, \dots, e^k$  is the local orthonormal basis of  $T^*(M|B)$ . Define then the relative supertrace

$$\text{str}_{\mathcal{E}/\mathcal{S}}: \text{End}_{C(M|B)}(\mathcal{E}) \rightarrow C^{\infty}(M) \text{ by } \text{str}_{\mathcal{E}/\mathcal{S}}(A) = 2^{-k/2} \text{str } c(\Gamma)A.$$

We fix from now on a Clifford connection  $\nabla^\mathcal{E}$  on  $\mathcal{E}$  and a descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  for the horizontally  $\mathcal{L}$ -twisted Clifford module  $\mathcal{E}$ . The connection  $\nabla^\mathcal{E}$  defines a connection  $(\nabla_\alpha^\mathcal{E}, \nabla_{\alpha\beta}, \omega_\alpha)$  (defined up to equivalence) on this descent datum. Each  $\mathcal{E}_\alpha$  is then a Clifford module on the fibration  $\pi^{-1}U_\alpha \rightarrow U_\alpha$ , and each connection  $\nabla_\alpha^\mathcal{E}$  is a Clifford connection.

Recall (see Proposition 2.12) that one defines  $\theta^\mathcal{E} \in \Omega^2(M, \text{End}(\mathcal{E}))$  by setting  $\theta^\mathcal{E}|_{\pi^{-1}U_\alpha} = \theta_\alpha^\mathcal{E} + \pi^*\omega_\alpha$ . Denote by  $c(R^{M|B})$  the action of the 2-form with values in the Clifford algebra obtained from  $R^{M|B}$  via the Lie algebra isomorphism  $\mathfrak{so}(T(M|B)) \rightarrow C^2(M|B)$ . Here  $C^2(M|B) \subset C(M|B)$  is a subspace consisting of elements  $\sum u_i v_i$ ,  $u_i, v_i \in T^*(M|B)$  with  $\sum \langle u_i, v_i \rangle = 0$ . Define  $\theta^{\mathcal{E}/\mathcal{S}} = \theta^\mathcal{E} - c(R^{M|B})$ .

The argument in [4], Proposition 3.43, shows that  $\theta^{\mathcal{E}/\mathcal{S}} \in \Omega^2(M, \text{End}_{C(M|B)}(\mathcal{E}))$ . Over an open set  $\pi^{-1}U_\alpha$  we have  $\theta^{\mathcal{E}/\mathcal{S}}|_{U_\alpha} = \theta_\alpha^{\mathcal{E}/\mathcal{S}} + \pi^*\omega_\alpha$  where  $\theta_\alpha^{\mathcal{E}/\mathcal{S}} \in \text{End}_{C(M|B)}(\mathcal{E}_\alpha)$  is defined via the equality  $\theta_\alpha^\mathcal{E} = \theta_\alpha^{\mathcal{E}/\mathcal{S}} + c(R^{M|B})$ . We can then define a differential form  $\text{Ch}_\mathcal{L}(\mathcal{E}/\mathcal{S})$  by

$$\text{Ch}_\mathcal{L}(\mathcal{E}/\mathcal{S}) = \text{str}_{\mathcal{E}/\mathcal{S}} e^{-\frac{u\theta^{\mathcal{E}/\mathcal{S}}}{2\pi i}} \in \Omega^*(M)[u].$$

The proof of the following result is standard and analogous to the proof of Proposition 2.13.

**Lemma 4.2.** *We have  $d_{\pi^*\Omega} \text{Ch}_\mathcal{L}(\mathcal{E}/\mathcal{S}) = 0$  and the corresponding class is  $H_{\pi^*\Omega}^*(M)$  is independent of the choice of Clifford connection  $\nabla^\mathcal{E}$ .*

We also introduce the fiberwise  $\widehat{A}$ -genus by  $\widehat{A}(TM|B) = \widehat{A}\left(\frac{u}{2\pi i} R^{M|B}\right) \in \Omega^*(M)[u]$ , where  $\widehat{A}(x)$  is the power series defined by

$$\widehat{A}(x) = \det^{1/2} \left( \frac{x/2}{\sinh x/2} \right).$$

Using the above data, we can define on each fibration  $\pi^{-1}U_\alpha \rightarrow U_\alpha$  a family of Dirac operators  $D_\alpha$  acting on the sections of the bundle  $\mathcal{E}_\alpha$ . Locally  $D_\alpha = \sum_i c(e^i) (\nabla_\alpha^\mathcal{E})_{e_i}$  where  $\{e_i\}, \{e^i\}$  are dual bases of  $T(M|B)$  and  $T^*(M|B)$  respectively. We leave it to the reader to check the following.

**Lemma 4.3.** *The collection  $D = \{D_\alpha\}$  defines an element in  $\Psi_\mathcal{L}^1(M|B; \mathcal{E})$ .*

With respect to the decomposition  $\mathcal{E} = \mathcal{E}^+ \oplus \mathcal{E}^-$  we have the decomposition  $\Psi_\mathcal{L}(M|B; \mathcal{E}) = \Psi_\mathcal{L}(M|B; \mathcal{E}^+, \mathcal{E}^+) \oplus \Psi_\mathcal{L}(M|B; \mathcal{E}^+, \mathcal{E}^-) \oplus \Psi_\mathcal{L}(M|B; \mathcal{E}^-, \mathcal{E}^+) \oplus \Psi_\mathcal{L}(M|B; \mathcal{E}^-, \mathcal{E}^-)$ . The Dirac operator then decomposes as  $D = D^+ \oplus D^-$  where  $D^+ \in \Psi_\mathcal{L}^1(M|B; \mathcal{E}^+, \mathcal{E}^-)$ ,  $D^- \in \Psi_\mathcal{L}^1(M|B; \mathcal{E}^-, \mathcal{E}^+)$ .

Classical arguments show that  $D^+$  is fiberwise elliptic and hence the analytical index  $\text{ind}(D^+)$  of  $D^+$  is well defined in  $K(\Psi_\mathcal{L}^{-\infty}(M|B; \mathcal{E}))$ .

We then have the following result

**Theorem 4.4.** *The following formula holds in  $H_\mathcal{L}^\bullet(B)$ :*

$$\Phi_{\nabla^\mathcal{H}}(\text{ch}(\text{ind } D^+)) = u^{-\frac{\dim M - \dim B}{2}} \left[ \int_{M/B} \widehat{A}\left(\frac{u}{2\pi i} R^{M|B}\right) \wedge \text{Ch}_\mathcal{L}(\mathcal{E}/\mathcal{S}) \right].$$

This index theorem is established in [20, 21] for general projective families however with the more restrictive conditions on the class  $[\mathcal{L}]$  of the gerbe  $\mathcal{L}$ . The superconnection proof of this result occupies the rest of the paper.

**4.2. Superconnections and index.** We continue in the notations of the previous section.

A twisted superconnection  $\mathbb{A}$  on the descent datum  $(U_\alpha, \mathcal{L}_{\alpha\beta}, \mu_{\alpha\beta\gamma}, \mathcal{E}_\alpha, \varphi_{\alpha\beta})$  is a collection  $(\mathbb{A}_\alpha)_{\alpha \in \Lambda}$  of superconnections on the vector bundles  $\pi_* \mathcal{E}_\alpha$  over the open sets  $U_\alpha$  such that when  $U_{\alpha\beta} \neq \emptyset$ ,

$$(4.2.1) \quad (\pi_* \varphi_{\alpha\beta})^* \mathbb{A}_\beta = \mathbb{A}_\alpha \otimes \text{id} + \text{id} \otimes \nabla_{\alpha\beta}.$$

We say that  $\mathbb{A}$  is a Bismut superconnection if each  $\mathbb{A}_\alpha$  is. Specifically we have

$$\mathbb{A}_\alpha = D_\alpha + \nabla_\alpha^{\mathcal{H}} - \frac{1}{4}c(T^{\mathcal{H}})$$

where  $T^{\mathcal{H}}$  is defined in (3.4.1).

The fact that  $\nabla^{\mathcal{H}}$  is a connection on the descent datum, and  $D \in \Psi^1(M|B, \mathcal{E})$ ,  $c(T^{\mathcal{H}}) \in \Omega^2(B, \Psi^0(M|B, \mathcal{E}))$  implies that the conditions (4.2.1) are satisfied. From now on  $\mathbb{A}$  will denote the Bismut superconnection.

We will also consider the rescaled Bismut superconnection  $\mathbb{A}_s$  defined by

$$(\mathbb{A}_s)_\alpha = \mathbb{A}_{\alpha,s} := s^{1/2}D_\alpha + \nabla_\alpha^{\mathcal{H}} - \frac{1}{4}s^{-1/2}c(T^{\mathcal{H}})$$

where  $s$  is either a positive number or a multiple of the formal variable  $u$ . Denote by  $\theta_\alpha^{\mathbb{A}_s}$  the curvature of the rescaled superconnection  $(\mathbb{A}_s)_\alpha$ . In particular we have forms  $u\theta_\alpha^{\mathbb{A}_s, u^{-1}} \in \Omega^*(U_\alpha, \Psi(\pi^{-1}U_\alpha|U_\alpha, \mathcal{E}_\alpha))[u^{1/2}]$ .

**Proposition 4.5.** *There exists a form  $u\theta^{\mathbb{A}_s, u^{-1}} \in \Omega^*(B, \Psi(M|B; \mathcal{E}))[u^{1/2}]$  such that*

$$u\theta^{\mathbb{A}_s, u^{-1}}|_{U_\alpha} = u(\theta_\alpha^{\mathbb{A}_s, u^{-1}} + \pi^*\omega_\alpha).$$

*Proof.* Recall that the curvature of  $\nabla_{\alpha\beta}$  is equal to  $\omega_\alpha - \omega_\beta$ . Therefore from the equation (4.2.1) we obtain

$$\phi_{\alpha\beta}(u\theta^{\mathbb{A}_s, u^{-1}}) = u(\theta_\alpha^{\mathbb{A}_s, u^{-1}} + \pi^*\omega_\alpha - \pi^*\omega_\beta).$$

The statement of the Proposition follows.  $\square$

Notice that  $u\theta^{\mathbb{A}_s, u^{-1}} = D^2 +$  forms of degree  $> 0$ . Therefore we can define  $e^{-\frac{u\theta^{\mathbb{A}_s, u^{-1}}}{2\pi i}} \in \Omega^*(B, \Psi^{-\infty}(M|B; \mathcal{E}))[u^{1/2}]$  by the usual Duhamel's formula.

Note that the parity considerations as in finite dimensional case show that the coefficients for the nonintegral powers of  $u$  are odd with respect to the grading and hence have a vanishing supertrace.

Given a superconnection  $\mathbb{A} = (\mathbb{A}_\alpha)_{\alpha \in \Lambda}$  on the  $\mathbb{Z}_2$ -graded horizontally  $\mathcal{L}$ -twisted Clifford module  $\mathcal{E}$  by the infinite-dimensional version of the Proposition 2.15 the the differential form

$$\text{Str}(e^{-\frac{u\theta^{\mathbb{A}_s, u^{-1}}}{2\pi i}}) \in \Omega^*(B)[u],$$

is closed with respect to the twisted de Rham differential  $d_\Omega$ . The proof is identical to the proof of the first part of the Proposition 2.13. The proof of the following Theorem is adapted from [15] and it uses ideas from [16]. The use of cyclic theory is inspired by [27].

**Theorem 4.6.** *The following equality holds in the  $H_{\mathcal{L}}^\bullet(B)$ :*

$$\Phi_{\nabla^{\mathcal{H}}}(\text{ch}(\text{ind } D^+)) = \left[ \text{Str}(e^{-\frac{u\theta^{\mathbb{A}_s, u^{-1}}}{2\pi i}}) \right]$$

*Proof.* Consider the twisted bundle  $\tilde{\mathcal{E}} = \mathcal{E} \oplus \mathcal{E}$  with the grading given by  $\Gamma = \begin{bmatrix} \gamma & 0 \\ 0 & -\gamma \end{bmatrix}$ .

The algebra  $\Psi_{\mathcal{L}}(M|B; \tilde{\mathcal{E}})$  of operators on  $\tilde{\mathcal{E}}$  is naturally  $\mathbb{Z}_2$  graded. When discussing cyclic complexes of this algebra and its subalgebras we always consider it as  $\mathbb{Z}_2$ -graded algebra. For an operator or (super) connection  $K$  on  $\mathcal{E}$  set  $\tilde{K} = K \oplus K$ ; so for example  $\tilde{\mathbb{A}} = \mathbb{A} \oplus \mathbb{A}$ ,  $\tilde{\nabla}^{\mathcal{H}} = \nabla^{\mathcal{H}} \oplus \nabla^{\mathcal{H}}$ , etc.

Let  $F = D(1 + D^2)^{-1/2}$ . Then it is immediate that  $F \in \Psi_{\mathcal{L}}^0(M|B; \mathcal{E})$  is odd with respect to the grading  $\Gamma$  and fiberwise elliptic. Construct the invertible operator  $U_D \in \Psi_{\mathcal{L}}^{-\infty}(M|B; \tilde{\mathcal{E}})$  by the same formula as before. Namely, choose a parametrix  $R$  for  $F$ . Let  $S_0 = 1 - RF$ ,  $S_1 = 1 - FR$ . Then set  $U_D = \begin{bmatrix} S_0 & -(1 + S_0)R \\ F & S_1 \end{bmatrix}$ . With such a choice the inverse is given by an explicit formula  $U_D^{-1} = \begin{bmatrix} S_0 & (1 + S_0)R \\ -F & S_1 \end{bmatrix}$ . Set  $P_D = U_D^{-1} \begin{bmatrix} 1_{\mathcal{E}} & 0 \\ 0 & 0 \end{bmatrix} U_D$ .

The choices in the constructions can be made so that we have  $(P_D)^{\pm} = P_{D^{\pm}}$ , see Definition 3.8.

Define the map  $\Phi_{\tilde{\nabla}^{\mathcal{H}}} : CC_{\bullet}^{-} \left( \Psi_{\mathcal{L}}^{-\infty}(M|B; \tilde{\mathcal{E}}) \right) \rightarrow (\Omega^*(M)[u], d_{\Omega})$  by

$$\Phi_{\tilde{\nabla}^{\mathcal{H}}}^k(A_0, \dots, A_k) := \int_{\Delta^k} \text{Str} \left( A_0 e^{-ut_0 \frac{\tilde{\theta}^{\mathcal{H}}}{2\pi i}} \tilde{\partial}^{\mathcal{H}}(A_1) e^{-ut_1 \frac{\tilde{\theta}^{\mathcal{H}}}{2\pi i}} \dots e^{-ut_{k-1} \frac{\tilde{\theta}^{\mathcal{H}}}{2\pi i}} \tilde{\partial}^{\mathcal{H}}(A_k) e^{-ut_k \frac{\tilde{\theta}^{\mathcal{H}}}{2\pi i}} \right) dt_1 \dots dt_k$$

for  $A_0 \otimes \dots \otimes A_k \in C_k \left( \Psi_{\mathcal{L}}^{-\infty}(M|B; \tilde{\mathcal{E}}) \right)$  and set again  $\Phi_{\tilde{\nabla}^{\mathcal{H}}} = \sum_{k=0}^{\infty} \Phi_{\tilde{\nabla}^{\mathcal{H}}}^k$ . Notice that

$$(4.2.2) \quad \left[ \Phi_{\tilde{\nabla}^{\mathcal{H}}} \left( \text{Ch} \left( P_D - \begin{bmatrix} 0 & 0 \\ 0 & 1_{\mathcal{E}} \end{bmatrix} \right) \right) \right] = \\ \Phi_{\nabla^{\mathcal{H}}}(\text{ch}(\text{ind } D^+)) - \Phi_{\nabla^{\mathcal{H}}}(\text{ch}(\text{ind } D^-)) = 2\Phi_{\nabla^{\mathcal{H}}}(\text{ch}(\text{ind } D^+))$$

Replacing in the formulas above  $\tilde{\theta}^{\mathcal{H}}$  by  $\tilde{\theta}^{\mathbb{A}_{u-1}}$  and  $\tilde{\partial}^{\mathcal{H}}$  by  $[\tilde{\mathbb{A}}_{u-1}, \cdot]$  we obtain the definition of the morphism  $\Phi_{\tilde{\mathbb{A}}} : CC_{\bullet}^{\text{entire}} \left( \Psi_{\mathcal{L}}^{-\infty}(M|B; \tilde{\mathcal{E}}) \right) \rightarrow (\Omega^*(M)[u], d_{\Omega})_{\bullet}$ .

**Lemma 4.7.** *The morphisms  $\Phi_{\tilde{\mathbb{A}}}, \Phi_{\tilde{\nabla}^{\mathcal{H}}} : CC_{\bullet}^{\text{entire}} \left( \Psi_{\mathcal{L}}^{-\infty}(M|B; \tilde{\mathcal{E}}) \right) \rightarrow (\Omega^*(B)[u], d_{\Omega})_{\bullet}$  are chain homotopic.*

*Proof.* This follows from the explicit formula for the chain homotopy, see Proposition 5.6 of [22]. This formula can be described as follows. Let  $\mathbb{A}_{u-1}(s) = s\mathbb{A}_{u-1} + (1-s)\nabla^{\mathcal{H}}$ . Then we can write

$$\mathbb{A}_{u-1}(s) = su^{-1/2}D + \nabla^{\mathcal{H}} - \frac{su^{1/2}}{4}c(T^{\mathcal{H}}), \\ \theta^{\mathbb{A}_{u-1}(s)} = \theta^{\mathcal{H}} + s[\nabla^{\mathcal{H}}, u^{-1/2}D - \frac{u^{1/2}}{4}c(T^{\mathcal{H}})] + s^2(u^{-1/2}D - \frac{u^{1/2}}{4}c(T^{\mathcal{H}}))^2 \\ \text{and } \frac{d}{ds}\mathbb{A}_{u-1}(s) = u^{-1/2}D - \frac{u^{1/2}}{4}c(T^{\mathcal{H}}).$$

Define  $H_k: C_k(\Psi_{\mathcal{L}}^{-\infty}(M|B, \mathcal{E} \oplus \mathcal{E})) \rightarrow \Omega^*(B)[u]$  by

$$(4.2.3) \quad H_k(A_0, \dots, A_k) = \int_0^1 ds \left( \sum_{m=0}^k (-1)^m \int_{\Delta^{k+1}} dt_1 \dots dt_{k+1} \text{Str} \left( A_0 e^{-ut_0 \frac{\tilde{\theta}^{\mathbb{A}}_{u-1}(s)}{2\pi i}} \right. \right. \\ \left. \left. [\tilde{\mathbb{A}}_{u-1}(s), A_1] e^{-ut_1 \frac{\tilde{\theta}^{\mathbb{A}}_{u-1}(s)}{2\pi i}} \dots [\tilde{\mathbb{A}}_{u-1}(s), A_m] e^{-ut_m \frac{\tilde{\theta}^{\mathbb{A}}_{u-1}(s)}{2\pi i}} \frac{\widetilde{d}}{ds} \mathbb{A}_{u-1} \right. \right. \\ \left. \left. e^{-ut_{m+1} \frac{\tilde{\theta}^{\mathbb{A}}_{u-1}(s)}{2\pi i}} \dots e^{-ut_{k-1} \frac{\tilde{\theta}^{\mathbb{A}}_{u-1}(s)}{2\pi i}} [\tilde{\mathbb{A}}_{u-1}(s), A_k] e^{-ut_{k+1} \frac{\tilde{\theta}^{\mathbb{A}}_{u-1}(s)}{2\pi i}} \right) \right)$$

and set  $H = \sum H_k: CC_{\bullet}^{\text{entire}}(\Psi_{\mathcal{L}}^{-\infty}(M|B; \tilde{\mathcal{E}})) \rightarrow \Omega^*(B)[u]$ . Then

$$\Phi_{\tilde{\mathbb{A}}} - \Phi_{\nabla \mathcal{H}} = d_{\Omega} \circ H + H \circ (b + uB).$$

□

Let  $\mathcal{F}$  be the algebra defined by

$$\mathcal{F} = \{F \in \Psi_{\mathcal{L}}^0(M|B; \tilde{\mathcal{E}})^{\text{even}} \mid [D, F] \in \Psi_{\mathcal{L}}^0(M|B; \mathcal{E})\}.$$

A simple modification of the argument in [13] as done in [2], cf. also [15] shows that  $\Phi_{\tilde{\mathbb{A}}}$  extends to a morphism, also denoted  $\Phi_{\tilde{\mathbb{A}}}: CC_{\bullet}^{\text{entire}}(\mathcal{F}) \rightarrow (\Omega^*(M)[u], d_{\Omega})$  defined by the same formula. Note that  $U_D \in \mathcal{F}$ ,  $P_D \in \mathcal{F}$  and  $\begin{bmatrix} 1_{\mathcal{E}} & 0 \\ 0 & 0 \end{bmatrix} \in \mathcal{F}$ .

Recall now that inner automorphisms act by identity on the entire cyclic homology, see e.g. [19] 4.1.3 or [18]. It follows that the chains  $\text{Ch}(P_D)$  and  $\text{Ch}\left(\begin{bmatrix} 1_{\mathcal{E}} & 0 \\ 0 & 0 \end{bmatrix}\right)$  are homologous in  $CC_{\bullet}^{\text{entire}}(\mathcal{F})$ .

We therefore obtain

$$\begin{aligned} \Phi(\text{ch}(\text{ind } D^+)) &= \frac{1}{2} \left[ \Phi_{\nabla \mathcal{H}} \left( \text{Ch} \left( P_D - \begin{bmatrix} 0 & 0 \\ 0 & 1_{\mathcal{E}} \end{bmatrix} \right) \right) \right] \\ &= \frac{1}{2} \left[ \Phi_{\tilde{\mathbb{A}}} \left( \text{Ch} \left( P_D - \begin{bmatrix} 0 & 0 \\ 0 & 1_{\mathcal{E}} \end{bmatrix} \right) \right) \right] \\ &= \frac{1}{2} \left[ \Phi_{\tilde{\mathbb{A}}}(\text{Ch}(P_D)) - \Phi_{\tilde{\mathbb{A}}} \left( \text{Ch} \left( \begin{bmatrix} 0 & 0 \\ 0 & 1_{\mathcal{E}} \end{bmatrix} \right) \right) \right] \\ &= \frac{1}{2} \left[ \Phi_{\tilde{\mathbb{A}}} \left( \text{Ch} \left( \begin{bmatrix} 1_{\mathcal{E}} & 0 \\ 0 & 0 \end{bmatrix} \right) \right) \right] - \frac{1}{2} \left[ \Phi_{\tilde{\mathbb{A}}} \left( \text{Ch} \left( \begin{bmatrix} 0 & 0 \\ 0 & 1_{\mathcal{E}} \end{bmatrix} \right) \right) \right] \\ &= \left[ \text{Str} \left( e^{-\frac{u\tilde{\theta}^{\mathbb{A}}_{u-1}}{2\pi i}} \right) \right]. \end{aligned}$$

□



### 4.3. The local index theorem.

**Theorem 4.8.** *Let  $D$  be a projective family of Dirac operators on a horizontally  $\mathcal{L}$ -twisted Clifford module  $\mathcal{E}$ . We have the following equality of classes in  $H_{\mathcal{L}}^{\bullet}(B)$ :*

$$\left[ \text{Str} \left( e^{-\frac{u}{2\pi i} \theta^{\mathbb{A}} u^{-1}} \right) \right] = \left[ u^{-\frac{k}{2}} \int_{M|B} \widehat{A} \left( \frac{u}{2\pi i} R^{M|B} \right) \text{Ch}_{\mathcal{L}}(\mathcal{E}/\mathcal{S}) \right].$$

*Proof.* Over each open set  $U_{\alpha}$  we have  $\theta^{\mathbb{A}} u^{-1} = \mathbb{A}_{\alpha, u^{-1}}^2 + \pi^* \omega_{\alpha}$ . Since  $\mathbb{A}_{\alpha, u^{-1}}^2$  and  $\pi^* \omega_{\alpha}$  commute, we have

$$\text{Str} e^{-\frac{u}{2\pi i} \theta^{\mathbb{A}} u^{-1}} = e^{-\frac{u}{2\pi i} \pi^* \omega_{\alpha}} \text{Str} e^{-\frac{u}{2\pi i} \mathbb{A}_{\alpha, u^{-1}}^2}.$$

According to Bismut's local index theorem for families [5]

$$\lim_{t \rightarrow 0} \text{Str} e^{-\frac{1}{2\pi i} \mathbb{A}_{\alpha, t}^2} = \int_{\pi^{-1} U_{\alpha} | U_{\alpha}} \widehat{A} \left( \frac{1}{2\pi i} R^{M|B} \right) \text{str}_{\mathcal{E}/\mathcal{S}} e^{-\frac{1}{2\pi i} \theta_{\alpha}^{\mathcal{E}/\mathcal{S}}}.$$

Moreover, by the result of Bismut and Fried [6]

$$(4.3.1) \quad \text{Str} e^{-\frac{1}{2\pi i} \mathbb{A}_{\alpha}^2} - \int_{\pi^{-1} U_{\alpha} | U_{\alpha}} \widehat{A} \left( \frac{1}{2\pi i} R^{M|B} \right) \text{str}_{\mathcal{E}/\mathcal{S}} e^{-\frac{1}{2\pi i} \theta_{\alpha}^{\mathcal{E}/\mathcal{S}}} = d \int_0^1 \xi_{\alpha}(t) dt$$

where

$$(4.3.2) \quad \xi_{\alpha}(t) = -\frac{1}{2\pi i} \text{Str} \frac{d\mathbb{A}_{\alpha, t}}{dt} e^{-\frac{1}{2\pi i} \mathbb{A}_{\alpha, t}^2}$$

is integrable at 0.

Let  $s > 0$  and let  $\delta_s^B$  be the operator on  $\Omega^*(B)$  which multiplies the forms of degree  $k$  by  $s^{k/2}$ . Then  $\delta_s^B \circ (\mathbb{A}_{\alpha, t}) \circ \delta_{s^{-1}}^B = s^{1/2} \mathbb{A}_{\alpha, t/s}$ . We therefore obtain

$$\begin{aligned} \lim_{t \rightarrow 0} \text{Str} e^{-\frac{s}{2\pi i} \mathbb{A}_{\alpha, t/s}^2} &= \delta_s^B \left( \lim_{t \rightarrow 0} \text{Str} e^{-\frac{1}{2\pi i} \mathbb{A}_{\alpha, t}^2} \right) \\ &= \delta_s^B \int_{\pi^{-1} U_{\alpha} | U_{\alpha}} \widehat{A} \left( \frac{1}{2\pi i} R^{M|B} \right) \text{str}_{\mathcal{E}/\mathcal{S}} e^{-\frac{1}{2\pi i} \theta_{\alpha}^{\mathcal{E}/\mathcal{S}}} \\ &= s^{-\frac{k}{2}} \int_{\pi^{-1} U_{\alpha} | U_{\alpha}} \widehat{A} \left( \frac{s}{2\pi i} R^{M|B} \right) \text{str}_{\mathcal{E}/\mathcal{S}} e^{-\frac{s}{2\pi i} \theta_{\alpha}^{\mathcal{E}/\mathcal{S}}} \end{aligned}$$

where, as before,  $k = \dim M - \dim B$ . Since both sides are polynomials in  $s$  we deduce that

$$\lim_{t \rightarrow 0} \text{Str} e^{-\frac{u}{2\pi i} \mathbb{A}_{\alpha, u^{-1}t}^2} = u^{-\frac{k}{2}} \int_{\pi^{-1} U_{\alpha} | U_{\alpha}} \widehat{A} \left( \frac{u}{2\pi i} R^{M|B} \right) \text{str}_{\mathcal{E}/\mathcal{S}} e^{-\frac{u}{2\pi i} \theta_{\alpha}^{\mathcal{E}/\mathcal{S}}}$$

Multiplying both sides by  $e^{-\frac{u}{2\pi i} \pi^* \omega_{\alpha}}$  we obtain

$$\lim_{t \rightarrow 0} \text{Str} \exp \left( -\frac{u}{2\pi i} \theta^{\mathbb{A}} u^{-1} t \right) \Big|_{U_{\alpha}} = u^{-\frac{k}{2}} \int_{\pi^{-1} U_{\alpha} | U_{\alpha}} \widehat{A} \left( \frac{u}{2\pi i} R^{M|B} \right) \text{Ch}_{\mathcal{L}}(\mathcal{E}/\mathcal{S})$$

and therefore

$$\lim_{t \rightarrow 0} \text{Str} \exp \left( -\frac{u}{2\pi i} \theta^{\mathbb{A}} u^{-1} t \right) = u^{-\frac{k}{2}} \int_{M|B} \widehat{A} \left( \frac{u}{2\pi i} R^{M|B} \right) \text{Ch}_{\mathcal{L}}(\mathcal{E}/\mathcal{S}).$$

Moreover by (4.3.1), we have

$$(4.3.3) \quad \text{Str} e^{-\frac{u}{2\pi i} \mathbb{A}_{\alpha, u^{-1}}^2} - u^{-\frac{k}{2}} \int_{\pi^{-1}U_\alpha|U_\alpha} \widehat{A}\left(\frac{u}{2\pi i} R^{M|B}\right) \text{str}_{\mathcal{E}/\mathcal{S}} e^{-\frac{u}{2\pi i} \theta_\alpha^{\mathcal{E}/\mathcal{S}}} = ud \int_0^1 u^{-\frac{1}{2}} \delta_u^B \xi_\alpha(t) dt$$

Here the right hand side is defined as follows. Write  $\xi_\alpha^{[l]} \in \Omega^l(B)$  for the component of degree  $l$  of  $\xi_\alpha$  from the equation (4.3.2). Then we have

$$u^{-\frac{1}{2}} \delta_u^B \xi_\alpha(t) = \sum u^{\frac{l-1}{2}} \xi_\alpha^{[l]}(t) = \frac{1}{2\pi i} \text{Str} \frac{d\mathbb{A}_{\alpha, u^{-1}t}}{dt} \exp\left(-\frac{u}{2\pi i} \mathbb{A}_{\alpha, u^{-1}t}^2\right) \in \Omega^*(B)[u].$$

Multiplying the identity (4.3.3) by  $e^{-\frac{u}{2\pi i} \pi^* \omega_\alpha}$  and using the equality

$$e^{-\frac{u}{2\pi i} \pi^* \omega_\alpha} ud(\cdot) = (ud + u^2 \Omega)(e^{-\frac{u}{2\pi i} \pi^* \omega_\alpha} \cdot)$$

we obtain

$$\text{Str} \exp\left(-\frac{u}{2\pi i} \theta_{u^{-1}}^{\mathbb{A}_{u^{-1}}}\right) \Big|_{U_\alpha} - u^{-\frac{k}{2}} \int_{\pi^{-1}U_\alpha|U_\alpha} \widehat{A}\left(\frac{u}{2\pi i} R^{M|B}\right) \text{Ch}_{\mathcal{L}}(\mathcal{E}/\mathcal{S}) = (ud + u^2 \Omega) \Xi_\alpha$$

where  $\Xi_\alpha = \int_0^1 \frac{1}{2\pi i} \text{Str} \frac{d\mathbb{A}_{\alpha, u^{-1}t}}{dt} \exp\left(-\frac{u}{2\pi i} \theta_{u^{-1}t}^{\mathbb{A}_{u^{-1}}}\right) dt$ .

From the equations (4.2.1) it follows that over  $U_{\alpha\beta}$  we have  $\frac{d\mathbb{A}_{\alpha, u^{-1}t}}{dt} = \phi_{\alpha\beta} \left(\frac{d\mathbb{A}_{\beta, u^{-1}t}}{dt}\right)$ . Hence there exists a form  $\dot{\mathbb{A}}_t \in u^{-1/2} \Omega^*(B; \Psi_{\mathcal{L}}(M|B, \mathcal{E}))[u]$  such that  $\dot{\mathbb{A}}_t \Big|_{U_\alpha} = \frac{d\mathbb{A}_{\alpha, u^{-1}t}}{dt}$ .

Therefore setting  $\Xi = \int_0^1 \frac{1}{2\pi i} \text{Str} \dot{\mathbb{A}}_t \exp\left(-\frac{u}{2\pi i} \theta_{u^{-1}t}^{\mathbb{A}_{u^{-1}}}\right) dt \in \Omega^*(B)[u]$  we can write

$$\text{Str} \exp\left(-\frac{u}{2\pi i} \theta_{u^{-1}}^{\mathbb{A}_{u^{-1}}}\right) - u^{-\frac{k}{2}} \int_{M|B} \widehat{A}\left(\frac{u}{2\pi i} R^{M|B}\right) \text{Ch}_{\mathcal{L}}(\mathcal{E}/\mathcal{S}) = d_\Omega \Xi$$

and the statement of the Theorem follows.  $\square$

Combining results of the Theorems 4.6 and 4.8 we obtain the main theorem of this paper

**Theorem 1.1.** *Let  $D$  be a projective family of Dirac operators on a horizontally  $\mathcal{L}$ -twisted Clifford module  $\mathcal{E}$  on a fibration  $\pi : M \rightarrow B$ . Then the following equality holds in  $H_{\mathcal{L}}^\bullet(B)$ :*

$$[\Phi_{\nabla^{\mathcal{H}}}(\text{ch}(\text{ind } D^+))] = \left[ u^{-\frac{k}{2}} \int_{M|B} \widehat{A}\left(\frac{u}{2\pi i} R^{M|B}\right) \text{Ch}_{\mathcal{L}}(\mathcal{E}/\mathcal{S}) \right],$$

where  $k = \dim M - \dim B$  is the dimension of the fibers.

## REFERENCES

- [1] Atiyah, M. F. and Singer, I. M. *The index of elliptic operators. IV.* Ann. of Math. (2) 93 1971 119–138.
- [2] Benameur, M.-T. and Carey A. *On the analyticity of the bivariant JLO cocycle.* Electron. Res. Announc. Math. Sci. 16 (2009), 37–43.
- [3] Benameur, M.-T. and Heitsch, J. L. *Index theory and non-commutative geometry. II. Dirac operators and index bundles.* J. K-Theory 1 (2008), no. 2, 305–356.
- [4] Berline, N., Getzler, E. and Vergne, M. *Heat kernels and Dirac operators.* Corrected reprint of the 1992 original. Grundlehren Text Editions. Springer-Verlag, Berlin, 2004.
- [5] Bismut, J.-M., *The Atiyah-Singer index theorem for families of Dirac operators: two heat equation proofs.* Invent. Math. 83 (1985), no. 1, 91–151.

- [6] Bismut, J.-M. and Freed, D. S., *The analysis of elliptic families. II. Dirac operators, eta invariants, and the holonomy theorem.* Comm. Math. Phys. 107 (1986), no. 1, 103–163.
- [7] Bott, R. and Tu, L. W. *Differential forms in algebraic topology.* Graduate Texts in Mathematics, 82. Springer-Verlag, New York-Berlin, 1982.
- [8] Brylinski, J.-L. *Loop spaces, characteristic classes and geometric quantization.* Modern Birkhuser Classics. Birkhuser Boston, Inc., Boston, MA, 2008.
- [9] Connes, A. *Entire cyclic cohomology of Banach algebras and characters of  $\theta$ -summable Fredholm modules.* K-Theory 1 (1988), no. 6, 519–548.
- [10] Connes, A. *Noncommutative differential geometry.* Inst. Hautes études Sci. Publ. Math. No. 62 (1985), 257–360.
- [11] Connes, A. *Noncommutative geometry.* Academic Press, Inc., San Diego, CA, 1994.
- [12] Dixmier, J. and Douady, A., *Champs continus d'espaces hilbertiens et de  $C^*$ -algèbres.* Bull. Soc. Math. France 91 1963 227–284.
- [13] Getzler, E. and Szenes, A. *On the Chern character of a theta-summable Fredholm module.* J. Funct. Anal. 84 (1989), no. 2, 343–357.
- [14] Gorokhovskiy, A. *Characters of cycles, equivariant characteristic classes and Fredholm modules.* Comm. Math. Phys. 208 (1999), no. 1, 1–23.
- [15] Gorokhovskiy, A. *Bivariant Chern character and longitudinal index.* J. of Funct. Analysis 237 (2006), 105–134.
- [16] Gorokhovskiy, A. and Lott, J. *Local index theory over étale groupoids.* J. Reine Angew. Math. 560 (2003), 151–198.
- [17] Hitchin, N. J. *The moduli space of complex Lagrangian submanifolds.* Surveys in differential geometry, 327–345, Surv. Differ. Geom., VII, Int. Press, Somerville, MA, 2000.
- [18] Khalkhali, M. *On the entire cyclic cohomology of Banach algebras.* Comm. Algebra 22 (1994), no. 14, 5861–5874.
- [19] Loday, J.-L., *Cyclic homology.* Grundlehren der Mathematischen Wissenschaften, 301. Springer-Verlag, Berlin, 1992.
- [20] Mathai, V.; Melrose, R. B. and Singer, I M. *The index of projective families of elliptic operators.* Geom. Topol. 9 (2005), 341–373.
- [21] Mathai, V.; Melrose, R. B. and Singer, I M. *The index of projective families of elliptic operators: the decomposable case.* Asterisque, 327 (2009) 251–292
- [22] Mathai, V. and Stevenson, D. *On a generalized Connes-Hochschild-Kostant-Rosenberg theorem.* Adv. Math. 200 (2006), no. 2, 303–335.
- [23] Meyer, R., *Local and analytic cyclic homology.* EMS Tracts in Mathematics, 3. European Mathematical Society (EMS), Zurich, 2007.
- [24] Murray, M. K. *Bundle gerbes.* J. London Math. Soc. (2) 54 (1996), no. 2, 403–416.
- [25] Nest R. and Tsygan B. *Algebraic index theorem for families.* Adv. Math. 113 (1995), no. 2, 151–205.
- [26] Nest R. and Tsygan B. *On the cohomology ring of an algebra.* Advances in geometry, 337–370, Progr. Math., 172, Birkhuser Boston, Boston, MA, 1999.
- [27] Nistor, V. *Super-connections and non-commutative geometry.* Cyclic cohomology and noncommutative geometry (Waterloo, ON, 1995), 115–136, Fields Inst. Commun., 17, Amer. Math. Soc., Providence, RI, 1997.

UMR 7122 DU CNRS, UNIVERSITÉ DE METZ, ILE DU SAULCY, METZ, FRANCE

*E-mail address:* `benameur@univ-metz.fr`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO, BOULDER, CO 80305-0395, USA

*E-mail address:* `Alexander.Gorokhovskiy@colorado.edu`