


Max-Planck-Institut für Mathematik Bonn

Quantum polynomial functors from e -Hecke algebras

by

Valentin Buciumas
Hankyung Ko



Quantum polynomial functors from e -Hecke algebras

Valentin Buciumas
Hankyung Ko

Max-Planck-Institut für Mathematik
Vivatsgasse 7
53111 Bonn
Germany

Mathematisches Institut
Universität Bonn
Endenicher Allee 60
53115 Bonn
Germany

QUANTUM POLYNOMIAL FUNCTORS FROM e -HECKE ALGEBRAS

VALENTIN BUCIUMAS AND HANKYUNG KO

ABSTRACT. We define a new category of quantum polynomial functors extending the quantum polynomials introduced by Hong and Yacobi. We show that our category has many properties of the category of Hong and Yacobi and is the natural setting in which one can define composition of quantum polynomial functors. Throughout the paper we highlight several key differences between the theory of classical and quantum polynomial functors.

1. INTRODUCTION

Hong and Yacobi [HY17] introduced a category of quantum polynomial functors which quantizes the strict polynomial functors of Friedlander and Suslin [FS97]. The purpose of this paper is to introduce higher level categories of quantum polynomial functors that extend the construction of Hong and Yacobi and explain why they are a natural candidate for the quantization of classical polynomial functors. The most visible advantage of our definition is that we are now able to compose quantum polynomial functors.

In order to explain why our quantization is natural, let us first describe what polynomial functors are. They were introduced by Friedlander and Suslin [FS97] and became an important tool in the study of rational cohomology of algebraic groups. Polynomial functors are defined as the category of functors between vector spaces which are polynomial on the space of morphisms. The definition can be formulated as follows. Let k be a field, and let \mathcal{V} be the category of finite dimensional vector spaces over k . The d -th symmetric group S_d acts on $V^{\otimes d}$ by permuting tensor factors. Let $\Gamma^d \mathcal{V}$ be the divided power category; it contains the same objects as \mathcal{V} and the morphisms are for $V, W \in \mathcal{V}$

$$(1) \quad \text{Hom}_{\Gamma^d \mathcal{V}}(V, W) = \Gamma^d \text{Hom}(V, W) = (\text{Hom}(V, W)^{\otimes d})^{S_d}.$$

The category \mathcal{P}^d of polynomial functors of degree d is the category of linear functors $F : \Gamma^d \mathcal{V} \rightarrow \mathcal{V}$. A polynomial functor is by definition an object in \mathcal{P} , the direct sum of \mathcal{P}^d over all $d \geq 0$.

The category \mathcal{P} has initially been introduced by [FS97] to prove the cohomological finite generation of finite group schemes over a field, where they use certain rational cohomology computations for the general linear groups effectively done in the category of polynomial functors. Another example of important rational cohomology result obtained from the polynomial functors is the untwisting of Frobenius due to Chałupnik [Cha15] and Touzé [Tou13]. Polynomial functors can also be used to compute cohomology for other classical algebraic groups (see [Tou10] and references therein). In a different direction, Hong, Touzé and Yacobi ([HTY14], [HY13]) showed that there is a categorical action of $\widehat{\mathfrak{sl}}_p$ on \mathcal{P} that categorifies the action of $\widehat{\mathfrak{sl}}_p$ on the Fock space, where p is the characteristic of the base field.

Two instrumental properties in the theory of polynomial functors are representability and composition. Representability allows one to prove that the category \mathcal{P}^d is equivalent to the category of modules of the Schur algebra $S(n; d)$ for any $n \geq d$, or the category of polynomial GL_n representations of degree d ([FS97, §3]). Composition is a natural property of functors, which is not so natural for modules over a Schur algebra, hence is a main advantage of studying polynomial functors. It is extremely useful in performing many cohomology calculations. Given $F, G \in \mathcal{P}$ and an exact

sequence in \mathcal{P} that represents an element of $\text{Ext}_{\mathcal{P}}^n(F, G)$, precomposing (respectively, postcomposing, when it is exact) the exact sequence with another polynomial functor $H \in \mathcal{P}$ gives a class in $\text{Ext}_{\mathcal{P}}^n(FH, GH)$ (resp., $\text{Ext}_{\mathcal{P}}^n(HF, HG)$). The special case of precomposing by the Frobenius functor, where the assignment is injective, is particularly interesting in many contexts (see [FFSS99]). In general, this can relate Ext spaces in different degrees and provides, for example, [FS97, Theorem 2.13] which is a key technique in main computations in [FS97].

A natural question is whether one can deform the polynomial functors into quantum polynomial functors. A first quantization of polynomial functors was developed by Hong and Yacobi [HY17]. They introduced a new category \mathcal{P}_q^d that is a q -deformation of the category \mathcal{P}^d and showed that it enjoys many properties that \mathcal{P}^d has. As in the classical case, \mathcal{P}_q^d is a monoidal category (which is now braided, as opposed to symmetric) that is equivalent to the category of modules for the quantum Schur algebra $S_q(n; d)$ of Dipper and James [DJ89]. They prove several applications to quantum invariant theory, including a quantum version of $(\text{GL}_n, \text{GL}_m)$ -Howe duality. However, their category of quantum polynomial functors does not allow for composition of the functors.

The underlying reason is that the action of S_d in equation (1) is replaced by a quantum action that depends on the extra structure given to the objects. In [HY17], the domain category of \mathcal{P}_q^d consists of pairs (V_n, R_n) , where one can think of V_n as the defining n -dimensional $U_q(\mathfrak{gl}_n)$ -module or the defining $A_q(n, n)$ -comodule (where $A_q(n, n)$ is the quantization of the coordinate ring of $n \times n$ matrices). The generators of the braid group \mathcal{B}_d act on $V_n^{\otimes d}$ via R -matrices R_n , the corresponding R -matrix of $U_q(\mathfrak{gl}_n)$ for the defining representation. When one applies a quantum polynomial F to V_n , there is an $U_q(\mathfrak{gl}_n)$ structure on $F(V_n)$ which produces an R -matrix $R_{F(V_n)}$ associated to $F(V_n)$. The problem is then that $R_{F(V_n)}$ and R_n are different, so the new pair $(F(V), R_{F(V)})$ is not in the domain category.

We can also see this in the classical case. The domain of a polynomial functor consists of vector spaces. One can endow such vector spaces with a GL_n -module structure. If V is a polynomial representation of degree e , then $F \in \mathcal{P}_q^d$ maps V to a polynomial representation $F(V)$ of degree de . So we can think of the domain in the classical case as containing finite dimensional polynomial representations of GL_n . The reason why this extra structure of V is not present in the definition of polynomial functors is because S_d acts on $V^{\otimes d}$ in the same way regardless of the GL_n structure of V . In other words, one can view a single polynomial functor as a functor between degree e modules and degree de modules for all $e \in \mathbb{N}$ at the same time. This is not true for quantum polynomial functors. The braid group acts on $V^{\otimes d}$ via R -matrices and different $U_q(\mathfrak{gl}_n)$ -modules have different associated R -matrices; therefore the action of \mathcal{B}_d on $V^{\otimes d}$ depends on the module structure of V . If V is the defining representation, the action of the braid group on $V^{\otimes d}$ factors through the Hecke algebra \mathcal{H}_d , but if V is a degree e polynomial representation, this action factors through the e -Hecke algebra $\mathcal{H}_{d,e}$ (see Definition 2.8). The e -Hecke algebras are quantizations of the symmetric group and together play the role the symmetric group plays in the classical case.

This extra structure should be taken into account in the quantum case. It is then natural to introduce the categories $\mathcal{P}_{q,e}^d$ of quantum polynomial functors that map degree e modules to degree de modules. In our setting we are able to compose quantum polynomials by Theorem 5.2. More precisely, given $G \in \mathcal{P}_{q,d_2}^{d_1}$ and $F \in \mathcal{P}_{q,d_2}^{d_1}$, their composition $F \circ G$ is a polynomial in $\mathcal{P}_{q,e}^{d_1 d_2}$.

We think of the categories $\mathcal{P}_{q,e}^d$ as “higher” analogues of \mathcal{P}_q^d . They satisfy many of the properties that \mathcal{P}_q^d satisfy. For example we show in Theorem 4.4 that the category $\mathcal{P}_{q,e}^d$ is a braided monoidal category. Another fundamental property we prove is that $\mathcal{P}_{q,e}^d$ is finitely generated when q is generic (Theorem 6.11). The reason why the root of unity case is significantly harder than the generic q case has to do with the domain category on which elements of $\mathcal{P}_{q,e}^d$ act which can be thought of as consisting of degree e polynomial representations of $U_q(\mathfrak{gl}_n)$. When q is a root of unity, this category is more complicated than the corresponding category for generic q . However, when $e = 1$, the

category of polynomial representations of degree 1 is “the same” for q a root of unity and for generic q ; it consists only of direct sums of the defining representation. Therefore our finite generation result holds for any q when $e = 1$ as we explain in Remark 6.12 and we get back the result of [HY17] in full generality. We also note that our proof for finite generation uses a different method than the one in [HY17]. We prefer this method since for $e > 1$ it allows one to clearly see why the requirement for $\Gamma_{q,e}^{d,V_m^{\otimes e}}$ to be a generator for $\mathcal{P}_{q,e}^d$ is $m \geq de$.

Another interesting difference between the quantum and the classical categories can be seen when taking into consideration e -Hecke pairs for all e at once. In Section 7 we define the category $\widetilde{\mathcal{P}}_q^d$ as the category of functors with domain all e -Hecke pairs for all e . For $q = 1$, this category is equivalent to the category $\mathcal{P}_{q,e}^d$ for any e . When q is generic, $\widetilde{\mathcal{P}}_q^d$ is not equivalent to $\mathcal{P}_{q,e}^d$ for any e .

We now briefly outline the content of the paper. In Section 2 we introduce the basics of quantum multilinear algebra which are of use throughout the paper. In Section 3 we define the categories $\mathcal{P}_{q,e}^d$, the main objects to be studied in this paper. We define a monoidal structure and a duality on the category \mathcal{P}_q^d . We then present several interesting examples of quantum polynomial functors. We focus on the quantum divided, symmetric and exterior power (due to Berenstein and Zwicknagl [BZ08]). We emphasize that the definition of these objects produce elements in $\mathcal{P}_{q,e}^d$ for each e , so their construction fits perfectly in our framework. In Section 4 we show the category of quantum polynomial functors is a braided monoidal category. In Section 5 we explain how two quantum polynomial functors can be composed in our setting. In Section 6 we prove that the category $\mathcal{P}_{q,e}^d$ has a (finite) projective generator for generic q . This immediately implies equivalence to the category of modules of a “generalized” q -Schur algebra. In Section 7 we consider a different category $\widetilde{\mathcal{P}}_q^d$ with domain all e -Hecke pairs for all $e \geq 0$ and we show that this category does not contain a projective generator for generic q . In Section 8 we present some remarks about quantum polynomial functors at roots of unity, including a conjecture for representability of $\mathcal{P}_{q,e}^d$ when $d = 2$.

Acknowledgements : We would like to thank Catharina Stroppel, Antoine Touzé and Daniel Tubbenhauer for many helpful discussions. We would like to thank Jiuzu Hong and Oded Yacobi for offering helpful suggestions to a preliminary version of the paper. We would like to thank the Max Planck Institute for Mathematics in Bonn for excellent working conditions where most of the work for this project was done.

2. PRELIMINARIES

Let k be a field. Let q be an element of k^\times . We say q is *generic* if q is not a root of unity. In this section we introduce several objects and prove some properties which will be of use in defining quantum polynomial functors. We note that several of these definitions and some of the properties are taken directly from [HY17].

2.1. Yang-Baxter spaces. Let \mathcal{V} be the category of finite dimensional vector spaces over the field k . Each $V \in \mathcal{V}$ comes with a chosen basis $\{v_1, \dots, v_n\}$ where $n = \dim(V)$. Even though our results are independent of the chosen basis, the exposition is more clear if we associate a fixed basis to each vector space.

Let $\tau : V \otimes W \rightarrow W \otimes V$ be the flip operator, namely $\tau(v \otimes w) = w \otimes v$. Let S_d be the symmetric group on d letters. Let \mathcal{B}_d be the Artin braid group generated by T_i , $1 \leq i \leq d - 1$ subject to the relations

$$(2) \quad \begin{aligned} T_i T_j &= T_j T_i \text{ if } |i - j| > 1 \\ T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1} \end{aligned}$$

The Hecke algebra \mathcal{H}_d is the quotient of the braid group \mathcal{B}_d by the relations

$$(T_i - q)(T_i + q^{-1}) = 0, \forall i.$$

For $V \in \mathcal{V}$, $R \in \text{End}(V \otimes V)$ is called an R -matrix if it satisfies the Yang-Baxter equation:

$$(3) \quad R_{12}R_{23}R_{12} = R_{23}R_{12}R_{23}$$

where $R_{12} = R \otimes 1_V \in \text{End}(V^{\otimes 3})$ and $R_{23} = 1_V \otimes R \in \text{End}(V^{\otimes 3})$.

If $R \in \text{End}(V \otimes V)$ is an R -matrix, we call the pair (V, R) a *Yang-Baxter space*. To each pair we can associate a right representation, $\rho_{d,V} : \mathcal{B}_d \rightarrow \text{End}(V^{\otimes d})$ that sends T_i to $1_{V^{\otimes i}} \otimes R \otimes 1_{V^{\otimes d-i-1}}$. We will most of the time use the short hand notation V for the Yang-Baxter space (V, R) and denote the R -matrix in the pair (V, R) by $R := R_V$.

We now define the quantum Hom-space algebra as it is defined by Hong and Yacobi [HY17]. Given two Yang-Baxter spaces V and W with basis $\{v_i\}$ and $\{w_j\}$, respectively, let $T(V, W)$ be the tensor algebra of $\text{Hom}(V, W)$, that is

$$T(V, W) = \bigoplus_{d \geq 0} T(V, W)_d$$

where $T(V, W)_d := \text{Hom}(V, W)^{\otimes d} \cong \text{Hom}(V^{\otimes d}, W^{\otimes d})$. Let $I(V, W)$ be the two sided ideal generated by $X \circ R_V - R_W \circ X$, for all $X \in \text{Hom}(V^{\otimes 2}, W^{\otimes 2})$. Define $A(V, W) := T(V, W)/I(V, W)$. The space $A(V, W)$ has a natural gradation

$$A(V, W) = \bigoplus_{d \geq 0} A(V, W)_d$$

where $A(V, W)_d = T(V, W)_d/I(V, W)_d$.

Denote by $x_{ji} : W \rightarrow V$ the map

$$x_{ji}(w_k) = \delta_{kj}v_i$$

with $x_{ji} \in \text{Hom}(W, V) \subset A(W, V)$.

Lemma 2.1. *The algebra $A(W, V)$ has a presentation by the generators x_{ji} and the relations generated by*

$$(4) \quad \sum_{k,l} (R_{W,kl}^{pq} x_{ki} x_{lj} - R_{V,ij}^{kl} x_{pk} x_{ql}),$$

where the coefficients $R_{V,ij}^{kl}$ are defined by the following equation:

$$R_V(v_i \otimes v_j) = \sum_{k,l} R_{V,ij}^{kl} v_k \otimes v_l.$$

Proof. Elements of the form $x_{ij}x_{kl}$ form a basis of $\text{Hom}(W^{\otimes 2}, V^{\otimes 2})$. The quadratic relations (4) are exactly the relations that generate $R(W, V)$. Since $R(W, V)$ generates $I(W, V)$, the result follows. \square

There is a degree preserving morphism of algebras

$$\Delta_{V,W,U} : A(V, U) \rightarrow A(V, W) \otimes A(W, U)$$

that is given on generators by $\Delta_{V,W,U}(x_{ij}) = \sum_k x_{ik} \otimes x_{kj}$. There is a map $V \rightarrow W \otimes \text{Hom}(W, V)$ given by

$$v_i \mapsto \sum_j w_j \otimes x_{ji}.$$

This extends to a map $\Delta_{V,W} : V \rightarrow W \otimes A(W, V)$.

Proposition 2.2. *The following diagram commutes:*

$$\begin{array}{ccc} V & \xrightarrow{\Delta_{V,W}} & W \otimes A(W, V) \\ \Delta_{V,U} \downarrow & & \downarrow \Delta_{W,U} \otimes 1 \\ U \otimes A(U, V) & \xrightarrow{1 \otimes \Delta_{U,W,V}} & U \otimes A(U, W) \otimes A(W, V) \end{array}$$

Proof. One can compute $(1 \otimes \Delta_{U,W,V})\Delta_{V,U}(v_i) = \sum_{k,j} u_j \otimes x_{jk} \otimes x_{ki} = (\Delta_{W,U} \otimes 1)\Delta_{V,W}(v_i)$ from which the commutativity of the diagram follows. \square

Let (V, R_V) and (W, R_W) be Yang-Baxter spaces. We define the generalized (q -)Schur algebra

$$S(V, W; d) := (A(W, V)_d)^*$$

as in [HY17]. The following is proved in [HY17]:

Proposition 2.3. *Let V, W be Yang-Baxter spaces. Then there is a natural isomorphism*

$$S(V, W; d) \cong \text{Hom}_{\mathcal{B}_d}(V^{\otimes d}, W^{\otimes d})$$

Proof. See Proposition 2.7 in [HY17]. \square

By taking the dual of $\Delta_{V,W,U}$ we obtain a map

$$m_{U,W,V} : S(W, V; d) \otimes S(U, W; d) \rightarrow S(U, V; d).$$

There is a natural map $m'_{U,W,V} : \text{Hom}_{\mathcal{B}_d}(W^{\otimes d}, V^{\otimes d}) \otimes \text{Hom}_{\mathcal{B}_d}(U^{\otimes d}, W^{\otimes d}) \rightarrow \text{Hom}_{\mathcal{B}_d}(U^{\otimes d}, V^{\otimes d})$ that takes $f \otimes g \mapsto f \circ g$. The following Proposition shows they are the same map under the isomorphism in Proposition 2.3.

Proposition 2.4. *Given three Yang-Baxter spaces V, U, W , the following diagram commutes:*

$$\begin{array}{ccc} S(W, V; d) \otimes S(U, W; d) & \xrightarrow{\cong} & \text{Hom}_{\mathcal{B}_d}(W^{\otimes d}, V^{\otimes d}) \otimes \text{Hom}_{\mathcal{B}_d}(U^{\otimes d}, W^{\otimes d}) \\ m_{U,W,V} \downarrow & & \downarrow m'_{U,W,V} \\ S(U, V; d) & \xrightarrow{\cong} & \text{Hom}_{\mathcal{B}_d}(U^{\otimes d}, V^{\otimes d}) \end{array}$$

Proof. See Proposition 2.8 in [HY17]. \square

Remark 2.5. If $W = V$, the quadratic relation (4) becomes the RTT relation due to Faddeev, Reshetikhin and Taktajan. The algebra $A(V, V)$ is then just the algebra denoted by A_{R_V} in [FRT88].

We record two properties of $A(V, V)$ which are standard results in the theory of quantum matrices; their proofs are nothing more than simple computations.

- (1) $A(V, V)$ is a bialgebra with comultiplication $\Delta_{V,V,V}(x_{ij}) = \sum_k x_{ik} \otimes x_{kj}$ and counit $e(x_{ij}) = e_{A(V,V)}(x_{ij}) = \delta_{ij}$.
- (2) V is an $A(V, V)$ -comodule with coaction given by $\Delta_{V,V}(v_i) = \sum_j v_j \otimes x_{ji}$.

We note that one of the two diagrams that need to commute for $\Delta_{V,V}$ to be a coaction is the diagram in Proposition 2.2 for $V = W = U$. Therefore we can think of the map $\Delta_{V,W} : V \rightarrow W \otimes A(W, V)$ as a generalization of the coaction.

Since the comultiplication is degree preserving (i.e., $\Delta_{V,V,V}$ maps $A(V, V)_d$ to $A(V, V)_d \otimes A(V, V)_d$), the map $m_{V,V,V}$ makes $S(V, V; d)$ into an algebra. The associativity of $m_{V,V,V}$ is equivalent to the coassociativity of $\Delta_{V,V,V}$. The unit of $S(V, V; d)$ is given by the counit e of $A(V, V)$.

2.2. Quantum matrix spaces and e -Hecke pairs. Let V_n denote an n -dimensional vector space. Let R_n be the R -matrix of $U_q(\mathfrak{gl}_n)$ for the defining representation, namely

$$(5) \quad R_n(v_i \otimes v_j) = \begin{cases} v_j \otimes v_i & \text{if } i < j \\ qv_i \otimes v_j & \text{if } i = j \\ (q - q^{-1})v_i \otimes v_j + v_j \otimes v_i & \text{if } i > j. \end{cases}$$

It is well known that (V_n, R_n) form a Yang-Baxter space. Define $A_q(n, n) := A(V_n, V_n)$.

The space $A_q(n, n)$ is the algebra of quantum $n \times n$ matrices (see [Tak02] and [FRT88]). It is a coquasitriangular bialgebra generated by elements $x_{ij}, 1 \leq i, j \leq n$ subject to the following RTT relations:

$$(6) \quad \sum_{k,l} (R_n)_{kl}^{pq} x_{ki} x_{lj} = \sum_{k,l} (R_n)_{ij}^{kl} x_{pk} x_{ql}.$$

We now present some standard properties of $A_q(n, n)$, see for example [LR97, Chapter 7]. We begin by reminding the reader about the coalgebra structure on $A_q(n, n)$. The comultiplication and counit are given on generators by

$$\Delta(x_{ij}) = \sum_k x_{ik} \otimes x_{kj}, \quad \epsilon(x_{ij}) = \delta_{ij}.$$

The vector space V_n is an $A_q(n, n)$ -comodule via the coaction $v_i \mapsto \sum_j v_j \otimes x_{ji}$.

The coalgebra structure on $A_q(n, n)$ allows one to endow the tensor product $V \otimes W$ of two $A_q(n, n)$ -comodules with the structure of an $A_q(n, n)$ -comodule. Therefore the category of finite dimensional $A_q(n, n)$ -comodules is a monoidal category. The unit is the trivial comodule k with the coaction $1 \in k \mapsto 1 \otimes 1 \in k \otimes A_q(n, n)$. There are standard isomorphisms $l_V : k \otimes V \rightarrow V$ and $r_V : V \otimes k \rightarrow V$.

The bialgebra $A_q(n, n)$ is coquasitriangular. This means that there is a map $\mathcal{R} : A_q(n, n) \otimes A_q(n, n) \rightarrow k$ that is invertible in the convolution algebra, satisfying the following conditions:

$$(7) \quad \begin{aligned} \mathcal{R}(ab, c) &= \mathcal{R}(a, c_{(1)})\mathcal{R}(b, c_{(2)}) \\ \mathcal{R}(a, bc) &= \mathcal{R}(a_{(1)}, b)\mathcal{R}(a_{(2)}, c) \\ b_{(1)}a_{(1)}\mathcal{R}(a_{(2)}, b_{(2)}) &= \mathcal{R}(a_{(1)}, b_{(1)})b_{(2)}a_{(2)} \end{aligned}$$

for all $a, b, c \in A_q(n, n)$. In the above formula we use Sweedler notation, namely we denote $\Delta(a) = a_{(1)} \otimes a_{(2)}$. The map \mathcal{R} is given on generators x_{ij} by the formula

$$(8) \quad \mathcal{R}(x_{ij} \otimes x_{kl}) = (R_n)_{lj}^{ik}.$$

The values of \mathcal{R} on higher order terms is given by repeated applications of the first two equalities in equation (7).

The existence of \mathcal{R} implies that for every $A_q(n, n)$ -comodules V, W there is an $A_q(n, n)$ -comodule isomorphism $R_{V,W} : V \otimes W \rightarrow W \otimes V$ given by the formula

$$(9) \quad R_{V,W} := (1 \otimes 1 \otimes \mathcal{R})(1 \otimes \tau \otimes 1)(\Delta_W \otimes \Delta_V)\tau.$$

This morphism makes the category of finite dimensional $A_q(n, n)$ -comodules into a braided monoidal category. A braided monoidal category \mathcal{C} is a monoidal category with braiding isomorphisms $\gamma_{V,W} : V \otimes W \rightarrow W \otimes V$ that satisfy

$$(10) \quad \begin{aligned} \gamma_{V \otimes W, U} &= (\gamma_{V,U} \otimes 1)(1 \otimes \gamma_{W,U}) \\ \gamma_{V, W \otimes U} &= (1 \otimes \gamma_{V,U})(\gamma_{V,W} \otimes 1) \\ \tilde{r}_V \gamma_{I,V} &= \tilde{l}_V, \quad \tilde{r}_V \gamma_{V,I} = \tilde{l}_V \end{aligned}$$

where I is the identity object in the monoidal category and \tilde{r}_V, \tilde{l}_V are the identity constraints in \mathcal{C} .

Proposition 2.6. *The category of finite dimensional $A_q(n, n)$ -comodules is a braided monoidal category with braiding isomorphisms given by $\gamma_{V,W} = R_{V,W}$.*

When $W = V$, the map $R_V := R_{V,V}$ satisfies the Yang-Baxter equation. This makes makes the pair (V, R_V) a Yang-Baxter space for every $A_q(n, n)$ -comodule V .

Definition 2.7. An *e-Hecke pair* V is an $A_q(n, n)$ -comodule for some $n \geq 1$ such that the image of the coaction $\Delta_V : V \rightarrow V \otimes A_q(n, n)$ lies in $V \otimes A_q(n, n)_e$.

We remark why we use the term “ e -Hecke pair”. Let V be an e -Hecke pair. First, we call it a “pair” because we think of V as the pair (V, R_V) . To explain the word “Hecke”, let us start with the case $e = 1$. Then V is a direct sum of defining comodules V_n and the action of the braid group \mathcal{B}_d on $V^{\otimes d}$ factors through the action of the Hecke algebra \mathcal{H}_d . This is not the case for general e . Instead, the action of \mathcal{B}_d on $V^{\otimes d}$ factors through a different deformation of the symmetric group S_d , which is realized as a subalgebra of \mathcal{H}_{de} as follows. Let w_i be the element in S_{de} such that

$$(11) \quad w_i(j) = \begin{cases} j + e & e(i-1) < j \leq ei \\ j - e & ei < j \leq e(i+1) \\ j & \text{otherwise.} \end{cases}$$

Definition 2.8. The e -Hecke algebra of rank d , denoted by $\mathcal{H}_{d,e}$, is the subalgebra of \mathcal{H}_{de} generated by $T_{w_1}, \dots, T_{w_{d-1}}$, where w_i are as in equation (11).

If V is an e -Hecke pair, the generator T_{w_i} acts on $V^{\otimes d}$ as the generator $T_i \in \mathcal{B}_d$ acts on $V^{\otimes d}$, and we have

$$\text{Hom}_{\mathcal{H}_{d,e}}(V^{\otimes d}, W^{\otimes d}) = \text{Hom}_{\mathcal{B}_d}(V^{\otimes d}, W^{\otimes d}) \cong (\text{Hom}(V, W)^{\otimes d})^{\mathcal{B}_d}.$$

Therefore, an e -Hecke pair has the property that its d -th tensor power is a module over the e -Hecke algebra $\mathcal{H}_{d,e}$.

The e -Hecke algebra arises when one considers an algebra version of the wreath product of finite groups. One can define the wreath product of finite groups G and $H \subseteq \mathcal{B}_d$, generalizing the usual construction using the symmetric group: Let the wreath product $G \wr H$ be the semidirect product $G^{\times d} \rtimes H$ (here the action of H is the braid group action permuting components, which factors through the symmetric group). If $G = \mathcal{B}_e$ and $H = \mathcal{B}_d$, then the wreath product $\mathcal{B}_e \wr \mathcal{B}_d$ is a subgroup of the larger braid group \mathcal{B}_{de} . Now we see how we replace the braid groups by Hecke algebras in constructing the wreath product. The group \mathcal{B}_e is replaced by the Hecke algebra H_e , and the product $\mathcal{B}_e^{\times d}$ is replaced by $H_e^{\otimes d}$. This latter algebra is a subalgebra of \mathcal{H}_{de} generated by $T_1, \dots, T_{e-1}, T_{e+1}, \dots, T_{2e-1}, \dots, T_{(d-1)e+1}, \dots, T_{de-1}$. The e -Hecke algebra $\mathcal{H}_{d,e}$ naturally acts on this; the generator T_{w_i} acts as the multiplication in H_{de} . Then the subalgebra in H_{de} generated by the above $H_e^{\otimes d}$ and our e -Hecke algebra $\mathcal{H}_{d,e}$, which is isomorphic to $H_e^{\otimes d} \otimes \mathcal{H}_{d,e}$ as a vector space, is an algebra version of the wreath product of $\mathcal{B}_e \wr \mathcal{B}_d$.

The last equality of equation (7) implies that R_V is an $A_q(n, n)$ -comodule homomorphism. Given an $A_q(n, n)$ -comodule V , write the coaction map as

$$\Delta_V : v_i \mapsto \sum_j v_j \otimes t_{ji}^V$$

for some $t_{ji}^V \in A_q(n, n)$. The equation above serves as the definition for t_{ji}^V .

Lemma 2.9. *The equation*

$$\sum_{k,l} R_{V,kl}^{pq} t_{ki}^V t_{lj}^V = \sum_{k,l} R_{V,ij}^{kl} t_{pk}^V t_{ql}^V.$$

is equivalent to the fact that R_V is an $A_q(n, n)$ -comodule homomorphism.

Proof. R_V is an $A_q(n, n)$ -comodule homomorphism if and only if

$$\Delta_{V \otimes V} R_V = (R_V \otimes 1) \Delta_{V \otimes V}.$$

Applying both sides to $v_i \otimes v_j$ and picking out the $A_q(n, n)$ -coefficients of $v_p \otimes v_q$ produces the the desired equation. □

Lemma 2.10. *The relation*

$$(12) \quad \sum_{k,l} R_{V,kl}^{pq} t_{ki}^V t_{lj}^V = \sum_{k,l} R_{V,ij}^{kl} t_{pk}^V t_{ql}^V.$$

holds in $A_q(n, n)$.

Proof. This follows from Lemma 2.9 and the fact that R_V is an $A_q(n, n)$ -comodule homomorphism. \square

Given an $A_q(n, n)$ -comodule V , we denote $A(V, V)$ by $A_q(V, V)$ and $S(V, V; d)$ by $S_q(V, V; d)$. If we define $S_q(n; d) := S_q(V_n, V_n; d)$, then $S_q(n; d)$ is the q -Schur algebra due to Dipper and James [DJ89].

Proposition 2.11. *Suppose q is generic or 1 and $\text{char } k = 0$. Then the category of finite dimensional $A_q(n, n)$ -comodules is semisimple.*

Proof. See Theorem 11.4.4 in [PW91] where our Proposition is proved for $\text{GL}_q(n)$. The same result for $A_q(n, n)$ follows in a similar way. \square

Proposition 2.12. *Suppose q is generic or 1 and $\text{char } k = 0$. Any irreducible $A_q(n, n)_e$ -comodule V is a direct summand of $V_n^{\otimes e}$.*

Proof. This is a well known result which follows for example from the work of Parshall and Wang [PW91]. \square

3. QUANTUM POLYNOMIAL FUNCTORS

3.1. Definition. In this section we propose a different definition of quantum polynomial functors that generalizes the definition in [HY17]. Our category of quantum polynomial functors enjoys many of the properties presented in [HY17], and additionally, it has a composition. Let d, e be non-negative integers.

Let us define the quantum divided power category $\Gamma_{q,e}^d \mathcal{V}$. Its objects are all e -Hecke pairs for all positive n . The morphisms are defined as

$$\text{Hom}_{\Gamma_{q,e}^d}(V, W) := \text{Hom}_{\mathcal{B}_d}(V^{\otimes d}, W^{\otimes d}).$$

Note that the category $\Gamma_{q,e}^d \mathcal{V}$ always contains a one dimensional e -Hecke pair that is obtained by tensoring the defining $A_q(1, 1)_1$ -comodule with itself e times. It then follows that the category contains an n -dimensional vector space for every positive integer n . When $d = 1$, the forgetful functor $(V, R) \mapsto V$ to finite dimensional vector spaces induces an equivalence of categories

$$(13) \quad \Gamma_{q,e}^1 \mathcal{V} \cong \mathcal{V}$$

for any q, e . When $q = 1$, the R -matrix R_V of any e -Hecke pair (V, R_V) is just the transpose map $V \otimes V \rightarrow V \otimes V$, $v \otimes w \mapsto w \otimes v$, thus we also have the equivalence

$$(14) \quad \Gamma_{1,e}^d \mathcal{V} \cong \Gamma^d \mathcal{V}$$

for any d, e , where $\Gamma^d \mathcal{V}$ is the domain category for the classical polynomial functors defined in the introduction.

Definition 3.1. A quantum polynomial functor of degree d on e -Hecke pairs is a linear functor

$$F : \Gamma_{q,e}^d \mathcal{V} \rightarrow \mathcal{V},$$

or equivalently (by equation (13)),

$$F : \Gamma_{q,e}^d \mathcal{V} \rightarrow \Gamma_{q,ed}^1 \mathcal{V}.$$

We denote by $\mathcal{P}_{q,e}^d$ the category of quantum polynomial functors of degree d on e -Hecke pairs. Morphisms are natural transformations of functors. The equivalence (14) tells us that we recover the category of polynomial functors \mathcal{P}^d of Friedlander-Suslin [FS97] (see Introduction) when $q = 1$.

There is an equivalent characterization of a polynomial functor both in the classical and Hong and Yacobi [HY17] setting, which directly applies to ours. Given $F \in \mathcal{P}_{q,e}^d$ and $V, W \in \Gamma_{q,e}^d \mathcal{V}$, we have a map

$$F_{V,W} : S_q(V, W; d) \rightarrow \text{Hom}(F(V), F(W))$$

which gives rise to two maps

$$F'_{V,W} : S_q(V, W; d) \otimes F(V) \rightarrow F(W)$$

$$F''_{V,W} : F(V) \rightarrow F(W) \otimes A_q(W, V)_d.$$

Proposition 3.2. ([HY17], Proposition 3.5) *A quantum polynomial functor F of degree d is equivalent to the following data:*

- (1) for each $V \in \Gamma_{q,e}^d$ a vector space $F(V)$;
- (2) given $V, W \in \Gamma_{q,e}^d$, a linear map

$$F''_{V,W} : F(V) \rightarrow F(W) \otimes A_q(W, V)_d$$

such that the following diagrams commute for any $V, W, U \in \Gamma_{q,e}^d$:

$$(15) \quad \begin{array}{ccc} F(V) & \xrightarrow{F''_{V,U}} & F(U) \otimes A_q(U, V)_d \\ F''_{V,W} \downarrow & & \downarrow 1 \otimes \Delta_{U,W,V} \\ F(W) \otimes A_q(W, V)_d & \xrightarrow{1 \otimes F''_{W,U}} & F(U) \otimes A_q(U, W)_d \otimes A_q(W, V)_d \end{array}$$

$$(16) \quad \begin{array}{ccc} F(V) & \xrightarrow{F''_{V,V}} & F(V) \otimes A_q(V, V)_d \\ \downarrow 1 & \swarrow 1 \otimes e & \\ F(V) \otimes k & & \end{array}$$

Proof. See Proposition 3.5 in [HY17]. We note that even though they prove it for when V, W, U are defining comodules, their proof goes through unchanged for general e -Hecke pairs as in our setting. \square

We can extend the map $F''_{V,V} : F(V) \rightarrow F(V) \otimes A_q(V, V)_d$ to a map $\Delta_{F(V)}^V : F(V) \rightarrow F(V) \otimes A_q(V, V)$ that satisfies the following property:

Lemma 3.3. *The map $\Delta_{F(V)}^V$ makes $F(V)$ into an $A_q(V, V)$ -comodule.*

Proof. The diagrams in Proposition 3.2 are exactly what is needed for $\Delta_{F(V)}^V$ to be a coaction map. \square

Let V be an e -Hecke pair. The bialgebra $A_q(V, V)$ is a quotient of the free algebra generated by $\{x_{ji}^V\}$ by the ideal generated by $\sum_{k,l} (R_{V,kl}^{pq} x_{ki}^V x_{lj}^V - R_{V,ij}^{kl} x_{pk}^V x_{ql}^V)$. The space V is an $A_q(V, V)$ -comodule via the coaction

$$\Delta_V^V : v_i \mapsto \sum_j v_j \otimes x_{ji}^V$$

and let

$$\Delta_V : v_i \mapsto \sum_j v_j \otimes t_{ji}^V$$

be the coaction map that makes V into an $A_q(n, n)$ -comodule.

Define the map $\psi_V : A_q(V, V) \rightarrow A_q(n, n)$ on the generators of $A_q(V, V)$ as follows:

$$(17) \quad \psi_V(x_{ji}^V) = t_{ji}^V.$$

Lemma 3.4. *The map ψ_V is a bialgebra homomorphism.*

Proof. We first need to check that ψ_V is well defined. We can do this by showing that :

$$\psi_V\left(\sum_{k,l} R_{V,kl}^{pq} x_{ki}^V x_{lj}^V - R_{V,ij}^{kl} x_{pk}^V x_{ql}^V\right) = \sum_{k,l} R_{V,kl}^{pq} t_{ki}^V t_{lj}^V - R_{V,ij}^{kl} t_{pk}^V t_{ql}^V = 0.$$

The first equality is by definition and the second one holds according to Lemma 2.10.

We now show that ψ_V is a coalgebra homomorphism. This is equivalent to showing that ψ_V commutes with the comultiplication and the counit, namely

$$\Delta_{A_q(n,n)} \psi_V = (\psi_V \otimes \psi_V) \Delta_{A_q(V,V)}.$$

$$e_{A_q(n,n)} \psi_V = e_{A_q(V,V)}$$

Both equations follows from the fact that Δ_V is a coaction. □

Let G be a quantum polynomial functor of degree d . We have maps:

$$G''_{V,W} : G(V) \rightarrow G(W) \otimes A_q(W, V)_d.$$

If we denote by $\{v_i^G\}$ and $\{w_j^G\}$ the bases of $G(V)$ and $G(W)$, respectively, the map $G''_{V,W}$ takes $v_i^G \mapsto w_j^G \otimes t_{ji}^{W,V}$ for $t_{ji}^{W,V} \in A_q(W, V)_d$. By the definition of $A_q(G(W), G(V))$ we get a map

$$G(V) \rightarrow G(W) \otimes A_q(G(W), G(V))$$

which takes

$$v_i^G \mapsto w_j^G \otimes x_{ji}^{W,V},$$

where $x_{ji}^{W,V}$ are the generators of $A_q(G(W), G(V))$.

Lemma 3.5. *The map*

$$(18) \quad \psi_{W,V}^G : A_q(G(W), G(V)) \rightarrow A_q(W, V)$$

defined on generators as $\psi_{W,V}^G(x_{ji}^{W,V}) := t_{ji}^{W,V}$ is well-defined. Therefore it is an algebra homomorphism.

Proof. The proof is similar to how we proved that ψ_V is well-defined in Lemma 3.4. We want

$$\psi_{W,V}^G\left(\sum_{k,l} R_{G(W),kl}^{pq} x_{ki}^{W,V} x_{lj}^{W,V} - R_{G(V),ij}^{kl} x_{pk}^{W,V} x_{ql}^{W,V}\right) = \sum_{k,l} R_{G(W),kl}^{pq} t_{ki}^{W,V} t_{lj}^{W,V} - R_{G(V),ij}^{kl} t_{pk}^{W,V} t_{ql}^{W,V} = 0.$$

The first equality is by definition. The second equality follows from the commutativity of the exterior square in the diagram

$$(19) \quad \begin{array}{ccccc} G(V) \otimes G(V) & \xrightarrow{(1 \otimes m)(1 \otimes \tau \otimes 1)G''_{V,W} \otimes 2} & G(W) \otimes G(W) \otimes A_q(G(W), G(V)) & \xrightarrow{1 \otimes \Psi_{W,V}^G} & G(W) \otimes G(W) \otimes A_q(W, V) \\ R_{G(V)} \downarrow & & R_{G(W)} \otimes 1 \downarrow & & R_{G(W)} \otimes 1 \downarrow \\ G(V) \otimes G(V) & \xrightarrow{(1 \otimes m)(1 \otimes \tau \otimes 1)G''_{V,W} \otimes 2} & G(W) \otimes G(W) \otimes A_q(G(W), G(V)) & \xrightarrow{1 \otimes \Psi_{W,V}^G} & G(W) \otimes G(W) \otimes A_q(W, V) \end{array}$$

where m is the multiplication in $A_q(GW, GV)$. The diagram is commutative because the two small squares are: the first by Lemma 2.9 and the second is trivial. \square

Lemma 3.6. *The maps $\psi_{W,V}^G$ satisfy the following two equations for all e -Hecke pairs U, V, W :*

$$(20) \quad \Delta_{U,W,V} \circ \psi_{U,V}^G = (\psi_{U,W}^G \otimes \psi_{W,V}^G) \circ \Delta_{G(U), G(W), G(V)}$$

$$(21) \quad e_{A_q(G(V), G(V))} = e_{A_q(V, V)} \psi_{V, V}^G$$

Proof. The first equation is equivalent to $\Delta_{U,W,V}(t_{ij}^{U,V}) = \sum_j t_{ik}^{U,W} \otimes t_{kj}^{W,V}$. This in turn follows from the commutativity of diagram (15).

The second equation is equivalent to $e_{A_q(V, V)}(t_{ij}^{W, V}) = \delta_{ij}$ which follows from the commutativity of diagram (16). \square

3.2. Basic operations on quantum polynomial functors. We denote by $\mathcal{P}_q^d := \bigoplus_e \mathcal{P}_{q,e}^d$ the category of quantum polynomial functors of degree d , by $\mathcal{P}_{q,e} := \bigoplus_d \mathcal{P}_{q,e}^d$ the category of quantum polynomial functors on e -Hecke pairs, and by $\mathcal{P}_q := \bigoplus_{e,d} \mathcal{P}_{q,e}^d$ the category of quantum polynomial functors.

3.2.1. Tensor product in $\mathcal{P}_{q,e}$. Given two quantum polynomials on e -Hecke pairs $F \in \mathcal{P}_{q,e}^d$ and $G \in \mathcal{P}_{q,e}^{d'}$, the (external) tensor product $F \otimes G \in \mathcal{P}_{q,e}^{d+d'}$ is defined in the same way as in [HY17]. For an e -Hecke pair V , we define it to be

$$F \otimes G(V) = F(V) \otimes G(V).$$

To define it on the morphisms one uses the inclusion $\mathcal{B}_d \times \mathcal{B}_{d'} \subset \mathcal{B}_{d+d'}$. To be more explicit, for two e -Hecke pairs V, W , the map $(F \otimes G)_{V,W}$ is the following composition:

$$\begin{aligned} \text{Hom}_{\mathcal{B}_{d+d'}}(V^{\otimes d+d'}, W^{\otimes d+d'}) &\hookrightarrow \text{Hom}_{\mathcal{B}_d \times \mathcal{B}_{d'}}(V^{\otimes d} \otimes V^{\otimes d'}, W^{\otimes d} \otimes W^{\otimes d'}) \\ &\rightarrow \text{Hom}_{\mathcal{B}_d}(V^{\otimes d}, W^{\otimes d}) \otimes \text{Hom}_{\mathcal{B}_{d'}}(V^{\otimes d'}, W^{\otimes d'}) \\ &\rightarrow \text{Hom}(F(V), F(W)) \otimes \text{Hom}(G(V), G(W)) \\ &\rightarrow \text{Hom}(F(V) \otimes G(V), F(W) \otimes G(W)) \end{aligned}$$

where the third map is $F_{V,W} \otimes G_{V,W}$.

Recall the constant functor $k \in \mathcal{P}_{q,e}^0$. It maps an e -Hecke pair $V \mapsto k$, where k is the trivial $A_q(n, n)$ -comodule. It is then an easy exercise using the definition above to show that $F \otimes k \cong F$ via the natural transformation $\eta : F \otimes k \rightarrow F$ given by the standard isomorphism $\eta_V : F(V) \otimes k \cong F(V)$. We similarly have that $k \otimes F \cong F$. This can be summarized as follows:

Proposition 3.7. *The category $\mathcal{P}_{q,e}$ is a monoidal category with the tensor product \otimes and the unit object k .*

3.2.2. *Duality in $\mathcal{P}_{q,e}^d$.* One defines a duality on the functor category using dualities on the domain and codomain categories. In \mathcal{V} , we have the linear dual $V \mapsto V^* = \text{Hom}_k(V, k)$. For our category $\Gamma_{q,e}^d \mathcal{V}$ where the objects have additional structures, we “lift” the linear dual to what is compatible with this structure, namely the twisted dual. Note that an $A_q(n, n)$ -comodule can be thought of as a polynomial representation for the corresponding quantum group $U_q(\mathfrak{gl}_n)$. For a $U_q(\mathfrak{gl}_n)$ -module V , one can define a twisted $U_q(\mathfrak{gl}_n)$ structure on the linear dual V^* of the underlying vector space by precomposing the $U_q(\mathfrak{gl}_n)$ action by an antiautomorphism τ_1 (see [Jan96, 9.20] for the definition of τ_1 and the twisted dual). We denote by ${}^\tau V$ the resulting $U_q(\mathfrak{gl}_n)$ -module. Then we have

$$(22) \quad {}^\tau({}^\tau V) \cong V$$

for any $U_q(\mathfrak{gl}_n)$ -module V . We remark that ${}^\tau -$ is a duality of a highest weight category, under which the irreducibles are self-dual and a standard module and a costandard module of the same highest weight are dual to each other. In particular, the duality preserves the degree of polynomial representations, that is, if V is an $U_q(\mathfrak{gl}_n)$ -module of degree e , then the dual ${}^\tau V$ is also a degree e $U_q(\mathfrak{gl}_n)$ -module, so we can extend this duality on the category of $A_q(n, n)$ -comodules. We therefore have a contravariant functor

$${}^\tau - : \Gamma_{q,e}^d \mathcal{V} \rightarrow \Gamma_{q,e}^d \mathcal{V}.$$

Now we define the duality $-^\#$ on $\mathcal{P}_{q,e}^d$ as

$$F^\# := {}^\tau F({}^\tau -).$$

It is not hard to show that $F^\#$ will satisfy the properties in Proposition 3.2 and therefore it is a quantum polynomial functor. Taking $q = 1$ in our setting agrees with the classical definition of the dual.

3.3. **Examples.** We now present several examples of quantum polynomial functors.

3.3.1. *Tensor powers.* For each $d \in \mathbb{N}$, the d -th tensor product functor $\otimes^d : V \mapsto V^{\otimes d}$ is a quantum polynomial of degree d . To be more precise, for each e , there is a functor \otimes_e^d in $\mathcal{P}_{q,e}^d$ which maps an e -Hecke pair V to the de -Hecke pair $V^{\otimes d}$. The map $\otimes_{V,W}^d$ on the morphisms is just the inclusion

$$\text{Hom}_{\mathcal{B}_d}(V, W) \hookrightarrow \text{Hom}(V^{\otimes d}, W^{\otimes d}).$$

We abusively denote this functor by \otimes^d for any e .

When $d = 0$ we get the constant polynomial functor $\otimes^0 : V \mapsto k$, which we denote abusively by k , and \otimes^1 is the identity functor which we prefer to denote by I .

3.3.2. *Divided powers and symmetric powers.* Given an e -Hecke pair W , the divided power $\Gamma_{q,e}^{d,W}$ is the object in $\mathcal{P}_{q,e}^d$ represented by W , namely, $\Gamma_{q,e}^{d,W} : V \mapsto \text{Hom}_{\mathcal{B}_d}(W^{\otimes d}, V^{\otimes d})$. The map on morphisms

$$\text{Hom}_{\mathcal{B}_d}(V_1^{\otimes d}, V_2^{\otimes d}) \rightarrow \text{Hom}(\text{Hom}_{\mathcal{B}_d}(W^{\otimes d}, V_1^{\otimes d}), \text{Hom}_{\mathcal{B}_d}(W^{\otimes d}, V_2^{\otimes d}))$$

is given by $f \mapsto f \circ -$. When W is the e -Hecke pair $(k^{\otimes e}, q^{e^2})$, we denote it by $\Gamma_{q,e}^d$ and call it the divided power. We may drop the index “ e ” in the notation because it is determined by V . That is, if V is an e -Hecke pair, then $\Gamma_q^{d,V}$ denotes the functor $\text{Hom}_{\Gamma_{q,e}^d \mathcal{V}}(V, -)$ in $\mathcal{P}_{q,e}^d$. Since a divided power is a representable functor, Yoneda’s lemma tells us that it is a projective object in $\mathcal{P}_{q,e}^d$ (see Proposition 6.1 for a detailed proof). Moreover, divided powers form a set of projective generators for $\mathcal{P}_{q,e}^d$ for generic q as we shall prove in Theorem 6.11.

For an e -Hecke pair V let $S_V^d \in \mathcal{P}_{q,e}^d$ be defined by

$$S_V^d := (\Gamma^{d,V})^\#$$

where the dual $F^\#$ of a functor F is defined in §3.2.2. These are the corepresentable objects in $\mathcal{P}_{q,e}^d$ and form a set of injective cogenerators for $\mathcal{P}_{q,e}^d$.

3.3.3. Quantum symmetric, divided and exterior powers. In this subsection we assume $\text{char}(k) = 0$. Let V be an e -Hecke pair. An important set of examples of quantum polynomial functors are the quantum symmetric and exterior powers for generic q due to Berenstein and Zwicknagl [BZ08] (they also require $\text{char}(k) = 0$). These are quantum deformations of the classical symmetric and exterior power, though their theory is significantly more complex. For example, it is not necessary that the dimension of the quantum symmetric power of an $U_q(\mathfrak{gl}_n)$ -module to be the same as the dimension of the classical symmetric power of the corresponding $U(\mathfrak{gl}_n)$ -module.

In this subsection we define the quantum symmetric, divided and exterior powers when q is generic or when it is a p -th root of unity for an odd integer p . In the generic case, the definition of quantum symmetric and exterior can easily be seen to be the same as in [BZ08].

The map $R_V : V^{\otimes 2} \rightarrow V^{\otimes 2}$ is diagonalizable for generic q and has eigenvalues $\pm q^r$, for a finite number of integers r (see [BZ08] for details). If q is a p -th root of unity and p is an odd integer, then $q^a \neq -q^b$ for any integers a, b . Therefore the following definition makes sense for this choice of q , even though the R -matrix R_V might not be diagonalizable. Define

$$(23) \quad \begin{aligned} \Lambda_q^2 V &:= \sum_{i \in \mathbb{Z}} \{w \in V^{\otimes 2} \mid (R_V + q^i)^N w = 0 \text{ for } N \gg 0\}, \\ \Gamma_q^2(V) &:= \sum_{i \in \mathbb{Z}} \{w \in V^{\otimes 2} \mid (R_V - q^i)^N w = 0 \text{ for } N \gg 0\}. \end{aligned}$$

We can now define

$$\begin{aligned} S_q^d(V) &:= \frac{V^{\otimes d}}{\sum_{1 \leq i \leq d-1} V \otimes \cdots \otimes \Lambda^2(V)_{i,i+1} \otimes \cdots \otimes V}, \\ \Lambda_q^d(V) &:= \cap_{1 \leq i \leq d-1} V \otimes \cdots \otimes \Lambda^2(V)_{i,i+1} \otimes \cdots \otimes V, \\ \Gamma_q^d(V) &:= \cap_{1 \leq i \leq d-1} V \otimes \cdots \otimes \Gamma^2(V)_{i,i+1} \otimes \cdots \otimes V. \end{aligned}$$

Remark 3.8. When q is generic, our quantum divided power agrees with the definition of quantum symmetric power in [BZ08]. Note that semisimplicity makes it unnecessary for [BZ08] to distinguish the two. We prefer to define the quantum symmetric power as a quotient since it is more natural.

Proposition 3.9. *The quantum symmetric, exterior and divided powers are quantum polynomial functors.*

Proof. Let $f \in \text{Hom}_{\mathcal{B}_d}(V^{\otimes d}, W^{\otimes d})$. In order to show that Λ_q^d and Γ_q^d are quantum polynomial functors we need to show that the restriction of f to $\Lambda_q^d(V)$ has image in $\Lambda_q^d(W)$ and the restriction of f to $\Gamma_q^d(V)$ has image in $\Gamma_q^d(W)$. Since f commutes with the action of $T_i \in \mathcal{B}_d$, we have $f(V \otimes \cdots \otimes \Lambda_q^2(V)_{i,i+1} \otimes \cdots \otimes V) \subseteq W \otimes \cdots \otimes \Lambda_q^2(W)_{i,i+1} \otimes \cdots \otimes W$. It then easily follows from the definition that f restricts to a map from $\Lambda_q^d(V)$ to $\Lambda_q^d(W)$. A similar proof works for Γ_q^d .

From the fact that f maps $V \otimes \cdots \otimes \Lambda_q^2(V)_{i,i+1} \otimes \cdots \otimes V$ to $W \otimes \cdots \otimes \Lambda_q^2(W)_{i,i+1} \otimes \cdots \otimes W$ it follows that f induces a map (which we denote by the same letter) $f : S_q^d(V) \rightarrow S_q^d(W)$ from which we deduce that S_q^d is a quantum polynomial functor. \square

Remark 3.10. Symmetric/divided power and quantum symmetric/divided power are different quantum polynomial functors. Consider for example the divided power functor $\Gamma^{2, V_1^{\otimes e}}$, where V_1 is the defining $A_q(1, 1)$ -comodule. For any $V \in \Gamma_{q,e}^2$, the image $\Gamma^{2, V_1^{\otimes e}}(V)$ is the eigenspace of $R_V : V \otimes V \rightarrow V \otimes V$ with eigenvalue q^e . The quantum divided power Γ_q^2 maps V to the direct sum of eigenspaces of R_V with eigenvalues $+q^r$ for all integers r .

Remark 3.11. Note that when $q = 1$, the quantum polynomials $S_{V_1^{\otimes e}}^d$, S_q^d , $\Gamma^{d, V_1^{\otimes e}}$, and Γ_q^d all return the classical symmetric power, since in that case $q^e = 1 = +q^r$, $\forall r$.

With our definition, the following statement is clear.

Proposition 3.12. *Let V be an e -Hecke pair. Let N be the maximal rank of the Jordan blocks of R_V . The sequence*

$$0 \rightarrow \Gamma_q^2 V \xrightarrow{p_1} V^{\otimes 2} \xrightarrow{p_2} \Lambda^2 V \rightarrow 0$$

is exact, where p_1 is the inclusion $\Gamma_q^2 V \hookrightarrow V^{\otimes 2}$ (recall that we defined $\Gamma_q^2 V$ as a subspace of $V^{\otimes 2}$) and p_2 is defined by

$$p_2 : w \mapsto \prod_{-e \leq i \leq e} (R_V - q^i)^N w.$$

Here we record two short exact sequences of degree 2 quantum polynomial functors

$$(24) \quad 0 \rightarrow \Lambda_q^2 \rightarrow \otimes^2 \rightarrow S_q^2 \rightarrow 0,$$

which is by definition, and

$$(25) \quad 0 \rightarrow \Gamma_q^2 \rightarrow \otimes^2 \rightarrow \Lambda_q^2 \rightarrow 0,$$

which follows from Proposition 3.12. Also note that the first map in (24) and the last map in (25) make an exterior power a direct summand of a tensor power.

Furthermore, we have the following.

Proposition 3.13. *There are isomorphisms of quantum polynomial functors*

$$\begin{aligned} (\otimes^n)^\# &\cong \otimes^n, \\ (S_q^n)^\# &\cong \Gamma_q^n, \\ (\Lambda_q^n)^\# &\cong \Lambda_q^n. \end{aligned}$$

Proof. Let V be an e -Hecke pair, or equivalently, a polynomial representation of $U_q(\mathfrak{gl}_n)$ of degree e . There is a canonical isomorphism

$$\phi_n : (\tau V)^{\otimes n} \rightarrow \tau(V^{\otimes n})$$

of $U_q(\mathfrak{gl}_n)$ -modules, since τ_1 and the comultiplication commute. (See [Jan96, 9.20]. We note that what Jantzen denotes by the twisted comultiplication Δ' is the usual comultiplication structure that makes $A_q(n, n)$ live in the dual of $U_q(\mathfrak{gl}_n)$.) This is then also an isomorphism of $A_q(n, n)$ -comodules. By (22), the dual of ϕ_n induces the desired isomorphism $\otimes^n \cong (\otimes^n)^\#$ of polynomial functors.

To prove the other statements, first consider the $n = 2$ case. The R -matrix of τV satisfies

$$R_{\tau V} = R_V.$$

This follows from the fact that the R -matrix of V^* is the transpose of R_V (follows easily from Proposition 4.2.7 in [CP94]) and the matrix of V twisted by τ_1 is also the transpose of R_V (follows from the formulas in [Jan96, 9.20]). The isomorphism ϕ_2 preserves eigenspaces of $R_{\tau V} = R_V$, and hence restricts to an isomorphism from $(\Lambda_q^2(\tau V))$, which is the direct sum of eigenspaces of $R_{\tau V}$ corresponding to eigenvalues $-q^i$, to $\tau(\Lambda_q^2 V)$, which is the direct sum of eigenspaces of R_V corresponding to eigenvalues $-q^i$. Thus, ϕ_2 induces an isomorphism

$$\Lambda_q^2(\tau V) \cong \tau(\Lambda_q^2 V).$$

The duality τ satisfies $\tau(V \otimes W) \cong \tau V \otimes \tau W$ because both τ_1 and the antipode are antiautomorphisms. So we have

$$\tau(V \otimes \cdots \otimes \Lambda_q^2(V)_{i,i+1} \otimes \cdots \otimes V) \cong \tau V \otimes \cdots \otimes \Lambda_q^2(\tau V)_{i,i+1} \otimes \cdots \otimes \tau V.$$

By intersecting we obtain $\tau(\Lambda_q^d(V)) \cong \Lambda_q^d(\tau V)$, from which $\Lambda_q^n \cong (\Lambda_q^n)^\#$ follows (again, we use (22)).

The second statement is then obtained by comparing the short exact sequence (24) and the dual of the short exact sequence (25) for $d = 2$. Then it is clear from the definition of the quantum symmetric and divided powers that the same is true for all d . \square

Remark 3.14. We now explain why we assume p to be an odd root of unity. Assume $q = i = \sqrt{-1}$ and V_2 is the defining $A_q(2, 2)$ -comodule. We want to decompose $V_2^{\otimes 2}$ into two parts $\Gamma_q^2(V_2)$ and $\Lambda_q^2(V_2)$. The problem with the construction above is that the R -matrix R_2 only has one eigenvalue $q = -q^{-1} = i$. It is not diagonalizable; it has a 2×2 Jordan block and two 1×1 blocks. Thus, we cannot separate the eigenvalues of R_{V_2} into “positive” and “negative” eigenvalues and therefore definition (23) doesn’t make sense.

4. BRAIDING ON $\mathcal{P}_{q,e}$

We can use the braiding structure on the category of $A_q(n, n)$ -comodules to endow the category $(\mathcal{P}_{q,e}, \otimes)$ with the structure of a braided monoidal category.

Let $F \in \mathcal{P}_{q,e}^d$ and $G \in \mathcal{P}_{q,e}^{d'}$. The tensor products $F \otimes G$ and $G \otimes F$ both live in $\mathcal{P}_{q,e}^{d+d'}$. For an e -Hecke pair V , recall that $F(V)$ and $G(V)$ are both $A_q(n, n)$ -comodules for some n according to Proposition 5.1. Let $R_{F(V),G(V)} : F(V) \otimes G(V) \rightarrow G(V) \otimes F(V)$ be the braiding isomorphism defined in equation (9). We use it to define the natural transformation $R_{F,G} : F \otimes G \rightarrow G \otimes F$ by

$$R_{F,G}(V) := R_{F(V),G(V)}.$$

This map turns $\mathcal{P}_{q,e}$ into a braided monoidal category. But before we can prove that we need two results which are interesting in their own right.

Let V be an e -Hecke Pair and let $F \in \mathcal{P}_{q,e}^d$ and $G \in \mathcal{P}_{q,e}^{d'}$. Then $F(V)$ and $G(V)$ are naturally $A_q(V, V)$ -comodules. They also have the structure of $A_q(n, n)$ -comodules by Proposition 5.1.

The coalgebra $A_q(n, n)$ is coquasitriangular, therefore the comodule structure supplies us with the braiding $R_{F(V),G(V)} : F(V) \otimes G(V) \rightarrow G(V) \otimes F(V)$ that we mentioned above. However the coalgebra $A_q(V, V)$ is also coquasitriangular. Therefore there is a universal R -matrix $\mathcal{R}^V \in A_q(V \otimes V) \otimes A_q(V \otimes V)$ that satisfies properties (7) and is defined on the generators of $A_q(V, V)$ by

$$\mathcal{R}^V(x_{ij}^V \otimes x_{kl}^V) = (R_V)_{lj}^{ik}.$$

Since $F(V), G(V)$ are $A_q(V, V)$ -comodules, there is an R -matrix $R'_{F(V),G(V)} : F(V) \otimes G(V) \rightarrow G(V) \otimes F(V)$.

Proposition 4.1. *The maps $R_{F(V),G(V)}$ and $R'_{F(V),G(V)}$ are equal.*

Proof. The map $\psi_V : A_q(V, V) \rightarrow A_q(n, n)$ defined in equation (17) has the property that

$$\mathcal{R}^V(x, y) = \mathcal{R}(\psi_V(x), \psi_V(y))$$

for all $x, y \in A_q(V, V)$. The equation holds when x, y are generators of $A_q(V, V)$ because we defined $\mathcal{R}'(x_{ij}^V \otimes x_{kl}^V) = (R_V)_{lj}^{ik}$. The R -matrix of the $A_q(n, n)$ -comodule V is R_V , therefore the following equation holds by definition $\mathcal{R}(t_{ij}^V \otimes t_{kl}^V) = (R_V)_{lj}^{ik}$. We have that $\psi_V(x_{ij}^V) = t_{ij}^V$ and therefore the equation above holds when x, y are generators.

Since the equation holds on generators, it holds for every $x, y \in A_q(V, V)$. It implies that the R -matrices $R_{F(V),G(V)}$ and $R'_{F(V),G(V)}$ are equal. \square

Lemma 4.2. *The following diagram is commutative:*

$$\begin{array}{ccc}
F(V) \otimes G(V) & \xrightarrow{(1 \otimes \tau \otimes 1)(F''_{V,W} \otimes G''_{V,W})} & F(W) \otimes G(W) \otimes A_q(W, V) \otimes A_q(W, V) \\
R_{F(V), G(V)} \downarrow & & \downarrow R_{F(W), G(W)} \otimes 1 \\
G(V) \otimes F(V) & \xrightarrow{(1 \otimes \tau \otimes 1)(G''_{V,W} \otimes F''_{V,W})} & G(W) \otimes F(W) \otimes A_q(W, V) \otimes A_q(W, V)
\end{array}$$

Proof. Note that by Proposition 4.1, we can replace in the equation above $R_{F(V), G(V)}$ and $R_{F(W), G(W)}$ by $R'_{F(V), G(V)}$ and $R'_{F(W), G(W)}$, respectively.

By considering the commutative diagram in equation (19) in the proof of Lemma 3.5 for the functor $F \oplus G$ we obtain that the diagram

$$\begin{array}{ccc}
F \oplus G(V) \otimes F \oplus G(V) & \xrightarrow{(1 \otimes \tau \otimes 1)((F \oplus G)''_{V,W} \otimes (F \oplus G)''_{V,W})} & F \oplus G(W) \otimes F \oplus G(W) \otimes A_q(W, V) \\
R_{F \oplus G(V)} \downarrow & & \downarrow R_{F \oplus G(W)} \otimes 1 \\
F \oplus G(V) \otimes F \oplus G(V) & \xrightarrow{(1 \otimes \tau \otimes 1)((F \oplus G)''_{V,W} \otimes (F \oplus G)''_{V,W})} & F \oplus G(W) \otimes F \oplus G(W) \otimes A_q(W, V)
\end{array}$$

is commutative. The result follows from the fact that the tensor product $F \oplus G(V) \otimes F \oplus G(V)$ can be written as

$$F(V) \otimes F(V) \oplus F(V) \otimes G(V) \oplus G(V) \otimes F(V) \oplus G(V) \otimes G(V)$$

and that the restriction of $R_{F \oplus G(V)}$ to $F(V) \otimes G(V)$ is just $R_{F(V), G(V)}$, while the restriction of $(1 \otimes \tau \otimes 1)((F \oplus G)''_{V,W} \otimes (F \oplus G)''_{V,W})$ to $F(V) \otimes G(V)$ and $G(V) \otimes F(V)$ is just $(1 \otimes \tau \otimes 1)F''_{V,U} \otimes G''_{V,U}$ and $(1 \otimes \tau \otimes 1)(G''_{V,U} \otimes F''_{V,U})$, respectively. \square

Proposition 4.3. *Let $F, F' \in \mathcal{P}_{q,e}^d$ be quantum polynomial functors and let $\alpha : F \rightarrow F'$ be a natural transformation. Then the maps $\alpha_V : F(V) \rightarrow F'(V)$ are $A_q(V, V)$ -comodule homomorphisms.*

Proof. The following diagram commutes for any $f \in \text{Hom}(V, V) \cong S_q(V, V; d)$ because α is a natural transformation:

$$\begin{array}{ccc}
F(V) & \xrightarrow{\alpha_V} & F'(V) \\
F(f) \downarrow & & \downarrow F'(f) \\
F(V) & \xrightarrow{\alpha_V} & F'(V)
\end{array}$$

This is the same as saying that the map α_V is an $S_q(V, V; d)$ -module homomorphism from which the conclusion follows. \square

Now we can prove the main result of this subsection.

Theorem 4.4. *The category $\mathcal{P}_{q,e}$ is a braided monoidal category with braiding isomorphism $R_{F,G} : F \otimes G \rightarrow G \otimes F$.*

Proof. We first show that the braiding is a well defined morphism in $\text{Hom}(F \otimes G, G \otimes F)$. This is equivalent to showing the commutativity of the following diagram

$$\begin{array}{ccc} F(V) \otimes G(V) & \xrightarrow{(F \otimes G)(f)} & F(W) \otimes G(W) \\ R_{F(V),G(V)} \downarrow & & \downarrow R_{F(W),G(W)} \\ G(V) \otimes F(V) & \xrightarrow{(G \otimes F)(f)} & G(W) \otimes F(W) \end{array}$$

for all e -Hecke pairs V, W and any $f \in \text{Hom}(V, W)$. The commutativity of the diagram above for all f is equivalent to the commutativity of the following diagram

$$\begin{array}{ccc} S_q(V, W; d + d') \otimes F(V) \otimes G(V) & \longrightarrow & F(W) \otimes G(W) \\ 1 \otimes R_{F(V),G(V)} \downarrow & & \downarrow R_{F(W),G(W)} \\ S_q(V, W; d + d') \otimes G(V) \otimes F(V) & \longrightarrow & G(W) \otimes F(W) \end{array}$$

where the horizontal maps are given by $f \otimes v^F \otimes v^G \mapsto (F \otimes G)(f)(v^F \otimes v^G)$.

The commutativity of the second diagram is now equivalent to the commutativity of the following diagram by the definition of $S_q(V, W; d + d')$ as the dual of $A_q(W, V)_{d+d'}$.

$$\begin{array}{ccc} F(V) \otimes G(V) & \xrightarrow{(1 \otimes \tau \otimes 1)F''_{V,U} \otimes G''_{V,U}} & F(W) \otimes G(W) \otimes A_q(W, V) \otimes A_q(W, V) \\ R_{F(V),G(V)} \downarrow & & \downarrow R_{F(W),G(W)} \otimes 1 \\ G(V) \otimes F(V) & \xrightarrow{(1 \otimes \tau \otimes 1)G''_{V,U} \otimes F''_{V,U}} & G(W) \otimes F(W) \otimes A_q(W, V) \otimes A_q(W, V) \end{array}$$

which is commutative by Lemma 4.2. This completes the proof of naturality of the braiding.

Now we show that $R_{F,G}$ is a natural transformation, namely we need to show that for $f : F \rightarrow F'$ and $g : G \rightarrow G'$ we have

$$R_{F',G'}(f \otimes g) = (g \otimes f)R_{F,G}$$

The relation above holds when applied to any V because we can write $R_{F,G}(V) = R_{F(V),G(V)}$ as the composition $(1 \otimes 1 \otimes \mathcal{R}^V)(1 \otimes \tau \otimes 1)(\Delta_{G(V)} \otimes \Delta_{F(V)})\tau$ and $f_V \otimes g_V$ commutes with each factor of that composition except for τ which switches f_V and g_V by Proposition 4.3. Notice that in the equation above we use \mathcal{R}^V , the universal R -matrix of the coalgebra $A_q(V, V)$, instead of \mathcal{R} . We can do this because of Proposition 4.1.

The natural transformation $R_{F,G}$ is an isomorphism because $R_{F(V),G(V)}$ is an isomorphism for every V . We show that it satisfies equations (10). To prove the first property $\gamma_{V \otimes W, U} = (\gamma_{V,U} \otimes 1)(1 \otimes \gamma_{W,U})$ we note that it is equivalent to

$$R_{F \otimes G, H}(V) = (R_{F,H}(V) \otimes 1_G)(1_F \otimes R_{G,H}(V))$$

for every e -Hecke pair V . This can be rewritten as

$$R_{F(V) \otimes G(V), H(V)} = (R_{F(V), H(V)} \otimes 1_{G(V)})(1_{F(V)} \otimes R_{G(V), H(V)})$$

and follows immediately from Proposition 2.6.

The third property $\tilde{r}_V \gamma_{I,V} = \tilde{l}_V$ can be rewritten as

$$r_V R_{k,F(V)} = l_{F(V)}$$

which again follows immediately from Proposition 2.6.

The rest of the properties follow by the same argument as above. \square

5. COMPOSITION OF QUANTUM POLYNOMIAL FUNCTORS

Given two linear functors F, G between arbitrary k -linear categories, one can define the composition $F \circ G$ if the domain of F agrees with the codomain of G , and then $F \circ G$ is a linear functor. The quantum polynomial functors, as in the definition presented, have (up to equivalence) \mathcal{V} as their codomain, which does not match the domains of quantum polynomial functors. We have to endow the image of F with some additional structure.

Proposition 5.1. *Given V an e -Hecke pair and F a quantum polynomial functor of degree d , then $F(V)$ is also an $A_q(n, n)$ -comodule. It is a de -Hecke pair.*

Proof. By Lemma 3.3, $F(V)$ is an $A_q(V, V)$ -comodule with coaction $\Delta_{F(V)}^V$. Since ψ is a coalgebra homomorphism by Lemma 3.4, the composition $(1 \otimes \psi) \Delta_{F(V)}^V$ makes $F(V)$ into an $A_q(n, n)$ -comodule. It is easy to see from the way $(1 \otimes \psi) \Delta_{F(V)}^V$ is defined that $F(V)$ is a de -Hecke pair. This completes the proof of the statement. \square

We now explain the composition of quantum polynomials functors. Let $G \in \mathcal{P}_{q,e}^{d_2}$ and $F \in \mathcal{P}_{q,d_2e}^{d_1}$. We define $F \circ G \in \mathcal{P}_{q,e}^{d_1 d_2}$ as follows: on objects $V \in \Gamma_{q,e}^{d_1 d_2} \mathcal{V}$ we let

$$F \circ G(V) = F(G(V)).$$

This composition makes sense because $G(V)$ is an $A_q(n, n)_{d_2e}$ -comodule by Proposition 5.1. Since F is a quantum polynomial functor of degree d_1 , we have maps

$$F''_{G(V),G(W)} : F(G(V)) \rightarrow F(G(W)) \otimes A_q(G(W), G(V))_{d_1}$$

that satisfy the commutation relations in Proposition 3.2. G is also a quantum polynomial functor so we have maps:

$$G''_{V,W} : G(V) \rightarrow G(W) \otimes A_q(W, V)_{d_2}.$$

Define $(F \circ G)''_{V,W} : F(G(V)) \rightarrow F(G(W)) \otimes A_q(W, V)$ as

$$(F \circ G)''_{V,W} := (1 \otimes \psi_{V,W}^G) \circ F''_{G(V),G(W)}.$$

Theorem 5.2. *$(F \circ G)''$ satisfies properties (15), (16) in Proposition 3.2. Therefore $F \circ G$ is a well-defined quantum polynomial functor in $\mathcal{P}_{q,e}^{d_1 d_2}$.*

Proof. Diagram (15) for $(F \circ G)$ is equivalent to the exterior square of the following diagram:

$$\begin{array}{ccccc}
FG(V) & \xrightarrow{F''_{GV,GU}} & FG(U) \otimes A_q(G(U), G(V)) & \xrightarrow{1 \otimes \psi} & FG(U) \otimes A_q(U, V) \\
F''_{GV,GW} \downarrow & & \downarrow 1 \otimes \Delta_{GU,GW,GV} & & \downarrow 1 \otimes \Delta_{U,W,V} \\
FG(W) \otimes A_q(G(W), G(V)) & \xrightarrow{F''_{GW,GU} \otimes 1} & FG(U) \otimes A_q(G(U), G(W)) \otimes A_q(G(W), G(V)) & \xrightarrow{1 \otimes \psi \otimes \psi} & FG(U) \otimes A_q(U, W) \otimes A_q(W, V) \\
1 \otimes \psi \downarrow & & & & \downarrow \\
FG(W) \otimes A_q(W, V) & \xrightarrow{(F \circ G)''_{W,U} \otimes 1} & & & FG(U) \otimes A_q(U, W) \otimes A_q(W, V)
\end{array}$$

We show the commutativity of the exterior square by showing the commutativity of all three interior quadrilaterals. The commutativity of the top left diagram follows from the fact that F is

a quantum polynomial functor. The bottom left diagram follows from the fact that the horizontal arrows modify only the left component of the tensor product, while the vertical arrows modify the right component only. The commutativity of the top right diagram follows from the fact that $\psi_{U,V}^G$ satisfies equation (20).

Diagram (16) for $F \circ G$ is the exterior triangle of the following diagram:

$$\begin{array}{ccccc}
FG(V) & \xrightarrow{F''_{GV,GV}} & FG(V) \otimes A_q(GV, GV) & \xrightarrow{1 \otimes \psi} & FG(V) \otimes A_q(V, V) \\
\downarrow 1 & & \swarrow 1 \otimes e_{A_q(GV, GV)} & & \swarrow 1 \otimes e_{A_q(V, V)} \\
FG(V) \otimes k & & & &
\end{array}$$

The left triangle commutes by (16) and the right triangle by (21). □

6. REPRESENTABILITY FOR $\mathcal{P}_{q,e}^d$

Recall the divided power functor $\Gamma_q^{d,V} \in \mathcal{P}_{q,e}^d$:

$$\Gamma_q^{d,V}(W) = \text{Hom}_{\mathcal{B}_d}(V^{\otimes d}, W^{\otimes d})$$

defined for each e -Hecke pair $V \in \Gamma_{q,e}^d \mathcal{V}$. We show in this section that a certain divided power functor is a projective generator in $\mathcal{P}_{q,e}^d$.

Proposition 6.1. (*Yoneda's Lemma*) *For any $n \geq 0$, the functor $\Gamma_{q,e}^{d,V_n^{\otimes e}} \in \mathcal{P}_{q,e}^d$ represents the evaluation functor $\mathcal{P}_{q,e}^d \rightarrow \mathcal{V}$ given by $F \mapsto F(V_n^{\otimes e})$ and therefore $\Gamma_{q,e}^{d,V_n^{\otimes e}}$ is a projective object in $\mathcal{P}_{q,e}^d$.*

Proof. We need to show the existence of an isomorphism between

$$\text{Hom}_{\mathcal{P}_{q,e}^d}(\Gamma_{q,e}^{d,V_n^{\otimes e}}, F) \rightarrow F(V_n^{\otimes e})$$

for any $F \in \mathcal{P}_{q,e}^d$. Define the map $\rho : \text{Hom}_{\mathcal{P}_{q,e}^d}(\Gamma_{q,e}^{d,V_n^{\otimes e}}, F) \rightarrow F(V_n^{\otimes e})$ by

$$\rho(f) = f_{V_n^{\otimes e}}(\text{id}) \in F(V_n^{\otimes e}).$$

Let $\phi : F(V_n^{\otimes e}) \rightarrow \text{Hom}_{\mathcal{P}_{q,e}^d}(\Gamma_{q,e}^{d,V_n^{\otimes e}}, F)$ be defined as follows: for any $v \in F(V_n^{\otimes e})$, $\phi(v)$ is the natural transformation such that $\phi(v)(U) : \Gamma_{q,e}^{d,V_n^{\otimes e}}(U) \rightarrow F(U)$ takes

$$g \in \text{Hom}_{\mathcal{B}_d}((V_n^{\otimes e})^{\otimes d}, U^{\otimes d}) \mapsto F_{V_n^{\otimes e}, U}(g)(v) \in F(U).$$

We now show that ρ and ϕ are inverses to each other. Start with $v \in F(V_n^{\otimes e})$. We have

$$\rho(\phi(v)) = \phi(v)(V_n^{\otimes e})(\text{id}) = F_{V_n^{\otimes e}, V_n^{\otimes e}}(\text{id})(v) = v$$

Now let $f \in \text{Hom}_{\mathcal{P}_{q,e}^d}(\Gamma_{q,e}^{d,V_n^{\otimes e}}, F)$. We want to show that

$$\phi(\rho(f))_U(g) = f_U(g)$$

for any $U \in \Gamma_{q,e}^d$ and $g \in \text{Hom}_{\mathcal{B}_d}((V_n^{\otimes e})^{\otimes d}, U^{\otimes d})$.

$$\begin{aligned}\phi(\rho(f))(U)(g) &= F_{V^{\otimes e}, U}(g)(f_{V^{\otimes e}}(\text{id})) \\ &= f_U(\Gamma_{V^{\otimes e}, U}^{V^{\otimes e}}(g)(\text{id})) \\ &= f_U(g)\end{aligned}$$

The second equality follows from the commutativity of the following diagram

$$(26) \quad \begin{array}{ccc} \Gamma_{q,e}^{d, V_n^{\otimes e}}(V_n^{\otimes e}) & \xrightarrow{(\Gamma_{q,e}^{d, V_n^{\otimes e}})_{V_n^{\otimes e}, U}(g)} & \Gamma_{q,e}^{d, V_n^{\otimes e}}(U) \\ f_{V_n^{\otimes e}} \downarrow & & \downarrow f_U \\ F(V_n^{\otimes e}) & \xrightarrow{F_{V_n^{\otimes e}, U}(g)} & F(U) \end{array}$$

which holds because f is a natural transformation.

It follows that $\Gamma_{q,e}^{d, V_n^{\otimes e}}$ is a projective object in $\mathcal{P}_{q,e}^d$. \square

Definition 6.2. Let W be an e -Hecke pair. The quantum polynomial functor $F \in \mathcal{P}_{q,e}^d$ is W -generated if for every e -Hecke pair U the map

$$F'_{W,U} : S_q(W, U; d) \otimes F(W) \rightarrow F(U)$$

is surjective. We say that the category $\mathcal{P}_{q,e}^d$ is *finitely generated* if there is an e -Hecke pair W such that every $F \in \mathcal{P}_{q,e}^d$ is W -generated.

Remark 6.3. Note that $S_q(W, U; d) = \Gamma_q^{d,W}(U)$. So the map $F'_{W,U}$ above gives a surjection

$$\Gamma_q^{d,W} \otimes F(W) \rightarrow F$$

in $\mathcal{P}_{q,e}^d$, where $\Gamma_q^{d,W} \otimes F(W)$ is interpreted as the direct sum of $\dim F(W)$ copies of $\Gamma_q^{d,W}$ in $\mathcal{P}_{q,e}^d$. Since $\Gamma_q^{d,W}$ is projective by Proposition 6.1, Definition 6.2 says that $\Gamma_q^{d,W}$ is a projective generator of the category $\mathcal{P}_{q,e}^d$.

Definition 6.4. Given two objects $V, W \in \Gamma_{q,e}^d \mathcal{V}$, we say V generates W if the identity $\text{Id}_{W^{\otimes d}}$ (which should be written Id_W if we view it as an identity in the category $\Gamma_{q,e}^d \mathcal{V}$) can be written as a linear combination of \mathcal{B}_d -homomorphisms which factor through $V^{\otimes d}$ (which should be stated as ‘in $\Gamma_{q,e}^d \mathcal{V}$, the identity on W can be written as a linear combination of endomorphisms on W that factor through V ’).

Proposition 6.5. For $V, W \in \Gamma_{q,e}^d \mathcal{V}$, the following conditions are equivalent.

- (1) V generates W ;
- (2) Any indecomposable summand of $W^{\otimes d}$ (as a \mathcal{B}_d -module) is isomorphic to a direct summand of $V^{\otimes d}$.

In that case, the natural map

$$F(V) \otimes \Gamma_q^{d,V}(W) \rightarrow F(W)$$

is surjective for any $F \in \mathcal{P}_{q,e}^d$.

Proof. 1. \rightarrow 2.: Let M be an indecomposable summand of $W^{\otimes d}$. Since M is indecomposable, there are \mathcal{B}_d -maps $f : W^{\otimes d} \rightarrow V^{\otimes d}$ and $g : V^{\otimes d} \rightarrow W^{\otimes d}$ such that $g \circ f$ is the identity on M . The image of M under f is therefore a summand of $V^{\otimes d}$ that is isomorphic to M .

2. \rightarrow 1.: Denote by M_i all the indecomposable summands of $W^{\otimes d}$. For each M_i , there is a summand N_i in $V^{\otimes d}$ that is isomorphic to M_i as \mathcal{B}_d -modules. Let f_i be a \mathcal{B}_d -map that maps the summand $M_i \subset W^{\otimes d} \rightarrow N_i \subset V^{\otimes d}$ and g_i a \mathcal{B}_d -map that maps $N_i \subset V^{\otimes d} \rightarrow M_i \subset W^{\otimes d}$. Then the sum of $g_i \circ f_i$ is the identity on $W^{\otimes d}$.

Now we assume the equivalent condition and prove the last statement. Since there are maps f_i, g_i such that $\sum_i g_i \circ f_i = \text{Id}_{W^{\otimes d}}$, by applying F we obtain $\sum_i F(g_i) \circ F(f_i) = \text{Id}_{F(W)}$. This implies that the natural map $F(V) \otimes \text{Hom}_{\mathcal{P}_{q,e}^d}(V, W) \rightarrow F(W)$ is surjective. \square

We obtain two important corollaries.

Corollary 6.6. *The functor $\Gamma_q^{d,V}$ is a projective generator in $\mathcal{P}_{q,e}^d$ if V generates W for all $W \in \Gamma_{q,e}^d \mathcal{V}$.*

Proof. This follows from the last sentence in Proposition 6.5. \square

Corollary 6.7. *If V generates W then V generates any finite direct sum of copies of the comodules that are direct summands of W as $A_q(n, n)$ -comodules.*

Proof. This is obvious from the equivalence in Proposition 6.5. \square

The following lemma is a standard fact in quantum theory.

Lemma 6.8. *Given a composition $\lambda = (\lambda_1, \dots, \lambda_n)$ of d , let $V_{q,\lambda}$ be the subspace of $V_n^{\otimes d}$ with basis given by the vectors $v_{\sigma(i_1)} \otimes \dots \otimes v_{\sigma(i_d)}$ where $v_{i_1} \otimes \dots \otimes v_{i_d} = v_1^{\otimes \lambda_1} \otimes \dots \otimes v_n^{\otimes \lambda_n}$, and $\sigma \in S_d$, where $\{v_i\}$ is the standard basis of V_n . Then $V_{q,\lambda}$ is a direct summand of $V_n^{\otimes d}$ as a \mathcal{B}_d -module.*

Proof. The generator T_j of the braid group maps the vector $v = \dots \otimes v_{i_j} \otimes v_{i_{j+1}} \otimes \dots$ to a linear combination of v and $v' = \dots \otimes v_{i_{j+1}} \otimes v_{i_j} \otimes \dots$. See equation (5). It follows that T_i leaves $V_{q,\lambda}$ invariant. Since this is true for all λ , the submodule $V_{q,\lambda}$ is a direct summand of $V_n^{\otimes d}$ as \mathcal{B}_d -modules. \square

Remark 6.9. As a vector space, $V_{q,\lambda}$ is the same as the permutation module M_λ in $V_n^{\otimes d}$ viewed as an \mathcal{S}_d -module.

If $W \in \Gamma_{q,e}^d \mathcal{V}$ is of the form $V_n^{\otimes e}$, then we can formulate an explicit sufficient condition for what generates W .

Proposition 6.10. *The e -Hecke pair $V_m^{\otimes e} \in \Gamma_{q,e}^d \mathcal{V}$ generates $V_n^{\otimes e}$ for any $n \in \mathbb{N}$ if $m \geq de$.*

Proof. First let $e = 1$. By Proposition 6.5, it is enough to prove that any \mathcal{B}_d -indecomposable summand of $V_n^{\otimes d}$ is a \mathcal{B}_d -summand of $V_m^{\otimes d}$. Let M be an indecomposable summand of $V_n^{\otimes d}$ as \mathcal{B}_d -modules. It follows from Lemma 6.8 that M is an indecomposable summand of $V_{q,\lambda}$ for some λ . If $m \geq d$, then $V_{q,\lambda}$ is isomorphic to a summand of $V_m^{\otimes d}$. The conclusion follows.

For general e , consider the \mathcal{B}_{de} -modules $V_n^{\otimes de}$ and $V_m^{\otimes de}$. The proof of the case $e = 1$ (where d is replaced by de), combined with Proposition 6.5, implies that there are \mathcal{B}_{de} -maps

$$f_i : V_n^{\otimes de} \rightarrow V_m^{\otimes de}, \quad g_i : V_m^{\otimes de} \rightarrow V_n^{\otimes de}$$

such that $\sum_i g_i \circ f_i = \text{Id}_{V_n^{\otimes de}}$, provided that $m \geq ed$. Consider the subgroup, call it $B_{d,e}$, of \mathcal{B}_{de} generated by $T_{w_1}, \dots, T_{w_{d-1}}$, where w_i are as in (11). The group $B_{d,e}$ is isomorphic to B_d by the assignment $T_{w_i} \mapsto T_i$, and the B_{de} action restricts to $B_{d,e}$ is, under this isomorphism, exactly the usual B_d action on $(V_n^{\otimes e})^{\otimes d}$ coming from the R-matrix of $V_n^{\otimes e}$. The \mathcal{B}_{de} -maps f_i and g_i can then be viewed as \mathcal{B}_d -maps

$$f_i : (V_n^{\otimes e})^{\otimes d} \rightarrow (V_m^{\otimes e})^{\otimes d}, \quad g_i : (V_m^{\otimes e})^{\otimes d} \rightarrow (V_n^{\otimes e})^{\otimes d}$$

with $\sum_i g_i \circ f_i = \text{Id}$. This gives the desired result. \square

The special case in the Proposition 6.10 is enough to guarantee a projective generator in the semisimple situation:

Theorem 6.11. *Suppose q is not a root of unity and $\text{char}(k) = 0$. Then the functor $\Gamma_{q,e}^{d,V_m^{\otimes e}}$ is a projective generator in $\mathcal{P}_{q,e}^d$ if $m \geq ed$.*

Proof. By Corollary 6.6, it is enough to show that any $W \in \Gamma_{q,e}^d \mathcal{V}$ is generated by $V_m^{\otimes e}$. But by Propositions 2.11 and 2.12, W is a direct sum of direct summands of some $V_n^{\otimes e}$. So Proposition 6.10 and Corollary 6.7 shows that $V_m^{\otimes e}$ generates W .

The fact that $V_m^{\otimes e}$ is projective is just Proposition 6.1. \square

Remark 6.12. For $e = 1$ we obtain Theorem 4.7 in Hong and Yacobi [HY17] when q is generic and $\text{char}(k) = 0$. But note that our proof for $e = 1$ works when q is a root of unity or $\text{char}(k) \neq 0$ with a minor addition which we now explain. Theorem 6.11 depends on Corollary 6.6, Proposition 6.10 and Corollary 6.7 which are true regardless if q is a root of unity or not and if $\text{char}(k)$ is 0 or not. It also depends on Propositions 2.11 and 2.12, which are not true in general for q a root of unity or $\text{char}(k) = 0$. However, when $e = 1$, Propositions 2.11 and 2.12 hold for q a root of unity and $\text{char}(k) = 0$ because 1-Hecke pairs are just direct sums of the defining comodule V_n for any n . Thus we obtain Theorem 4.7 in [HY17] with no restrictions on q or the characteristic of k .

Corollary 6.13. *The evaluation functor $\mathcal{P}_{q,e}^d \rightarrow \text{mod}(S_q(V_n^{\otimes e}, V_n^{\otimes e}, d))$ is an equivalence of categories for q not a root of unity and $\text{char}(k) = 0$. It follows that $S_q(V_n^{\otimes e}, V_n^{\otimes e}, d)$ and $S_q(V_m^{\otimes e}, V_m^{\otimes e}, d)$ are Morita equivalent when $m, n \geq de$.*

Remark 6.14. We can think of the generalized Schur algebras $S_q(V_n^{\otimes e}, V_n^{\otimes e}; d)$ for generic q as follows. Quantum Schur-Weyl duality (due to Jimbo) says that there is a commuting action of the Hecke algebra \mathcal{H}_d and the q -Schur algebra $S_q(V_n, V_n; d)$ on the space $V_n^{\otimes d}$ and these two actions satisfy a double centralizer property. The e -Hecke algebra $\mathcal{H}_{d,e} \subset \mathcal{H}_{de}$ acts on $(V_n^{\otimes e})^{\otimes d}$ as explained before. Then $S_q(V_n^{\otimes e}, V_n^{\otimes e}; d) \supset S_q(V_n, V_n; de)$ is the largest object that makes the following two actions commute:

$$S_q(V_n^{\otimes e}, V_n^{\otimes e}; d) \curvearrowright (V_n^{\otimes e})^{\otimes d} \curvearrowleft \mathcal{H}_{d,e}$$

and one can ask if it's the object that preserves a double centralizer property.

7. THE CATEGORY $\widetilde{\mathcal{P}}_q^d$

We now define a category that ‘‘lives inbetween’’ the category $\mathcal{P}_{q,e}^d$ for any e and the category \mathcal{P}_q^d . Let d be a positive integer. The quantum divided power category $\Gamma_q^d \mathcal{V}$ is the category with objects formal finite direct sums $\bigoplus_i V_i$ where each V_i is an e -Hecke for some e . The morphisms are defined on homogeneous objects as follows:

$$\text{Hom}_{\Gamma_q^d}(V_i, W_j) := \text{Hom}_{\mathcal{B}_d}(V_i^{\otimes d}, W_j^{\otimes d}).$$

The Hom extends naturally to all objects via the formula

$$\text{Hom}_{\Gamma_q^d}\left(\bigoplus_i V_i, \bigoplus_j W_j\right) = \bigoplus_{i,j} \text{Hom}_{\Gamma_q^d}(V_i, W_j).$$

Definition 7.1. The category $\widetilde{\mathcal{P}}_q^d$ is the category of linear functors

$$F : \Gamma_q^d \mathcal{V} \rightarrow \Gamma_q^1 \mathcal{V}.$$

Morphisms are natural transformations of functors. Note that $\Gamma_q^1 \mathcal{V}$ is equivalent to \mathcal{V} .

Remark 7.2. Given a linear functor $F : \Gamma_q^d \mathcal{V} \rightarrow \Gamma_q^1 \mathcal{V}$, denote by F_e its restriction to $\Gamma_{q,e}^d \mathcal{V}$. Then it is not hard to see that $F_e \in \mathcal{P}_{q,e}^d$.

We can define a composition on $\widetilde{\mathcal{P}}_q^d$ similarly to how we defined composition between a functor in $\mathcal{P}_{q,d_2e}^{d_1}$ and a functor in $\mathcal{P}_{q,e}^{d_2}$.

It is interesting to note that when $q = 1$, the category $\widetilde{\mathcal{P}}_q^d$ becomes the category of classical polynomial functors \mathcal{P}^d due to Friedlander and Suslin. Therefore when $q = 1$, the category $\widetilde{\mathcal{P}}_q^d$ and the categories $\mathcal{P}_{q,e}^d$ for any e are all equivalent. We show that for generic q , $\widetilde{\mathcal{P}}_q^d$ is not equivalent to $\mathcal{P}_{q,e}^d$ for any e . In fact in the quantum case $\widetilde{\mathcal{P}}_q^d$ is “closer” to \mathcal{P}_q^d than to $\mathcal{P}_{q,e}^d$ since both $\widetilde{\mathcal{P}}_q^d$ and \mathcal{P}_q^d are *not* finitely generated while $\mathcal{P}_{q,e}^d$ is, when q is generic. This is another reason why the theory of quantum polynomial functors is richer in the quantum case than it is in the classical case.

Proposition 7.3. *Suppose q is not a root of unity. Then the category $\widetilde{\mathcal{P}}_q^d$ is not finitely generated. In particular, $\widetilde{\mathcal{P}}_q^d$ is not equivalent to $\mathcal{P}_{q,e}^d$ for any e .*

Proof. Consider the sequence of objects $k^{\otimes f}$ in $\Gamma_q^d \mathcal{V}$, where $f \in \mathbb{N}$ and $k = (k, q)$ is the trivial $A_q(1, 1)$ -comodule of degree 1. Then each $(k^{\otimes f})^{\otimes d}$ is a one-dimensional \mathcal{B}_d -module on which each T_i acts as multiplication by q^{f^2} . These form an infinite collection of irreducibles \mathcal{B}_d -modules since q is not a root of unity. Now take $F = \bigotimes^d$ in and we show that no V makes the evaluation map

$$V^{\otimes d} \otimes \Gamma_q^{d,V}(k^{\otimes f}) = V^{\otimes d} \otimes \text{Hom}_{\mathcal{B}_d}(V^{\otimes d}, (k^{\otimes f})^{\otimes d}) \rightarrow (k^{\otimes f})^{\otimes d}$$

surjective for all f . If there is such V , on the contrary, then the head (as \mathcal{B}_d -module) of $V^{\otimes d}$ contains all the $k^{\otimes f}$ above. This is impossible since V is finite dimensional. \square

Remark 7.4. Note that the proof makes use of the normalization of the R -matrix R_n defined in equation (5). In particular, we use that $R_n(v_1 \otimes v_1) = qv_1 \otimes v_1$. For a different normalization (for example where $R_n(v_1 \otimes v_1) = v_1 \otimes v_1$) the proof above doesn’t work. The result stays true, while the argument becomes computationally more complicated. One needs to look at R_2 (which will have an eigenvalue $-q^{-2}$ for the normalization mentioned above) and modify the proof of Proposition 7.3 accordingly.

8. REMARKS ON $\mathcal{P}_{q,e}^d$ WHEN q IS A ROOT OF UNITY

Let q be an l -th root of unity where $l > 1$ is an odd integer.

8.1. Representability. The proof of Theorem 6.11 is not valid in this case, since an indecomposable e -Hecke pair $W \in \text{comod}(A_q(n, n))$ is not necessarily a summand of $V_n^{\otimes e}$. But it can still be true that $\mathcal{P}_{q,e}^d$ is finitely generated, hence equivalent to the module category of a finite dimensional algebra (possibly the (q, e) -Schur algebra as in the q -generic case). For example in the classical case ($q = 1$) when the field k has characteristic p , the category of polynomial functors is not semisimple, but it does have a projective generator just as in the case when the characteristic of the field is 0. In this section we present some remarks on the case when q is a root of unity not equal to 1.

First recall that it is only at the last step of the proof that we use the semisimplicity. In particular, Corollaries 4.4, 4.5 and Proposition 4.8 are valid when q is non-generic. We summarize them as a separate statement.

Proposition 8.1. *Let V be an e -Hecke pair. The category $\mathcal{P}_{q,e}^d$ has a (finite) projective generator $\Gamma_q^{d,V}$ if for any e -Hecke pair W , every indecomposable \mathcal{B}_d -summand of $W^{\otimes d}$ is isomorphic, as a \mathcal{B}_d -module, to a direct summand of $V^{\otimes d}$.*

Note that the set of divided powers generates $\mathcal{P}_{q,e}^d$, thus if $\mathcal{P}_{q,e}^d$ is finitely generated then one can find a functor of the form $\Gamma_q^{d,V}$ which is a projective generator.

The condition in Proposition 8.1 is reduced to an elementary statement about Jordan block decomposition of R -matrices if $d = 2$.

Proposition 8.2. *Let $n \in \mathbb{N}$, and let U be an indecomposable $A_q(m, m)$ -comodule of degree e . Denote the Jordan blocks of $R_{V_n^{\otimes e}}$ by $B(n_i, a_i)$, where i runs through some finite index set I , $n_i \in \mathbb{N}$ is the rank of the block and $a_i \in k$ is the generalized eigenvalue of the block. Similarly, name the Jordan blocks of R_U by $B(m_j, b_j)$, where $j \in J$.*

Then the identity map on $U \otimes U$ factors through $V_n^{\otimes e} \otimes V_n^{\otimes e}$ as a \mathcal{B}_2 -map if and only if for each $j \in J$, there exists $i \in I$ such that $a_i = b_j$ and $n_i = m_j$.

Proof. It is enough to factor the identity on each Jordan block of U . But we can do it by embedding $B(m_j, b_j)$ onto the block $B(n_i, a_i)$. \square

Remark 8.3. The condition in Proposition 8.2 is trivially true if the R -matrix of any e -Hecke pair is diagonalizable. This is the case when q is not a root of unity, but it is not when q is a root of unity. For example R_q is not diagonalizable when $q = \pm i$. If we restrict us to 1-Hecke pairs, these are the only values for q where R_V is not diagonalizable. For general e -Hecke pair V , we expect there are other roots of unity where R_V is not diagonalizable.

Remark 8.4. It is possible that the functor cohomology for quantum polynomial functors agrees with the corresponding quantum group cohomology even if the category $\mathcal{P}_{q,e}^d$ does not have a finite generator. A similar approach is found in Suslin's appendix in [FFSS99].

8.2. An additional structure. An important feature when q is a root of unity is the existence of the Frobenius twist. When $e = 1$, this structure comes from the module categories of the q -Schur algebra or quantum group. Let us write this in terms of polynomial functors. This is explained in a previous version of [HY17], which we now repeat. The algebra map

$$\text{Fr}_{n,m} : A_1(n, m) \rightarrow A_q(n, m)$$

defined on the standard generators by

$$x_{ij} \mapsto x_{ij}^l$$

is also a coalgebra map. Using this map, we can define a functor

$$(-)^{[1]} : \mathcal{P}^d \rightarrow \mathcal{P}_{q,1}^{ld}.$$

We call it the Frobenius twist. (We remind the reader that the classical polynomial functor category \mathcal{P}^d can be viewed as $\mathcal{P}_{1,e}^d$ for any e , and that $\mathcal{P}_{q,1}^{ld}$ is equivalent to the category \mathcal{P}_q^{ld} of Hong-Yacobi [HY17]. See Remark 6.12.) Given $F \in \mathcal{P}^d$, its Frobenius twist $F^{[1]}$ is defined to be a functor from $\Gamma_{q,1}^d \mathcal{V}$ to $\Gamma_{q,d}^1 \mathcal{V} \cong \mathcal{V}$ that sends a 1-Hecke pair V to the vector space $F(V)$ (forgetting the 1-Hecke structure, V is viewed as an object in $\Gamma^d \mathcal{V}$). To define what $F^{[1]}$ does to morphisms, it is enough to specify the map

$$(F^{[1]})''_{V,W} : F(V) \rightarrow F(W) \otimes A_q(W, V).$$

Note that V and W are 1-Hecke pairs, hence direct sum of standard $A_q(n, n)$ -comodules and $A_q(m, m)$ -comodules, respectively, for certain n, m . We define the map $\text{Fr} : A_1(W, V) \rightarrow A_q(W, V)$ as a straightforward generalization of $\text{Fr}_{n,m}$. It maps

$$x_{ij} \in \bigoplus A_1(V_n, V_m) \cong A_1(V, W) \mapsto x_{ij}^l \in \bigoplus A_q(V_n, V_m) \cong A_q(V, W).$$

This extends to a bialgebra map. Now we define the map $(F^{[1]})''_{V,W}$; it is the composition

$$(1_{F(W)} \otimes \text{Fr}) \circ F''_{V,W} : F(V) \rightarrow F(W) \otimes A_1(W, V) \rightarrow F(W) \otimes A_q(W, V).$$

It is not hard to see that $(F^{[1]})''_{V,W}$ satisfies the properties needed to make $F^{[1]}$ into a quantum polynomial functor.

Consider $I^{[1]} \in \mathcal{P}_{q,1}^l$ where $I \in \mathcal{P}^1$ is the identity polynomial $\mathcal{V} = \Gamma^1 \mathcal{V} \rightarrow \mathcal{V}$. Then for any d , the composition

$$- \circ - : \mathcal{P}_{q,l}^d \otimes \mathcal{P}_{q,1}^l \rightarrow \mathcal{P}_{q,1}^{dl}$$

induces the functor

$$- \circ I^{[1]} : \mathcal{P}_{q,l}^d \rightarrow \mathcal{P}_{q,1}^{dl}.$$

Since the Frobenius $I^{[1]}$ lives only in the category $\mathcal{P}_{q,1}^l$, we cannot compose the Frobenius with itself multiple times (as in the classical case).

One may ask if we can define a class of objects $I_e^{[1]} \in \mathcal{P}_{q,e}^l$ for all e which are reasonable analogues of the Frobenius. This will supply higher Frobenius twists using composition:

$$I_e^{[r]} := I_{l^{r-1}e}^{[1]} \circ \cdots \circ I_e^{[1]}.$$

Then precomposing or postcomposing $I_e^{[r]}$ provides functors between various polynomial functor categories.

However, $I_e^{[1]}$ does not come easily from the structure of the q -Schur algebras as in the $e = 1$ case; if $e > 1$ we do not have an analogous coalgebra map

$$A_1, (V, W) \rightarrow A_q(V, W)$$

where V, W are e -Hecke pairs. Thus we must take a different approach to define $I_e^{[1]}$. One may try to define the Frobenius as the cohomology of certain complexes that are of interest by themselves.

Classically (by this we mean $q = 1$ and $\text{char}(k) = p$), we have the following exact sequence of polynomial functors:

$$(27) \quad 0 \rightarrow I^{[1]} \rightarrow S^p \rightarrow \Gamma^p \rightarrow I^{[1]} \rightarrow 0.$$

That is, one can define the Frobenius polynomial $I^{[1]} \in \mathcal{P}_{1,e}^p$ as either the kernel or the cokernel of the middle map in (27). Alternatively, the following complex, called the d -th de Rham complex, has nontrivial cohomology when p divides d .

$$(28) \quad 0 \rightarrow S^d \rightarrow S^{d-1} \otimes \Lambda^1 \rightarrow S^{d-2} \otimes \Lambda^2 \rightarrow \cdots \rightarrow \Lambda^d \rightarrow 0$$

If $d = ap$, its cohomology is given by the Frobenius twist of the $(a-1)p$ -th de Rham complex.

One can try to quantize these complexes. That gives a way to define $I_e^{[1]}$. Note that all the objects in (27) and (28) except the Frobenius are defined in $\mathcal{P}_{q,e}^d$ for all e . Also note that the quantum symmetric powers and quantum divided powers usually behave very differently when we move away from the degree 1 case to the degree e case. For example, the dimension of $S_q^d(V)$ depends on the $A_q(n, n)$ -comodule structure of V and not only on the dimension of V (this phenomena is investigated in [BZ08]). Therefore the dimension of $I_e^{[1]}(V)$ for an e -Hecke pair V is different from the dimension of V , in contrary to the classical case. In particular, we cannot obtain $I_e^{[1]}(V)$ from the underlying space of V by just twisting the module structure.

This discussion can be the starting point of further investigations. One can try to define the quantum Frobenius twist as mentioned above, and try to understand its properties. Then, one can try to understand its role in the representation theory of quantum groups, or its uses in cohomology theory.

REFERENCES

- [BZ08] Arkady Berenstein and Sebastian Zwicknagl. Braided symmetric and exterior algebras. *Trans. Amer. Math. Soc.*, 360(7):3429–3472, 2008.
- [Cha15] Marcin Chalupnik. Derived Kan extension for strict polynomial functors. *Int. Math. Res. Not. IMRN*, (20):10017–10040, 2015.

- [CP94] Vyjayanthi Chari and Andrew Pressley. *A guide to quantum groups*. Cambridge University Press, Cambridge, 1994.
- [DJ89] Richard Dipper and Gordon James. The q -Schur algebra. *Proc. London Math. Soc. (3)*, 59(1):23–50, 1989.
- [FFSS99] Vincent Franjou, Eric M. Friedlander, Alexander Scorichenko, and Andrei Suslin. General linear and functor cohomology over finite fields. *Ann. of Math. (2)*, 150(2):663–728, 1999.
- [FRT88] L. D. Faddeev, N. Yu. Reshetikhin, and L. A. Takhtajan. Quantization of Lie groups and Lie algebras. In *Algebraic analysis, Vol. I*, pages 129–139. Academic Press, Boston, MA, 1988.
- [FS97] Eric M. Friedlander and Andrei Suslin. Cohomology of finite group schemes over a field. *Invent. Math.*, 127(2):209–270, 1997.
- [HTY14] Jiuzu Hong, Antoine Touzé, and Oded Yacobi. Polynomial functors and categorifications of Fock space. In *Symmetry: representation theory and its applications*, volume 257 of *Progr. Math.*, pages 327–352. Birkhäuser/Springer, New York, 2014.
- [HY13] Jiuzu Hong and Oded Yacobi. Polynomial functors and categorifications of Fock space II. *Adv. Math.*, 237:360–403, 2013.
- [HY17] Jiuzu Hong and Oded Yacobi. Quantum polynomial functors. *J. Algebra*, 479:326–367, 2017.
- [Jan96] Jens Carsten Jantzen. *Lectures on quantum groups*, volume 6 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1996.
- [LR97] Larry A. Lambe and David E. Radford. *Introduction to the quantum Yang-Baxter equation and quantum groups: an algebraic approach*, volume 423 of *Mathematics and its Applications*. Kluwer Academic Publishers, Dordrecht, 1997.
- [PW91] Brian Parshall and Jian Pan Wang. Quantum linear groups. *Mem. Amer. Math. Soc.*, 89(439):vi+157, 1991.
- [Tak02] Mitsuhiro Takeuchi. A short course on quantum matrices. In *New directions in Hopf algebras*, volume 43 of *Math. Sci. Res. Inst. Publ.*, pages 383–435. Cambridge Univ. Press, Cambridge, 2002. Notes taken by Bernd Strüber.
- [Tou10] Antoine Touzé. Cohomology of classical algebraic groups from the functorial viewpoint. *Adv. Math.*, 225(1):33–68, 2010.
- [Tou13] A. Touzé. A construction of the universal classes for algebraic groups with the twisting spectral sequence. *Transform. Groups*, 18(2):539–556, 2013.

MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY
E-mail address: valentin.buciumas@gmail.com

MATHEMATISCHES INSTITUT, UNIVERSITÄT BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY
E-mail address: hankyung@math.uni-bonn.de