


Max-Planck-Institut für Mathematik Bonn

KP hierarchy for Hurwitz-type cohomological field theories

by

Reinier Kramer



Max-Planck-Institut für Mathematik
Preprint Series 2021 (42)

Date of submission: July 13, 2021

KP hierarchy for Hurwitz-type cohomological field theories

by

Reinier Kramer

Max-Planck-Institut für Mathematik
Vivatsgasse 7
53111 Bonn
Germany

KP HIERARCHY FOR HURWITZ-TYPE COHOMOLOGICAL FIELD THEORIES

REINIER KRAMER

ABSTRACT. We generalise a result of Kazarian regarding Kadomtsev-Petviashvili integrability for single Hodge integrals to general cohomological field theories related to Hurwitz-type counting problems or hypergeometric tau-functions. The proof uses recent results on the relations between hypergeometric tau-functions and topological recursion, as well as the Eynard-DOSS correspondence between topological recursion and cohomological field theories. In particular, we recover the result of Alexandrov of KP integrability for triple Hodge integrals with a Calabi-Yau condition.

CONTENTS

1. Introduction	1
Results of this paper	2
Open questions	2
Outline of the paper	3
Notation	3
On the origin of this paper	3
Acknowledgments	3
2. Prerequisites on the KP hierarchy and topological recursion	3
2.1. The KP hierarchy	3
2.2. Space of tau-functions and Lie action	4
2.3. Single Hodge integrals	5
2.4. Hypergeometric KP tau-functions and topological recursion	5
2.5. Topological recursion and cohomological field theories	7
2.6. The Mariño-Vafa formula and KP for topological vertex amplitudes	8
3. KP hierarchy for intersection numbers	10
3.1. The change of variables	10
3.2. KP for intersection numbers	12
3.3. Finiteness of the transformation	13
3.4. The case of triple Hodge integrals	14
References	16

I. INTRODUCTION

The moduli spaces of curves are a central object in modern algebraic geometry, and have been studied intensively. In particular, their intersection theory is a subject of ongoing research. The space $\overline{\mathcal{M}}_{g,n}$ has n line bundles \mathbb{L}_i whose fibres at a point are the cotangent lines at the i th point of the represented curve, and a rank- g Hodge bundle \mathbb{E} whose fibres are the space of one-forms at the curve. Their Chern classes are defined to be $\psi_i := c_1(\mathbb{L}_i)$ and $\lambda_j := c_j(\mathbb{E})$, respectively. Moreover, the spaces $\overline{\mathcal{M}}_{g,n}$ for different g and n have many structure maps between them, and many classes behave well under these maps. A collection of classes on all $\overline{\mathcal{M}}_{g,n}$ satisfying certain coherence axioms with respect to the structure maps are called cohomological field theories (CohFTs), and these play an important role in enumerative geometry of curves. One well-known example is the total Hodge class $\Lambda(t) = \sum \lambda_i t^i$.

By the Witten-Kontsevich theorem [Wit91; Kon92], moduli spaces of curves have many relations to areas of mathematical physics and integrable hierarchies. In particular, this theorem proves that a generating function of the intersection numbers of ψ -classes is a tau-function of the Korteweg-de Vries hierarchy.

Furthermore, the Ekedahl-Lando-Shapiro-Vainshtein formula [ELSV01] relates single Hodge integrals, i.e. intersection numbers of $\Lambda(-1)$ with ψ -classes, to simple single Hurwitz numbers, counting ramified coverings of $\mathbb{P}_{\mathbb{C}}^1$ with only simple ramifications (with profile $(2, 1, 1, 1, \dots)$) except for one point. Hurwitz numbers themselves also give a large class of tau-functions of Toda or Kadomtsev-Petviashvili hierarchies (of which the KdV hierarchy is a reduction), as noted by Okounkov [Ok00].

Kazarian [Kaz09] interpreted the ELSV formula as a change of variables from the generating function of single Hodge integrals to a tau-function of the Kadomtsev-Petviashvili hierarchy, using the result of Okounkov on simple single Hurwitz numbers.

All of these results have strong relations to Chekhov-Eynard-Orantin topological recursion [CEO06; EO07], a succesful way of encoding many counting problem with a natural genus expansion into a spectral curve with a recursively defined collection of multidifferentials, which should be generating functions of the counts. The Witten-Kontsevich ψ -intersection numbers can be encoded this way, and this is somehow the base case of the theory. Many types of Hurwitz numbers obey topological recursion as well, starting with [BM08; BEMS11] for the first case of simple Hurwitz numbers, and culminating in the works of Bychkov-Dunin-Barkwoski-Kazarian-Shadrin [BDKS20a; BDKS20b], which prove topological recursion for two large families of hypergeometric KP tau-functions, encompassing nearly all previously-studied cases of Hurwitz numbers.

In another direction, there is a general correspondence between topological recursion and intersection numbers of CohFTs [Eyn14; DOSS14], which vastly generalises the ELSV formula when combined with the results on topological recursion for Hurwitz numbers.

Another related direction is the conjecture of Mariño and Vafa [MV02] on a further generalisation of the ELSV formula, proved independently in [LLZ03; OP04]. This Mariño-Vafa formula relates triple Hodge integrals with a Calabi-Yau condition to topological vertex amplitudes, i.e. Gromov-Witten invariants of \mathbb{C}^3 . Topological recursion was conjectured for toric Calabi-Yau threefolds by Bouchard-Klemm-Mariño-Pasquetti [BKMP09]. It was first proved in [Che18; Zho09] for \mathbb{C}^3 , as well as in [Eyn11] as an example of the general correspondence of theorem 2.11, while the general BKMP conjecture was proved in [EO15]. Although the Mariño-Vafa formula fits in the framework of hypergeometric tau-functions, this case is not subsumed by the proof scheme of [BDKS20b].

Both the space of CohFTs and the space of KP tau-functions have an action of an infinite-dimensional group, respectively the Givental group and the Heisenberg-Virasoro group. As certain elements of these spaces have been identified by Witten-Kontsevich and Kazarian, and different integrable hierarchies have been constructed for general CohFTs by Dubrovin-Zhang [DZ01] and Buryak [Bur15a], one may ask how general the relation is with KP specifically, and the group actions are a natural tool to study this question.

Alexandrov [Ale20] showed that in the case of a rank-one CohFT, the orbits of the Witten-Kontsevich CohFT/tau-function under these two different group actions have an intersection which is only two-dimensional, and contains exactly the triple Hodge integrals that appear in the Mariño-Vafa formula. As a consequence, Alexandrov generalises Kazarian’s result to show that the generating function of Calabi-Yau triple Hodge integrals satisfies the KP hierarchy after a linear change of variables.

Results of this paper. We give a new viewpoint on the relation found by Alexandrov, by generalising Kazarian’s proof in [Kaz09] to all hypergeometric KP tau-functions satisfying topological recursion, using the above results. This yields a change of variables coming from the function X for any hypergeometric tau-function preserving the KP hierarchy after removing the unstable terms of the tau-function. When topological recursion holds, this resulting tau-function can be interpreted as the generating function of the cohomological field theory.

In general, the change of variables contains infinite linear combinations. However, we identify when the linear combinations are actually finite, and find a finite-dimensional family for each CohFT rank / number of ramification points. In the rank one case, this recovers exactly the triple Hodge integrals, in a particular parametrisation. For higher rank, this family seems to fit within Alexandrov’s deformed generalised Kontsevich model [Ale21].

Interestingly, the function X may also be a Möbius transformation. In this case, there is no correction term, and this can be interpreted as certain independence of the parametrisation of the spectral curve. This also resolves the meaning behind Kazarian’s change of coordinates, as voiced in [Kaz09, Remark 2.6]: “The definition for the change (6) looks unmotivated. [...] The only motivation that we can provide here is that ‘it works’.” There is quite a freedom of choice, but the particular choice Kazarian made reduces to the finite-dimensional family indicated above.

Open questions. Single and triple Hodge integrals have been studied intensively in relation to Dubrovin-Zhang hierarchies, yielding relations to the intermediate long wave hierarchy and the fractional Volterra hierarchy, cf. [Bur15b; Bur16; LYZZ21]. The relation between those results and the current work are still unclear, and will be discussed elsewhere.

The family where the linear change of variables is finite seems like an interesting and natural deformation of Witten’s r -spin class, keeping a single ramification point, but splitting the pole of dx . However, this family

seems mostly unknown, with the exception of Alexandrov's work mentioned above. It may be interesting to investigate it more closely, in order to better understand the deformation of higher-order zeroes of dx .

Currently, there is a gap in the literature on limits of spectral curves, which in particular limits the validity of the proof theorem 2.6, and hence the applicability of the main theorem of this paper, to dx with simple zeroes. Future work with Borot, Bouchard, Chidambaram, and Shadrin will fix this, and will investigate more generally the applicability of limit arguments for topological recursion.

For the BKP hierarchy, similar results should hold. In particular, Alexandrov and Shadrin [AS21] proved an adapted topological recursion for a large class of hypergeometric BKP tau-functions, analogous to theorem 2.6. The analogous ELSV-Eynard-DOSS correspondence between this kind of topological recursion and cohomological field theories has not appeared in the literature, but the special case of completed cycles spin Hurwitz numbers is treated in work of the author with Giacchetto and Lewański [GKL21].

Outline of the paper. Section 2 contains prerequisites. In sections 2.1 and 2.2, we give a short introduction to the Kadomtsev-Petviashvili hierarchy and its space of solutions. In section 2.3, we recall the main ideas from [Kaz09], which we will generalise. In sections 2.4 and 2.5, we recall recent results on hypergeometric tau-functions and their relations to topological recursion and cohomological field theories, and state our main theorem, which is theorem 2.14. We also introduce, in section 2.6, the generating function of triple Hodge numbers, which is the main motivating example of this paper.

In section 3, we prove the main result. Firstly, in section 3.1, we find a change of variables, for any hypergeometric tau-function, that preserves this tau-function after removal of unstable terms, corollary 3.5. In section 3.2, we restrict to the case where topological recursion holds, and use this machinery to obtain tau-functions of intersection numbers, proving our main result. We also determine, in section 3.3, the exact conditions for the change of variables to be finite, in a specific sense. Finally, in section 3.4, we return to the triple Hodge integrals, and prove an explicit version of the main theorem for this case.

Notation. We work over the field of complex numbers \mathbb{C} . We will use the function $\zeta(z) := e^{\frac{z}{2}} - e^{-\frac{z}{2}}$, μ and ν will denote partitions, and $z_\mu := \prod_{i=1}^{\mu_1} i^{m_i(\mu)} m_i(\mu)!$, where $m_i(\mu)$ is the number of parts of μ of size i . We will also consistently write $n := \ell(\mu)$ and $[[n]] := \{1, \dots, n\}$.

On the origin of this paper. An earlier version of this text, only concerning triple Hodge integrals, was written in 2018, shortly after A. Alexandrov informed me of his result. That version appeared in my PhD dissertation [Kra19, Chapter 10]. This paper is an updated and extended version of that chapter.

Acknowledgments. I would like to thank A. Alexandrov for informing me of his theorem on triple Hodge integrals, S. Shadrin for introducing me to the subject and suggesting generalising Kazarian's method to this case, and both of them, G. Carlet, N. Chidambaram, and A. Giacchetto for many interesting discussions.

The author was supported by the Netherlands Organization for Scientific Research and by the Max-Planck-Gesellschaft.

2. PREREQUISITES ON THE KP HIERARCHY AND TOPOLOGICAL RECURSION

In this section, we review some standard notions on the KP hierarchy and its relations to the infinite Grassmannian. We give the main outline of Kazarian's proof of KP for single Hodge integrals, which we will use as a blueprint for our results. We also recall the class of hypergeometric tau-functions, which fulfills a central role in this paper, as well as its relation to topological recursion and cohomological field theories. Finally, we recall the Mariño-Vafa formula for triple Hodge integrals and show it fits in the setup.

2.1. The KP hierarchy. The Kadomtsev-Petviashvili hierarchy is an infinite set of evolutionary differential equations in infinitely many variables. It is a very well-studied system, and some introductions into the subject can be found in [Dico3; Kha99; MJDo0].

Let $\underline{t} = \{t_i\}_{i \geq 1}$ be a set of independent variables and $\partial := \frac{\partial}{\partial t_1}$. Define the pseudo-differential operator (i.e. a Laurent series in ∂^{-1} with coefficients functions in \underline{t} with composition defined formally)

$$(1) \quad L = \partial + u_1 \partial^{-1} + u_2 \partial^{-2} + \dots$$

where the u_j are dependent variables in the t_i . For a pseudo-differential operator O , define O_+ to be its purely differential part, the part without powers of ∂^{-1} . The *Lax formulation of the KP hierarchy* is given by the system of equations

$$(2) \quad \frac{\partial L}{\partial t_i} = [(L^i)_+, L].$$

This is a system of partial differential equations for the u_j , and they can be interpreted as the compatibility equations for the system

$$(3) \quad L\Psi = z\Psi \quad \frac{\partial\Psi}{\partial t_i} = (L^i)_+\Psi.$$

The function Ψ is called the *Baker-Akhiezer function*. The first non-trivial equation, the KP equation, is

$$(4) \quad 3\frac{\partial^2 u_1}{\partial t_2^2} - \frac{\partial}{\partial t_1} \left(4\frac{\partial u_1}{\partial t_3} - 12u_1\frac{\partial u_1}{\partial t_1} - \frac{\partial^3 u_1}{\partial t_1^3} \right) = 0.$$

The Baker-Akhiezer function can be written in the form

$$(5) \quad \Psi = \frac{\tau(\{t_k - \frac{z^{-k}}{k}\})}{\tau(\{t_k\})} \exp\left(\sum_{k=1}^{\infty} t_k z^k\right).$$

Here τ is a single function, called a tau-function, dependent on the t_k , and all dependent variables can be expressed in terms of this one function. This way, the entire hierarchy can be rewritten as bilinear equations for τ called *Hirota equations*.

There is an enrichment of the KP hierarchy, introduced by Takasaki-Takebe [TT95], and called the \hbar -KP hierarchy, which is given by rescaling all $\frac{\partial}{\partial t_i} \rightarrow \hbar \frac{\partial}{\partial t_i}$ in equations (1) and (2), and allowing the u_i to also be power series in \hbar .

2.2. Space of tau-functions and Lie action. The space of solutions of the KP hierarchy is an infinite-dimensional Grassmannian [SS83], which is usually Plücker embedded in a Fock space, i.e. a highest weight module of a certain Clifford algebra. The Hirota equations are then the Plücker relations defining the Grassmannian inside the Fock space. By the boson-fermion correspondence, this can also be expressed in terms of symmetric functions, which is the viewpoint we will adopt here.

Definition 2.1. We write $\Lambda := \mathbb{C}[[p_1, p_2, \dots]]$ for the *space of symmetric functions*, also called the *bosonic Fock space* (of type A). Here the p_k are power-sum functions $p_k = \sum_i X_i^k$ in some countably infinite variable set $\underline{X} = \{X_i\}$.

For other symmetric functions in \underline{X} , e.g. the Schur functions s_λ , we write $s_\lambda(\underline{p}) := s_\lambda(\underline{X})$.

The space of symmetric functions has a projective action of the Lie algebra $\mathfrak{gl}(\infty)$, the algebra of infinite square matrices $(a_{ij})_{i,j \in \mathbb{Z} + \frac{1}{2}}$.¹ This space has a standard basis given by $E_{kl} = (\delta_{ik}\delta_{jl})_{ij}$. Define the *vertex operator*

$$(6) \quad Z(z, w) = \frac{1}{z-w} \left(\exp\left(\sum_{j=1}^{\infty} (z^j - w^j)p_j\right) \exp\left(-\sum_{k=1}^{\infty} (z^{-k} - w^{-k})\frac{1}{k}\frac{\partial}{\partial p_k}\right) - 1 \right).$$

Then expanding this vertex operator as

$$(7) \quad Z(z, w) = \sum_{i,j \in \mathbb{Z} + \frac{1}{2}} Z_{ij} z^{i+1/2} w^{-j-1/2},$$

the assignment $E_{ij} \mapsto Z_{ij}$ is a projective representation of $\mathfrak{gl}(\infty)$, i.e. a representation of a central extension $\widehat{\mathfrak{gl}(\infty)}$.

The matrices $\alpha_k = \sum_{l \in \mathbb{Z} + \frac{1}{2}} E_{l-k, l}$ give rise to the following operators on Λ :

$$(8) \quad a_k := \begin{cases} p_k & k > 0 \\ -k \frac{\partial}{\partial p_{-k}} & k < 0 \\ 0 & k = 0 \end{cases}.$$

We also define the following operators:

$$(9) \quad L_m := \frac{1}{2} \sum_{i=-\infty}^{\infty} :a_i a_{m-i}:, \quad M_l := \frac{1}{6} \sum_{i,j=-\infty}^{\infty} :a_i a_j a_{l-i-j}:,$$

where the $: \cdot :$, the *normal ordering*, means one should order the operators inside in order of decreasing index. All of these operators are in $\widehat{\mathfrak{gl}(\infty)}$.

¹In order to make the Lie bracket well-defined, some decay condition is needed. A common choice is restriction to finitely many diagonals, but there are other options, see e.g. [SS83]. We will remain agnostic on this choice, as in this paper, the required convergence is guaranteed by our constructions.

Theorem 2.2 ([SS83]). *Under the identification $t_k = \frac{p_k}{k}$, the space of KP tau-functions is the orbit of $1 \in \Lambda$ under the action of $\widehat{\mathfrak{gl}(\infty)}$.*

2.3. Single Hodge integrals. In [Kaz09], Kazarian considered the generating function for single Hodge integrals,

$$(10) \quad F_H(u; T_0, T_1, T_2, \dots) := \sum_{g,n} \frac{1}{n!} \sum_{d_1, \dots, d_n \geq 0} \int_{\mathcal{M}_{g,n}} \Lambda(-u^2) \prod_{i=1}^n \psi_i^{d_i} T_{d_i},$$

and showed that its exponent, $Z_H := \exp(F_H)$, is a tau-function for the KP hierarchy, after a certain change of coordinates. Explicitly, this change of coordinates is given as follows: define

$$(11) \quad D = (u+z)^2 z \frac{\partial}{\partial z}.$$

Then we define the T_d in terms of other coordinates q_k by the linear correspondence

$$(12) \quad q_k \leftrightarrow z^k, \quad T_d \leftrightarrow D^d z.$$

The proof consists of three steps, and makes essential use of the ELSV formula [ELSV01] to transform this generating function into a generating function of Hurwitz numbers.

The first step, [Kaz09, Theorem 2.2], is the observation that the generating function for single simple Hurwitz numbers is a tau-function for the KP hierarchy. This is a well-known result, see [Okooo]. In fact, the single simple Hurwitz generating function can be obtained from the trivial τ -function 1 by the action of two very explicit elements of the Lie group associated to $\widehat{\mathfrak{gl}(\infty)}$. The second step, [Kaz09, Theorem 2.3], uses the ELSV formula to rewrite the Hurwitz generating function (after subtracting the unstable geometries) as a generating function for single Hodge integrals. This introduces certain combinatorial factors, that suggest a certain change of coordinates. After this change of coordinates, we obtain Z_H , viewed as a function in q 's.

The third step, [Kaz09, Theorem 2.5] shows that a certain class of coordinate changes preserves solutions of the KP hierarchy, after they are modified with a quadratic function. In essence, this coordinate change is given infinitesimally by the flow along the differential part of an $A \in \widehat{\mathfrak{gl}(\infty)}$, whose polynomial part is exactly the added quadratic function. In this specific case, this quadratic function is exactly the $(0, 2)$ part of the Hurwitz generating function.

In this paper, we will generalise this proof scheme to a more general setting. We will start from a general hypergeometric tau-function in the sense of theorem 2.6 below, corresponding to the first point of the proof.

We obtain a change of coordinates coming from this formalism that can always be completed to an automorphism of KP when correcting with the $H_{0,2}$ of equation (23), without any further assumption, corresponding to the third point of the proof.

If we restrict to the class of hypergeometric tau-functions satisfying topological recursion, we can use the correspondence between topological recursion and cohomological field theories of Eynard and Dunin-Barkowski-Orantin-Shadrin-Spitz [Eyn14; DOSS14], which generalises the ELSV formula and hence gives the second step.

In the particular case of triple Hodge integrals, the role of the ELSV formula is taken by the Mariño-Vafa formula. For explanations on all the required notions and notation, see the following sections.

2.4. Hypergeometric KP tau-functions and topological recursion. An important class of KP tau-functions is given by the hypergeometric tau-functions [KMMM95; OS01a; OS01b], for which we will use the results and notation of [BDKS20a]. In two large families of examples, these satisfy Eynard-Orantin topological recursion [EO07] (or its generalisation to non-simple ramification given by Bouchard-Eynard [BE13]), which we define first. We will confine ourselves to the case of rational spectral curves, as this is the appropriate setting for the Hurwitz-type problems covered.

Definition 2.3 ([EO07; BE13]). A *rational spectral curve* is a quadruple $\mathcal{C} = (\Sigma = \mathbb{P}^1, dx, dy, B)$, where dx and dy are meromorphic one-forms on Σ with no common zeroes, only simple poles of dx , and without poles of dy at zeroes of dx , and $B = B(z_1, z_2) = \frac{dz_1 dz_2}{(z_1 - z_2)^2}$ is a symmetric $(1, 1)$ -form on $\Sigma \times \Sigma$. Write $R \subset \Sigma$ for the set of zeroes of dx , and r_a for the order of vanishing of dx at $a \in R$.

On a rational spectral curve, define a set of symmetric multidifferentials $\{\omega_{g,n}\}_{g \geq 0, n \geq 1}$ on Σ^n via *topological recursion* as follows: first, define the unstable cases by $\omega_{0,1} := y dx$ (this need only be defined locally near the a_i using any primitive of y) and $\omega_{0,2} := B$. Then, for $2g - 2 + (n + 1) > 0$, the stable range, define

$$(13) \quad \omega_{g,n+1}(z_{[n]}, z_{n+1}) := \sum_{a \in R} \sum_{\substack{I \subset \{0, \dots, r_a - 1\} \\ \# I = n}} \operatorname{Res}_{z=a} \frac{\int_a^z \omega_{0,2}(\cdot, z_{n+1})}{\prod_{i=2}^m (\omega_{0,1}(z) - \omega_{0,1}(\sigma_a^i(z)))} \mathcal{W}_{g, |I|+1, n}(\sigma_a^I(z); z_{[n]}),$$

where σ_a is a generator of the local deck transformations of a primitive of dx at a , and

$$(14) \quad \mathcal{W}_{g,m,n}(\zeta_{[m]}; z_{[n]}) := \sum_{\substack{\mu \vdash [m] \\ \bigsqcup_{k=1}^{l(\mu)} N_k = [n] \\ \sum g_k = g + l(\mu) - n}} \prod_{k=1}^{l(\mu)} \omega_{g_k, |\mu_k| + |N_k|}(\zeta_{\mu_k}, z_{N_k})$$

where the prime on the summation means exclusion of any $(g_k, |\mu_k| + |N_k|) = (0, 1)$.

Remark 2.4. Often, the definition of spectral curves involves functions x and y , in stead of their derivatives. However, these functions may not be defined globally on x , e.g. they may – and in this paper will – contain logarithmic terms. As most of the theory of topological recursion (with the notable exception of the global topological recursion of Bouchard-Eynard [BE13]) only depends on the derivatives, I have chosen to use this as a definition.

Theorem 2.5 ([BEO15; BS17]). *Let \mathcal{C} be a rational spectral curve with simple zeroes of dx . A collection $\{\omega_{g,n}\}_{g \geq 0, n \geq 1}$ with $\omega_{0,1} = ydx$ and $\omega_{0,2} = B$ satisfies topological recursion if and only if the following hold:*

- Meromorphicity: For $2g - 2 + n > 0$, $\omega_{g,n}$ extends to a meromorphic form on Σ^n ;
- Linear loop equation: For any g, n , and $a \in R$,

$$(15) \quad \omega_{g,n+1}(z, z_{[n]}) + \omega_{g,n+1}(\sigma_a(z), z_{[n]})$$

is holomorphic near $z = a$ and has a simple zero at $z = a$;

- Quadratic loop equation: For any g, n , and $a \in R$,

$$(16) \quad \omega_{g-1,n+2}(z, \sigma_a(z), z_{[n]}) + \sum_{\substack{g_1 + g_2 = g \\ I \sqcup J = [n]}} \omega_{g_1, |I|+1}(z, z_I) \omega_{g_2, |J|+1}(\sigma_a(z), z_J)$$

is holomorphic near $z = a$ and has a double zero at $z = a$;

- Projection property: For $2g - 2 + n > 0$,

$$(17) \quad \omega_{g,n}(z_{[n]}) = \sum_{a_1, \dots, a_n \in R} \left(\prod_{j=1}^n \operatorname{Res}_{\zeta_j = a_j} \int_{a_j}^{z_j} \omega_{0,2}(z_j, \cdot) \right) \omega_{g,n}(\zeta_{[n]}).$$

If only the meromorphicity and linear and quadratic loop equations hold, the problem is said to satisfy blobbed topological recursion, cf. [BS17]. In this case, the $\omega_{g,n}$ are determined by the spectral curve along with their holomorphic parts at ramification points.

The ‘point’ of topological recursion is that the $\omega_{g,n}$ will often encode enumerative invariants in their Taylor series expansion around a given point of the spectral curve in a given coordinate. For us, this is also the case, as we consider the class given by the following theorem:

Theorem 2.6 ([BDKS20a; BDKS20b]). *Consider two formal power series*

$$(18) \quad \hat{\psi}(\hbar^2, y) := \sum_{k=1}^{\infty} \sum_{m=0}^{\infty} c_{k,m} y^k \hbar^{2m}, \quad \hat{y}(\hbar^2, z) := \sum_{k=1}^{\infty} \hat{y}_k(\hbar^2) z^k := \sum_{k=1}^{\infty} \sum_{m=0}^{\infty} s_{k,m} z^k \hbar^{2m},$$

and their associated hypergeometric \hbar -KP tau-function or Orlov-Scherbin partition function

$$(19) \quad Z(\underline{p}) = e^{F(\underline{p})} = \sum_{\nu \in \mathcal{P}} \exp \left(\sum_{\square \in \nu} \hat{\psi}(-\hbar c_{\square}) \right) s_{\nu}(\underline{p}) s_{\nu} \left(\left\{ \frac{\hat{y}_k}{\hbar} \right\} \right).$$

Define

$$(20) \quad \begin{aligned} \psi(y) &:= \hat{\psi}(0, y), & y(z) &:= \hat{y}(0, z), & x(z) &:= \log z - \psi(y(z)), \\ X(z) &:= e^{x(z)}, & D &:= \frac{\partial}{\partial x}, & Q &:= z \frac{dx}{dz} \end{aligned}$$

and write

$$(21) \quad H_n := \sum_{k_1, \dots, k_n=1}^{\infty} \frac{\partial^n F}{\partial p_{k_1} \cdots \partial p_{k_n}} \Big|_{p=0} X_1^{k_1} \cdots X_n^{k_n}.$$

Then these can be decomposed as

$$(22) \quad H_n = \sum_{g=0}^{\infty} \hbar^{2g-2+n} H_{g,n},$$

with $H_{g,n}$ independent of \hbar , and

$$(23) \quad DH_{0,1}(X(z)) = y(z), \quad H_{0,2}(X(z_1), X(z_2)) = \log \left(\frac{z_1^{-1} - z_2^{-1}}{X_1^{-1} - X_2^{-1}} \right).$$

If moreover $\frac{d\psi(y)}{dy}|_{y=y(z)}$ and $\frac{dy(z)}{dz}$ have analytic continuations to meromorphic functions in z and all coefficients of positive powers of \hbar^2 in $\hat{\psi}(\hbar^2, y(z))$ and $\hat{y}(\hbar^2, z)$ are rational functions of z whose singular points are disjoint from the zeroes of dx , then the n -point differentials

$$(24) \quad \omega_{g,n} := d_1 \cdots d_n H_{g,n} + \delta_{g,0} \delta_{n,2} \frac{dX_1 dX_2}{(X_1 - X_2)^2}$$

can be extended analytically to $(\mathbb{P}^1)^n$ as global rational forms, and the collection of n -point differentials satisfies meromorphicity and the linear and quadratic loop equations, i.e. blobbed topological recursion, for the curve $(\mathbb{P}^1, dx(z), dy(z), B = \frac{dz_1 dz_2}{(z_1 - z_2)^2})$.

Finally, if $\hat{\psi}$ and \hat{y} belong to one of the two families

$$\text{Family I} \quad \hat{\psi}(\hbar^2, y) = \mathcal{S}(\hbar \partial_y) P_1(y) + \log \left(\frac{P_2(y)}{P_3(y)} \right); \quad \hat{y}(\hbar^2, z) = \frac{R_1(z)}{R_2(z)}$$

$$\text{Family II} \quad \hat{\psi}(\hbar^2, y) = \alpha y; \quad \hat{y}(\hbar^2, z) = \frac{R_1(z)}{R_2(z)} + \mathcal{S}(\hbar z \partial_z)^{-1} \log \left(\frac{R_3(z)}{R_4(z)} \right),$$

where $\alpha \in \mathbb{C}^\times$ and the P_i and R_j are arbitrary polynomials such that $\psi(y)$ and $y(z)$ are non-zero, but vanishing at zero, and no singular points of y are mapped to branch points by x , then the n -point differentials also satisfy the projection property, and hence topological recursion, for the curve above.

Remark 2.7. The proof of theorem 2.6 for higher order zeroes of dx uses a limit argument which, although used more often, is currently not fully justified by the literature (in particular, invoking [BE13] is not sufficient). However, this gap will be filled soon.

Remark 2.8. It is possible to allow for constant terms in $\hat{\psi}$ in equation (18), but using quasihomogeneity of the s_ν in equation (19), one can see this can be absorbed in a recaling of the argument of \hat{y} . From the spectral curve point of view, this follows from the fact that the two curves

$$(25) \quad \begin{cases} X(z) &= z e^{-\psi \circ y(z) + \log a} = a z e^{-\psi \circ y(z)} \\ y(z) &= z \end{cases} \quad \text{and} \quad \begin{cases} X(z') &= z' e^{-\psi \circ y(\frac{z'}{a})} \\ y(z') &= \frac{z'}{a} \end{cases}$$

can be identified via $z' = az$.

Remark 2.9. We will consistently use the symbol x for the function which is part of the spectral curve data and X for its exponential, which is the expansion parameter for this class of Hurwitz problems.

We will need different parts of this theorem for the different parts of the proof. In particular, topological recursion is needed to obtain intersection numbers.

2.5. Topological recursion and cohomological field theories. Topological recursion is strongly related to intersection theory of the moduli spaces of curves: there is a quite general correspondence between spectral curves and certain coherent collections of intersection classes in the moduli spaces. These coherent collections are cohomological field theories, which were originally defined by Kontsevich and Manin [KM94] to axiomatise Gromov-Witten theory.

Definition 2.10 ([KM94]). Let V be a vector space with a non-degenerate bilinear form η . A *cohomological field theory (CohFT)* on (V, η) is a collection of maps

$$(26) \quad \Omega_{g,n}: V^{\otimes n} \rightarrow H^*(\overline{\mathcal{M}}_{g,n}),$$

for all $g \geq 0, n \geq 1$ such that $2g - 2 + n > 0$, such that

- $\Omega_{g,n}$ is \mathfrak{S}_n -equivariant with respect to simultaneous permutation of the factors and the marked points;
- with respect to the glueing maps

$$(27) \quad \rho: \overline{\mathcal{M}}_{g-1, n+2} \rightarrow \overline{\mathcal{M}}_{g,n}, \quad \sigma: \overline{\mathcal{M}}_{g, |I|+1} \times \overline{\mathcal{M}}_{h, |J|+1} \rightarrow \overline{\mathcal{M}}_{g+h, |I \sqcup J|},$$

we get

$$\begin{aligned} \rho^* \Omega_{g,n}(v_{[n]}) &= \Omega_{g-1, n+2}(v_{[n]} \otimes \eta^\dagger), \\ \sigma^* \Omega_{g+h, |I|+|J|}(v_I \otimes v_J) &= \Omega_{g, |I|+1} \otimes \Omega_{h, |J|+1}(v_I \otimes \eta^\dagger \otimes v_J); \end{aligned}$$

There is a large group acting on the space of CohFTs, called the Givental group [Giv01; Sha09; Tel12]. It consists of $R(u) \in \text{Id} + u \text{End}(V)[[u]]$ such that $R(u)R^\dagger(-u) = \text{Id}$. There is also a group of translation actions by $T(u) \in u^2 V[[u]]$. For an exposition which is well-adapted to the current setting, but assumes simple ramification of x , see [GKL21, Section 2.2].

Theorem 2.11 ([Eyn14; DOSS14; BKS20]). *Consider a compact rational spectral curve $(\mathbb{P}^1, dx, dy, B)$, and define V^* to be the space of residueless meromorphic one-forms on \mathbb{P}^1 with poles only at $a \in R$ of order at most $r_a + 1$. Choose a basis $\{d\xi^j\}_{j \in J}$ of V^* with dual basis e_j and define $d\xi_k^j = (d \circ \frac{1}{dx})^k d\xi^j$. Then*

$$(28) \quad \omega_{g,n}(z_1, \dots, z_n) = \sum_{j_1, \dots, j_n \in J} \int_{\mathcal{M}_{g,n}} \Omega_{g,n}(e_{j_1} \otimes \dots \otimes e_{j_n}) \prod_{i=1}^n \sum_{k_i=0}^{\infty} \psi_i^{k_i} d\xi_{k_i}^{j_i}(z_i),$$

where Ω is a cohomological field theory on V , given explicitly by acting on a direct sum of Witten r_a -spin classes for all ramification points of order r_a by an R and T determined, respectively, by B and y .

Remark 2.12. The space V^* is naturally related to the projection property of theorem 2.5: the $d\xi_k^j$ span the image of the projection operator. Its dimension, the rank of the CohFT, equals the degree of the divisor of zeroes of dx .

There are two common choices for the basis $d\xi^j$, depending on a local coordinate ζ_a around a ramification point a such that $x(z) = \zeta_a(z)^{r_a} + x(a)$. One is $d\xi^{a,k}(z) = \text{Res}_{z'=a} \left(\int_a^{z'} B(z, \cdot) \right) \frac{d\zeta(z')}{\zeta(z')^k}$, with $1 \leq k \leq r_a - 1$, cf. [BKS20, Equation (80)], while the other is $\xi^a(z) = \int^z \frac{B(\zeta_a, \cdot)}{d\zeta_a} \Big|_{\zeta_a=0}$, in case $r_a = 2$, cf. [GKL21, Equation (2.23)]. Both have merit, depending on the situation, but they are not compatible.

Furthermore, several normalisation conventions exist for the recursion operator linking $d\xi_k^j$ to $d\xi_{k+1}^j$. These different conventions can be related by rescaling Ω and the correlators, using that the integrand must be of degree $3g - 3 + n$.

So the $\omega_{g,n}$ we are concerned with can be expanded in different ways: as a formal series around $X = 0$ by theorem 2.6, and on a basis of meromorphic differentials with poles at the zeroes of dx by theorem 2.11. The change of variables we require is found by relating these different expansions.

In order to apply the Eynard-DOSS correspondence to get a good change of variables, we will want to take a different basis of V^* . It turns out to be useful to relate to powers of our preferred coordinate z , so the basis we take is $\xi^j := \left(\frac{dx}{dz}\right)^{-1} z^j = \frac{d}{dx} \frac{z^{j+1}}{j+1}$.

Definition 2.13. Let Ω be a cohomological field theory on a space (V, η) with a basis $\{e_j\}_{j \in J}$. Its *generating function* F_Ω is defined as

$$(29) \quad F_\Omega(\{T_k^j \mid k \geq 0, j \in J\}) := \sum_{\substack{g,n \\ 2g-2+n>0}} \frac{\hbar^{2g-2+n}}{n!} \sum_{j_1, \dots, j_n \in J} \int_{\mathcal{M}_{g,n}} \Omega(e_{j_1} \otimes \dots \otimes e_{j_n}) \prod_{i=1}^n \sum_{k_i=1}^{\infty} \psi_i^{k_i} T_{k_i}^{j_i},$$

where we write $\{T^j\}$ for the basis of V^* dual to $\{e_j\}$ and $T_k^j = T^j \otimes p_k$.

The main theorem of this paper is the following:

Theorem 2.14. *If a cohomological field theory Ω is obtained from theorem 2.11 applied to either family in theorem 2.6, then the exponential of $F_\Omega(\mathcal{I}(q))$ is an \hbar -KP tau-function in $\{t_d = \frac{q_d}{d}\}$, where the $T_k^j(q)$ are defined by*

$$(30) \quad T_{-1}^j = \frac{1}{j+1} q_{j+1}, \quad T_{k+1}^j = \sum_{m=1}^{\infty} \sum_{l=0}^{\infty} \mathcal{T}_l q_{m+l} \frac{\partial}{\partial q_m} T_k^j, \quad \text{with } \mathcal{T}_l \text{ given by } -Q(z)^{-1} = \sum_{l=0}^{\infty} \mathcal{T}_l z^l.$$

The proof of this theorem is given in proposition 3.8.

Remark 2.15. The proof of this theorem does not use anything specific to the families mentioned, it just requires topological recursion to obtain a cohomological field theory. As soon as topological recursion is proved for another hypergeometric tau-function and the spectral curve fits in the scope of theorem 2.11, this theorem generalises. For an example, see the next section.

2.6. The Mariño-Vafa formula and KP for topological vertex amplitudes. A particularly interesting family of hypergeometric tau-functions is given by the theory of the topological vertex, or triple Hodge integrals. For the triple Hodge integrals, the ELSV-type formula required is the Mariño-Vafa formula [MV02]. This theory is the particular case for \mathbb{C}^3 of the Gromov-Witten theory of toric Calabi-Yau threefolds, conjectured by Bouchard-Klemm-Mariño-Pasquetti [BKMP09] to satisfy topological recursion. The case we are interested

in was proved in [Cher8; Zoo9], as well as in [Eyn1] as an example of the general correspondence of theorem 2.11, while the general BKMP conjecture was proved in [EO15]. Interestingly, this family does not quite fit in the families of theorem 2.6 for general parameters.

In this section, we use the triple Hodge integrals as an example of our general theory, using methods slightly adapted to this special case. We will see in section 3.3 why this case is particularly nice.

Definition 2.16. *The triple Hodge cohomological field theory with Calabi-Yau condition is the one-dimensional CohFT $\text{TH}_{g,n}(a, b, c) = \Lambda(a)\Lambda(b)\Lambda(c)$, where the parameters a, b, c satisfy $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} = 0$.*

We write

$$(31) \quad F_{\text{TH}}(a, b, c; \underline{T}) := F_{\text{TH}(a,b,c)}(\underline{T}).$$

An adapted application of theorem 2.14 is given in the following theorem. This theorem has already been proved by Alexandrov [Ale20], here we give a new proof.

Theorem 2.17 ([Ale20, Theorem 2]). *Define $T_0(\underline{q}) := q_1$, $T_{k+1}(\underline{q}) := \sum_{m=1}^{\infty} m(u^2 q_m + u \frac{w+2}{\sqrt{w+1}} q_{m+1} + q_{m+2}) \frac{\partial}{\partial q_m} T_k$. Then*

$$(32) \quad F_{\text{TH}}(-u^2, -u^2 w, \frac{u^2 w}{w+1}; \{T_k(\underline{q})\})$$

is a solution of the KP hierarchy with respect to the variables $\{t_d = \frac{q_d}{d}\}$, identically in u and w .

In this particular case, we will make slightly different choices to end up with the formulation above.

Remark 2.18. Note that the triple $a = -u^2$, $b = -u^2 w$, $c = \frac{u^2 w}{w+1}$ does indeed satisfy $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} = 0$, and moreover any triple satisfying this condition can be written this way.

Remark 2.19. In the limit $w \rightarrow 0$, this theorem reduces to the main theorem, 2.1, of [Kaz09]. In the limit $u \rightarrow 0$, it reduces to the Witten-Kontsevich theorem [Wit91; Kon92]: in that limit $T_d \rightarrow (2d-1)!! q_{2d+1}$ and independence of even parameters reduces the KP hierarchy to the KdV hierarchy.

Before giving the Mariño-Vafa formula, note that in genus zero

$$(33) \quad \int_{\mathcal{M}_{0,n}} \frac{\Lambda(a)\Lambda(b)\Lambda(c)}{\prod_{i=1}^n (1 - \mu_i \psi_i^{d_i})} = |\mu|^{n-3}$$

for $n \geq 3$, and this serves as a definition for $n = 1, 2$. These terms are not included in F_{TH} .

Theorem 2.20 (Mariño-Vafa formula, [MV02; LLZ03; OP04]). *There is a relation between triple Hodge integrals and characters of symmetric groups, as follows:*

$$(34) \quad \exp \left(\sum_{\mu} \sum_{g=0}^{\infty} \frac{(w+1)^{g+n-1}}{|\text{Aut } \mu|} \prod_{i=1}^n \frac{\prod_{j=1}^{\mu_i-1} (\mu_i + jw)}{(\mu_i - 1)!} \int_{\mathcal{M}_{g,n}} \frac{\Lambda(-1)\Lambda(-w)\Lambda(\frac{w}{w+1})}{\prod_{i=1}^n (1 - \mu_i \psi_i)} \beta^{2g-2+n+|\mu|} p_{\mu} \right) \\ = \sum_{m=0}^{\infty} \sum_{\mu, \nu \vdash m} \frac{\chi_{\mu}^{\nu}}{z_{\mu}} e^{(1+\frac{w}{2})\beta f_2(\nu)} \prod_{\square \in \nu} \frac{\beta w}{\varsigma(\beta w h_{\square})} p_{\mu}.$$

On the right-hand side the sum is over all partitions ν of size equal to $|\mu|$, the product is over all boxes in the Young diagram of ν , and h_{\square} is the hook length of the box \square . Furthermore, $f_2(\nu) = \frac{1}{2} \sum_j (\nu_j - j + \frac{1}{2})^2 - (-j + \frac{1}{2})^2$ is the shifted symmetric sum of squares.

Remark 2.21. Even though it seems the triple Hodge class in this formula only depends on one parameter, w , the parameter β can be interpreted in this way as well, entering as a cohomological grading parameter. Hence, the formula does govern the entire generating function of triple Hodge integrals.

In the limit $w \rightarrow 0$, the Mariño-Vafa formula reduces to the ELSV formula, as the product over boxes simplifies to the hook length formula for the dimension of the $\mathfrak{S}_{|\mu|}$ -representation associated to ν .

Remark 2.22. This formula is perfectly well-behaved for $w = -1$, but theorem 2.17 does not make sense in this case. From the general theorem 2.14, we will see that in this case X is a Möbius transformation, and hence conforms to corollary 3.6.

By symmetry in the arguments of the Λ -classes, the point $w = -1$ is equivalent to the limit $w \rightarrow \infty$, which in the conventional formulation of the Mariño-Vafa formula is the initial condition for the cut-and-join equation used to prove the formula, see [Zho03, Theorem 3.3]. In this case, the integral reduces to $\int_{\overline{\mathcal{M}}_{g,1}} \lambda_g \psi^{2g-2}$ by Mumford's relation. These integrals were calculated by Faber and Pandharipande [FP00].

The right-hand side of the Mariño-Vafa formula is a hypergeometric KP tau-function, which can be seen explicitly by the following lemma. In essence this lemma was used by both [LLZ03; OP04] to prove the Mariño-Vafa formula.

Lemma 2.23. *Introduce an extra parameter \hbar in equation (34) by rescaling $\beta \rightarrow \hbar\beta$ and $p_k \rightarrow \hbar^{-k}p_k$ to obtain*

$$(35) \quad \exp \left(\sum_{\mu} \sum_{g=0}^{\infty} \frac{(w+1)^{g+n-1}}{|\text{Aut } \mu|} \prod_{i=1}^n \frac{\prod_{j=1}^{\mu_i-1} (\mu_i + jw)}{(\mu_i - 1)!} \int_{\mathcal{M}_{g,n}} \frac{\Lambda(-1)\Lambda(-w)\Lambda(\frac{w}{w+1})}{\prod_{i=1}^n (1 - \mu_i \psi_i)} \hbar^{2g-2+n} \beta^{2g-2+n+|\mu|} p_{\mu} \right) \\ = \sum_{m=0}^{\infty} \sum_{\mu, \nu \vdash m} \frac{\chi_{\mu}^{\nu}}{z_{\mu}} e^{(1+\frac{w}{2})\hbar\beta f_2(\nu)} \prod_{\square \in \nu} \frac{\beta w}{\varsigma(\hbar\beta w \hbar_{\square})} p_{\mu}.$$

This right-hand side may alternatively be written as a hypergeometric \hbar -KP tau-function in the shape of theorem 2.6, with

$$\hat{\psi}(\hbar^2, y) = -\beta y, \quad \hat{y}(\hbar^2, z) = \sum_{k=1}^{\infty} \frac{\hbar}{\varsigma(\hbar k \beta w)} (\beta w z)^k \quad X(z) = z(1 - \beta w z)^{1/w}.$$

Proof. By basic theory of symmetric functions, $\sum_{\mu \vdash m} \frac{\chi_{\mu}^{\nu}}{z_{\mu}} p_{\mu} = s_{\nu}(\underline{p})$. Also, by [OP04, Equations (o.6), (o.7)],

$$\frac{1}{\prod_{\square \in \nu} q^{\hbar_{\square}/2} - q^{-\hbar_{\square}/2}} = q^{-|\nu|/2 - f_2(\nu)/2} s_{\nu}(1, q^{-1}, q^{-2}, \dots)$$

where here the q^{-k} are the ‘usual’ variables, i.e. the ones in which s_{ν} is symmetric, not the power sum variables.

Writing $q = e^{\hbar\beta w}$ and using that $f_2(\nu) = \sum_{\square \in \nu} c_{\square}$ gives

$$\sum_{m=0}^{\infty} \sum_{\mu, \nu \vdash m} \frac{\chi_{\mu}^{\nu}}{z_{\mu}} e^{(1+\frac{w}{2})\hbar\beta f_2(\nu)} \prod_{\square \in \nu} \frac{\beta w}{\varsigma(\hbar\beta w \hbar_{\square})} p_{\mu} = \sum_{m=0}^{\infty} \sum_{\nu \vdash m} s_{\nu}(\underline{p}) e^{\sum_{\square \in \nu} \hbar\beta c_{\square}} s_{\nu} \left(\left\{ \frac{\beta w}{e^{\hbar\beta w(k+\frac{1}{2})}} \right\}_{k=0}^{\infty} \right).$$

To revert to power-sum variables, we use that $p_k \left(\left\{ \frac{\beta w}{e^{\hbar\beta w(k+\frac{1}{2})}} \right\}_{k=0}^{\infty} \right) = \frac{(\beta w)^k}{\varsigma(\hbar k \beta w)}$, and inserting this in the definition of \hat{y} yields the result. \square

Zhou [Zho10] also explored this relation between triple Hodge integrals and integrable hierarchies, extending it to the 2-Toda hierarchy and to certain relative Gromov-Witten theories.

3. KP HIERARCHY FOR INTERSECTION NUMBERS

In this section, we will formulate and prove the main theorem, generalising Kazarian’s method to the generating functions of intersection numbers coming from hypergeometric tau-functions.

3.1. The change of variables. We will interpret any $X(z)$ defined by equations (18) and (20) as giving a change of coordinates. For this, define a linear correspondence Θ between power series in X or z on the one hand and linear series in p or q on the other by

$$(36) \quad p_k \leftrightarrow X^k, \quad q_m \leftrightarrow z^m.$$

This defines a change of coordinates as follows:

Definition 3.1. We define a linear morphism between power series in $\{p_m\}_{m \geq 1}$ and $\{q_d\}_{d \geq 1}$ by

$$(37) \quad p_k(\underline{q}) = \sum_{m=k}^{\infty} c_k^m q_m \quad \text{with } c_k^m \text{ given by} \quad X^k = \sum_{m=k}^{\infty} c_k^m z^m.$$

In order to make this change of coordinates and remain within the realm of solutions of the KP hierarchy, we should flow along the action of the infinite general linear algebra. Hence, we should find the infinitesimal flow associated to this change. For this, we introduce a flow parameter β by

$$(38) \quad X_{\beta}(z) := \frac{1}{\beta} X(\beta z) = z e^{-\psi(y(\beta z))},$$

such that $X_0(z) = z$ and $X_1(z) = X(z)$.

Lemma 3.2. For $X_\beta(z) := \frac{1}{\beta}X(\beta z)$, where $X(z) = z + \mathcal{O}(z^2)$, and with $Q(z) := \frac{z}{X} \frac{dX}{dz}$, the flow along β of the function X_β is given by

$$(39) \quad \frac{\partial X_\beta}{\partial \beta} = \left(1 - \frac{1}{Q(\beta z)}\right) \frac{z}{\beta} \frac{\partial X_\beta}{\partial z} = \frac{1}{\beta} (Q(\beta z) - 1) X_\beta.$$

Proof. By definition of Q , $X = \frac{z}{Q(z)} \frac{dX}{dz}$. Therefore,

$$\begin{aligned} \frac{\partial X_\beta}{\partial \beta} &= \frac{\partial}{\partial \beta} \left(\frac{1}{\beta} X(\beta z) \right) = \frac{z}{\beta} \frac{dX}{dz} \Big|_{z \rightarrow \beta z} - \frac{1}{\beta^2} X(\beta z) \\ &= \frac{z}{\beta} \frac{dX}{dz} \Big|_{z \rightarrow \beta z} - \frac{1}{\beta^2} \left(\frac{z}{Q(z)} \frac{dX}{dz} \right) \Big|_{z \rightarrow \beta z} = \left(1 - \frac{1}{Q(\beta z)}\right) \frac{z}{\beta} \frac{\partial X_\beta}{\partial z}. \quad \square \end{aligned}$$

We will use this with [Kaz09, Theorem 2.5], which uses the $\widehat{\mathfrak{gl}(\infty)}$ action on τ -functions:

Theorem 3.3 ([Kaz09]). *In the situation of a correspondence like equation (37), there is a quadratic function $Q(p)$ such that the transformation sending an arbitrary series $\Phi(\underline{p})$ to the series $\Psi(\underline{q}) = (\Phi + Q)|_{p \rightarrow p(\underline{q})}$ is an automorphism of the KP hierarchy: it sends solutions to solutions.*

The function $Q(\underline{p})$ is not unique.

Proposition 3.4. *In the general situation of theorem 2.6, without analytic assumptions, the quadratic function for the change of definition 3.1 can be taken to be $-\frac{1}{2}\Theta(H_{0,2})$.*

Proof. Consider the more general linear correspondence Θ_β between power series in X_β or z on the one hand and linear series in p or q on the other by

$$p_k \leftrightarrow X_\beta^k, \quad q_m \leftrightarrow z^m.$$

This gives a linear morphism between power series in $\{p_m\}_{m \geq 1}$ and $\{q_d\}_{d \geq 1}$ by

$$p_k(\beta; \underline{q}) = \sum_{m=k}^{\infty} c_k^m q_m \quad \text{with } c_k^m \text{ given by} \quad X_\beta^k = \sum_{m=k}^{\infty} c_k^m z^m,$$

such that $p_k(0; \underline{q}) = q_k$.

Under Θ_β , the operator $z^{m+1} \frac{\partial}{\partial z}$ transforms into $\sum_{k=1}^{\infty} k q_{m+k} \frac{\partial}{\partial q_k}$, which is the differential part of $L_m(\underline{q})$. The polynomial part of this operator is

$$\frac{1}{2} \sum_{k=1}^{m-1} q_k q_{m-k},$$

which under the correspondence transforms into

$$\frac{1}{2} \sum_{k=1}^{m-1} z_1^k z_2^{m-k} = \frac{1}{2} z_1 z_2 \frac{z_1^{m-1} - z_2^{m-1}}{z_1 - z_2} = -\frac{1}{2} \frac{z_1^{m-1} - z_2^{m-1}}{z_1^{-1} - z_2^{-1}}.$$

Therefore, the correction to be made to lemma 3.2 to obtain a KP-preserving flow is found by the substitution $f(z) z \frac{\partial}{\partial z} \rightarrow -\frac{1}{2} \frac{1}{z_1^{-1} - z_2^{-1}} (z_1^{-1} f(z_1) - z_2^{-1} f(z_2))$ for a series $f(z) \in z\mathbb{C}[[z]]$. Note that $\frac{1}{\beta} \left(1 - \frac{1}{Q(\beta z)}\right)$ satisfies these requirements, and we find that the differential operator of lemma 3.2 needs to be completed by

$$-\frac{1}{2\beta(z_1^{-1} - z_2^{-1})} \left(z_1^{-1} \left(1 - \frac{1}{Q(\beta z_1)}\right) - z_2^{-1} \left(1 - \frac{1}{Q(\beta z_2)}\right) \right) = \frac{1}{2\beta(z_1^{-1} - z_2^{-1})} \left(\frac{1}{z_1 Q(\beta z_1)} - \frac{1}{z_2 Q(\beta z_2)} \right) - \frac{1}{2\beta}.$$

By a similar calculation as for lemma 3.2,

$$\frac{\partial z}{\partial \beta} \Big|_{X \text{ const.}} = \frac{1}{\beta} \left(\frac{1}{Q(\beta z)} - 1 \right) z,$$

from which it follows that

$$\begin{aligned} -\frac{\partial H_{0,2}}{\partial \beta} \Big|_{X \text{ const.}} &= -\frac{\partial}{\partial \beta} \log \left(\frac{z_1^{-1} - z_2^{-1}}{X_1^{-1} - X_2^{-1}} \right) \Big|_{X \text{ const.}} \\ &= \frac{1}{z_1^{-1} - z_2^{-1}} \left(z_1^{-2} \frac{\partial z_1}{\partial \beta} \Big|_{X \text{ const.}} - z_2^{-2} \frac{\partial z_2}{\partial \beta} \Big|_{X \text{ const.}} \right) \\ &= \frac{1}{\beta(z_1^{-1} - z_2^{-1})} \left(\frac{1}{z_1 Q(\beta z_1)} - \frac{1}{z_2 Q(\beta z_2)} \right) - \frac{1}{\beta}, \end{aligned}$$

which, up to a factor 2, is exactly the polynomial correction needed.

From these calculations, we find that

$$A := \left(1 - \frac{1}{Q(\beta z)}\right) \frac{z}{\beta} \frac{\partial}{\partial z} - \frac{1}{2} \frac{\partial H_{0,2}}{\partial \beta} \Big|_{X \text{ const.}}$$

corresponds to a linear combination of L_m under Θ_β , and hence preserves KP. Now consider a KP tau-function $\Phi(\underline{p})$ and define the function $Z(\beta, \underline{q}) = \exp(\Phi(\underline{p}(\beta, \underline{q}) - \frac{1}{2}\Theta(H_{0,2}))$. Then

$$(40) \quad \frac{\partial}{\partial \beta} Z = \left(\sum_{k=1}^{\infty} \frac{\partial p_k(\beta, \underline{q})}{\partial \beta} \frac{\partial}{\partial p_k} - \frac{1}{2} \Theta \left(\frac{\partial H_{0,2}}{\partial \beta} \Big|_{X \text{ const.}} \right) \right) Z$$

$$(41) \quad = \Theta \left(\left(1 - \frac{1}{Q(\beta z)}\right) \frac{z}{\beta} \frac{\partial}{\partial z} - \frac{1}{2} \frac{\partial H_{0,2}}{\partial \beta} \Big|_{X \text{ const.}} \right) Z(\beta)$$

$$(42) \quad = \Theta(A) Z(\beta)$$

As $Z(0) = Z$, and $\Theta(A)$ preserves τ -functions of KP, this automorphism does indeed preserve solutions. \square

Corollary 3.5. *For $Z(\underline{p})$ defined by equation (19), $Z(\underline{p}) \exp(-\Theta(\hbar^{-1}H_{0,1} + \frac{1}{2}H_{0,2})) \Big|_{\underline{p} \rightarrow \underline{p}(\underline{q})}$ is also an \hbar -KP tau-function, whose logarithm does not contain unstable terms.*

Proof. As all equations in the KP hierarchy only contain at least second derivatives of F , addition of a linear term $-\Theta(\hbar^{-1}H_{0,1})$ preserves solutions. By proposition 3.4, subtracting the $(0, 2)$ term and changing $p \mapsto p(\underline{q})$ is an automorphism as well. \square

Corollary 3.6. *In case $X(z)$ is a Möbius transformation with the shape of equation (20), i.e. $X(z) = \frac{az}{1+bz}$ (taking into account remark 2.8), this quadratic function can be taken to be 0.*

Proof. By direct calculation,

$$H_{0,2}(z_1, z_2) = \log \left(\frac{z_1^{-1} - z_2^{-1}}{X(z_1)^{-1} - X(z_2)^{-1}} \right) = \log a.$$

Comparing this with the proof of proposition 3.4, the quadratic correction is needed to complete the operator A , which only depends on $\frac{\partial H_{0,2}}{\partial \beta}$. As this vanishes in the present case, we may as well omit the entire correction. \square

Remark 3.7. The usual B -function of topological recursion,

$$(43) \quad B(z_1, z_2) = \frac{dz_1 dz_2}{(z_1 - z_2)^2} = d_1 d_2 \log(z_1^{-1} - z_2^{-1}),$$

is invariant under all Möbius transformations, so $d_1 d_2 H_{0,2}$ vanishes if X is any Möbius transformation. However, this is not the case for $H_{0,2}$ itself: it is invariant under a one-dimensional subgroup, changes by a constant under the two-dimensional subgroup above, but under other Möbius transformations also changes by addition of terms like $\log z_i$.

Viewed another way, these more general Möbius transformations would take us out of the realm of formal power series in z . However, in a space of functions, a shift $z \mapsto z + c$ does preserve the KP hierarchy, so if the formal power series converges to a function on a large enough domain, this shift does preserve KP. This argument is essentially taken from [Kaz09, Section 8]. In particular, under the ‘natural analytic assumptions’ of [BDKS20b, section 1.3], i.e. the assumptions in the second part of theorem 2.6, the $H_{g,n}$ do extend to rational functions on all of \mathbb{P}^1 , so this shift is well-defined.

3.2. KP for intersection numbers. Now we will restrict to the cases where topological recursion, and hence theorem 2.11, can be used, in order to relate to intersection numbers. In this case, the following holds from equation (28).

$$(44) \quad F(\underline{p}) = \left(\hbar^{-1}H_{0,1} + \frac{1}{2}H_{0,2} + \sum_{2g-2+n>0} \frac{\hbar^{2g-2+n}}{n!} \sum_{j_1, \dots, j_n \in J} \int_{\mathcal{M}_{g,n}} \Omega_{g,n}(e_{j_1} \otimes \dots \otimes e_{j_n}) \prod_{i=1}^n \sum_{k_i=0}^{\infty} \psi_i^{k_i} \xi_{k_i}^{j_i}(z_i) \right) \Big|_{X_i^{k_i} \rightarrow p_{k_i}},$$

if we define $\xi_k^j(z) := \int_{z'=r_j}^z d\xi_k^j(z')$, noting that due to the shape of the $H_{g,n}$ in equations (21) and (22) and $X(z)$ in equation (20), the $H_{g,n}$ have no constant terms in z_i .

Under the correspondence $p_k \leftrightarrow X^k$, $q_m \leftrightarrow z^m$ of definition 3.1, we define T_k^j by

$$(45) \quad T_k^j(\underline{p}) \leftrightarrow \frac{1}{dx} d\xi_k^j(z) = (-D)^{k+1} \frac{z^{j+1}}{j+1},$$

with D as in equation (20). Explicitly,

$$(46) \quad T_{-1}^j = \frac{1}{j+1} q_{j+1}, \quad T_{k+1}^j = \sum_{m=1}^{\infty} \sum_{l=0}^{\infty} \mathcal{T}_l q_{m+l} \frac{\partial}{\partial q_m} T_k^j, \quad \text{with } \mathcal{T}_l \text{ given by } -Q(z)^{-1} = \sum_{l=0}^{\infty} \mathcal{T}_l z^l.$$

Note that, even though the recursion operator for the T_k^j may have infinitely many terms, its alternate description via equation (45) ensures they are well-defined in Λ .

Therefore, by definition,

$$(47) \quad F_{\Omega}(T(\underline{p})) = \Theta(\hbar^{-1}H_{0,1} + \frac{1}{2}H_{0,2}) + \sum_{2g-2+n>0} \frac{\hbar^{2g-2+n}}{n!} \sum_{j_1, \dots, j_n \in J} \int_{\mathcal{M}_{g,n}} \Omega_{g,n}(e_{j_1} \otimes \dots \otimes e_{j_n}) \prod_{i=1}^n \sum_{k_i=0}^{\infty} \psi_i^{k_i} T_{k_i}^{j_i}(\underline{p})$$

is the logarithm of a tau-function, where $\{e_j\}$ is the dual basis to the basis $\{d\xi_0^j\}$ of V^* .

Proposition 3.8. *Suppose the pair of functions $(\hat{\psi}, \hat{y})$ lies in family I or II of theorem 2.6, and let Ω be the cohomological field theory associated to the related topological recursion via theorem 2.11. Then*

$$(48) \quad Z_{\Omega}(\underline{q}) = \exp(F_{\Omega}(T(p(\underline{q}))))$$

is an \hbar -KP tau-function.

Proof. Apply corollary 3.5 to $\exp(F_{\Omega}(T(p)))$ using equation (47). \square

3.3. Finiteness of the transformation. The operator A in the proof of proposition 3.4 corresponds to a finite sum of L_m if and only if $Q(z)^{-1}$ is a polynomial in z . As this case seems particularly nice, we will investigate it here.

Write $P(z) = Q(z)^{-1}$ for this polynomial, and write $r+1$ for its degree. From equation (20), it follows that $P(0) = 1$, so we may write

$$(49) \quad P(z) = \prod_{j=1}^{r+1} (1 - c_j z).$$

We immediately see that $dx(z) = \frac{dz}{zP(z)}$, and hence the spectral curve has a unique ramification point, ∞ , of ramification index r . This is also the rank of the associated Frobenius algebra. But we can do better. By calculating the residues in v of

$$(50) \quad \frac{v^{r+1} dv}{(1 - vz) \prod_{k=1}^{r+1} (v - c_k)}$$

and using that they sum to zero, one can check that (if all c_j are distinct)²

$$(51) \quad \frac{dx}{dz} = \frac{1}{z} + \sum_{j=1}^{r+1} \frac{c_j^{r+1}}{\prod_{k \neq j} (c_j - c_k)} \frac{1}{1 - c_j z},$$

from which we see that

$$(52) \quad x(z) = \log z - \sum_{j=1}^{r+1} \prod_{k \neq j} \left(1 - \frac{c_k}{c_j}\right)^{-1} \log(1 - c_j z).$$

If $r = 0$, dx has two (simple) poles, and hence no zeroes. In fact, in this case, X is a Möbius transformation.

If $r = 1$, this recovers the triple Hodge curve, studied in section 3.4 below.

If $r > 1$, the Frobenius algebra is not semi-simple: it seems to be a deformation of the algebra corresponding to Witten's $r+1$ -spin cohomological field theory, which is given by $x = y^{r+1}$, cf. [Wit93; DNOPS16; BCEG21]. This class fits in Alexandrov's theory of the deformed generalised Kontsevich model [Ale21]: it seems like it is a complementary subspace of the polynomial deformations of the Witten $r+1$ -spin theory.

Interestingly, except for special choices of c_j , these cases seem not to be covered in the two families in theorem 2.6 for which topological recursion is proved (for any choice of y). Even the $r = 1$ case does not fall in that scope, unless $\frac{c_1}{c_2} \in \mathbb{Q}$.

²If some c_j coincide, the residue argument still holds, but the result changes.

3.4. The case of triple Hodge integrals. Let us now consider the special case of triple Hodge integrals. The approach taken in this section overlaps with the previous results, but is also slightly different in details, adapted to this specific problem. In this case, the ELSV-type formula is completely explicit, and there is no need to take the detour via topological recursion.

The coordinate change we want to perform is inspired by the Mariño-Vafa formula.

$$\begin{aligned}
 (53) \quad F(w, \beta; \underline{p}) &= \log \left(\sum_{m=0}^{\infty} \sum_{\mu, \nu \vdash m} \frac{\chi_{\mu}^{\nu}}{z_{\mu}} e^{(1+\frac{w}{2})\beta f_2(\nu)} \prod_{\square \in \nu} \frac{\beta w}{\zeta(\beta w h_{\square})} p_{\mu} \right) \\
 &= \sum_{\mu} \sum_{g=0}^{\infty} \frac{(w+1)^{g+n-1}}{|\text{Aut } \mu|} \prod_{i=1}^n \frac{\prod_{j=1}^{\mu_i-1} (\mu_i + jw)}{(\mu_i - 1)!} \int_{\overline{\mathcal{M}}_{g,n}} \frac{\Lambda(-1)\Lambda(-w)\Lambda(\frac{w}{w+1})}{\prod_{i=1}^n (1 - \mu_i \psi_i)} \beta^{2g-2+n+|\mu|} p_{\mu}.
 \end{aligned}$$

As $2g - 2 + n + |\mu| = \frac{2}{3} \dim \overline{\mathcal{M}}_{g,n} + \sum_{i=1}^n (\mu_i + \frac{1}{3})$ and $g + n - 1 = \frac{1}{3} \dim \overline{\mathcal{M}}_{g,n} + \sum_{i=1}^n \frac{2}{3}$, we get after rewriting $u := \beta^{\frac{1}{3}}(w+1)^{\frac{1}{6}}$

$$\begin{aligned}
 (54) \quad F(w, \beta; \underline{p}) &= \sum_{\mu} \frac{1}{|\text{Aut } \mu|} \sum_{g=0}^{\infty} \prod_{i=1}^n u^4 \frac{\prod_{j=1}^{\mu_i-1} (\mu_i + jw)\beta}{(\mu_i - 1)!} \int_{\overline{\mathcal{M}}_{g,n}} \frac{\Lambda(-u^2)\Lambda(-u^2w)\Lambda(\frac{u^2w}{w+1})}{\prod_{i=1}^n (1 - \mu_i u^2 \psi_i)} p_{\mu} \\
 &= \sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \int_{\overline{\mathcal{M}}_{g,n}} \Lambda(-u^2)\Lambda(-u^2w)\Lambda(\frac{u^2w}{w+1}) \prod_{i=1}^n \sum_{d=0}^{\infty} T_d(\underline{p}) \psi_i^d \\
 &= F_{\text{TH}}\left(-u^2, -u^2w, \frac{u^2w}{w+1}; T(\underline{p})\right) + H_{0,1} + H_{0,2},
 \end{aligned}$$

where

$$(55) \quad T_d(\underline{p}) := \sum_{m=1}^{\infty} \frac{\prod_{j=1}^{m-1} (m + jw)}{(m-1)!} m^d u^{2d+4} \beta^{m-1} p_m.$$

Hence, our goal is to show that this change of variables and addition of the unstable terms preserves solutions of the KP hierarchy.

Lemma 3.9. *The following two expressions are inverse to each other:*

$$(56) \quad X(z) = \frac{z}{1 + (w+1)\beta z} \left(\frac{1 + \beta z}{1 + (w+1)\beta z} \right)^{\frac{1}{w}}; \quad z(X) = \sum_{m=1}^{\infty} \frac{\prod_{j=1}^{m-1} (m + jw)}{(m-1)!} \beta^{m-1} X^m.$$

Proof. This can be proved by a residue calculation. Start from the formula for $X(z)$ with $\beta = 1$ and write $z(X) = \sum_{m=1}^{\infty} C_m X^m$. Then $C_m = \text{Res}_{X=0} z X^{-m} \frac{dX}{X}$, and

$$\begin{aligned}
 \frac{dX}{X} &= \frac{dz}{z} + \frac{d(1+z)^{\frac{1}{w}}}{(1+z)^{\frac{1}{w}}} + \frac{d(1+(w+1)z)^{-\frac{w+1}{w}}}{(1+(w+1)z)^{-\frac{w+1}{w}}} \\
 &= \frac{dz}{z} + \frac{1}{w} \frac{dz}{1+z} - \frac{(w+1)^2}{w} \frac{dz}{1+(w+1)z} \\
 &= \frac{dz}{z(1+z)(1+(w+1)z)}.
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 C_m &= \text{Res}_{X=0} z X^{-m} \frac{dz}{z(1+z)(1+(w+1)z)} \\
 &= \text{Res}_{z=0} z^{-m} (1+z)^{-\frac{m}{w}-1} (1+(w+1)z)^{m\frac{w+1}{w}-1} dz \\
 &= \text{Res}_{z=0} z^{-m} \sum_{k=0}^{\infty} \frac{\prod_{i=0}^{k-1} (-\frac{m}{w} - 1 - i)}{k!} z^k \sum_{l=0}^{\infty} \frac{\prod_{j=0}^{l-1} (m\frac{w+1}{w} - 1 - j)}{l!} (w+1)^l z^l dz \\
 &= \text{Res}_{z=0} z^{-m} \sum_{k=0}^{\infty} \frac{\prod_{i=1}^k (\frac{m}{w} + i)}{k!} (-1)^k z^k \sum_{l=0}^{\infty} \frac{\prod_{j=m-l}^{m-1} (\frac{m}{w} + j)}{l!} (w+1)^l z^l dz \\
 &= \sum_{k=0}^{m-1} \frac{\prod_{i=1}^k (\frac{m}{w} + i)}{k!} \frac{\prod_{j=k+1}^{m-1} (\frac{m}{w} + j)}{(m-k-1)!} (-1)^k (w+1)^{m-k-1}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{\prod_{j=1}^{m-1} \left(\frac{m}{w} + j\right)}{(m-1)!} \sum_{k=0}^{m-1} \binom{m-1}{k} (-1)^k (w+1)^{m-k-1} \\
 &= \frac{\prod_{j=1}^{m-1} \left(\frac{m}{w} + j\right)}{(m-1)!} w^{m-1} = \frac{\prod_{j=1}^{m-1} (m+jw)}{(m-1)!}.
 \end{aligned}$$

Finally, β can be introduced in this formula by scaling $z \rightarrow \beta z$, $X \rightarrow \beta X$. \square

Corollary 3.10. *The expressions for $X(z)$ in lemma 3.9 and lemma 2.23 are related by a Möbius transformation*

$$(57) \quad z \mapsto \frac{z}{1 + (w+1)\beta z}.$$

Hence, by corollary 3.6, they require the same correction term for their induced linear change of variables.

We see that in this particular case we may obtain the function X in two different ways: from the general theory of theorem 2.6, or from the specific shape of the Mariño-Vafa formula, theorem 2.20. In fact, the second choice is nothing but choosing the spectral curve coordinate z to equal ξ (which is unique in this case), or in other words $T_0 = q_1$.

Remark 3.11. Under the correspondence of theorem 2.11, the rank of the cohomological field theory corresponds to the number of zeroes of dx , counted with multiplicities. So for rank one, dx can only have one zero, and hence must have three poles. By Möbius transformation, we may place the zero at infinity, and two of the poles at 0 and -1 , from which we find that dx must correspond to the $\frac{dX}{X}$ found in the proof of lemma 3.9. This may explain in part why Alexandrov [Ale20] finds only the triple Hodge CohFT in the intersection of the orbits of the Givental and Heisenberg-Virasoro groups. However, dx is not the only datum of a spectral curve, and while \mathbb{P}^1 is rigid and has a unique B , it is not clear why there is no freedom in the choice of dy .

Lemma 3.12. *The series $X(z)$ from lemma 3.9 satisfies the differential equation*

$$(58) \quad \frac{\partial X}{\partial \beta}(z) = -((w+2)z + (w+1)\beta z^2)z \frac{\partial X}{\partial z}(z).$$

Proof. For $X(z) = \frac{z}{1+(w+1)z} \left(\frac{1+z}{1+(w+1)z}\right)^{\frac{1}{w}}$, we get $Q(z)^{-1} = (1+z)(1+(w+1)z)$, which using lemma 3.2 immediately yields the result. \square

We use this lemma in combination with the linear correspondence of definition 3.1, slightly adapted as follows: define a linear correspondence Θ between power series in X or z on the one hand and linear series in p or \tilde{q} on the other by

$$(59) \quad p_k \leftrightarrow X^k, \quad \tilde{q}_m \leftrightarrow z^m.$$

Definition 3.13. We define a linear morphism between power series in $\{p_m\}_{m \geq 1}$ and $\{\tilde{q}_d\}_{d \geq 1}$ by

$$(60) \quad p_k(\tilde{q}) = \sum_{m=k}^{\infty} c_k^m \tilde{q}_m \quad \text{with } c_k^m \text{ given by} \quad X^k = \sum_{m=k}^{\infty} c_k^m z^m.$$

Under the correspondence $p_k \leftrightarrow X^k$, $\tilde{q}_m \leftrightarrow z^m$, we have

$$(61) \quad T_d(p) \leftrightarrow (u^2 D)^d u^4 z; \quad D := X \frac{\partial}{\partial X} = (1 + \beta z)(1 + (w+1)\beta z)z \frac{\partial}{\partial z}.$$

In terms of \tilde{q} -variables, this gives

$$(62) \quad T_d = u^2 \sum_{m=1}^{\infty} m(\tilde{q}_m + (w+2)\beta \tilde{q}_{m+1} + (w+1)\beta^2 \tilde{q}_{m+2}) \frac{\partial}{\partial \tilde{q}_m} T_{d-1}; \quad T_0 = u^4 \tilde{q}_1.$$

If we write $q_m := u^{4m} \tilde{q}_m$, and using $\beta = \frac{u^3}{\sqrt{w+1}}$, this becomes

$$(63) \quad T_d = \sum_{m=1}^{\infty} m \left(u^{4m+2} \tilde{q}_m + (w+2) \frac{u^3}{\sqrt{w+1}} u^{4m+2} \tilde{q}_{m+1} + u^6 u^{4m+2} \tilde{q}_{m+2} \right) \frac{1}{u^{4m}} \frac{\partial}{\partial \tilde{q}_m} T_{d-1}$$

$$(64) \quad = \sum_{m=1}^{\infty} m \left(u^2 q_m + \frac{u(w+2)}{\sqrt{w+1}} q_{m+1} + q_{m+2} \right) \frac{\partial}{\partial q_m} T_{d-1};$$

$$(65) \quad T_0 = q_1.$$

This is exactly the definition given in theorem 2.17.

Corollary 3.14. For $X(z) = \frac{z}{1+(w+1)\beta z} \left(\frac{1+\beta z}{1+(w+1)\beta z} \right)^{\frac{1}{w}}$, the quadratic correction of theorem 3.3 is $Q = -\frac{1}{2}\Theta(H_{0,2})$.

Proof. The function $X(z)$ satisfies the conditions of theorem 2.6, so we may apply proposition 3.4. \square

Now we are ready to prove the main result on KP integrability of triple Hodge integrals.

Proof of theorem 2.17. By lemma 2.23, F is a solution of the KP hierarchy in the variables $t_k := \frac{pk}{k}$. In the same way as for proposition 3.8, now using corollary 3.14 and equation (54),

$$(66) \quad F_{\text{TH}} \left(-u, -wu, \frac{wu}{w+1}; \{T_d(p(\tilde{q}))\} \right)$$

is a solution of the KP hierarchy in the variables $\frac{\tilde{q}_m}{m}$. As the KP hierarchy is quasi-homogeneous, rescaling $\tilde{q}_m \rightarrow q_m$ preserves solutions. This completes the proof. \square

Remark 3.15. The result in this subsection do hold for $w = -1$ (ignoring powers of u), but in this specific case $X(z)$ is a Möbius transformation, so it reduces to the setting of corollary 3.6. From another point of view, in this case the change of coordinates equation (37) is an isomorphism, whereas it gives a half-dimensional subspace in all other cases. Equations for this half-dimensional space, in the linear Hodge case, were found in [Ale15], cf. also [GW17] for a reformulation. These can be viewed as a deformation of the reduction from KP to KdV. Similar equations should exist for triple Hodge integrals as well, but clearly none of this works for $w = -1$.

In light of section 3.3, one may expect a deformation of the reduction from KP to r -KdV or r -Gelfand-Dickey for the families found there.

REFERENCES

- [Ale15] Alexander Alexandrov. “Enumerative Geometry, Tau-Functions and Heisenberg-Virasoro Algebra.” In: *Comm. Math. Phys.* 338.1 (2015), pp. 195–249. ISSN: 0010-3616. DOI: [10.1007/s00220-015-2379-8](https://doi.org/10.1007/s00220-015-2379-8). arXiv: [1404.3402](https://arxiv.org/abs/1404.3402).
- [Ale20] Alexander Alexandrov. “KP integrability of triple Hodge integrals. I. From Givental group to hierarchy symmetries.” In: (2020). arXiv: [2009.01615](https://arxiv.org/abs/2009.01615).
- [Ale21] Alexander Alexandrov. “KP integrability of triple Hodge integrals. II. Generalized Kontsevich matrix model.” In: *Anal. Math. Phys.* 11.1 (2021), Paper No. 24, 82. ISSN: 1664-2368. DOI: [10.1007/s13324-020-00451-7](https://doi.org/10.1007/s13324-020-00451-7). arXiv: [2009.10961](https://arxiv.org/abs/2009.10961).
- [AS21] Alexander Alexandrov and Sergey Shadrin. “Elements of spin Hurwitz theory: closed algebraic formulas, blobbed topological recursion, and a proof of the Giacchetto-Kramer-Lewanski conjecture.” In: (2021). arXiv: [2105.12493](https://arxiv.org/abs/2105.12493).
- [BCEG21] Raphaël Belliard, Séverin Charbonnier, Bertrand Eynard, and Elba Garcia-Failde. “Topological recursion for generalised Kontsevich graphs and r -spin intersection numbers.” In: (2021). arXiv: [2105.08035](https://arxiv.org/abs/2105.08035).
- [BDKS20a] Boris Bychkov, Petr Dunin-Barkowski, Maxim Kazarian, and Sergey Shadrin. “Explicit closed algebraic formulas for Orlov-Scherbin n -point functions.” In: (2020). arXiv: [2008.13123](https://arxiv.org/abs/2008.13123).
- [BDKS20b] Boris Bychkov, Petr Dunin-Barkowski, Maxim Kazarian, and Sergey Shadrin. “Topological recursion for Kadomtsev-Petviashvili tau functions of hypergeometric type.” In: (2020). arXiv: [2012.14723](https://arxiv.org/abs/2012.14723).
- [BE13] Vincent Bouchard and Bertrand Eynard. “Think globally, compute locally.” In: *Journal of High Energy Physics* 2013.2 (2013), pp. 1–34. DOI: [10.1007/JHEP02\(2013\)143](https://doi.org/10.1007/JHEP02(2013)143). arXiv: [1211.2302](https://arxiv.org/abs/1211.2302).
- [BEMS11] Gaëtan Borot, Bertrand Eynard, Motohico Mulase, and Brad Safnuk. “A matrix model for simple Hurwitz numbers, and topological recursion.” In: *Journal of Geometry and Physics* 61.2 (2011), pp. 522–540. DOI: [10.1016/j.geomphys.2010.10.017](https://doi.org/10.1016/j.geomphys.2010.10.017). arXiv: [0906.1206](https://arxiv.org/abs/0906.1206).
- [BEO15] Gaëtan Borot, Bertrand Eynard, and Nicolas Orantin. “Abstract loop equations, topological recursion, and applications.” In: *Commun. Number Theory Phys.* 9.1 (2015), pp. 1–89. DOI: [10.4310/CNTP.2015.v9.n1.a2](https://doi.org/10.4310/CNTP.2015.v9.n1.a2). arXiv: [1303.5808](https://arxiv.org/abs/1303.5808).
- [BKMP09] Vincent Bouchard, Albrecht Klemm, Marcos Mariño, and Sara Pasquetti. “Remodeling the B-model.” In: *Comm. Math. Phys.* 287.1 (2009), pp. 117–178. ISSN: 0010-3616. DOI: [10.1007/s00220-008-0620-4](https://doi.org/10.1007/s00220-008-0620-4).
- [BKS20] Gaëtan Borot, Reinier Kramer, and Yannik Schüler. “Higher Airy structures and topological recursion for singular spectral curves.” In: (2020). arXiv: [2010.03512](https://arxiv.org/abs/2010.03512).

- [BMo8] Vincent Bouchard and Marcos Mariño. “Hurwitz numbers, matrix models and enumerative geometry.” In: *From Hodge theory to integrability and TQFT tt^* -geometry*. Vol. 78. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 2008, pp. 263–283. DOI: [10.1090/pspum/078/2483754](https://doi.org/10.1090/pspum/078/2483754). arXiv: [0709.1458](https://arxiv.org/abs/0709.1458).
- [BS17] Gaëtan Borot and Sergey Shadrin. “Blobbed topological recursion: properties and applications.” In: *Math. Proc. Camb. Phil. Soc.* 162 (2017), pp. 39–87. DOI: [10.1017/S0305004116000323](https://doi.org/10.1017/S0305004116000323). arXiv: [1502.00981](https://arxiv.org/abs/1502.00981).
- [Bur15a] Alexandr Buryak. “Double ramification cycles and integrable hierarchies.” In: *Comm. Math. Phys.* 336.3 (2015), pp. 1085–1107. DOI: [10.1007/s00220-014-2235-2](https://doi.org/10.1007/s00220-014-2235-2). arXiv: [1403.1719](https://arxiv.org/abs/1403.1719).
- [Bur15b] Alexandr Buryak. “Dubrovin-Zhang hierarchy for the Hodge integrals.” In: *Commun. Number Theory Phys.* 9.2 (2015), pp. 239–272. DOI: [10.4310/CNTP.2015.v9.n2.a1](https://doi.org/10.4310/CNTP.2015.v9.n2.a1). arXiv: [1308.5716](https://arxiv.org/abs/1308.5716).
- [Bur16] Alexandr Buryak. “ILW equation for the Hodge integrals revisited.” In: *Math. Res. Lett.* 23.3 (2016), pp. 675–683. DOI: [10.4310/MRL.2016.v23.n3.a5](https://doi.org/10.4310/MRL.2016.v23.n3.a5). arXiv: [1505.05911](https://arxiv.org/abs/1505.05911).
- [CEO06] Leonid Chekhov, Bertrand Eynard, and Nicolas Orantin. “Free energy topological expansion for the 2-matrix model.” In: *J. High Energy Phys.* 2006.12 (2006), pp. 1–31. DOI: [10.1088/1126-6708/2006/12/053](https://doi.org/10.1088/1126-6708/2006/12/053). arXiv: [math-ph/0603003](https://arxiv.org/abs/math-ph/0603003).
- [Che18] Lin Chen. “Bouchard-Klemm-Marino-Pasquetti conjecture for \mathbb{C}^3 .” In: *Topological recursion and its influence in analysis, geometry, and topology*. Vol. 100. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 2018, pp. 83–102.
- [Dic03] Leonid A. Dickey. *Soliton equations and Hamiltonian systems*. Second. Vol. 26. Advanced Series in Mathematical Physics. World Scientific Publishing Co., Inc., River Edge, NJ, 2003, pp. xii+408. ISBN: 981-238-173-2. DOI: [10.1142/5108](https://doi.org/10.1142/5108).
- [DNOPS16] Petr Dunin-Barkowski, Paul Norbury, Nicolas Orantin, Alexandr Popolitov, and Sergey Shadrin. “Primary invariants of Hurwitz Frobenius manifolds.” In: (2016), pp. 1–24. arXiv: [1605.07644](https://arxiv.org/abs/1605.07644).
- [DOSS14] Petr Dunin-Barkowski, Nicolas Orantin, Sergey Shadrin, and Loek Spitz. “Identification of the Givental formula with the spectral curve topological recursion procedure.” In: *Comm. Math. Phys.* 328.2 (2014), pp. 669–700. DOI: [10.1007/s00220-014-1887-2](https://doi.org/10.1007/s00220-014-1887-2). arXiv: [1211.4021](https://arxiv.org/abs/1211.4021).
- [DZ01] Boris Dubrovin and Youjin Zhang. “Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov–Witten invariants.” In: (Aug. 2001). arXiv: [math/0108160](https://arxiv.org/abs/math/0108160).
- [ELSV01] Torsten Ekedahl, Sergei Lando, Michael Shapiro, and Alek Vainshtein. “Hurwitz numbers and intersections on moduli spaces of curves.” In: *Invent. Math.* 146.2 (2001), pp. 297–327. DOI: [10.1007/s002220100164](https://doi.org/10.1007/s002220100164).
- [EO07] Bertrand Eynard and Nicolas Orantin. “Invariants of algebraic curves and topological expansion.” In: *Commun. Number Theory Phys.* 1.2 (2007), pp. 347–452. DOI: [10.4310/CNTP.2007.v1.n2.a4](https://doi.org/10.4310/CNTP.2007.v1.n2.a4). arXiv: [math-ph/0702045](https://arxiv.org/abs/math-ph/0702045).
- [EO15] Bertrand Eynard and Nicolas Orantin. “Computation of open Gromov–Witten invariants for toric Calabi–Yau 3-folds by topological recursion, a proof of the BKMP conjecture.” In: *Commun. Math. Phys.* 337.2 (2015), pp. 483–567. DOI: [10.1007/s00220-015-2361-5](https://doi.org/10.1007/s00220-015-2361-5). arXiv: [1205.1103](https://arxiv.org/abs/1205.1103).
- [Eyn11] Bertrand Eynard. “Intersection numbers of spectral curves.” In: (Apr. 2011). arXiv: [1104.0176](https://arxiv.org/abs/1104.0176).
- [Eyn14] Bertrand Eynard. “Invariants of spectral curves and intersection theory of moduli spaces of complex curves.” In: *Communications in Number Theory and Physics* 8.3 (2014), pp. 541–588. DOI: [10.4310/CNTP.2014.v8.n3.a4](https://doi.org/10.4310/CNTP.2014.v8.n3.a4). arXiv: [1110.2949](https://arxiv.org/abs/1110.2949).
- [FP00] Carel Faber and Rahul Pandharipande. “Hodge integrals and Gromov–Witten theory.” In: *Invent. Math.* 139 (2000), pp. 173–199. DOI: [10.1007/s002229900028](https://doi.org/10.1007/s002229900028). arXiv: [math/9810173](https://arxiv.org/abs/math/9810173).
- [Giv01] Alexander B. Givental. “Gromov–Witten invariants and quantization of quadratic hamiltonians.” In: *Mosc. Math. J.* 1.4 (2001), pp. 1–17. DOI: [10.17323/1609-4514-2001-1-4-551-568](https://doi.org/10.17323/1609-4514-2001-1-4-551-568). arXiv: [math/0108100](https://arxiv.org/abs/math/0108100).
- [GKL21] Alessandro Giacchetto, Reinier Kramer, and Danilo Lewański. “A new spin on Hurwitz theory and ELSV via theta characteristics.” In: (2021). arXiv: [2104.05697](https://arxiv.org/abs/2104.05697).

- [GW17] Shuai Guo and Gehao Wang. “Virasoro constraints and polynomial recursion for the linear Hodge integrals.” In: *Lett. Math. Phys.* 107.4 (2017), pp. 757–791. DOI: [10.1007/s11005-016-0923-x](https://doi.org/10.1007/s11005-016-0923-x). arXiv: [1608.02077](https://arxiv.org/abs/1608.02077).
- [Kaz09] Maxim Kazarian. “KP hierarchy for Hodge integrals.” In: *Adv. Math.* 221.1 (2009), pp. 1–21. DOI: [10.1016/j.aim.2008.10.017](https://doi.org/10.1016/j.aim.2008.10.017). arXiv: [0809.3263](https://arxiv.org/abs/0809.3263).
- [Kha99] Sergey Kharchev. “Kadomtsev–Petviashvili hierarchy and generalized Kontsevich model.” In: *Moscow Seminar in Mathematical Physics*. Vol. 191. Amer. Math. Soc. Transl. Ser. 2. Amer. Math. Soc., Providence, RI, 1999, pp. 119–162. DOI: [10.1090/trans2/191/05](https://doi.org/10.1090/trans2/191/05). arXiv: [hep-th/9810091](https://arxiv.org/abs/hep-th/9810091).
- [KM94] Maxim Kontsevich and Yuri I. Manin. “Gromov–Witten classes, quantum cohomology, and enumerative geometry.” In: *Communications in Mathematical Physics* 164.3 (1994), pp. 525–562. DOI: [10/dmfxd6](https://doi.org/10/dmfxd6). arXiv: [hep-th/9402147](https://arxiv.org/abs/hep-th/9402147).
- [KMMM95] Sergey M. Kharchev, Andrei V. Marshakov, Andrei D. Mironov, and Alexei Yu. Morozov. “Generalized Kazakov–Migdal–Kontsevich model: group theory aspects.” In: *Internat. J. Modern Phys. A* 10.14 (1995), pp. 2015–2051. ISSN: 0217-751X. DOI: [10.1142/S0217751X9500098X](https://doi.org/10.1142/S0217751X9500098X).
- [Kon92] Maxim Kontsevich. “Intersection theory on the moduli space of curves and the matrix Airy function.” In: *Comm. Math. Phys.* 147.1 (1992), pp. 1–23. DOI: [10.1007/BF02099526](https://doi.org/10.1007/BF02099526).
- [Kra19] Reinier Kramer. “Cycles of curves, cover counts, and central invariants.” Ph.D. thesis. Universiteit van Amsterdam, 2019. URL: <https://hdl.handle.net/11245.1/588b7706-082d-4836-9cf7-7ca5025f3d64>.
- [LLZ03] Chiu-Chu Melissa Liu, Kefeng Liu, and Jian Zhou. “A Proof of a Conjecture of Mariño–Vafa on Hodge Integrals.” In: *J. Differential Geometry* 65 (2003), pp. 289–340. DOI: [10.4310/jdg/1090511689](https://doi.org/10.4310/jdg/1090511689). eprint: [math/0306434](https://arxiv.org/abs/math/0306434).
- [LYZZ21] Si-Qi Liu, Di Yang, Youjin Zhang, and Chunhui Zhou. “The Hodge–FVH correspondence.” In: *J. Reine Angew. Math.* 775 (2021), pp. 259–300. ISSN: 0075-4102. DOI: [10.1515/crelle-2020-0051](https://doi.org/10.1515/crelle-2020-0051).
- [MJD00] Tetsuji Miwa, Michio Jimbo, and Etsuro Date. *Solitons: Differential equations, symmetries and infinite dimensional algebras*. Vol. 135. Cambridge Tracts in Mathematics. Cambridge University Press, 2000.
- [MV02] Marcos Mariño and Cumrun Vafa. “Framed knots at large N .” In: *Contemporary Mathematics* 310 (2002), pp. 185–205. DOI: [10.1090/conm/310/05404](https://doi.org/10.1090/conm/310/05404).
- [Ok000] Andrei Okounkov. “Toda equations for Hurwitz numbers.” In: *Math. Res. Lett.* 7 (2000), pp. 447–453. DOI: [10.4310/MRL.2000.v7.n4.a10](https://doi.org/10.4310/MRL.2000.v7.n4.a10).
- [OP04] Andrei Okounkov and Rahul Pandharipande. “Hodge integrals and invariants of the unknot.” In: *Geometry and Topology* 8 (2004), pp. 675–699. DOI: [10.2140/gt.2004.8.675](https://doi.org/10.2140/gt.2004.8.675). arXiv: [math/0307209](https://arxiv.org/abs/math/0307209).
- [OS01a] Alexandr Yu. Orlov and D. M. Scherbin. “Multivariate hypergeometric functions as τ -functions of Toda lattice and Kadomtsev–Petviashvili equation.” In: vol. 152/153. *Advances in nonlinear mathematics and science*. 2001, pp. 51–65. DOI: [10.1016/S0167-2789\(01\)00158-0](https://doi.org/10.1016/S0167-2789(01)00158-0).
- [OS01b] Alexandr. Yu. Orlov and D. M. Shcherbin. “Hypergeometric solutions of soliton equations.” In: *Teoret. Mat. Fiz.* 128.1 (2001), pp. 84–108. ISSN: 0564-6162. DOI: [10.1023/A:1010402200567](https://doi.org/10.1023/A:1010402200567).
- [Sha09] Sergey Shadrin. “BCOV theory via Givental group action on cohomological fields theories.” In: *Mosc. Math. J.* 9.2 (2009), pp. 411–429. DOI: [10.17323/1609-4514-2009-9-2-411-429](https://doi.org/10.17323/1609-4514-2009-9-2-411-429). arXiv: [0810.0725](https://arxiv.org/abs/0810.0725).
- [SS83] Mikio Sato and Yasuko Sato. “Soliton equations as dynamical systems on infinite-dimensional Grassmann manifold.” In: *North-Holland Math. Stud.* U.S.–Japan seminar on Nonlinear partial differential equations in applied science (Tokyo, July 1982). Vol. 81. Lecture Notes Numer. Appl. Anal. 5. 1983, pp. 259–271.
- [Tel12] Constantin Teleman. “The structure of 2D semi-simple field theories.” In: *Invent. Math.* 188.3 (2012), pp. 525–588. DOI: [10.1007/s00222-011-0352-5](https://doi.org/10.1007/s00222-011-0352-5). arXiv: [0712.0160](https://arxiv.org/abs/0712.0160).
- [TT95] Kanehisa Takasaki and Takashi Takebe. “Integrable hierarchies and dispersionless limit.” In: *Rev. Math. Phys.* 7.5 (1995), pp. 743–808. ISSN: 0129-055X. DOI: [10.1142/S0129055X9500030X](https://doi.org/10.1142/S0129055X9500030X).
- [Wit91] Edward Witten. “Two-dimensional gravity and intersection theory on moduli space.” In: *Surveys in differential geometry*. Conference on geometry and topology (Harvard University,

- Cambridge, MA, USA, Apr. 27–29, 1990). Vol. 1. Lehigh University, Bethlehem, PA, 1991, pp. 243–310.
- [Wit93] Edward Witten. “Algebraic geometry associated with matrix models of two-dimensional gravity.” In: *Topological methods in modern mathematics (Stony Brook, NY, 1991)*. Publish or Perish, Houston, TX, 1993, pp. 235–269.
- [Zho03] Jian Zhou. “Hodge integrals, Hurwitz numbers, and Symmetric Groups.” In: (2003), p. 24. arXiv: [math/0308024](https://arxiv.org/abs/math/0308024).
- [Zho09] Jian Zhou. “Local mirror symmetry for the topological vertex.” In: (2009). arXiv: [0911.2343](https://arxiv.org/abs/0911.2343).
- [Zho10] Jian Zhou. “Hodge integrals and integrable hierarchies.” In: *Lett Math Phys* 93 (2010), pp. 55–71. DOI: [10.1007/s11005-010-0397-1](https://doi.org/10.1007/s11005-010-0397-1). arXiv: [math/0310408](https://arxiv.org/abs/math/0310408).

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111 BONN, GERMANY.
Email address: rkramer@mpim-bonn.mpg.de