


Max-Planck-Institut für Mathematik Bonn

The second shifted difference of partitions and its applications

by

Kevin Gomez
Joshua Males
Larry Rolen



Max-Planck-Institut für Mathematik
Preprint Series 2022 (19)

Date of submission: March 22, 2022

The second shifted difference of partitions and its applications

by

Kevin Gomez
Joshua Males
Larry Rolen

Max-Planck-Institut für Mathematik
Vivatsgasse 7
53111 Bonn
Germany

Department of Mathematics
Vanderbilt University
1420 Stevenson Center
Nashville, TN 37240
USA

Department of Mathematics
University of Manitoba
450 Machray Hall
Winnipeg
Canada

THE SECOND SHIFTED DIFFERENCE OF PARTITIONS AND ITS APPLICATIONS

KEVIN GOMEZ, JOSHUA MALES, AND LARRY ROLEN

ABSTRACT. A number of recent papers have estimated ratios of the partition function $p(n-j)/p(n)$, which appears in many applications. Here, we prove an easy-to-use effective bound on these ratios. Using this, we then study second shifted difference of partitions, $f(j, n) := p(n) - 2p(n-j) + p(n-2j)$, and give another easy-to-use estimate of $f(j, n)$. As applications of these, we prove a shifted convexity property of $p(n)$, as well as giving new estimates of the k -rank partition function $N_k(m, n)$ and non- k -ary partitions along with their differences.

1. INTRODUCTION AND STATEMENT OF RESULTS

The study of the values of the partition function $p(n)$, which counts the number of partitions of a positive integer n , has a long history. Recall that a partition λ of n is a non-increasing list $(\lambda_1, \lambda_2, \dots, \lambda_s)$ such that $\sum_{j=1}^s \lambda_j = n$. In their famed collaboration a century ago, Hardy and Ramanujan [7] proved that as $n \rightarrow \infty$ we have the asymptotic formula

$$p(n) \sim \frac{1}{4\sqrt{3}n} e^{\pi\sqrt{\frac{2n}{3}}}. \quad (1.1)$$

Their proof gave birth to the Circle Method, which is a highly important tool used throughout analytic number theory today. Following their discovery, Rademacher [13] improved Hardy and Ramanujan's application of the Circle Method to prove an exact formula for $p(n)$. Over the past 100 years, there have been a plethora of investigations into estimates and asymptotics for partitions and their extensions in the literature.

Here, we will study differences of partition values in detail. To this end, let Δ be the backward difference operator defined on sequences $f(n)$ by

$$\Delta(f(n)) := f(n) - f(n-1),$$

and its recursive counterpart

$$\Delta^k(f(n)) := \Delta\left(\Delta^{k-1}(f(n))\right).$$

One of the simplest properties of $p(n)$ to prove is that it is convex for $n \geq 2$ (see e.g. [6]), i.e.

$$p(n) + p(n-2) \geq 2p(n-1).$$

1991 *Mathematics Subject Classification.* 11P82.

Key words and phrases. Partitions, convexity, asymptotic expansions.

The research of the second author conducted for this paper is supported by the Pacific Institute for the Mathematical Sciences (PIMS). The research and findings may not reflect those of the Institute. The third author is grateful for support from a grant from the Simons Foundation (853830, LR), support from a Dean's Faculty Fellowship from Vanderbilt University, and to the Max Planck Institute for Mathematics in Bonn for its hospitality and financial support.

Recast using the operator Δ this is the same as proving that

$$\Delta^2(p(n)) \geq 0$$

for all $n \geq 2$. Gupta [6] also investigated higher powers of Δ applied to p , proving that there exist constants n_r for all $r > 0$ such that $\Delta^r(p(n)) \geq 0$ for all $n \geq n_r$.

We initiate the investigation of what we call j -shifted differences, defined for $1 \leq j < n$ on sequences $f(n)$ by

$$\Delta_j(f(n)) := f(n) - f(n - j).$$

In analogy to Gupta, it is clear using (1.1) that there exist constants n_j such that for all $n \geq n_j$ one has that $\Delta_j^2(p(n)) \geq 0$. Let $N := n - \frac{1}{24}$. Our methods rely on a careful study of the value of the function

$$f(j, n) := \Delta_j^2(p(n)) = p(n) - 2p(n - j) + p(n - 2j),$$

and in Theorem 2.2 we prove a precise estimation of $f(j, n)$ with $j \leq \frac{\sqrt{N}}{4}$, in particular providing a strict error term allowing us to closely control the precision of the formula by taking N large enough. In doing so, we provide an easy-to-use estimate for the ratio of partition numbers. Throughout, we use the notation $f(x) = O_{\leq}(g(x))$ to mean that $|f(x)| \leq g(x)$ for x in the appropriate domain.

Theorem 1.1. *Let $n \geq 14$ and $j < \frac{\sqrt{N}}{2}$. Then*

$$\frac{p(n-j)}{p(n)} = e^{\frac{\pi j}{\sqrt{6N}}} \left(1 + \frac{j}{N} - \frac{\pi j^2}{4\sqrt{6N}} - \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N}} + O_{\leq} \left(\frac{2.71}{N} \right) \right) \left(1 + \frac{\sqrt{3}}{\sqrt{2N}\pi} + O_{\leq} \left(\frac{1350}{N} \right) \right).$$

Theorems of a similar flavour to Theorem 1.1 are abundant in the literature. Lehmer [10, 11] used Rademacher's exact formula for $p(n)$ [13] to provide bounds on the value of $p(n)$ that have seen many applications. More recently, estimates for the ratio of partition values have played a prominent role in proving that the associated Jensen polynomial is eventually hyperbolic [5, 9], a problem intricately linked with variants of the Riemann hypothesis.

Theorem 1.1 thus applies to many interesting situations. In the remainder of the introduction, we will highlight a few of particular interest. Our first main result using Theorem 2.2 gives an explicit formula for n_j for ranges of j .

Theorem 1.2. *Let $n \geq 3922$ and $j \leq \frac{\sqrt{N}}{4}$. Then we have that*

$$\Delta_j^2(p(n)) \geq 0.$$

Equivalently, $p(n)$ satisfies the extended convexity result

$$p(n) + p(n - 2j) \geq 2p(n - j).$$

Remark. The methods here should extend to finding formula for $n_{r,j}$ such that for all $n \geq n_{r,j}$ one has $\Delta_j^r(p(n)) \geq 0$, however this would quickly become very lengthy and so we do not pursue this here.

Our results also apply outside of proving new properties of the partition function itself. We consider the k -rank function $N_k(m, n)$ which counts the number of partitions of n into at least $(k - 1)$ successive Durfee squares with k -rank equal to m [4]. When $k = 1$ we recover the number

of partitions of n whose Andrews–Garvan crank equals m , and when $k = 2$ we recover Dyson’s partition rank function. Then for $m > \frac{n}{2}$ we have that (see e.g. [12, page 6])

$$N_k(m, n) = p(n - k - m + 1) - p(n - k - m), \quad N_k(m, n) - N_k(m + 1, n) = f(1, n - k - m).$$

We give precise formulae both $N_k(m, n)$ and for the differences of k -ranks in certain ranges of m in the following theorems, improving on [12, Theorem 1.4] in this range. The proof follows from a direct application of Theorem 1.1.

Theorem 1.3. *Let $m > \frac{n}{2}$ and $\ell := n - k - m + \frac{23}{24} > 16$. Then we have that*

$$\frac{N_k(m, n)}{p(n - k - m + 1)} = 1 - e^{-\frac{\pi}{\sqrt{6\ell}}} \left(1 - \frac{\sqrt{3}}{\sqrt{2\pi\ell}} + O_{\leq} \left(\frac{4.04}{\ell} \right) \right) \left(1 + \frac{\sqrt{3}}{\sqrt{2\ell}\pi} + O_{\leq} \left(\frac{1350}{\ell} \right) \right).$$

We then turn to obtaining a precise estimate for the differences of k -ranks, with the proof following from a direct application of Theorem 2.2.

Theorem 1.4. *Let $m > \frac{n}{2}$ and $\ell = n - k - m + \frac{23}{24} > 16$. Then we have that*

$$\begin{aligned} \frac{N_k(m, n) - N_k(m + 1, n)}{p(n - k - m + 1)} &= 1 + e^{\frac{\sqrt{2}\pi}{\sqrt{3\ell}}} \left(1 + \left(\frac{\sqrt{3}}{\sqrt{2\pi}} - \frac{\sqrt{3}}{\sqrt{2\pi}} \right) \frac{1}{\sqrt{\ell}} + O_{\leq} \left(\frac{2079}{\ell} \right) \right) \\ &\quad - e^{-\frac{\pi}{\sqrt{6\ell}}} \left(2 + \left(\frac{2\sqrt{3}}{\sqrt{2\pi}} - \frac{2\sqrt{3}}{\sqrt{2\pi}} \right) \frac{1}{\sqrt{\ell}} + O_{\leq} \left(\frac{3929}{\ell} \right) \right). \end{aligned}$$

As a direct implication, we recover positivity of the differences of k -ranks in these cases, as in [12, Corollary 1.5].

Our final application is to so-called non- k -ary partitions ($k \in \mathbb{N}$), recently defined by Schneider [14] as partitions of n with no parts equal to k ¹. Letting $\nu_k(n)$ be the number of non- k -ary partitions of n , it is clear that $\nu_k(n) = p(n) - p(n - k)$. By Theorem 1.1 we immediately obtain an effective estimate for the ratio $\nu_k(n)/p(n)$, improving on [1, Theorem 1]. We also have

$$\nu_k(n) - \nu_k(n - k) = f(k, n),$$

and so we also obtain precise estimates for differences of non- k -ary partitions using Theorem 2.2 for $k < \frac{\sqrt{N}}{4}$, with a direct implication being the following theorem.

Theorem 1.5. *For $n \geq 3922$ and $k \leq \frac{\sqrt{N}}{4}$ we have*

$$\nu_k(n) - \nu_k(n - k) > 0.$$

ACKNOWLEDGMENTS

The authors are very grateful to Kathrin Bringmann who shared many calculations and insightful comments invaluable for this paper.

¹While [14] uses the terminology “ k -nuclear”, Schneider has recommended the authors use the term non- k -ary based on advice of G. Andrews to better fit the case of $k = 1$, classically called the non-unitary partitions.

2. THE PROOFS

In this section we prove the main results of the paper. We begin by proving a technical estimate of the value of $p(n-j)$, utilising Rademacher's exact formula for the partition function.

Proposition 2.1. *Let $N := n - \frac{1}{24}$ and $j \in \mathbb{N}_0$. Then*

$$p(n-j) = \frac{e^{\pi\sqrt{\frac{2(N-j)}{3}}}}{4\sqrt{3}(N-j)} \times \left(1 - \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N-j}} + O_{\leq} \left(\frac{2\pi^2(N-j)e^{-\pi\sqrt{\frac{2(N-j)}{3}}}}{3} + 2^3 3^{-\frac{1}{2}} \pi(N-j)^{\frac{1}{2}} e^{-\frac{\pi}{2}\sqrt{\frac{N-j}{2}}} \right) \right).$$

Proof. We first recall the following result from [8, Theorem 1.1] with $\alpha = 1$, which is simply Rademacher's exact formula for the partition function [13],

$$p(n) = \frac{\pi}{2^{\frac{5}{4}} 3^{\frac{3}{4}} N^{\frac{3}{4}}} \sum_{k=1}^{\infty} \frac{A_k(n)}{k} I_{\frac{3}{2}} \left(\frac{\pi}{k} \sqrt{\frac{2N}{3}} \right), \quad (2.1)$$

where I_{ν} is the usual I -Bessel function and

$$A_k(n) := \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} e^{\pi i s(h,k) - \frac{2\pi i n h}{k}}$$

is a Kloosterman sum with $s(h,k)$ the usual Dedekind sum. By page 172 of [15], we have the following representation of the I -Bessel function

$$I_{\frac{3}{2}}(x) = \frac{x^{\frac{3}{2}}}{2\sqrt{2\pi}} \int_{-1}^1 (1-t^2) e^{xt} dt. \quad (2.2)$$

We now bound the integrand for $-1 \leq t \leq 0$ by 1 and find that

$$\int_{-1}^0 (1-t^2) e^{xt} dt = O_{\leq}(1). \quad (2.3)$$

Here, the notation $f(x) = O_{\leq}(g(x))$ means that $|f(x)| \leq g(x)$, that is, that there is no implied constant in the big-Oh estimate.

Next we compute the integral for $0 \leq t \leq 1$. To do so, we make the change of variables $u = 1-t$ to find that

$$\int_0^1 (1-t^2) e^{xt} dt = e^x \left(2 \int_0^1 u^2 e^{-xu} \frac{du}{u} - \int_0^1 u^3 e^{-xu} \frac{du}{u} \right).$$

Under the change of variables $w = ux$, it is easy to show that this is equal to

$$\frac{2e^x}{x^2} \left(1 - \frac{1}{x} - \Gamma(2, x) + \frac{\Gamma(3, x)}{2x} \right),$$

where $\Gamma(a, b)$ is the usual incomplete Γ -function. Since $\Gamma(2, x) = (x + 1)e^{-x}$ and $\Gamma(3, x) = (x^2 + 2x + 2)e^{-x}$, we find that

$$\int_0^1 (1 - t^2) e^{xt} dt = \frac{2e^x}{x^2} \left(1 - \frac{1}{x} - \frac{x+1}{e^x} + \frac{x^2 + 2x + 2}{2xe^x} \right). \quad (2.4)$$

Plugging (2.3) and (2.4) into (2.2), we obtain

$$\begin{aligned} I_{\frac{3}{2}}(x) &= \frac{x^{\frac{3}{2}}}{2\sqrt{2\pi}} \left(\frac{2e^x}{x^2} \left(1 - \frac{1}{x} - \frac{x+1}{e^x} + \frac{x^2 + 2x + 2}{2xe^x} \right) + O_{\leq}(1) \right) \\ &= \frac{e^x}{\sqrt{2\pi x}} \left(1 - \frac{1}{x} + \left(\frac{1}{x} - \frac{x}{2} \right) e^{-x} + O_{\leq} \left(\frac{x^2}{2e^x} \right) \right). \end{aligned}$$

Noting that for $x \geq 1$, one has $|\frac{1}{x} - \frac{x}{2}| \leq \frac{x^2}{2}$ we have

$$I_{\frac{3}{2}}(x) = \frac{e^x}{\sqrt{2\pi x}} \left(1 - \frac{1}{x} + O_{\leq} \left(x^2 e^{-x} \right) \right)$$

for $x \geq 1$. In particular,

$$I_{\frac{3}{2}} \left(\pi \sqrt{\frac{2N}{3}} \right) = \frac{3^{\frac{1}{4}} e^{\pi \sqrt{\frac{2N}{3}}}}{2^{\frac{3}{4}} \pi N^{\frac{1}{4}}} \left(1 - \frac{\sqrt{3}}{\sqrt{2\pi}} \frac{1}{\sqrt{N}} + O_{\leq} \left(\frac{2\pi^2 N e^{-\pi \sqrt{\frac{2N}{3}}}}{3} \right) \right) \quad (2.5)$$

for $n \geq 1$. This corresponds to the term $k = 1$ in the sum in (2.1), and we need to bound the remaining terms of the sum. Note that $|A_k(n)| \leq k$, so we may bound the remaining terms in the sum by

$$\left| \sum_{k=2}^{\infty} \frac{A_k(n)}{k} I_{\frac{3}{2}} \left(\frac{\pi}{k} \sqrt{\frac{2N}{3}} \right) \right| \leq \sum_{k=2}^{\infty} I_{\frac{3}{2}} \left(\frac{\pi}{k} \sqrt{\frac{2N}{3}} \right).$$

We now emulate (3.20) of [2]. By (3.18) of [2] we have

$$\sum_{2 \leq k \leq \lfloor X \rfloor} I_{\frac{3}{2}} \left(\frac{X}{k} \right) \leq 2\sqrt{\frac{X}{\pi}} e^{\frac{X}{2}}.$$

The remaining terms are

$$\sum_{k \geq \lfloor X \rfloor + 1} I_{\frac{3}{2}} \left(\frac{X}{k} \right) \leq \frac{X^{\frac{3}{2}}}{\Gamma(\frac{5}{2}) \sqrt{2}} \sum_{k \geq \lfloor X \rfloor + 1} \frac{1}{k^{\frac{3}{2}}},$$

where we use Lemma 2.2 (3) of [2]. We thus have the bound

$$\sum_{k \geq 2} I_{\frac{3}{2}} \left(\frac{X}{k} \right) \leq 2\sqrt{\frac{X}{\pi}} e^{\frac{X}{2}} + \frac{2X^{\frac{3}{2}}}{\Gamma(\frac{5}{2}) \sqrt{2}}.$$

It remains to bound the final sum with basic calculus by $2\sqrt{\frac{X}{\pi}}e^{\frac{X}{2}}$, yielding

$$\sum_{k \geq 2} I_{\frac{3}{2}}\left(\frac{X}{k}\right) \leq 4\sqrt{\frac{X}{\pi}}e^{\frac{X}{2}}.$$

In our application, this yields

$$\sum_{k=2}^{\infty} I_{\frac{3}{2}}\left(\frac{\pi}{k}\sqrt{\frac{2N}{3}}\right) \leq 4\sqrt[3]{\frac{2N}{3}}e^{\frac{\pi}{2}\sqrt{\frac{2N}{3}}} = \frac{2^{\frac{9}{4}}N^{\frac{1}{4}}}{3^{\frac{1}{4}}}e^{\frac{\pi}{2}\sqrt{\frac{2N}{3}}}. \quad (2.6)$$

Combining (2.5) and (2.6) in (2.1), we find that

$$\begin{aligned} p(n) &= \frac{\pi}{2^{\frac{5}{4}}3^{\frac{3}{4}}N^{\frac{3}{4}}}\left(\frac{3^{\frac{1}{4}}e^{\pi\sqrt{\frac{2N}{3}}}}{2^{\frac{3}{4}}\pi N^{\frac{1}{4}}}\left(1 - \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} + O_{\leq}\left(\frac{2\pi^2Ne^{-\pi\sqrt{\frac{2N}{3}}}}{3}\right)\right)\right) + O_{\leq}\left(\frac{2^{\frac{9}{4}}N^{\frac{1}{4}}}{3^{\frac{1}{4}}}e^{\frac{\pi}{2}\sqrt{\frac{2N}{3}}}\right) \\ &= \frac{e^{\pi\sqrt{\frac{2N}{3}}}}{4\sqrt{3}N}\left(1 - \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} + O_{\leq}\left(\frac{2\pi^2Ne^{-\pi\sqrt{\frac{2N}{3}}}}{3} + 2^33^{-\frac{1}{2}}\pi N^{\frac{1}{2}}e^{-\frac{\pi}{2}\sqrt{\frac{N}{2}}}\right)\right). \end{aligned}$$

Note that $N = N(n)$ is implicitly a function of n and $N(n-j) = N(n) - j$ and so the claim follows. \square

Next we want to estimate the functions $f(j, n)$. The first step in the proof is to obtain estimates of $\frac{p(n-j)}{p(n)}$ analogous to those of [2], proving Theorem 1.1 en-route.

Theorem 2.2. *Let $j < \frac{\sqrt{N}}{4}$ and $n \geq 14$. Then*

$$\begin{aligned} \frac{f(j, n)}{p(n)} &= 1 + e^{\frac{\sqrt{2\pi}j}{\sqrt{3}N}}\left(1 + \left(\frac{\sqrt{3}}{\sqrt{2\pi}} - \frac{\sqrt{3}}{\sqrt{2\pi}}\right)\frac{1}{\sqrt{N}} + \frac{2j}{N} - \frac{\pi j^2}{\sqrt{6}N^{\frac{3}{2}}} + O_{\leq}\left(\frac{2075}{N}\right)\right) \\ &\quad - e^{\frac{\pi j}{\sqrt{6}N}}\left(2 + \left(\frac{2\sqrt{3}}{\sqrt{2\pi}} - \frac{2\sqrt{3}}{\sqrt{2\pi}}\right)\frac{1}{\sqrt{N}} + \frac{2j}{N} - \frac{\pi j^2}{2\sqrt{6}N^{\frac{3}{2}}} + O_{\leq}\left(\frac{3926}{N}\right)\right). \end{aligned}$$

Proof. Recall that

$$\frac{f(j, n)}{p(n)} = 1 - 2\frac{p(n-j)}{p(n)} + \frac{p(n-2j)}{p(n)}.$$

We first bound $\frac{1}{p(n)}$. By Proposition 2.1 we have

$$p(n) = \frac{e^{\pi\sqrt{\frac{2N}{3}}}}{4\sqrt{3}N}\left(1 - \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} + g(N)\right),$$

where

$$|g(N)| \leq \frac{2\pi^2Ne^{-\pi\sqrt{\frac{2N}{3}}}}{3} + 2^3 \cdot 3^{-\frac{1}{2}}\pi N^{\frac{1}{2}}e^{-\frac{\pi}{2}\sqrt{\frac{N}{2}}} =: h(N).$$

Now we want to approximate

$$\frac{1}{1 - \left(\frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} - g(N)\right)}.$$

We have for N

$$\left| \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} - g(N) \right| < \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} + |g(N)| \leq \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} + h(N) < 0.99,$$

which can be seen by taking Taylor series. Now we claim that for $0 < |z| < 0.99$

$$\frac{1}{1-z} = 1 + z + O_{\leq} \left(100|z|^2 \right).$$

To see this, we bound

$$\left| \frac{1}{1-z} - 1 - z \right| = \left| \frac{1 - (1+z)(1-z)}{1-z} \right| = \frac{|z|^2}{|1-z|} \leq \frac{|z|^2}{1-|z|} < \frac{1}{0.01}|z|^2 = 100|z|^2.$$

Thus

$$\begin{aligned} \frac{1}{p(n)} &= \frac{4\sqrt{3}Ne^{-\pi\sqrt{\frac{2N}{3}}}}{1 - \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} + g(N)} \\ &= 4\sqrt{3}Ne^{-\pi\sqrt{\frac{2N}{3}}} \left(1 + \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} - g(N) + O_{\leq} \left(100 \left| \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} - g(N) \right|^2 \right) \right) \\ &= 4\sqrt{3}Ne^{-\pi\sqrt{\frac{2N}{3}}} \left(1 + \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} + O_{\leq} \left(|g(N)| + 100 \left| \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} - g(N) \right|^2 \right) \right). \end{aligned}$$

Now the error may be bound against

$$h(N) + 100 \left(\frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} + h(N) \right)^2 = O_{\leq} \left(\frac{1350}{N} \right)$$

again by basic calculus. Thus

$$\frac{1}{p(n)} = 4\sqrt{3}Ne^{-\pi\sqrt{\frac{2N}{3}}} \left(1 + \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N}}} + O_{\leq} \left(\frac{1350}{N} \right) \right).$$

Consider $p(n-J)$, $J \in \{j, 2j\}$. Then by Proposition 2.1, we have

$$p(N-J) = \frac{e^{\pi\sqrt{\frac{2}{3}}\sqrt{N-J}}}{4\sqrt{3}(N-J)} \left(1 - \frac{\sqrt{3}}{\sqrt{2\pi\sqrt{N-J}}} + O_{\leq} (h(N-J)) \right). \quad (2.7)$$

First note that $j < \frac{\sqrt{N}}{4}$ implies that (note that $N \geq 14 - \frac{1}{24}$)

$$N - J \geq N - 2j \geq N - \frac{\sqrt{N}}{2} \geq 12.$$

Moreover since $1 \leq j \leq \frac{\sqrt{N}}{4}$

$$\frac{J}{N} \leq \frac{2j}{N} \leq \frac{\sqrt{N}}{2N} = \frac{1}{2\sqrt{N}} \leq \frac{1}{2\sqrt{14 - \frac{1}{24}}} < 0.2.$$

We now first approximate the exponential in (2.7). We claim that

$$e^{\pi\sqrt{\frac{2}{3}}\sqrt{N-J}} = e^{\pi\sqrt{\frac{2}{3}}\sqrt{N}-\frac{\pi J}{\sqrt{6N}}} \left(1 - \frac{\pi J^2}{4\sqrt{6}N^{\frac{3}{2}}} + O_{\leq} \left(\frac{0.1}{N} \right) \right).$$

To see this define

$$f(x) := \sqrt{1-x} - 1 + \frac{x}{2} + \frac{x^2}{8}.$$

It is straightforward to show that for $0 \leq x < 0.2$ we have

$$|f(x)| \leq 0.1x^3.$$

We now use this to write

$$\sqrt{N-J} = \sqrt{N}\sqrt{1-\frac{J}{N}} = \sqrt{N} \left(f\left(\frac{J}{N}\right) + 1 - \frac{J}{2N} - \frac{1}{8}\left(\frac{J}{N}\right)^2 \right)$$

and obtain

$$e^{\pi\sqrt{\frac{2}{3}}\sqrt{N-J}-\pi\sqrt{\frac{2}{3}}\sqrt{N}+\frac{\pi J}{\sqrt{6N}}} = e^{-\pi\sqrt{\frac{2}{3}}\sqrt{N}\left(-f\left(\frac{J}{N}\right)+\frac{J^2}{8N^2}\right)}.$$

Note that

$$-f(x) + \frac{x^2}{8} = 1 - \frac{x}{2} - \sqrt{1-x} = \frac{x^2}{8} + \dots > 0.$$

Also note that for $x > 0$

$$e^{-x} - 1 + x = \sum_{n \geq 2} \frac{(-1)^n x^n}{n!} \leq \frac{x^2}{2}$$

by Leibnitz's criterium. Thus,

$$\begin{aligned} & e^{-\pi\sqrt{\frac{2}{3}}\sqrt{N}\left(-f\left(\frac{J}{N}\right)+\frac{J^2}{8N^2}\right)} \\ &= 1 - \pi\sqrt{\frac{2}{3}}\sqrt{N} \left(-f\left(\frac{J}{N}\right) + \frac{J^2}{8N^2} \right) + O_{\leq} \left(\frac{1}{2} \left(\pi\sqrt{\frac{2}{3}}\sqrt{N} \left(-f\left(\frac{J}{N}\right) + \frac{J^2}{8N^2} \right) \right)^2 \right) \\ &= 1 - \frac{\pi J^2}{\sqrt{6}N^{\frac{3}{2}}} + O_{\leq} \left(\pi\sqrt{\frac{2}{3}}\sqrt{N} \left| f\left(\frac{J}{N}\right) \right| + \frac{1}{2} \left(\pi\sqrt{\frac{2}{3}}\sqrt{N} \left(\left| f\left(\frac{J}{N}\right) \right| + \frac{J^2}{8N^2} \right) \right)^2 \right). \end{aligned}$$

We now bound the error against

$$\begin{aligned} & \pi\sqrt{\frac{2}{3}}\sqrt{N} \cdot 0.1 \left(\frac{J}{N} \right)^3 + \frac{\pi^2}{3} N \left(0.1 \left(\frac{J}{N} \right)^3 + \frac{J^2}{8N^2} \right)^2 \\ & \leq \pi\sqrt{\frac{2}{3}} \cdot 0.1 \frac{1}{2^3 N} + \frac{\pi^2}{3} N \left(0.1 \left(\frac{1}{2\sqrt{N}} \right)^3 + \frac{1}{8} \left(\frac{1}{2\sqrt{N}} \right)^2 \right)^2 \\ & = \frac{0.1\pi}{4\sqrt{6}} \frac{1}{N} + \frac{\pi^2}{24} \frac{1}{N} \left(\frac{0.1}{\sqrt{N}} + \frac{1}{4} \right)^2 = \left(\frac{0.1\pi}{4\sqrt{6}} + \frac{\pi^2}{24} \left(\frac{0.1}{\sqrt{N}} + \frac{1}{4} \right)^2 \right) \frac{1}{N} \leq \frac{0.1}{N}, \end{aligned}$$

where in the final inequality we use that $N \geq 14 - \frac{1}{24}$. This gives the claim.

Next we claim that for $x < 0.2$

$$\frac{1}{1-x} = 1 + x + O_{\leq} \left(1.25x^2 \right).$$

To see this, we bound

$$\left| \frac{1}{1-x} - 1 - x \right| = \frac{x^2}{|1-x|} \leq \frac{x^2}{1-|x|} \leq \frac{x^2}{0.8}.$$

We use this for

$$\frac{1}{N-J} = \frac{1}{N} \frac{1}{1-\frac{J}{N}} = \frac{1}{N} \left(1 + \frac{J}{N} + O_{\leq} \left(1.25 \left(\frac{J}{N} \right)^2 \right) \right) = \frac{1}{N} \left(1 + \frac{J}{N} + O_{\leq} \left(\frac{0.4}{N} \right) \right).$$

Next we use that for $0 \leq x < 0.2$

$$\frac{1}{\sqrt{1-x}} - 1 \leq 0.6x.$$

Thus

$$\begin{aligned} \frac{1}{\sqrt{N-J}} &= \frac{1}{\sqrt{N}} \frac{1}{\sqrt{1-\frac{J}{N}}} = \frac{1}{\sqrt{N}} \left(1 + O_{\leq} \left(0.6 \frac{J}{N} \right) \right) = \frac{1}{\sqrt{N}} + O_{\leq} \left(0.6 \frac{J}{N^{\frac{3}{2}}} \right) \\ &= \frac{1}{\sqrt{N}} + O_{\leq} \left(\frac{0.3}{N} \right). \end{aligned} \tag{2.8}$$

Finally, by basic calculus we find the bound

$$\frac{2\pi^2 x e^{-\pi\sqrt{\frac{2x}{3}}}}{3} + 2^3 3^{-\frac{1}{2}} \pi x^{\frac{1}{2}} e^{-\frac{\pi}{2}\sqrt{\frac{x}{2}}} \leq 15x^{\frac{1}{2}} e^{-\frac{\pi}{2}\sqrt{\frac{x}{2}}}$$

for $x \geq 12$. Thus

$$|f(N-J)| \leq 15(N-J)^{\frac{1}{2}} e^{-\frac{\pi}{2\sqrt{2}}\sqrt{N-J}}.$$

Now note that

$$x^{\frac{1}{2}} e^{-\frac{\pi}{2}\sqrt{\frac{x}{2}}}$$

is decreasing for $x \geq 1$. We then use the bound

$$N-J \geq N - \frac{\sqrt{N}}{2}$$

and thus

$$(N-J)^{\frac{1}{2}} e^{-\frac{\pi}{2}\sqrt{\frac{N-J}{2}}} \leq \left(N - \frac{\sqrt{N}}{2} \right)^{\frac{1}{2}} e^{-\frac{\pi}{2}\sqrt{\frac{N-\sqrt{N}}{2}}} = O_{\leq} \left(\frac{1.1}{N} \right),$$

Thus

$$\begin{aligned} p(n-J) &= \frac{1}{4\sqrt{3}} e^{\pi\sqrt{\frac{2N}{3}} - \frac{\pi J}{\sqrt{6N}}} \left(1 - \frac{\pi J^2}{4\sqrt{6}N^{\frac{3}{2}}} + O_{\leq} \left(\frac{0.1}{N} \right) \right) \frac{1}{N} \left(1 + \frac{J}{N} + O_{\leq} \left(\frac{0.4}{N} \right) \right) \\ &\quad \times \left(1 - \frac{\sqrt{3}}{\sqrt{2\pi}} \left(\frac{1}{\sqrt{N}} + O_{\leq} \left(\frac{0.3}{N} \right) \right) + O_{\leq} \left(\frac{1.1}{N} \right) \right). \end{aligned}$$

We combine

$$\begin{aligned} & \left(1 - \frac{\pi J^2}{4\sqrt{6}N^{\frac{3}{2}}} + O_{\leq} \left(\frac{0.1}{N}\right)\right) \left(1 + \frac{J}{N} + O_{\leq} \left(\frac{0.4}{N}\right)\right) \\ &= 1 + \frac{J}{N} - \frac{\pi J^2}{4\sqrt{6}N^{\frac{3}{2}}} + O_{\leq} \left(\frac{0.4}{N} + \frac{\pi J^3}{4\sqrt{6}N^{\frac{5}{2}}} + \frac{0.4\pi J^2}{4\sqrt{6}N^{\frac{5}{2}}} + \frac{0.1}{N} + 0.1 \frac{J}{N^2} + 0.1 \cdot \frac{0.4}{N^2}\right). \end{aligned}$$

The error may be bounded against $\frac{0.56}{N}$.

Next, we estimate

$$1 - \frac{\sqrt{3}}{\sqrt{2\pi}} \left(\frac{1}{\sqrt{N}} + O_{\leq} \left(\frac{0.3}{N}\right) + O_{\leq} \left(\frac{1.1}{N}\right)\right) = 1 - \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N}} + O_{\leq} \left(\frac{1.31}{N}\right).$$

Thus,

$$\begin{aligned} & 4\sqrt{3}Ne^{-\pi\sqrt{\frac{2N}{3}} + \frac{\pi J}{\sqrt{6}N}} p(n-J) \\ &= \left(1 + \frac{J}{N} - \frac{\pi J^2}{4\sqrt{6}N^{\frac{3}{2}}} + O_{\leq} \left(\frac{0.56}{N}\right)\right) \left(1 - \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N}} + O_{\leq} \left(\frac{1.31}{N}\right)\right) \\ &= 1 + \frac{J}{N} - \frac{\pi J^2}{4\sqrt{6}N^{\frac{3}{2}}} - \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N}} + O_{\leq} \left(\frac{2.71}{N}\right). \end{aligned}$$

Note that this calculation also proves Theorem 1.1. Thus, overall, we obtain

$$\begin{aligned} \frac{f(n,j)}{p(n)} &= 1 + \frac{1}{p(n)}(p(n-2j) - 2p(n-j)) \\ &= 1 + 4\sqrt{3}Ne^{-\pi\sqrt{\frac{2N}{3}}} \left(1 + \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N}} + O_{\leq} \left(\frac{1350}{N}\right)\right) \frac{e^{\pi\sqrt{\frac{2N}{3}}}}{4\sqrt{3}N} \\ &\quad \times \left(e^{\frac{\sqrt{2}\pi j}{\sqrt{3}N}} \left(1 + \frac{2j}{N} - \frac{\pi j^2}{\sqrt{6}N^{\frac{3}{2}}} - \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N}} + O_{\leq} \left(\frac{2.71}{N}\right)\right) \right. \\ &\quad \left. - 2e^{\frac{\pi j}{\sqrt{6}N}} \left(1 + \frac{j}{N} - \frac{\pi j^2}{4\sqrt{6}N^{\frac{3}{2}}} - \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N}} + O_{\leq} \left(\frac{2.71}{N}\right)\right)\right). \end{aligned} \tag{2.9}$$

Using that $j \leq \frac{\sqrt{N}}{4}$ and $n \geq 14$ a straightforward calculation gives

$$\begin{aligned} & \left(1 + \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N}} + O_{\leq} \left(\frac{1350}{N}\right)\right) \left(1 + \frac{2j}{N} - \frac{\pi j^2}{\sqrt{6}N^{\frac{3}{2}}} - \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N}} + O_{\leq} \left(\frac{2.71}{N}\right)\right) \\ &= 1 + \left(\frac{\sqrt{3}}{\sqrt{2\pi}} - \frac{\sqrt{3}}{\sqrt{2\pi}}\right) \frac{1}{\sqrt{N}} + \frac{2j}{N} - \frac{\pi j^2}{\sqrt{6}N^{\frac{3}{2}}} + O_{\leq} \left(\frac{2075}{N}\right). \end{aligned}$$

We turn to the final product in (2.9) given by

$$2 \left(1 + \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N}} + O_{\leq} \left(\frac{1350}{N}\right)\right) \left(1 + \frac{j}{N} - \frac{\pi j^2}{4\sqrt{6}N^{\frac{3}{2}}} - \frac{\sqrt{3}}{\sqrt{2\pi}\sqrt{N}} + O_{\leq} \left(\frac{2.71}{N}\right)\right).$$

Again using $j \leq \frac{\sqrt{N}}{4}$ and $n \geq 14$ it is not hard to show that this is equal to

$$2 + \left(\frac{2\sqrt{3}}{\sqrt{2\pi}} - \frac{2\sqrt{3}}{\sqrt{2\pi}} \right) \frac{1}{\sqrt{N}} + \frac{2j}{N} - \frac{\pi j^2}{2\sqrt{6}N^{\frac{3}{2}}} + O_{\leq} \left(\frac{3926}{N} \right)$$

Combining everything together, we obtain the statement of the theorem. \square

We end by proving the eventual positivity of the ratio $\frac{f(j,n)}{p(n)}$, which is crucial to the applications in the introduction.

Theorem 2.3. *Let $j \leq \frac{\sqrt{N}}{4}$. Then we have that $\frac{f(j,n)}{p(n)} > 0$ for all $n \geq 3922$.*

Proof. Let $X = e^{\frac{\sqrt{2\pi}j}{\sqrt{3N}}} - 2e^{\frac{\pi j}{\sqrt{6N}}}$. We have by Theorem 2.2 that the result follows if

$$\begin{aligned} \frac{f(j,n)}{p(n)} - 1 &= \left(1 + \left(\frac{\sqrt{3}}{\sqrt{2\pi}} - \frac{\sqrt{3}}{\sqrt{2\pi}} \right) \frac{1}{\sqrt{N}} + \frac{j}{N} - \frac{\pi j^2}{4\sqrt{6}N^{\frac{3}{2}}} + \frac{3926}{N} \right) X \\ &\quad + e^{\frac{\sqrt{2\pi}j}{\sqrt{3N}}} \left(\frac{j}{N} - \frac{3\pi j^2}{4\sqrt{6}N^{\frac{3}{2}}} \right) > -1 \end{aligned} \quad (2.10)$$

for all $n \geq 3922$.

We first observe using $j \leq \frac{\sqrt{N}}{4}$ that

$$\frac{3\pi j^2}{4\sqrt{6}N^{\frac{3}{2}}} < \frac{j}{N},$$

implying that the final term in (2.10) is positive. Thus, the claim follows if

$$\left(1 + \left(\frac{\sqrt{3}}{\sqrt{2\pi}} - \frac{\sqrt{3}}{\sqrt{2\pi}} \right) \frac{1}{\sqrt{N}} + \frac{j}{N} - \frac{\pi j^2}{4\sqrt{6}N^{\frac{3}{2}}} + \frac{3926}{N} \right) X > -1.$$

Since $-1 < X < 0$ for all $n \geq 4$, this follows if

$$\left(\frac{\sqrt{3}}{\sqrt{2\pi}} - \frac{\sqrt{3}}{\sqrt{2\pi}} \right) \frac{1}{\sqrt{N}} + \frac{j}{N} - \frac{\pi j^2}{4\sqrt{6}N^{\frac{3}{2}}} + \frac{3926}{N} < 0.$$

It is simple to check that this is always satisfied for $n \geq 3922$, and the theorem follows. \square

REFERENCES

- [1] A. Akande, T. Genao, S. Haag, M. Hendon, N. Pulagam, R. Schneider, and A. Sills, *Computational study of non-unitary partitions*, <https://arxiv.org/abs/2112.03264>, preprint.
- [2] K. Bringmann, B. Kane, L. Rolén, and Z. Tripp, *Fractional partitions and conjectures of Chern-Fu-Tang and Heim-Neuhauser*, Trans. Am. Math. Soc., Series B, accepted for publication.
- [3] F. Dyson, *A new symmetry of partitions*, J. Combinatorial Theory **7** (1969), 56–61.
- [4] Garvan, F., *Generalizations of Dyson's rank and non-Rogers-Ramanujan partitions*, Manuscripta Math., **84** (1994), n0. 3-4, 343–359.
- [5] M. Griffin, K. Ono, Ken, L. Rolén, and D. Zagier, *Jensen polynomials for the Riemann zeta function and other sequences*, Proc. Natl. Acad. Sci. USA, **116** (2019), no. 23, 11103–11110.
- [6] H. Gupta, *Finite differences of the partition function*, Math. Comp. **32** (1978), 144 1241–1243.

- [7] G. Hardy and S. Ramanujan, *Asymptotic formulae in combinatory analysis*, Proc. London Math. Soc. Ser. 2 **17** (1918), 75–115.
- [8] J. Iskander, V. Jain, and V. Talvola, *Exact formulae for the fractional partition functions*, Res. Number Theory **6** (2020), 201–215.
- [9] H. Larson and I. Wagner, *Hyperbolicity of the partition Jensen polynomials*, Res. Number Theory, **5** (2019), no. 2, Paper No. 19, 12.
- [10] D. Lehmer, *On the remainders and convergence of the series for the partition function*, Trans. Amer. Math. Soc., **46** (1939), 362–373.
- [11] D. Lehmer, *On the series for the partition function*, Trans. Amer. Math. Soc., **43** (1938), no. 2, 271–295.
- [12] Z. Liu and N. H. Zhou, *Uniform asymptotic formulas for the Fourier coefficients of the inverse of theta functions*, <https://arxiv.org/abs/1903.00835>, preprint.
- [13] H. Rademacher, *A convergent series for the partition function $p(n)$* , PNAS February 1, 1937 **23** (2), 78–84.
- [14] R. Schneider, *Nuclear partitions and a formula for $p(n)$* , <https://arxiv.org/abs/1912.00575>, preprint.
- [15] G. Watson, *A treatise on the theory of Bessel functions*, Cambridge University Press (1995).

DEPARTMENT OF MATHEMATICS, 1420 STEVENSON CENTER, VANDERBILT UNIVERSITY, NASHVILLE, TN 37240
Email address: kevin.j.gomez@vanderbilt.edu

450 MACHRAY HALL, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANITOBA, WINNIPEG, CANADA
Email address: joshua.males@umanitoba.ca

DEPARTMENT OF MATHEMATICS, 1420 STEVENSON CENTER, VANDERBILT UNIVERSITY, NASHVILLE, TN 37240
Email address: larry.rolen@vanderbilt.edu