


# Max-Planck-Institut für Mathematik Bonn

## Anderson finiteness for RCD spaces

by

Sergio Zamora



Max-Planck-Institut für Mathematik  
Preprint Series 2023 (4)

Date of submission: February 2, 2023

# Anderson finiteness for RCD spaces

by

Sergio Zamora

Max-Planck-Institut für Mathematik  
Vivatsgasse 7  
53111 Bonn  
Germany

# Anderson finiteness for RCD spaces

Sergio Zamora

zamora@mpim-bonn.mpg.de

## 1 Introduction

For  $K \in \mathbb{R}$ ,  $N \in [1, \infty)$ , the class of  $RCD^*(K, N)$  spaces consists of proper metric measure spaces that satisfy a synthetic condition of having Ricci curvature bounded below by  $K$  and dimension bounded above by  $N$ . This class is closed under measured Gromov–Hausdorff convergence and contains the class of complete Riemannian manifolds of Ricci curvature  $\geq K$  and dimension  $\leq N$ .

$RCD^*(K, N)$  spaces have a well defined notion of dimension called *rectifiable dimension* (see Theorem 16), which is always an integer between 0 and  $N$ , and is lower semi-continuous with respect to pointed Gromov–Hausdorff convergence (see Theorem 20). This motivates the following definition.

**Definition 1.** Let  $K \in \mathbb{R}$ ,  $N \in [1, \infty)$ , and  $(X, p)$  a pointed  $RCD^*(K, N)$  space. We define the *collapsing volume* of  $(X, p)$  as

$$\text{vol}_{K,N}^*(X, p) := \inf d_{GH}((X, p), (Y, q)),$$

where the infimum is taken among all pointed  $RCD^*(K, N)$  spaces  $(Y, q)$  whose rectifiable dimension is strictly less than the one of  $X$ .

**Proposition 2.** Let  $(X_i, p_i)$  be a sequence of pointed  $RCD^*(K, N)$  spaces that converges in the Gromov–Hausdorff sense to a pointed  $RCD^*(K, N)$  space  $(X, p)$ . If the rectifiable dimension of  $X_i$  is  $n$  for each  $i$ , then the following are equivalent:

1. The rectifiable dimension of  $X$  is strictly less than  $n$ .
2.  $\text{vol}_{K,N}^*(X_i, p_i) \rightarrow 0$ .
3.  $\text{vol}_{K,N}^*(X_{i_k}, p_{i_k}) \rightarrow 0$  for a subsequence.

*Proof.* (3  $\Rightarrow$  1) By hypothesis, there is a sequence of  $RCD^*(K, N)$  spaces  $(Y_{i_k}, q_{i_k})$  of rectifiable dimension strictly less than  $n$  and converging to  $(X, p)$  as  $k \rightarrow \infty$ . From the fact that rectifiable dimension is lower semi-continuous, we deduce 1. The implications 1  $\Rightarrow$  2  $\Rightarrow$  3 are tautological.  $\square$

**Corollary 3.** Let  $X_i$  be a sequence of  $RCD^*(K, N)$  spaces, and  $p_i, p'_i \in X_i$  pairs of points with  $\limsup_{i \rightarrow \infty} d(p_i, p'_i) < \infty$ . Then  $\text{vol}_{K,N}^*(X_i, p_i) \rightarrow 0$  if and only if  $\text{vol}_{K,N}^*(X_i, p'_i) \rightarrow 0$ .

The main result of this note is a generalization to  $RCD^*(K, N)$  spaces of a classical finiteness result of Anderson [1].

**Theorem 4.** For each  $K \in \mathbb{R}$ ,  $N \in [1, \infty)$ ,  $D > 0$ ,  $\nu > 0$ , the class of pointed  $RCD^*(K, N)$  spaces of diameter  $\leq D$  and  $\text{vol}_{K,N}^* \geq \nu$  contains finitely many fundamental group isomorphism types.

Theorem 4 will be obtained from the following result which states that a lower bound on the collapsing volume of the quotient of an  $RCD^*(K, N)$  space by a discrete group yields a uniform discreteness gap on the corresponding group (see Equation 1 below for the definition of  $d_p$ ).

**Theorem 5.** For each  $K \in \mathbb{R}$ ,  $N \in [1, \infty)$ ,  $\nu > 0$ , there is  $\varepsilon > 0$  such that the following holds. If  $(X, p)$  is a pointed  $RCD^*(K, N)$  space and  $\Gamma \leq \text{Iso}(X)$  is a discrete group of measure preserving isometries with  $\text{vol}_{K,N}^*(X/\Gamma, [p]) \geq \nu$ , then

$$\{g \in \Gamma \mid d_p(g, Id_X) \leq \varepsilon\} = \{Id_X\}.$$

The remainder of this note contains the proof of theorems 4 and 5. In Section 2 we cover the required material and in Section 3 we present the proofs.

## 2 Preliminaries

### 2.1 Notation

For a metric space  $X$ ,  $p \in X$ ,  $r > 0$ , the closed ball of radius  $r$  centered at  $p$  will be denoted as  $B(p, r, X)$ . We denote by  $X \cup \{*\}$  the metric space obtained by adjoining to  $X$  a point  $*$  with  $d(x, *) = \infty$  for all  $x \in X$ .

### 2.2 RCD spaces and isometries

A  $CD^*(K, N)$  space is a proper metric space  $(X, d)$  equipped with a fully supported Radon measure  $\mathbf{m}$  for which an appropriate entropy in its space of probability measures is in a suitable sense concave with respect to the  $L^2$ -Wasserstein distance. For a  $CD^*(K, N)$  space  $(X, d, \mathbf{m})$ , if its Sobolev space  $W^{1,2}$  is a Hilbert space, we say that it is an  $RCD^*(K, N)$  space. See [7] for a precise definition and different reformulations.

**Remark 6.** If  $(X, d, \mathbf{m})$  is an  $RCD^*(K, N)$  space, then for any  $c > 0$ ,  $(X, d, c\mathbf{m})$  is also an  $RCD^*(K, N)$  space, and for any  $\lambda > 0$ ,  $(X, \lambda d, \mathbf{m})$  is an  $RCD^*(\lambda^{-2}K, N)$  space. Also, if a metric measure space  $(X, d, \mathbf{m})$  is an  $RCD^*(K - \varepsilon, N)$  space for all  $\varepsilon > 0$ , then it is also an  $RCD^*(K, N)$  space.

Along this note we are interested only in topological properties of  $RCD^*(K, N)$  spaces, so by an abuse of notation, we say that a proper metric space  $(X, d)$  is an  $RCD^*(K, N)$  space if it admits a Radon measure that makes it an  $RCD^*(K, N)$  space. Any two such

measures are equivalent, so we can still talk about *full (or zero) measure sets* even after this abuse.

Even though we don't know much about the global topology of  $RCD^*(K, N)$  spaces, we know they are semi-locally-simply-connected [16], and their universal cover is still an  $RCD^*(K, N)$  space [14].

**Theorem 7.** (Wang) Let  $X$  be an  $RCD^*(K, N)$  space. Then  $X$  is semi-locally-simply-connected, so for any  $p \in X$  we can identify the fundamental group  $\pi_1(X, p)$  with the group of deck transformations of the universal cover  $\tilde{X}$ .

**Theorem 8.** (Mondino–Wei) If  $X$  is an  $RCD^*(K, N)$  space, then its universal cover  $\tilde{X}$  admits a  $\pi_1(X)$ -invariant measure that makes it an  $RCD^*(K, N)$  space.

Recall that for Riemannian manifolds, if an isometry coincides with the identity up to first order at a point, then it is necessarily the identity. An analogue of this statement for  $RCD^*(K, N)$  spaces is the following [11].

**Lemma 9.** Let  $X$  be an  $RCD^*(K, N)$  space and  $f : X \rightarrow X$  a non-trivial isometry. Then the set of fixed points of  $f$  has zero measure.

For a pointed proper metric space  $(X, p)$ , we define the compact-open distance between two functions  $h_1, h_2 : X \rightarrow X$  as

$$d_p(h_1, h_2) := \inf_{r>0} \left\{ \frac{1}{r} + \sup_{x \in B(p, r, X)} d(h_1 x, h_2 x) \right\}. \quad (1)$$

When we restrict this metric to the group of isometries  $Iso(X)$ , we get a left invariant (not necessarily geodesic) metric that induces the compact open topology (independent of  $p$ ) and makes  $Iso(X)$  a proper metric group. However, this distance is defined on the full class of functions  $X \rightarrow X$ , where it is not left invariant nor proper anymore.

Recall that if  $X$  is a proper geodesic space and  $\Gamma \leq Iso(X)$  is a closed group of isometries, the metric  $d'$  on  $X/\Gamma$  defined as  $d'([x], [y]) := \inf_{g \in \Gamma} (d(gx, y))$  makes it a proper geodesic space. By the work of Galaz–Kell–Mondino–Sosa, the class of  $RCD^*(K, N)$  spaces is closed under quotients by discrete groups [9].

**Theorem 10.** (Galaz–Kell–Mondino–Sosa) Let  $(X, d, \mathbf{m})$  be an  $RCD^*(K, N)$  space and  $\Gamma \leq Iso(X)$  a discrete group of measure preserving isometries. Then the metric space  $(X/\Gamma, d')$  admits a measure  $\mathbf{m}'$  that makes it an  $RCD^*(K, N)$  space. Moreover, if  $\rho : X \rightarrow X/\Gamma$  denotes the projection,  $\mathbf{m}'$  can be taken so that  $\mathbf{m}(A) = \mathbf{m}'(\rho(A))$  for all Borel subsets  $A$  of  $X$  sent isometrically to  $X/\Gamma$  by  $\rho$ .

## 2.3 Gromov–Hausdorff topology

The Gromov–Hausdorff topology in the class of pointed proper metric spaces quantifies how much two spaces are from being isometric.

**Definition 11.** Let  $(X, p), (Y, q)$  be pointed proper metric spaces and  $\varepsilon > 0$ . We say that a function  $f : X \rightarrow Y \cup \{*\}$  is an  $\varepsilon$ -approximation if  $d(f(p), q) \leq \varepsilon$  and for  $R = 1/\varepsilon$  one has

$$f^{-1}(B(q, R, Y)) \subset B(p, 2R, X), \quad (2)$$

$$\sup_{x_1, x_2 \in B(p, 2R, X)} |d(f(x_1), f(x_2)) - d(x_1, x_2)| \leq \varepsilon, \quad (3)$$

$$\sup_{y \in B(q, R, Y)} \inf_{x \in B(p, 2R, X)} d(f(x), y) \leq \varepsilon. \quad (4)$$

The *pointed Gromov–Hausdorff distance* between  $(X, p)$  and  $(Y, q)$  is defined as

$$d_{GH}((X, p), (Y, q)) := \inf\{\varepsilon > 0 \mid \text{there is an } \varepsilon\text{-approximation } f : X \rightarrow Y \cup \{*\}\}.$$

Strictly speaking,  $d_{GH}$  as defined above is not a distance as it is not symmetric. However, it still generates a first countable Hausdorff topology in the class of pointed proper metric spaces (see [4], Chapter 8). This is called the *Gromov–Hausdorff topology*.

**Proposition 12.** (Gromov) For pointed proper metric spaces  $(X, p)$  and  $(X_i, p_i)$ , we have  $d_{GH}((X_i, p_i), (X, p)) \rightarrow 0$  as  $i \rightarrow \infty$  if and only if  $d_{GH}((X, p), (X_i, p_i)) \rightarrow 0$  as  $i \rightarrow \infty$ . Moreover, in either case there are sequences  $\phi_i : X_i \rightarrow X \cup \{*\}$  and  $\psi_i : X \rightarrow X_i \cup \{*\}$  of  $\varepsilon_i$ -approximations with  $\varepsilon_i \rightarrow 0$  and such that

$$\lim_{i \rightarrow \infty} d_p(\phi_i \circ \psi_i, Id_X) = 0. \quad (5)$$

**Corollary 13.** Let  $(X_i, p_i)$  be a sequence of pointed proper metric spaces that converges in the Gromov–Hausdorff sense to a pointed proper metric space  $(X, p)$ . Then for each  $R > 0$ ,  $\varepsilon > 0$ , there is  $M \in \mathbb{N}$  such that any set  $S \subset B(p_i, R, X_i)$  with  $d(s_1, s_2) \geq \varepsilon$  for each  $s_1, s_2 \in S$ , one has  $|S| \leq M$ .

One of the main features of the class of  $RCD^*(K, N)$  spaces is the Gromov–Hausdorff compactness property [2].

**Theorem 14.** (Bacher–Sturm) If  $(X_i, p_i)$  is a sequence of pointed  $RCD^*(K, N)$  spaces, then one can find a subsequence that converges in the Gromov–Hausdorff sense to a pointed  $RCD^*(K, N)$  space  $(X, p)$ .

**Definition 15.** Let  $X$  be an  $RCD^*(K, N)$  space and  $n \in \mathbb{N}$ . We say that  $p \in X$  is an  $n$ -regular point if for each  $\lambda_i \rightarrow \infty$ , the sequence  $(\lambda_i X, p)$  converges in the Gromov–Hausdorff sense to  $(\mathbb{R}^n, 0)$ .

Mondino–Naber showed that the set of regular points in an  $RCD^*(K, N)$  space has full measure [13]. This result was refined by Brué–Semola who showed that almost all points have the same local dimension [3].

**Theorem 16.** (Brué–Semola) Let  $X$  be an  $RCD^*(K, N)$  space. Then there is a unique  $n \in \mathbb{N} \cap [0, N]$  such that the set of  $n$ -regular points in  $X$  has full measure. This number  $n$  is called the *rectifiable dimension* of  $X$ .

**Definition 17.** Let  $X_i$  be a sequence of  $RCD^*(K, N)$  spaces of rectifiable dimension  $n$ . A choice of points  $x_i \in X_i$  is said to be a *Reifenberg sequence* if for any  $\lambda_i \rightarrow \infty$ , the sequence  $(\lambda_i X_i, x_i)$  converges in the Gromov–Hausdorff sense to  $(\mathbb{R}^n, 0)$ .

**Theorem 18.** (Mondino–Naber) For each  $i \in \mathbb{N}$ , let  $(X_i, d_i, \mathbf{m}_i)$  be an  $RCD^*(-\varepsilon_i, N)$  space with  $\varepsilon_i \rightarrow 0$  of rectifiable dimension  $n$ . Assume that for some choice  $p_i \in X_i$ , the sequence  $(X_i, p_i)$  converges in the Gromov–Hausdorff sense to  $(\mathbb{R}^n, 0)$ . Then there is a sequence of subsets  $U_i \subset B(p_i, 1, X_i)$  with  $\mathbf{m}_i(U_i)/\mathbf{m}_i(B(p_i, 1, X_i)) \rightarrow 1$  such that any sequence  $x_i \in U_i$  is a Reifenberg sequence.

From the work of Mondino–Naber and Brué–Semola, we can conclude that the rectifiable dimension is a topological invariant.

**Corollary 19.** Let  $X$  be an  $RCD^*(K, N)$  space of rectifiable dimension  $n$ . There is an open dense subset  $U \subset X$  locally bi-Lipschitz homeomorphic to  $\mathbb{R}^n$ .

Using the results above, Kitabeppu showed that the rectifiable dimension is lower semi-continuous with respect to Gromov–Hausdorff convergence [12].

**Theorem 20.** (Kitabeppu) Let  $(X_i, p_i)$  be a sequence of pointed  $RCD^*(K, N)$  spaces of rectifiable dimension  $n$  converging in the Gromov–Hausdorff sense to the space  $(X, p)$ . Then the rectifiable dimension of  $X$  is at most  $n$ .

The well known Cheeger–Gromoll splitting theorem [6] was extended by Cheeger–Colding for limits of Riemannian manifolds with lower Ricci curvature bounds [5], and later by Gigli to this setting [10].

**Theorem 21.** (Gigli) Let  $(X, d, \mathbf{m})$  be an  $RCD^*(0, N)$  space of rectifiable dimension  $n$ . If  $(X, d)$  contains an isometric copy of  $\mathbb{R}^m$ , then there is  $c > 0$  and a metric measure space  $(Y, d^Y, \mathbf{n})$  such that  $(X, d, c\mathbf{m})$  is isomorphic to the product  $(\mathbb{R}^m \times Y, d^{\mathbb{R}^m} \times d^Y, \mathcal{H}^m \otimes \mathbf{n})$ . Moreover,  $(Y, d^Y, \mathbf{n})$  is an  $RCD^*(0, N - m)$  space of rectifiable dimension  $n - m$ .

## 2.4 Equivariant Gromov–Hausdorff convergence

Recall from Proposition 12 that if a sequence of pointed proper metric spaces  $(X_i, p_i)$  converges in the Gromov–Hausdorff sense to the pointed proper metric space  $(X, p)$ , one has  $\varepsilon_i$ -approximations  $\phi_i : X_i \rightarrow X \cup \{*\}$  and  $\psi_i : X \rightarrow X_i \cup \{*\}$  with  $\varepsilon_i \rightarrow 0$  and satisfying Equation 5.

**Definition 22.** Consider a sequence of pointed proper metric spaces  $(X_i, p_i)$  that converges in the Gromov–Hausdorff sense to a pointed proper metric space  $(X, p)$  and a sequence of closed groups of isometries  $\Gamma_i \leq Iso(X_i)$ . We say that the sequence  $\Gamma_i$  *converges equivariantly* to a closed group  $\Gamma \leq Iso(X)$  if:

- For each  $g \in \Gamma$ , there is a sequence  $g_i \in \Gamma_i$  with  $d_p(\psi_i \circ g_i \circ \phi_i, g) \rightarrow 0$  as  $i \rightarrow \infty$ .
- For a sequence  $g_i \in \Gamma_i$  and  $g \in Iso(X)$ , if there is a subsequence  $g_{i_k}$  with  $d_p(\psi_{i_k} \circ g_{i_k} \circ \phi_{i_k}, g) \rightarrow 0$  as  $k \rightarrow \infty$ , then  $g \in \Gamma$ .

We say that a sequence of isometries  $g_i \in Iso(X_i)$  *converges* to an isometry  $g \in Iso(X)$  if

$$d_p(\psi_i \circ g_i \circ \phi_i, g) \rightarrow 0 \text{ as } i \rightarrow \infty.$$

Equivariant convergence allows one to take limits before or after taking quotients [8].

**Lemma 23.** Let  $(Y_i, q_i)$  be a sequence of proper metric spaces that converges in the Gromov–Hausdorff sense to a proper space  $(Y, q)$ , and  $\Gamma_i \leq Iso(Y_i)$  a sequence of closed groups of isometries that converges equivariantly to a closed group  $\Gamma \leq Iso(Y)$ . Then the sequence  $(Y_i/\Gamma_i, [q_i])$  converges in the Gromov–Hausdorff sense to  $(Y/\Gamma, [q])$ .

Fukaya–Yamaguchi obtained an Arzelá–Ascoli type result for equivariant convergence ([8], Proposition 3.6).

**Theorem 24.** (Fukaya–Yamaguchi) Let  $(Y_i, q_i)$  be a sequence of proper metric spaces that converges in the pointed Gromov–Hausdorff sense to a proper space  $(Y, q)$ , and take a sequence  $\Gamma_i \leq Iso(Y_i)$  of closed groups of isometries. Then there is a subsequence  $(Y_{i_k}, q_{i_k}, \Gamma_{i_k})_{k \in \mathbb{N}}$  such that  $\Gamma_{i_k}$  converges equivariantly to a closed group  $\Gamma \leq Iso(Y)$ .

**Proposition 25.** Let  $(X_i, p_i)$  be a sequence of pointed proper metric spaces that converges in the Gromov–Hausdorff sense to a pointed proper metric space  $(X, p)$ . Assume a sequence of closed groups  $\Gamma_i \leq Iso(X_i)$  converges equivariantly to a closed group  $\Gamma \leq Iso(X)$ . Then the sequence of pointed metric spaces  $(\Gamma_i, d_{p_i}, Id_{X_i})$  converges in the Gromov–Hausdorff sense to  $(\Gamma, d_p, Id_X)$ .

*Proof.* Recall that one has  $\varepsilon_i$ -approximations  $\phi_i : X_i \rightarrow X \cup \{*\}$  and  $\psi_i : X \rightarrow X_i \cup \{*\}$  with  $\varepsilon_i \rightarrow 0$  and satisfying Equation 5. With these functions, one could define  $f_i : \Gamma_i \rightarrow \Gamma \cup \{*\}$  in the following way: for each  $g \in \Gamma_i$ , if there is an element  $\gamma \in \Gamma$  with  $d_p(\phi_i \circ g \circ \psi_i, \gamma) \leq 1$ , choose  $f_i(g)$  to be an element of  $\Gamma$  that minimizes  $d_p(\phi_i \circ g \circ \psi_i, f_i(g))$ . Otherwise, set  $f_i(g) = *$ . Now we verify that  $f_i$  are  $\delta_i$ -approximations for some  $\delta_i \rightarrow 0$ . The fact that  $f_i(Id_{X_i}) \rightarrow Id_X$  follows from Equation 5.

The fourth condition, corresponding to Equation 4, follows directly from the first condition in the definition of equivariant convergence and our construction.

To check that  $f_i$  satisfy the second condition, corresponding to Equation 2, assume by contradiction that after taking a subsequence we can find  $g_i \in \Gamma_i$  such that  $d_{p_i}(g_i, Id_{X_i}) \rightarrow$



$\infty$  but  $d_p(f_i(g_i), Id_X)$  is bounded. This implies that  $d(g_i(p_i), p_i) \rightarrow \infty$  and since  $g_i$  is an isometry,  $d(g_i \circ \psi_i(p), p_i) \rightarrow \infty$ . As  $\phi_i$  are  $\varepsilon_i$ -approximations for  $\varepsilon_i \rightarrow 0$ ,  $d(\phi_i \circ g_i \circ \psi_i(p), p) \rightarrow \infty$  as well. On the other hand, as  $d_p(f_i(g_i), Id_X)$  is bounded,  $f_i(g_i) \neq *$  and  $d_p(\phi_i \circ g_i \circ \psi_i, f_i(g_i)) \leq 1$  for all  $i$ , meaning that  $d_p(\phi_i \circ g_i \circ \psi_i, Id_X)$  is bounded, which is a contradiction.

To verify the third condition, corresponding to Equation 3, assume by contradiction that after taking a subsequence we can find two sequences  $g_i, h_i \in \Gamma_i$  such that the sequences  $d_{p_i}(g_i, Id_{X_i}), d_{p_i}(h_i, Id_{X_i})$  are bounded but  $|d_{p_i}(g_i, h_i) - d_p(f_i(g_i), f_i(h_i))| \geq \eta$  for some  $\eta > 0$ . After again taking a subsequence we can assume  $g_i$  converges to  $g \in \Gamma$  and  $h_i$  converges to  $h \in \Gamma$ . This means that  $d_p(f_i(g_i), \phi_i \circ g_i \circ \psi_i) \rightarrow 0$ ,  $d_p(f_i(h_i), \phi_i \circ h_i \circ \psi_i) \rightarrow 0$ . Hence for  $i$  large enough one has  $|d_{p_i}(g_i, h_i) - d_p(\phi_i \circ g_i \circ \psi_i, \phi_i \circ h_i \circ \psi_i)| \geq \eta/2$ .

We first deal with the case when

$$d_p(\phi_i \circ g_i \circ \psi_i, \phi_i \circ h_i \circ \psi_i) \geq d_{p_i}(g_i, h_i) + \eta/2 \quad (6)$$

for infinitely many  $i$ . By definition of  $d_{p_i}$ , there is a sequence  $\rho_i > 0$  with

$$d_{p_i}(g_i, h_i) + \frac{\eta}{4} \geq \frac{1}{\rho_i} + \sup_{x \in B(p_i, \rho_i, X_i)} d(g_i x, h_i x).$$

Setting  $r_i := \min\{\rho_i, 12/\eta\}$ , we obtain a bounded sequence such that

$$d_{p_i}(g_i, h_i) + \frac{\eta}{3} \geq \frac{1}{r_i} + \sup_{x \in B(p_i, r_i, X_i)} d(g_i x, h_i x).$$

For  $i$  large enough and  $x \in B(p, r_i - 2\varepsilon_i, X)$ ,

$$\begin{aligned} d(\phi_i \circ g_i \circ \psi_i(x), \phi_i \circ h_i \circ \psi_i(x)) &\leq \varepsilon_i + d(g_i(\psi_i(x)), h_i(\psi_i(x))) \\ &\leq \varepsilon_i - \frac{1}{r_i} + d_{p_i}(g_i, h_i) + \frac{\eta}{3}. \end{aligned}$$

Again from the definition of  $d_p$  we deduce

$$\begin{aligned} d_p(\phi_i \circ g_i \circ \psi_i, \phi_i \circ h_i \circ \psi_i) &\leq \frac{1}{r_i - 2\varepsilon_i} + \sup_{x \in B(p, r_i - 2\varepsilon_i, X)} d(\phi_i \circ g_i \circ \psi_i(x), \phi_i \circ h_i \circ \psi_i(x)) \\ &\leq \frac{1}{r_i - 2\varepsilon_i} - \frac{1}{r_i} + d_{p_i}(g_i, h_i) + \frac{\eta}{3} + \varepsilon_i. \end{aligned}$$

From the fact that  $d_{p_i}(g_i, Id_{X_i}), d_{p_i}(h_i, Id_{X_i})$  are bounded, we get that  $r_i$  is bounded away from 0. Then the right hand side is less than  $d_{p_i}(g_i, h_i) + \eta/2$  for  $i$  large enough, contradicting Equation 6. The other case when

$$d_{p_i}(g_i, h_i) \geq d_p(\phi_i \circ g_i \circ \psi_i, \phi_i \circ h_i \circ \psi_i) + \eta/2$$

for infinitely many  $i$  is analogous. □

As a consequence of Theorem 21, it is easy to understand the situation when the quotients of a sequence converge to  $\mathbb{R}^n$ .

**Lemma 26.** For each  $i \in \mathbb{N}$ , let  $(X_i, p_i)$  be a pointed  $RCD^*(-\varepsilon_i, N)$  space of rectifiable dimension  $n$  with  $\varepsilon_i \rightarrow 0$ . Assume  $(X_i, p_i)$  converges in the Gromov–Hausdorff sense to a pointed  $RCD^*(0, N)$  space  $(X, p)$ , there is a sequence of closed groups of isometries  $\Gamma_i \leq Iso(X_i)$  that converges equivariantly to  $\Gamma \leq Iso(X)$ , and the sequence of pointed proper metric spaces  $(X_i/\Gamma_i, [p_i])$  converges in the Gromov–Hausdorff sense to  $(\mathbb{R}^n, 0)$ . Then  $\Gamma$  is trivial.

*Proof.* One can use the submetry  $X \rightarrow X/\Gamma = \mathbb{R}^n$  to lift the lines of  $\mathbb{R}^n$  to lines in  $X$  passing through  $p$ . By iterated applications of Theorem 21, we get that  $X = \mathbb{R}^n \times Y$  for some  $RCD^*(0, N - n)$  space  $Y$ . But from Theorem 20, the rectifiable dimension of  $X$  is at most  $n$ , so  $Y$  is a point. Since  $\Gamma \leq Iso(\mathbb{R}^n)$  satisfies  $\mathbb{R}^n/\Gamma = \mathbb{R}^n$ , it must be trivial.  $\square$

## 2.5 Group norms

Let  $(X, p)$  be a pointed proper geodesic space and  $\Gamma \leq Iso(X)$  a group of isometries. The norm  $\|\cdot\|_p : \Gamma \rightarrow \mathbb{R}$  associated to  $p$  is defined as  $\|g\|_p := d(gp, p)$ . The spectrum  $\sigma(\Gamma, X, p)$  is defined as the set of  $r \geq 0$  such that

$$\langle \{g \in \Gamma \mid \|g\|_p \leq r\} \rangle \neq \langle \{g \in \Gamma \mid \|g\|_p \leq r - \varepsilon\} \rangle \text{ for all } \varepsilon > 0.$$

This spectrum is closely related to the covering spectrum introduced by Sormani–Wei in [15], and it also satisfies a continuity property.

**Proposition 27.** Let  $(X_i, p_i)$  be a sequence of pointed proper metric spaces that converges in the Gromov–Hausdorff sense to  $(X, p)$  and consider a sequence of closed isometry groups  $\Gamma_i \leq Iso(X_i)$  that converges equivariantly to a closed group  $\Gamma \leq Iso(X)$ . Then for any convergent sequence  $r_i \in \sigma(\Gamma_i, X_i, p_i)$ , we have  $\lim_{i \rightarrow \infty} r_i \in \sigma(\Gamma, X, p)$ .

*Proof.* Let  $r = \lim_{i \rightarrow \infty} r_i$ . By definition, there is a sequence  $g_i \in \Gamma_i$  with  $\|g_i\|_p = r_i$ , and  $g_i \notin \langle \{\gamma \in \Gamma_i \mid \|\gamma\|_{p_i} \leq r_i - \varepsilon\} \rangle$  for all  $\varepsilon > 0$ . Up to subsequence, we can assume that  $g_i$  converges to some  $g \in Iso(X)$  with  $\|g\|_p = r$ .

If  $r \notin \sigma(\Gamma, X, p)$ , it would mean there are  $h_1, \dots, h_k \in \Gamma$  with  $\|h_j\|_p < r$  for each  $j \in \{1, \dots, k\}$ , and  $h_1 \cdots h_k = g$ . For each  $j$ , choose sequences  $h_j^i \in \Gamma_i$  that converge to  $h_j$ . As the norm is continuous with respect to convergence of isometries, for  $i$  large enough one has  $\|h_j^i\|_p < r_i$  for each  $j$ .

The sequence  $g_i(h_1^i \cdots h_k^i)^{-1} \in \Gamma_i$  converges to  $g(h_1 \cdots h_k)^{-1} = e \in \Gamma$ , so its norm is less than  $r_i$  for  $i$  large enough, allowing us to write  $g_i$  as a product of  $k + 1$  elements with norm  $< r_i$ , thus a contradiction.  $\square$

**Definition 28.** Let  $G$  be a group and  $S \subset G$  a generating subset containing the identity. We say that  $S$  is a *determining set* if  $G$  has a presentation  $G = \langle S \mid R \rangle$  with  $R$  consisting of words of length 3 using as letters the elements of  $S \cup S^{-1}$ .

**Proposition 29.** For each  $M \in \mathbb{N}$ , there are only finitely many isomorphism types of groups admitting a determining set of size  $\leq M$ .

The following lemma can be found in ([17], Section 2.12).

**Lemma 30.** Let  $D > 0$ ,  $(Y, q)$  a pointed proper geodesic space and  $G \leq Iso(Y)$  a closed group of isometries with  $\text{diam}(Y/G) \leq D$ . If  $\{g \in G \mid \|g\|_q \leq 10D\}$  is not a determining set, then there is a non-trivial covering map  $\tilde{Y} \rightarrow Y$ .

### 3 Proof of main theorems

*Proof of Theorem 5.* By contradiction, assume there is a sequence  $(X_i, p_i)$  of pointed  $RCD^*(K, N)$  spaces, discrete groups  $\Gamma_i \leq Iso(X_i)$  of measure preserving isometries such that  $\text{vol}_{K, N}^*(X_i/\Gamma_i, [p_i]) \geq \nu$ , and elements  $g_i \in \Gamma_i \setminus \{Id_{X_i}\}$  with  $d_{p_i}(g_i, Id_{X_i}) \rightarrow 0$ . After taking a subsequence, we can assume the spaces  $X_i$  have dimension  $n$  for some  $n \in \mathbb{N} \cap [0, N]$ , the sequence  $(X_i/\Gamma_i, [p_i])$  converges to a pointed  $RCD^*(K, N)$  space  $(Y, q)$  of rectifiable dimension  $n$ . By Corollary 3, we can also assume that  $q$  is  $n$ -regular.

Choose  $\eta_i \rightarrow \infty$  diverging so slowly that  $(\eta_i X_i/\Gamma_i, [p_i])$  converges to  $(\mathbb{R}^n, 0)$  and  $\eta_i d_{p_i}(g_i, Id_{X_i}) \rightarrow 0$ , and set  $Y_i := \eta_i X_i$ . By Theorem 18, we can find a Reifenberg sequence  $y_i \in Y_i/\Gamma_i$  and  $\tilde{y}_i \in B(p_i, 1, Y_i)$  with  $[\tilde{y}_i] = y_i$ , which by Lemma 9 can be taken so that  $\|g_i\|_{\tilde{y}_i} \neq 0$ .

Notice that by Corollary 3, we still have  $d_{\tilde{y}_i}(g_i, Id_{Y_i}) \rightarrow 0$ . Hence we can find  $1/\lambda_i \in \sigma(\Gamma_i, Y_i, \tilde{y}_i)$  with  $\lambda_i \rightarrow \infty$ . As  $y_i$  is a Reifenberg sequence,  $(\lambda_i Y_i/\Gamma_i, y_i)$  converges to  $(\mathbb{R}^n, 0)$ , so by Lemma 26, the actions of  $\Gamma_i$  on  $\lambda_i Y_i$  converge equivariantly to the trivial group. This contradicts Proposition 27, as by construction we have  $1 \in \sigma(\Gamma_i, \lambda_i Y_i, \tilde{y}_i)$  for all  $i$ .  $\square$

*Proof of Theorem 4:* Assuming the contrary, we could find a sequence  $(X_i, p_i)$  of pointed  $RCD^*(K, N)$  spaces of diameter  $\leq D$  and collapsing volume  $\geq \nu$  whose fundamental groups are pairwise non-isomorphic. After taking a subsequence, we may assume their universal covers  $(\tilde{X}_i, \tilde{p}_i)$  converge to an  $RCD^*(K, N)$  space  $(\tilde{X}, \tilde{p})$ , and the actions of  $\pi_1(X_i)$  converge to the action of a group  $\Gamma$  in  $\tilde{X}$ .

By Theorem 5, there is  $\varepsilon > 0$  such that the elements of  $\Gamma_i$  are at pairwise  $d_{\tilde{p}_i}$ -distance  $\geq \varepsilon$ . By Lemma 30, for each  $i$  the set  $S_i := \{g \in \Gamma_i \mid \|g\|_{\tilde{p}_i} \leq 10D\}$  is determining in  $\Gamma_i$ , and by plugging  $r = 1$  in Equation 1, we get  $S_i \subset \{g \in \Gamma_i \mid d_{\tilde{p}_i}(g, Id_{\tilde{X}_i}) \leq 10D + 3\}$ . As  $(\Gamma_i, d_{\tilde{p}_i}, Id_{\tilde{X}_i})$  converges in the Gromov–Hausdorff sense to  $(\Gamma, d_{\tilde{p}}, Id_{\tilde{X}})$ , Corollary 13 implies that  $|S_i| \leq M$  for some  $M \in \mathbb{N}$ , so by Proposition 29 there are only finitely many isomorphism types in the sequence  $\{\pi_1(X_i)\}_{i \in \mathbb{N}}$ , contradicting our initial assumption.  $\square$

## References

- [1] Anderson, M.T., *Short geodesics and gravitational instantons*. J. Differential Geom. 31 (1990), no. 1, 265–275.
- [2] Bacher, K.; Sturm, K.-T., *Localization and tensorization properties of the curvature–dimension condition for metric measure spaces*. J. Funct. Anal. 259 (2010), no. 1, 28–56.
- [3] Brué, E.; Semola, D., *Constancy of the dimension for  $RCD(K,N)$  spaces via regularity of Lagrangian flows*. Comm. Pure Appl. Math. 73 (2020), no. 6, 1141–1204.
- [4] Burago, D., Burago, Y., Ivanov, D., *A course in metric geometry*. Graduate Studies in Mathematics, 33. American Mathematical Society, Providence, RI, 2001. xiv+415 pp. ISBN: 0-8218-2129-6
- [5] Cheeger, J.; Colding, T. H., *On the structure of spaces with Ricci curvature bounded below*. I. J. Differential Geom. 46 (1997), no. 3, 406–480.
- [6] Cheeger, J.; Gromoll, D., *The splitting theorem for manifolds of nonnegative Ricci curvature*. J. Differential Geometry 6 (1971/72), 119–128.
- [7] Erbar, M.; Kuwada, K.; Sturm, K.-T., *On the equivalence of the entropic curvature–dimension condition and Bochner’s inequality on metric measure spaces*. Invent. Math. 201 (2015), no. 3, 993–1071
- [8] Fukaya, K.; Yamaguchi, T., *The fundamental groups of almost non–negatively curved manifolds*. Ann. of Math. (2) 136 (1992), no. 2, 253–333.
- [9] Galaz-García, F.; Kell, M.; Mondino, A.; Sosa, G., *On quotients of spaces with Ricci curvature bounded below*. J. Funct. Anal. 275 (2018), no. 6, 1368–1446.
- [10] Gigli, N., *An overview of the proof of the splitting theorem in spaces with non-negative Ricci curvature*. Anal. Geom. Metr. Spaces 2 (2014), no. 1, 169–213.
- [11] Guijarro, L.; Santos-Rodríguez, J., *On the isometry group of  $RCD^*(K, N)$ –spaces*. Manuscripta Math. 158 (2019), no. 3–4, 441–461.
- [12] Kitabeppu, Y., *A sufficient condition to a regular set being of positive measure on  $RCD$  spaces*. Potential Anal. 51 (2019), no. 2, 179–196.
- [13] Mondino, A.; Naber, A., *Structure theory of metric measure spaces with lower Ricci curvature bounds*. J. Eur. Math. Soc. (JEMS) 21 (2019), no. 6, 1809–1854.
- [14] Mondino, A.; Wei, G., *On the universal cover and the fundamental group of an  $RCD^*(K, N)$ –space*. J. Reine Angew. Math. 753 (2019), 211–237.

- [15] Sormani, C.; Wei, G., *The covering spectrum of a compact length space*. J. Differential Geom. 67 (2004), no. 1, 35–77.
- [16] Wang, J, *RCD spaces are semi-locally simply connected*. (to appear).
- [17] Zamora, S., *Fundamental Groups and Limits of Almost Homogeneous Spaces*. (2020) arXiv preprint [arXiv:2007.01985](https://arxiv.org/abs/2007.01985).