


Max-Planck-Institut für Mathematik Bonn

Combinatorics and large genus asymptotics of the Brézin-Gross-Witten numbers

by

Jindong Guo
Paul Norbury
Di Yang
Don Zagier



Max-Planck-Institut für Mathematik
Preprint Series 2024 (38)

Date of submission: December 31, 2024

Combinatorics and large genus asymptotics of the Brézin-Gross-Witten numbers

by

Jindong Guo
Paul Norbury
Di Yang
Don Zagier

Max-Planck-Institut für Mathematik
Vivatsgasse 7
53111 Bonn
Germany

School of Mathematical Sciences
University of Science and Technology of China
230026 Hefei
P. R. China

School of Mathematics and Statistics
University of Melbourne
VIC 3010 Melbourne
Australia

International Centre for Theoretical Physics
34151 Trieste
Italy

COMBINATORICS AND LARGE GENUS ASYMPTOTICS OF THE BRÉZIN–GROSS–WITTEN NUMBERS

JINDONG GUO, PAUL NORBURY, DI YANG, DON ZAGIER

ABSTRACT. In this paper, we study combinatorial and asymptotic properties of some interesting rational numbers called the Brézin–Gross–Witten (BGW) numbers, which can be represented as the intersection numbers of psi and Theta classes on the moduli space of stable algebraic curves. In particular, we discover and prove the uniform large genus asymptotics of certain normalized BGW numbers, and give a new proof of the polynomiality phenomenon for the large genus. We also propose several new conjectures including monotonicity and integrality on the BGW numbers. Applications to the Painlevé II hierarchy and to the BGW-kappa numbers are given.

CONTENTS

1.	Introduction	1
2.	Review of general theory of BGW numbers	9
3.	Exact formulas for n -point BGW numbers	11
4.	Rational functions, asymptotics and integrality	14
5.	Uniform large genus asymptotics	18
6.	Polynomiality in large genus	23
7.	Further asymptotic formulas	27
8.	Subexponential asymptotic terms	30
9.	Application to the Painlevé II hierarchy	32
10.	Application to BGW-kappa numbers	35
	References	39

1. INTRODUCTION

In this paper, we study some interesting and important rational numbers called the *Brézin–Gross–Witten (BGW) numbers* [2, 5, 6, 14, 17, 23]. Originally, the BGW numbers were defined via matrix models [6, 23], and specifically are proportional to the Taylor coefficients with respect to the so-called Miwa variables (cf. [3, 6, 22, 23, 30]) of the logarithm of the integral

$$\int_{U_n} e^{\frac{1}{\beta} \operatorname{tr}(J^\dagger U + JU^\dagger)} dU, \tag{1}$$

where dU denotes the normalized Haar measure on the unitary group U_n , and J and J^\dagger are arbitrary $n \times n$ matrices. Later, alternative definitions and properties of the BGW numbers were given in a number of further papers (cf. [2, 5, 7, 14, 17, 29, 36, 37, 42]).

As customary in the literature, denote by $\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g^\Theta$ the BGW numbers, where $g \geq 1$ (genus), $n \geq 1$ and $d_1, \dots, d_n \geq 0$ satisfy

$$d_1 + \cdots + d_n = g - 1 \quad (2)$$

(see e.g. [2, 14, 42]).

For a long time, no topological or combinatorial meaning for the BGW numbers was known, but recently, two ways were found to define these numbers topologically. First of all, they are equal to the following integrals on the moduli space of stable curves:

$$\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g^\Theta = \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \cdots \psi_n^{d_n} \Theta_{g,n}, \quad (3)$$

as it was conjectured in [37] with later a complete proof given in [7]. Here $\overline{\mathcal{M}}_{g,n}$ denotes the Deligne–Mumford moduli space of stable algebraic curves of genus g with n distinct marked points, ψ_j denotes the first Chern class of the j th cotangent line bundle, and $\Theta_{g,n}$ denotes the Theta-class introduced by the second author of the present paper [37] (cf. [7, 29, 41]). This definition, which is the reason for the notation we use, is in exact analogy with Witten’s notation for his intersection numbers [40] and elucidate (2) simply as the degree-dimension counting. Secondly, it was proved in [42] that the BGW numbers can be given by an ELSV-like formula

$$\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g^\Theta = \frac{(-1)^{g-1+n} 2^{2g-2+n}}{\prod_{i=1}^n d_i!} \int_{\overline{\mathcal{M}}_{g,n}} \frac{\Lambda(-1)^2 \Lambda(\frac{1}{2}) \exp(\sum_{d=1}^{\infty} \frac{(-1)^{d-1} \kappa_d}{2^d d})}{\prod_{i=1}^n (1 + \frac{2d_i+1}{2} \psi_i)}, \quad (4)$$

where $\Lambda(z)$ denotes the Chern polynomial of the Hodge bundle. Two efficient algorithms of computing the BGW numbers will be reviewed in Section 2.

The first few BGW numbers are given by

$$\begin{aligned} \langle \tau_0 \rangle_1^\Theta &= \frac{1}{8}, & \langle \tau_1 \rangle_2^\Theta &= \frac{3}{128}, & \langle \tau_1^2 \rangle_3^\Theta &= \frac{63}{512}, & \langle \tau_2 \rangle_3^\Theta &= \frac{15}{1024}, \\ \langle \tau_1^3 \rangle_4^\Theta &= \frac{7221}{2048}, & \langle \tau_1 \tau_2 \rangle_4^\Theta &= \frac{8625}{32768}, & \langle \tau_3 \rangle_4^\Theta &= \frac{525}{32768}. \end{aligned} \quad (5)$$

(Here we have omitted the numbers containing τ_0 except for $\langle \tau_0 \rangle_1^\Theta$ because of equation (8) below.) From these and many further examples, we observe that the BGW numbers $\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g^\Theta$, $d_1, \dots, d_n \geq 0$, are integral away from the prime 2, and we conjecture that this is true in general. We call this the Integrality Conjecture. Furthermore, they seem to have many small factors, e.g., $\langle \tau_2 \tau_3 \rangle_6^\Theta$ equals $2^{-21} 3^2 5^2 7^3 103^1$, and $\langle \tau_2^3 \tau_3^2 \tau_4 \tau_5 \rangle_{22}^\Theta$ is divisible by $2^{-71} 3^{11} 5^2 7^2 11^2$. A precise conjecture that at least partially explains these factorizations will be given in Section 4 (see Conjecture 3).

To proceed, let us introduce the *normalized BGW numbers* $C(\mathbf{d})$ by

$$C(\mathbf{d}) := \frac{2^{2g(\mathbf{d})-1} \prod_{j=1}^n (2d_j + 1)!!}{(X(\mathbf{d}) - 1)!} \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_{g(\mathbf{d})}^\Theta, \quad (6)$$

where $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$, $g(\mathbf{d}) = |\mathbf{d}| + 1$ with $|\mathbf{d}| := d_1 + \cdots + d_n$, and

$$X(\mathbf{d}) := \sum_{j=1}^n (2d_j + 1) = 2g(\mathbf{d}) - 2 + n. \quad (7)$$

Obviously, $C(0) = 1/4$. Using a relation given in Section 2.1 (see (45)), we know that

$$C(\mathbf{d}) = C(0, \mathbf{d}). \quad (8)$$

This property would of course also be true without the factor $2^{2g(\mathbf{d})-1}$ in (6), but the normalization given here will make the asymptotic properties of the numbers nicer.

Because of (8), in the study of $C(\mathbf{d})$, it is sufficient to consider the case when d_1, \dots, d_n are all positive, in other words, when \mathbf{d} is a partition. From now on, we will usually restrict to this case. In particular, we do this in the following table, which gives the values of the normalized BGW numbers for $g = 2, \dots, 7$.

From Table 1 we observe that the normalized BGW numbers $C(\mathbf{d})$ for partitions of $g - 1$ with $2 \leq g \leq 7$ all lie between the values for the crudest and finest partitions $(g - 1)$ and (1^{g-1}) of $g - 1$. (Here we use the standard convention of writing d^m to mean that the argument d is repeated m times.) By a computer program using an algorithm given in Section 2.1, we also checked that this is true up to $g = 40$. For example, for $g = 40$, all values lie between the two numbers

$$C(39) = 0.316326705 \dots, \quad C(1^{39}) = 0.316963758 \dots. \quad (9)$$

The following conjecture states that this nesting property holds for all g .

Conjecture 1. *We have $C(g - 1) \leq C(\mathbf{d}) \leq C(1^{g-1})$ for any partition \mathbf{d} of $g - 1$.*

But in fact much more is true. We denote by $\ell(\mathbf{d})$ the length of a partition \mathbf{d} and for any fixed $g \geq 1$ we define an ordering for all partitions of $g - 1$ first by increasing length and then lexicographically for a given length, i.e., $\mathbf{d} \prec \mathbf{d}'$ if either $\ell(\mathbf{d}) < \ell(\mathbf{d}')$ or $\ell(\mathbf{d}) = \ell(\mathbf{d}')$ and $\mathbf{d}_i < \mathbf{d}'_i$, where the non-zero entries of both \mathbf{d} and \mathbf{d}' are arranged in increasing order and i is the first index for which $\mathbf{d}_i \neq \mathbf{d}'_i$. Purely by chance—simply because the calculations of tables of $C(\mathbf{d})$ up to $g = 40$ using the recursion (46) were done using the software package GP-PARI, which happens to order partitions in the way just described—we noticed empirically the following

Conjecture 2. *The function $\mathbf{d} \mapsto C(\mathbf{d})$ from partitions of $g - 1$ to \mathbb{Q} is strictly monotone increasing with respect to the above ordering for every g .*

To make this property more visible, we have given the numbers $C(\mathbf{d})$ in Table 1 both as rational numbers and as real numbers to 6 significant digits. For ease of reading, we have also listed the smallest common denominator $D = D_g$ of these numbers for each g and then tabulated the integers $DC(\mathbf{d})$ in the last column.

From the numerical tables we see a different property: the values of the normalized BGW numbers for a fixed g are very close to each other, e.g. the minimum and maximum values for $g = 40$ given in (9) differ by less than a third of a percent. In view of the nesting property, we can concentrate on only the two values $C(g - 1)$ and $C(1^{g-1})$, and indeed we can verify that these two numbers are close to each other for all g . On one hand, the value of $C(g - 1)$ is given by the explicit formula [17, 5]

$$C(g - 1) = \frac{g}{4^{2g-1}} \binom{2g-1}{g}^2 = \frac{(2g-1)!!^3}{2^{g+1}(2g-1)!g!}, \quad g \geq 1, \quad (10)$$

which by Stirling's formula has the asymptotics

$$C(g - 1) \sim \frac{1}{\pi} \left(1 - \frac{1}{4g} + \frac{1}{32g^2} + \frac{1}{128g^3} - \frac{5}{2048g^4} + \dots \right), \quad g \rightarrow \infty. \quad (11)$$

On the other hand, as we will see in Section 9, the value of $C(1^{g-1})$ is given by

$$C(1^{g-1}) = \frac{3^{g-1}(g-1)!}{(3g-2)!} y_g, \quad (12)$$

$g = 2, \quad D = 32$			
(1)	$\frac{9}{32}$	0.281250	9
$g = 3, \quad D = 1280$			
(2)	$\frac{75}{256}$	0.292969	375
(1, 1)	$\frac{189}{640}$	0.295313	378
$g = 4, \quad D = 143360$			
(3)	$\frac{1225}{4096}$	0.299072	42875
(1, 2)	$\frac{8625}{28672}$	0.300816	43125
(1, 1, 1)	$\frac{21663}{71680}$	0.302218	43326
$g = 5, \quad D = 378470400$			
(4)	$\frac{19845}{65536}$	0.302811	114604875
(1, 3)	$\frac{14945}{49152}$	0.304057	115076500
(2, 2)	$\frac{209275}{688128}$	0.304122	115101250
(1, 1, 2)	$\frac{34995}{114688}$	0.305132	115483500
(1, 1, 1, 1)	$\frac{4825971}{15769600}$	0.306030	115823304
$g = 6, \quad D = 91842150400$			
(5)	$\frac{160083}{524288}$	0.305334	28042539525
(1, 4)	$\frac{1766205}{5767168}$	0.306252	28126814625
(2, 3)	$\frac{883225}{2883584}$	0.306294	28130716250
(1, 1, 3)	$\frac{442715}{1441792}$	0.307059	28200945500
(1, 2, 2)	$\frac{6198625}{20185088}$	0.307089	28203743750
(1, 1, 1, 2)	$\frac{5768625}{18743296}$	0.307770	28266262500
(1, 1, 1, 1, 1)	$\frac{3540311739}{11480268800}$	0.308382	28322493912
$g = 7, \quad D = 37471597363200$			
(6)	$\frac{1288287}{4194304}$	0.307152	11509459436475
(1, 5)	$\frac{8392923}{27262976}$	0.307851	11535653017350
(2, 4)	$\frac{184659615}{599785472}$	0.307876	11536609447125
(3, 3)	$\frac{138495805}{449839104}$	0.307879	11536700556500
(1, 1, 4)	$\frac{92508885}{299892736}$	0.308473	11558985180750
(1, 2, 3)	$\frac{46257505}{149946368}$	0.308494	11559750499500
(2, 2, 2)	$\frac{4533499725}{14694744064}$	0.308512	11560424298750
(1, 1, 1, 3)	$\frac{23168971}{74973184}$	0.309030	11579851705800
(1, 1, 2, 2)	$\frac{2270671055}{7347372032}$	0.309045	11580422380500
(1, 1, 1, 1, 2)	$\frac{1137113661}{3673686016}$	0.309529	11598559342200
(1, 1, 1, 1, 1, 1)	$\frac{34568613873}{111522611200}$	0.309970	11615054261328

TABLE 1. Numerical data for $C(\mathbf{d})$ with $g \leq 7$

where the y_g are defined by requiring that the generating series

$$Y := \sum_{g \geq 1} y_g X^{1-3g} = \frac{1}{4X^2} + \frac{9}{4X^5} + \frac{1323}{16X^8} + \frac{108315}{16X^{11}} + \frac{62737623}{64X^{14}} + \dots \quad (13)$$

satisfies the following third-order nonlinear ODE:

$$Y''' + 6Y Y' - 2Y - X Y' = 0, \quad ' = \frac{d}{dX}. \quad (14)$$

This equation can be referred to as the *Painlevé XXXIV equation* (cf. [5, 9, 20, 26]). From (13) and (14) we know that the coefficients y_g satisfy the recursion

$$y_g = (3g-2)(3g-4)y_{g-1} + \frac{2}{g-1} \sum_{h=1}^{g-1} (3h-1)y_h y_{g-h} \quad (g \geq 2) \quad (15)$$

with the initial value $y_1 = 1/4$, and from this one can obtain the large g asymptotics

$$y_g \sim A \frac{(3g-2)!}{3^{g-1}(g-1)!} \left(1 - \frac{1}{6g} - \frac{7}{72g^2} - \frac{41}{432g^3} - \frac{1789}{10368g^4} + \dots \right), \quad (16)$$

where A is some positive constant. The determination of this constant is difficult. With the help of the relationship between the Painlevé XXXIV equation and the Painlevé II equation [9, 20] and a method given in [8], one can obtain, by employing a deep result of Fokas–Its–Kapaev–Novokshenov [21], that $A = 1/\pi$. Now using (12) we get

$$C(1^{g-1}) \sim \frac{1}{\pi} \left(1 - \frac{1}{6g} - \frac{7}{72g^2} - \frac{41}{432g^3} - \frac{1789}{10368g^4} + \dots \right), \quad g \rightarrow \infty. \quad (17)$$

We now note that Conjecture 1 together with formulas (11) and (17) implies that

$$\frac{1}{\pi} - \frac{1}{4\pi g(\mathbf{d})} + O\left(\frac{1}{g(\mathbf{d})^2}\right) \leq C(\mathbf{d}) \leq \frac{1}{\pi} - \frac{1}{6\pi g(\mathbf{d})} + O\left(\frac{1}{g(\mathbf{d})^2}\right) \quad (18)$$

as $g(\mathbf{d})$ tends to infinity. The following theorem, which will be proved in Section 5, gives a slight weakening of this, with an unspecified (though effective) O -constant.

Theorem 1. *For $\mathbf{d} \in (\mathbb{Z}_{\geq 0})^n$, we have*

$$C(\mathbf{d}) = \frac{1}{\pi} + O\left(\frac{1}{g(\mathbf{d})}\right) \quad (19)$$

uniformly as the genus $g(\mathbf{d}) = |\mathbf{d}| + 1$ goes to ∞ .

Explicitly, this theorem says that there exists an absolute constant K such that

$$\left| C(\mathbf{d}) - \frac{1}{\pi} \right| \leq \frac{K}{g(\mathbf{d})} \quad \text{for all } \mathbf{d} \in (\mathbb{Z}_{\geq 0})^n. \quad (20)$$

We note that the proof of Theorem 1 does not use the fact that the constant A in (16) equals $1/\pi$ and therefore provides a new proof of this fact, independent of [21].

The monotonicity conjecture (Conjecture 2) says in particular that the normalized numbers $C(\mathbf{d})$ with partitions \mathbf{d} of a fixed length are smaller than those of greater length, so if $I_{g,n}$ denotes the smallest interval containing the normalized BGW numbers of length n for a given g , then $I_{g,n}$ lies strictly to the left of $I_{g,n+1}$. The numerical data (up to $g = 40$) shows that much more is true. The following picture shows all 31185 normalized BGW numbers with $g = 40$. At this resolution, what one sees are just 39 intervals that look like points, meaning that the normalized BGW numbers of length n



are much closer to each other than to those of length $n - 1$ or $n + 1$. More precisely, we find that each interval $I_{g,n}$ has length $O(g^{-3})$, even though the gaps between the intervals have an average length of the order $O(g^{-2})$ (because these $g - 1$ intervals lie in an interval of total length $O(1/g)$ by Theorem 1). A conjectural statement giving a much more precise result is stated at the end of Section 6.

From the numerical data we observe for some small values of n that $C(\mathbf{d})/C(g(\mathbf{d}) - 1)$ is a rational function of d_n if $d_1, \dots, d_{n-1} \geq 1$ are fixed. In fact this is always true, as we will prove in Section 4. Equation (11) and Theorem 1 then imply that $C(\mathbf{d})$ for $\mathbf{d} = (\mathbf{d}', d_n)$ with $\mathbf{d}' = (d_1, \dots, d_{n-1})$ fixed has an asymptotic expansion of the form

$$C(\mathbf{d}) \sim \frac{1}{\pi} \sum_{k=0}^{\infty} \frac{A_k(\mathbf{d}')}{g(\mathbf{d})^k}, \quad d_n \rightarrow \infty, \quad (21)$$

where the $A_k(\mathbf{d}')$ are rational numbers with $A_0(\mathbf{d}') = 1$. Now looking at the explicit formulas for small n (see Section 4), we find that as d_1, \dots, d_{n-1} grow, the asymptotic expansion of $C(\mathbf{d})$ stabilizes to a well-defined power series in $1/g(\mathbf{d})$ depending on n . And then we discover that if we rewrite them as power series in $1/X(\mathbf{d})$, where $X(\mathbf{d}) = 2g - 2 + n$ as usual, we get a power series independent of n and beginning

$$\frac{1}{\pi} \left(1 - \frac{1}{2X} + \frac{5}{8X^2} - \frac{11}{16X^3} + \frac{83}{128X^4} - \frac{143}{256X^5} + \dots \right). \quad (22)$$

We can recognize this power series as the large- X expansion of the function

$$\gamma(X) = \frac{\Gamma\left(\frac{X}{2} + 1\right)^2}{\pi \Gamma\left(\frac{X+1}{2}\right) \Gamma\left(\frac{X+3}{2}\right)}. \quad (23)$$

We now find that the difference of $C(\mathbf{d})$ and $\gamma(X(\mathbf{d}))$ is of the order $O(X(\mathbf{d})^{-2 \min\{d_i\} - 2})$, and also that $C(\mathbf{d}) \leq \gamma(X(\mathbf{d}))$ in all cases, with strict inequality unless $n = 1$.

We also find that sometimes two normalized BGW numbers with the same g and n are extremely close to each other, a numerical example being given the two numbers

$$\begin{aligned} C(1, 18, 20) &\approx 0.3163749000332518760707893046, \\ C(1, 19, 19) &\approx 0.3163749000332518760707893073, \end{aligned} \quad (24)$$

which agree to 26 significant digits. These phenomena and many others of the same kind will be discussed in Sections 4, 6, 7 and 8.

We now turn to the second main theme of this paper, which will shed light on all aspects of the discussion so far.

In the study of Witten's intersection numbers, two of the authors [25] discovered, and stated as a conjecture, that for each k the coefficient of $1/g^k$ in the large genus asymptotics of normalized Witten's intersection numbers is a polynomial of n and the multiplicities in the arguments, and also that only the multiplicities of $0, 1, \dots, [3k/2] - 1$ are involved. In the computations for the current paper, we discovered that the same phenomenon holds also for the normalized BGW numbers $C(\mathbf{d})$, now with the k th coefficient depending on n and the multiplicities of $0, 1, \dots, [k/2] - 1$. Both of these conjectural statements were proved by Eynard *et al* [18].

There is a further discovery. We already know that for each k the coefficient $A_k(\mathbf{d}')$ in (21), $\mathbf{d}' \in (\mathbb{Z}_{\geq 1})^{n-1}$, is a polynomial a_k of n and the multiplicities. As we will see from Section 6, the DVV-type relations for BGW numbers (cf. Section 2.1) imply that the power series $\sum_{k=0}^{\infty} a_k/g^k$ is unchanged by $(g \rightarrow g-1, n \rightarrow n+2)$. So, if we write this power series in terms of X^{-1} instead of g^{-1} with $X = 2g - 2 + n$, then the coefficients are polynomials of the multiplicities of $1, \dots, [k/2] - 1$, independent of n . Namely,

$$C(\mathbf{d}) \sim \frac{1}{\pi} \sum_{k=0}^{\infty} \frac{c_k(p_1(\mathbf{d}'), \dots, p_{[k/2]-1}(\mathbf{d}'))}{X(\mathbf{d})^k}, \quad d_n \rightarrow \infty, \quad (25)$$

for some polynomials c_k , where $p_r(\mathbf{d}')$ denotes the multiplicity of r in \mathbf{d}' . The first few polynomials c_k are given by

$$c_0 = 1, \quad c_1 = -\frac{1}{2}, \quad c_2 = \frac{5}{8}, \quad c_3 = -\frac{11}{16}, \quad c_4 = \frac{83}{128} - \frac{27}{8}p_1.$$

The constant terms of c_k agree (necessarily) with the coefficients of $\gamma(X)$ (cf. (22)). This makes it very natural to introduce the renormalized BGW numbers $\widehat{C}(\mathbf{d})$ by

$$\widehat{C}(\mathbf{d}) := \frac{C(\mathbf{d})}{\gamma(X(\mathbf{d}))}, \quad \mathbf{d} \in (\mathbb{Z}_{\geq 1})^n. \quad (26)$$

Theorem 2. *For any fixed n and fixed $\mathbf{d}' \in (\mathbb{Z}_{\geq 1})^{n-1}$, the numbers $\widehat{C}(\mathbf{d})$ satisfy*

$$\widehat{C}(\mathbf{d}) \sim \sum_{k=0}^{\infty} \frac{\widehat{c}_k(p_1(\mathbf{d}'), p_2(\mathbf{d}'), \dots)}{X(\mathbf{d})^k}, \quad d_n \rightarrow \infty, \quad (27)$$

where $\mathbf{d} = (\mathbf{d}', d_n)$, \widehat{c}_k are universal polynomials of p_1, p_2, \dots having rational coefficients, with $\widehat{c}_0 \equiv 1$ and $\widehat{c}_k|_{p_b=0} = 0$ ($k \geq 1$). Moreover, under the degree assignments

$$\deg p_d = 2d + 1 \quad (d \geq 1), \quad (28)$$

the polynomials \widehat{c}_k , $k \geq 1$, satisfy the degree estimates

$$\deg \widehat{c}_k \leq k - 1. \quad (29)$$

We will give a proof of this theorem, independent of [18], in Section 6.

From (29), we know that \widehat{c}_k does not depend on p_d with $d \geq (k-1)/2$. In particular, $\widehat{c}_k = 0$ for $k = 1, 2, 3$. We list a few more \widehat{c}_k below:

$$\widehat{c}_4 = -\frac{27}{8}p_1, \quad \widehat{c}_5 = -\frac{27}{4}p_1, \quad \widehat{c}_6 = -\frac{45}{4}p_1 - \frac{1125}{16}p_2.$$

Several more coefficients for both c_k and \widehat{c}_k are given in Table 2 of Section 6. We are also able to give explicit expressions for all $\widehat{c}_k|_{p_b=\delta_{b,d}}$; see Section 7.

By Theorem 2 and the above-mentioned result of Eynard *et al* [18, Theorem 4.3], we also know that for any fixed $L \geq 1$ and fixed $n \geq 1$, and for $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 1})^n$,

$$\widehat{C}(\mathbf{d}) = \sum_{k=0}^{L-1} \frac{\widehat{c}_k(p_1(\mathbf{d}), p_2(\mathbf{d}), \dots)}{X(\mathbf{d})^k} + O\left(\frac{1}{X(\mathbf{d})^L}\right), \quad g(\mathbf{d}) \rightarrow \infty, \quad (30)$$

where the implied O-constant only depends on n and L .

Based on an algorithm given in [15, 16] (see equations (52)–(53) of Section 2.2) we have computed $\widehat{C}(d^n)$ for genera far bigger than 40, from which we also see the phenomenon that $\widehat{C}(d^n)$ rapidly tends to 1 as $d \rightarrow \infty$. (For example, $1 - \widehat{C}(100^{10})$ is

roughly 1.8×10^{-285} .) This together with Theorem 2 leads us to the discovery of the conjectural asymptotic formula

$$1 - \widehat{C}(d^n) \sim \left(\frac{1}{2}\right)^{\delta_{n,2}} \sqrt{\frac{4(n-1)}{\pi n d}} \left(\frac{(n-1)^{n-1}}{n^n}\right)^{2d+1}, \quad \text{as } d \rightarrow \infty. \quad (31)$$

More details and generalizations of this will be discussed in Section 8.

We end this section by presenting two applications of Theorem 1.

The first one is an application for the Painlevé II hierarchy. Following [4, 9, 12, 34], define a sequence of polynomials $m_d = m_d(u_0, u_1, u_2, \dots, u_{2d})$, $d \geq 0$, by means of generating series as follows:

$$b \partial^2(b) - \frac{1}{2} \partial(b)^2 - 2(\lambda - 2u_0) b^2 = -2\lambda, \quad (32)$$

where $\partial := \sum_{i \geq 0} u_{i+1} \partial / \partial u_i$, and

$$b(\lambda) = 1 + \sum_{d \geq 0} \frac{(2d+1)!! m_d}{\lambda^{d+1}}. \quad (33)$$

The first few m_d are $m_0 = u_0$, $m_1 = \frac{1}{2}u_0^2 + \frac{1}{12}u_2$. By the *Painlevé II hierarchy* we mean the following family of ODEs:

$$2^{2d-1}(2d-1)!! (\partial_X + 2V) \left(m_{d-1} \left(\frac{V_X - V^2}{2}, \frac{(V_X - V^2)_X}{2}, \dots \right) \right) - VX - \alpha_d = 0, \quad (34)$$

where $d \geq 1$ and α_d are constants. We will focus on the case when $\alpha_d = 1/2$, $d \geq 1$. In this case, it can be shown that, for each $d \geq 1$, there exists a unique formal solution $V(X)$ to (34) of the form

$$V(X) = - \sum_{n=0}^{\infty} \frac{v_{d,n}}{X^{(2d+1)n+1}}, \quad v_{d,n} \in \mathbb{C}, \quad v_{d,0} = \frac{1}{2}. \quad (35)$$

In Section 9 we will use Theorem 1 to prove the following theorem.

Theorem 3. *For each $d \geq 1$, the coefficients $v_{d,n}$ of the formal solution $V(X)$ to the d th member of the Painlevé II hierarchy have the following asymptotics:*

$$v_{d,n} \sim \frac{1}{\pi} \frac{((2d+1)n-1)!}{(2d+1)^{n-1} (n-1)!}, \quad n \rightarrow \infty. \quad (36)$$

We note that for the particular case when $d = 1$ the above theorem was proved in [21] by using a deep Riemann–Hilbert analysis, and we now achieve a new proof. As far as we know, the cases with $d \geq 2$ are new.

The second application that we will present is to use Theorem 1 to study the large genus asymptotics of the more general integrals, which we call the *BGW-kappa numbers*, where the Theta-class is coupled with powers of κ_1 -class as well as psi-classes

$$\int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \cdots \psi_n^{d_n} \Theta_{g,n} \kappa_1^m =: \langle \kappa_1^m \prod_{j=1}^n \tau_{d_j} \rangle_g^\Theta. \quad (37)$$

By the degree-dimension matching, these numbers vanish unless $m + d_1 + \cdots + d_n = g - 1$. A small table of these BGW-kappa numbers is provided in Section 10.

Like the numbers $C(\mathbf{d})$, we introduce the *normalized BGW-kappa numbers* as follows:

$$C(m; \mathbf{d}) := \frac{3^m 2^{2g-1} \prod_{j=1}^n (2d_j + 1)!!}{(X(m; \mathbf{d}) - 1)!} \langle \kappa_1^m \tau_{d_1} \cdots \tau_{d_n} \rangle_g^\ominus, \quad X(m; \mathbf{d}) := X(\mathbf{d}) + 3m. \quad (38)$$

Obviously, $C(0; \mathbf{d}) = C(\mathbf{d})$. In Section 10 we will use Theorem 1 to prove the following

Proposition 1. *For any fixed $m \geq 0$, there exists a constant $K(m)$ such that*

$$\left| C(m; \mathbf{d}) - \frac{1}{\pi} \right| \leq \frac{K(m)}{g(m; \mathbf{d})} \quad \text{for all } \mathbf{d} \in (\mathbb{Z}_{\geq 0})^n, \quad (39)$$

where $g(m; \mathbf{d}) = g(\mathbf{d}) + m = |\mathbf{d}| + m + 1$.

We also give in Section 10 an application of Theorem 2 to BGW-kappa numbers.

The paper is organized as follows. In Section 2 we review a recursive definition of BGW numbers as well as an explicit formula for their n -point generating series. In Section 3 we give closed formulas for BGW numbers. In Section 4 we present several results on structures for BGW numbers. In Section 5 we prove Theorem 1, and in Section 6 we prove Theorem 2. Further asymptotic formulas and conjectural subexponential asymptotics are given in Sections 7, 8, respectively. In Sections 9, 10 we present applications of the main theorems.

Acknowledgements The work was partially supported by NSFC No. 12371254, the CAS No. YSBR-032, the National Key R and D Program of China 2020YFA0713100, and the China Scholarship Council. Parts of the work of J.G., P.N. and D.Y. were done during their visits in MPIM, Bonn; they thank MPIM for excellent working conditions.

2. REVIEW OF GENERAL THEORY OF BGW NUMBERS

The definitions for BGW numbers given in Section 1 are not directly calculable since the integrals over the unitary group or moduli space are not algorithmically defined. In this section we review two algorithms that can be used to effectively compute the BGW numbers and that can and have been implemented on a computer.

2.1. Recursive definition of the BGW numbers. It is known from [14, 36] (cf. also [2, 7, 37]) that the partition function Z of the BGW numbers, defined by

$$Z = \exp \left(\sum_{g \geq 1} \sum_{n \geq 1} \frac{1}{n!} \sum_{\substack{d_1, \dots, d_n \geq 0 \\ d_1 + \dots + d_n = g-1}} \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g^\ominus t_{d_1} \cdots t_{d_n} \right) \quad (40)$$

(cf. (2)), is a particular tau-function for the celebrated KdV hierarchy. In particular, $u := \partial^2 \log Z / \partial t_0^2$ satisfies the KdV hierarchy

$$u_{t_d} = \partial_x (m_d(u, u_x, u_{xx}, \dots, u_{2dx})), \quad d \geq 0, \quad (41)$$

with $t_0 \equiv x$, where m_d is defined in (32), (33). It is also known that Z satisfies the following infinite set of linear equations called the *Virasoro constraints*:

$$L_m Z = 0, \quad m \geq 0, \quad (42)$$

where L_m , $m \geq 0$, are operators defined by

$$\begin{aligned} L_m &:= -(2m+1)!! \frac{\partial}{\partial t_m} + \sum_{d \geq 0} \frac{(2d+2m+1)!!}{(2d-1)!!} t_d \frac{\partial}{\partial t_{d+m}} \\ &+ \frac{1}{2} \sum_{a+b=m-1} (2a+1)!! (2b+1)!! \frac{\partial^2}{\partial t_a \partial t_b} + \frac{1}{8} \delta_{m,0}. \end{aligned} \quad (43)$$

See [2, 14, 22, 36] (cf. also [3, 5, 7, 17, 37]).

For $n \geq 1$, $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$, it is convenient to denote

$$B(\mathbf{d}) := \langle \tau_{d_1} \dots \tau_{d_n} \rangle_{g(\mathbf{d})}^{\Theta} \prod_{j=1}^n (2d_j + 1)!!, \quad (44)$$

where we recall that $g(\mathbf{d}) = |\mathbf{d}| + 1$. Using the $m = 0$ case of (42), we obtain

$$B(0, \mathbf{d}) = (2g(\mathbf{d}) + n - 2) B(\mathbf{d}), \quad 2g(\mathbf{d}) - 2 + n > 0. \quad (45)$$

In general, a recursion for the BGW numbers that is equivalent to the Virasoro constraints was derived in [14] by Do and the second author of the present paper:

$$\begin{aligned} B(d, \mathbf{d}) &= \sum_{i=1}^n (2d_i + 1) B(d_1, \dots, d_i + d, \dots, d_n) \\ &+ \frac{1}{2} \sum_{a+b=d-1} \left(B(a, b, \mathbf{d}) + \sum_{I \sqcup J = \{1, \dots, n\}} B(a, \{d_i\}_{i \in I}) B(b, \{d_j\}_{j \in J}) \right), \end{aligned} \quad (46)$$

where $d \geq 0$. We refer to (46) as the DVV-type relation (here ‘‘DVV’’ stands for Dijkgraaf–Verlinde–Verlinde), because it is analogous to the DVV relation for Witten’s intersection numbers [13]. It is also closely related to the topological recursion [14, 19]. Note that originally the DVV-type relation for BGW numbers was written in another normalization, denoted $U_{g,n}(2d_1 + 1, \dots, 2d_n + 1)$ [14], which is related to $B(\mathbf{d})$ by

$$U_{g,n}(2d_1 + 1, \dots, 2d_n + 1) = B(\mathbf{d}) / \prod_{j=1}^n (2d_j + 1).$$

By induction on the sum $\sum_{i=1}^n (2d_i + 1) = 2g(\mathbf{d}) + n - 2$, we know that the numbers $B(\mathbf{d})$ can be uniquely determined by (46) along with the initial value

$$B(0) = \frac{1}{8} \quad (47)$$

(see (5), (44)). However, it is not at all obvious from (46) that B is symmetric in its arguments. In other words, if we force this symmetry by defining B as a function on unordered multisets, then (46) is an overdetermined system because we can choose any of the $n + 1$ arguments of $B(d_1, \dots, d_{n+1})$ as the ‘‘ d ’’ of (46) and it is non-trivial that the right-hand side will be independent of this choice.

Using the DVV-type relation (46), we can in principle compute the numbers $C(\mathbf{d})$ for partitions $\mathbf{d} = (d_1, \dots, d_n)$ of $g - 1$ for any g , and we have done so for all g up to 40.

2.2. Explicit formulas of n -point generating series. Following [4, 5, 17], consider the following n -point generating series of the BGW numbers:

$$F_n(\lambda_1, \dots, \lambda_n) = \sum_{d_1, \dots, d_n \geq 0} \frac{B(d_1, \dots, d_n)}{\lambda_1^{d_1+1} \dots \lambda_n^{d_n+1}}. \quad (48)$$

Using the matrix-resolvent method [4], an explicit formula for $F_n(\lambda_1, \dots, \lambda_n)$ was obtained in [17] (see [5] for a different proof)

$$F_1(\lambda) = \sum_{d \geq 0} \frac{(2d+1)!!^3}{8^{d+1}(d+1)!(2d+1)} \frac{1}{\lambda^{d+1}}, \quad (49)$$

$$F_n(\lambda_1, \dots, \lambda_n) = -\frac{1}{n} \sum_{\sigma \in S_n} \frac{\text{tr}(M(\lambda_{\sigma(1)}) \cdots M(\lambda_{\sigma(n)}))}{\prod_{i=1}^n (\lambda_{\sigma(i+1)} - \lambda_{\sigma(i)})} - \delta_{n,2} \frac{\lambda_1 + \lambda_2}{(\lambda_1 - \lambda_2)^2}, \quad n \geq 2, \quad (50)$$

where

$$M(\lambda) := \sum_{k \geq -1} \left(\frac{(2k-1)!!}{2^k} \right)^3 \begin{pmatrix} k(k+1) & k+1 \\ -\frac{8k^3+12k^2+4k+1}{8} & -k(k+1) \end{pmatrix} \frac{\lambda^{-k}}{(k+1)!}, \quad (51)$$

with the usual conventions $(-1)!! = 1$ and $(-3)!! = -1$.

There is a useful variant of formulas like (50) given by Dubrovin and the third author of the present paper (see [15, Proposition 3.2.3]). A special case of the variant gives the numbers $\langle \tau_{d_1} \dots \tau_{d_n} \rangle_g^\ominus$ efficiently when all but at most two of the d 's are equal. Indeed, define a sequence of traceless 2×2 matrix-valued functions $M_{m,d}(\lambda)$ ($m \geq 0$) by setting $M_{0,d}(\lambda) = M(\lambda)$ as above (independent of d) and then inductively defining

$$M_{m,d}(\lambda) = \frac{1}{m} \sum_{i+j=m-1} [(\lambda^d M_{i,d}(\lambda))^- , M_{j,d}(\lambda)] \quad (52)$$

for $m \geq 1$, where $A(\lambda)^-$ denotes the sum of the terms with strictly negative exponents of a Laurent series $A(\lambda)$ in $1/\lambda$. Then we have the following generating function formula:

$$\frac{(\lambda_1 - \lambda_2)^2}{m!} \sum_{a,b \geq 0} \frac{B(a,b,d^m)}{\lambda_1^{a+1} \lambda_2^{b+1}} = \sum_{k=0}^m \text{tr}(M_{k,d}(\lambda_1) M_{m-k,d}(\lambda_2)) - \delta_{m,0}. \quad (53)$$

This formula allows us to calculate all of the numbers $C(a,b,d^{m-2})$, and in particular the numbers $C(d^m)$, quite efficiently even when the genus is large, in which case the recursive formula (46) would be useless because it requires one to have computed and stored the BGW numbers for all smaller genera. Using it, we computed the rational numbers $C(d^m)$ for $1 \leq d \leq 100$ and $1 \leq m \leq 10$. This computation took about 20 hours on a relatively fast desktop computer, which sounds like a lot until one realizes that, for example, the numerator and denominator of the rational number $C(100^{10})$ each has 3020 digits.

3. EXACT FORMULAS FOR n -POINT BGW NUMBERS

For $n \geq 2$, expanding the right-hand side of (50) in the region $|\lambda_1| > \dots > |\lambda_n| \gg 0$, one gets a formula for n -point BGW numbers which is similar to a formula for Witten's

intersection numbers given in [25]. For $\sigma \in S_n$, introduce the notation

$$S_\sigma^+ = \{1 \leq r \leq n \mid \sigma(r+1) > \sigma(r)\}, \quad S_\sigma^- = \{1 \leq r \leq n \mid \sigma(r+1) < \sigma(r)\}.$$

Here σ is considered to be cyclic, so that we have the convention $\sigma(n+1) = \sigma(1)$. For $k_1, \dots, k_n \geq -1$, introduce the notation

$$a_{k_1, \dots, k_n} := \text{tr}(A_{k_1} \cdots A_{k_n}), \quad (54)$$

where $A_k := f(k)R(k)$, $k \geq -1$, with

$$f(k) := \frac{(2k-1)!!^3}{2^{3k}(k+1)!}, \quad R(k) := \begin{pmatrix} k(k+1) & k+1 \\ -\frac{8k^3+12k^2+4k+1}{8} & -k(k+1) \end{pmatrix}, \quad (55)$$

and we make the convention that $a_{k_1, \dots, k_n} = 0$ if any of the k_i is less than or equal to -2 . Note that A_k is just the coefficient of λ^{-k} in the Laurent series $M(\lambda)$ defined in (51).

Proposition 2. For $n \geq 2$ and $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$, we have

$$B(\mathbf{d}) = \sum_{\substack{\sigma \in S_n \\ \sigma(n)=n}} (-1)^{|S_\sigma^+|+1} \sum_{\substack{J \in (\mathbb{Z} + \frac{1}{2})^n \\ \{1 \leq q \leq n \mid J_q > 0\} = S_\sigma^+}} a_{d_{\sigma(1)+J_1-J_n, \dots, d_{\sigma(n)+J_n-J_{n-1}}}. \quad (56)$$

Proof. In the region $|\lambda_1| > \cdots > |\lambda_n| \gg 0$, we have the Laurent expansion

$$\prod_{q=1}^n \frac{1}{\lambda_{\sigma(q+1)} - \lambda_{\sigma(q)}} = (-1)^{|S_\sigma^+|} \sum_{j_1, \dots, j_n \geq 0} \prod_{q=1}^n \lambda_{\sigma(q)}^{J_{\sigma, q}(j_q) - J_{\sigma, q-1}(j_{q-1}) - 1}, \quad (57)$$

where for $q = 1, \dots, n$,

$$J_{\sigma, q}(j) := \begin{cases} -j-1, & \sigma(q) < \sigma(q+1), \\ j, & \sigma(q) > \sigma(q+1). \end{cases} \quad (58)$$

Expanding both sides of (50) and using (57), we get

$$B(\mathbf{d}) = \sum_{\substack{\sigma \in S_n \\ \sigma(n)=n}} (-1)^{|S_\sigma^+|+1} \sum_{j \in (\mathbb{Z}_{\geq 0})^n} a_{d_{\sigma(1)+J_{\sigma, 1}(j_1)-J_{\sigma, n}(j_n), \dots, d_{\sigma(n)+J_{\sigma, n}(j_n)-J_{\sigma, n-1}(j_{n-1})}}. \quad (59)$$

For each $\sigma \in S_n$ with $\sigma(n) = n$, by changing the variable $J_q = \frac{1}{2} + J_{\sigma, q}(j_q)$, $q = 1, \dots, n$, we obtain formula (56). \square

Proposition 2 could be rewritten in a more elegant way as follows:

Proposition 3. For $n \geq 2$ and $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$, we have

$$B(\mathbf{d}) = \sum_{\substack{\sigma \in S_n \\ \sigma(n)=n}} (-1)^{|S_\sigma^+|+1} \sum_{\substack{k_1, \dots, k_n \geq -1 \\ k_1 + \dots + k_n = d_1 + \dots + d_n}} a_{k_1, \dots, k_n} \omega_{\mathbf{d}, \sigma, \mathbf{k}}, \quad (60)$$

where the numbers $\omega_{\mathbf{d}, \sigma, \mathbf{k}}$ have the following explicit expression

$$\omega_{\mathbf{d}, \sigma, \mathbf{k}} = \max \left\{ 0, \min_{r \in S_\sigma^+} \left\{ \sum_{q=1}^r (d_{\sigma(q)} - k_q) \right\} + \min_{r \in S_\sigma^-} \left\{ \sum_{q=1}^r (k_q - d_{\sigma(q)}) \right\} \right\}. \quad (61)$$

Proof. For each $\sigma \in S_n$ with $\sigma(n) = n$ we change the variable $k_q = d_{\sigma(q)} + J_{\sigma,q}(j_q) - J_{\sigma,q-1}(j_{q-1})$, $q = 1, \dots, n$, in formula (59) and we get

$$B(\mathbf{d}) = \sum_{\substack{\sigma \in S_n \\ \sigma(n)=n}} (-1)^{|S_\sigma^+|+1} \sum_{\substack{k_1, \dots, k_n \geq -1 \\ k_1 + \dots + k_n = d_1 + \dots + d_n}} a_{k_1, \dots, k_n} \omega_{\mathbf{d}, \sigma, \mathbf{k}}, \quad (62)$$

where the numbers $\omega_{\mathbf{d}, \sigma, \mathbf{k}}$ are the number of solutions $\underline{j} \in (\mathbb{Z}^{\geq 0})^n$ for the linear equations:

$$d_{\sigma(q)} + J_{\sigma,q}(j_q) - J_{\sigma,q-1}(j_{q-1}) = k_q, \quad q = 1, \dots, n. \quad (63)$$

Now it suffices to prove that these numbers $\omega_{\mathbf{d}, \sigma, \mathbf{k}}$ have the expression (61). Indeed, by using (58), equations (63) can be solved in terms of j_n by

$$j_r = \begin{cases} -j_n - 1 + \sum_{q=1}^r (d_{\sigma(q)} - k_q), & r \in S_\sigma^+, \\ j_n + \sum_{q=1}^r (k_q - d_{\sigma(q)}), & r \in S_\sigma^-. \end{cases} \quad (64)$$

Here we use $\sigma(n) = n$ to obtain $J_{\sigma,n}(j_n) = j_n$. Therefore, $\omega_{\mathbf{d}, \sigma, \mathbf{k}}$ is equal to the number of $j_n \in \mathbb{Z}_{\geq 0}$ such that $j_r \geq 0$ in (64) for all $r = 1, \dots, n-1$, and hence is equal to the right-hand side of (61). This finishes the proof. \square

We note that when $n \geq 3$ and some of d_j are less than zero, then formula (56) or formula (60) still holds true (where both sides are 0), since both sides are the coefficients of $\lambda_1^{-d_1-1} \dots \lambda_n^{-d_n-1}$ in the power series $F_n(\lambda_1, \dots, \lambda_n)$ defined in (48).

Let us give some examples for Proposition 3. First we introduce the following notation: for $n \geq 1$ and $\mathbf{e} \in \mathbb{Z}^n$, write

$$M(\mathbf{e}) = \max\{0, \min_{1 \leq i \leq n} \{e_i\}\} \quad (65)$$

which can be written in terms of a generating function by

$$\sum_{e_1, \dots, e_n \geq 0} x_1^{e_1-1} \dots x_n^{e_n-1} M(e_1, \dots, e_n) = \frac{1}{(1-x_1 \dots x_n) \prod_{i=1}^n (1-x_i)}. \quad (66)$$

For $n = 2$, Proposition 3 reads that

$$B(d_1, d_2) = \sum_{k_1+k_2=d_1+d_2} M(d_1 - k_1) a_{k_1, k_2}, \quad (67)$$

where a_{k_1, k_2} can be explicitly given as follows:

$$a_{k_1, k_2} = -f(k_1)f(k_2) \left(((k_1 - k_2)^2 + \frac{k_1+k_2}{2})(k_1+1)(k_2+1) - \frac{k_1+k_2+2}{8} \right),$$

with $f(k)$ defined in (55). Actually, we also have a simpler formula for $B(d_1, d_2)$:

$$B(d_1, d_2) = \frac{1}{g} \sum_{h=0}^{d_1} (g-2h) F_h F_{g-h}, \quad \text{where } F_h := \frac{(2h-1)!!^3}{2^{3h} h!}. \quad (68)$$

The equivalence of (67) and (68) can be proved as in [24]. Since $\sum_{h=0}^g (g-2h) F_h F_{g-h}$ vanishes by antisymmetry, we see the RHS of (68) is indeed symmetric in d_1 and d_2 . We also note that formula (68) also holds for $d_1 = -1$, $d_2 \geq 1$ and $d_2 = -1$, $d_1 \geq 1$ (where both sides are 0). With the normalization $C(\mathbf{d})$ the above formula becomes

$$C(d_1, d_2) = \frac{2^{2g}}{(2g)!} \sum_{h=0}^{d_1} (g-2h) F_h F_{g-h}. \quad (69)$$

For $n = 3$, Proposition 3 reads

$$B(d_1, d_2, d_3) = -2 \sum_{k_1+k_2+k_3=d_1+d_2+d_3} a_{k_1, k_2, k_3} M(d_1 - k_1, d_1 + d_2 - k_1 - k_2), \quad (70)$$

where a_{k_1, k_2, k_3} can be given more explicitly by

$$a_{k_1, k_2, k_3} = f(k_1) f(k_2) f(k_3) (k_1 - k_2) (k_2 - k_3) (k_3 - k_1) \left((k_1 + 1)(k_2 + 1)(k_3 + 1) + \frac{1}{8} \right). \quad (71)$$

For $n = 4$, Proposition 3 reads that

$$\begin{aligned} B(d_1, d_2, d_3, d_4) &= 2 \sum_{k_1+k_2+k_3+k_4=d_1+d_2+d_3+d_4} a_{k_1, k_2, k_3, k_4} \\ &\times \left(M(d_1 - k_1, d_1 + d_2 - k_1 - k_2, k_4 - d_4) \right. \\ &\quad - M(d_1 - k_2, d_1 + d_2 - k_2 - k_3, d_1 + d_3 - k_1 - k_2, k_4 - d_4) \\ &\quad \left. - M(d_1 - k_1, d_2 - k_3, k_2 - d_3, k_4 - d_4) \right), \end{aligned} \quad (72)$$

where we have used the fact that $a_{k_1, k_2, k_3, k_4} = a_{k_2, k_3, k_4, k_1} = a_{k_1, k_4, k_3, k_2}$.

4. RATIONAL FUNCTIONS, ASYMPTOTICS AND INTEGRALITY

In this section we consider the numbers $C(\lambda, g - 1 - |\lambda|)$, where λ is a given fixed partition and we allow g to vary.

When $|\lambda| = 0$, the formula for $C(g - 1)$ is known; see (10). Based on the DVV-type relation (46), we can find that for $|\lambda| = 1, 2, 3$ the quotient of $C(\lambda, g - 1 - |\lambda|)$ by $C(g - 1)$ is a rational function of g . Explicitly,

$$\begin{aligned} \frac{C(1, g - 2)}{C(g - 1)} &= \frac{g - 1}{(2g - 1)^3} Q_1(g), \\ \frac{C(\lambda, g - 3)}{C(g - 1)} &= \frac{(g - 1)(g - 2)}{(2g - 1)^3(2g - 3)^3} Q_\lambda(g), \quad |\lambda| = 2, \\ \frac{C(\lambda, g - 4)}{C(g - 1)} &= \frac{(g - 1)(g - 2)(g - 3)}{(2g + 1)(2g - 1)^3(2g - 3)^3(2g - 5)^3} Q_\lambda(g), \quad |\lambda| = 3, \end{aligned}$$

where

$$\begin{aligned} Q_1(g) &= 8g^2 - 4g + 3, \\ Q_{1,1}(g) &= 64g^4 - 192g^3 + 224g^2 - 144g + 117, \\ Q_2(g) &= 64g^4 - 192g^3 + 216g^2 - 108g + \frac{135}{2}, \\ Q_{1,1,1}(g) &= 1024g^7 - 7168g^6 + 19072g^5 - 24256g^4 + 18832g^3 - 15520g^2 + 11418g + 14823, \\ Q_{1,2}(g) &= 1024g^7 - 7168g^6 + 18944g^5 - 23040g^4 + 14056g^3 - 6272g^2 + 5411g + \frac{16365}{2}, \\ Q_3(g) &= 1024g^7 - 7168g^6 + 18816g^5 - 21952g^4 + 10360g^3 + 1125g + \frac{7875}{2}. \end{aligned}$$

The general situation is described in the following two propositions.

Proposition 4. *For $n \geq 1$ and for a given fixed partition $\lambda = (\lambda_1, \dots, \lambda_{n-1})$, we have*

$$\frac{C(\lambda, g - 1 - |\lambda|)}{C(g - 1)} = \frac{\tilde{Q}_\lambda(g)}{(2g - 2|\lambda| - 2)_{2|\lambda|+n} \prod_{i=1}^{|\lambda|} (2g - 2i + 1)^2}, \quad (73)$$

where $\tilde{Q}_\lambda(g) \in \mathbb{Z}[1/2][g]$ is a polynomial, and $(a)_b := a(a+1)\cdots(a+b-1)$ denotes the ordinary Pochhammer symbol.

Proof. Write the left-hand side of (73) as $\tilde{\Phi}_\lambda(g)$. In terms of $\tilde{\Phi}_\lambda(g)$, recursion (46) reads

$$\begin{aligned} \tilde{\Phi}_{d,\lambda}(g) &= \sum_{j=1}^{n-1} \frac{2\lambda_j + 1}{2g + n - 2} \tilde{\Phi}_{\lambda_1, \dots, \lambda_j + d, \dots, \lambda_{n-1}}(g) + \left(1 - \frac{2d + 2|\lambda| + n - 1}{2g + n - 2}\right) \tilde{\Phi}_\lambda(g) \\ &+ \sum_{\substack{a+b=d-1 \\ a,b \geq 0}} \left[\frac{8g(g-1)}{(2g+n-2)(2g-1)^2} \tilde{\Phi}_{a,b,\lambda}(g-1) \right. \\ &+ \sum_{I \sqcup J = \{1, \dots, n-1\}} 2(X(a, \lambda_I) - 1)! C(a + |\lambda_I|) \tilde{\Phi}_{a, \lambda_I}(a + 1 + |\lambda_I|) \tilde{\Phi}_{b, \lambda_J}(g - 1 - a - |\lambda_I|) \\ &\left. \times \frac{(g - a - |\lambda_I|)_{a+1+|\lambda_I|} (g - 1 - a - |\lambda_I|)_{a+1+|\lambda_I|}}{\left((2g + n - 2 - X(a, \lambda_I))_{X(a, \lambda_I)+1} \left(g - \frac{1}{2} - a - |\lambda_I|\right)_{a+1+|\lambda_I|}\right)^2} \right]. \end{aligned} \quad (74)$$

Note that $\tilde{\Phi}_\emptyset(g) = 1$. Then Proposition 4 is proved by induction in $|\lambda|$. Indeed, using the induction assumption, one can show that each term in the right-hand side of (74) multiplying the factor $(2g - 2|\lambda| - 2d - 2)_{2|\lambda|+2d+n+1} \prod_{i=1}^{|\lambda|+d} (2g - 2i + 1)^2$ is a polynomial in $\mathbb{Z}[1/2][g]$. This finishes the proof. \square

Proposition 5. For $n \geq 1$ and for a given fixed partition $\lambda = (\lambda_1, \dots, \lambda_{n-1})$, we have

$$\frac{C(\lambda, g - 1 - |\lambda|)}{C(g - 1)} = \frac{Q_\lambda(g)}{\prod_{j=1}^{n-3} (2g + j)} \prod_{i=1}^{|\lambda|} \frac{2g - 2i}{(2g - 2i + 1)^3}, \quad (75)$$

where $Q_\lambda(g) \in \mathbb{Z}[1/2][g]$ are polynomials.

Proof. For $n = 1$ we have $Q_\emptyset(g) = 1$. Assume that $n \geq 3$ and n is odd. Write $\mathbf{d} = (\lambda, g - 1 - |\lambda|)$. From Proposition 3 we know that

$$\frac{(2g + n - 3)!}{2^{2g-1}} C(\mathbf{d}) = \sum_{\substack{\sigma \in S_n \\ \sigma(n)=n}} (-1)^{|S_\sigma^+|+1} \sum_{\substack{k_1, \dots, k_n \geq -1 \\ k_1 + \dots + k_n = d_1 + \dots + d_n}} a_{k_1, \dots, k_n} \omega_{\mathbf{d}, \sigma, \mathbf{k}}, \quad (76)$$

where a_{k_1, \dots, k_n} is defined in (54), and $\omega_{\mathbf{d}, \sigma, \mathbf{k}}$ is defined in (61). By the definition (54) of a_{k_1, \dots, k_n} , we know that

$$\frac{8^{k_1 + \dots + k_n} (k_n + 1)! a_{k_1, \dots, k_n}}{(2k_n - 1)!!^3} \in \mathbb{Z}[1/2][k_n]. \quad (77)$$

Now we claim that all nonzero summands in the RHS of (76) correspond to

$$g - |\lambda| \leq k_n \leq g + \frac{n-3}{2}. \quad (78)$$

Indeed, one can show that $k_n < g - |\lambda|$ implies $\omega_{\mathbf{d}, \sigma, \mathbf{k}} = 0$, and that $k_n > g + \frac{n-3}{2}$ implies $a_{k_1, \dots, k_n} = 0$. Therefore, we obtain from (77) and (78) that

$$\frac{8^{g-1} (g + \frac{n-1}{2})!}{(2g - 2|\lambda| - 1)!!^3} \sum_{\substack{\sigma \in S_n \\ \sigma(n)=n}} (-1)^{|S_\sigma^+|+1} \sum_{\substack{k_1, \dots, k_n \geq -1 \\ k_1 + \dots + k_n = d_1 + \dots + d_n}} a_{k_1, \dots, k_n} \omega_{\mathbf{d}, \sigma, \mathbf{k}} \in \mathbb{Z}[1/2][g]. \quad (79)$$

Moreover, we notice that the polynomial (79) has zeros at $g = |\lambda|, |\lambda| - 1, \dots, -\frac{n-1}{2}$, (since at these points, $g - 1 - |\lambda|$ is a negative integer and the polynomials (79) equals 0 by the discussion in Section 3), so we obtain that

$$\frac{(2g+n-3)! 8^{g-1} (g-|\lambda|+1)!}{2^{2g-1} (2g-2|\lambda|-1)!!^3} C(\mathbf{d}) \in \mathbb{Z}[1/2][g]. \quad (80)$$

Using (80) and the one-point formula (10), we deduce (75) for $n \geq 3$ odd. For n even and $n \geq 4$, the proof is similar and for $n = 2$, the proof is based on the fact that $C(d_1, d_2) = C(0, d_1, d_2)$. \square

We note that the above two propositions are analogous to results of Liu–Xu [33] on Witten’s intersection numbers, and that in [25] we used the matrix-resolvent formula to prove rationality for Witten’s intersection numbers.

We make one further remark. In the formulas given before Proposition 4 we see that the polynomials $Q_\lambda(g)$ occurring for different λ of the same length m are very close for g large, e.g., the three polynomials $Q_\lambda(g)$ for $m = 3$ all start $1024g^7 - 7168g^6$, and even their g^5 coefficients are near each other. We will return to this point in more detail later.

It will be convenient to write the polynomials $Q_\lambda(g)$ as polynomials of the variable X , i.e., $Q_\lambda(g) = P_\lambda(X)$, $X = 2g - 2 + n$. For example,

$$\begin{aligned} P_1(X) &= X^2 - X + \frac{3}{2}, \\ P_2(X) &= X^4 - 6X^3 + \frac{27}{2}X^2 - \frac{27}{2}X + \frac{135}{8}, \\ P_{1,1}(X) &= X^4 - 10X^3 + 38X^2 - 68X + \frac{273}{4}, \\ P_3(X) &= X^6 - 15X^5 + \frac{177}{2}X^4 - 260X^3 + \frac{3375}{8}X^2 - \frac{3375}{8}X + \frac{7875}{16}, \\ P_{1,2}(X) &= X^6 - 21X^5 + 179X^4 - 795X^3 + \frac{15957}{8}X^2 - \frac{23247}{8}X + \frac{41121}{16}, \\ P_{1,1,1}(X) &= X^7 - 28X^6 + \frac{653}{2}X^5 - \frac{4109}{2}X^4 + \frac{30361}{4}X^3 - \frac{33581}{2}X^2 + \frac{170757}{8}X - \frac{82467}{8}. \end{aligned}$$

The statement of Proposition 5 can then be written equivalently as

$$\frac{C(\lambda, g-1-|\lambda|)}{C(g-1)} = \frac{P_\lambda(2g+n-2)}{\prod_{j=1}^{n-3} (2g+j)} \prod_{i=1}^{|\lambda|} \frac{2g-2i}{(2g-2i+1)^3}, \quad (81)$$

where $P_\lambda(X) \in \mathbb{Z}[1/2][X]$ is a monic polynomial.

By using formula (10) and formula (81) we arrive at the following proposition.

Proposition 6. *For any fixed $n \geq 2$, fixed $\lambda = (\lambda_1, \dots, \lambda_{n-1}) \in (\mathbb{Z}_{\geq 1})^{n-1}$, and for d_n being an indeterminate, we have*

$$C(\lambda, d_n) = \frac{1}{X(\lambda, d_n)^{\delta_{n,2}} (X(\lambda, d_n) - 1)!} \frac{(2d_n + 1)!!^3}{2^{d_n+1} d_n!} P_\lambda(X(\lambda, d_n)), \quad (82)$$

where $P_\lambda(X) \in \mathbb{Z}[1/2][X]$ is a monic polynomial of degree $\sum_{j=1}^{n-1} (2\lambda_j + 1) + \delta_{n,2} - 2$.

We observe from the above examples that the polynomials $P_\lambda(X)$ for $|\lambda| = 1, 2$ are irreducible over \mathbb{Q} , and we have checked that it is still true for $|\lambda| = 3, \dots, 9$. We expect that this irreducibility holds for all partitions λ , and this is consistent with an observation that the BGW numbers often contain large prime factors.

Remark 1. For $\mathbf{d}' = (d_1, \dots, d_{n-1}) \in (\mathbb{Z}_{\geq 0})^{n-1}$ fixed and d_n an indeterminate, write $\mathbf{d} = (\mathbf{d}', d_n)$. Then from Proposition 6 and (8) we have

$$C(\mathbf{d}) = \frac{1}{X(\mathbf{d})^{\delta_{n,2}} (X(\mathbf{d}) - 1)!} \frac{(2d_n + 1)!!^3}{2^{d_n+1} d_n!} P_{\mathbf{d}'}(X(\mathbf{d})), \quad (83)$$

Here $P_{\mathbf{d}'}(X)$ is defined via

$$P_{0, \mathbf{d}'}(X) := (X - 1)^{1 - \delta_{n,2}} P_{\mathbf{d}'}(X - 1), \quad \mathbf{d}' \in (\mathbb{Z}_{\geq 0})^{n-1}. \quad (84)$$

Formula (83) (or Proposition 6) implies the following corollary, which is similar to a result of Liu–Xu [31, 33] for Witten’s intersection numbers.

Corollary 1. For $g, n \geq 1$, $d_1, \dots, d_n \geq 0$ satisfying $d_1 + \dots + d_n = g - 1$, we have

$$g^{\delta_{n,2}} \frac{d_n! \prod_{j=1}^n (2d_j + 1)!!}{(2d_n + 1)!!^3} \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g \in \mathbb{Z}[1/2]. \quad (85)$$

In the introduction we formulated the “Integrality Conjecture” that the numbers $\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g$ are integral away from 2, and also made the observation that they are often highly factorized. Corollary 1 does not imply the Integrality Conjecture, but does give both some bounds on the denominators of $\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g$ and nice information about prime factors of $\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g$. The following statement, based on the numerical data up to genus 40, gives a stronger version of the Integrality Conjecture.

Conjecture 3. For $n \geq 1$ and $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$, we have both

$$\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g^\ominus \in \frac{\prod_{j=1}^n d_j!}{2^{4g}} \mathbb{Z} \quad (86)$$

and

$$\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_g^\ominus \in \frac{\max_{1 \leq j \leq n} \{(2d_j + 1)!!\} \prod_{r \geq 0, p_r(\mathbf{d}) \geq 1} (p_r(\mathbf{d}) - 1)!}{2^{4g}} \mathbb{Z}, \quad (87)$$

where $g = |\mathbf{d}| + 1$, and $p_r(\mathbf{d})$ denotes the multiplicity of r in \mathbf{d} .

We notice that none of (85), (86), (87) imply either of the others, and also that each is weaker than the best possible factorization. For instance,

$$\langle \tau_6 \tau_7 \tau_8 \tau_{18} \rangle_{40}^\ominus = 2^{-150} 3^{18} 5^{11} 7^5 11^2 13^3 17^1 19^2 23^2 29^2 31^2 37^2 101^1 M,$$

where M has no prime factors less than 1000, whereas formula (85) implies the divisibility of $\langle \tau_6 \tau_7 \tau_8 \tau_{18} \rangle_{40}^\ominus$ (away from 2) by $3^{-1} 13^{-2} 5^2 7^1 19^2 23^2 29^2 31^2 37^2$, formula (86) implies the divisibility by $3^{14} 5^6 7^4 11^1 13^1 17^1$, and formula (87) implies the divisibility by $3^9 5^5 7^3 11^2 13^1 17^1 19^1 23^1 29^1 31^1 37^1$.

5. UNIFORM LARGE GENUS ASYMPTOTICS

In this section, we prove Theorem 1. Our proof will mainly use the recursion (46), and techniques introduced by Aggarwal [1] in the study of the large genus asymptotics of Witten's intersection numbers. Before entering into the details, it is convenient to rewrite the DVV-type relation (46) in terms of $C(\mathbf{d})$ as follows:

$$\begin{aligned} C(\mathbf{d}) &= \sum_{j=2}^n \frac{2d_j + 1}{X(\mathbf{d}) - 1} C(d_2, \dots, d_j + d_1, \dots, d_n) \\ &+ \sum_{\substack{a, b \geq 0 \\ a+b=d_1-1}} \left[\frac{2}{X(\mathbf{d}) - 1} C(a, b, d_2, \dots, d_n) \right. \\ &+ \left. \sum_{I \sqcup J = \{2, \dots, n\}} \frac{(X(a, \mathbf{d}_I) - 1)! (X(b, \mathbf{d}_J) - 1)!}{(X(\mathbf{d}) - 1)!} C(a, \mathbf{d}_I) C(b, \mathbf{d}_J) \right], \end{aligned} \quad (88)$$

where $n \geq 1$, $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$, and $X(\cdot)$ is as in (7).

5.1. Lower bound. Let us first show the following lemma on positivity of $C(\mathbf{d})$.

Lemma 1. *For every $n \geq 1$ and every $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$, we have*

$$C(\mathbf{d}) > 0. \quad (89)$$

Proof. By using the DVV-type relation (88) and by recalling that $C(0) = 1/4$. \square

Now we give in the following lemma a better lower bound for $C(\mathbf{d})$.

Lemma 2. *For every $n \geq 1$ and every $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$, we have*

$$C(\mathbf{d}) \geq C(|\mathbf{d}|). \quad (90)$$

Proof. Noticing that the statement is trivial when $n = 1$ and the fact that $C(\mathbf{d})$ is unchanged by removing 0's in \mathbf{d} , we can assume $n \geq 2$ and $d_1, \dots, d_n \geq 1$. Because of the symmetry of $C(\mathbf{d})$ in its arguments, we can also assume that d_1 is the smallest of the d_j 's. We now do mathematical induction with respect to $X(\mathbf{d}) = 2|\mathbf{d}| + n \geq 3$. For $X(\mathbf{d}) = 3$, from Table 1 we see that (90) is true. By Lemma 1 and by the induction hypothesis we get

$$C(\mathbf{d}) - C(|\mathbf{d}|) \geq -2d_1 \frac{C(|\mathbf{d}|) - C(|\mathbf{d}| - 1)}{X(\mathbf{d}) - 1} + 2 \frac{C(0) C(|\mathbf{d}| - 1)}{(X(\mathbf{d}) - 1)(X(\mathbf{d}) - 2)}. \quad (91)$$

Using formula (10), we can write the right-hand side of (91) as

$$\frac{1}{2} \left(\frac{1}{X(\mathbf{d}) - 2} - \frac{d_1}{|\mathbf{d}|(|\mathbf{d}| + 1)} \right) \frac{C(|\mathbf{d}| - 1)}{X(\mathbf{d}) - 1}, \quad (92)$$

which is positive because $d_1 \leq \frac{|\mathbf{d}|}{n}$ and $n \geq 2$. This finishes the proof. \square

5.2. Upper bound. To give an upper bound of $C(\mathbf{d})$, we first give in the following lemma an estimate related to the quadratic-in- C terms of (88), analogous to [1, Lemma 3.1].

Lemma 3. For $n \geq 1$ and for $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 1})^n$, we have

$$\sum_{\substack{a, b \geq 0 \\ a+b=d_1-1}} \sum_{I \sqcup J = \{2, \dots, n\}} \frac{(X(a, \mathbf{d}_I) - 1)! (X(b, \mathbf{d}_J) - 1)!}{(X(\mathbf{d}) - 1)!} \leq \frac{4}{(X(\mathbf{d}) - 1)(X(\mathbf{d}) - 2)}. \quad (93)$$

Proof. For $n = 1$, the inequality (93) is trivial. Assume that $n \geq 2$. For each a, b, I, J satisfying $a + b = d_1 - 1$ and $I \sqcup J = \{2, \dots, n\}$, we set $n_1 = |I|$, $g_1 = a + 1 + |\mathbf{d}_I|$, $n_2 = |J|$, $g_2 = b + 1 + |\mathbf{d}_J|$. Then $X(a, \mathbf{d}_I) = 2g_1 + n_1 - 1$ and $X(b, \mathbf{d}_J) = 2g_2 + n_2 - 1$. By counting the number of 4-tuples (a, b, I, J) with given values of n_i and g_i , we find

$$\begin{aligned} & \sum_{\substack{a, b \geq 0 \\ a+b=d_1-1}} \sum_{I \sqcup J = \{2, \dots, n\}} \frac{(X(a, \mathbf{d}_I) - 1)! (X(b, \mathbf{d}_J) - 1)!}{(X(\mathbf{d}) - 1)!} \\ & \leq \sum_{n_1+n_2=n-1} \sum_{\substack{g_1, g_2 \geq 1 \\ g_1+g_2=g}} \binom{n-1}{n_1} \frac{(2g_1 + n_1 - 2)! (2g_2 + n_2 - 2)!}{(X(\mathbf{d}) - 1)!} \\ & = \sum_{n_1+n_2=n-1} \binom{n-1}{n_1} \left(2 \frac{n_1! (X(\mathbf{d}) - n_1 - 3)!}{(X(\mathbf{d}) - 1)!} + \sum_{\substack{g_1, g_2 \geq 2 \\ g_1+g_2=g}} \frac{(2g_1 + n_1 - 2)! (2g_2 + n_2 - 2)!}{(X(\mathbf{d}) - 1)!} \right), \end{aligned} \quad (94)$$

where $g = g(\mathbf{d}) = |d| + 1$ as usual. We estimate the two terms on the right-hand side of (94) separately. For the first term we have

$$\begin{aligned} 2 \sum_{n_1+n_2=n-1} \binom{n-1}{n_1} \frac{n_1! (X(\mathbf{d}) - n_1 - 3)!}{(X(\mathbf{d}) - 1)!} &= \frac{2}{(X(\mathbf{d}) - 1)(X(\mathbf{d}) - 2)} \sum_{n_1=0}^{n-1} \prod_{j=1}^{n_1} \frac{n-j}{X(\mathbf{d}) - 2 - j} \\ &\leq \frac{2}{(X(\mathbf{d}) - 1)(X(\mathbf{d}) - 2)} \sum_{n_1=0}^{\infty} \left(\frac{1}{3}\right)^{n_1} = \frac{3}{(X(\mathbf{d}) - 1)(X(\mathbf{d}) - 2)}, \end{aligned} \quad (95)$$

where in the inequality we used the fact that $n \leq \frac{X(\mathbf{d})}{3}$ (implied by $\mathbf{d} \in (\mathbb{Z}_{\geq 1})^n$). For the second term we have

$$\begin{aligned} & \sum_{n_1+n_2=n-1} \sum_{\substack{g_1, g_2 \geq 2 \\ g_1+g_2=g}} \binom{n-1}{n_1} \frac{(2g_1 + n_1 - 2)! (2g_2 + n_2 - 2)!}{(X(\mathbf{d}) - 1)!} \\ & \leq \sum_{\substack{g_1, g_2 \geq 2 \\ g_1+g_2=g \\ n_1+n_2=n-1}} \frac{\binom{2g-4}{2g_1-2}^{-1}}{(X(\mathbf{d}) - 1)(X(\mathbf{d}) - 2)} \leq \frac{n(g-3) \binom{2g-4}{2}^{-1}}{(X(\mathbf{d}) - 1)(X(\mathbf{d}) - 2)} \leq \frac{1}{(X(\mathbf{d}) - 1)(X(\mathbf{d}) - 2)}, \end{aligned} \quad (96)$$

where for the first inequality we used the fact that $X(\mathbf{d}) = (2g_1 + n_1 - 2) + (2g_2 + n_2 - 2)$ and that

$$\binom{a_1}{b_1} \binom{a_2}{b_2} \leq \binom{a_1 + a_2}{b_1 + b_2}, \quad (97)$$

and for the last inequality we used $g \geq 3$ and $n \leq g - 1$. Combining (94), (95), (96), we obtain the lemma. \square

Following Aggarwal [1], for $\mathcal{X}, n \geq 1$, introduce

$$\theta_{\mathcal{X}, n} := \max_{\substack{\mathbf{d} \in (\mathbb{Z}_{\geq 0})^n \\ X(\mathbf{d}) = \mathcal{X}}} C(\mathbf{d}). \quad (98)$$

Before continuing, we introduce a number-theoretic function $f(\mathcal{X}, n)$, defined through the recursion

$$f(\mathcal{X}, n) = \frac{2}{3}f(\mathcal{X} - 1, n - 1) + \frac{1}{3}f(\mathcal{X} - 1, n + 1) + \frac{4}{(\mathcal{X} - 1)(\mathcal{X} - 2)}, \quad \forall n \geq 3, \mathcal{X} \geq 8, \quad (99)$$

together with the initial data $f(\mathcal{X}, n) = 1/\pi$ for $1 \leq \mathcal{X} \leq 7$ or $n = 1, 2$. By induction, we know that $f(\mathcal{X}, n)$ is monotone increasing with respect to n , and

$$f(\mathcal{X}, n) \leq \frac{1}{\pi} + \sum_{k=8}^{\mathcal{X}} \frac{4}{(k-1)(k-2)}. \quad (100)$$

Hence $f(\mathcal{X}, n)$ is bounded by 1. Let us also prove the following lemma.

Lemma 4. *For $\mathcal{X} \geq 1$, and for $1 \leq n \leq \frac{\mathcal{X}}{5}$, we have the uniform estimate*

$$f(\mathcal{X}, n) = \frac{1}{\pi} + O\left(\frac{1}{\mathcal{X}}\right), \quad \mathcal{X} \rightarrow \infty. \quad (101)$$

Proof. It is not difficult to show, either directly or using generating function, that the solution of the recursion (99) with the given initial condition, is given for $n \geq 2$ and $\mathcal{X} \geq 7$ by

$$f(\mathcal{X}, n) = \frac{1}{\pi} + \frac{2(\mathcal{X} - 7)}{3(\mathcal{X} - 1)} - \sum_{k=8}^{\mathcal{X}} \frac{2(k-7)}{3(k-1)} P(n, \mathcal{X} - k), \quad (102)$$

where the coefficients $P(n, j)$ are defined by

$$\left(\frac{3 - \sqrt{9 - 8t^2}}{2t}\right)^{n-2} =: \sum_{j=0}^{\infty} P(n, j) t^j. \quad (103)$$

These coefficients can be estimated by the residue theorem:

$$P(n, j) = \frac{1}{2\pi i} \int_C t^{-j-1} \left(\frac{3 - \sqrt{9 - 8t^2}}{2t}\right)^{n-2} dt \leq 1.05^{-j} \times 1.23^{n-2},$$

where we have taken the contour C to be the circle $|t| = 1.05 < \sqrt{9/8}$, on which we have $\left| \frac{3 - \sqrt{9 - 8t^2}}{2t} \right| < 1.23$. Let us now estimate the right-hand side of (102):

$$\sum_{k=8}^{\mathcal{X}} \frac{2(k-7)}{3(k-1)} P(n, \mathcal{X} - k) \geq \sum_{k=M+1}^{\mathcal{X}} \frac{2(k-7)}{3(k-1)} P(n, \mathcal{X} - k) \geq \frac{2(M-6)}{3M} \sum_{j=0}^{\mathcal{X}-M-1} P(n, j), \quad (104)$$

for any $7 \leq M \leq \mathcal{X}$. Taking $M = \lceil 0.1\mathcal{X} \rceil$ and using $n \leq \frac{\mathcal{X}}{5}$, we have

$$\sum_{j=\mathcal{X}-M}^{\infty} P(n, j) \leq 14 \times 1.05^{-\mathcal{X}+M} \times 1.23^n \leq 14 \times 0.998^{\mathcal{X}}. \quad (105)$$

By using $\sum_{j=0}^{\infty} P(n, j) = 1$ and concluding formula (102) and the estimates (104), (105), we finish the proof of the lemma. \square

The significance of the function $f(\mathcal{X}, n)$ is given by the following important lemma.

Lemma 5. *For $n \geq 1$, $\mathcal{X} \geq 1$, the numbers $\theta_{\mathcal{X}, n}$ have the upper bound*

$$\theta_{\mathcal{X}, n} \leq f(\mathcal{X}, n). \quad (106)$$

Proof. For $1 \leq \mathcal{X} \leq 7$, we check from Table 1 that inequality (106) holds. For $n = 1$, inequality (106) is implied by (10). For $n = 2$, let $0 \leq d_1 \leq d_2$ and $g = d_1 + d_2 + 1$. Using (69) we get

$$\begin{aligned} C(d_1, d_2) &= \frac{2^{2g}}{(2g)!} \sum_{h=0}^{d_1} (g-2h) F_h F_{g-h} \\ &\leq \frac{2^{2g}}{(2g)!} \left(gF_g + \frac{(g-1)(g-2)}{2} F_1 F_{g-1} \right) = \frac{(16g^2 - 23g + 9) \Gamma(g - \frac{1}{2})^2}{8\pi(2g-1)\Gamma(g)^2} \leq \frac{1}{\pi}, \end{aligned} \quad (107)$$

where in the first inequality we used that F_{d+1}/F_d is monotone increasing for $d \geq 0$.

Now consider the case that $n \geq 3$, $\mathcal{X} \geq 8$. Let us prove inequality (106) by induction on \mathcal{X} . For every $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$ satisfying $X(\mathbf{d}) = \mathcal{X}$, assume $\mathbf{d} \in (\mathbb{Z}_{\geq 1})^n$ and without loss of generality assume $d_1 = \min\{d_j\}$. Then by applying Lemma 3 to (88) and by induction, we obtain

$$C(\mathbf{d}) \leq \left(1 - \frac{2d_1}{\mathcal{X}-1}\right) f(\mathcal{X}-1, n-1) + \frac{2d_1}{\mathcal{X}-1} f(\mathcal{X}-1, n+1) + \frac{4}{(\mathcal{X}-1)(\mathcal{X}-2)}, \quad (108)$$

where we have used the fact that $f(\mathcal{X}, n)$ is bounded by 1. This gives

$$C(\mathbf{d}) \leq \frac{2}{3} f(\mathcal{X}-1, n-1) + \frac{1}{3} f(\mathcal{X}-1, n+1) + \frac{4}{(\mathcal{X}-1)(\mathcal{X}-2)} = f(\mathcal{X}, n),$$

where we have used $n \geq 3$, $d_1 \leq \frac{|\mathbf{d}|}{n} = \frac{\mathcal{X}-n}{2n}$, and the fact that the function $f(\mathcal{X}, n)$ is increasing with respect to n . For the case that some of d_j equal zero, by (8) and by induction we have

$$C(\mathbf{d}) \leq f(\mathcal{X}-1, n-1), \quad (109)$$

where the right-hand side is less than $f(\mathcal{X}, n)$ because of (99) and again the monotonicity of $f(\mathcal{X}, n)$. Combining both cases, we finish the proof of (106). \square

We are ready to prove Theorem 1.

Proof of Theorem 1. By Lemma 2 and (98), we have the lower and upper bound

$$C(|\mathbf{d}|) \leq C(\mathbf{d}) \leq \theta_{X(\mathbf{d}),n}, \quad (110)$$

for every $n \geq 1$, $\mathbf{d} \in (\mathbb{Z}_{\geq 0})^n$. We know from (10) that the lower bound equals $\frac{1}{\pi} + O(\frac{1}{g(\mathbf{d})})$ with an absolute constant in $O(\frac{1}{g(\mathbf{d})})$. For the upper bound, we know from Lemma 4 we know that when $n \leq \frac{X(\mathbf{d})}{5}$, $\theta_{X,n}$ is bounded by $\frac{1}{\pi} + O(\frac{1}{X})$ with an absolute constant in $O(\frac{1}{X})$. Consider the case that $\mathbf{d} \in (\mathbb{Z}_{\geq 0})^n$ with $\frac{X(\mathbf{d})}{5} < n \leq \frac{X(\mathbf{d})}{3}$. Assume $\mathbf{d} \in (\mathbb{Z}_{\geq 1})^n$. Then there must exist some j such that $d_j = 1$. Assuming that d_1 equals 1, then recursion (88) reads

$$\begin{aligned} C(1, d_2, \dots, d_n) &= \sum_{j=2}^n \frac{2d_j + 1}{X(\mathbf{d}) - 1} C(d_2, \dots, d_j + 1, \dots, d_n) \\ &+ \frac{2}{X(\mathbf{d}) - 1} C(d_2, \dots, d_n) + \sum_{I \sqcup J = \{2, \dots, n\}} \frac{(X(0, \mathbf{d}_I) - 1)! (X(0, \mathbf{d}_J) - 1)!}{(X(\mathbf{d}) - 1)!} C(\mathbf{d}_I) C(\mathbf{d}_J). \end{aligned} \quad (111)$$

By applying Lemma 3 and by taking maximum in both sides of (111), we get

$$C(\mathbf{d}) \leq \theta_{X(\mathbf{d})-1, n-1} + \frac{4}{(X(\mathbf{d}) - 1)(X(\mathbf{d}) - 2)}. \quad (112)$$

When some d_j is 0, the inequality (112) is still true according to (8), so we get

$$\theta_{X,n} \leq \theta_{X-1, n-1} + \frac{4}{(X-1)(X-2)}, \quad (113)$$

for every $\frac{X}{5} < n \leq \frac{X}{3}$. Writing this as

$$\theta_{X,n} + \frac{4}{X-1} \leq \theta_{X-1, n-1} + \frac{4}{X-2}, \quad (114)$$

and iterating t times we find

$$\theta_{X,n} \leq \theta_{X-t, n-t} + \frac{4}{X-1} - \frac{4}{X-1-t} \leq f(X-t, n-t) + \frac{4}{X-1} - \frac{4}{X-1-t}, \quad (115)$$

for any $t \leq \lfloor \frac{5n-X+1}{4} \rfloor$. Applying this with $t = \lfloor \frac{5n-X+1}{4} \rfloor$ and using Lemma 4 we obtain that $\theta_{X,n}$ is bounded by $\frac{1}{\pi} + O(\frac{1}{X})$ uniformly when $\frac{X}{5} < n \leq \frac{X}{3}$. This together with (110) implies that formula (19) holds for $\frac{X(\mathbf{d})}{5} \leq n \leq \frac{X(\mathbf{d})}{3}$. For $n > \frac{X(\mathbf{d})}{3}$, we have $\theta_{X(\mathbf{d}),n} = \theta_{X(\mathbf{d})-1, n-1}$ (indeed, since $C(\mathbf{d})$ is unchanged by removing any 0 argument, we have by definition $\theta_{X,n} = \theta_{X-1, n-1}$ for $n > \frac{X}{3}$), which implies that formula (19) still holds. Combining all three cases, we obtain the statement of Theorem 1. \square

Remark 2. Although our proof for BGW numbers is similar to Aggarwal's for Witten's intersection numbers [1], we have made several improvements and simplifications. For examples, the technique of random walks used in [1] is avoid here, and our estimates are completely uniform instead of requiring $n = o(\sqrt{g})$ in [1]. Actually, as it was shown in [10], it is not possible to extend the asymptotics of the normalized Witten's intersection numbers [1, 10, 11, 25] beyond the range $n = o(\sqrt{g})$. We hope to generalize Theorem 1 to Witten's intersection numbers with a better normalization.

6. POLYNOMIALITY IN LARGE GENUS

In this section, we prove Theorem 2 by using the recursion (88).

Proof of Theorem 2. We first allow the fixed $\mathbf{d}' = (d_1, \dots, d_{n-1}) \in (\mathbb{Z}_{\geq 0})^{n-1}$. Write $\mathbf{d} = (\mathbf{d}', d_n)$ with $d_n \geq 0$. By using (83) and Stirling's formula we have

$$C(\mathbf{d}) = \frac{1}{\pi} \sum_{k=0}^{\infty} \frac{C_k(\mathbf{d}')}{X(\mathbf{d})^k}, \quad \text{as } d_n \rightarrow \infty, \quad (116)$$

where C_k are functions of \mathbf{d}' . By using the recursion (88) and by performing Laurent expansions, we obtain

$$\begin{aligned} C_k(d, \mathbf{d}') - C_k(\mathbf{d}') &= - \sum_{l=1}^{k-1} (-1)^{k-l} \binom{k-1}{l-1} C_l(d, \mathbf{d}') \\ &+ \sum_{j=1}^{n-1} (2d_j + 1) (C_{k-1}(d_1, \dots, d_j + d, \dots, d_{n-1}) - C_{k-1}(\mathbf{d}')) \\ &+ \sum_{\substack{a, b \geq 0 \\ a+b=d-1}} \left[2 (C_{k-1}(a, b, \mathbf{d}') - C_{k-1}(\mathbf{d}')) + \sum_{I \sqcup J = \{1, \dots, n-1\}} \sum_{l=0}^{k-2} \mathbf{a}_{(a, \mathbf{d}'_I), k, l} C_l(b, \mathbf{d}'_J) \right], \end{aligned} \quad (117)$$

where $d \geq 0$ and $\mathbf{a}_{\mathbf{w}, k, l}$ are numbers defined by

$$\mathbf{a}_{\mathbf{w}, k, l} := \sum_{u=l}^{k-2} \left(4^{|\mathbf{w}|+1} \binom{u-1}{l-1} X(\mathbf{w})^{u-l} S(k-u-1, X(\mathbf{w})) B(\mathbf{w}) \right).$$

Here $B(\mathbf{w})$ is defined in (44) and $S(n, k)$ denotes the Stirling number of the second kind. Write

$$C_k(\mathbf{d}') =: \tilde{c}_k(p_0(\mathbf{d}'), p_1(\mathbf{d}'), \dots), \quad (118)$$

where $p_r(\mathbf{d}')$ denotes the multiplicity of r in \mathbf{d}' . This defines functions $\tilde{c}_k(\mathbf{p})$, $k \geq 0$, where $\mathbf{p} = (p_0, p_1, p_2, \dots)$. Then formula (117) becomes

$$\begin{aligned} \tilde{c}_k(\mathbf{p} + \mathbf{e}_d) - \tilde{c}_k(\mathbf{p}) &= - \sum_{l=1}^{k-1} (-1)^{k-l} \binom{k-1}{l-1} \tilde{c}_l(\mathbf{p} + \mathbf{e}_d) \\ &+ \sum_{i \geq 0} (2i + 1) p_i (\tilde{c}_{k-1}(\mathbf{p} - \mathbf{e}_i + \mathbf{e}_{i+d}) - \tilde{c}_{k-1}(\mathbf{p})) \\ &+ \sum_{\substack{a, b \geq 0 \\ a+b=d-1}} \left[2 \sum_{l=0}^{k-1} (\tilde{c}_{k-1}(\mathbf{p} + \mathbf{e}_a + \mathbf{e}_b) - \tilde{c}_{k-1}(\mathbf{p})) \right. \\ &\left. + \sum_{\substack{\mathcal{E}(\mathbf{t} + \mathbf{e}_a) \leq k-1 \\ 0 \leq t_r \leq p_r, r \geq 0}} \sum_{l=0}^{k-2} \left(\tilde{c}_l(\mathbf{p} - \mathbf{t} + \mathbf{e}_b) \alpha_{\mathbf{t} + \mathbf{e}_a, k, l} \prod_{i \geq 0} \binom{p_i}{t_i} \right) \right], \quad d \geq 0, \end{aligned} \quad (119)$$

where $\mathcal{E}(\mathbf{t}) = \sum_{j=0}^{\infty} (2j+1)t_j$, $\alpha_{\mathbf{t}, k, l} = \mathbf{a}_{(0^{t_0} 1^{t_1} 2^{t_2} \dots), k, l}$, and \mathbf{e}_d denotes $(0, \dots, 0, 1, 0, 0, \dots)$ with "1" appearing in the $(d+1)$ th place.

Let us now prove by induction that $\tilde{c}_k(\mathbf{p})$, $k \geq 0$, belong to $\mathbb{Q}[p_0, p_1, \dots]$ and satisfy the degree estimates

$$\deg \tilde{c}_k(\mathbf{p}) \leq k - 1, \quad k \geq 1, \quad (120)$$

under the degree assignments $\deg p_d = 2d + 1$, $d \geq 0$. For $k = 0$, by using Theorem 1 we know that $\tilde{c}_0(\mathbf{p}) \equiv 1$. Assume that for $1 \leq l \leq k - 1$, $\tilde{c}_l(\mathbf{p}) \in \mathbb{Q}[p_0, p_1, \dots]$ are polynomials satisfying $\deg \tilde{c}_l(\mathbf{p}) \leq l - 1$ ($l \geq 1$). Then for k and for every $d \geq 0$, the RHS of equation (119) are polynomials in $p_0, \dots, p_{\lfloor (k-3)/2 \rfloor}$. Moreover, by the inductive assumption these polynomials are independent of d for every $d \geq k + 1$, i.e.,

$$\tilde{c}_k(\mathbf{p} + \mathbf{e}_d) - \tilde{c}_k(\mathbf{p}) = \begin{cases} f_d(p_0, \dots, p_{\lfloor (k-3)/2 \rfloor}), & d \leq k, \\ g(p_0, \dots, p_{\lfloor (k-3)/2 \rfloor}), & d \geq k + 1 \end{cases} \quad (121)$$

for some f_d ($d \leq k$), g belonging to $\mathbb{Q}[p_0, \dots, p_{\lfloor (k-3)/2 \rfloor}]$. The compatibility of (121) implies that $g(p_0, \dots, p_{\lfloor (k-3)/2 \rfloor}) \equiv A$ is a constant. Solving (121) we obtain that \tilde{c}_k have the form

$$\tilde{c}_k(\mathbf{p}) = h(p_0, \dots, p_k) + A n'(\mathbf{p}), \quad (122)$$

where $h \in \mathbb{Q}[p_0, \dots, p_k]$ and $n'(\mathbf{p}) := \sum_{i \geq 0} p_i$. Now we aim to show that $A = 0$. Consider equation (119) with k replaced by $k + 1$. Using a similar analysis and using (122), we obtain that for every $d \geq 2k + 2$,

$$\tilde{c}_{k+1}(\mathbf{p} + \mathbf{e}_d) - \tilde{c}_{k+1}(\mathbf{p}) = 4Ad + A' n'(\mathbf{p}) + g(p_0, \dots, p_k). \quad (123)$$

where $A' \in \mathbb{Q}$ is a constant, and $g(p_0, \dots, p_k)$ is some polynomial in $\mathbb{Q}[p_0, \dots, p_k]$. This contradicts with (69) unless $A = 0$. Therefore,

$$\tilde{c}_k(\mathbf{p}) \in \mathbb{Q}[p_0, \dots, p_k]. \quad (124)$$

Then taking $d \geq k + 1$ in equation (119) gives

$$\begin{aligned} 0 = & - \sum_{l=1}^{k-1} (-1)^{k-l} \binom{k-1}{l-1} \tilde{c}_l(\mathbf{p}) + \sum_{i \geq 0} (2i+1) p_i (\tilde{c}_{k-1}(\mathbf{p} - \mathbf{e}_i) - \tilde{c}_{k-1}(\mathbf{p})) \\ & + \sum_{a=0}^{\lfloor \frac{k-3}{2} \rfloor} \left(4 (\tilde{c}_{k-1}(\mathbf{p} + \mathbf{e}_a) - \tilde{c}_{k-1}(\mathbf{p})) + \sum_{\substack{\mathcal{E}(\mathbf{t} + \mathbf{e}_a) \leq k-1 \\ 0 \leq t_r \leq p_r, r \geq 0}} \sum_{l=0}^{k-2} \left(\tilde{c}_l(\mathbf{p} - \mathbf{t}) \alpha_{\mathbf{t} + \mathbf{e}_a, k, l} \prod_{i \geq 0} \binom{p_i}{t_i} \right) \right). \end{aligned} \quad (125)$$

Using (125) and (119), we obtain

$$\begin{aligned}
 \Delta_{p_d} \tilde{c}_k(\mathbf{p}) &= - \sum_{l=1}^{k-1} (-1)^{k-l} \binom{k-1}{l-1} \Delta_{p_d} \tilde{c}_l(\mathbf{p}) + \sum_{i \geq 0} (2i+1) p_i \Delta_{p_{i+d}} \tilde{c}_{k-1}(\mathbf{p} - \mathbf{e}_i) \\
 &+ 2 \sum_{a=0}^{d-1} \Delta_{p_a} \Delta_{p_{d-1-a}} \tilde{c}_{k-1}(\mathbf{p}) - 4 \sum_{a=d}^{[(k-3)/2]} \Delta_{p_a} \tilde{c}_{k-1}(\mathbf{p}) \\
 &+ \sum_{a=0}^{d-1} \sum_{\substack{\mathcal{E}(\mathbf{t}+\mathbf{e}_a) \leq k-1 \\ 0 \leq t_r \leq p_r (r \geq 0)}} \sum_{l=0}^{k-2} \left(\Delta_{p_{d-1-a}} \tilde{c}_l(\mathbf{p} - \mathbf{t}) \alpha_{\mathbf{t}+\mathbf{e}_a, k, l} \prod_{i \geq 0} \binom{p_i}{t_i} \right) \\
 &+ \sum_{a=d}^{[(k-3)/2]} \sum_{\substack{\mathcal{E}(\mathbf{t}+\mathbf{e}_a) \leq k-1 \\ 0 \leq t_r \leq p_r (r \geq 0)}} \sum_{l=0}^{k-2} \left(\tilde{c}_l(\mathbf{p} - \mathbf{t}) \alpha_{\mathbf{t}+\mathbf{e}_a, k, l} \prod_{i \geq 0} \binom{p_i}{t_i} \right), \quad d \geq 0. \tag{126}
 \end{aligned}$$

We find that each term of the RHS of (126) is of degree less than or equal to $k-2-2d$ for every $d \geq 0$, which implies $\deg \tilde{c}_k \leq k-1$. In particular, \tilde{c}_k is a polynomial that only depends on $p_0, \dots, p_{[k/2]-1}$.

Now restrict to the case when \mathbf{d}' is a fixed partition. From (116), (29) we know that

$$\widehat{C}(\mathbf{d}) \sim \sum_{k=0}^{\infty} \frac{\widehat{C}_k(\mathbf{d}')}{X(\mathbf{d})^k}, \quad d_n \rightarrow \infty, \tag{127}$$

where \widehat{C}_k are functions of \mathbf{d}' with $\widehat{C}_0 \equiv 1$. Define $\widehat{c}_k(p_1, p_2, \dots) \in \mathbb{Q}[p_1, p_2, \dots]$, $k \geq 0$, via

$$\gamma(X) \sum_{k=0}^{\infty} \frac{\widehat{C}_k(p_1, p_2, \dots)}{X^k} = \sum_{k=0}^{\infty} \frac{\tilde{c}_k(0, p_1, p_2, \dots)}{X^k}, \tag{128}$$

where the left-hand side is understood as a power series in X^{-1} . It then follows from $\deg \tilde{c}_k \leq k-1$ that $\deg \widehat{c}_k \leq k-1$. Using (127), (116), (118), we know that

$$\widehat{C}_k(\mathbf{d}') = \widehat{c}_k(p_1(\mathbf{d}'), p_2(\mathbf{d}'), \dots), \tag{129}$$

for all \mathbf{d}' . The statement that $\widehat{c}_k(0, 0, \dots) = 0$ follows from the fact that $C(d) = \gamma(2d+1)$. This finishes the proof of Theorem 2. \square

Remark 3. Formula (117) is analogous to a formula given in [33, Corollary 3.6].

For $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$, we extend the definition of $\widehat{C}(\mathbf{d})$ in (26) by

$$\widehat{C}(\mathbf{d}) := \frac{C(\mathbf{d})}{\gamma(X(\mathbf{d}) - p_0(\mathbf{d}))}. \tag{130}$$

Then the following corollary easily follows from Theorem 2.

Corollary 2. For any fixed $n \geq 1$ and fixed $\mathbf{d}' = (d_1, \dots, d_{n-1}) \in (\mathbb{Z}_{\geq 0})^{n-1}$, we have

$$\widehat{C}(\mathbf{d}) \sim \sum_{k=0}^{\infty} \frac{\widehat{c}_k(p_1(\mathbf{d}'), p_2(\mathbf{d}'), \dots)}{(X(\mathbf{d}) - p_0(\mathbf{d}'))^k}, \quad X(\mathbf{d}) \rightarrow \infty, \tag{131}$$

where $\mathbf{d} = (\mathbf{d}', d_n)$, and $\widehat{c}_k(p_1, p_2, \dots)$ are the same polynomials as those in Theorem 2.

We note that, since for fixed $n \geq 1$ and fixed $d_1, \dots, d_{n-1} \geq 0$, $C(\mathbf{d})/C(|\mathbf{d}|)$ is a rational function of $X(\mathbf{d})$ whose asymptotic expansion is convergent, the polynomials $\widehat{c}_k(p_1, p_2, \dots)$, $k \geq 0$, contain all information of BGW numbers. We provide in Table 2 some explicit values of $\widehat{c}_k(p_1, p_2, \dots)$, $c_k(p_1, p_2, \dots)$.

k	$c_k(p_1, p_2, \dots)$	$\widehat{c}_k(p_1, p_2, \dots)$
0	1	1
1	$-\frac{1}{2}$	0
2	$\frac{5}{8}$	0
3	$-\frac{11}{16}$	0
4	$\frac{83}{128} - \frac{27}{8} p_1$	$-\frac{27}{8} p_1$
5	$-\frac{143}{256} - \frac{81}{16} p_1$	$-\frac{27}{4} p_1$
6	$\frac{625}{1024} - \frac{639}{64} p_1 - \frac{1125}{16} p_2$	$-\frac{45}{4} p_1 - \frac{1125}{16} p_2$
7	$-\frac{1843}{2048} + \frac{25533}{128} p_1 - \frac{1701}{8} p_1^2 - \frac{19125}{32} p_2$	$\frac{783}{4} p_1 - \frac{1701}{8} p_1^2 - \frac{10125}{16} p_2$

TABLE 2. Expressions for $c_k(p_1, p_2, \dots)$ and $\widehat{c}_k(p_1, p_2, \dots)$ with $k \leq 7$

Similar to (30), for fixed $L \geq 0$ and fixed $n \geq 1$, and for $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$,

$$\widehat{C}(\mathbf{d}) = \sum_{k=0}^{L-1} \frac{\widehat{c}_k(p_1(\mathbf{d}), p_2(\mathbf{d}), \dots)}{(X(\mathbf{d}) - p_0(\mathbf{d}))^k} + O\left(\frac{1}{(X(\mathbf{d}) - p_0(\mathbf{d}))^L}\right), \quad g(\mathbf{d}) \rightarrow \infty, \quad (132)$$

where the implied O-constant only depends on n and L .

Remark 4. We have the following conjectural statement, which is stronger than (132), that for any fixed $L \geq 0$,

$$\widehat{C}(\mathbf{d}) = \sum_{k=0}^{L-1} \frac{\widehat{c}_k(p_1(\mathbf{d}), p_2(\mathbf{d}), \dots)}{(X(\mathbf{d}) - p_0(\mathbf{d}))^k} + O\left(\frac{\widehat{c}_L(p_1(\mathbf{d}), p_2(\mathbf{d}), \dots)}{(X(\mathbf{d}) - p_0(\mathbf{d}))^L}\right), \quad g(\mathbf{d}) \rightarrow \infty, \quad (133)$$

where the implied constant only depends on L . In terms of $C(\mathbf{d})$, this conjecture states that for any fixed $L \geq 0$,

$$C(\mathbf{d}) = \frac{1}{\pi} \sum_{k=0}^{L-1} \frac{c_k(p_1(\mathbf{d}), p_2(\mathbf{d}), \dots)}{(X(\mathbf{d}) - p_0(\mathbf{d}))^k} + O\left(\frac{c_L(p_1(\mathbf{d}), p_2(\mathbf{d}), \dots)}{(X(\mathbf{d}) - p_0(\mathbf{d}))^L}\right), \quad (134)$$

as $g(\mathbf{d}) \rightarrow \infty$, where the implied constant only depends on L .

We end this section by giving some information about the interval $I_{g,n}$ defined in the introduction. Denote by $m(g, n)$ and $M(g, n)$ its endpoints, i.e., the minimum and maximum of all $C(\mathbf{d})$ with $\mathbf{d} \in (\mathbb{Z}_{\geq 1})^n$ and $g = |\mathbf{d}| + 1$. Conjecture 2 implies that $m(g, n) = C(1^n, g - 1 - n)$ and $M(g, n) = C(d^p(d+1)^{n-p})$, where $d = \lfloor \frac{g-1}{n} \rfloor$ and $p = (d+1)n - g + 1$. Using (133), we then find the conjectural asymptotic formulas

$$\gamma(2g - 2 + n) - m(g, n) = \frac{27n}{8\pi(2g - 2 + n)^4} + O\left(\frac{1}{g^4}\right), \quad (135)$$

$$\gamma(2g - 2 + n) - M(g, n) = \frac{(2d+1)!!^3}{2^{d+1}\pi(d+1)!} \frac{(d+1)n - g}{(2g - 2 + n)^{2d+2}} + O\left(\frac{1}{g^{2d+2}}\right), \quad (136)$$

where $d = \lfloor \frac{g-1}{n} \rfloor$ and both O -constants are absolute. In particular $M(g, n)$ differs from $\gamma(2g - 2 + n)$ by a quantity of the order of g^{-2d-1} . These two formulas are only to leading order in $1/X$, but the full analysis gives several more terms. These statements, which greatly refine Theorem 1, are still only conjectural, but we are currently trying to prove some version of them and hope to return to this in a later publication.

7. FURTHER ASYMPTOTIC FORMULAS

In this section, we will give asymptotic formulas of another type for BGW numbers, including and based on a closed asymptotic formula for two-point BGW numbers.

For every $d \geq 1$, define the power series $W_d(X) \in X^{-2d-2} \mathbb{Q}[[X^{-1}]]$ by the formula

$$1 - \widehat{C}(d, \frac{X}{2} - 1 - d) \sim W_d(X), \quad \text{as } X \rightarrow \infty. \quad (137)$$

(The fact that $W_d(X)$ is well defined is because of Theorem 2.) In terms of the polynomials $\widehat{c}_k(\mathbf{p})$, we have

$$W_d(X) = - \sum_{k=2d+2}^{\infty} \frac{\widehat{c}_k(\mathbf{e}_d)}{X^k}. \quad (138)$$

Formula (69) implies that, for every $d \geq 2$,

$$W_{d-1}(X) - W_d(X) = \frac{(2d-1)!!^3 \left(\frac{X}{2} - 2d\right) \Gamma\left(\frac{X+3}{2}\right) \Gamma\left(\frac{X+1}{2} - d\right)^3}{8^d d! \Gamma\left(\frac{X}{2} + 1\right)^3 \Gamma\left(\frac{X}{2} + 1 - d\right)}, \quad (139)$$

where the right-hand side is interpreted as a power series of X^{-1} by Stirling's formula. Together with the limiting condition $W_\infty(X) = 0$, formula (139) determines $W_d(X)$, $d \geq 1$, completely, the first few terms being

$$\begin{aligned} W_d(X) = & \frac{(2d+1)!!^3}{2^{d+1} (d+1)!} \left(\frac{1}{X^{2d+2}} + \frac{(2d-1)(d+1)}{X^{2d+3}} \right. \\ & \left. + \frac{(2d+3)(d+1)(6d^3 + 7d^2 - 8d + 1)}{6(d+2) X^{2d+4}} + \dots \right). \end{aligned} \quad (140)$$

In the following proposition, we give an explicit formula for $W_d(X)$.

Proposition 7. *The power series $W_d(X)$ for any fixed $d \geq 1$ is given by*

$$W_d(X) = \frac{(2d+1)!!^3}{2^{d+1} d!} \sum_{j \geq 1} \frac{A_j(d)}{d+j} \frac{1}{(X - 2d - j + 1)_{2d+2j}}. \quad (141)$$

Here the inverse Pochhammer symbols $1/(X - 2d - j + 1)_{2d+2j}$ are interpreted as elements of $\mathbb{Q}[[1/X]]$, and $A_j(d)$ ($j \geq 1$) are polynomials defined by the asymptotic formula

$$2^{-2d-4} \frac{\Gamma\left(\frac{X+1}{2} - d\right)^3 \Gamma\left(\frac{X+3}{2}\right)}{\Gamma\left(\frac{X}{2} + 2\right)^3 \Gamma\left(\frac{X}{2} - d + 1\right)} = \sum_{j=1}^{\infty} \frac{A_j(d)}{(X - 2d - j + 1)_{2d+2j+2}}, \quad (142)$$

in which both sides are interpreted as power series of X^{-1} . More explicitly,

$$A_j(d) = (-1)^{j-1} (j-1)! \sum_{0 \leq l \leq \lfloor \frac{j+1}{2} \rfloor} \frac{(2l-1)!!}{8^l l!^3} (j-2l)_{2l} \left(d + \frac{3}{2} - l\right)_{j-1}. \quad (143)$$

Proof. It is easy to verify that the left-hand side of (142) (as a power series in X^{-1}) is invariant under $X \rightarrow 2d - 3 - X$, so $A_j(d)$ is well defined from (142).

Let us first use the mathematical induction to prove the following equality:

$$\widehat{C}(d_1, d_2 + 1) - \widehat{C}(d_1, d_2) = \frac{4^{(d_1+d_2+2)} (2d_1 + 4d_2 + 7) (d_1 + 1) F_{d_1+1} F_{d_2+1}}{(2d_1 + 2d_2 + 5)! \gamma(2d_1 + 2d_2 + 4)}, \quad (144)$$

with F_d and $\gamma(X)$ given by (68) and (23), respectively. Denote by $G(d_1, d_2)$ the right-hand side of (144). For $d_1 = 0$, (144) follows directly from (10). If we assume that (144) is true for $d_1 = k - 1$, then for $d_1 = k$

$$\begin{aligned} \widehat{C}(k, d_2 + 1) - \widehat{C}(k, d_2) &= (\widehat{C}(k - 1, d_2 + 2) - \widehat{C}(k - 1, d_2 + 1)) \\ &\quad + (\widehat{C}(k, d_2 + 1) - \widehat{C}(k - 1, d_2 + 2)) - (\widehat{C}(k, d_2) - \widehat{C}(k - 1, d_2 + 1)) \\ &= G(k - 1, d_2 + 1) + \frac{4^{k+d_2+2} (d_2 + 2 - k) F_k F_{d_2+2}}{(2k + 2d_2 + 4)!} - \frac{4^{k+d_2+1} (d_2 + 1 - k) F_k F_{d_2+1}}{(2k + 2d_2 + 2)!} \\ &= G(k, d_2), \end{aligned}$$

where for the second equality we used formula (69). This completes the proof of (144).

By (137) and (144) we find the identity

$$W_d(X) - W_d(X + 2) = \frac{(2d + 1)!!^3}{8^{d+1} d!} (X - d + \frac{3}{2}) \frac{\Gamma(\frac{X+1}{2} - d)^3 \Gamma(\frac{X+3}{2})}{\Gamma(\frac{X+4}{2})^3 \Gamma(\frac{X+2}{2} - d)}, \quad (145)$$

where the right-hand side is understood as its asymptotic expansion in X^{-1} as $X \rightarrow \infty$. This formula together with $W_d(X) \in X^{-2d-2} \mathbb{Q}[[X^{-1}]]$ uniquely determines $W_d(X)$. It is easy to verify that the right-hand side of (141), with $A_j(d)$ defined by (142), has the same recursive property. Hence the first statement of Proposition 7 is proved.

Let us now prove (143). Denote by $r_d(X) \in \mathbb{Q}[[X^{-1}]]$ the asymptotic expansion of the left-hand side of (142). Using the property $\Gamma(z + 1) = z\Gamma(z)$, we see that

$$(X - 2d - 1)^3 r_{d+1}(X) = (X - 2d) r_d(X). \quad (146)$$

From (142) and (146), we obtain the following two recursions for $A_j(d)$:

$$\begin{aligned} A_{j+1}(d + 1) - A_{j+1}(d) &= 2(d + j + 1)(2d + j + 2) A_j(d) \\ &\quad - ((2d + j + 3)(2d + 3) + j^2) A_j(d + 1), \end{aligned} \quad (147)$$

$$-j^3 A_j(d + 1) + (2d - j + 3) A_{j+1}(d + 1) - (2d + j + 3) A_{j+1}(d) = 0. \quad (148)$$

Here $j \geq 0$, and we make the convention that $A_0(d) \equiv 0$. Notice that equations (147)–(148), together with the initial value $A_1(d) \equiv 1$, uniquely determine all $A_j(d)$. It is easy to verify that the right-hand side of (143) satisfies (147)–(148) and takes value 1 when $j = 1$. This completes the proof of (143). \square

It follows from (138) and (141) that

$$\widehat{c}_k(\mathbf{e}_d) = -\frac{(2d + 1)!!^3}{2^{d+1} d!} \sum_{j=1}^{\lfloor \frac{k}{2} \rfloor - d} \frac{A_j(d)}{d + j} \sum_{l=0}^{k-2d-2j} \binom{k-1}{l} (-j)^l S(k - l - 1, 2d + 2j - 1), \quad (149)$$

where $S(n, k)$ are the Stirling numbers of the second kind.

It is interesting to notice that the polynomial $A_j(d)$ is the product of $(d + 3/2)_{\lfloor j/2 \rfloor}$ and a polynomial of degree $\lfloor (j-1)/2 \rfloor$, which can be easily proved by using (143). For the reader's convenience we provide the first few $A_j(d)$:

$$\begin{aligned} A_1(d) &= 1, & A_2(d) &= -\frac{1}{2}(2d+3), & A_3(d) &= \frac{1}{8}(2d+3)(10d+21), \\ A_4(d) &= -\frac{3}{16}(2d+3)(2d+5)(14d+31). \end{aligned}$$

We also remark that although $W_d(X)$ is defined as the asymptotics of two-point BGW numbers it also gives information about multi-point BGW numbers. Indeed, from (127) and Theorem 2 we can deduce that for a given $n \geq 2$ and for $\mathbf{d} = (d_1, \dots, d_n)$ with $1 \leq d_1 \leq \dots \leq d_{n-1}$ fixed,

$$\widehat{C}(\mathbf{d}) = 1 - \sum_{j=1}^{n-1} W_{d_j}(X(\mathbf{d})) + O(X(\mathbf{d})^{-2d_1-2d_2-3}), \quad d_n \rightarrow \infty. \quad (150)$$

Similar to (137), for $n \geq 1$, $\lambda = (\lambda_1, \dots, \lambda_{n-1})$, define $W_\lambda(X) \in \mathbb{Q}[[X^{-1}]]$ via

$$\widehat{C}(\lambda, d_n) \sim - \sum_{I \subset \{1, \dots, n-1\}} W_{\lambda_I}(X(\lambda, d_n)), \quad d_n \rightarrow \infty, \quad (151)$$

with $W_\emptyset(X) = -1$. Then we have the following proposition.

Proposition 8. *We have*

$$\sum_{k \geq 0} \frac{\widehat{c}_k(\mathbf{p})}{X^k} = - \sum_{q_1, q_2, \dots \geq 0} W_{1^{q_1}, 2^{q_2}, \dots}(X) \prod_{i=1}^{\infty} \binom{p_i}{q_i}, \quad (152)$$

where both sides are understood as elements in $\mathbb{Q}[p_1, p_2, \dots][[X^{-1}]]$. Moreover, the power series $W_\lambda(X) \in X^{-2|\lambda|-\ell(\lambda)-1} \mathbb{Q}[[X^{-1}]]$.

Proof. Using (127), Theorem 2 and (151), we obtain (152). Using (28)–(29) and comparing degrees on both sides of (152), we obtain that $W_\lambda(X) \in X^{-2|\lambda|-\ell(\lambda)-1} \mathbb{Q}[[X^{-1}]]$. \square

We list a few examples of $W_\lambda(X)$ below:

$$\begin{aligned} W_{1,1}(X) &= \frac{1701}{4X^7} + \frac{380295}{64X^8} + \frac{832815}{16X^9} + \frac{2935197}{8X^{10}} + \dots, \\ W_{1,2}(X) &= \frac{388125}{16X^9} + \frac{83804625}{128X^{10}} + \frac{1336975875}{128X^{11}} + \frac{131751025875}{1024X^{12}} + \dots, \\ W_{1,1,1}(X) &= \frac{1754703}{8X^{10}} + \frac{245520639}{32X^{11}} + \frac{79688662083}{512X^{12}} + \frac{615031348329}{256X^{13}} + \dots, \\ W_{1,1,2}(X) &= \frac{779513625}{32X^{12}} + \frac{21043769625}{16X^{13}} + \frac{41031922798125}{1024X^{14}} + \dots. \end{aligned}$$

We note that Proposition 8 together with (127) and Theorem 2 implies (150) and formulas like

$$\widehat{C}(\mathbf{d}) = 1 - \sum_{i=1}^{n-1} W_{d_i}(X(\mathbf{d})) - \sum_{1 \leq i < j \leq n-1} W_{d_i, d_j}(X(\mathbf{d})) + O(X(\mathbf{d})^{-2d_1-2d_2-2d_3-4}), \quad (153)$$

for $n \geq 1$, $1 \leq d_1 \leq \dots \leq d_{n-1}$ fixed and $d_n \rightarrow \infty$. Based on numerical experiments, we conjecture that the leading term of $W_\lambda(X)$ is

$$2(2|\lambda| + \ell(\lambda))! C(\lambda) X^{-2|\lambda| - \ell(\lambda) - 1}. \quad (154)$$

8. SUBEXPONENTIAL ASYMPTOTIC TERMS

In Section 1 we have described the discovery of the conjectural asymptotic formula (31). In this section we study subexponential asymptotics more systematically.

Let us look, for a fixed partition $\boldsymbol{\mu} = (1 \leq \mu_1 \leq \dots \leq \mu_n)$, at the asymptotics of $1 - \widehat{C}(\mathbf{d})$ with $\mathbf{d} = \boldsymbol{\mu}d = (\mu_1 d, \dots, \mu_n d)$ as $d \rightarrow \infty$. Similar to (31), we find (based on computations using formula (60)) the following conjectural asymptotic formulas as $d \rightarrow \infty$:

$$\begin{aligned} 1 - \widehat{C}(\mu_1 d, \mu_2 d) &\sim \frac{2}{\sqrt{\pi d}} \frac{\mu_1^{2\mu_1 d + 3/2} \mu_2^{2\mu_2 d + 3/2}}{|\boldsymbol{\mu}|^{2|\boldsymbol{\mu}|d + 7/2}}, \\ 1 - \widehat{C}(\mu_1 d, \mu_2 d, \mu_3 d) &\sim \frac{2 p_{\mu_1}}{\sqrt{\pi d}} \frac{(\mu_2 + \mu_3)^{2(\mu_2 + \mu_3)d + 5/2} \mu_1^{2\mu_1 d + 3/2}}{|\boldsymbol{\mu}|^{2|\boldsymbol{\mu}|d + 9/2}}, \\ 1 - \widehat{C}(\mu_1 d, \mu_2 d, \mu_3 d, \mu_4 d) &\sim \frac{2 p_{\mu_1}}{\sqrt{\pi d}} \frac{(\mu_2 + \mu_3 + \mu_4)^{2(\mu_2 + \mu_3 + \mu_4)d + 7/2} \mu_1^{2\mu_1 d + 3/2}}{|\boldsymbol{\mu}|^{2|\boldsymbol{\mu}|d + 11/2}}. \end{aligned}$$

With the help of these formulas and based on more computations, we obtain the following conjectural asymptotic formula: for any fixed $n \geq 2$,

$$1 - \widehat{C}(d_1, \dots, d_n) \sim \left(\frac{1}{2}\right)^{\delta_{n,2}} \sum_{j=1}^n \frac{2}{\pi X(\mathbf{d})} \left(\frac{X(\mathbf{d})}{2d_j + 1}\right)^{-1}, \quad \min_{1 \leq j \leq n} \{d_j\} \rightarrow \infty. \quad (155)$$

Since

$$\frac{2}{\pi} (2d+1)! \sim \frac{(2d+1)!!^3}{2^{d+1} (d+1)!}, \quad d \rightarrow \infty,$$

we can rewrite the conjectural formula (155) equivalently as

$$1 - \widehat{C}(d_1, \dots, d_n) \sim \left(\frac{1}{2}\right)^{\delta_{n,2}} \sum_{j=1}^n \frac{(2d_j+1)!!^3}{2^{d_j+1} (d_j+1)!} \frac{1}{(X(\mathbf{d}) - 2d_j)_{2d_j+2}}, \quad \min_{1 \leq j \leq n} \{d_j\} \rightarrow \infty. \quad (156)$$

Remark 5. Notice that the form $\frac{(2d+1)!!^3}{2^{d+1} (d+1)!}$ in (156) also appears in the leading coefficients of $W_d(X)$ in (141), so we obtain from (132) that formula (156) holds true even if some d_j 's are not large (but requires $X(\mathbf{d}) \rightarrow \infty$). So we guess that

$$1 - \widehat{C}(\mathbf{d}) = \left(\sum_{j=1}^n \frac{(2d_j+1)!!^3}{2^{d_j+1} (d_j+1)!} \frac{1}{(X(\mathbf{d}) - 2d_j)_{2d_j+2}} \right) (1 + o(1)) \quad (157)$$

for any fixed $n \geq 3$ and for $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$, uniformly as $X(\mathbf{d}) \rightarrow \infty$. The conjectural formula (157) implies that, for any two partitions \mathbf{d} and \mathbf{d}' of the same

length and the same weight,

$$\widehat{C}(\mathbf{d}) - \widehat{C}(\mathbf{d}') = \sum_{m \geq 0} \frac{(2m+1)!!^3}{2^{m+1}(m+1)!} \frac{p_m - p'_m}{(X(\mathbf{d}) - 2m)_{2m+2}} (1 + o(1)), \quad (158)$$

as $X(\mathbf{d}) = X(\mathbf{d}') \rightarrow \infty$. We note that there is a coherent consistence between formula (158) and Conjecture 2. Another point is that formula (158) also explains the phenomenon (cf. (24)) described in Section 1. As a further example, we have

$$\begin{aligned} \widehat{C}(2, 3, 14, 19) &= 0.99999999969689849693814650552875212296 \dots, \\ \widehat{C}(2, 3, 15, 18) &= 0.99999999969689849693814752270899360002 \dots, \end{aligned}$$

whose difference is about 1.01718×10^{-24} and the prediction gives 0.92432×10^{-24} with error less than 10 percent.

Now we compute the subleading terms for the subexponential asymptotics (31), (156). Based on the numerical experiments, we find that the error between 1 and $\widehat{C}(d^n)$ for fixed $n \geq 2$ is of the form

$$1 - \widehat{C}(d^n) \sim Y_n(d) \left(1 + \frac{b_1(n)}{d} + \frac{b_2(n)}{d^2} + \dots \right), \quad d \rightarrow \infty, \quad (159)$$

where

$$Y_n(d) = \left(\frac{1}{2} \right)^{\delta_{n,2}} \sqrt{\frac{4(n-1)}{\pi n d}} \left(\frac{(n-1)^{n-1}}{n^n} \right)^{2d+1}, \quad (160)$$

and $b_1(n), b_2(n), \dots$ are rational number whose numerical values become a little simpler if we set

$$L_n := 24 \log(1 + b_1(n)x + b_2(n)x^2 + \dots), \quad (161)$$

in which the first few values are given by

$$\begin{aligned} L_n &= \frac{-11n^2 - n + 7}{n^2 - n} x + \frac{14n^3 - 16n^2 + n + 7}{2n(n-1)^2} x^2 \\ &\quad + \frac{-721n^6 + 1803n^5 - 1953n^4 + 901n^3 - 243n^2 + 93n - 31}{120n^3(n-1)^3} x^3 + \dots \end{aligned} \quad (162)$$

Recall the discovery in Section 7 that the asymptotics for two-point BGW numbers are building blocks for the higher-point numbers, at least when all but one d_j 's are fixed. The following observation generalizes this for the subexponential asymptotics.

Observation. First of all, we find that the ratio of $\widehat{C}(d, d, d, d) - 1$ and $\widehat{C}(d, 3d+1) - 1$, which are both exponentially small, is asymptotically equal to 4 to all orders in d^{-1} . Here the number $3d+1$ is chosen such that these two BGW numbers have the same argument $X(\mathbf{d})$. More generally, we conjecture that the ratio of $\widehat{C}(d^n) - 1$ and $\widehat{C}(d, d') - 1$ is asymptotically equal to n for $n \geq 3$, where d' is defined by $2d' + 1 = (n-1)(2d+1)$. We notice that if n is odd, then d' is not an integer, but we can still define $C(d, d')$ by the two point formula (69) with the definition of F_h there replaced by

$$F_h := \frac{\Gamma(h + \frac{1}{2})^3}{\pi^{3/2} \Gamma(h+1)}. \quad (163)$$

Now let us focus on the asymptotics for $1 - \widehat{C}(d_1, d_2)$ again. Recall that if d_1 is fixed, this asymptotics is exactly $W_{d_1}(X)$ explicitly given in (141). Another important observation is that, the right-hand side of (141) makes sense (i.e. each term has lower order than the previous one) even if X and d are proportional and are in the region $X - 2d \gg 0$. Note that this property will not be true if we write $W_d(X)$ in another basis, e.g. the powers of X^{-1} as in (140) or the basis $1/(X-1)_k^-$ ($k = 1, 2, \dots$) used in [18]. Therefore, for any fixed integer $N \geq 1$, we define a function $W(N; d, X)$ by

$$W(N; d, X) = \frac{(2d+1)!!^3}{2^{d+1}d!} \sum_{j=1}^N \frac{A_j(d)}{d+j} \frac{1}{(X-2d-j+1)_{2d+2j}}, \quad (164)$$

where $A_j(d)$ are polynomials defined in (143). We note that $W(N; d, X)$ is well defined in the region $X - 2d > N$. According to the previous observations, we conjecture that

$$1 - \widehat{C}(d_1, d_2) = W(N; d_1, X(d_1, d_2))(1 + O(X(d_1, d_2)^{-N})), \quad X(d_1, d_2) \rightarrow \infty \quad (165)$$

holds for any fixed $N \geq 1$ and for $d_1 \leq d_2$. Now for general $\mathbf{d} = (d_1, \dots, d_n)$ of length $n \geq 3$, we have the following conjecture.

Conjecture 4. *For any fixed $n \geq 3$, any fixed $N \geq 0$ and for $\mathbf{d} = (d_1, \dots, d_n)$ satisfying $\min_{1 \leq i < j \leq n} \{d_i + d_j\} \geq N/2$, we have the asymptotics:*

$$1 - \widehat{C}(\mathbf{d}) = \sum_{i=1}^n W(N; d_i, X(\mathbf{d})) (1 + O(X(\mathbf{d})^{-N})), \quad X(\mathbf{d}) \rightarrow \infty. \quad (166)$$

As a consequence of Conjecture 4, for fixed $n \geq 0$, the asymptotic expansion of $1 - \widehat{C}(d^n)$ is explicitly given by

$$1 - \widehat{C}(d^n) = n W(N; d, n(2d+1))(1 + O(d^{-N})), \quad d \rightarrow \infty, \quad (167)$$

for any fixed $N \geq 0$. This explicitly gives all the numbers $b_1(n), b_2(n), \dots$ in (159), and the first three of them coincide with those in (162).

9. APPLICATION TO THE PAINLEVÉ II HIERARCHY

In this section, we will discuss the connections between the BGW numbers and two famous Painlevé hierarchies. We begin with the Painlevé XXXIV hierarchy (cf. [5, 9]), by which we mean the following family of ODEs:

$$2u + t \frac{du}{dt} - (2d+1) \frac{d}{dt} \left(m_d \left(u, \frac{du}{dt}, \frac{d^2u}{dt^2}, \dots, \frac{d^{2d}u}{dt^{2d}} \right) \right) = 0, \quad (168)$$

where $d \geq 1$, and m_d are the polynomials defined in (32), (33). The case with $d = 1$ agrees with the Painlevé XXXIV equation (14).

Lemma 6. *For each $d \geq 1$, there exists a unique formal solution to equation (168) of the form*

$$u(t) = \sum_{n \geq 0} \frac{A_{d,n}}{t^{(2d+1)n+2}}, \quad A_{d,0} = \frac{1}{8}. \quad (169)$$

Proof. If we assign degrees

$$\deg u_i = i + 2, \quad i \geq 0, \quad (170)$$

then the polynomials m_d are homogeneous of degree $2d + 2$. The lemma follows. \square

Lemma 7. *The coefficients $A_{d,n}$ in (169) are related to the BGW correlators by*

$$A_{d,n} = \frac{((2d+1)n+1)!}{2^{2nd+1}(2d+1)!!^n n!} C(d^n), \quad (171)$$

where $C(d_1, \dots, d_n)$ is defined in (6).

Proof. As it was done in [5], dividing both sides of the $m = 0$ case of (42) by Z and differentiating the resulting equality twice with respect to t_0 and using (41), one obtains

$$2u - (1-t_0)u_{t_0} + \sum_{k \geq 1} (2k+1)t_k \partial_{t_0}(m_k) = 0, \quad (172)$$

where we recall that $u := \partial_{t_0}^2(\log Z)$. Specializing $\mathbf{t} = \mathbf{t}^* = (t_0^*, t_1^*, t_2^*, \dots)$ in (172) with

$$t_d^* = 1, \quad t_i^* = 0 \ (i \neq 0, i \neq d), \quad (173)$$

we find

$$2u(\mathbf{t}^*) - (1-t_0)u_{t_0}(\mathbf{t}^*) + (2d+1)\partial_{t_0}(m_d(u(\mathbf{t}^*), u_{t_0}(\mathbf{t}^*), \dots)) = 0. \quad (174)$$

The lemma is proved by noticing that

$$u|_{\mathbf{t}=\mathbf{t}^*} = \sum_{n \geq 1} \frac{C(d^n)((2d+1)n+1)!}{2^{2nd+1}(1-t_0)^{(2d+1)n+2}(2d+1)!!^n n!} + \frac{1}{8(1-t_0)^2} \quad (175)$$

and by putting $t = 1 - t_0$. \square

It is convenient to work with another normalization of the Painlevé XXXIV hierarchy:

$$2^{2d+1}(2d+1)!! \partial_X(m_d(Y/2, Y_X/2, Y_{XX}/2, \dots)) - XY_X - 2Y = 0, \quad d \geq 1, \quad (176)$$

which is related to (168) by the rescalings

$$t = \frac{1}{2} \left(\frac{(2d-1)!!}{2} \right)^{-1/(2d+1)} X, \quad u = 2 \left(\frac{(2d-1)!!}{2} \right)^{2/(2d+1)} Y. \quad (177)$$

The formal solution of interest (a solution to (176)) now has the form

$$Y(X) = \sum_{n \geq 0} \frac{y_{d,n}}{X^{(2d+1)n+2}}, \quad y_{d,0} = \frac{1}{4}. \quad (178)$$

Theorem 4. *For each $d \geq 1$, the coefficients $y_{d,n}$ of the unique formal solution $Y(X)$ given in (178) to the Painlevé XXXIV hierarchy (176) have the following asymptotics:*

$$y_{d,n} \sim \frac{1}{\pi} \frac{((2d+1)n+1)!}{(2d+1)^n n!}, \quad n \rightarrow \infty. \quad (179)$$

Proof. By using (171), (177) and Theorem 1 we obtain (179). \square

Corollary 3. *For each $d \geq 1$, we have*

$$y_{d,n} \sim \frac{1}{\pi} \frac{((2d+1)n+1)!}{(2d+1)^n n!} \left(1 + \frac{r_1(d)}{n} + \frac{r_2(d)}{n^2} + \dots \right) \quad (n \rightarrow \infty), \quad (180)$$

with explicitly computable coefficients $r_k(d) \in \mathbb{Q}$.

Proof. Let us first show, without using Theorem 4, that

$$y_{d,n} \sim A \frac{((2d+1)n+1)!}{(2d+1)^n n!} \left(1 + \frac{r_1(d)}{n} + \frac{r_2(d)}{n^2} + \dots\right), \quad n \rightarrow \infty, \quad (181)$$

for some nonzero constant A . Using (176), (178) and using the homogeneity of m_d , we obtain a recursion that expresses $(2d+1)n y_{d,n}$ by an element in

$$\text{span}_{\mathbb{Q}} \left\{ \sum_{n_1+\dots+n_k=n-1} \prod_{j=1}^k (((2d+1)n_j+2)_{i_j} y_{d,n_j}) \mid k \geq 1, \mathbf{i} \in (\mathbb{Z}_{\geq 0})^k, |\mathbf{i}| + 2k = 2d+3 \right\}.$$

The leading asymptotics in (181) of $y_{d,n}$ can be deduced from the linear terms

$$(2d+1)n y_{d,n} - ((2d+1)(n-1)+2)_{2d+1} y_{d,n-1}.$$

It follows that

$$\prod_{j=1}^k (y_{d,n_j} ((2d+1)n_j+2)_{i_j}) = O\left(y_{d,n} \frac{n!}{((2d+1)n+1)!} \prod_{j=1}^k \frac{((2d+1)n_j+1+i_j)!}{n_j!}\right) \quad (182)$$

for each $k \geq 0$, $i_1 + \dots + i_k + 2k = 2d+2$ and $n_1 + \dots + n_k = n-1$. By using the logarithmic convexity of the function $((2d+1)n+1+i_j)!/n!$, we obtain the right-hand side of (182) is $O(y_{d,n} n^{-2d(h-1)-2k+2})$ when $n_1, \dots, n_k \leq n-h$. This shows that up to any relative power in n^{-1} , the recursion that $y_{d,n}$ satisfies is linear and of finite order. This proves (181). The determination of $A = 1/\pi$ follows from Theorem 4. \square

Corollary 4. *For each $d \geq 1$, we have the asymptotic expansion*

$$\widehat{C}(d^n) \sim 1 + \frac{a_1(d)}{X(d^n)} + \frac{a_2(d)}{X(d^n)^2} + \dots \quad (n \rightarrow \infty) \quad (183)$$

with explicitly computable $a_k(d) \in \mathbb{Q}$, the first three cases being

$$\begin{aligned} \widehat{C}(1^n) &\sim 1 - \frac{9}{8(3n)^3} - \frac{9}{4(3n)^4} - \frac{219}{8(3n)^5} + \dots, \\ \widehat{C}(2^n) &\sim 1 - \frac{225}{16(5n)^5} - \frac{2025}{16(5n)^6} - \frac{96075}{128(5n)^7} + \dots, \\ \widehat{C}(3^n) &\sim 1 - \frac{55125}{128(7n)^7} - \frac{275625}{32(7n)^8} - \frac{3340575}{32(7n)^9} + \dots. \end{aligned}$$

Proof. Formula (183) is proved by using (26), (171), (177), (178), Corollary 3 and Stirling's formula. \square

By (149) and the conjectural formula (134), one can deduce that for any fixed $d \geq 1$,

$$1 - \widehat{C}(d^n) \sim \frac{(2d+1)!!^2 (2d-1)!!}{2^{d+1} (d+1)! X(d^n)^{2d+1}} \quad (X(d^n) = (2d+1)n) \quad (184)$$

to leading order as $n \rightarrow \infty$. This conjectural formula is consistent with Corollary 4. Compare also with the asymptotic formula (31) when n is fixed and d tends to infinity.

Proof of Theorem 3. According to [9] (cf. [20]), performing the following invertible transformation

$$Y = V_X - V^2, \quad (185)$$

$$V = -\frac{2^{2d-1}(2d-1)!! \partial_X(m_{d-1}(\frac{Y}{2}, \frac{Y_X}{2}, \dots)) - \alpha_d}{2^{2d}(2d-1)!! m_{d-1}(\frac{Y}{2}, \frac{Y_X}{2}, \dots) - X} \quad (186)$$

on the Painlevé XXXIV hierarchy yields the Painlevé II hierarchy (34).

We note that in general, α_d could be an arbitrary constant. But the particular solution $V(X)$ derived by the above transformation of the power series in (178) only solves the Painlevé II hierarchy for the parameter $\alpha_d = \frac{1}{2}$. This can be seen by comparing the coefficients of X^{-2} on both sides of (185), and by noticing that $V(X)$ has the leading term $-\alpha_d/X$. So $Y(X)$ defined in (178) corresponds to the formal solution (35) to (34) with $\alpha_d = 1/2$.

To prove formula (36), we note that the transformation (185) gives the following relations between $v_{d,n}$ and $y_{d,n}$:

$$y_{d,n} = ((2d+1)n+1)v_{d,n} - \sum_{n_1+n_2=n} v_{d,n_1} v_{d,n_2}. \quad (187)$$

Using the asymptotics (179) of $y_{d,n}$ and facts about asymptotics of very rapidly divergent series (cf. [8]), it is easy to show that $v_{d,n}$ is asymptotically equal to $y_{d,n}/((2d+1)n-1)$, as claimed in (36). \square

Similar to Corollary 3, we have the following

Corollary 5. *For each $d \geq 1$, the coefficients $v_{d,n}$ of the formal solution to the Painlevé II hierarchy has the following asymptotic expansion:*

$$v_{d,n} \sim \frac{1}{\pi} \frac{((2d+1)n+1)!}{(2d+1)^n n!} \left(1 + \frac{s_1(d)}{n} + \frac{s_2(d)}{n^2} + \dots\right), \quad n \rightarrow \infty, \quad (188)$$

with explicitly computable coefficients $s_k(d) \in \mathbb{Q}$.

Proof. Similar to the proof of Corollary 3. \square

10. APPLICATION TO BGW-KAPPA NUMBERS

For $g, m \geq 1$, $n \geq 0$, $d_1, \dots, d_n \geq 0$, define the BGW-kappa numbers by

$$\left\langle \kappa_1^m \prod_{j=1}^n \tau_{d_j} \right\rangle_g^\Theta := \int_{\mathcal{M}_{g,n}} \kappa_1^m \psi_1^{d_1} \dots \psi_n^{d_n} \Theta_{g,n}, \quad (189)$$

which vanishes unless $m + d_1 + \dots + d_n = g - 1$. One could of course also include powers of other kappa's in (189), but we will study only the integrals with a power of κ_1 and here we use the terminology ‘‘BGW-kappa numbers’’ for convenience. Another name that can be found in the literature is super JT gravity. According to [38], these numbers are related to the volumes of moduli spaces of super hyperbolic surfaces, called Stanford–Witten volumes $V_{g,n}^\Theta(\mathbf{L})$. This relation is given by

$$V_{g,n}^\Theta(L_1, \dots, L_n) = \sum_{\substack{m, d_1, \dots, d_n \geq 0 \\ m + d_1 + \dots + d_n = g - 1}} \left\langle \kappa_1^m \tau_{d_1} \dots \tau_{d_n} \right\rangle_g^\Theta \frac{(2\pi^2)^m}{m!} \prod_{j=1}^n \frac{L_j^{2d_j}}{2^{d_j} d_j!}. \quad (190)$$

As proved in [38] (cf. [28, 32, 35]), the BGW-kappa numbers can be expressed in terms of the BGW numbers as follows:

$$\langle \kappa_1^m \prod_{j=1}^n \tau_{d_j} \rangle_g^\Theta = \sum_{l=1}^m \frac{(-1)^{m-l}}{l!} \sum_{\substack{m_1, \dots, m_l \geq 1 \\ m_1 + \dots + m_l = m}} \binom{m}{m_1, \dots, m_l} \left\langle \prod_{j=1}^n \tau_{d_j} \prod_{i=1}^l \tau_{m_i} \right\rangle_g^\Theta. \quad (191)$$

Using (191) we compute a few BGW-kappa numbers in Table 3. It follows from (191) and the Integrality Conjecture for the BGW numbers that the BGW-kappa numbers are integral away from powers of 2. We refer to the latter statement as the Integrality Conjecture for the BGW-kappa numbers.

$g = 2$	
$\langle \kappa_1 \rangle^\Theta = \frac{3}{128} \approx 2.34 \times 10^{-2}$	
$g = 3$	
$\langle \kappa_1 \tau_1 \rangle^\Theta = \frac{63}{512} \approx 1.23 \times 10^{-1}$	$\langle \kappa_1^2 \rangle^\Theta = \frac{111}{1024} \approx 1.08 \times 10^{-1}$
$g = 4$	
$\langle \kappa_1 \tau_1^2 \rangle^\Theta = \frac{7221}{2048} \approx 3.53$	$\langle \kappa_1^2 \tau_1 \rangle^\Theta = \frac{106911}{32768} \approx 3.26$
$\langle \kappa_1 \tau_2 \rangle^\Theta = \frac{8625}{32768} \approx 2.63 \times 10^{-1}$	$\langle \kappa_1^3 \rangle^\Theta = \frac{45093}{16384} \approx 2.75$
$g = 5$	
$\langle \kappa_1 \tau_1^3 \rangle^\Theta = \frac{4825971}{16384} \approx 2.95 \times 10^2$	$\langle \kappa_1^2 \tau_2 \rangle^\Theta = \frac{1974135}{131072} \approx 1.51 \times 10$
$\langle \kappa_1 \tau_1 \tau_2 \rangle^\Theta = \frac{524925}{32768} \approx 1.60 \times 10$	$\langle \kappa_1^3 \tau_1 \rangle^\Theta = \frac{16199169}{65536} \approx 2.47 \times 10^2$
$\langle \kappa_1 \tau_3 \rangle^\Theta = \frac{44835}{65536} \approx 6.84 \times 10^{-1}$	$\langle \kappa_1^4 \rangle^\Theta = \frac{53483271}{262144} \approx 2.04 \times 10^2$
$\langle \kappa_1^2 \tau_1^2 \rangle^\Theta = \frac{9127017}{32768} \approx 2.79 \times 10^2$	
$g = 6$	
$\langle \kappa_1 \tau_1^4 \rangle^\Theta = \frac{3540311739}{65536} \approx 5.40 \times 10^4$	$\langle \kappa_1^2 \tau_1 \tau_2 \rangle^\Theta = \frac{1155623625}{524288} \approx 2.20 \times 10^3$
$\langle \kappa_1 \tau_1^2 \tau_2 \rangle^\Theta = \frac{605705625}{262144} \approx 2.31 \times 10^3$	$\langle \kappa_1^2 \tau_3 \rangle^\Theta = \frac{151428375}{2097152} \approx 7.22 \times 10$
$\langle \kappa_1 \tau_2^2 \rangle^\Theta = \frac{55787625}{524288} \approx 1.06 \times 10^2$	$\langle \kappa_1^3 \tau_1^2 \rangle^\Theta = \frac{386376633}{8192} \approx 4.72 \times 10^4$
$\langle \kappa_1 \tau_1 \tau_3 \rangle^\Theta = \frac{19922175}{262144} \approx 7.60 \times 10$	$\langle \kappa_1^3 \tau_2 \rangle^\Theta = \frac{4184142525}{2097152} \approx 2.00 \times 10^3$
$\langle \kappa_1 \tau_4 \rangle^\Theta = \frac{8831025}{4194304} \approx 2.11$	$\langle \kappa_1^4 \tau_1 \rangle^\Theta = \frac{171037302471}{4194304} \approx 4.08 \times 10^4$
$\langle \kappa_1^2 \tau_1^3 \rangle^\Theta = \frac{13555541331}{262144} \approx 5.17 \times 10^4$	$\langle \kappa_1^5 \rangle^\Theta = \frac{69673098483}{2097152} \approx 3.32 \times 10^4$

TABLE 3. BGW-kappa numbers up to genus 6

We also provide a table (see Table 4) for the normalized BGW-kappa numbers $C(m; \mathbf{d})$ defined in (38) with $g \leq 7$. The data for $C(1; \mathbf{d})$ is omitted since $C(1; \mathbf{d}) = C(1, \mathbf{d})$. As before, we have listed in Table 4 the smallest common denominator $D = D_g$ of these numbers for each g and then tabulated the integer $D C(m; \mathbf{d})$ in the last column. Inspired by Conjecture 2, and based on Table 4 and further numerical experiments, we also conjecture that the function $(m, \mathbf{d}) \mapsto C(m; \mathbf{d})$ for partitions (m, \mathbf{d}) of $g - 1$ is monotone with respect to the ordering that $(m, \mathbf{d}) \prec (m', \mathbf{d}')$ if $m > m'$, or $m = m'$ and $\mathbf{d} \prec \mathbf{d}'$.

We now give a proof of Proposition 1.

$g = 3, \quad D = 1280$			
$C(2; \emptyset)$	$\frac{333}{1280}$	0.260156	333
$g = 4, \quad D = 1146880$			
$C(3; \emptyset)$	$\frac{135279}{573440}$	0.235908	270558
$C(2; 1)$	$\frac{45819}{163840}$	0.279657	320733
$g = 5, \quad D = 252313600$			
$C(4; \emptyset)$	$\frac{53483271}{252313600}$	0.211971	53483271
$C(3; 1)$	$\frac{2314167}{9011200}$	0.256810	64796676
$C(2; 2)$	$\frac{131609}{458752}$	0.286885	72384950
$C(2; 1, 1)$	$\frac{9127017}{31539200}$	0.289386	73016136
$g = 6, \quad D = 734737203200$			
$C(5; \emptyset)$	$\frac{69673098483}{367368601600}$	0.189654	139346196966
$C(4; 1)$	$\frac{24433900353}{104962457600}$	0.232787	171037302471
$C(3; 2)$	$\frac{278942835}{1049624576}$	0.265755	195259984500
$C(3; 1, 1)$	$\frac{386376633}{1435033600}$	0.269246	197824836096
$C(2; 3)$	$\frac{3365075}{11534336}$	0.291744	214355277500
$C(2; 1, 2)$	$\frac{5926275}{20185088}$	0.293597	215716410000
$C(2; 1, 1, 1)$	$\frac{13555541331}{45921075200}$	0.295192	216888661296
$g = 7, \quad D = 399697038540800$			
$C(6; \emptyset)$	$\frac{1057428386631}{6245266227200}$	0.169317	67675416744384
$C(5; 1)$	$\frac{1196989428069}{5709957693440}$	0.209632	83789259964830
$C(4; 2)$	$\frac{103748833683}{427483463680}$	0.242697	97005159493605
$C(4; 1, 1)$	$\frac{2242040330133}{9084023603200}$	0.246811	98649774525852
$C(3; 3)$	$\frac{31418131}{115343360}$	0.272388	108872620991680
$C(3; 1, 2)$	$\frac{80848213893}{293894881280}$	0.275092	109953570894480
$C(3; 1, 1, 1)$	$\frac{6931945897497}{24981064908800}$	0.277488	110911134359952
$C(2; 4)$	$\frac{354207573}{1199570944}$	0.295279	118021963323600
$C(2; 1, 3)$	$\frac{222438209}{749731840}$	0.296690	118586257982080
$C(2; 2, 2)$	$\frac{4360002121}{14694744064}$	0.296705	118592057691200
$C(2; 1, 1, 2)$	$\frac{3184112229}{10687086592}$	0.297940	119085797364600
$C(2; 1, 1, 1, 1)$	$\frac{466903889307}{1561316556800}$	0.299045	119527395662592

TABLE 4. Numerical data for $C(m, \mathbf{d})$ with $g \leq 7$

Proof of Proposition 1. It follows from (191) and (38) that

$$C(m; \mathbf{d}) = 3^m \sum_{l=1}^m \frac{(-1)^{m-l}}{l!} \sum_{\substack{m_1, \dots, m_l \geq 1 \\ m_1 + \dots + m_l = m}} \binom{m}{m_1, \dots, m_l} \\ \times \frac{C(d_1, \dots, d_n, m_1, \dots, m_l)}{(X(m; \mathbf{d}) - m + l)_{m-l} \prod_{i=1}^l (2m_i + 1)!!}. \quad (192)$$

For fixed $m \geq 0$, the summation in the RHS of (192) is finite. The summand corresponding to $l = m$, $m_1 = \dots = m_l = 1$ contributes the leading term $C(\mathbf{d}, 1^m)$, which by Theorem 1 equals $1/\pi + O(1/g(m; \mathbf{d}))$ uniformly as $g(m; \mathbf{d}) \rightarrow \infty$. The rest summands equal $O(1/(X(\mathbf{m}; \mathbf{d}) - m + l)_{m-l})$, which also equals $O(1/g(m; \mathbf{d}))$ uniformly as $g(m; \mathbf{d}) \rightarrow \infty$. This proves the proposition. \square

For the special case when $n \geq 1$ and $d_1, \dots, d_{n-1} \geq 0$ are all fixed, Proposition 1 is analogous to a result given by Liu–Xu [33] for Witten’s intersection numbers.

Remark 6. In another direction, we also study the large- g asymptotics when m is no longer fixed. In particular, we find that $\langle \kappa_1^m \tau_{d_1} \dots \tau_{d_n} \rangle_g$ has a simple asymptotic formula when $\mathbf{d} = (d_1, \dots, d_n) \in (\mathbb{Z}_{\geq 0})^n$ is fixed and $g \rightarrow \infty$:

$$\langle \kappa_1^{g-1-|\mathbf{d}|} \tau_{d_1} \dots \tau_{d_n} \rangle_g^\Theta \sim \frac{\pi^{2|\mathbf{d}|+n-2} 2^{g-1-3|\mathbf{d}|}}{3^{3g-\frac{7}{2}-|\mathbf{d}|+n} \prod_{j=1}^n (2d_j + 1)!!} (3g - 4 - |\mathbf{d}| + n)!, \quad (193)$$

with the first four cases being

$$\langle \kappa_1^{g-1} \rangle_g^\Theta \sim \frac{2^{g-1}}{\pi^2 3^{3g-\frac{7}{2}}} (3g - 4)!, \quad \langle \kappa_1^{g-2} \tau_1 \rangle_g^\Theta \sim \frac{\pi 2^{g-4}}{3^{3g-\frac{5}{2}}} (3g - 4)!, \\ \langle \kappa_1^{g-3} \tau_2 \rangle_g^\Theta \sim \frac{\pi^3 2^{g-7}}{5 \cdot 3^{3g-\frac{7}{2}}} (3g - 5)!, \quad \langle \kappa_1^{g-3} \tau_1^2 \rangle_g^\Theta \sim \frac{\pi^4 2^{g-7}}{3^{3g-\frac{3}{2}}} (3g - 4)!,$$

as $g \rightarrow \infty$.

The following proposition generalizes Proposition 6.

Proposition 9. *For fixed $m \geq 1$, fixed $n \geq 2$, fixed $d_1, \dots, d_{n-1} \geq 1$, and for d_n being an indeterminate, we have*

$$C(m; \mathbf{d}) = \frac{1}{(X(m; \mathbf{d}) - 1)!} \frac{(2d_n + 1)!!^3}{2^{d_n+1} d_n!} P_{m; d_1, \dots, d_{n-1}}(X(m; \mathbf{d})), \quad (194)$$

where $X(m; \mathbf{d}) = X(\mathbf{d}) + 3m$ as before and $P_{m; d_1, \dots, d_{n-1}}(X) \in \mathbb{Q}[X]$. Moreover, $\frac{C(m; \mathbf{d})}{C(m+|\mathbf{d}|)}$ is a rational function of $X(m; \mathbf{d})$.

Proof. Substituting (82) in (192), we get (194). The second statement that $\frac{C(m; \mathbf{d})}{C(m+|\mathbf{d}|)}$ is a rational function of $X(m; \mathbf{d})$ can then be deduced from (10). \square

The following proposition generalizes Theorem 2.

Proposition 10. *For every $m \geq 0$, for fixed $n \geq 1$, fixed $\mathbf{d}' = (d_1, \dots, d_{n-1}) \in (\mathbb{Z}_{\geq 1})^n$, and $d_n \geq 0$, we have*

$$C(m; \mathbf{d}) \sim \frac{1}{\pi} \sum_{k=0}^{\infty} \frac{C_k(m; \mathbf{d}')}{X(m; \mathbf{d})^k} \quad (X(m; \mathbf{d}) \rightarrow \infty), \quad (195)$$

where $\mathbf{d} = (\mathbf{d}', d_n)$, $C_k(m; \mathbf{d}')$ are functions of m, d_1, \dots, d_{n-1} with $C_0 \equiv 1$. Moreover, there exist a sequence of polynomials

$$c_k(m; p_1, p_2, \dots) \in \mathbb{Q}[p_1, p_2, \dots], \quad k \geq 0, \quad (196)$$

with $c_0(m; p_1, p_2, \dots) \equiv 1$, such that

$$C_k(m; \mathbf{d}') = c_k(m; p_1(\mathbf{d}'), p_2(\mathbf{d}'), \dots), \quad k \geq 0. \quad (197)$$

Furthermore, under the degree assignments

$$\deg p_i = 2i + 1 \quad (i \geq 1), \quad (198)$$

the polynomials $c_k(m; p_1, p_2, \dots)$, $k \geq 1$, satisfy the degree estimates

$$\deg c_k(m; p_1, p_2, \dots) \leq k - 1; \quad (199)$$

in particular, $c_k(m; p_1, p_2, \dots)$ does not depend on any p_d with $d \geq (k - 1)/2$.

Proof. Expanding both sides of (192) with respect to $1/X(m; \mathbf{d})$, we obtain

$$\begin{aligned} C_k(m; d_1, \dots, d_{n-1}) &= 3^m \sum_{l=1}^m \frac{(-1)^{m-l}}{l!} \sum_{\substack{m_1, \dots, m_l \geq 1 \\ m_1 + \dots + m_l = m}} \binom{m}{m_1, \dots, m_l} \frac{1}{\prod_{i=1}^l (2m_i + 1)!!} \\ &\quad \times \sum_{j=0}^k C_j(d_1, \dots, d_{n-1}, m_1, \dots, m_l) S(k - j, m - l - 1), \end{aligned}$$

where $S(n, k)$ are the Stirling numbers of the second kind. The proposition is then proved by using Theorem 2. \square

REFERENCES

- [1] A. Aggarwal, Large genus asymptotics for intersection numbers and principal strata volumes of quadratic differentials. *Invent. Math.*, **226** (2021), 897–1010.
- [2] A. Alexandrov, Cut-and-join description of generalized Brezin-Gross-Witten model. *Adv. Theor. Math. Phys.*, **22** (2018), 1347–1399.
- [3] J. Ambjørn, L. Chekhov, The NBI matrix model of IIB superstrings. *J. High Energy Phys.*, **12** (1998), 7–13.
- [4] M. Bertola, B. Dubrovin, D. Yang, Correlation functions of the KdV hierarchy and applications to intersection numbers over $\overline{\mathcal{M}}_{g,n}$. *Phys. D*, **327** (2016), 30–57.
- [5] M. Bertola, G. Ruzza, Brezin-Gross-Witten tau function and isomonodromic deformations. *Commun. Number Theory Phys.*, **13** (2019), 827–883.
- [6] E. Brézin, D.J. Gross, The external field problem in the large N limit of QCD. *Phys. Lett. B*, **97** (1980), 120–124.
- [7] N. K. Chidambaram, E. Garcia-Failde, A. Giacchetto, Relations on $\overline{\mathcal{M}}_{g,n}$ and the negative r -spin Witten conjecture. [arXiv:2205.15621](https://arxiv.org/abs/2205.15621).
- [8] D. Chen, M. Möller, D. Zagier, Quasimodularity and large genus limits of Siegel-Veech constants. *J. Amer. Math. Soc.*, **31** (2018), 1059–1163.
- [9] P. A. Clarkson, N. Joshi, A. Pickering, Bäcklund transformations for the second Painlevé hierarchy: a modified truncation approach. *Inverse Problems*, **15** (1999), 175–187.
- [10] V. Delecroix, É. Goujard, P. Zograf, A. Zorich, Uniform lower bound for intersection numbers of psi-classes. *Symmetry Integrability Geom. Methods Appl.*, **16** (2020), 86–98.
- [11] V. Delecroix, É. Goujard, P. Zograf, A. Zorich, Large genus asymptotic geometry of random square-tiled surfaces and of random multicurves. *Invent. Math.*, **230** (2022), 123–224.
- [12] L. A. Dickey, *Soliton Equations and Hamiltonian Systems*, 2nd. World Scientific Publishing, 2003.
- [13] R. Dijkgraaf, E. Verlinde, H. Verlinde, Loop equations and Virasoro constraints in non-perturbative 2-D quantum gravity. *Nucl. Phys. B*, **348** (1991), 435–456.

- [14] N. Do, P. Norbury, Topological recursion on the Bessel curve. *Commun. Number Theory Phys.* **12** (2018), 53–73.
- [15] B. Dubrovin, D. Yang, Generating series for GUE correlators. *Lett. Math. Phys.*, **107** (2017), 1971–2012.
- [16] B. Dubrovin, D. Yang, On Gromov-Witten invariants of \mathbb{P}^1 . *Math. Res. Lett.*, **26** (2019), 729–748.
- [17] B. Dubrovin, D. Yang, D. Zagier, On tau-functions for the KdV hierarchy. *Selecta Math.*, **27** (2021), paper No. 12, 47 pp.
- [18] B. Eynard, E. Garcia-Falde, A. Giacchetto, P. Gregori, D. Lewański, Resurgent large genus asymptotics of intersection numbers. [arXiv:2309.03143](https://arxiv.org/abs/2309.03143).
- [19] B. Eynard, N. Orantin, Invariants of algebraic curves and topological expansion, *Commun. Number Theory Phys.* **1** (2007), 347–452.
- [20] A. S. Fokas, M. J. Ablowitz, On a unified approach to transformations and elementary solutions of Painlevé equations. *J. Math. Phys.* **23** (1982), 2033–2042.
- [21] A. S. Fokas, A. R. Its, A. A. Kapaev, V. Yu. Novokshenov, *Painlevé Transcendents: The Riemann-Hilbert Approach*. *Mathematical Surveys and Monographs* **128** (2006).
- [22] D. J. Gross, M. J. Newman, Unitary and hermitian matrices in an external field. II. The Kontsevich model and continuum Virasoro constraints. *Nuclear Phys. B*, **380** (1992), 168–180.
- [23] D. J. Gross, E. Witten, Possible Third Order Phase Transition in the Large N Lattice Gauge Theory. *Phys. Rev. D*, **21** (1980), 446–453.
- [24] J. Guo, A remark on equivalence between two formulas of the two-point Witten-Kontsevich correlators. *Chin. Q. J. Math.*, **39** (2024), 82–85.
- [25] J. Guo, D. Yang, On the large genus asymptotics of psi-class intersection numbers. *Math. Ann.*, **388** (2024), 61–97.
- [26] E. L. Ince, *Ordinary Differential Equations*, reprint of the 1926 edition (Lonmans, Green & Co., London), Dover Publications, NY, 1956.
- [27] C. Itzykson, J. B. Zuber, Combinatorics of the modular group. II. The Kontsevich integrals. *Internat. J. Modern Phys. A*, **7** (1992), 5661–5705.
- [28] R. Kaufmann, Y. Manin, D. Zagier, Higher Weil-Petersson volumes of moduli spaces of stable n -pointed curves. *Comm. Math. Phys.*, **181** (1996), 763–787.
- [29] M. Kazarian, P. Norbury, Polynomial relations among kappa classes on the moduli space of curves. *IMRN*, **2024** (2024), 1825–1867.
- [30] M. Kontsevich, Intersection theory on the moduli space of curves and the matrix Airy function. *Comm. Math. Phys.*, **147** (1992), 1–23.
- [31] K. Liu, H. Xu, Intersection numbers and automorphisms of stable curves. *Michigan Math. J.*, **58** (2009), 385–400.
- [32] K. Liu, H. Xu, Recursion formulae of higher Weil-Petersson volumes. *IMRN*, **2009** (2009), 835–859.
- [33] K. Liu, H. Xu, A remark on Mirzakhani’s asymptotic formulae. *Asian J. Math.*, **18** (2014), 29–52.
- [34] F. Magri, A simple model of the integrable Hamiltonian equation. *J. Math. Phys.*, **19** (1978), 1156–1162.
- [35] Y. Manin, P. Zograf, Invertible cohomological field theories and Weil-Petersson volumes. *Ann. Inst. Fourier (Grenoble)*, **50** (2000), 519–535.
- [36] A. Mironov, A. Morozov, G. W. Semenoff, Unitary matrix integrals in the framework of generalized Kontsevich model. 1. Brezin-Gross-Witten model. *Int. J. Mod. Phys. A*, **11** (1996), 5031–5080.
- [37] P. Norbury, A new cohomology class on the moduli space of curves. *Geom. Topol.*, **27** (2023), 2695–2761.
- [38] P. Norbury, Enumerative geometry via the moduli space of super Riemann surface. [arXiv:2005.04378](https://arxiv.org/abs/2005.04378).
- [39] D. Stanford, E. Witten, JT gravity and the ensembles of random matrix theory. *Adv. Theor. Math. Phys.*, **24** (2020), 1475–1680.
- [40] E. Witten, Two-dimensional gravity and intersection theory on moduli space. *Surveys Diff. Geom.*, **1** (1991), 243–310.
- [41] J. Xu, D. Yang, Galilean symmetry of the KdV hierarchy. [arXiv:2403.04631](https://arxiv.org/abs/2403.04631).
- [42] D. Yang, Q. Zhang, On the Hodge-BGW correspondence. *Commun. Number Theory Phys.*, **18** (2024), 611–651.
- [43] P. G. Zograf, An explicit formula for Witten’s 2-correlators. *J. Math. Sci.*, **240** (2019), 535–538.

JINDONG GUO, SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, 230026 HEFEI, P.R. CHINA

E-mail address: guojindong@mail.ustc.edu.cn

PAUL NORBURY, SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, VIC 3010 MELBOURNE, AUSTRALIA

E-mail address: norbury@unimelb.edu.au

DI YANG, SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, 230026 HEFEI, P.R. CHINA

E-mail address: diyang@ustc.edu.cn

DON ZAGIER, MAX PLANCK INSTITUTE FOR MATHEMATICS, 53111 BONN, GERMANY, AND INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS, TRIESTE, ITALY

E-mail address: dbz@mpim-bonn.mpg.de