


Max-Planck-Institut für Mathematik Bonn

Nonstationary smooth geometric structures for
contracting measurable cocycles

by

Karin Melnick



Nonstationary smooth geometric structures for contracting measurable cocycles

Karin Melnick

Max-Planck-Institut für Mathematik
Vivatsgasse 7
53111 Bonn
Germany

Department of Mathematics
University of Maryland
4176 Campus Drive
College Park, MD 20742
USA

NONSTATIONARY SMOOTH GEOMETRIC STRUCTURES FOR CONTRACTING MEASURABLE COCYCLES

KARIN MELNICK

ABSTRACT. We implement a differential-geometric approach to normal forms for contracting measurable cocycles to $\text{Diff}^q(\mathbf{R}^n, \mathbf{0})$, $q \geq 2$. We obtain resonance polynomial normal forms for the contracting cocycle and its centralizer, via C^q changes of coordinates. These are interpreted as nonstationary invariant differential-geometric structures. We also consider the case of contracted foliations in a manifold, and obtain C^q homogeneous structures on leaves for an action of the group of subresonance polynomial diffeomorphisms together with translations.

1. INTRODUCTION

Normal forms for differentiable contractions have a long history. Sternberg proved in [17] that a C^q -smooth diffeomorphism f of \mathbf{R}^n fixing $\mathbf{0}$ and satisfying $\|D_{\mathbf{0}}f\| < 1$ is C^q conjugate on a neighborhood of $\mathbf{0}$ to a polynomial of *resonance* type, obtained from the Taylor series of f at $\mathbf{0}$. The degree of this polynomial is bounded in terms of the spectrum of $D_{\mathbf{0}}f$.

This article concerns a generalization of this normal forms theorem to a flow acting on a product $M \times \mathbf{R}^n$, where M is a probability space, via a measurable cocycle to $\text{Diff}^q(\mathbf{R}^n, \mathbf{0})$.

1.1. Previous results on nonstationary normal forms. Guysinsky and Katok found nonstationary normal forms in the C^0 setting in [8] and [7]. They consider a transformation F of a continuous \mathbf{R}^n -bundle $\mathcal{E} \rightarrow M$, where M is a compact metric space, acting along fibers by C^q diffeomorphisms preserving the zero section. The contracting assumption is that $\sup_{x \in M} \|D_{\mathbf{0}}F_x\| < 1$. Assuming that the Mather spectrum of F acting on $\Gamma(\mathcal{E})$ has the *narrow bands* property, they prove that a continuous family of C^q coordinate changes leaves F acting along fibers by *subresonance polynomials*. Their degree is bounded in terms of the Mather spectrum.

Date: July 18, 2019.

This article was in large part written during a visit to the Max Planck Institut für Mathematik in Bonn, whose support I gratefully acknowledge, along with partial support from NSF grant DMS 1255462. I wish to acknowledge the important role of David Fisher in this project, as the one who proposed using the results of Ruelle for the measurable setting, and who also introduced me to Feres' paper. I furthermore thank him for numerous helpful discussions and insights.

The outline of their proof resembles Sternberg's. They first solve formally and then make a fixed point argument on the space of coordinate changes to obtain the solution. An important element in their conclusions is that the centralizer of F also acts along fibers by subresonance polynomials in the chosen coordinates.

In [4], Feres gives an interpretation of Guysinsky and Katok's normal forms in the case \mathcal{E} represents an F -invariant lamination in M , as a continuous F -invariant family of geometric structures: they are reductions of the frame bundles of some sufficiently high order of the leaves to the group of subresonance polynomials. He further obtains *generalized connections* on these reductions, from which he constructs the polynomial charts on leaves using Gromov's Frobenius theorem for partial differential relations—all under the additional assumption that the Mather spectral partition is differentiable along leaves.

This differentiability assumption on the spectral partition is problematic, because it does not usually hold. In a recent paper, Kalinin and Sadovskaya show [12] that differentiability along leaves of the Mather spectral *filtration*, which does hold in general, suffices to obtain smoothness along leaves of the normal forms.

These C^0 nonstationary normal forms have been used to prove numerous outstanding theorems, primarily on local rigidity of discrete group actions on homogeneous spaces (to cite just a few: [5], [6], [13], [11]). The initial action satisfies the narrow band condition on the spectrum, and sufficiently small deformations retain the property. We are interested in other situations in which one cannot necessarily verify the narrow band condition. In this case, an alternative approach is to consider measurable contractions.

1.2. The approach in this paper. We construct polynomial normal forms for contracting measurable cocycles to $\text{Diff}^q(\mathbf{R}^n, \mathbf{0})$, $q \geq 2$. Contracting in this context means negative Lyapunov spectrum. Our primary tool is the theorem of Ruelle in [16], rather than an explicit fixed point argument. We take a differential-geometric point of view and apply Ruelle's theorem to a prolongation of the initial cocycle to higher-order frame bundles $\mathcal{E}^{(r)}$ of \mathcal{E} . We then find a measurable invariant family of smooth reductions of $\mathcal{E}^{(r)}$ to the group of subresonance polynomials. From these reductions, at sufficiently high order, we construct, by methods similar to Feres', coordinate atlases on fibers in which our cocycle and its centralizer have values in *resonance polynomials*; these are defined in terms of the Lyapunov spectrum.

A statement of resonance polynomial normal forms for measurable contracting cocycles and their centralizers appears in [9, Thms 6.1, 6.2]. They briefly sketch the steps of a proof, following the scheme of [8]. A detailed proof for $q = \infty$ was given in 2005 by W. Li and K. Lu [15]. They in fact treat *hyperbolic* measurable cocycles—that is, they assume only that 0 is not in

the spectrum; they do not, however, extend their polynomial normal forms to the centralizer of the initial transformation.

Our preprint appeared online essentially simultaneously with one by Kalinin and Sadovskaya [10] in which they also obtain, by a different approach from ours, *subresonance* polynomial normal forms for contracting measurable cocycles and their centralizers for finite q . They can treat $q = r + \alpha$ for integral $r \geq 1$ and $0 < \alpha < 1$; they require $q + \alpha$ greater than the ratio of minimum and maximum Lyapunov exponents, while we require q greater than this ratio. They require a temperedness assumption on the C^q norms of their cocycle, while we, in order to apply Ruelle's theorem, make a stronger assumption of finite first moment. They apply the normal forms theorem to contracted foliations and obtain a homogeneous structure along leaves, as in our theorem 3.15 below.

The group of resonance polynomials is significantly smaller than the corresponding subresonance group. On the other hand, we lose the continuity between fibers present in the uniform case. In the important special case that the fibers are plaques of an invariant C^0 foliation, however, we obtain C^q smoothness along leaves, and homogeneity of the atlas. A recent paper [1] proves Hölder continuity of Lyapunov decompositions for diffeomorphisms of compact manifolds on sets of arbitrarily large measure, as in Lusin's theorem; as the regularity of our frame bundle reductions and resulting atlases seems to be controlled by regularity of the Lyapunov filtration, one could hope for a similar regularity result for them.

We provide an additional differential-geometric interpretation of the invariant structures on fibers, which generalizes the well-known flat affine connections on fibers in the case of 1/2-pinch spectrum: let r be the maximum degree of the resonance polynomials associated to the Lyapunov spectrum, and assume $r \leq q - 1$. Then in almost every fiber there is a C^{q-1} flag of submanifolds $\nu_x^1 \subset \dots \subset \nu_x^l$, $l \leq r$, and a C^{q-2} flat connection ∇_x^i on the normal bundle of ν_x^i in ν_x^{i+1} , for each $i = 1, \dots, l - 1$. In the case of a foliation, these flags of submanifolds fit together to a C^{q-1} flag of foliations, along with their connections, in almost-every leaf.

1.3. Statement of main results. Let $(\{\varphi^t\}, M, \mu)$ be an ergodic flow preserving a probability measure. Let $\mathcal{E} \cong M \times \mathbf{R}^n \rightarrow M$ be a measurable \mathbf{R}^n -bundle to which $\{\varphi^t\}$ lifts, preserving $M \times \{\mathbf{0}\}$, and acting C^q -smoothly on the fibers for $q \geq 2$. Denote by $F(t, x) = F_x^t$ the corresponding cocycle to $\text{Diff}^q(\mathbf{R}^n, \mathbf{0})$, and by $T(t, x) = T_x^t$ the cocycle $D_{\mathbf{0}}F_x^t$. Assume that

$$(MET) \quad \sup_{0 \leq t \leq 1} \ln^+ \|T^t\|, \sup_{0 \leq t \leq 1} \ln^+ \|T^{-t}\| \in L^1(M, \mu)$$

and let

$$-\infty < \lambda^{(1)} < \dots < \lambda^{(s)} < 0 \leq \lambda^{(s+1)} < \dots < \lambda^{(m)}$$

be the Lyapunov spectrum of T , given by the Multiplicative Ergodic Theorem. The cocycle $\{F_x^t\}$ will be called a *measurable contraction* if $m = s$ —that is, if the Lyapunov spectrum is all negative.

See definition 3.2 below for the group of *resonance polynomials* $\mathcal{H}^{(r),0}$ associated to the Lyapunov spectrum. The *centralizer* $Z(F_x^t)$ of the $\{\varphi^t\}$ action on \mathcal{E} is given in definition 2.5. For $F \in \text{Diff}^q(\mathbf{R}^n)$, write $\|F\|_q$ for the C^q -norm of F restricted to the unit disk $\overline{B(1)}$.

Theorem 3.13: *Assume $\{F_x^t\}$ is a measurable contraction, satisfying*

$$\sup_{0 \leq t \leq 1} \ln^+ \|F_x^{\pm t}\|_q \in L^1(M, \mu).$$

Let $r = \lfloor \lambda^{(1)}/\lambda^{(s)} \rfloor$, and assume $r \leq q - 1$. Then there exist the following differential-geometric structures on \mathcal{E}_x , for all x in a φ^t -invariant, μ -conull subset:

- (1) *A family \mathcal{A}_x of C^q charts, with transitions in $\mathcal{H}^{(r),0}$ (restricted to a neighborhood of $\mathbf{0}$). The collection $\cup_x \mathcal{A}_x$ is $Z(F_x^t)$ -invariant.*
- (2) *A C^{q-1} filtration $\mathcal{V}_x^1 \subset \dots \subset \mathcal{V}_x^l = T\mathcal{E}_x$, with $l \leq r$, equipped with C^{q-2} flat connections ∇_x^i on $\mathcal{V}_x^i/\mathcal{V}_x^{i-1}$, $i = 1, \dots, l$. The filtrations and connections are invariant by $Z(F_x^t)$.*

Elements of $Z(F_x^t)$ act by resonance polynomials in the coordinates given by (1). Part (1) above should be compared with [15, Thm 1.1(v)], while noting that they assume $q = \infty$, and that they do not extend their result to the centralizer. The filtration in (2) restricted to the zero section of \mathcal{E} is a conglomeration of the Lyapunov filtration of $\{T_x^t\}$.

In the important case that M is a compact manifold and $\{\varphi^t\}$ a C^0 flow, and where the fibers of \mathcal{E} are plaques of a C^0 foliation L by n -dimensional submanifolds admitting a φ^t -invariant C^q -structure (see section 2.2 below for details), we have the following smooth geometric structures on the leaves:

Theorem (compare Thm 3.15 below) *Assume $q \geq \lfloor \lambda^{(1)}/\lambda^{(s)} \rfloor + 1$ and that for fixed t , the q -jet $J_x^{(q)}(\varphi^t|_L)$ is continuous in x . For all x in a φ^t -invariant, μ -conull subset, for all $y \in L_x$, there is a family of C^q charts on L_x at y , varying measurably between leaves, with the additional properties:*

- (1) *They are global diffeomorphisms $\mathbf{R}^n \rightarrow L_x$;*
- (2) *φ^t and its centralizer act by resonance polynomials in these charts;*
- (3) *they make L_x into a homogeneous space for the group generated by translations and subresonance polynomials (see definition 3.2).*

If the spectrum Σ^0 is 1/2-pinned, then L carries an invariant family of C^{q-1} flat affine structures, measurable in x .

The Lyapunov-Ruelle foliations tangent to the Lyapunov filtration along L carry the following differential-geometric structure:

Theorem 3.16: *The foliation $\cup_x L_x$ contains a $Z(\varphi^t)$ -invariant filtered family of C^{q-1} subfoliations*

$$L^{i_1} \subset \dots \subset L^{i_l} = L \quad l \leq r,$$

in which, for all $1 \leq j \leq l$, the subfoliations $L^{i_{j-1}} \subset L^{i_j}$ carry a $Z(\varphi^t)$ -invariant, flat transverse connection, which is measurable in x and C^{q-2} inside almost every L_x ; in particular, almost every leaf $L_x^{i_1}$ carries an invariant flat connection.

2. SMOOTH DYNAMICAL FOLIATIONS FROM RUELLE

This section contains results from Ruelle [16], in their general form in subsection 2.1, and in the special case of foliations on compact C^0 manifolds in subsection 2.2. Subsection 2.3 contains some key results restricting the possible values of F_x^t -invariant tensors. The formulations of Ruelle's results are not exactly the same as in his paper. In particular, we work with a flow, and we keep track of invariance by the full centralizer of this flow. Proposition 2.13 in section 2.3 requires a simple modification of the innards of Ruelle's proof. The present section thus contains references to and some recapitulation of Ruelle, as well as some further proofs.

In this section, $(\{\varphi^t\}, M, \mu)$ and the cocycle $\{F_x^t\}$ are as in the introduction, but the action is not assumed to be a measurable contraction; that is, the Lyapunov spectrum of T has the general form

$$-\infty < \lambda^{(1)} < \dots < \lambda^{(s)} < 0 \leq \lambda^{(s+1)} < \dots < \lambda^{(m)}$$

Let $M_T \subset M$ be the μ -conull set where the conclusions of the MET hold for the cocycle T . For $x \in M_T$, denote the filtration corresponding to the negative portion of the Lyapunov decomposition of $T_0 \mathcal{E}_x \cong \mathbf{R}^n$ by

$$\mathbf{0} \subset V_x^1 \subset \dots \subset V_x^s \subseteq \mathbf{R}^n.$$

A key feature of Ruelle's theorem is that it applies to a noninvertible system. We are interested here in cocycles over invertible transformations or flows, and invertibility will be an essential assumption in section 2.3. A crucial role will also be played, however, by Ruelle's perturbation theorem [16, Thm 4.1], which applies to arbitrary sequences of linear transformations that are sufficiently close to $\{T_x^k\}$; see proposition 2.13 below.

2.1. Submanifolds in fibers. Following are results on the existence of the Lyapunov-Ruelle submanifolds in the fibers of \mathcal{E} and their invariance by the centralizer of $\{F_x^t\}$.

Theorem 2.1 (Ruelle 1979). *Assume $\sup_{0 \leq t \leq 1} \ln^+ \|F_x^{\pm t}\|_q \in L^1(M, \mu)$. Let $M_F \subseteq M$ be the set where the conclusion of the ergodic theorem holds for this function under the flow $\{\varphi^t\}$. Let $M_0 = M_F \cap M_T$.*

For all $x \in M_0$, there is a nested family of C^q injectively immersed submanifolds

$$\mathbf{0} \subset \nu_x^1 \subset \cdots \subset \nu_x^s \subset \mathcal{E}_x.$$

For $i = 1, \dots, s$, the submanifolds ν_x^i are tangent at $\mathbf{0}$ to V_x^i , and are characterized by

$$\nu_x^i = \{u \in \mathcal{E}_x : \limsup_{t \rightarrow \infty} \frac{1}{t} \ln \|F_x^t u\| \leq \lambda^{(i)}\}$$

The union $\cup_x \nu_x^i$ is F_x^t -invariant: $F_x^t(\nu_x^i) = \nu_{\varphi^t(x)}^i$.

Ruelle's original statement is for sequences rather than flows. The following proposition explicates the relation between his theorem and the above statement modulo this difference.

Proposition 2.2. *The conclusions of the theorem hold for the sequences $\{F_x^k = F(k, x)\}$ and $\{T_x^k = T(k, x)\}$ corresponding to iteration of $\varphi = \varphi^1$.*

Proof: Let $\lambda^{(i)} < \lambda < \lambda^{(i+1)}$; if $i = s$, replace $\lambda^{(i+1)}$ with 0. Ruelle's theorem 5.1 a) [16] asserts the existence of measurable functions $\beta(x) > \alpha(x) > 0$ such that the sets

$$\hat{\nu}_x^\lambda = \{u \in B(\alpha(x)) : \|F_x^k(u)\| \leq \beta(x)e^{k\lambda} \forall k \geq 0\}$$

are C^q submanifolds of \mathcal{E}_x , tangent at $\mathbf{0}$ to V_x^i , for all x in a μ -conull subset Γ . The set Γ is given by [16] (5.2)–(5.4) and, under our hypotheses, contains M_0 .

Remark 5.2 c) of [16] (see also (6.2) on p 52) concerns an important lower bound on the shrinkage of α along $\{\varphi^k\}$ -orbits: given $\lambda^{(s)} < \zeta < 0$, one can arrange that $\alpha(\varphi^k(x))$ decreases less rapidly than $e^{k\zeta}$.

Set

$$\nu_x^i = \bigcup_{k \geq 0} (F_x^k)^{-1}(\hat{\nu}_{\varphi^k(x)}^\lambda)$$

For $u \in \hat{\nu}_x^\lambda$ and $\lambda^{(i)} < \lambda' < \lambda$, part b') of [16, Thm 5.1] gives

$$\limsup_{k \rightarrow \infty} \frac{1}{k} \ln \|F_x^k u\| \leq \lambda'$$

It follows that

$$\nu_x^i \subseteq \{u \in \mathcal{E}_x : \limsup_{k \rightarrow \infty} \frac{1}{k} \ln \|F_x^k u\| \leq \lambda^{(i)}\}$$

Proof of the reverse containment follows an argument on the bottom of p 52 in [16]: if u belongs to the right-hand set above, then for $\lambda^{(i)} < \lambda < \min\{\lambda^{(i+1)}, \zeta\}$, and sufficiently large C , $\|F_x^k(u)\| \leq Ce^{k\lambda}$. For sufficiently large l , $\alpha(\varphi^l(x)) > Ce^{l\zeta}$. Then, for all $k \geq 0$,

$$\|F_x^{k+l}(u)\| \leq Ce^{l\zeta}e^{k\lambda} < \alpha(\varphi^l(x))e^{k\lambda} < \beta(\varphi^l(x))e^{k\lambda}$$

so $F_x^l(u) \in \hat{\nu}_{\varphi^l(x)}^\lambda$. Therefore $u \in \nu_x^i$.

Now F_x^k -invariance (both forwards and backwards) of the $\cup_x \nu_x^i$ follows easily from the lim sup characterization.

On p 53 of [16] is the further argument showing ν_x^i is an injectively immersed submanifold: namely, each $(F_x^m)^{-1}(\overset{\circ}{\nu}_{\varphi^m(x)}^\lambda)$ is contained in $(F_x^{l_0})^{-1}(\overset{\circ}{\nu}_{\varphi^{l_0}(x)}^\lambda)$ for some $l_0 > m$, in fact, for all $l \geq l_0$. Now ν_x^i is an increasing union of C^q submanifolds which are topological disks, tangent at $\mathbf{0}$ to V_x^i ; the remaining conclusions follow. \diamond

Remark 2.3. *In several key places below, reference will be made to the smaller submanifolds $\overset{\circ}{\nu}_x^\lambda$ in the foregoing proof, and the functions α and β in their definitions. In a few places, we will have to recall and tinker with some details of their construction.*

Remark 2.4. *By part b) of [16, Thm 5.1], the derivatives of F_x^k become arbitrarily small for sufficiently large k , uniformly on compact subsets of $\overset{\circ}{\nu}_x^s$. For $i \leq q - 1$, consider the cocycle $\tilde{F}^{(i)}(k, x) \in \text{Diff}(\mathbf{R}^n \times \mathbf{R}^n, \mathbf{0})$ mapping (u, v) to $(F_x^k(u), \mathcal{S}(D_u^{(i)} F_x^k)(v))$, where \mathcal{S} converts an i -symmetric tensor to the corresponding homogeneous polynomial of degree i . Then $\tilde{F}^{(i)}$ satisfies the finite first moment condition of theorem 2.1 with respect to the C^{q-i} -norm. (See section 3.3 for a similar but somewhat more demanding verification.) Then [16, Thm 5.1 b)] implies that $D_u(\mathcal{S}(D_u^{(i)} F_x^k))$ becomes arbitrarily small uniformly in $u \in \overset{\circ}{\nu}_x^s$, and it follows that $D^{(i+1)} F_x^k$ grows arbitrarily small for sufficiently large k , uniformly on compact subsets of $\overset{\circ}{\nu}_x^s$.*

The following proposition asserts that a bundle automorphism centralizing the initial cocycle preserves its Lyapunov-Ruelle submanifolds. We first define the centralizer.

Definition 2.5. *Let $\{F_x^k\}$ satisfy the discrete-time version of the standing assumptions of this section. The centralizer of $\{F_x^k\}$, denoted $Z(F_x^k)$, comprises all $(\{G_x^k\}, \psi)$, where*

- (1) ψ is a measurable automorphism of (M, μ) commuting with φ
- (2) $G(k, x) = G_x^k \in \text{Diff}^q(\mathbf{R}^n, \mathbf{0})$ is a measurable cocycle over ψ satisfying for all $k, l \in \mathbf{Z}$,

$$G(k, \varphi^l(x)) \circ F(l, x) = F(l, \psi^k(x)) \circ G(k, x)$$

- (3) $\ln^+ \|G_x^{\pm 1}\|_q \in L^1(M, \mu)$.

The centralizer of $\{F_x^t\}$, $t \in \mathbf{R}$, is defined similarly, and denoted $Z(F_x^t)$.

Proposition 2.6. *Let φ , M_0 , and $\{\nu_x^i\}$ be as in proposition 2.2, and let $(\{G_x^k\}, \psi) \in Z(F_x^k)$. Let $M_G \subseteq M$ comprise the points for which the conclusion of the ergodic theorem holds for both functions $\ln^+ \|G_x^{\pm 1}\|_q$ under the flow $\{\varphi^t\}$. Then G_x^k preserves $\cup_x \nu_x^i$ for all $i = 1, \dots, s$, for all $x \in \cap_{i \in \mathbf{Z}} \psi^i(M_0 \cap M_G)$.*

Proof: For $x \in M_0$, and $1 \leq i \leq s$, take $u \in \nu_x^i$, and let $u' = G_x^1(u)$. By the centralizing assumption,

$$(1) \quad \limsup_{k \rightarrow \infty} \frac{1}{k} \ln \|F_{\psi(x)}^k(u')\| = \limsup_{k \rightarrow \infty} \frac{1}{k} \ln \|G_{\varphi^k(x)}^1 F_x^k(u)\|$$

From proposition 2.2,

$$\limsup_{k \rightarrow \infty} \frac{1}{k} \ln \|F_x^k(u)\| \leq \lambda^{(i)};$$

in particular, for sufficiently large k , the image $F_x^k(u) \in \overline{B(1)}$. For such k , the norm $\|G_{\varphi^k(x)}^1 F_x^k(u)\|$ is bounded by the C^1 -norm $\|G_{\varphi^k(x)}^1\|_1$, times $\|F_x^k(u)\|$. Then we can also bound

$$\|G_{\varphi^k(x)}^1 F_x^k(u)\| \leq \|G_{\varphi^k(x)}^1\|_q \cdot \|F_x^k(u)\|$$

Now (1) is bounded above by

$$(2) \quad \limsup_{k \rightarrow \infty} \frac{1}{k} \ln \left(\|G_{\varphi^k(x)}^1\|_q \cdot \|F_x^k(u)\| \right)$$

For $x \in M_G$,

$$\limsup_{k \rightarrow \infty} \frac{1}{k} \ln \|G_{\varphi^k(x)}^1\|_q \leq \lim_{k \rightarrow \infty} \frac{1}{k} \ln^+ \|G_{\varphi^k(x)}^1\|_q = 0$$

Therefore (2) is bounded by $\lambda^{(i)}$, as is (1), for any $u' \in G_x^1(\nu_x^i)$. Thus $G_x^1(\nu_x^i) \subseteq \nu_{\psi(x)}^i$ for $x \in M_0 \cap M_G \cap \psi^{-1}(M_0)$.

A similar argument gives $G_x^{-1}(\nu_x^i) \subseteq \nu_{\psi^{-1}(x)}^i$ for $x \in M_0 \cap M_G \cap \psi(M_0)$.

Then for $x \in \cap_{i \in \mathbf{Z}} \psi^i(M_0 \cap M_G)$, we conclude $G_x^k(\nu_x^i) = \nu_{\psi(x)}^i$ for all $k \in \mathbf{Z}$.

◇

Now we can finish the

Proof: (of Theorem 2.1) Take the submanifolds ν_x^i associated to $\{F_x^k\}$, as in proposition 2.2. The F_x^t -invariance for all $t \in \mathbf{R}$ follows from proposition 2.6; note that M_0 is φ^t -invariant.

To complete the proof, it suffices to show that for $u \in \nu_x^i$,

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \ln \|F_x^t(u)\| \leq \lambda^{(i)}$$

Any t can be written $n_t + \epsilon_t$ with $n_t \in \mathbf{N}$ and $0 \leq \epsilon_t < 1$. For sufficiently large t , the image $F_x^{n_t}(u) \in \overline{B(1)}$. Then

$$\|F_x^t(u)\| = \|F_{\varphi^{n_t}(x)}^{\epsilon_t} F_x^{n_t}(u)\| \leq \|F_{\varphi^{n_t}(x)}^{\epsilon_t}\|_q \cdot \|F_x^{n_t}(u)\|$$

Now

$$\frac{1}{t} \ln \|F_x^t(u)\| \leq \frac{1}{n_t} \sup_{0 \leq \epsilon \leq 1} \ln \left(\|F_{\varphi^{n_t}(x)}^{\epsilon}\|_q \cdot \|F_x^{n_t}(u)\| \right)$$

As in the centralizer proposition, the ergodic theorem gives

$$\lim_{k \rightarrow \infty} \frac{1}{k} \sup_{0 \leq \epsilon \leq 1} \ln^+ \|F_{\varphi^k(x)}^\epsilon\|_q = 0$$

so

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \ln \|F_x^t(u)\| \leq \limsup_{t \rightarrow \infty} \frac{1}{n_t} \ln \|F_x^{n_t}(u)\| \leq \lambda^{(i)}$$

◇

Remark 2.7. *On regularity: If F_x^t acts by C^∞ diffeomorphisms of \mathbf{R}^n and satisfies the finiteness condition that*

$$\sup_{0 \leq t \leq 1} \ln^+ \|F_x^{\pm t}\|_q \in L^1(M, \mu) \quad \text{for all } q \geq 1,$$

then the submanifolds ν_x^r in theorem 2.1 are C^∞ . Section 5.3 of [16] verifies the smoothness of the ν_x^λ . (Note that the conull set M_0 can vary between the C^q and C^∞ cases). The arguments in the proof of proposition 2.2 are also valid in the C^∞ setting. The hypothesis of proposition 2.6 should then be replaced with $G_x^k \in \text{Diff}^\infty(\mathbf{R}^n, \mathbf{0})$, and $\ln^+ \|G_x^{\pm 1}\|_q \in L^1(M, \mu)$ for all $q \geq 1$. We note that there is also a C^ω version of these submanifolds (see [16, Sec. 5.4]).

Ruelle's regularity assumption is actually that the F_x^k are of class $C^{q,\theta}$ with $q \geq 1$ and $\theta \in (0, 1]$ —that is, the derivatives of order q are Hölder continuous of exponent θ . The submanifolds ν_x^i are then also $C^{q,\theta}$ -regular. The results of this section can be verified in this greater generality. See also [10]. We will, however, use $q \geq 2$ for theorems 3.13 and 3.15 below.

Remark 2.8. *One need not assume that μ is φ^t -ergodic. An arbitrary invariant probability measure can be decomposed into ergodic components. We make this assumption mostly to simplify statements and proofs.*

2.2. Foliations by filtered stable manifolds. We now let M be a compact C^0 manifold and $\{\varphi^t\}$ a C^0 flow. We will assume that the fibers of \mathcal{E} are plaques of a C^0 foliation L of M by n -dimensional submanifolds, admitting a φ^t -invariant C^q smooth structure. Thus M admits a C^0 foliated atlas in which the transitions are also C^q along the leaves of L . Fix a C^0 norm on $TL \rightarrow M$; we will assume $\{\varphi^t\}$ is nonuniformly contracting along L .

We will call a C^q atlas along L a family of C^q smooth charts $\theta_x : (\mathcal{E}_x, \mathbf{0}) \rightarrow (L_x, x)$, varying measurably in x , and with C^q norms of θ_x on $\overline{B(1)}$ and of $(\theta_x|_{\overline{B(1)}})^{-1}$ bounded in x . The cocycle $\{F_x^t\}$ is the local action of $\{\varphi^t\}$ on L_x in these charts:

$$F_x^t = (\theta_{\varphi^t(x)})^{-1} \circ \varphi^t \circ \theta_x.$$

Each F_x^t is defined on a neighborhood of the origin in $\mathcal{E}_x \cong \mathbf{R}^n$.

A C^q atlas along L will be called *uniformly biLipschitz* if there is $\kappa \geq 1$ such that for some continuous metric d on M , for almost all x ,

$$\frac{1}{\kappa} \|u - v\| \leq d(\theta_x(u), \theta_x(v)) \leq \kappa \|u - v\| \quad \forall u, v \in \overline{B(1)}$$

It is not hard to find a uniformly biLipschitz atlas along L : take a finite collection of C^0 foliated charts $\varphi_i : V_i \rightarrow U_i, i = 1, \dots, N$, for which V_i are open balls of radius 2 and $\cup_i \varphi_i(\frac{1}{2}\overline{V}_i)$ form a finite cover of M . Then make a piecewise continuous assignment $x \mapsto i$ with $x \in \varphi_i(\frac{1}{2}\overline{V}_i)$ and set $\theta_x = \varphi_i$ restricted to the leaf in V_i through $\varphi_i^{-1}(x)$ and recentered at x .

Here is the main result of this section, which is very close to [16, Thm 6.3].

Theorem 2.9 (Ruelle 1979). *Let $\{\varphi^t\}$ be a C^0 , measure-preserving flow on a compact manifold (M, μ) , preserving a C^0 foliation L and a C^q structure on the leaves. Assume $\sup_{-1 \leq t \leq 1} J_x^{(q)}(\varphi^t|_L)$ is bounded in x and that $D(\varphi^t|_L)$ has all negative Lyapunov exponents in a uniformly biLipschitz C^q atlas $\{(\mathcal{E}_x, \theta_x)\}$ along L . Then the corresponding cocycle satisfies the hypotheses of theorem 2.1; the resulting submanifolds $\nu_x^1 \subset \dots \subset \nu_x^s = \mathcal{E}_x$ satisfy, for $x \in M_0$:*

- (1) *Suppose $L_x = L_y$ and $y \in M_0 \cap \theta_x(\nu_x^i)$ for some $1 \leq i \leq s$. Then $\theta_y(\nu_y^i) \cap \theta_x(\nu_x^i)$ is relatively open in each term.*
- (2) *The resulting foliations $L^1 \subset \dots \subset L^s = L$ are φ^t -invariant, and can be defined at all points of L_x^s , for all $x \in M_0$.*
- (3) *The distributions $\mathcal{V}_x^i = T_x(L_x^i)$ vary C^{q-1} -smoothly within L_x^s for $1 \leq i < s$.*

Proof: The assumption of negative Lyapunov exponents along L implies $\{\varphi^t\}$ is nonuniformly contracting along L , so $\{F_x^t\}$ is defined on $\overline{B(1)}$ for sufficiently large t . The boundedness assumption on q -jets of φ^t along L implies $\{F_x^t\}$ satisfies the finite first moment assumption of theorem 2.1.

Now recall that ν_x^s is the increasing union $\cup_{k \geq 0} (F_x^k)^{-1} \overset{\circ}{\nu}_{\varphi^k(x)}^\lambda$ for $\lambda^{(s)} < \lambda < 0$. For any compact $D \subset \nu_x^s$, there is k_0 such that F_x^k is guaranteed to be defined on D for all $k \geq k_0$.

Let κ be the biLipschitz constant relating $\{\theta_x\}$ with a distance d on M . Let $u \in \nu_x^i$ with $\theta_y^{-1}\theta_x(u) = v$. Then

$$\begin{aligned} \|F_y^k(v)\| &= \|\theta_{\varphi^k(y)}^{-1} \circ \varphi^k(\theta_x(u))\| \\ &= \|\theta_{\varphi^k(y)}^{-1} \circ \theta_{\varphi^k(x)} \circ F_x^k(u)\| \\ &\leq \kappa \left(d(\varphi^k(y), \varphi^k(x)) + d(\theta_{\varphi^k(x)} \circ F_x^k(u), \varphi^k(x)) \right) \\ &\leq \kappa^2 \left(\|F_x^k(\theta_x^{-1}(y))\| + \|F_x^k(u)\| \right) \end{aligned}$$

so

$$\limsup_{k \rightarrow \infty} \frac{1}{k} \ln \|F_y^k(v)\| \leq \limsup_{k \rightarrow \infty} \frac{1}{k} \ln \kappa^2 \max\{\|F_x^k(\theta_x^{-1}(y))\|, \|F_x^k(u)\|\} \leq \lambda^{(i)}$$

From the characterization in theorem 2.1/proposition 2.2, we conclude $v \in \nu_y^i$. Thus

$$\theta_x(\nu_x^i) \cap \theta_y(\mathcal{E}_y) = \theta_x(\nu_x^i) \cap \theta_y(\nu_y^i)$$

which, similarly, equals $\theta_y(\nu_y^i) \cap \theta_x(\mathcal{E}_x)$. Now (1) is proved.

The φ^t -invariance of the foliations L^i , $i = 1, \dots, s$, follows immediately from their definition in terms of the atlas $(\mathcal{E}_x, \theta_x)$ and the F_x^t -invariant submanifolds ν_x^i . Given $x \in M_0$ and $y \in L_x^s$, the trajectories $\varphi^t(x)$ and $\varphi^t(y)$ come arbitrarily close as $t \rightarrow \infty$: one can choose a path between them in L_x^s and cover this path with finitely many open sets of the form $\theta_{x_i}(\nu_{x_i}^s)$. Each such open interval is uniformly exponentially contracted under φ^t as $t \rightarrow \infty$. Let t_0 be a time at which the images of x and y are contained in the image of $\theta_{\varphi^{t_0}(x)}$. All foliations are defined on this set, so $L_{\varphi^{t_0}(y)}^1 \subset \dots \subset L_{\varphi^{t_0}(y)}^s$ can be defined on a neighborhood of $\varphi^{t_0}(y)$. Then to prove (2), let $L_y^i = \varphi^{-t_0}(L_{\varphi^{t_0}(y)}^i)$ for each $1 \leq i < s$.

Given $x \in M_0$ and $\lambda^{(1)} < \lambda_1 < \lambda^{(2)} < \dots < \lambda = \lambda_s < 0$, let α be the minimum over $i = 1, \dots, s$ of the radii $\alpha(x)$ in the definition of $\nu_x^{\lambda_i}$. We will show that the distribution \mathcal{V}_y^i varies smoothly over $y \in \theta_x(B(\alpha)) \cap L_x^s$, for any $1 \leq i < s$. Let $u = \theta_x^{-1}(y)$. Note that

$$\theta_x^{-1}(L_y^i) \subseteq \{v \in \mathcal{E}_x : \limsup_{t \rightarrow \infty} \frac{1}{t} \ln \|F_x^t(v) - F_x^t(u)\| \leq \lambda^{(i)}\}$$

and $\theta_x^{-1}(L_y^i)$ contains a neighborhood of u in the right-hand set. Remark 5.2 (b) of [16] shows that the tangent space at u of the latter set varies C^{q-1} -smoothly in $u \in \nu_x^\lambda$. Mapping forward by θ_x gives (3). \diamond

Remark 2.10. Let $\{\psi^k\}$ be a continuous transformation commuting with $\{\varphi^t\}$, preserving the foliation L , acting C^q differentiably along leaves, with $J_x^{(q)}(\psi|_L)$ bounded in x . It follows from proposition 2.6 that the foliations $L^1 \subset \dots \subset L^s \subset L$ are ψ^k -invariant. We will denote the collection of such ψ by $Z(\varphi^t)$.

2.3. Stability of invariant tensors. The aim of this section is to establish restrictions on values of F_x^t -invariant tensors on the Lyapunov-Ruelle manifolds ν_x^i , for $1 \leq i \leq s$. Sequences of the form $\{D_u F_x^k\}_{k \in \mathbb{N}}$, for $u \in \nu_x^i$, have the same spectrum as $\{T_x^k\}_{k \in \mathbb{N}}$, by Ruelle's perturbation theorem. We will show that in fact the push-forward by $\{F_x^k\}$ of a tensor at $u \in \nu_x^i$ has the same asymptotic expansion rates as the push-forward at $\mathbf{0}$ by $\{T_x^k\}$.

2.3.1. *Digression on Ruelle's proof: constants in the construction.* Here we must expose some internal constants in Ruelle's construction of the $\hat{\nu}_x^\lambda$ (see remark 2.3), in order to introduce the proof of proposition 2.13.

Ruelle's perturbation theorem [16, Thm 4.1] says that given a linear sequence $\{T_k\}$ satisfying

- $\limsup_{k \rightarrow \infty} \ln \|T_k\| \leq 0$;
- $\lim_{k \rightarrow \infty} (T^{k*} T^k)^{1/2k} = \Lambda$ where $T^k = T_k \cdots T_1$;

and given $\eta > 0$, there exists $\delta > 0$, such that for any linear sequence $\{T'_k\}$ with

$$\|T - T'\| = \sup_k \|T_k - T'_k\| e^{3k\eta} < \delta,$$

$\{T'^k\}$ has a well-defined spectrum, equal to that of $\{T^k\}$. For $\lambda^{(i)}$ in this spectrum, and $P^{(i)}$ the projection to the corresponding asymptotic eigenspace for $\{T'^k\}$, equation (4.5) of [16] additionally says that given $\epsilon > 0$, there is $B'_\epsilon > 1$ such that $\|T'^k \circ P^{(i)}\| \leq B'_\epsilon \cdot e^{k(\lambda^{(i)} + \epsilon)}$.

This theorem is applied several times in the construction of $\hat{\nu}_x^\lambda$, with constants defined as follows. First, $\lambda = \lambda^{(i)} + \epsilon$, with $\lambda < \lambda^{(i+1)}$ (if $i = s$, take $\lambda^{(i+1)} = 0$). Then η is such that $0 < 4\eta \leq -\lambda$. Now δ is given by the perturbation theorem applied to the sequence $\{T_k = T(1, \varphi^{k-1}(x))\}$ (He further requires $\delta < 1/\sqrt{2}A$, where A is another constant given by [16, Thm 4.1]). Next, $0 < \beta(x) < \min\{1, \delta/G\}$, where

$$(3) \quad G(x) = \sup_k \|F_{\varphi^{k-1}(x)}^1\|_q \cdot e^{-k\eta - \lambda}$$

Note $G(\varphi^l(x)) \leq e^{l\eta} G(x)$. Last, $\alpha(x) = \beta(x)/B'_\epsilon$. End of digression.

Before proceeding, we state some basic facts about behavior of Lyapunov exponents under restriction to invariant subspaces and quotients.

Proposition 2.11. *Let $\{T_x^t\}$ be a linear cocycle over an ergodic, measure-preserving flow $(\{\varphi^t\}, M, \mu)$, satisfying (MET). Let Σ be the Lyapunov spectrum of $\{T_x^t\}$, with Lyapunov decomposition $W_x^1 \oplus \cdots \oplus W_x^m$. Suppose that $\{U_x\}_{x \in M}$ form a measurable T_x^t -invariant subbundle. Then*

- (1) *The spectrum of $\{T_x^t\}$ restricted to U is a subset of Σ , and $U_x = \oplus_i (U_x \cap W_x^i)$.*
- (2) *The spectrum of $\{\bar{T}_x^t\}$, the cocycle on the quotient by U , is a subset of Σ , and each W^i maps surjectively to the corresponding Lyapunov distribution in the quotient.*

Item (1) follows quickly from the definition of the Lyapunov decomposition for an invertible system. For (2), the Oseledec-Pesin reduction theorem [2, Thm 6.10] is helpful. It gives tempered equivalences (see [2, p 103]) between $\{T_x^1\}$ and block ϵ -approximately conformal matrices. The blocks

correspond to the Lyapunov decomposition. If $W_x^i \cap U_x = 0$, then the angle $\angle(W_{\varphi^t(x)}^i, U_{\varphi^t(x)})$ decreases subexponentially as $t \rightarrow \infty$. Thus for $v \in W_x^i$, the projection of $T_x^t v$ to $U_{\varphi^t(x)}^\perp$ has norm shrinking like $e^{t\lambda^{(i)}}$. For $\mathbf{0} \neq W_x^i \cap U_x \neq W_x^i$, the restriction of T_x^t to W_x^i is tempered equivalent to an ϵ -approximately scalar matrix, for any ϵ , which again implies that $\angle(W_{\varphi^t(x)}^i \cap U_{\varphi^t(x)}, W_{\varphi^t(x)}^i \cap T_x^t(U_x^\perp))$ decreases subexponentially.

Now let $\mathcal{T} \rightarrow M$ be a tensor bundle derived from $T_0\mathcal{E} \rightarrow M$ via T_x^t -invariant subbundles, quotients, and tensor operations. Write $\{T_{x*}^t\}$ for the linear cocycle induced from $\{T_x^t\}$ on \mathcal{T} . Note that as $\{T_x^t\}$ satisfies (MET), so does $\{T_{x*}^t\}$. Let Σ be the spectrum of $\{T_{x*}^t\}$ on \mathcal{T} , and denote by \mathcal{T}_x^σ the Lyapunov distribution corresponding to $\sigma \in \Sigma$.

Proposition 2.12. *If $\tau : M \rightarrow \mathcal{T}$ is a measurable, T_{x*}^t -invariant section, then $\tau(x) \in \mathcal{T}_x^0$ for almost-every $x \in M_0$.*

Proof: Suppose that $\tau(x)$ has a nontrivial component on \mathcal{T}_x^σ for some $\sigma > 0$. Then $\|(T_x^t)_*(\tau(x))\| = \|\tau(\varphi^t(x))\| \rightarrow \infty$ as $t \rightarrow \infty$. On the other hand, $\|\tau\|$ agrees with a continuous and bounded function on a set of measure $1 - \epsilon$ by Lusin's theorem. By ergodicity, for almost every x , the trajectory $\varphi^t(x)$ visits this set infinitely many times. We conclude that for almost every x , the component of $\tau(x)$ on \mathcal{T}_x^σ is zero.

If $\tau(x)$ has a nontrivial component on \mathcal{T}_x^σ with $\sigma < 0$, then the same argument with $t \rightarrow -\infty$ leads to the same contradiction. \diamond

Proposition 2.13. *Let $\mathcal{S} \rightarrow \cup_{x \in M_0} \nu_x^i$ be a tensor bundle constructed from $\oplus_x T\nu_x^i$ via F_x^k -invariant subbundles, quotients, and tensor operations, smooth in each ν_x^i and measurable in x . Denote $\{T_{\mathcal{S}}(k, x)\}$ the linear cocycle for $\{T_{x*}^k\}$ on \mathcal{S}_0 . Fix $u \in \nu_x^i$, and consider the following sequence mapping \mathcal{S}_u to $\mathcal{S}_{F_x^k(u)}$:*

$$T'_{\mathcal{S}}(k) = (F_x^k)_{*u} = (F_{\varphi^{k-1}(x)}^1)_{*F_x^{k-1}(u)} \cdots (F_x^1)_{*u}$$

Then $\{T'_{\mathcal{S}}(k)\}$ has a well-defined spectrum, equal to that of $\{T_{\mathcal{S}}(k, x)\}$.

Remark 2.14. *In the proof below, we will modify G in Ruelle's proof (see [16, (5.5)]). The functions $\beta(x)$ and $\alpha(x)$ will be modified accordingly; see subsection 2.3.1. The bound on shrinkage along orbits $e^{k\zeta} = O(\alpha(\varphi^k(x)))$, for $\lambda^{(s)} < \zeta < 0$, will remain intact, and the $\{\nu_x^i\}$ will be unchanged.*

Proof: It suffices to prove the statement when $u \in \mathring{\nu}_x^\lambda$, for λ as in subsection 2.3.1 above, because the spectrum of $T'_{\mathcal{S}}(k)$ is not changed by precomposition with an invertible linear map. By definition, $\|F_x^k(u)\| \leq \beta(x)e^{\lambda k}$ for all $k \geq 0$. As $q \geq 2$,

$$\|D_{F_x^{k-1}(u)} F_{\varphi^{k-1}(x)}^1 - T_{\varphi^{k-1}(x)}^1\| \leq \|F_{\varphi^{k-1}(x)}^1\|_q \beta(x) e^{(k-1)\lambda}$$

Similarly,

$$\|(D_{F_x^{k-1}(u)} F_{\varphi^{k-1}(x)}^1)^{-1} - (T_{\varphi^{k-1}(x)}^1)^{-1}\| \leq \|(F_{\varphi^{k-1}(x)}^1)^{-1}\|_q \beta(x) e^{(k-1)\lambda}$$

Because $x \in M_F$,

$$\lim_{k \rightarrow \infty} \frac{1}{k} \ln^+ \|(F_{\varphi^{k-1}(x)}^1)^{\pm 1}\|_q = 0$$

Let \mathcal{S} be a tensor bundle of type (a, b) . Let ϵ and η be as in subsection 2.3.1. Let δ be given by the perturbation theorem applied to the linear sequence $\{T_k = T(1, \varphi^{k-1}(x))_*\}$, acting on \mathcal{S}_0 . Instead of the function in (3), set

$$G(x) = \sup_k (\|F_{\varphi^{k-1}(x)}^1\|_q)^b \cdot (\|(F_{\varphi^{k-1}(x)}^1)^{-1}\|_q)^a \cdot e^{-k\eta - \lambda} < \infty$$

Then choose $\beta(x) < \min\{\delta/((a+b)G(x)), 1\}$. Now

$$\begin{aligned} & \| (F_{\varphi^{k-1}(x)}^1)_{*F_x^{k-1}(u)} - (T_{\varphi^{k-1}(x)}^1)_* \| \\ & \leq b \cdot \|D_{F_x^{k-1}(u)} F_{\varphi^{k-1}(x)}^1 - T_{\varphi^{k-1}(x)}^1\| \cdot \|F_{\varphi^{k-1}(x)}^1\|_q^{b-1} \cdot \|(F_{\varphi^{k-1}(x)}^1)^{-1}\|_q^a \\ & \quad + a \cdot \|(D_{F_x^{k-1}(u)} F_{\varphi^{k-1}(x)}^1)^{-1} - (T_{\varphi^{k-1}(x)}^1)^{-1}\| \cdot \|F_{\varphi^{k-1}(x)}^1\|_q^b \cdot \|(F_{\varphi^{k-1}(x)}^1)^{-1}\|_q^{a-1} \\ & \leq (a+b) \cdot \|(F_{\varphi^{k-1}(x)}^1)^{-1}\|_q^a \cdot \|F_{\varphi^{k-1}(x)}^1\|_q^b \cdot \beta(x) e^{(k-1)\lambda} \end{aligned}$$

Then (see [16, (5.5)]),

$$\|(F_{\varphi^{k-1}(x)}^1)_{*F_x^{k-1}(u)} - (T_{\varphi^{k-1}(x)}^1)_*\| e^{3k\eta} \leq (a+b) \cdot \beta(x) G < \delta$$

Now by [16, Thm 4.1], $\{T_{\mathcal{S}}(k, x)\}$ and $\{T'_{\mathcal{S}}(k)\}$ have the same spectrum. As in (3), $G(\varphi^l(x)) \leq e^{l\eta} G(x)$. Then α and β decrease along φ -orbits as in [16, Rmk 5.2 c)] (see also [16, Sec 4.7, p 43]). Referring to the proof of proposition 2.2, one can now see that our modifications of α and β do not ultimately alter the manifolds $\{\nu_x^i\}$. \diamond

Proposition 2.15. *Let $\oplus_x \tau_x$ be a family of smooth tensors belonging to a bundle $\mathcal{S} \rightarrow \cup_{x \in M_0} \nu_x^i$, varying measurably in x . Assume that \mathcal{S} is F_x^k -invariant, as in proposition 2.13, and that τ is F_x^t -invariant—that is,*

$$(F_x^t)_*(\tau_x(u)) = \tau_{\varphi^t(x)}(F_x^t u).$$

Let Σ be the spectrum of $T_{\mathcal{S}}$ on \mathcal{S}_0 , and denote $\Sigma^{\leq 0}$ the subset of nonpositive Lyapunov exponents. Then, for all x in a φ^t -invariant, μ -conull subset,

$$\lim_{t \rightarrow \infty} \frac{1}{t} \ln \|(F_x^t)_*(\tau_x(u))\| \in \Sigma^{\leq 0} \quad \forall u \in \nu_x^i.$$

Proof: By proposition 2.13 above, $\{T'_{\mathcal{S}}(k)\}$ and $\{T_{\mathcal{S}}(k, x)\}$ have the same spectrum, so

$$\lim_{k \rightarrow \infty} \frac{1}{k} \ln \|(F_x^k)_{*u}(\tau_x(u))\| \in \Sigma$$

To extend the limit to $t \rightarrow \infty$ in \mathbf{R} , the argument is the same as in the proof of theorem 2.1.

For $\lambda^{(i)} < \lambda < \lambda^{(i+1)}$, consider the measurable function

$$S(x) = \sup_{u \in \nu_x^\lambda} \|\tau_x(u)\|$$

By Lusin's theorem, $S(x)$ agrees with a continuous and bounded function on a set of measure $1 - \epsilon$, visited by $\varphi^t(x)$ for infinitely many t . On the other hand, if

$$\lim_{t \rightarrow \infty} \frac{1}{t} \ln \|(F_x^t)_*(\tau_x(u))\| > 0,$$

then $\|\tau_{\varphi^t(x)}(F_x^t u)\| \rightarrow \infty$ as $t \rightarrow \infty$. This would be a contradiction. \diamond

3. GEOMETRIC STRUCTURES ON FIBERS FOR CONTRACTIONS

Here $(\{\varphi^t\}, M, \mu)$ and $\mathcal{E} \rightarrow M$ are as in the previous section. Also as above, $F(t, x) = F_x^t \in \text{Diff}^q(\mathbf{R}^n, \mathbf{0})$ and $T_x^t = D_{\mathbf{0}}F(t, x)$ are the cocycles associated to the $\{\varphi^t\}$ -action on \mathcal{E} . In this section the Lyapunov exponents Σ^0 of $\{T_x^t\}$ are assumed to *all be negative*:

$$-\infty < \lambda^{(1)} < \dots < \lambda^{(s)} < 0,$$

so $\{F_x^t\}$ is asymptotically infinitesimally contracting on the fibers of \mathcal{E} as $t \rightarrow \infty$. We will use Ruelle's dynamical foliations to construct a family of F_x^t -invariant geometric structures on the fibers of \mathcal{E} .

3.1. Prolongation of F . Denote the general linear group of \mathbf{R}^n by $\text{GL}(n)$. The group $\text{GL}^{(r)}(n)$ comprises the r -jets at $\mathbf{0}$ of local diffeomorphisms of \mathbf{R}^n fixing $\mathbf{0}$. It can be identified with truncated Taylor series, or polynomials of degree r , with zero constant term and invertible linear component. The Lie algebra $\mathfrak{gl}^{(r)}(n)$ comprises the r -jets at $\mathbf{0}$ of local vector fields vanishing at $\mathbf{0}$, and can be identified with the polynomials of degree r with zero constant term. The bracket of $X, Y \in \mathfrak{gl}^{(r)}(n)$ can be computed as $[X, Y] = J_{\mathbf{0}}^{(r)}[\tilde{X}, \tilde{Y}]$, where \tilde{X} and \tilde{Y} are corresponding representative local vector fields. We will denote ρ_s^r the projection $\text{GL}^{(r)}(n) \rightarrow \text{GL}^{(s)}(n)$, for $r > s$, and also the projection $\mathfrak{gl}^{(r)}(n) \rightarrow \mathfrak{gl}^{(s)}(n)$. We set $S^{(r)}(n) = \ker \rho_{r-1}^r$; it is isomorphic to the abelian group $\text{Sym}^r(\mathbf{R}^{n*}) \otimes \mathbf{R}^n$.

We will make use below of a norm on $\mathfrak{gl}^{(r)}(n)$. View an element X as an ordered n -tuple of polynomials of degree at most r , and define $\|X\|$ to be the maximum norm of the monomial coefficients in X . Note that when X is viewed as a smooth map of \mathbf{R}^n , then this norm is bounded above by the C^r norm $\|X\|_r$. For $X \in \mathfrak{gl}(n)$, it is also bounded above by the usual linear operator norm (and below by $\|X\|/n$). We will need a bound on the growth of $\|\cdot\|$ under composition (followed by truncation above degree r).

Proposition 3.1. *Let $X, Y \in \mathfrak{gl}^{(r)}(n)$. Then*

$$\|Y \circ X\| \leq \|Y\| \cdot \max\{\|X\|, \|X\|^r\} \cdot c(r, n)$$

where $c(r, n)$ is a combinatorial constant depending on r and n .

Observe that for a monomial $Y(u_1, \dots, u_n)$ of degree j and a polynomial $X = (X_1, \dots, X_n)$, the maximum coefficient of $Y \circ X$ is bounded by $c'(j, n) \cdot \|Y\| \cdot \|X\|^j$, where $c'(j, n)$ is a combinatorial constant. The claimed bound for arbitrary $Y, X \in \mathfrak{gl}^{(r)}(n)$ follows by a similar estimate.

In a manifold M^n , the order- r frame bundle $\mathcal{F}^{(r)}(M)$ is a principal $\mathrm{GL}^{(r)}(n)$ -bundle, comprising r -jets at $\mathbf{0}$ of coordinate parametrizations of M . The order- r frame bundle of \mathbf{R}^n can be trivialized $\mathcal{F}^{(r)}(\mathbf{R}^n) \cong \mathbf{R}^n \times \mathrm{GL}^{(r)}(n)$ by identifying (u, g) with the r -jet at $\mathbf{0}$ of g composed with translation by u .

Define the r th prolongation of \mathcal{E} by $\mathcal{E}_x^{(r)} = \mathcal{F}^{(r)}(\mathcal{E}_x)$. Denote by π_s^r the projection $\mathcal{E}^{(r)} \rightarrow \mathcal{E}^{(s)}$, for $r > s$; this map of principal bundles is ρ_s^r -equivariant. We have a measurable trivialization $\mathcal{E}^{(r)} \cong M \times (\mathbf{R}^n \times \mathrm{GL}^{(r)}(n))$.

The fibers of π_0^r , each equivariantly diffeomorphic to $\mathrm{GL}^{(r)}(n)$, come equipped with a $\mathfrak{gl}^{(r)}(n)$ -valued 1-form, which we will denote ω . For $g \in \mathrm{GL}^{(r)}(n)$, this form satisfies $R_g^* \omega = \mathrm{Ad} g^{-1} \circ \omega$, where R_g is the right translation, and Ad is the adjoint representation.

The r th prolongation of F reflects the $\{F_x^t\}$ -action on $\mathcal{E}^{(r)}$ but is defined so that it leaves invariant the section $(\mathbf{0}, \mathrm{Id})$:

$$\begin{aligned} F^{(r)} &: \mathbf{R} \times M \rightarrow \mathrm{Diff}^{q-r}(\mathbf{R}^n \times \mathrm{GL}^{(r)}(n)) \\ F^{(r)}(t, x) &: (u, g) \mapsto (F_x^t(u), (J_u^{(r)} F_x^t) \cdot g \cdot (J_{\mathbf{0}}^{(r)} F_x^t)^{-1}) \end{aligned}$$

Here $J_u^{(r)} f$ denotes the r -jet of f at u , identified with an element of $\mathrm{GL}^{(r)}(n)$ via pre- and post-composition with appropriate translations; thus the product in the second coordinate takes place in $\mathrm{GL}^{(r)}(n)$. It is easy to verify that $F^{(r)}$ is a cocycle.

There is a corresponding r th prolongation of T :

$$T^{(r)}(t, x) = D_{(\mathbf{0}, \mathrm{Id})}(F^{(r)}(t, x)) = D_{\mathbf{0}} F_x^t \oplus \mathrm{Ad}(J_{\mathbf{0}}^{(r)} F_x^t)$$

In terms of ω , the derivative of the $F^{(r)}(t, x)$ -action on fibers of π_0^r is by $\mathrm{Ad} J_{\mathbf{0}}^{(r)} F_x^t$: If $v \in T_{(u, g)} \mathcal{E}_x^{(r)} \cap \ker(\pi_0^r)_*$ with $\omega(v) = X$, then we can write $v = (\mathbf{0}, gX)$. Under the derivative of $F^{(r)}(t, x)$, the first coordinate remains $\mathbf{0}$, and the second coordinate becomes

$$(J_u^{(r)} F_x^t) g X (J_{\mathbf{0}}^{(r)} F_x^t)^{-1} = (J_u^{(r)} F_x^t) g (J_{\mathbf{0}}^{(r)} F_x^t)^{-1} (J_{\mathbf{0}}^{(r)} F_x^t) X (J_{\mathbf{0}}^{(r)} F_x^t)^{-1}$$

which evaluates under ω to $(\mathrm{Ad} J_{\mathbf{0}}^{(r)} F_x^t)(X)$.

3.2. Subresonance Polynomials. Subresonance polynomials arise naturally in our context. We will refer to [4, Sec 3]; see also [8, Prop 1.1]. An important role will be played by a nilpotent proper subgroup of the subresonance polynomials, to be defined below as the *strict subresonance polynomials*.

Recall that $\lambda^{(s)} < 0$ denotes the greatest element of the Lyapunov spectrum Σ^0 of the cocycle $\{T_x^t\}$. Let $d_i, i = 1, \dots, s$, be the dimensions of the Lyapunov distributions corresponding to $\lambda^{(1)}, \dots, \lambda^{(s)}$, respectively; these dimensions are constant on M_T by ergodicity of $\{\varphi^t\}$. Let $W^1 \oplus \dots \oplus W^s$ be a decomposition of \mathbf{R}^n into subspaces of dimension d_1, \dots, d_s , respectively.

We first establish some notation. Given $l, r \geq 1$ and $\sigma \in \mathbf{R}$,

$$\begin{aligned} \widehat{W}_l^\sigma &:= \sum \{(W^{p_1^*} \otimes \dots \otimes W^{p_l^*}) \otimes W^i : \lambda^{(i)} - \sum_{j=1}^l \lambda^{(p_j)} = \sigma\} \\ \widehat{W}^{(r),\sigma} &:= \bigoplus_{l=1}^r \widehat{W}_l^\sigma; \quad \widehat{V}_l^\sigma := \bigoplus_{\lambda \leq \sigma} \widehat{W}_l^\lambda; \quad \widehat{V}^{(r),\sigma} = \bigoplus_{l=1}^r \widehat{V}_l^\sigma \end{aligned}$$

Let

$$\Lambda = e^{\lambda^{(1)}} \text{Id}_{W_1} \oplus \dots \oplus e^{\lambda^{(s)}} \text{Id}_{W_s} \in \text{GL}(n).$$

Definition 3.2. *The subresonance polynomials in $\text{GL}^{(r)}(n)$ are*

$$\mathcal{H}^{(r)}(\Lambda) = \widehat{V}^{(r),0} \cap \text{GL}^{(r)}(n)$$

The corresponding strict subresonance polynomials and resonance polynomials are, respectively,

$$\mathcal{X}^{(r)}(\Lambda) = \text{Id} + \widehat{V}^{(r),\lambda^{(s)}} \quad \text{and} \quad \mathcal{H}^{(r),0}(\Lambda) = \widehat{W}^{(r),0} \cap \text{GL}^{(r)}(n)$$

Often $\mathcal{H}^{(r)}(\Lambda)$ will simply be written $\mathcal{H}^{(r)}$ when Σ^0 and Λ are clear, and similarly for $\mathcal{X}^{(r)}$ and $\mathcal{H}^{(r),0}$.

Note that subresonance polynomials have bounded degree: for $r \geq \lfloor \lambda^{(1)}/\lambda^{(s)} \rfloor$, the groups $\mathcal{H}^{(r)} = \mathcal{H}^{(r+1)}$. Note also that all derivatives at any $u \in \mathbf{R}^n$ of a subresonance polynomial are strictly subresonance. Similarly, the translation by $u \in \mathbf{R}^n$ of a subresonance polynomial, $\tau_{-h(u)} \circ h \circ \tau_u$, is subresonance.

Proposition 3.3. *The polynomial germs in $\mathcal{H}^{(r)}$, $\mathcal{X}^{(r)}$, and $\mathcal{H}^{(r),0}$ form subgroups of $\text{GL}^{(r)}(n)$, with Lie algebras*

$$\mathfrak{h}^{(r)} = \widehat{V}^{(r),0} \quad \mathfrak{x}^{(r)} = \widehat{V}^{(r),\lambda^{(s)}} \quad \mathfrak{h}^{(r),0} = \widehat{W}^{(r),0}$$

The group $\mathcal{X}^{(r)}$ is a nilpotent normal subgroup of $\mathcal{H}^{(r)}$. When $r \geq \lfloor \lambda^{(1)}/\lambda^{(s)} \rfloor$, then polynomials in $\mathcal{H}^{(r)}$ are global diffeomorphisms of \mathbf{R}^n .

Proof: It suffices to prove the statements for $r \geq \lfloor \lambda^{(1)}/\lambda^{(s)} \rfloor$, as the homomorphism ρ_l^r , $l \leq r$, preserves all group-theoretic properties and maps $\widehat{W}^{(r),\sigma}$ to $\widehat{W}^{(l),\sigma}$ and $\widehat{V}^{(r),\sigma}$ to $\widehat{V}^{(l),\sigma}$.

The Λ^k -stable polynomials in $\mathfrak{gl}^{(r)}(n)$ are those X for which $\|\Lambda^k X \Lambda^{-k}\|$ is bounded as $k \rightarrow \infty$. The Λ^k -stable polynomials with invertible first derivative at $\mathbf{0}$ form a subgroup of $\text{Diff}(\mathbf{R}^n, \mathbf{0})$ by [4, Prop 8]; by [4, Lemma

7 (6)], this group equals $\mathcal{H}^{(r)}$. The Lie algebra $\mathfrak{h}^{(r)}$ comprises all Λ^k -stable polynomials in $\mathfrak{gl}^{(r)}(n)$, which are $\widehat{V}^{(r),0}$.

We can express $\mathcal{X}^{(r)} \subset \mathcal{H}^{(r)}$ as

$$\mathcal{X}^{(r)} = \{g \in \mathcal{H}^{(r)} : \lim_{k \rightarrow \infty} \frac{1}{k} \ln \|\Lambda^k g \Lambda^{-k} - \text{Id}\| \leq \lambda^{(s)}\}$$

By bilinearity of the bracket,

$$(1) \text{ For } X, Y \in \sum_{\sigma \leq \lambda^{(s)}} \widehat{W}^{(r),\sigma} = \widehat{V}^{(r),\lambda^{(s)}},$$

$$\lim_{k \rightarrow \infty} \frac{1}{k} \ln \|\Lambda^k [X, Y] \Lambda^{-k}\| \leq \lim_{k \rightarrow \infty} \frac{1}{k} \left(\ln \|\Lambda^k X \Lambda^{-k}\| + \ln \|\Lambda^k Y \Lambda^{-k}\| \right)$$

Using proposition 3.1, one can check the following inequalities:

$$(2) \text{ For } g, h \in \mathcal{H}^{(r)},$$

$$\lim_{k \rightarrow \infty} \frac{1}{k} \ln \|\Lambda^k g h \Lambda^{-k} - \text{Id}\| \leq \lim_{k \rightarrow \infty} \frac{1}{k} \sup \{ \ln \|\Lambda^k g \Lambda^{-k} - \text{Id}\|, \ln \|\Lambda^k h \Lambda^{-k} - \text{Id}\| \}$$

$$(3) \text{ For } X \in \widehat{V}^{(r),\lambda^{(s)}} \text{ and } g \in \mathcal{H}^{(r)},$$

$$\lim_{k \rightarrow \infty} \frac{1}{k} \ln \|\Lambda^k g X g^{-1} \Lambda^{-k}\| \leq \lim_{k \rightarrow \infty} \frac{1}{k} \ln \|\Lambda^k X \Lambda^{-k}\|$$

Item (2) with $g, h \in \mathcal{X}^{(r)}$ shows this subset is closed under multiplication. Similarly,

$$\|\Lambda^k h^{-1} \Lambda^{-k} - \text{Id}\| = \|(\text{Id} - \Lambda^k h \Lambda^{-k})(\Lambda^k h^{-1} \Lambda^{-k})\|$$

and $\|\Lambda^k h^{-1} \Lambda^{-k}\|$ is bounded, so $\mathcal{X}^{(r)}$ is closed under inversion. Now $\mathcal{X}^{(r)}$ is a subgroup, and it is clear that the Lie algebra $\mathfrak{x}^{(r)}$ is as claimed.

One deduces from item (1) that $\mathfrak{x}^{(r)}$ is nilpotent.

Finally, from (3), $\mathfrak{x}^{(r)}$ is a Lie algebra ideal, so $\mathcal{X}^{(r)} \triangleleft \mathcal{H}^{(r)}$.

We will prove inductively on r that $\mathcal{H}^{(r),0}$ forms a subgroup (dropping the assumption $r \geq \lfloor \lambda^{(s)}/\lambda^{(1)} \rfloor$). It is easy to see that (truncations of) compositions of polynomials in $\widehat{W}^{(r),0}$ are again in $\widehat{W}^{(r),0}$, so we will just check closure under inverse.

For $r = 1$, we have

$$\mathcal{H}^{(1),0} = \text{GL}(W^1) \oplus \dots \oplus \text{GL}(W^s)$$

with Lie algebra

$$\mathfrak{h}^{(1),0} = \text{End}(W^1) \oplus \dots \oplus \text{End}(W^s) = \widehat{W}^{(1),0}$$

Assume that $\mathcal{H}^{(j-1),0}$ is closed under inversion. Let $g, h \in \mathcal{H}^{(j)}$ with $\bar{g} = \rho_{j-1}^j(g), \bar{h} = \rho_{j-1}^j(h) \in \mathcal{H}^{(j-1),0}$. Write $g = \bar{g} + \gamma$ and $h = \bar{h} + \delta$ with $\gamma, \delta \in S^{(j)}(n)$. Then the composition in $\mathcal{H}^{(j)}$ can be written

$$(4) \quad g \circ h = \bar{g} \circ \bar{h} + (D_0 g) \circ \delta + \gamma \circ (D_0 h) + R$$

where $\bar{g} \circ \bar{h} \in \mathcal{H}^{(j-1),0}$, and R is a sum of degree- j compositions of terms of \bar{g} and \bar{h} . Assume $g \in \mathcal{H}^{(j),0}$, and let $h = g^{-1} = \bar{g}^{-1} + \delta$. Then

$$(5) \quad \delta = -D_0 g^{-1} \circ \gamma \circ D_0 g^{-1} - D_0 g^{-1} \circ R \in \widehat{W}_j^0$$

We conclude that $\mathcal{H}^{(r),0} \leq \mathcal{H}^{(r)}$ with Lie algebra $\mathfrak{h}^{(r),0} = \widehat{W}^{(r),0}$. \diamond

Proposition 3.4. (compare [4, Prop 7]) *Let $\{F_k\}$ be a sequence in $GL^{(r)}(n)$, $r \geq 2$, with $\rho_1^r(F_k) = T_k$. Let $F^k = F_k \circ \dots \circ F_1$ and $T^k = T_k \circ \dots \circ T_1$. Suppose that $(T^{k*} T^k)^{1/2k} \rightarrow \Lambda$, with spectrum Σ^0 and eigenspace decomposition $W^1 \oplus \dots \oplus W^s$. Then:*

- (1) *Denote by A^k the restriction of $Ad(F^k)$ to $S^{(r)}(n)$. The limit $\lim_{k \rightarrow \infty} (A^{k*} A^k)^{1/2k}$ exists, and has spectrum comprising all*

$$\exp(\lambda^{(i)} - \sum_{j=1}^r \lambda^{(p_j)}) \quad \text{where} \quad \lambda^{(i)}, \lambda^{(p_j)} \in \Sigma^0$$

- (2) *For $X \in S^{(r)}(n)$, the limit $\lim_{k \rightarrow \infty} \frac{1}{k} \ln \|F^k \circ X \circ F^{-k}\|$ exists, and equals σ if and only if $X \in \widehat{W}_r^\sigma$.*

Proof: For $X \in S^{(r)}(n)$ and $F \in GL^{(r)}(n)$ with $\rho_1^r(F) = T$, the conjugation is

$$F \circ X \circ F^{-1} = T \circ X \circ T^{-1} \quad (\text{see [4, Eqn. 27]})$$

Then

$$(A^{*k} A^k)^{1/2k}(X) = ((Ad T^k)^*(Ad T^k))^{1/2k}(X)$$

The symmetrization of $Ad T^k$ on $S^{(r)}(n)$ is simply $Ad(T^{k*} T^k)^{1/2}$. Then the above expression is

$$(T^{k*} T^k)^{1/2k} \circ X \circ (T^{k*} T^k)^{-1/2k} \rightarrow \Lambda \circ X \circ \Lambda^{-1}$$

The remaining claims of (1) and (2) now follow. \diamond

3.3. $T^{(r)}$ spectrum, Lyapunov decomposition, and algebraic hull.

Let $T^{(r)}(t, x) = D_{(\mathbf{0}, \text{Id})} F^{(r)}(t, x)$ as in section 3.1 above. We have assumed that $q \geq 2$ and

$$\sup_{0 \leq \epsilon \leq 1} \ln^+ \|F_x^{\pm \epsilon}\|_q \in L^1(M, \mu).$$

We will deduce that, for $r \leq q$, the linear cocycle $T^{(r)}$ satisfies (MET). The two components of $T^{(r)}(t, x)$ are $D_0 F_x^t$ and $Ad J_0^{(r)} F_x^t$. As $\|D_0 F_x^t\| \leq \|F_x^t\|_q$, we can focus on the second factor. Fix $x \in M$ and ϵ between -1 and 1 , and let $X \in \mathfrak{gl}^{(r)}(n)$ with $\|X\| = 1$. By proposition 3.1,

$$\begin{aligned} \ln^+ \|(Ad J_0^{(r)} F_x^\epsilon)(X)\| &\leq \ln^+ \left(\|J_0^{(r)} F_x^\epsilon\| \cdot \max_{i=1, r} \|X \circ (J_0^{(r)} F_x^\epsilon)^{-1}\| \cdot c(r, n) \right) \\ &\leq \ln^+ \|J_0^{(r)} F_x^\epsilon\| + \ln^+ \|(J_0^{(r)} F_x^\epsilon)^{-1}\|^{r^2} + \ln c(r, n)^{r+1} \end{aligned}$$

Let $C_0 = (r+1) \ln c(r, n)$. As the above expression is independent of the unit vector X , we obtain with respect to the linear norm on $\mathrm{GL}(\mathfrak{gl}^{(r)}(n))$,

$$\begin{aligned} & \int \ln^+ \|(\mathrm{Ad} J_{\mathbf{0}}^{(r)} F_x^\epsilon)\| dx \\ & \leq \int \ln^+ \|J_{\mathbf{0}}^{(r)} F_x^\epsilon\| dx + \int r^2 \ln^+ \|(J_{\mathbf{0}}^{(r)} F_x^\epsilon)^{-1}\| dx + C_0 \\ & \leq \int \ln^+ \|J_{\mathbf{0}}^{(r)} F_x^\epsilon\| dx + \int r^2 \ln^+ \|J_{\mathbf{0}}^{(r)} F_x^{-\epsilon}\| dx + C_0 \\ & \leq \int \ln^+ \|F_x^\epsilon\|_r dx + \int r^2 \ln^+ \|F_x^{-\epsilon}\|_r dx + C_0 \end{aligned}$$

Inserting \sup_ϵ in the integrands in the above chain of inequalities leads to

$$\sup_{0 \leq \epsilon \leq 1} \ln^+ \|\mathrm{Ad} J_{\mathbf{0}}^{(r)} F_x^{\pm\epsilon}\| \in L^1(M, \mu),$$

as desired. The Multiplicative Ergodic Theorem thus applies to $\{T^{(r)}(t, x)\}$.

Denote by $M_T \subseteq M$ the set on which the conclusions of the MET hold for $\{T^{(r)}(t, x)\}$. For $\sigma \in \mathrm{Spec} T^{(r)}$, denote by $V_x^{(r), \sigma}$ and $W_x^{(r), \sigma}$ the corresponding terms in the Lyapunov filtration and Lyapunov decomposition, respectively. We can assume, after a tempered linear cocycle equivalence, that the Lyapunov decomposition for $\{T_x^t\}$ is constant $W_x^i = W^i$, where W^i is as in section 3.2 (see [2, Thm 6.1]). The following proposition implies that now the Lyapunov decomposition for $\{T^{(r)}(t, x)\}$ is also constant.

Proposition 3.5. *Let $1 \leq r \leq q$.*

(1) *The Lyapunov exponents of $\{T^{(r)}(t, x)\}$ are $\cup_{l=0}^r \Sigma^l$, where,*

$$\Sigma^l = \left\{ \lambda^{(i)} - \sum_{j=1}^l \lambda^{(p_j)} : \lambda^{(i)}, \lambda^{(p_j)} \in \Sigma^0 \right\} \quad \text{for } l \geq 1$$

(2) *For $\sigma \in \cup_{l=0}^r \Sigma^l$, for x in a φ^t -invariant μ -conull set, the $\{T^{(r)}(t, x)\}$ Lyapunov space at x comprises $W^\sigma \subset \mathbf{R}^n$, direct sum*

$$\widehat{W}^{(r), \sigma} = W_x^{(r), \sigma} \cap \mathfrak{gl}^{(r)}(n)$$

Proof: For $r \geq 2$, proposition 3.4 says that $\Sigma^r \subset \mathrm{Spec} T^{(r)}$ and that, for each $\sigma \in \Sigma^r$,

$$W_x^{(r), \sigma} \cap S^{(r)}(n) = \widehat{W}_r^\sigma.$$

Thus for almost all $x \in M$, by proposition 2.11 (1) and proposition 3.4 (1),

$$S^{(r)}(n) = \bigoplus_{\sigma \in \Sigma^r} \widehat{W}_r^\sigma$$

The quotient vector bundle is

$$T_{(\mathbf{0}, \mathrm{Id})} \mathcal{E}^{(r-1)} = T_{(\mathbf{0}, \mathrm{Id})} \mathcal{E}^{(r)} / S^{(r)}(n)$$

The spectrum of $T^{(r-1)}$ is a subset of $\text{Spec } T^{(r)}$, and $W_x^{(r),\sigma}$ projects onto $W_x^{(r-1),\sigma}$, by proposition 2.11 (2). Projecting $\mathcal{E}^{(j)} \rightarrow \mathcal{E}^{(j-1)}$ for $j = r, \dots, 2$, yields $\text{Spec } T^{(r)} = \cup_{l=2}^r \Sigma^l \cup \text{Spec } T^{(1)}$. It is easy to see that $\text{Spec } T^{(1)} = \Sigma^1 \cup \Sigma^0$, as the action here only involves T_x^t and $\text{Ad } T_x^t$; moreover, the Lyapunov space

$$W_x^{(1),\sigma} = W^\sigma \oplus \widehat{W}_1^\sigma.$$

We conclude $\text{Spec } T^{(r)} = \cup_{l=0}^r \Sigma^l$, as claimed in (1).

With respect to the filtration induced by $\ker(\rho_j^r)_*$, $j = r-1, \dots, 0$, the space $W_x^{(r),\sigma} \cap \mathfrak{gl}^{(r)}(n)$ has the same associated graded space as $\widehat{W}^{(r),\sigma}$; in particular, $W_x^{(1),\sigma} \cap \mathfrak{gl}(n) = \widehat{W}_1^\sigma$. Suppose that $W_x^{(j-1),\sigma} \cap \mathfrak{gl}^{(j-1)}(n) = \widehat{W}^{(j-1),\sigma}$ for some $j \geq 2$. Form the tensor bundle

$$\begin{aligned} \mathcal{T}_x &= (W_x^{(j-1),\sigma} \cap \ker(\rho_0^{j-1})_*)^* \otimes \left(\ker(\rho_{j-1}^j)_* / (W_x^{(j),\sigma} \cap \ker(\rho_{j-1}^j)_*) \right) \\ &= (\widehat{W}^{(j-1),\sigma})^* \otimes (S^{(j)}(n) / \widehat{W}_j^\sigma) \end{aligned}$$

Let $\{T^{(j)}(t, x)_*\}$ be the linear cocycle on \mathcal{T} induced from $\{T^{(j)}(t, x)\}$, and define an invariant tensor $\{\tau_x \in \mathcal{T}_x\}$ by

$$W_x^{(j),\sigma} = \text{graph}(\tau_x) + \widehat{W}_j^\sigma$$

Proposition 2.12 says that τ_x belongs, for almost every x , to the $\{T^{(j)}(t, x)_*\}$ Lyapunov space \mathcal{T}_x^0 . But this space is trivial, so $\tau_x = 0$, and $W_x^{(j),\sigma} \cap \mathfrak{gl}^{(j)}(n) = \widehat{W}^{(j),\sigma}$ for almost every x . Induction on j yields (2). \diamond

Henceforth denote $W^{(r),\sigma} = W^\sigma \oplus \widehat{W}^{(r),\sigma}$ and similarly for $V^{(r),\sigma}$.

Proposition 3.6. *For all $x \in M_T$ and all $t \in \mathbf{R}$,*

$$J_{\mathbf{0}}^{(r)} F_x^t \in \mathcal{H}^{(r),0}.$$

If $(\{G_x^k\}, \psi) \in Z(F_x^t)$ then also for all k and almost all x ,

$$J_{\mathbf{0}}^{(r)} G_x^k \in \mathcal{H}^{(r),0}.$$

Remark 3.7. *In fact, the conclusion holds for any $\{G_x^k\}$ satisfying (1) and (3) of definition 2.5, for which the prolonged linear cocycle*

$$U^{(r)}(k, x) = D_{\mathbf{0}} G_x^k \oplus \text{Ad}(J_{\mathbf{0}}^{(r)} G_x^k)$$

preserves the Lyapunov decomposition of $T^{(r)}$.

Remark 3.8. *Viewing $J_{\mathbf{0}}^{(r)} F_x^k$ as a polynomial map in $GL^{(r)}(n)$, we see from remark 2.4 that it tends to $\mathbf{0}$ as $k \rightarrow \infty$.*

Proof: The proof only makes use of the Lyapunov decomposition of $\{T^{(r)}(t, x)\}$; recall from proposition 3.5 (2),

$$W_x^{(r),0} \cap \mathfrak{gl}^{(r)}(n) = \mathfrak{h}^{(r),0}$$

The proof will apply essentially verbatim to $\{G_x^k\}$.

First, $T_x^t = D_0 F_x^t$ preserves the Lyapunov decomposition $W^1 \oplus \cdots \oplus W^s$, which implies that, for all $x \in M_T$,

$$(6) \quad J_0^{(1)} F_x^t \in \mathcal{H}^{(1),0} = \mathrm{GL}(W^1) \times \cdots \times \mathrm{GL}(W^s)$$

For $j \geq 2$, let $g = J_0^{(j)} F_x^t$, and $\bar{g} = \rho_{j-1}^j(g)$. Write $g = \bar{g} + \gamma$, with $\gamma \in S^{(j)}(n)$, and suppose $\bar{g} \in \mathcal{H}^{(j-1),0}$. Let Λ be as in section 3.2, and note that $\Lambda \in \mathcal{H}^{(1),0} \subset \mathfrak{h}^{(j),0}$. By (6), Λ commutes with $D_0 F_x^t$ for all $t \in \mathbf{R}$, $x \in M_T$. By similar computations to equations (4) and (5),

$$g \circ \Lambda \circ g^{-1} \equiv \gamma \circ \Lambda \circ D_0 g^{-1} - \Lambda \circ \gamma \circ D_0 g^{-1} \pmod{\mathfrak{h}^{(j),0}}$$

On the other hand, this conjugate belongs to $\mathfrak{h}^{(j),0}$, and both Λ and $D_0 g^{-1}$ preserve all the Lyapunov spaces W^i , so

$$\Lambda \circ \gamma \circ \Lambda^{-1} \equiv \gamma \pmod{\mathfrak{h}_j^0},$$

which implies $\gamma \in \mathfrak{h}_j^0$. Proceeding inductively gives $J_0^{(r)} F_x^t \in \mathcal{H}^{(r),0}$ for all t and all $x \in M_T$. \diamond

3.4. Dynamical submanifolds as reductions of $\mathcal{E}^{(r)}$. In this section we apply theorem 2.1 to $\{F^{(r)}(t, x)\}$ and interpret the resulting submanifolds as invariant $\mathcal{X}^{(r)}$ -reductions of $\mathcal{E}^{(r)}$. Combining this information with proposition 3.6 gives an $\mathcal{H}^{(r)}$ -reduction of $\mathcal{E}^{(r)}$ invariant by the natural action of $\{F_x^t\}$.

First we must recall an aspect of the construction of the $\hat{\nu}_x^\lambda$ that will play a role in our upcoming proof.

3.4.1. Digression on Ruelle's proof: expressing as graphs. For $\lambda^{(i)} < \lambda < \lambda^{(i+1)}$, the submanifold $\hat{\nu}_x^\lambda$ is defined on [16, p 47] as the image under a certain map Φ of the graph of a C^1 function

$$\varphi : V_x^i \cap B(\alpha(x)) \rightarrow (V_x^i)^\perp \cap B(\alpha(x))$$

Write $\alpha = \alpha(x)$. The equation [16, (5.15)] gives the bound

$$\|D_u \varphi\| \leq \frac{A\delta\sqrt{1 - (A\delta)^2}}{1 - 2(A\delta)^2} \quad \forall u \in V_x^i \cap B(\alpha)$$

where A and δ are the constants given by the perturbation theorem as described in subsection 2.3.1. In equation (5.9) of [16], the additional bound $\delta < 1/\sqrt{2}A$ was imposed. There is no problem in assuming $\delta < 1/2A$, which makes $\|D_u \varphi\| < 1$.

The map $\Phi : (V_x^i \cap B(\alpha)) \oplus ((V_x^i)^\perp \cap B(\alpha)) \rightarrow B(\alpha)$ is

$$\Phi(u, v) = \frac{u}{\alpha} \sqrt{\alpha^2 - \|v\|^2} + v$$

For a differentiable path $\gamma(t) = (u(t), v(t))$ in $\hat{L}_b = \Phi^{-1}(\{u = b\})$,

$$\dot{u} = \frac{\langle v, \dot{v} \rangle}{\alpha^2 - \|v\|^2} \cdot u$$

If γ is contained in $B(\alpha/\sqrt{2})$, then $\|\dot{v}(t)\| \geq \|\dot{u}(t)\|$ along γ and $\|v(1) - v(0)\| \geq \|u(1) - u(0)\|$. Now graph φ intersects $\hat{L}_b \cap B(\alpha/\sqrt{2})$ in at most one point, and

$$\dot{v}_x^\lambda \cap B(\alpha/2) \subseteq \Phi(\text{graph } \varphi \cap B(\alpha/\sqrt{2})) \subseteq \dot{v}_x^\lambda \cap B(\alpha/\sqrt{2})$$

is also the graph of a C^1 function. End of digression.

A fiber $\mathcal{E}_x^{(r)} = \mathbf{R}^n \times \text{GL}^{(r)}(n)$ can be viewed as an open subset of \mathbf{R}^N with origin at $(\mathbf{0}, \text{Id})$. Pulling back the norm $\|\cdot\|$ on $\mathfrak{gl}^{(r)}(n)$ by ω yields a Finsler metric on $\text{GL}^{(r)}(n)$, and thus a product Finsler metric on each $\mathcal{E}_x^{(r)}$. This metric is comparable to the Euclidean metric on \mathbf{R}^N on a compact neighborhood of the origin.

The verification that, for $r \leq q$,

$$\sup_{0 \leq \epsilon \leq 1} \ln^+ \|F^{(r)}(\pm\epsilon, x)\|_{q-r} \in L^1(M, \mu)$$

is left to the reader, with the indication that it resembles the verification at the beginning of section 3.3 that $\{T^{(r)}(t, x)\}$ satisfies (MET). Let M_F be the set where the conclusion of the Ergodic Theorem holds for this function under $\{\varphi^t\}$. Let $\sigma < 0$ in Σ^l , $l \leq r$. Theorem 2.1 gives C^{q-r} -smooth submanifolds $\nu_x^{(r), \sigma} \subset \mathcal{E}_x^{(r)}$ for all $x \in M_0 = M_F \cap M_T$. Denote by $\nu_x^{(r), s}$ the submanifolds associated to $\sigma = \lambda^{(s)}$. The following key result is a nonlinear analogue of proposition 3.5 (2).

Proposition 3.9. *For all x in a φ^t -invariant, μ -conull subset of M_0 and $1 \leq r \leq q$, $\nu_x^{(r), s}$ is a C^{q-r} -smooth reduction of $\mathcal{F}^{(r)}\mathcal{E}_x$ to $\mathcal{X}^{(r)}$, projecting onto ν_x^s under π_0^r .*

Proof:

Step 1: Fibers under π_{r-1}^r tangent to $\mathfrak{r}^{(r)}$. For $(w, v) \in T_{(u, g)}\mathcal{E}_x^{(r)}$ the effect of the action of $F^{(r)}(t, x)$ in the Finsler metric $\|\cdot\|_{(u, g)}$ is

$$\|D_{(u, g)}F^{(r)}(t, x) \cdot (w, v)\|^2 = \|D_u F_x^t(w)\|^2 + \|(\text{Ad } J_0^{(r)} F_x^t)(\omega(v))\|^2$$

If $(u, g) \in \nu_x^{(r), s}$, then $F^{(r)}(k, x) \cdot (u, g)$ is in the Euclidean ball $\overline{B(1)}$ for all sufficiently large k . The characterization of $\nu_x^{(r), s}$ in theorem 2.1 can equivalently be expressed in terms of our Finsler distance d .

Let $r \geq 2$ and $v \in T_{(u, g)}\nu_x^{(r), s} \cap \ker(\pi_{r-1}^r)_*$. Then

$$\lim_{k \rightarrow \infty} \frac{1}{k} \ln \|F^{(r)}(k, x)_* v\| = \lim_{k \rightarrow \infty} \frac{1}{k} \ln \|(\text{Ad } J_0^{(r)} F_x^k) \cdot (\omega(v))\| \leq \lambda^{(s)}$$

which implies, by proposition 3.4 (2),

$$\omega(v) \in \widehat{V}_r^{\lambda^{(s)}} = \mathfrak{r}^{(r)} \cap S^{(r)}(n)$$

We denote this last space by \widehat{V}_r^s below. The above implication is also clear when $r = 1$, with the conclusion $\omega(v) \in \widehat{V}_1^s$.

Conversely, if $X \in \widehat{V}_r^s$ with $\|X\| = 1$, then, by proposition 3.4 (2),

$$\lim_{k \rightarrow \infty} \frac{1}{k} \ln \|(J_{\mathbf{0}}^{(r)} F_x^k) \circ X \circ (J_{\mathbf{0}}^{(r)} F_x^k)^{-1}\| \leq \lambda^{(s)}.$$

Again, the same bound holds in the case $r = 1$. For $g' = g \cdot e^{\eta X}$, $\eta \in \mathbf{R}$,

$$\begin{aligned} d(F^{(r)}(k, x) \cdot (u, g'), F^{(r)}(k, x) \cdot (u, g)) &\leq |\eta| \cdot \|D_{(u, h)} F^{(r)}(k, x) \cdot (0, \omega^{-1}(X))\| \\ &= |\eta| \cdot \|(\text{Ad } J_{\mathbf{0}}^{(r)} F_x^k)(X)\| \end{aligned}$$

where h is some point in $\{g e^{t\eta X}\}_{t=0}^1$. As (u, g) belongs to $\nu_x^{(r), s}$,

$$\lim_{k \rightarrow \infty} \frac{1}{k} \ln d\left(F^{(r)}(k, x) \cdot (u, g'), (\mathbf{0}, \text{Id})\right) \leq \lambda^{(s)}$$

and $(u, g') \in \nu_x^{(r), s}$ for $\eta \in \mathbf{R}$ (see proposition 2.2). Thus for all r ,

$$\widehat{V}_r^s = \omega(T_{(u, g)} \nu_x^{(r), s} \cap \ker(\pi_{r-1}^r)_*).$$

Moreover, $\nu_x^{(r), s}$ is saturated by the principal action of $\mathcal{X}^{(r)} \cap \ker \rho_{r-1}^r$.

Step 2: Projection under π_{r-1}^r is onto.

The image $\pi_{r-1}^r(\nu_x^{(r), s})$ is clearly contained in $\nu_x^{(r-1), s}$. The image of $T_{(\mathbf{0}, \text{Id})} \nu_x^{(r), s}$ under $(\pi_{r-1}^r)_*$ is, by proposition 3.5 (2) and theorem 2.1,

$$V^{(r), s} / \widehat{V}_r^s = V^{(r-1), s} = T_{(\mathbf{0}, \text{Id})} \nu_x^{(r-1), s}$$

Thus for x in a φ^t -invariant, conull set, $\pi_{r-1}^r(\nu_x^{(r), s})$ and $\nu_x^{(r-1), s}$ coincide in a neighborhood of $(\mathbf{0}, \text{Id})$.

Choose λ between $\lambda^{(s)}$ and the next greater element of $\text{Spec } T^{(r)}$; choose $\lambda/3 < \zeta < 0$. We claim that for all $k \geq 0$, there exists $m \geq 0$ such that

$$F^{(r-1)}(m, \varphi^k(x)) \cdot \mathring{\nu}_{\varphi^k(x)}^{(r-1), \lambda} \subset \pi_{r-1}^r \cdot \nu_{\varphi^{k+m}(x)}^{(r), s}$$

The intersection of these contains a neighborhood in each of $(\mathbf{0}, \text{Id})$. From the proof of proposition 2.2, the radii of the left-hand terms are bounded above by $\beta(\varphi^k(x)) e^{m\lambda}$. Also, $\nu_{\varphi^{k+m}(x)}^{(r), s}$ contains a ball of radius $\alpha(\varphi^{k+m}(x)) > C e^{(k+m)\zeta}$; as the restriction of π_{r-1}^r is the quotient by the proper action of $\mathcal{X}^{(r)} \cap \ker \rho_{r-1}^r$, the projections contain balls of comparable radius. The

claim follows. Finally,

$$\begin{aligned}
\pi_{r-1}^r \cdot \nu_x^{(r),s} &= \pi_{r-1}^r \cdot \bigcup_{k \geq 0} F^{(r)}(k, x)^{-1} \cdot \mathring{\nu}_{\varphi^k(x)}^{(r),\lambda} \\
&= \pi_{r-1}^r \cdot \bigcup_{k \geq 0} F^{(r)}(k, x)^{-1} \cdot \nu_{\varphi^k(x)}^{(r),s} \\
&= \bigcup_{k \geq 0} F^{(r-1)}(k, x)^{-1} \cdot \pi_{r-1}^r \cdot \nu_{\varphi^k(x)}^{(r),s} \\
&\supset \bigcup_{k \geq 0} F^{(r-1)}(k, x)^{-1} \cdot \mathring{\nu}_{\varphi^k(x)}^{(r-1),\lambda} \\
&= \nu_x^{(r-1),s}
\end{aligned}$$

Step 3: Fibers under π_0^r tangent to $\mathfrak{r}^{(r)}$.

Now $\pi_0^r = \pi_0^1 \circ \dots \circ \pi_{r-1}^r$ maps $\nu_x^{(r),s}$ onto ν_x^s . The vertical tangent subspaces of $\nu_x^{(r),s}$ evaluate under ω to subspaces

$$\mathfrak{h}_x^{(r)}(u, g) = \omega(T_{(u,g)} \nu_x^{(r),s} \cap \ker(\pi_0^r)_*) \subset \mathfrak{gl}^{(r)}(n)$$

with the same associated graded algebra as $\mathfrak{r}^{(r)}$. Note that

$$\mathfrak{h}_{\varphi^t(x)}^{(j)}(F^{(j)}(t, x) \cdot (u, g)) = (\text{Ad } J_{\mathbf{0}}^{(j)} F_x^t) \cdot \mathfrak{h}_x^{(j)}(u, g)$$

for $1 \leq j \leq r$, and they vary smoothly in (u, g) and measurably in x .

Suppose that $\mathfrak{h}_x^{(j-1)}(u, g) = \mathfrak{r}^{(j-1)}$ for almost all x , for all u, g . As $\mathfrak{h}_x^{(j)}(u, g) \cap S^{(j)}(n) = \widehat{V}_j^s$ from step 1, we can define linear functions

$$\tau_x(u, g) : \mathfrak{r}^{(j-1)} \rightarrow S^j(n)/\widehat{V}_j^s \quad \text{by} \quad \mathfrak{h}_x^{(j)}(u, g) = \text{graph}(\tau_x(u, g)) + \widehat{V}_j^s$$

Note that τ corresponds to a tensor on $\cup_x \nu_x^{(r),s}$, smooth in (u, g) and $F^{(j)}$ -invariant—that is,

$$\tau_{\varphi^t(x)}(F^{(j)}(t, x) \cdot (u, g)) = (\text{Ad } J_{\mathbf{0}}^{(j)} F_x^t) \circ \tau_x(u, g) \circ (\text{Ad } J_{\mathbf{0}}^{(j-1)} F_x^t)^{-1}$$

Under this action, $\tau_x(u, g)$ is stable as $t \rightarrow \infty$ by proposition 2.15. The stable subspace is trivial, so $\tau_x(u, g) = 0$ for almost-every $x \in M_0$, for all $(u, g) \in \nu_x^{(j),s}$. Induction on j yields $\mathfrak{h}_x^{(r)}(u, g) = \mathfrak{r}^{(r)}$.

Step 4: Fibers of π_0^r are connected, equal $\mathcal{X}^{(r)}$ -orbits.

For $x \in M_0$, write $B_x = B(\alpha(x)/2) \subset \mathcal{E}_x^{(r)}$. From subsection 3.4.1, $\mathring{\nu}_x^{(r),\lambda} \cap B_x$ is the graph of a function from a neighborhood of $\mathbf{0}$ in $V_x^{(r),s}$ to $(V_x^{(r),s})^\perp$. From proposition 3.5 (2), $V_x^{(r),s} = \mathbf{R}^n \oplus \widehat{V}^{(r),s}$. Then $(\pi_0^r)^{-1}(u) \cap \mathring{\nu}_x^{(r),\lambda} \cap B_x$ is the graph of a function on $\widehat{V}^{(r),s}$, so is connected.

Given $k \geq 0$, there exists $m \geq 0$ such that

$$\beta(\varphi^k(x))e^{m\lambda} < \frac{Ce^{(k+m)\zeta}}{2} < \frac{\alpha(\varphi^{k+m}(x))}{2}$$

so

$$F^{(r)}(m, \varphi^k(x)) \cdot \mathring{\nu}_{\varphi^k(x)}^{(r),\lambda} \subset B_{\varphi^{k+m}(x)}$$

Then

$$\nu_x^{(r),s} \cap (\pi_0^r)^{-1}(u) = \bigcup_{k \geq 0} F^{(r)}(k, x)^{-1} \cdot \left((\pi_0^r)^{-1}(F_x^k u) \cap \mathring{\nu}_{\varphi^k(x)}^{(r),\lambda} \cap B_{\varphi^k(x)} \right)$$

This is an increasing union of connected sets (see the proof of proposition 2.2), so it is connected. Now the fibers of $\nu_x^{(r),s}$ are connected $\mathcal{X}^{(r)}$ -orbits, so $\nu_x^{(r),s}$ is a reduction to $\mathcal{X}^{(r)}$, invariant by $F^{(r)}(t, x)$. \diamond

The prolongation $F^{(r)}(t, x)$ is defined so that it preserves the section $(\mathbf{0}, \text{Id})$. The natural action of F_x^t on $\mathcal{E}^{(r)}$ is by $(u, g) \mapsto (F_x^t(u), J_u^{(r)} F_x^t \cdot g)$. Now we can interpret the $F^{(r)}(t, x)$ -invariant submanifolds $\nu_x^{(r),s}$ in terms of this natural action.

Corollary 3.10. *For any r with $1 \leq r \leq q$, the natural action of $\{F_x^t\}$ on $\mathcal{E}^{(r)}$, and of any $(\{G_x^k\}, \psi) \in Z(F_x^t)$, preserves a reduction $\mathcal{R}^{(r)}$ to $\mathcal{H}^{(r)}$.*

Proof: Let $\mathcal{R}^{(r)}$ be the saturation of $\cup_x \nu_x^{(r),s}$ by the right $\mathcal{H}^{(r)}$ -action. Recall from proposition 3.3 that $\mathcal{X}^{(r)} \triangleleft \mathcal{H}^{(r)}$. By propositions 3.9 and 3.6, $\mathcal{R}^{(r)}$ is the desired F_x^t -invariant reduction.

For $(\{G_x^k\}, \psi) \in Z(F_x^t)$, the prolongation $(\{G^{(r)}(k, x)\}, \psi) \in Z(F^{(r)}(t, x))$. Thus by proposition 2.6, the family $\cup_x \nu_x^{(r),s}$, restricted to an appropriate ψ -invariant, full measure subset, are invariant by $G^{(r)}(k, x)$. By proposition 3.6, $J_0^{(r)} G_x^k \in \mathcal{H}^{(r),0}$, so $\mathcal{R}^{(r)}$ is also G_x^k -invariant. \diamond

Fix $r = \lfloor \lambda^{(1)}/\lambda^{(s)} \rfloor$. We will next interpret the submanifolds $\cup_x \nu_x^{(r),s}$ as a φ^t -invariant family of rigid geometric structures on the fibers of \mathcal{E} .

3.5. Dynamical submanifolds as geometric structures. The reductions obtained in proposition 3.9 and corollary 3.10 above are not generally geometric structures of finite type in the sense of Cartan, because $\mathcal{X}^{(r)}$ contains rank-one elements (see [14, Prop I.1.4]). In this section, we implement a measurable version of Feres' approach in [4] to construct invariant generalized connections on the fibers \mathcal{E}_x and coordinate atlases in which F_x^t acts by resonance polynomials. The automorphism group of each atlas on \mathcal{E}_x is a finite-dimensional Lie group. We further interpret these structures as an invariant family of flat connections on vector bundles over \mathcal{E}_x .

Before stating the main theorem in general, we describe two important cases in detail. Let $M_{\mathcal{R}}$ be the φ^t -invariant μ -conull subset on which the conclusions of proposition 3.9 hold.

Example: $r = 1$. This case arises when $\Sigma^0 = \{\lambda\}$, or when $\lambda^{(s)} < \lambda^{(1)}/2$. In the latter case the spectrum is said to be *1/2-pinched*. Then $\nu_x^{(1),s}$ is a C^{q-1} reduction of $\mathcal{F}^{(1)}(\mathcal{E}_x)$ to $\{\text{Id}\}$. It determines a framing $g_x : \nu_x^s \rightarrow \text{GL}(n)$ for all $x \in M_{\mathcal{R}}$. Invariance by $F^{(1)}(t, x)$ means

$$g_{\varphi^t(x)}(F_x^t(u)) = (J_u^{(1)} F_x^t) \cdot g_x(u) \cdot (J_0^{(1)} F_x^t)^{-1},$$

in other words,

$$g_{\varphi^t(x)} \cdot T_x^t = (F_x^t)_* g_x$$

The tangent spaces $T_\xi(\nu_x^{(1),s})$ project isomorphically to $T_{\pi_0^{-1}(\xi)} \mathcal{E}_x$. The union of right translates of these spaces by $\text{GL}(n)$ is a C^{q-2} horizontal distribution \mathcal{D}_x on $\mathcal{F}^{(1)}(\mathcal{E}_x)$. The $\oplus_x \mathcal{D}_x$ are principal connections, equivalent to C^{q-2} affine connections $\oplus_x \nabla_x$ on $\cup_x \mathcal{E}_x$, invariant by F_x^t .

The torsion $\oplus_x \tau_x$ of $\oplus_x \nabla_x$ is an F_x^t -invariant tensor. The spectrum of $T_{\mathcal{S}}$ on $\mathcal{S}_0 = \oplus_x \wedge^2 T_0^*(\mathcal{E}_x) \otimes T_0(\mathcal{E}_x)$ is bounded below by $\lambda^{(1)} - 2\lambda^{(s)} > 0$. Propositions 2.13 and 2.15 then imply that the connections ∇_x are torsion-free for all x in a φ^t -invariant, μ -conull subset of $M_{\mathcal{R}}$.

One could make a similar argument to show vanishing of the curvature of ∇_x assuming $q > 2$. However, it is clear in any case that they are flat: the sections $\nu_x^{(1),s} \subset \mathcal{F}^{(1)}(\mathcal{E}_x)$ are parallel by construction. Thus $\nu_x^{(1),s}$ correspond to coordinate charts θ_x on \mathcal{E}_x for which $\theta_x^* \nabla_x$ equals the flat connection on \mathbf{R}^n . The resulting flat affine structures on $\cup_x \mathcal{E}_x$ are F_x^t -invariant; in fact, from the F_x^t -invariant framings above, we obtain a family of \mathbf{R}^n -structures—framings by commuting vector fields—invariant by F_x^t up to the linear action of the algebraic hull of $\{T_x^t\}$ in $\mathcal{H}^{(1),0}$.

If $(\{G_x^k\}, \psi) \in Z(F_x^t)$, then $G^{(1)}(k, x)$ preserves $\cup_x \nu_x^{(1),s}$ and $\oplus_x \mathcal{D}_x$ (as usual, over $\cap_{i \in \mathbf{Z}} \psi^i(M_{\mathcal{R}} \cap M_G)$), so the natural action of $\{G_x^k\}$ preserves the flat connections $\oplus_x \nabla_x$. By proposition 3.6, $\{G_x^k\}$ also preserves the \mathbf{R}^n -structures on $\cup_x \mathcal{E}_x$, up to the algebraic hull of $J_0^{(1)} G_x^k$, which is contained in $\mathcal{H}^{(1),0}$.

Remark 3.11. *We sketch how Ruelle's perturbation theorem implies that according to the frames in $\nu_x^{(1),s}$, the Lyapunov filtration is $V^1 \subset \dots \subset V^s$ at every point of ν_x^s : given $\xi \in \nu_x^{(1),s}(u)$, consider the sequences $\{T_k\} = \{T_{\varphi^{k-1}(x)}^1\}$ and $\{T'_k\} = \{D_{F_x^{k-1}(u)} F_{\varphi^{k-1}(x)}^1\}$. For λ and η as in section 2.3.1*

$$\|T'_k - T_k\| e^{k\eta} < A e^{k\zeta}$$

for some constant A and $\lambda < \zeta < 0$, for all $k \geq 0$. Now [16, (4.4)] implies that the i th Lyapunov projections P^i for $\{T^k\} = \{T_k \circ \dots \circ T_1\}$ and P^i for $\{T'^k\} = \{D_u F_x^k\}$ satisfy

$$\|D_u F_x^\ell \circ P'^i \circ (D_u F_x^\ell)^{-1} - P^i\| \leq A' e^{\ell\zeta}$$

for some constant A' , for all $\ell \geq 0$. For $Q'^i = \xi^{-1} \circ P'^i \circ \xi$,

$$\|D_0 F_x^\ell \circ Q'^i \circ (D_0 F_x^\ell)^{-1} - P^i\| \leq B' e^{\ell\zeta}$$

for some B' , where we have used that $F^{(1)}(\ell, x) \cdot \xi$ tends to Id at least as fast as $e^{\ell\lambda}$. It follows that the i th Lyapunov space at u in the frame ξ is congruent to W^i modulo V^i for all i , so the filtrations are equal.

Finally, we remark that \mathcal{D}_x is tangent to the $\mathcal{H}^{(1),0}$ -saturation of $\nu_x^{(1),s}$. As $\mathcal{H}^{(1),0}$ preserves the Lyapunov filtration $V^1 \subset \dots \subset V^s$, the Lyapunov filtration of $T\nu_x^{(1),s}$ is ∇_x -parallel for all $x \in M_{\mathcal{R}}$.

Example: $r = 2$. In this case $3\lambda^{(s)} < \lambda^{(1)} \leq 2\lambda^{(s)}$.

3.5.1. *Conglomeration of Lyapunov filtration to length 2.* Let i_* be the minimal element of $\{1, \dots, s\}$ such that $\lambda^{(1)} - \lambda^{(i_*)} \leq \lambda^{(s)}$. Whenever $j, l < i_*$ or $j, l \geq i_*$, then $\lambda^{(l)} - \lambda^{(j)} > \lambda^{(s)}$. The first case is clear from the definition of i_* . For the second, note

$$\lambda^{(l)} - \lambda^{(j)} \geq \lambda^{(i_*)} - \lambda^{(s)} \geq \lambda^{(1)} - 2\lambda^{(s)} > \lambda^{(s)}$$

If we write $\widetilde{W}^1 = \bigoplus_{j=1}^{i_*-1} W^j$ and $\widetilde{W}^2 = \bigoplus_{j=i_*}^s W^j$, then

$$\mathcal{X}^{(1)} \subset \text{Id} + (\widetilde{W}^2)^* \otimes \widetilde{W}^1$$

3.5.2. *Framings of subquotients.* The $\mathcal{X}^{(1)}$ -reductions $\nu_x^{(1),s} \subset \mathcal{E}_x^{(1)}$ induce distributions \mathcal{V}_x^1 on \mathcal{E}_x corresponding to \widetilde{W}^1 . For $\xi = (u, g) \in \nu_x^{(1),s}$ and $\xi' = F^{(1)}(t, x) \cdot \xi \in \nu_{\varphi^t(x)}^{(1),s}$,

$$\xi' \cdot T_x^t = F_x^t \cdot \xi$$

(the right-hand side is the natural action of F_x^t on $\mathcal{E}^{(1)}$). Since T_x^t preserves \widetilde{W}^1 and \widetilde{W}^2 , the distributions $\bigoplus_x \mathcal{V}_x^1$ are F_x^t -invariant.

Under restriction to \widetilde{W}^1 or projection modulo \widetilde{W}^1 ,

$$\text{Res}_{\widetilde{W}^1}(\mathcal{X}^{(1)}) = \{\text{Id}\} \quad \text{Proj}_{\mathbf{R}^n/\widetilde{W}^1}(\mathcal{X}^{(1)}) = \{\text{Id}\}$$

Therefore, $\bigcup_x \nu_x^{(1),s}$ also induces $F^{(1)}(t, x)$ -invariant framings of $\bigoplus_x \mathcal{V}_x^1$ and $\bigoplus_x (T\mathcal{E}_x)/\mathcal{V}_x^1$. If we denote g_x the framing of \mathcal{V}_x^1 , then

$$(F_x^t)_* g_x = g_{\varphi^t(x)} \circ T_x^t$$

A similar identity holds for the framing of $T\nu_x^s/\mathcal{V}_x^1$.

3.5.3. *Flat connections on $\bigcup_x \mathcal{E}_x$.* As $\mathcal{X}^{(2)} = \mathcal{X}^{(1)}$, the restriction $\pi_1^2 : \nu_x^{(2),s} \rightarrow \nu_x^{(1),s}$ is a diffeomorphism. The inverse induces a C^{q-2} horizontal distribution \mathcal{D}_x on $\nu_x^{(1),s}$ as follows. Suppose that $\hat{\xi} = (u, J_0^{(2)}\varphi) \in \nu_x^{(2),s}$ projects to $\xi \in \nu_x^{(1),s}$. Define a map $\mathbf{R}^n \rightarrow \mathcal{E}_x^{(1)}$

$$(\tau_u \circ \varphi, J^{(1)}\varphi) : v \mapsto (\varphi(v) + u, J_v^{(1)}\varphi)$$

The subspace

$$S(\hat{\xi}) = \text{im } D_0(\tau_u \circ \varphi, J^{(1)}\varphi) \subset T_{\xi} \mathcal{E}_x^{(1)}$$

depends only on $\hat{\xi}$. Now set $\mathcal{D}_x(\xi) = S(\hat{\xi})$. Note that this space projects under $(\pi_0)_*$ onto $T_u\mathcal{E}_x \cong \mathbf{R}^n$. Because $\mathcal{D}_x(\xi)$ comes from a diffeomorphism φ , it is called *holonomic* (see [3, p 29]).

For $h \in GL(n)$, write R_{h*} for the derivative of the right action on $\mathcal{E}^{(1)}$.

Proposition 3.12. *The distribution \mathcal{D}_x defines a C^{q-2} principal connection ∇_x on $\nu_x^{(1),s}$ varying measurably in x ; more precisely, for almost all $x \in M$,*

- (1) $\mathcal{D}_x(\xi) \subset T_\xi \nu_x^{(1),s}$
- (2) $(R_h)_* \mathcal{D}_x(\xi) = \mathcal{D}_x(\xi \cdot h)$ for all $h \in \mathcal{X}^{(1)}$.

The resulting connections ∇_x on \mathcal{E}_x are flat.

Proof: Write $\xi = (u, g)$ and $\hat{\xi} = (u, \hat{g}) = (u, J_0^{(2)}\varphi)$; set $\xi' = F^{(1)}(t, x) \cdot \xi$ and $\hat{\xi}' = F^{(2)}(t, x) \cdot \hat{\xi}$. First we compare $\mathcal{D}_{\varphi^t(x)}(\xi') = S(\hat{\xi}')$ with $F^{(1)}(t, x)_* \cdot \mathcal{D}_x(\xi)$. Write $(F_x^t)_*$ for the derivative of the natural action on $\mathcal{E}^{(1)}$ as in subsection 3.5.2 (although it's ambiguous). Set $\hat{h} = (J_0^{(2)} F_x^t)^{-1}$ and $h = \rho_1^2(\hat{h})$. For the natural action, one can check

$$S(F_x^t \cdot \hat{\xi}) = (F_x^t)_* S(\hat{\xi})$$

For the principal right action, on the other hand, $S(\hat{\xi} \cdot \hat{h})$ is given by

$$(7) \quad \begin{aligned} D_0(\tau_u \circ \varphi \circ \hat{h}, J^{(1)}(\varphi \circ \hat{h})) &= (R_h)_* \circ D_0(\tau_u \circ \varphi, J^{(1)}\varphi) \\ &\quad + \omega^{-1} \circ D_0(J^{(1)}\hat{h}) \end{aligned}$$

(see [4, eqn (54)]). Because $\hat{h} \in \mathcal{H}^{(2),0}$ by proposition 3.6, the image of $D_0(J^{(1)}\hat{h})$ must be strict subresonance—that is, in $\mathfrak{r}^{(1)}$. In conclusion,

$$S(\hat{\xi}') \equiv (F_x^t)_* \circ R_{h*} \cdot S(\hat{\xi}) \pmod{\omega^{-1}(\mathfrak{r}^{(1)})}$$

and thus

$$(8) \quad \mathcal{D}_{\varphi^t(x)}(\xi') \equiv F^{(1)}(t, x)_* (\mathcal{D}_x(\xi)) \pmod{\omega^{-1}(\mathfrak{r}^{(1)})}.$$

Now define an $F^{(1)}(t, x)$ -invariant family of sections τ_x of $(T\mathcal{E}_x)^* \otimes (\text{End}(T\mathcal{E}_x)/\omega^{-1}(\mathfrak{r}^{(1)}))$, restricted to $\nu_x^{(1),s}$, by

$$\mathcal{D}_x(\xi) \pmod{\omega^{-1}(\mathfrak{r}^{(1)})} = \text{graph}(\tau_x(\xi))$$

The stability of τ given by proposition 2.15 (together with proposition 2.13) implies that τ_x vanishes for almost every x . Thus \mathcal{D}_x is tangent to $\nu_x^{(1),s}$ for almost every $x \in M$ and (1) is verified.

Point (2) follows from the fact that $\nu_x^{(2),s}$ and $\nu_x^{(1),s}$ are diffeomorphic reductions to $\mathcal{X}^{(1)} \subset GL^{(1)}(n)$. If $\hat{\xi} \in \nu_x^{(2),s}$ projects to $\xi \in \nu_x^{(1),s}$, and $h \in \mathcal{X}^{(1)}$, then by equation (7),

$$\mathcal{D}_x(\xi \cdot h) = S(\hat{\xi} \cdot h) = (R_h)_* \mathcal{D}_x(\xi).$$

Now that we have \mathcal{D}_x tangent to $\nu_x^{(1),s}$ for almost every x , and given that $\nu_x^{(2),s}$ is at least C^1 , we can argue as in [4, sec 5.1], [3] that the reductions $\nu_x^{(2),s}$ define a complete and consistent partial differential relation, and therefore, the distributions \mathcal{D}_x are integrable. It follows that the corresponding connections ∇_x on \mathcal{E}_x are flat. The integral leaves of \mathcal{D}_x give a family of affine charts on \mathcal{E}_x with transitions in the group $\mathcal{X}^{(1)}$. \diamond

3.5.4. *Flat connections on $\oplus_x \mathcal{V}_x^1$ and $\oplus_x (T\mathcal{E}_x)/\mathcal{V}_x^1$ invariant by $Z(F_x^t)$.* The connections $\oplus_x \nabla_x$ are not invariant by the natural action of F_x^t on $\cup_x \mathcal{E}_x$. By proposition 3.10, the natural action preserves an $\mathcal{H}^{(2)}$ -reduction $\mathcal{R}^{(2)} \subseteq \mathcal{E}^{(2)}$, containing the saturation of $\cup_x \nu_x^{(2),s}$ by the right action of $J_0^{(2)} F_x^t$. We can extend \mathcal{D} to a connection on $\mathcal{R}^{(1)} = \pi_1^2(\mathcal{R}^{(2)})$ using the right action of $\mathcal{H}^{(1)}$ (it is still not F_x^t -invariant).

We remark that there is an F_x^t -invariant horizontal distribution on $\mathcal{R}^{(2)}$. It does not give a connection, but it is integrable. See the general statements in theorem 3.13 (1) below.

From section 3.5.2, the distributions $\oplus_x \mathcal{V}_x^1$ are F_x^t -invariant. There are thus F_x^t -equivariant maps $\text{Res}_{\mathcal{V}^1} : \mathcal{R}^{(1)} \rightarrow \mathcal{F}\mathcal{V}^1$ and $\text{Proj}_{T\mathcal{E}/\mathcal{V}^1} : \mathcal{R}^{(1)} \rightarrow \mathcal{F}(T\mathcal{E}/\mathcal{V}^1)$. Note that $\omega^{-1}(\mathfrak{r}^{(1)})$ is in the kernel of both $(\text{Res}_{\mathcal{V}^1})_*$ and $(\text{Proj}_{T\mathcal{E}/\mathcal{V}^1})_*$.

Let $\xi \in \mathcal{R}^{(1)}$, with $F_x^t \cdot \xi = \xi'$. Writing $h = J_0^{(1)} F_x^t$, we have

$$(F_x^t)_* \mathcal{D}_x(\xi) = (R_h)_* \circ F^{(1)}(t, x)_* \mathcal{D}_x(\xi)$$

From equation (8) and the fact that h normalizes $\mathcal{X}^{(1)}$,

$$(F_x^t)_* \mathcal{D}_x(\xi) \bmod \omega^{-1}(\mathfrak{r}^{(1)}) \equiv (R_h)_* \mathcal{D}_{\varphi^t(x)}(\xi' \cdot h^{-1}) = \mathcal{D}_{\varphi^t(x)}(\xi')$$

Pushing forward \mathcal{D} by $(\text{Red}_{\mathcal{V}^1})_*$ or $(\text{Proj}_{T\mathcal{E}/\mathcal{V}^1})_*$ thus gives F_x^t -invariant, flat connections on the vector bundles $\oplus_x \mathcal{V}_x^1$ and $\oplus_x T\mathcal{E}_x/\mathcal{V}_x^1$. Remark that by corollary 3.10, proposition 2.6, and proposition 3.6, the distributions $\oplus_x \mathcal{V}_x^1$ and these flat connections are moreover invariant by all of $Z(F_x^t)$. Because \mathcal{D}_x is tangent to $\mathcal{R}_x^{(1)}$ for almost all x and $\mathcal{H}^{(1)}$ preserves the filtrations of \widetilde{W}^1 and $\mathbf{R}^n/\widetilde{W}^1$ determined by $V^1 \subset \dots \subset V^{i_*}$ and $V^{i_*}/\widetilde{W}^1 \subset \dots \subset V^s/\widetilde{W}^1$, respectively, the Lyapunov filtrations of \mathcal{V}_x^1 and $T\mathcal{E}_x/\mathcal{V}_x^1$ are parallel for these connections, for almost all x (see remark 3.11).

Main Theorem: Let $r = \lfloor \lambda^{(1)}/\lambda^{(s)} \rfloor$ and $q \geq r + 1$.

Theorem 3.13. *There exist the following differential-geometric structures on \mathcal{E}_x , for all x in a φ^t -invariant, μ -conull subset:*

- (1) *A family \mathcal{A}_x of C^q charts, with transitions in $\mathcal{H}^{(r),0}$ (restricted to a neighborhood of $\mathbf{0}$). The collection $\cup_x \mathcal{A}_x$ is $Z(F_x^t)$ -invariant.*

- (2) A C^{q-1} filtration $\mathcal{V}_x^1 \subset \cdots \subset \mathcal{V}_x^l = T\mathcal{E}_x$, with $l \leq r$, equipped with C^{q-2} -smooth flat connections ∇_x^i on $\mathcal{V}_x^i/\mathcal{V}_x^{i-1}$, $i = 1, \dots, l$. The filtrations and connections are invariant by $Z(F_x^t)$.

Proof: Denote $\mathcal{R}^{(r)} = \left(\cup_x \nu_x^{(r),s} \right) \cdot \mathcal{H}^{(r)}$ the $Z(F_x^t)$ -invariant $\mathcal{H}^{(r)}$ -reduction of $\mathcal{E}^{(r)}$ given by corollary 3.10. We similarly have an invariant $\mathcal{R}^{(r+1)} \subset \mathcal{E}^{(r+1)}$, which is C^{q-r-1} -smooth. As $\mathcal{H}^{(r)} = \mathcal{H}^{(r+1)}$, the restriction $\pi_r^{r+1} : \mathcal{R}^{(r+1)} \rightarrow \mathcal{R}^{(r)}$ is a diffeomorphism.

We first construct a $Z(F_x^t)$ -invariant family of holonomic distributions $\mathcal{D}_x(\xi) = S(\hat{\xi})$ on $\cup_x \nu_x^{(r),s}$ as in subsection 3.5.3; here $\hat{\xi}$ is the unique lift of $\xi \in \cup_x \nu_x^{(r),s} \subset \mathcal{R}^{(r)}$ to $\mathcal{R}^{(r+1)}$. If $\hat{h} = J_{\mathbf{0}}^{(r+1)} F_x^t \in \mathcal{H}^{(r+1),0}$, viewed as a local diffeomorphism fixing $\mathbf{0}$, then $D_{\mathbf{0}}(J^{(r)} \hat{h})$ has image in $\mathfrak{r}^{(r)}$. We follow the proof of proposition 3.12 (1), and equation (8) becomes

$$\mathcal{D}_{\varphi^t(x)}(\xi') \equiv F^{(r)}(t, x)_*(\mathcal{D}_x(\xi)) \quad \text{mod } \omega^{-1}(\mathfrak{r}^{(r)})$$

Then we can construct an invariant tensor σ with values in the restriction to $\cup_x \nu_x^{(r),s}$ of

$$\mathcal{T} = (T\mathcal{E}_x)^* \otimes (\mathfrak{gl}^{(r)}(\mathcal{E}_x)/\omega^{-1}(\mathfrak{r}^{(r)}))$$

Here $\mathfrak{gl}^{(r)}(\mathcal{E}_x)$ denotes the tensor bundle over $\mathcal{E}_x^{(r)}$ with fiber

$$\mathfrak{gl}_{(u,g)}^{(r)}(\mathcal{E}_x) = (\tau_u \circ g) \cdot \mathfrak{gl}^{(r)}(n) \cdot (\tau_u \circ g)^{-1}$$

Vanishing of σ is implied by proposition 2.15, and we conclude that the restriction to $\nu_x^{(r),s}$ of \mathcal{D}_x is tangent to $\nu_x^{(r),s}$ for all x in a φ^t -invariant, μ -conull set.

The distributions $\oplus_x \mathcal{D}_x$ are integrable, again by Gromov's Frobenius theorem, because $\cup_x \nu_x^{(r),s}$ is a complete and consistent partial differential relation; moreover, because they are holonomic, each leaf represents the r -jets of (the germ of) a diffeomorphism $(\mathbf{R}^n, \mathbf{0}) \rightarrow (\mathcal{E}_x, \mathbf{0})$.

Now we extend these distributions to $\mathcal{R}^{(r)}$ by the same formula: $\mathcal{D}_x(\xi) = S(\hat{\xi})$. Any $\xi \in \mathcal{R}_x^{(r)}$ equals $\xi' \cdot h$ for $\xi' \in \nu_x^{(r),s}$ and $h \in \mathcal{H}^{(r)}$. Equation (7) gives that $S(\hat{\xi}' \cdot \hat{h}) = \mathcal{D}_x(\xi)$ is tangent to $\mathcal{R}_x^{(r)}$. Note that $\oplus_x \mathcal{D}_x$ are invariant by the natural $Z(F_x^t)$ -action. Given $\xi = (\mathbf{0}, g) \in \mathcal{R}_x^{(r)}(\mathbf{0})$, denote α_ξ the germ of a coordinate parametrization of \mathcal{E}_x with $\alpha_\xi(\mathbf{0}) = \mathbf{0}$, $J_{\mathbf{0}}^{(r)} \alpha_\xi = g$ and

$$\text{im } D_u(\alpha_\xi, J^{(r)} \alpha_\xi) = \mathcal{D}_x(\alpha_\xi(u), J_u^{(r)} \alpha_\xi)$$

for all u in the domain of α_ξ . Note that this exists for all $\xi \in \nu_x^{(r),s}(\mathbf{0})$.

Given α_ξ as above for $\xi \in \mathcal{R}_x^{(r)}(\mathbf{0})$, for $\eta = (\alpha_\xi(u), J_u^{(r)} \alpha_\xi)$, the lift $\hat{\eta}$ to $\mathcal{R}^{(r+1)}$ equals $(\alpha_\xi(u), J_u^{(r+1)} \alpha_\xi)$: let φ be a representative of $\hat{\eta}$. Then $\eta = (\varphi(\mathbf{0}), J_{\mathbf{0}}^{(r)} \varphi)$; in particular, $J_{\mathbf{0}}^{(1)} \varphi = J_u^{(1)} \alpha_\xi$. Also, $S(\hat{\eta})$ equals $\text{im } D_u(\alpha_\xi, J^{(r)} \alpha_\xi)$.

The two equalities together imply $J_{\mathbf{0}}^{(r+1)}\varphi = J_u^{(r+1)}\alpha_\xi$, which proves the claim.

Now given $h \in \mathcal{H}^{(r)} = \mathcal{H}^{(r+1)}$ and u in the domain of $\alpha_\xi \circ h$, let $\eta = (\alpha_\xi(h(u)), J_{h(u)}^{(r)}\alpha_\xi) \in \mathcal{R}_x^{(r)}$. We can write

$$((\alpha_\xi \circ h)(u), J_u^{(r)}(\alpha_\xi \circ h)) = \eta \cdot (\tau_{-h(u)} \circ h \circ \tau_u)$$

Note that $\tau_{-h(u)} \circ h \circ \tau_u \in \mathcal{H}^{(r+1)}$. The lift of our point to $\mathcal{R}^{(r+1)}$ is

$$(\alpha_\xi(h(u)), J_{h(u)}^{(r+1)}\alpha_\xi) \cdot (\tau_{-h(u)} \circ h \circ \tau_u) = ((\alpha_\xi \circ h)(u), J_u^{(r+1)}(\alpha_\xi \circ h))$$

The r -jets of $\alpha_\xi \circ h$ thus lie in the integral leaf of \mathcal{D}_x through $\xi \cdot h$. We conclude $\alpha_{\xi h} = \alpha_\xi \circ h$ for all $\xi \in \mathcal{R}_x^{(r)}(\mathbf{0})$ and $h \in \mathcal{H}^{(r)}$. In particular, given α_ξ for $\xi \in \nu_x^{(r),s}(\mathbf{0})$, we can define $\alpha_{\xi \cdot h} = \alpha_\xi \circ h$.

Now $Z(F_x^t)$ preserves $\cup_x \mathcal{A}_x$, and each \mathcal{A}_x is parametrized by $\mathcal{R}_x^{(r)}(\mathbf{0}) = \{\mathbf{0}\} \times \mathcal{H}^{(r)}$, so the action of F_x^t on \mathcal{E}_x is determined by $J_{\mathbf{0}}^{(r)}F_x^t$ (the same is true for any $\{G_x^k\}$ in $Z(F_x^t)$). By proposition 3.6, the subatlas

$$\mathcal{A}_x^0 = \{\alpha_\xi : \xi \in \{\mathbf{0}\} \times \mathcal{H}^{(r),0}\}$$

is $Z(F_x^t)$ -invariant. The transitions between the charts in \mathcal{A}_x^0 belong to $\mathcal{H}^{(r),0}$. Point (1) is proved.

The first step to prove part (2) is to conglomerate the Lyapunov filtration as in subsection 3.5.1. Recall that $(r+1)\lambda^{(s)} < \lambda^{(1)} \leq r\lambda^{(s)}$. Set $i_0 = 1$, and recursively define i_k to be the minimal $i > i_{k-1}$ with $\lambda^{(i_{k-1})} - \lambda^{(i_k)} \leq \lambda^{(s)}$. It is a simple induction argument to see that $\lambda^{(i_k)} - (r-k+1)\lambda^{(s)} > 0$. It follows that the process terminates with i_{r_*-1} where $r_* - 1 < r$. Set $i_{r_*} = s+1$, and

$$\widetilde{W}^l = \bigoplus_{j=i_{l-1}}^{i_l-1} W^j$$

Then $\widetilde{W}^1 \oplus \cdots \oplus \widetilde{W}^{r_*}$ is an $\mathcal{H}^{(1),0}$ -invariant decomposition, of length at most r , with respect to which $\mathcal{X}^{(1)}$ is block upper-triangular. Then, as in section 3.5.2, we obtain C^{q-1} filtrations

$$0 \subset \mathcal{V}_x^1 \subset \cdots \subset \mathcal{V}_x^{r_*} = T\mathcal{E}_x$$

invariant by $Z(F_x^t)$. For $1 \leq l \leq r$, define a map on $\mathcal{R}^{(r)}$

$$Q_l = \text{Proj}_{\mathcal{V}^l/\mathcal{V}^{l-1}} \circ \text{Res}_{\mathcal{V}^l} \circ \pi_1^r$$

Note that \mathcal{D}_x is not projectible under $(\pi_1^r)_*$ because it is not right $\mathcal{H}^{(r)}$ -invariant; however, it does descend under $(Q_l)_*$ to a horizontal distribution on $\mathcal{F}(\mathcal{V}^l/\mathcal{V}^{l-1})$: The degree r analogue of equation (7) says that \mathcal{D}_x is right- $\mathcal{H}^{(r)}$ -invariant modulo $\omega^{-1}(\mathfrak{r}^{(r)})$. But $\omega^{-1}(\mathfrak{r}^{(1)})$ is in the kernel of $(\text{Proj}_{\mathcal{V}^l/\mathcal{V}^{l-1}} \circ \text{Res}_{\mathcal{V}^l})_*$. By the same reasoning, the projection of \mathcal{D}_x to a

horizontal distribution on $\mathcal{F}(\mathcal{V}^\ell/\mathcal{V}^{\ell-1})$ gives a C^{q-2} principal connection $\bar{\nabla}^\ell$. The principal group here is $\mathrm{GL}(\widetilde{W}^\ell)$, which can be obtained as a quotient of $\mathcal{H}^{(r)}$. Because \mathcal{D}_x is integrable, so is the projection by $(Q_l)_*$; thus $\bar{\nabla}^\ell$ is flat.

As \mathcal{D}_x is tangent to \mathcal{R}_x , the Lyapunov filtration of $\mathcal{V}_x^\ell/\mathcal{V}_x^{\ell-1}$ is $\bar{\nabla}^\ell$ -parallel for all ℓ , for almost all x , as in remark 3.11 and subsection 3.5.3. \diamond

3.6. Foliated case: Smooth geometric structures on leaves. We return to the important special case that M is a compact C^0 manifold and the fibers \mathcal{E}_x are plaques of a φ^t -invariant foliation; we assume that the action on this foliation is contracting. More precisely, let L , as in section 2.2, be a C^0 foliation of M by n -dimensional submanifolds admitting a φ^t -invariant C^q smooth structure, and assume $\sup_{0 \leq t \leq 1} \|\|J_x^q(\varphi^{\pm t}|_L)\|\|$ is bounded in x (for example, it is continuous in x). Let $\{(\mathcal{E}_x, \theta_x)\}$ be a uniformly biLipschitz C^q atlas along L . Define the cocycle $\{F_x^t\}$ as in section 2.2, and assume that the Lyapunov exponents for $\{T_x^t = D_0 F_x^t\}$ are all negative.

Before interpreting the results of the previous section in this setting, we must assemble the manifolds $\nu_x^{(r),s}$ associated to the prolongation of $\{F_x^t\}$ into an invariant submanifold of the r -frame bundle $\cup_x \mathcal{F}^{(r)} L_x = \cup_x L_x^{(r)}$ along L .

Proposition 3.14. *Let $r \leq q - 1$. For all x in a φ^t -invariant, μ -conull subset of M ,*

- (1) *there exists a neighborhood $x \in U_x \subset L_x$, and a C^{q-r} -smooth reduction $\mu_x^{(r),s}$ of $L_x^{(r)}|_{U_x}$ to $\mathcal{X}^{(r)}$, such that*
- (2) *$\cup_{y \in L_x} \mu_y^{(r),s} \cdot \mathcal{H}^{(r)}$ is a C^{q-r} -smooth $\mathcal{H}^{(r)}$ -reduction $\mathcal{P}_x^{(r)} \subset L_x^{(r)}$, and $\cup_x \mathcal{P}_x^{(r)}$ is invariant by $Z(\varphi^t)$.*

Proof: *Step 1: Bounded atlas along $L^{(r)}$.*

Some extension of the uniformly biLipschitz property is required, although the fibers of $L^{(r)}$ are not compact. We choose a C^q atlas $\{(\mathcal{E}_x, \tilde{\theta}_x)\}$ along L with the properties:

- The prolongations $\tilde{\theta}_x^{(r)} : \mathcal{E}_x^{(r)} \rightarrow L_x^{(r)}$ are such that $\cup_x \tilde{\theta}_x^{(r)}(\overline{B(1)})$ lies in a compact subset K of $\cup_x L_x^{(r)}$;
- $\{(\mathcal{E}_x^{(r)}, \tilde{\theta}_x^{(r)})\}$ is uniformly biLipschitz with respect to the Finsler metric d on $\mathcal{E}^{(r)}$ and some continuous metric d_K on K .

(As in section 2.2, such an atlas can be obtained from an appropriately chosen finite cover of M by foliated charts.) Note that the resulting C^{q-r} atlas along $L^{(r)}$ is $\mathrm{GL}^{(r)}(n)$ -equivariant. Let $\{\tilde{F}_x^t\}$ be the cocycle determined by $\{(\mathcal{E}_x, \tilde{\theta}_x)\}$. The prolonged cocycle satisfies (compare sections 2.2, 3.4)

$$\sup_{0 \leq \epsilon \leq 1} \ln^+ \|\tilde{F}^{(r)}(\pm\epsilon, x)\|_{q-r} \in L^1(M, \mu).$$

The set $\{g : Kg \cap K \neq \emptyset\}$ lies in a compact $C \subset \mathrm{GL}^{(r)}(n)$. Let m be a Lipschitz constant valid for the atlas and for all $g \in C$ restricted to K .

Step 2: Construction of $\tilde{\mu}_x^{(r),s}$ from bounded atlas.

Let $\tilde{\nu}_x^{(r),s}$ be the submanifolds given by theorem 2.1 applied to $\{\tilde{F}^{(r)}(t, x)\}$. Write $B = \overline{B(1)}$. Let $(u, g) \in \tilde{\nu}_x^{(r),s} \cap B$ with $\tilde{\theta}_x(u) = y \in L_x$, and

$$\tilde{\theta}_x^{(r)}(u, g) = \eta = \tilde{\theta}_y^{(r)}(\mathbf{0}, \mathrm{Id}) \cdot h \quad h \in C \subset \mathrm{GL}^{(r)}(n)$$

We wish to show that

$$(9) \quad \tilde{\theta}_x^{(r)}\left(\tilde{\nu}_x^{(r),s} \cap B\right) \cap \tilde{\theta}_y^{(r)}(B \cdot h) \subset \tilde{\theta}_y^{(r)}\left(\tilde{\nu}_y^{(r),s} \cdot h\right)$$

Let $\xi = \tilde{\theta}_x^{(r)}(u', g') = \tilde{\theta}_y^{(r)}(v, h')$ with $(u', g') \in \tilde{\nu}_x^{(r),s} \cap B$ and $(v, h'h^{-1}) \in B$. By the biLipschitz property of $\tilde{\theta}_{\varphi^t(y)}^{(r)}$,

$$(10) \quad d\left(\left(\tilde{F}_y^t(v), (J_v^{(r)} \tilde{F}_y^t) \cdot h'h^{-1} \cdot (J_{\mathbf{0}}^{(r)} \tilde{F}_y^t)^{-1}\right), (\mathbf{0}, \mathrm{Id})\right) \\ \leq md_K\left(\varphi^t \xi \cdot h^{-1} (J_{\mathbf{0}}^{(r)} \tilde{F}_y^t)^{-1}, J_{\mathbf{0}}^{(r)} \tilde{\theta}_{\varphi^t(y)}^{(r)}\right)$$

The following points are both in K (for sufficiently large $t > 0$):

$$\varphi^t \eta \cdot h^{-1} (J_{\mathbf{0}}^{(r)} \tilde{F}_y^t)^{-1} = J_{\mathbf{0}}^{(r)} \tilde{\theta}_{\varphi^t(y)}^{(r)} \quad ; \quad \varphi^t \eta \cdot (J_{\mathbf{0}}^{(r)} \tilde{F}_x^t)^{-1} = \tilde{\theta}_{\varphi^t x}^{(r)}\left(\tilde{F}^{(r)}(t, x)(u, g)\right)$$

Now right translate by $(J_{\mathbf{0}} \tilde{F}_y^t) \cdot h \cdot (J_{\mathbf{0}} \tilde{F}_x^t)^{-1} \in C$ to obtain the bound

$$(10) \leq m^2 d_K\left(\varphi^t \xi \cdot (J_{\mathbf{0}}^{(r)} \tilde{F}_x^t)^{-1}, \varphi^t \eta \cdot (J_{\mathbf{0}}^{(r)} \tilde{F}_x^t)^{-1}\right),$$

and apply $(\tilde{\theta}_{\varphi^t(x)}^{(r)})^{-1}$ to obtain

$$\leq m^3 d\left(\tilde{F}^{(r)}(t, x)(u', g'), \tilde{F}^{(r)}(t, x)(u, g)\right)$$

Thus finally

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \ln d\left(\tilde{F}^{(r)}(t, y)(v, h'h^{-1}), (\mathbf{0}, \mathrm{Id})\right) \leq \lambda^{(s)}$$

and $(v, h'h^{-1}) \in \tilde{\nu}_y^{(r),s}$. A similar argument shows

$$(11) \quad \tilde{\theta}_y^{(r)}\left(\tilde{\nu}_y^{(r),s} \cap B\right) \cdot h \cap \tilde{\theta}_x^{(r)}(B) \subset \tilde{\theta}_x^{(r)}\left(\tilde{\nu}_x^{(r),s}\right)$$

In conclusion, the manifolds $\tilde{\nu}_x^{(r),s} \cap B$ map forward under $\tilde{\theta}_x^{(r)}$ to submanifolds $\tilde{\mu}_x^{(r),s} \subset L_x^{(r)}$ which smoothly fit together with $\tilde{\mu}_y^{(r),s}$, for $y \in L_x$, after a vertical translation, according to (9) and (11).

Step 3: Tempered cocycle equivalence, construction of $\mu_x^{(r),s}$

The Oseledec-Pesin reduction theorem [2, Thm 6.10] gives a linear tempered equivalence $\{g_x\}$ of $\{\tilde{T}_x^t\}$ with a cocycle $\{T_x^t\}$ having constant Lyapunov

decomposition (as we assumed in section 3.3). By proposition 3.5, the Lyapunov decomposition of $\{T^{(r)}(t, x)\}$ is also constant, for $x \in M_T$, defined as in section 3.3. Recall that tempered means

$$\lim_{t \rightarrow \pm\infty} \frac{1}{t} \ln \|g_{\varphi^t(x)}^{\pm 1}\| = 0$$

Now set $\theta_x = \tilde{\theta}_x \circ g_x$. The new cocycles are $F_x^t = g_{\varphi^t(x)}^{-1} \circ \tilde{F}_x^t \circ g_x$ and

$$F^{(r)}(t, x)(u, h) = \left(g_{\varphi^t(x)}^{-1} \circ \tilde{F}^{(r)}(t, x)(g_x u, g_x h g_x^{-1}) \right) \cdot g_{\varphi^t(x)}$$

By proposition 3.1 and the tempered property,

$$\begin{aligned} \limsup_{t \rightarrow \infty} \frac{1}{t} \ln d \left(\tilde{F}^{(r)}(t, x)(g_x u, g_x h g_x^{-1}), (\mathbf{0}, \text{Id}) \right) &\leq \lambda^{(s)} \Leftrightarrow \\ \limsup_{t \rightarrow \infty} \frac{1}{t} \ln d \left(F^{(r)}(t, x)(u, h), (\mathbf{0}, \text{Id}) \right) &\leq \lambda^{(s)} \end{aligned}$$

Then

$$\nu_x^{(r),s} = (g_x^{-1}, \text{Ad } g_x^{-1}) \cdot \tilde{\nu}_x^{(r),s} \quad \text{and} \quad \theta_x^{(r)} \left(\nu_x^{(r),s} \right) = \tilde{\mu}_x^{(r),s} \cdot g_x$$

By proposition 3.9, the submanifolds $\mu_x^{(r),s} = \tilde{\mu}_x^{(r),s} \cdot g_x$ are C^{q-r} -smooth $\mathcal{X}^{(r)}$ -reductions of $L_x^{(r)}$ over $U_x = \theta_x(\mathcal{E}_x)$, for all x in a φ^t -invariant, μ -conull set $M_{\mathcal{R}}$. They smoothly fit together when appropriately vertically translated. Point (1) is proved.

Step 4: $\mathcal{H}^{(r)}$ saturation, construction of $\mathcal{P}_x^{(r)}$

Let $y \in \theta_x(\mathcal{E}_x)$. From above, there exists $g \in \text{GL}^{(r)}(n)$ such that $\mu_x^{(r),s} \cap \mu_y^{(r),s} \cdot g$ is open in each term. A point $\xi \in \mu_x^{(r),s}$ determines a frame of $\mu_x^{(r-1),s}$ at $\bar{\xi} = \pi_{r-1}^r(\xi)$ in which, as in remark 3.11, the Lyapunov filtration comprises the subspaces $V^{(r-1),\sigma}$ for $\sigma \leq \lambda^{(s)}$. On the other hand, the arguments of step 2 above show that the derivative of right translation by $\bar{g} = \rho_{r-1}^r(g)$ carries the Lyapunov filtration of $\mu_y^{(r-1),s}$ at $\bar{\xi}$ to that of $\mu_x^{(r-1),s}$ at $\bar{\xi} \cdot \bar{g}$. It follows (see formula (7) and its higher-order analogues) that $g \in \mathcal{H}^{(r)}$. For $x \in M_{\mathcal{R}}$, set $\mathcal{P}_x^{(r)} = \cup_{y \in L_x} \mu_y^{(r),s} \cdot \mathcal{H}^{(r)}$, a C^{q-r} reduction of $L_x^{(r)}$ to $\mathcal{H}^{(r)}$.

Invariance of $\cup_x \nu_x^{(r),s}$ by $F^{(r)}(t, x)$ gives

$$\varphi^t \mu_x^{(r),s} = \mu_{\varphi^t(x)}^{(r),s} \cdot (J_{\mathbf{0}}^{(r)} F_x^t)$$

By proposition 3.6, $J_{\mathbf{0}}^{(r)} F_x^t \in \mathcal{H}^{(r),0}$, so $\cup_x \mathcal{P}_x^{(r)}$ are φ^t -invariant. Any $\psi \in Z(\varphi^t)$ (see remark 2.10) enjoys a similar invariance of the $\mu_x^{(r),s}$, and by proposition 3.6, preserves $\cup_x \mathcal{P}_x^{(r)}$ where $x \in \cap_{i \in \mathbf{Z}} \psi^i(M_{\mathcal{R}} \cap M_G)$. \diamond

Our first theorem for contracted foliations concerns the atlas from theorem 3.13 (1): it gives rise to a homogeneous structure on leaves.

Theorem 3.15. *Assume $q \geq \lfloor \lambda^{(1)}/\lambda^{(s)} \rfloor + 1$. For all $x \in M_{\mathcal{R}}$, for all $y \in L_x$, there is a family $\mathcal{B}^0(y)$ of C^q charts on L_x , varying measurably between leaves and with the following additional properties:*

- (1) *Global diffeomorphisms: For all x in a φ^t -invariant, μ -conull set, each $\beta \in \mathcal{B}^0(y)$ is a diffeomorphism $\beta : (\mathbf{R}^n, \mathbf{0}) \rightarrow (L_x, y)$.*
- (2) *Homogeneity: Given $z \in L_x$, $\beta \in \mathcal{B}^0(y)$, and $\gamma \in \mathcal{B}^0(z)$, we have $(\gamma \circ \beta^{-1}) \cdot \mathcal{B}^0(y) = \mathcal{B}^0(z)$.*
- (3) *Invariance: Let $\mathcal{B}_x^0 = \cup_{y \in L_x} \mathcal{B}^0(y)$. The collection $\cup_x \mathcal{B}_x^0$ is $Z(\varphi^t)$ -invariant, and in these coordinates, $Z(\varphi^t)$ acts by resonance polynomials—that is, elements of $\mathcal{H}^{(r),0}$.*
- (4) *Structure group: For $\beta, \gamma \in \mathcal{B}_x^0$, the transition $\gamma^{-1} \circ \beta$ acts on \mathbf{R}^n by a translation composed with the an element of $\mathcal{H}^{(r)}$; in particular, \mathcal{B}_x^0 forms a C^q atlas on L_x .*

If the spectrum Σ^0 is 1/2-pinchd, then L carries an invariant family of C^{q-1} flat affine structures, measurable in x .

As usual, the $Z(\varphi^t)$ -invariance in (3) holds for a given centralizing ψ on an appropriate conull, ψ -invariant subset of $M_{\mathcal{R}}$.

Proof: Set $r = \lfloor \lambda^{(1)}/\lambda^{(s)} \rfloor$, and let $\cup_x \mathcal{P}_x^{(r)} = \cup_x \mu_x^{(r),s} \cdot \mathcal{H}^{(r)}$ be the reduction of $L^{(r)}$ to $\mathcal{H}^{(r)} \cong \mathcal{H}^{(r+1)}$ given by proposition 3.14 (2). Let $\{(\mathcal{E}_x, \theta_x)\}$ be the foliated atlas along L from the proof.

As in the proof of theorem 3.13 (1), define integrable horizontal distributions on $\mu_x^{(r),s}$ and extent to integrable, $Z(\varphi^t)$ -invariant, horizontal distributions on $\mathcal{P}_x^{(r)}$. Given $\xi \in \mathcal{P}_x^{(r)}$, denote \widehat{L}_ξ the integral leaf through ξ . For $\xi = \theta_y^{(r)}(\mathbf{0}, g)$, let $\beta_\xi = \theta_y \circ \alpha(\mathbf{0}, g)$, where $\alpha(\mathbf{0}, g)$ is as in the proof of theorem 3.13 (1). Note that the image of $J^{(r)}\beta_\xi$ is in \widehat{L}_ξ . Set

$$\mathcal{B}^0(y) = \{\beta_\xi : \xi \in \theta_y^{(r)}(\{\mathbf{0}\} \times \mathcal{H}^{(r),0})\}$$

For $x \in M_{\mathcal{R}}$, write $\hat{x} = J_{\mathbf{0}}^{(r)}\theta_x$. The C^{q-r} leaves $\widehat{L}_{\hat{x}} \subset L_x^{(r)}$ vary measurably in x , and therefore so does the maximum domain of definition of $\beta_{\hat{x}}$. Lusin's theorem gives $\epsilon > 0$ and a compact $C \subset M$, visited infinitely many times by $\{\varphi^t(x)\}$ for almost all x , and such that $\beta_{\hat{y}}$ is defined on $B(\epsilon)$ for all $y \in C$.

Let $J_{\mathbf{0}}^{(r)}F_x^t = h_t \in \mathcal{H}^{(r),0}$, which are diffeomorphisms by proposition 3.3. By remark 3.8, $h_k \rightarrow \mathbf{0}$ uniformly on compact sets as $k \rightarrow \infty$ in \mathbf{N} . Next,

$$\beta_{\hat{x}} = \varphi^{-k} \circ \beta_{\widehat{\varphi^k(x)}} \circ h_k$$

Let $x \in M_{\mathcal{R}}$ and $k_m \rightarrow \infty$ be such that $\varphi^{k_m}(x) \in C$ for all $m \geq 1$. The domain of $\beta_{\hat{x}}$ contains $\cup_m h_{k_m}^{-1}(B(\epsilon)) = \mathbf{R}^n$. Given $y \in L_x$, there is m such that $\varphi^{k_m}(y) \in \text{im } \beta_{\widehat{\varphi^{k_m}(x)}}$. Thus $y \in \text{im } \beta_{\hat{x}}$. Now $\beta_{\hat{x}}$ is a diffeomorphism as

claimed in (1). As in the proof of theorem 3.13 (1), $\beta_{\xi \cdot h} = \beta_{\xi} \circ h$, so β_{ξ} is a diffeomorphism for all ξ in $\mathcal{P}_x^{(r)}$, for all $x \in M_{\mathcal{R}}$.

Let $\beta = \beta_{\xi} \in \mathcal{B}_x^0(y)$, $\gamma = \beta_{\omega} \in \mathcal{B}_x^0(z)$. Let $\beta_{\xi}(u) = z$, so $J_u^{(r)}\beta_{\xi} = \eta \in \widehat{L}_{\xi}(z)$. Now compare $J^{(r)}(\beta_{\xi} \circ \tau_u)$ and $J^{(r)}\beta_{\eta}$: both map the origin to η and have image in $\widehat{L}_{\xi} \subset \mathcal{P}_x^{(r)}$. Then $\beta_{\xi} \circ \tau_u = \beta_{\eta}$. Next, $\eta = \omega \cdot h$ for a unique $h \in \mathcal{H}^{(r)}$. Thus $\gamma^{-1}\beta = \beta_{\omega}^{-1}\beta_{\xi} = h \circ \tau_{-u}$. Now (4) is proved. The same argument also shows that $\mathcal{B}^0(y)$ can be defined for all $y \in L_x$, assuming $x \in M_{\mathcal{R}}$. This completes the proof of (1).

Given β, γ as in point (2), one can write

$$\mathcal{B}_x^0(y) = \beta \cdot \mathcal{H}^{(r),0} \quad \mathcal{B}_x^0(z) = \gamma \cdot \mathcal{H}^{(r),0}$$

and the conclusion follows immediately.

Point (3) is just as in the proof of theorem 3.13 (1).

If the spectrum is 1/2 pinched, then $r = 1$, and $\mathcal{H}^{(1)}$ comprises linear transformations, so the atlas \mathcal{B}_x^0 is a flat affine structure on L_x . \diamond

Theorem 3.16. *The foliation $\cup_x L_x$ contains a $Z(\varphi^t)$ -invariant filtered family of C^{q-1} subfoliations*

$$L^{i_1} \subset \dots \subset L^{i_l} = L \quad l \leq r,$$

a subfamily of the filtered family in theorem 2.9, equipped with flat connections ∇_x^j on the normal bundles of $L^{i_{j-1}} \subset L^{i_j}$ for all $1 \leq j \leq l$. The $\oplus_x \nabla_x^j$ are $Z(\varphi^t)$ -invariant, measurable in x , and C^{q-2} inside almost every L_x ; in particular, almost every leaf $L_x^{i_1}$ carries a C^{q-2} invariant flat connection.

Proof: Extract the components of the Lyapunov filtration corresponding to the conglomerated distributions:

$$V^{i_j} = \sum_{m \leq j} \widetilde{W}^m = \sum_{m \leq i_j} W^m$$

where $i_1, \dots, i_{r_*} = i_l$ and the subspaces \widetilde{W}^m are as in the proof of theorem 3.13 (2). The $Z(F_x^t)$ -invariant distributions \mathcal{V}_x^j , $1 \leq j \leq l$, on \mathcal{E}_x are tangent at $\mathbf{0}$ to V^{i_j} , and thus to the submanifolds $\mathcal{V}_x^{i_j}$ given by theorem 2.1. Then for almost all $x \in M$, the images $(\theta_x)_* \mathcal{V}_x^j$ are tangent to $L_x^{i_j}$, where L^{i_j} is the foliation given by theorem 2.9.

The $Z(F_x^t)$ -invariant flat connections ∇_x^j on $\mathcal{V}_x^j/\mathcal{V}_x^{j-1}$ given by theorem 3.13 (2) push forward under θ_x to flat connections on $T_x L^{i_j}/T_x L^{i_{j-1}}$, which are $Z(\varphi^t)$ -invariant and C^{q-2} inside \mathcal{E}_x . \diamond

REFERENCES

1. V. Araujo, A. Bufetov, and S. Filip, *On Hölder-continuity of Oseledets subspaces*, J. London Math. Soc. (2) **93** (2016), no. 1, 194–218.

2. L. Barreira and Y. Pesin, *Introduction to smooth ergodic theory*, Amer. Math. Soc., 2013.
3. R. Feres, *Rigid geometric structures and actions of semisimple Lie groups*, Rigidité, Groupe Fondamentale, et Dynamique, vol. 13, Société Mathématique de France, 2002, pp. 121–166.
4. ———, *A differential-geometric view of normal forms of contractions*, Modern dynamical systems and applications, Cambridge University Press, Cambridge, UK, 2004, pp. 103–121.
5. D. Fisher and G. Margulis, *Almost isometric actions, property (T), and local rigidity*, Invent. Math. **162** (2005), 19–80.
6. ———, *Local rigidity of affine actions of higher rank groups and lattices*, Ann. of Math. (2) **170** (2009), no. 1, 67–122.
7. M. Guysinsky, *The theory of non-stationary normal forms*, Erg. Thy. Dyn. Sys. **21** (2001), 845–862.
8. M. Guysinsky and A. Katok, *Normal forms and invariant geometric structures for dynamical systems with invariant contracting foliations*, Math. Res. Lett. **5** (1998), 149–163.
9. B. Kalinin and A. Katok, *Invariant measures for actions of higher rank abelian groups*, Smooth ergodic theory and its applications (Seattle, WA, 1999), Proc. Sympos. Pure Math., vol. 69, Amer. Math. Soc., 2001, pp. 593–637.
10. B. Kalinin and V. Sadovskaya, *Normal forms for non-uniform contractions*, www.personal.psu.edu/~bvk102/Papers/papersindex.html.
11. ———, *Global rigidity for totally nonsymplectic Anosov \mathbf{z}^k actions*, Geom. Topol. **10** (2006), 929–954.
12. ———, *Non-stationary normal forms on contracting foliations: smoothness and homogeneity*, Geom. Ded. **183** (2016), 181–194.
13. A. Katok and R. Spatzier, *Differential rigidity of Anosov actions of higher rank abelian groups and algebraic lattice actions*, Tr. Math. Inst. Steklova **216** (1997).
14. S. Kobayashi, *Transformation groups in differential geometry*, Springer, Berlin, 1995.
15. W. Li and K. Lu, *Sternberg theorems for random dynamical systems*, Comm. Pure Appl. Math. **58** (2005), 941–988.
16. D. Ruelle, *Ergodic theory of differentiable dynamical systems*, Publ. Math. IHÉS **50** (1978), 27–58.
17. S. Sternberg, *Local contractions and a theorem of Poincaré*, Amer. J. Math. **79** (1957), no. 4, 809–824.

Karin Melnick
 Department of Mathematics
 4176 Campus Drive
 University of Maryland
 College Park, MD 20742
 USA
karin@math.umd.edu