


Max-Planck-Institut für Mathematik Bonn

Representation of unity by binary forms

by

Shabnam Akhtari



Representation of unity by binary forms

Shabnam Akhtari

Max-Planck-Institut für Mathematik
Vivatsgasse 7
53111 Bonn
Germany

REPRESENTATION OF UNITY BY BINARY FORMS

SHABNAM AKHTARI

ABSTRACT. In this paper, it is shown that if $F(x, y)$ is an irreducible binary form with integral coefficients and degree $n \geq 3$, then provided that the absolute value of the discriminant of F is large enough, the equation $F(x, y) = \pm 1$ has at most $11n - 2$ solutions in integers x and y . We will also establish some sharper bounds when more restrictions are assumed. These upper bounds are derived by combining methods from classical analysis and geometry of numbers. The theory of linear forms in logarithms plays an essential role in studying the geometry of our Diophantine equations.

1. INTRODUCTION

Let $F(x, y) = a_n x^n + a_{n-1} x^{n-1} y + \dots + a_0 y^n$ be an irreducible binary form with rational integer coefficients and $n \geq 3$. We will study N_n , the number of solutions to the equation

$$(1) \quad F(x, y) = \pm 1,$$

in integers x and y . We will regard (x, y) and $(-x, -y)$ as one solution. So we may only count the solutions with $y \geq 0$. But how large can N_n be? Let p be a prime and consider the following irreducible form

$$F_1(x, y) = x^n + p(x - y)(2x - y) \dots (nx - y).$$

It is easy to see that $F_1(x, y) = 1$ has the following n solutions

$$(1, 1), (1, 2), \dots, (1, n).$$

Thus a linear upper bound of the shape cn is best possible except for the determination of c . We will show that

Theorem 1.1. *Let $F(x, y)$ be an irreducible binary form with integral coefficients and degree $n \geq 3$. Then the Diophantine equation $|F(x, y)| = 1$ has at most $11n - 2$ solutions in integers x and y , provided that the absolute value of the discriminant of F is greater than D_0 , where $D_0 = D_0(n)$ is an effectively computable constant. Moreover, assume that the polynomial $F(x, 1)$ has r real roots and $2s$ non-real roots ($r + 2s = n$). Then $|F(x, y)| = 1$ has at most $11r + 4s - 1$ solutions in integers x and y .*

We remark here that D_0 can be computed effectively in terms of n , the degree of F . Indeed, we may take $D_0 = 2^{22}(n + 1)^{10}n^n$. Theorem 4.6 gives an algorithm to compute D_0 .

In the above theorem, we supposed that F is irreducible. We will see in Section 2, that when F is reducible, the situation is simpler. Let D be the discriminant of form

2000 *Mathematics Subject Classification.* 11D45.

Key words and phrases. Thue Equations, Linear Forms in Logarithms.

F (it is defined in Section 3). Note that the condition $|D| > D_0(n)$ is a restriction, because we know for binary form $F \in \mathbb{Z}[x, y]$ of degree n and discriminant $D \neq 0$, we have the following sharp bound (see [13]):

$$(2) \quad n \leq 3 + 2(\log |D|) / \log 3.$$

In Section 3, we will see how Theorem 1.1 gives an upper bound for the number of integral solutions to $F(x, y) = \pm 1$ when F has a small discriminant.

One may conjecture that the number of solutions may be estimated in terms of r the number of real solutions of $F(x, 1) = 0$. This is not the case. For example, let n be an even integer and p a prime number. If we put

$$F(x, y) = x^n + p(x - y)^2(2x - y)^2 \dots \left(\frac{n}{2}x - y\right)^2$$

then $F(x, y)$ is irreducible and $F(x, 1) = 0$ has no real root. However, $F(x, y) = 1$ has the following solutions:

$$(1, 1), (1, 2), \dots, \left(1, \frac{n}{2}\right).$$

In Proposition 5.1, we will show that the number of solutions (x, y) with large enough y can be estimated in terms of r .

In 1909, Thue [23] derived the first general sharpening of Liouville's theorem on rational approximation to algebraic numbers, proving, if θ is algebraic of degree $n \geq 3$ and $\epsilon > 0$, that there exists a constant $c(\theta, \epsilon)$ such that

$$\left| \theta - \frac{p}{q} \right| > \frac{c(\theta, \epsilon)}{q^{\frac{n}{2} + 1 + \epsilon}}$$

for all $p \in \mathbb{Z}$ and $q \in \mathbb{N}$. It follows almost immediately, if $F(x, y)$ is an irreducible binary form in $\mathbb{Z}[x, y]$ of degree at least three and h a nonzero integer, that the equation

$$(3) \quad F(x, y) = h$$

has only finitely many solutions in integers x and y . Equation (3) is called a Thue equation.

For any nonzero integer h let $\omega(h)$ denote the number of distinct prime factors of h . In 1933, Mahler [17] proved that equation (3) has at most $C_1^{1+\omega(h)}$ solutions in co-prime integers x and y , where C_1 is a positive number that depends on F only. In 1987, Bombieri and Schmidt [5] showed that the number of solutions of $F(x, y) = h$ in co-prime integers x and y is at most

$$C_2 n^{1+\omega(h)},$$

where C_2 is an absolute constant. Further they showed that C_2 may be taken 215 if n is sufficiently large. Note that this upper bound is independent of the coefficients of the form F ; a result of this flavour was first deduced in 1983 by Evertse [9]. In the introduction of [5], Bombieri and Schmidt comment that their argument can be used to prove a more general result. For example, if N_n is the corresponding bound in the special case $h = 1$, one obtains $N_n n^{\omega(h)}$ as a bound in the general case. For this reason we will focus on the equation $|F(x, y)| = 1$.

The effective solution of an arbitrary Thue equation has its origin in Baker's [3] theorem that says that if $\kappa > n + 1$, then every integer solution (x, y) of equation (3) satisfies

$$\max\{|x|, |y|\} < C_3 \exp \log^\kappa |h|$$

where C_3 is an effectively computable constant depending only on n , κ and the coefficients of F .

Evertse and Győry (see [9] and [11]) have studied the Thue inequality

$$(4) \quad 0 < |F(x, y)| \leq h.$$

Define, for $3 \leq n < 400$

$$(N(n), \delta(n)) = \left(6n7^{\binom{n}{3}}, \frac{5}{6}n(n-1) \right)$$

and for $n > 400$

$$(N(n), \delta(n)) = (6n, 120(n-1)).$$

They prove that if

$$|D| > h^{\delta n} \exp(80n(n-1)),$$

then the number of solutions to (4) in co-prime integers x and y is at most $N(n)$.

Győry [14] also shows, for binary form F of degree $n \geq 3$, that if $0 < a < 1$ and

$$|D| \geq n^n (3.5^n h^2)^{(2(n-1)/(1-a))},$$

then the number of solutions to (4) in co-prime integers x and y is at most $25n + (n+2)\left(\frac{2}{a} + \frac{1}{4}\right)$, and if F is reducible then at most $5n + (n+2)\left(\frac{2}{a} + \frac{1}{4}\right)$.

A great reference in this field is a work of Stewart [22]. We will follow many arguments from [22] here. A consequence of Stewart's main theorem in [22] is that if the discriminant D of F is non-zero and

$$|D|^{1/n(n-1)} \geq |h|_{\frac{2}{n+\epsilon}},$$

then the number of pairs of co-prime integers (x, y) for which $F(x, y) = h$ holds is at most

$$1400 \left(1 + \frac{1}{8\epsilon n} \right) n.$$

Bennett [4] and Okazaki [20] have obtained very good upper bounds for the number of solutions to cubic Thue equations. Some upper bounds are given for the number of integral solutions to quartic Thue equations in [1] and [2]. Throughout this paper we may assume n , the degree of our binary form, is greater than 4.

We will use methods from [22] to give upper bounds on the number of "small" solutions to (1). Then, in Section 6, we will generalize some ideas from [20, 2] to associate a transcendental curve $\phi(x, y)$ to the binary form $F(x, y)$. Introducing this curve will give us the opportunity to bring the theory of linear forms in logarithms in.

2. REDUCIBLE FORMS

Let us take a brief interlude from the principal matter at hand to discuss the much simpler situation where the form $F(x, y)$ is reducible over $\mathbb{Z}[x, y]$. In general, equation (1) may have infinitely many integral solutions; $F(x, y)$ could, for instance, be a power of a linear or indefinite binary quadratic form that represents unity. If $F(x, y)$ is a reducible form, however, we may very easily derive a stronger version of our main theorem under the assumption that $F(x, 1)$ has at least two distinct zeros.

Suppose that $F(x, y)$ is reducible and can be factored over $\mathbb{Z}[x, y]$ as follows

$$F(x, y) = F_1(x, y)F_2(x, y),$$

with $\deg(F_1) \leq \deg(F_2)$ and F_1 irreducible over $\mathbb{Z}[x, y]$. Therefore, the following equations must be satisfied:

$$(5) \quad F_1(x, y) = \pm 1$$

and

$$(6) \quad F_2(x, y) = \pm 1.$$

This means the number of solutions to (1) is no more than the minimum of number of solutions to (5) and (6).

First suppose that F_1 is a linear form. Then the equation (6) can be written as a polynomial of degree at most $n - 1$ in x and therefore there are no more than $2(n - 1)$ complex solutions to above equations.

Now let us suppose that F_1 is a quadratic form. Using Bézout's theorem from classical algebraic geometry concerning the number of common points of two plane algebraic curves, we conclude that (1) has at most $4(n - 2)$ integral solutions.

If $\deg(F_1) \geq 3$ then Theorem 1.1 will give us an upper bound for the number of integral solutions to (5), and therefore to (1).

3. EQUIVALENT FORMS

Our approach depends on the fact that if we transform F by the action of an element of $GL_2(\mathbb{Z})$ the problem of counting solutions remains unchanged, while the Diophantine approximation properties of F can change very drastically. Let

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

and define the binary form F_A by

$$F_A(x, y) = F(ax + by, cx + dy).$$

If the determinant of matrix A is equal to ± 1 then we say that F_A and $-F_A$ are equivalent to F .

Suppose that $A \in GL_2(\mathbb{Z})$ and (x, y) is a solution of (1) in integers x and y . Then

$$A \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} ax + by \\ cx + dy \end{pmatrix}$$

and $(ax + by, cx + dy)$ is a solution of $F_{A^{-1}}(x, y) = \pm 1$ in integers x and y .

Let F be a binary form that factors in \mathbb{C} as

$$\prod_{i=1}^n (\alpha_i x - \beta_i y).$$

The discriminant D_F of F is given by

$$D_F = \prod_{i < j} (\alpha_i \beta_j - \alpha_j \beta_i)^2.$$

Observe that for any 2×2 matrix A with integer entries

$$(7) \quad D_{F_A} = (\det A)^{n(n-1)} D_F.$$

We denote by N_F the number of solutions in integers x and y of the Diophantine equation (1). If F_1 and F_2 are equivalent then $N_{F_1} = N_{F_2}$ and $D_{F_1} = D_{F_2}$.

Let p be a prime number and put

$$A_0 = \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}, \quad A_j = \begin{pmatrix} 0 & -1 \\ p & j \end{pmatrix}$$

for $j = 1, \dots, p$. Then we have

$$\mathbb{Z}^2 = \cup_{j=0}^p A_j \mathbb{Z}^2.$$

Therefore the number of solutions of (1) is at most $N_{F_0} + N_{F_1} + \dots + N_{F_p}$, where

$$F_j(x, y) = F_{A_j}(x, y).$$

Note that by (7),

$$\left| D_{F_{A_j}} \right| \geq p^{n(n-1)}.$$

Therefore, if N is an upper bound for the number of solutions to (1) for binary forms F with $|D_F| \geq p^{n(n-1)}$ then $(p+1)N$ will be an upper bound for the number of solutions to $|F(x, y)| = 1$ when F has a nonzero discriminant.

Assume that $F(x, y) = \pm 1$ has a solution (x_0, y_0) . Then there is a matrix A in $GL_2(\mathbb{Z})$ for which $A^{-1}(x_0, y_0)$ is $(1, 0)$. Therefore, $(1, 0)$ is a solution to

$$F_A(x, y) = \pm 1.$$

We conclude that either F_A or $-F_A$ is a monic form. From now on we will assume that the binary form $F(x, y)$ in Theorem 1.1 is monic.

4. HEIGHTS

In this section we give a brief review of the theory of height functions of polynomials and binary forms.

For the polynomial $G(x) = c(x - \beta_1) \dots (x - \beta_n)$ with $c \neq 0$, the Mahler measure $M(G)$ is defined by

$$M(G) = |c| \prod_{i=1}^n \max(1, |\beta_i|).$$

Mahler [16] showed, for polynomial G of degree n and discriminant D , that

$$(8) \quad M(G) \geq \left(\frac{D}{n^n} \right)^{\frac{1}{2n-2}}.$$

The Mahler measure of an algebraic number α is defined as the Mahler measure of the minimal polynomial of α over \mathbb{Q} .

For an algebraic number α , the (naive) height of α , denoted by $H(\alpha)$, is defined by the following identities.

$$H(\alpha) = H(f(x)) = \max(|a_n|, |a_{n-1}|, \dots, |a_0|)$$

where $f(x) = a_n x^n + \dots + a_1 x + a_0$ is the minimal polynomial of α over \mathbb{Z} .

We have

$$(9) \quad \binom{n}{\lfloor n/2 \rfloor}^{-1} H(\alpha) \leq M(\alpha) \leq (n+1)^{1/2} H(\alpha).$$

We will use transformations in $GL_2(\mathbb{Z})$ to dispense with a technical hypothesis about the height of F . We call the polynomials $f(x)$ and $f^*(x) \in \mathbb{Z}$ strongly equivalent if $f^*(x) = f(x+a)$ for some $a \in \mathbb{Z}$. Two algebraic integers α and α' are called strongly equivalent if their minimal polynomials are strongly equivalent.

Proposition 4.1. (*Győry [13]*) *Suppose that $f(x)$ is a monic polynomial in $\mathbb{Z}[x]$ with degree $n \geq 2$ and non-zero discriminant D . There is a polynomial $f^*(x) \in \mathbb{Z}$ strongly equivalent to $f(x)$ so that*

$$H(f^*(x)) < \exp\{n^{4n^{12}}|D|^{6n^8}\} < \exp \exp\{4(\log |D|)^{13}\}.$$

For polynomial $f(x) = a_n x^n + \dots + a_1 x + a_0$ with degree n and integer coefficients, put

$$L(f) = |a_n| + \dots + |a_1| + |a_0|.$$

Mahler [15] showed that

$$(10) \quad 2^{-n} L(f) \leq M(f) \leq L(f).$$

Define the absolute logarithmic height of an algebraic number as follows. Let α_1 be a root of $F(x, 1) = 0$ and $\mathbb{Q}(\alpha)^\sigma$ the embeddings of $\mathbb{Q}(\alpha)$ in \mathbb{C} , $1 \leq \sigma \leq n$. For $\rho \in \mathbb{Q}(\alpha)$, we respectively have n Archimedean valuations of $\mathbb{Q}(\alpha)$:

$$|\rho|_\sigma = \left| \rho^{(\sigma)} \right|, \quad 1 \leq \sigma \leq n.$$

We enumerate simple ideals of $\mathbb{Q}(\alpha)$ by indices $\sigma > n$ and define non-Archimedean valuations of $\mathbb{Q}(\alpha)$ by the formulas

$$|\rho|_\sigma = (\text{Norm } \mathfrak{p})^{-k},$$

where

$$k = \text{ord}_{\mathfrak{p}}(\alpha), \quad \mathfrak{p} = \mathfrak{p}_\sigma, \quad \sigma > n,$$

for any $\rho \in \mathbb{Q}(\alpha)^*$. Then we have the *product formula* :

$$\prod_1^\infty |\rho|_\sigma = 1, \quad \rho \in \mathbb{Q}(\alpha)^*.$$

Note that $|\rho|_\sigma \neq 1$ for only finitely many ρ . We should also remark that if $\sigma_2 = \bar{\sigma}_1$, i.e.,

$$\sigma_2(x) = \bar{\sigma}_1(x) \quad \text{for} \quad x \in \mathbb{Q}(\alpha),$$

then the valuations $|\cdot|_{\sigma_1}$ and $|\cdot|_{\sigma_2}$ are identical. We define the *absolute logarithmic height* of α as

$$h(\alpha) = \frac{1}{2n} \sum_{\sigma=1}^\infty |\log |\alpha|_\sigma|.$$

This height is called absolute because it is independent of the field in which the number α lies.

The following Lemmata about the height of algebraic numbers will be helpful later.

Lemma 4.2. *For every non-zero algebraic number α , we have $h(\alpha^{-1}) = h(\alpha)$. For algebraic numbers $\alpha_1, \dots, \alpha_n$, we have*

$$h(\alpha_1 \dots \alpha_n) \leq h(\alpha_1) + \dots + h(\alpha_n)$$

and

$$h(\alpha_1 + \dots + \alpha_n) \leq \log n + h(\alpha_1) + \dots + h(\alpha_n).$$

Proof. See [7] for a proof. □

Lemma 4.3. (Voutier [24]) *Suppose α is a non-zero algebraic number of degree n which is not a root of unity. If $n \geq 2$ then*

$$h(\alpha) = \frac{1}{n} \log M(\alpha) > \frac{1}{4n} \left(\frac{\log \log n}{\log n} \right)^3.$$

Lemma 4.4. (Mahler [16]) *If a and b are distinct zeros of polynomial $P(x)$ with degree n , then we have*

$$|a - b| \geq \sqrt{3}(n+1)^{-n} M(P)^{-n+1},$$

where $M(P)$ is the Mahler measure of P .

In the following lemma we approximate the size of $f'(\alpha)$ in terms of the discriminant and heights of f , where f' is the derivative of the polynomial f and α is a root of $f = 0$.

Lemma 4.5. *Let $f(x) = a_n x^n + \dots + a_1 x + a_0$ be an irreducible polynomial of degree n and with integral coefficients. Suppose that α_m is a root of $f(x) = 0$. For $f'(x)$ the derivative of f , we have*

$$2^{-(n-1)^2} \frac{|D_f|}{M(f)^{2n-2}} \leq |f'(\alpha_m)| \leq \frac{n(n+1)}{2} H(f) (\max(1, |\alpha_m|))^{n-1},$$

where D_f is the discriminant, $M(f)$ is the Mahler measure and $H(f)$ is the naive height of f .

Proof. The right hand side inequality is trivial by noticing that

$$f'(x) = n a_n x^{n-1} + \dots + a_1 x.$$

To see the left hand side inequality, observe that for α_i, α_j , two distinct roots of $f(x)$, we have

$$|\alpha_i - \alpha_j| \leq 2 \max(1, |\alpha_i|) \max(1, |\alpha_j|).$$

Then

$$\begin{aligned} |f'(\alpha_m)| &= \prod_{i=1, i \neq m}^n |\alpha_i - \alpha_m| \geq \prod_{i=1, i \neq m}^n \frac{|\alpha_i - \alpha_m|}{\max(1, |\alpha_i|) \max(1, |\alpha_m|)} \\ &\geq 2^{n-1-n(n-1)} \prod_{j=1}^n \prod_{i=1, i \neq j}^n \frac{|\alpha_i - \alpha_j|}{\max(1, |\alpha_i|) \max(1, |\alpha_j|)} \\ &= 2^{-(n-1)^2} \frac{|D_F|}{M(F)^{2n-2}}. \end{aligned}$$

□

Suppose that \mathbb{K} is an algebraic number field of degree d over \mathbb{Q} embedded in \mathbb{C} . If $\mathbb{K} \subset \mathbb{R}$, we put $\chi = 1$, and otherwise $\chi = 2$. We are given numbers $\gamma_1, \dots, \gamma_n \in \mathbb{K}^*$ with absolute logarithmic heights $h(\gamma_j)$, $1 \leq j \leq n$. Let $\log \gamma_1, \dots, \log \gamma_n$ be arbitrary fixed non-zero values of the logarithms. Suppose that

$$A_j \geq \max\{dh(\gamma_j), |\log \gamma_j|\}, \quad 1 \leq j \leq n.$$

Now consider the linear form

$$\mathfrak{L} = b_1 \log \gamma_1 + \dots + b_n \log \gamma_n,$$

with $b_1, \dots, b_n \in \mathbb{Z}$ and with the parameter

$$B = \max\{1, \max\{b_j A_j / A_n : 1 \leq j \leq n\}\}.$$

For brevity we put

$$\begin{aligned} \Omega &= A_1 \dots A_n, \\ C(n) &= C(n, \chi) = \frac{16}{n! \chi} e^n (2n + 1 + 2\chi)(n + 2)(4n + 4)^{n+1} \left(\frac{1}{2} en\right)^\chi, \\ C_0 &= \log(e^{4.4n+7} n^{5.5} d^2 \log(en)), \\ W_0 &= \log(1.5eBd \log(ed)). \end{aligned}$$

The following is the main result of [19].

Proposition 4.6 (Matveev [19]). *If $\log \gamma_1, \dots, \log \gamma_n$ are linearly independent over \mathbb{Z} and $b_n \neq 0$, then*

$$\log |\mathfrak{L}| > -C(n)C_0W_0d^2\Omega.$$

5. STEPS OF THE PROOF OF THEOREM 1.1

Suppose that (x, y) is an integral solution to (1). We will assume that F is monic, as we may. Then we have

$$(x - \alpha_1 y)(x - \alpha_2 y) \dots (x - \alpha_n y) = \pm 1.$$

Therefore, for some $\alpha \in \{\alpha_1, \alpha_2, \dots, \alpha_n\}$,

$$|x - \alpha y| \leq 1.$$

Definition. We say the pair of solution (x, y) is related to α if

$$\alpha \in \{\alpha_1, \alpha_2, \dots, \alpha_n\}$$

and

$$|x - \alpha y| = \min_{1 \leq j \leq n} |x - \alpha_j y|.$$

Let $F(x, y)$ be a binary form of degree $n \geq 5$, discriminant D , with $|D| > D_0$ and Mahler measure $M(F)$, where D_0 is an effectively computable constant depending only on n (see the statement of Theorem 1.1). We will assume that all coefficients of F are integer and $F(x, 1) = 0$ has r real roots and $2s$ non-real roots ($r + 2s = n$). Here we describe briefly the steps of our proof to the main result of this manuscript, Theorem 1.1.

In the following steps, we fix a root of $F(x, 1) = 0$ and estimate the number of solutions related to that root from above. Let α be a complex root of $F(x, 1) = 0$ and $\bar{\alpha}$ be its complex conjugate. For integers x and y we have

$$|x - \alpha y| = |x - \bar{\alpha} y|.$$

Hence, a solution (x, y) of (1) is related to α if and only if it is related to $\bar{\alpha}$. It is, therefore, sufficient to count the number of solutions related to one of α and $\bar{\alpha}$.

Proposition 5.1. *For binary form $F(x, y)$ with integer coefficients and degree n , let α be a non-real root of $F(x, 1) = 0$. If a pair of integer (x, y) satisfies $F(x, y) = \pm 1$ and is related to α then*

$$(11) \quad |y| \leq \frac{(n+1)2^{\frac{(n-1)^2}{n}}}{(\sqrt{3}|D|)^{1/n}} M(F)^{3-3/n},$$

Proof. Let $\alpha = \tau + it$, with $t \neq 0$, be a non-real root of $F(x, 1) = 0$. If a solution (x, y) of (1) is related to α then $\bar{\alpha}$, the complex conjugate of α is also a root of $F(x, 1) = 0$ and we have

$$\left| \frac{x}{y} - \alpha \right| = \frac{\left| \frac{x}{y} - \alpha \right| + \left| \frac{x}{y} - \bar{\alpha} \right|}{2} \geq \frac{|\alpha - \bar{\alpha}|}{2}.$$

Moreover, if $\beta \neq \alpha$ is a root of $F(x, 1) = 0$ then

$$\left| \frac{x}{y} - \beta \right| \geq \frac{\left| \frac{x}{y} - \alpha \right| + \left| \frac{x}{y} - \beta \right|}{2} \geq \frac{|\beta - \alpha|}{2}.$$

Thus

$$\begin{aligned} \frac{1}{|y|^n} &= \left| \frac{x}{y} - \alpha \right| \prod_{\alpha_i \neq \alpha} \left| \frac{x}{y} - \alpha_i \right| \\ &\geq \frac{|\alpha - \bar{\alpha}|}{2} \prod_{\alpha_i \neq \alpha} \frac{|\alpha - \alpha_i|}{2} \\ &= |\alpha - \bar{\alpha}| |f'(\alpha)| 2^{-n}. \end{aligned}$$

By Lemma 4.4,

$$|\alpha - \bar{\alpha}| \geq \sqrt{3}(n+1)^{-n} M(F)^{-n+1}.$$

This, together with Lemma 4.5, shows that

$$\frac{1}{|y|^n} \geq \sqrt{3}(n+1)^{-n} 2^{-(n-1)^2} \frac{|D|}{M(f)^{3n-3}}.$$

This completes our proof. \square

Repeating an argument of Stewart [22] and using our assumption that absolute value of the discriminant of F is large in terms of its degree, in Section 7 we will show that there are at most $5(r+s)$ solutions (x, y) with $0 < y \leq M(F)^2$.

Lemma 7.5 and 7.6 give an upper bound $2r+s$ for the number of solutions (x, y) with $M(F)^2 < y < M(F)^{1+(n-1)^2}$. To prove Lemma 7.5 we will appeal to a classical inequality of Lewis and Mahler (see Lemma 7.4).

For a non-real root α of $F(x, 1) = 0$, Proposition 5.1 says that we only need to count the solutions (x, y) related to α with

$$|y| \leq \frac{(n+1)2^{(n-1)^2/n}}{\sqrt{3}|D|^{1/n}} M(F)^{3-3/n}.$$

The solutions with larger y must be related to a real root of $F(x, 1) = 0$.

Our approach to count the number of possibly remaining solutions differs from the approach of Bombieri-Schmidt [5] and Stewart [22]. In Section 6, we will define a logarithmic map $\phi(x, y)$. Some geometric properties of this curve lead us to obtain an exponential gap principle in Section 9. This new type of gap principle, together with Baker theory of linear forms in logarithms (see Proposition 4.6), will be used in Section 10 to establish an upper bound $2r$ for the number of solutions (x, y) with $y \geq M(F)^{1+(n-1)^2}$.

For some technical reasons, particularly to estimate quantities in Proposition 4.6 while counting the number of solutions (x, y) with $y \geq M(F)^{1+(n-1)^2}$, we will need

to exclude a set of solutions from our search. This set is called \mathfrak{A} and is defined in section 7. The set \mathfrak{A} contains $2r + 2s - 2$ “small” solutions.

Hence, under the assumption of Theorem 1.1, there can not exist more than $11r + 4s - 2$ to equation (1).

6. THE LOGARITHMIC CURVE $\phi(x, y)$

In order to count the number of “large” solutions to $F(x, y) = 1$, many mathematicians including Bombieri and Schmidt [5] and Stewart [22] followed and refined a general method inaugurated by Siegel and Mahler. The general line of attack to the problem of counting “large” solutions deals rather efficiently with solutions x, y to $F(x, y) = 1$, provided that $\max(|x|, |y|)$ is larger than a certain power of the height of F . We will, in contrast, associate a transcendental curve $\phi(x, y)$ to the binary form $F(x, y)$. However, the reason in success of both our method and the more classical method of Siegel and Mahler lies in the fact that $\frac{x}{y}$ is a good approximation to a root of the equation $F(x, 1) = 0$ when either x or y is large enough.

Let D be the discriminant of the binary form $F(x, y)$ and $f(x) = F(x, 1)$. Define, for $m \in \{1, 2, \dots, n\}$,

$$(12) \quad \phi_m(x, y) = \log \left| \frac{D^{\frac{1}{n(n-2)}} (x - y\alpha_m)}{(f'(\alpha_m))^{\frac{1}{n-2}}} \right|$$

and

$$(13) \quad \phi(x, y) = (\phi_1(x, y), \phi_2(x, y), \dots, \phi_n(x, y)).$$

We will estimate the size of $f'(\alpha_m)$ from below in order to give an upper bound on the size of $\phi(x, y)$.

Lemma 6.1. *Suppose that F is a monic binary form satisfying the conditions in Theorem 1.1. Then $(1, 0)$ is a solution to the equation $|F(x, y)| = 1$ and*

$$\|\phi(1, 0)\| \leq n \log \left(|D|^{\frac{1}{n(n-2)}} M(F)^{\frac{2n-2}{n-2}} \right),$$

Proof. By the definition of ϕ in (13),

$$\|\phi(1, 0)\| \leq \sum_{m=1}^n \log \left| \frac{D^{\frac{1}{n(n-2)}}}{|f'(\alpha_m)|^{\frac{1}{n-2}}} \right|$$

Lemma 4.5 estimates $|f'(\alpha_m)|^{\frac{1}{n-2}}$ as follows,

$$|f'(\alpha_m)| \geq 2^{-(n-1)^2} \frac{|D|}{M(F)^{2n-2}}.$$

Since D_F is large, definitely larger than $2^{-(n-1)^2}$, we have

$$|f'(\alpha_m)| \geq \frac{1}{M(F)^{2n-2}}.$$

This completes our proof. \square

Lemma 6.2. *Suppose that (x, y) is a solution to the equation $|F(x, y)| = 1$ for the binary form F in Theorem 1.1. Suppose that*

$$|x - \alpha_i y| = \min_{1 \leq j \leq n} |x - \alpha_j y|.$$

Then

$$\|\phi(x, y)\| \leq \frac{(n+1)^2}{4} \log \frac{1}{|x - \alpha_i y|} + n \log \left(|D|^{\frac{1}{n(n-2)}} M(F)^{\frac{2n-2}{n-2}} \right),$$

where $\|\cdot\|$ is the Euclidean norm.

Proof. Since $|F(x, y)| = \prod_{1 \leq j \leq n} |x - \alpha_j y| = 1$ and $|x - \alpha_i y| = \min_{1 \leq j \leq n} |x - \alpha_j y|$, we have $|x - \alpha_i y| \leq 1$. Let us assume that

$$|x - \alpha_{s_j} y| \leq 1, \quad \text{for } 1 \leq j \leq p$$

and

$$|x - \alpha_{b_k} y| > 1, \quad \text{for } 1 \leq k \leq n - p,$$

where $1 \leq p, s_j, b_k \leq n$. Since

$$|x - \alpha_i y| = \min_{1 \leq j \leq n} |x - \alpha_j y|,$$

we have

$$|\log |x - \alpha_{s_j} y|| \leq |\log |x - \alpha_i y||.$$

We also have

$$\prod_k |x - \alpha_{b_k} y| = \frac{1}{\prod_j |x - \alpha_{s_j} y|}.$$

Therefore, for any $1 \leq k \leq n - p$, we have

$$\log |x - \alpha_{b_k} y| \leq p \log \frac{1}{|x - \alpha_i y|}.$$

From here and the definition of $\phi(x, y)$ (see (13)), we conclude that

$$\begin{aligned} \|\phi(x, y)\| &\leq \sum_{m=1}^n \log \left| \frac{D^{\frac{1}{n(n-2)}}}{|f'(\alpha_m)|^{\frac{1}{n-2}}} \right| + (n-p)p |\phi_i(x, y)| + p |\phi_i(x, y)| \\ &= \sum_{m=1}^n \log \left| \frac{D^{\frac{1}{n(n-2)}}}{|f'(\alpha_m)|^{\frac{1}{n-2}}} \right| + ((n+1)p - p^2) |\phi_i(x, y)|. \end{aligned}$$

The function $f(p) = (n+1)p - p^2$ assumes its maximum value $\frac{(n+1)^2}{4}$ at $p = \frac{n+1}{2}$. To complete the proof we use our estimate in Lemma 6.1. \square

Lemma 6.3. *Let F be an irreducible monic binary form of degree n . Suppose that (x, y) is a solution to the Thue equation $F(x, y) = \pm 1$ with $y \geq M(F)^{1+(n-1)^2}$. Then*

$$\|\phi(1, 0)\| < \|\phi(x, y)\|.$$

Proof. Let $\alpha_1, \dots, \alpha_n$ be the roots of $F(z, 1) = 0$. Then

$$\left(\frac{x}{y} - \alpha_1\right) \dots \left(\frac{x}{y} - \alpha_n\right) = \frac{\pm 1}{y^n}.$$

There must exist a root α_j so that $\left|\frac{x}{y} - \alpha_j\right| \geq \frac{1}{y}$. By Lemma 4.5 and since $y \geq M(F)^{1+(n-1)^2}$, the absolute value of the term $\phi_j(x, y)$ alone exceeds $n \log \left(|D|^{\frac{1}{n(n-2)}} M(F)^{\frac{2n-2}{n-2}} \right)$. By Lemma 6.1, our proof is complete. \square

Let U be the unit group of the algebraic number field $\mathbb{Q}(\alpha)$. We define the mapping τ on U to be the obvious restriction of the embedding of $\mathbb{Q}(\alpha)$ in \mathbb{C}^n ; i.e. $\tau : u \mapsto (\sigma_1(u), \sigma_2(u) \dots \sigma_n(u))$, where $\sigma_i(u)$ are algebraic conjugates of u . By Dirichlet's unit theorem, we have a sequence of mappings

$$(14) \quad \tau : U \rightarrow V \subset \mathbb{C}^n$$

and

$$(15) \quad \log : V \rightarrow \Lambda,$$

where Λ is a $(r + s - 1)$ -dimensional lattice in \mathbb{R}^n and the mapping \log is defined as follows.

For $(x_1, \dots, x_n) \in V$, let

$$\log(x_1, x_2, \dots, x_n) := (\log |x_1|, \log |x_2|, \dots, \log |x_n|).$$

Suppose that $\{\lambda_2, \dots, \lambda_{r+s}\}$ is a system of fundamental units of $\mathbb{Q}(\alpha)$. Then $\log(\tau(\lambda_2)), \dots, \log(\tau(\lambda_{r+s}))$ form a basis for the lattice Λ . Moreover, every basis for Λ is associated with a system of fundamental units of $\mathbb{Q}(\alpha)$. So we will fix a system of fundamental units $\{\lambda_2, \dots, \lambda_{r+s}\}$ so that $\log(\tau(\lambda_2)), \dots, \log(\tau(\lambda_{r+s}))$ are respectively first to $r + s - 1$ -th successive minima of the lattice Λ (see [6], for the definition of successive minima). Therefore,

$$\|\log(\tau(\lambda_2))\| \leq \dots \leq \|\log(\tau(\lambda_{r+s}))\|,$$

where $\|\cdot\|$ is the Euclidean norm. If (x, y) is a pair of solution to (1) then $\frac{x - \alpha_i y}{x - \alpha_j y}$ is a unit in $\mathbb{Q}(\alpha_i, \alpha_j)$ and we may write

$$(16) \quad \phi(x, y) = \phi(1, 0) + \sum_{k=2}^{r+s} m_k \log(\tau(\lambda_k)), \quad m_k \in \mathbb{Z}.$$

7. LAYERS OF SOLUTIONS

As we defined in Section 5, a solution (x, y) is said to be related to α_i if

$$|x - \alpha_i y| = \min_{1 \leq j \leq n} |x - \alpha_j y|.$$

Fix a positive real number Y_0 . Let us first find a bound for the number of solutions (x, y) with $0 < y \leq Y_0$. We may suppose that $F(x, y)$ is a monic form with integral coefficients and has the smallest Mahler measure among all equivalent monic forms. Following Stewart [22] and Bombieri and Schmidt [5], we will estimate the number of solutions (x, y) to (1) for which $0 < y \leq Y_0$. For binary form

$$F(x, y) = (x - \alpha_1 y) \dots (x - \alpha_n y)$$

put

$$L_i(x, y) = x - \alpha_i y$$

for $i = 1, \dots, n$. Then

Lemma 7.1. *Suppose F is a monic binary form with integral coefficients. Then for every solution (x, y) of (1) we have*

$$\frac{1}{L_i(x, y)} - \frac{1}{L_j(x, y)} = (\beta_j - \beta_i)y,$$

where β_1, \dots, β_n are such that the form

$$J(u, w) = (u - \beta_1 w) \dots (u - \beta_n w)$$

is equivalent to F .

Proof. This is Lemma 4 of [22] and Lemma 3 of [5], by taking $(x_0, y_0) = (1, 0)$. \square

For every solution $(x, y) \neq (1, 0)$ of (1), fix $j = j(x, y)$ with

$$|L_j(x, y)| \geq 1.$$

Then, by Lemma 7.1,

$$(17) \quad \frac{1}{|L_i(x, y)|} \geq |\beta_j - \beta_i||y| - 1.$$

For complex conjugate $\bar{\beta}_j$ of β_j , where $j = j(x, y)$, we also have

$$\frac{1}{|L_i(x, y)|} \geq |\bar{\beta}_j - \beta_i||y| - 1.$$

Hence

$$\frac{1}{|L_i(x, y)|} \geq |\operatorname{Re}(\beta_j) - \beta_i||y| - 1,$$

where $\operatorname{Re}(\beta_j)$ is the real part of β_j . We now choose an integer $m = m(x, y)$ with $|\operatorname{Re}(\beta_j) - \beta_j| \leq 1/2$, and we obtain

$$(18) \quad \frac{1}{|L_i(x, y)|} \geq \left(|m - \beta_i| - \frac{1}{2} \right) |y| - 1,$$

for $i = 1, \dots, n$.

For $1 \leq i \leq n$, Let \mathfrak{X}_i be the set of solutions to (1) with $1 \leq y \leq Y_0$ and $|L_i(x, y)| \leq \frac{1}{2y}$.

Remark 1. When α_k and α_l are complex conjugates, $\mathfrak{X}_l = \mathfrak{X}_k$ and therefore we only need to consider $r + s$ different sets \mathfrak{X}_i .

Remark 2. If a solution (x, y) with $1 \leq y \leq Y_0$ is related to α_i then $(x, y) \in \mathfrak{X}_i$.

Remark 3. A solution (x, y) may belong to more than one set \mathfrak{X}_i .

Lemma 7.2. Suppose (x_1, y_1) and (x_2, y_2) are two distinct solutions in \mathfrak{X}_i with $y_1 \leq y_2$. Then

$$\frac{y_2}{y_1} \geq \frac{2}{7} \max(1, |\beta_i(x_1, y_1) - m(x_1, y_1)|).$$

Proof. This is Lemma 5 of [22] and Lemma 4 of [5]. \square

Lemma 7.3. Suppose (x, y) is a solution to (1) with $y > 0$ and $|L_i(x, y)| > \frac{1}{2y}$. Then

$$|m(x, y) - \beta_i(x, y)| \leq \frac{7}{2}.$$

Proof. This is Lemma 6 of [22]. \square

By Lemma 7.1 the form

$$J(u, w) = (u - \beta_1 w) \dots (u - \beta_n w)$$

is equivalent to $F(x, y)$ and therefore the form

$$\hat{J}(u, w) = (u - (\beta_1 - m)w) \dots (u - (\beta_n - m)w)$$

is also equivalent to $F(x, y)$. Therefore, since we assumed that F has the smallest Mahler measure among its equivalent forms, we get

$$(19) \quad \prod_{i=1}^n \max(1, |\beta_1(x, y) - m(x, y)|) \geq M(F).$$

For each set \mathfrak{X}_i that is not empty, let $(x^{(i)}, y^{(i)})$ be the element with the largest value of y . Let \mathfrak{X} be the set of solutions of (1) with $1 \leq y \leq Y_0$ minus the elements $(x^{(1)}, y^{(1)}), \dots, (x^{(r+s)}, y^{(r+s)})$. Suppose that, for integer i , the set \mathfrak{X}_i is non-empty. Index the elements of \mathfrak{X}_i as

$$(x_1^{(i)}, y_1^{(i)}), \dots, (x_v^{(i)}, y_v^{(i)}),$$

so that $y_1^{(i)} \leq \dots \leq y_v^{(i)}$ (note that $(x_v^{(i)}, y_v^{(i)}) = (x^{(i)}, y^{(i)})$). By Lemma 7.2

$$\frac{2}{7} \max\left(1, \left|\beta_i(x_k^{(i)}, y_k^{(i)})\right|\right) \leq \frac{y_{k+1}^{(i)}}{y_k^{(i)}}$$

for $k = 1, \dots, v-1$. Hence

$$\prod_{(x, y) \in \mathfrak{X} \cap \mathfrak{X}_i} \frac{2}{7} \max\left(1, \left|\beta_i(x_k^{(i)}, y_k^{(i)})\right|\right) \leq Y_0.$$

For (x, y) in \mathfrak{X} but not in \mathfrak{X}_i we have, by Lemma 7.3,

$$\frac{2}{7} \max\left(1, \left|\beta_i(x_k^{(i)}, y_k^{(i)})\right|\right) \leq 1.$$

Thus

$$\prod_{(x, y) \in \mathfrak{X}} \frac{2}{7} \max\left(1, \left|\beta_i(x_k^{(i)}, y_k^{(i)})\right|\right) \leq Y_0.$$

Let $|\mathfrak{X}|$ be the cardinality of \mathfrak{X} . Comparing the above inequality with (19), we obtain

$$(20) \quad \left(\left(\frac{2}{7}\right)^n M(F)\right)^{|\mathfrak{X}|} \leq Y_0^{r+s},$$

for we have $r+s$ different \mathfrak{X}_i . Therefore, by (8), we have

$$\left(\frac{2}{7}\right)^n M(F) \geq M(F)^\theta.$$

Here $\theta = \theta(D)$ may be taken equal to $\frac{1}{2}$, for the discriminant D is assumed to be very large. From here and by (20),

$$|\mathfrak{X}| \leq \frac{(r+s) \log Y_0}{\theta \log M(F)}.$$

Thus, when $Y_0 = M(F)^2$ and D_F is large enough, we have $|\mathfrak{X}| < 4(r+s)$. Consequently, there are at most $5(r+s)$ solutions (x, y) with $0 < y \leq M(F)^2$. We should remark here that we repeat Stewart's [22] approach for counting solutions with small y and no improvement has taken place in estimating θ . The reason that our value for θ is smaller is that we are working with forms with larger discriminant.

In order to count the number of solutions (x, y) with $M(F)^2 < y < M(F)^{1+(n-1)^2}$, we will need the following refinement of an inequality of Lewis and Mahler:

Lemma 7.4. *Let F be a binary form of degree $n \geq 3$ with integer coefficients and nonzero discriminant D . For every pair of integers (x, y) with $y \neq 0$*

$$\min_{\alpha} \left| \alpha - \frac{x}{y} \right| \leq \frac{2^{n-1} n^{n-1/2} (M(F))^{n-2} |F(x, y)|}{|D|^{1/2} |y|^n},$$

where the minimum is taken over the zeros α of $F(z, 1)$.

Proof. This is Lemma 3 of [22]. \square

Lemma 7.5. *Let $F(x, y)$ be a binary form with integral coefficients, degree n and discriminant D , where $|D| \geq D_0(n)$. Suppose that α_i is a real root of $F(z, 1) = 0$. Then related to α_i , there are at most 2 solutions for equation (1) in integers x and y with $M(F)^2 < y < M(F)^{1+(n-1)^2}$.*

Proof. Assume that (x_1, y_1) , (x_2, y_2) and (x_3, y_3) are three distinct solutions to (1) and all related to α_i with $y_3 > y_2 > y_1 > M(F)^2$. By Lemma 7.4, for $j = 1, 2$, we have

$$\left| \frac{x_{j+1}}{y_{j+1}} - \frac{x_j}{y_j} \right| \leq \frac{2^n n^{n-1/2} (M(F))^{n-2}}{|D|^{1/2} |y_j|^n}.$$

Since (x_1, y_1) , (x_2, y_2) and (x_3, y_3) are distinct solutions, for $j = 1, 2$, we have $|x_{j+1}y_j - x_jy_{j+1}| \geq 1$. Therefore,

$$\left| \frac{1}{y_j y_{j+1}} \right| \leq \left| \frac{x_{j+1}}{y_{j+1}} - \frac{x_j}{y_j} \right| \leq \frac{M(F)^{n-2}}{|y_j|^n}.$$

This is because we assumed that $|D|$ is large. Thus,

$$(21) \quad \frac{y_j^{n-1}}{M(F)^{n-2}} \leq y_{j+1}.$$

Following Stewart [22], we define δ_j , for $j = 1, 2, 3$, by

$$y_j = M(F)^{1+\delta_j}.$$

By (8), $M(F) > 1$ and so (21) implies that

$$(n-1)\delta_j \leq \delta_{j+1}.$$

From here, we conclude that

$$y_3 \geq M(F)^{1+(n-1)^2}.$$

In other words, related to each real root α_i , there are at most 2 solutions in x and y with $M(F)^2 < y < M(F)^{1+(n-1)^2}$. \square

Lemma 7.6. *Let $F(x, y)$ be a binary form with integral coefficients, degree n and discriminant D , where $|D| \geq D_0(n)$. Suppose that α_i is a non-real root of $F(z, 1) = 0$. Then related to α_i , there exists at most 1 solution to equation (1) in integers x and y with $M(F)^2 < y < M(F)^{1+(n-1)^2}$.*

Proof. Assume that (x_1, y_1) and (x_2, y_2) are two distinct solutions to (1) and all related to α_i , a non-real root of $F(z, 1) = 0$, with $y_2 > y_1 > M(F)^2$. Similar to (21) in the proof of Lemma 7.5, we have

$$\frac{y_1^{n-1}}{M(F)^{n-2}} \leq y_2.$$

This contradicts (11), since $y_1 > M(F)^2$ and $M(F)$ is large. Therefore, related to each non-real α_i , there is at most 1 solutions in x and y with $M(F)^2 < y < M(F)^{1+(n-1)^2}$. \square

So we conclude that there are at most $7r + 6s$ solutions (x, y) with $0 < y < M(F)^{1+(n-1)^2}$ to equation (1) when $F(z, 1) = 0$ has r real roots and $2s$ non-real ones.

Stewart [22] invented the above method to count all solutions with $y > M(F)^2$. He obtained the bound

$$n \left(4 + \frac{\log 331890}{\log(n-1)} \right)$$

for the number of solutions to (1) with $y > M(F)^2$ (see page 815 of [22]). Our method allows us to save the summand $\frac{\log 331890}{\log(n-1)}$. This gives us a better bound for binary forms with smaller degree.

The rest of paper is devoted to count the number of solutions (x, y) with $y \geq M(F)^{1+(n-1)^2}$. As we commented in Section 5, we need to consider this case only when we study the solutions (x, y) related to the real roots of $F(x, 1) = 0$.

Lemma 7.7. *For every fixed integer m , there are at most $2r + 2s - 2$ solutions (x, y) to (1) for which in (16), $m_{r+s} = m$.*

Proof. Let S be the $(r + s - 1)$ -dimensional affine space of all vectors

$$\phi(1, 0) + \sum_{i=2}^{r+s} \mu_i \log(\tau(\lambda_i)) \quad (\mu_i \in \mathbb{R}).$$

Let $\mu_{r+s} = m$. Then the points

$$\phi(1, 0) + \sum_{i=2}^{r+s-1} \mu_i \log(\tau(\lambda_i)) + m \log(\tau(\lambda_{r+s}))$$

form an $(r + s - 2)$ -dimensional hyperplane S_1 of S . Put $f(t) = F(t, 1)$. For $t \in \mathbb{R}$, define $y(t)$ and $x(t)$ as follows:

$$\begin{aligned} y(t) &:= |f(t)|^{-1/n}, \\ x(t) &:= ty(t). \end{aligned}$$

Similar to $\phi(x, y)$, we define the curve $\phi(t)$ on \mathbb{R} :

$$\phi(t) = (\phi_1(t), \phi_2(t), \dots, \phi_n(t)),$$

where, for $1 \leq m \leq n$,

$$\phi_m(t) = \log \left| \frac{D^{\frac{1}{n(n-2)}}(x(t) - \alpha_m y(t))}{|f'(\alpha_m)|^{\frac{1}{n-2}}} \right|.$$

Observe that for an integral solution (x, y) to (1) and $\phi(x, y)$ defined in (13), we have

$$\phi(x, y) = \phi \left(\frac{x}{y} \right).$$

Let $\vec{N} = (N_1, \dots, N_n) \in S$ be the normal vector of S_1 . Then the number of times that the curve $\phi(t)$ intersects S_1 equals the number of solutions in t to

$$(22) \quad \vec{N} \cdot \phi(t) = 0.$$

We have

$$\lim_{t \rightarrow \alpha_i^+} \log |t - \alpha_i| = -\infty$$

and

$$\lim_{t \rightarrow \alpha_i^-} \log |t - \alpha_i| = -\infty.$$

Note that if α_i is a non-real root of $F(x, 1)$ then $\bar{\alpha}_i$, the complex conjugate of α_i is also a root and we have

$$\log |t - \alpha_i| = \log |t - \bar{\alpha}_i|.$$

If $\alpha_1, \dots, \alpha_r$ are the real roots and $\alpha_{r+1}, \dots, \alpha_{r+s}, \alpha_{r+s+1}, \dots, \alpha_{r+2s}$ are non-real roots with $\alpha_{r+s+k} = \bar{\alpha}_{r+k}$, then the derivative $\frac{d}{dt} \left(\vec{N} \cdot \phi(t) \right)$ can be written as $\frac{P(t)}{Q(t)}$, where $Q(t) = (t - \alpha_1) \dots (t - \alpha_r)(t - \alpha_{r+1}) \dots (t - \alpha_{r+s})$ and $P(t)$ is a polynomial of degree $r + s - 1$. Therefore, the derivative has at most $r + s - 1$ zeros and consequently, the equation (22) can not have more than $2r + 2s - 2$ solutions. \square

Definition of the set \mathfrak{A} . Assume that equation (1) has more than $2r + 2s - 2$ solutions. Then we can list $(1, 0)$ and $2r + 2s - 3$ other solutions (x_i, y_i) ($1 \leq i \leq 2r + 2s - 3$), so that $\|\phi(x_i, y_i)\|$ are the smallest among all $\|\phi(x, y)\|$, where (x, y) varies over all non-trivial pairs of solutions. We denote the set of all these $2r + 2s - 2$ solutions by \mathfrak{A} .

The important property of \mathfrak{A} is that for every solution $(x_0, y_0) \in \mathfrak{A}$ and every solution $(x, y) \notin \mathfrak{A}$ to (1) with $y \geq M(F)^{1+(n-1)^2}$, by Lemma 6.3 and the definition, we have

$$\|\phi(x_0, y_0)\| \leq \|\phi(x, y)\|.$$

Corollary 7.8. *Let $(x, y) \notin \mathfrak{A}$ be a solution to (1) with $y \geq M(F)^{1+(n-1)^2}$. Then*

$$\|\log(\tau(\lambda_2))\| \leq \dots \leq \|\log(\tau(\lambda_{r+s}))\| \leq 2 \|\phi(x, y)\|.$$

Proof. Since we have assumed that $\|\log(\tau(\lambda_2))\| \leq \dots \leq \|\log(\tau(\lambda_{r+s}))\|$, it is enough to show that $\|\log(\tau(\lambda_{r+s}))\| \leq 2 \|\phi(x, y)\|$. By Lemma 7.7, there is at least one small solution $(x_0, y_0) \in \mathfrak{A}$ so that

$$\phi(x, y) - \phi(x_0, y_0) = \sum_{i=2}^{r+s} k_i \log(\tau(\lambda_i)),$$

with $k_n \neq 0$. Since $\{\log(\tau(\lambda_i))\}$ is a reduced basis for the lattice Λ in (15), by Lemma 6.3 and from the definition of \mathfrak{A} we conclude that

$$\|\log(\tau(\lambda_{r+s}))\| \leq \|\phi(x, y) - \phi(x_0, y_0)\| \leq 2 \|\phi(x, y)\|.$$

\square

Lemma 7.9. *Suppose $(x, y) \notin \mathfrak{A}$. Then*

$$\|\phi(x, y)\| \geq \frac{1}{2} \log \left(\frac{|D|^{\frac{1}{n(n-1)}}}{2} \right).$$

Proof. Let $(x', y') \in \mathfrak{A}$ be a pair of solutions to equation (1) and α_i and α_j be two distinct roots of the polynomial $F(x, 1)$. We have

$$\begin{aligned} \left| e^{\phi_i(x', y') - \phi_i(x, y)} - e^{\phi_j(x', y') - \phi_j(x, y)} \right| &= \left| \frac{x' - y'\alpha_i}{x - y\alpha_i} - \frac{x' - y'\alpha_j}{x - y\alpha_j} \right| \\ &= \frac{|\alpha_i - \alpha_j| |xy' - yx'|}{|x - y\alpha_i| |x - y\alpha_j|} \\ &\geq \frac{|\alpha_i - \alpha_j|}{|x - y\alpha_i| |x - y\alpha_j|}. \end{aligned}$$

The last inequality follows from the fact that $|xy' - yx'|$ is a non-zero integer. Since $|\phi_i| < \|\phi\|$ and $\|\phi(x', y')\| < \|\phi(x, y)\|$, we may conclude

$$\begin{aligned} \left(2e^{2\|\phi(x, y)\|} \right)^{\frac{n(n-1)}{2}} &\geq \prod_{1 \leq i < j \leq n} \left| e^{\phi_i(x', y') - \phi_i(x, y)} - e^{\phi_j(x', y') - \phi_j(x, y)} \right| \\ &\geq \prod_{1 \leq i < j \leq n} \left| \frac{x' - y'\alpha_i}{x - y\alpha_i} - \frac{x' - y'\alpha_j}{x - y\alpha_j} \right| \\ &\geq \prod_{1 \leq i < j \leq n} \frac{|\alpha_i - \alpha_j|}{|x - y\alpha_i| |x - y\alpha_j|} = \sqrt{|D|}. \end{aligned}$$

□

8. DISTANCE FUNCTIONS

Suppose that $(x, y) \neq (1, 0)$ is a solution to (1) and let $t = \frac{x}{y}$. We have

$$\phi(x, y) = \phi(t) = \sum_{i=1}^n \log \frac{|t - \alpha_i|}{|f'(\alpha_i)|^{\frac{1}{n-2}}} \mathbf{b}_i,$$

where,

$$\mathbf{b}_i = \frac{1}{n} (-1, \dots, -1, n-1, -1, \dots, -1).$$

Without loss of generality, we will suppose that the pair of solution (x, y) is related to α_n ;

$$|x - \alpha_n y| = \min_{1 \leq j \leq n} |x - \alpha_j y|.$$

We may write

$$(23) \quad \phi(x, y) = \phi(t) = \sum_{i=1}^{n-1} \log \frac{|t - \alpha_i|}{|f'(\alpha_i)|^{\frac{1}{n-2}}} \mathbf{c}_i + E_n \mathbf{b}_n,$$

where, for $1 \leq i \leq n-1$,

$$(24) \quad \mathbf{c}_i = \mathbf{b}_i + \frac{1}{n-1} \mathbf{b}_n, \quad E_n = \log \frac{|t - \alpha_n|}{|f'(\alpha_n)|^{\frac{1}{n-2}}} - \frac{1}{n-1} \sum_{i=1}^{n-1} \log \frac{|t - \alpha_i|}{|f'(\alpha_i)|^{\frac{1}{n-2}}}$$

One can easily observe that, for $1 \leq i \leq n$,

$$(25) \quad \mathbf{c}_i \perp \mathbf{b}_n, \quad \text{and } \|\mathbf{c}_i\| = \frac{\sqrt{n^2 - 3n + 2}}{n-1}.$$

Lemma 8.1. *Let*

$$\mathbf{L}_n = \left\{ \sum_{i=1}^{n-1} \log \frac{|\alpha_n - \alpha_i|}{|f'(\alpha_i)|^{\frac{1}{n-1}}} \mathbf{c}_i + z \mathbf{b}_n, \quad z \in \mathbb{R} \right\}.$$

Suppose that (x, y) is a solution to (1) with

$$|x - \alpha_n y| = \min_{1 \leq j \leq n} |x - \alpha_j y| \text{ and } y > M(F)^{1+(n-1)^2}.$$

Then the distance between $\phi(x, y)$ and the line \mathbf{L}_n is less than

$$\frac{1}{M(F)^{n(n-1)}} \exp\left(\frac{-4 \|\phi(x, y)\|}{(n+1)^2}\right).$$

Proof. The distance between $\phi(x, y)$ and \mathbf{L}_n is equal to

$$\left\| \sum_{i=1}^{n-1} \log \frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \mathbf{c}_i \right\|,$$

where $t = \frac{x}{y}$. We will show that when $i \neq n-1$,

$$\left| \log \frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \right| < \frac{|t - \alpha_n|}{\min_{i \neq j} \{|\alpha_j - \alpha_i|\}},$$

We will consider two cases $|t - \alpha_i| > |\alpha_n - \alpha_i|$ and $|t - \alpha_i| \leq |\alpha_n - \alpha_i|$. First assume that $|t - \alpha_i| > |\alpha_n - \alpha_i|$. We have

$$\left| \log \frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \right| = \log \frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \leq \log \left(\frac{|t - \alpha_n|}{|\alpha_n - \alpha_i|} + 1 \right) < \frac{|t - \alpha_n|}{|\alpha_i - \alpha_n|}.$$

Now assume that $|t - \alpha_i| \leq |\alpha_n - \alpha_i|$. Then

$$\left| \log \frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \right| = \log \frac{|\alpha_n - \alpha_i|}{|t - \alpha_i|} \leq \log \left(\frac{|t - \alpha_n|}{|t - \alpha_i|} + 1 \right) < \frac{|t - \alpha_n|}{|\alpha_i - t|}.$$

Note that, since we assumed t is closer to α_n ,

$$|\alpha_i - t| \geq \frac{|\alpha_i - t| + |\alpha_n - t|}{2} \geq \frac{|\alpha_i - \alpha_n|}{2}.$$

Hence, we obtain

$$(26) \quad \left| \log \frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \right| < 2 \frac{|t - \alpha_n|}{m},$$

where $m = \min_{i \neq j} \{|\alpha_j - \alpha_i|\}$. This, together with (25), gives

$$\left\| \sum_{i=1}^{n-1} \log \frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \mathbf{c}_i \right\| < \frac{2\sqrt{n(n^2 - 3n + 2)}}{n-1} \frac{|u|}{m},$$

where $u = t - \alpha_n$. Using (8), we obtain

$$(27) \quad \left\| \sum_{i=1}^{n-1} \log \frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \mathbf{c}_i \right\| < \frac{2M(F)^{n-1}(n+1)^n \sqrt{n(n^2 - 3n + 2)}}{\sqrt{3}(n-1)} |u|.$$

We shall estimate $|u|$ now. From Lemma 6.2 we have

$$\|\phi(x, y)\| - n \log \left(|D|^{\frac{1}{n(n-2)}} M(F)^{\frac{2n-2}{n-2}} \right) \leq \frac{(n+1)^2}{4} \log \frac{1}{|x - \alpha_n y|},$$

which implies

$$\log |yu| < \frac{-4 \|\phi(x, y)\|}{(n+1)^2} + \frac{4n}{(n+1)^2} \log \left(|D|^{\frac{1}{n(n-2)}} M(F)^{\frac{2n-2}{n-2}} \right).$$

Therefore,

$$|u| < \exp \left(\frac{-4 \|\phi(x, y)\|}{(n+1)^2} \right) \frac{\exp \left(\frac{4n}{(n+1)^2} \log \left(|D|^{\frac{1}{n(n-2)}} M(F)^{\frac{2n-2}{n-2}} \right) \right)}{|y|}$$

Comparing this with (27), since we took $n \geq 5$ and $|y| > M(F)^{1+(n-1)^2}$, our proof is complete. \square

For 3 distinct roots of $F(x, 1) = 0$, say α_i, α_j and α_n , let us define

$$T_{i,j}(t) := \log \left| \frac{(t - \alpha_i)(\alpha_n - \alpha_j)}{(t - \alpha_j)(\alpha_n - \alpha_i)} \right|,$$

so that for a pair of solution $(x, y) \neq (1, 0)$,

$$\begin{aligned} T_{i,j}(x, y) = T_{i,j}(t) &= \log \left| \frac{\alpha_n - \alpha_i}{\alpha_n - \alpha_j} \right| + \log \left| \frac{t - \alpha_j}{t - \alpha_i} \right| \\ (28) \qquad \qquad \qquad &= \log |\lambda_{i,j}| + \sum_{k=2}^{r+s} m_k \log \left| \frac{\lambda_k}{\lambda'_k} \right|, \end{aligned}$$

where $t = \frac{x}{y}$, $\lambda_{i,j} = \frac{\alpha_n - \alpha_i}{\alpha_n - \alpha_j}$, $m_k = m_k(x, y) \in \mathbb{Z}$, and for $2 \leq k \leq r+s$, λ_k are the fundamental units of number field $\mathbb{Q}(\alpha_i)$ and $\sigma(\lambda_k) = \lambda'_k$ are the fundamental units of the number field $\mathbb{Q}(\alpha_j)$ and index σ is the \mathbb{Q} -isomorphism from $\mathbb{Q}(\alpha_i)$ to $\mathbb{Q}(\alpha_j)$ such that $\sigma(\alpha_i) = \alpha_j$. The function $T(x, y)$ cries out to be treated by Baker's theory of linear forms in logarithms. For this we will wait till the very last part of the paper, Section 10, where we estimate $|T_{i,j}|$ from below. The following lemma gives an upper bound upon $|T_{i,j}|$.

Lemma 8.2. *Let (x, y) be a pair of solution to (1) with $|y| > M(F)^{1+(n-1)^2}$. Then there exists a pair (i, j) for which*

$$|T_{i,j}(x, y)| < \frac{\sqrt{\frac{2}{n-2}}}{M(F)^{n(n-1)}} \exp \left(\frac{-4 \|\phi(x, y)\|}{(n+1)^2} \right).$$

Proof. Let us define

$$\beta_i = \begin{cases} \alpha_i & \text{if } i \leq n-1 \\ \beta_{i-n+1} & \text{if } i \geq n. \end{cases}$$

Note that

$$\begin{aligned}
& \sum_{k=1}^{n-2} \sum_{i=1}^{n-1} \log^2 \left| \frac{(t - \beta_i)(\alpha_n - \beta_{i+k})}{(\alpha_n - \beta_i)(t - \beta_{i+k})} \right| \\
&= 2(n-2) \sum_{i=1}^{n-1} \log^2 \left| \frac{t - \alpha_i}{\alpha_n - \alpha_i} \right| - 4 \sum_{\substack{j \neq i \\ j \neq n}} \log \left| \frac{t - \alpha_i}{\alpha_n - \alpha_i} \right| \log \left| \frac{t - \alpha_j}{\alpha_n - \alpha_j} \right| \\
&= 2(n-2) \sum_{i=1}^{n-1} \log^2 \left| \frac{t - \alpha_i}{\alpha_n - \alpha_i} \right| - 2 \sum_{i=1}^{n-1} \log \left| \frac{t - \alpha_i}{\alpha_n - \alpha_i} \right| \left| \sum_{\substack{j \neq i \\ j \neq n}} \log \left| \frac{t - \alpha_j}{\alpha_n - \alpha_j} \right| \right| \\
&= 2(n-2) \sum_{i=1}^{n-1} \log^2 \left| \frac{t - \alpha_i}{\alpha_n - \alpha_i} \right| - 2 \sum_{i=1}^{n-1} \log \left| \frac{t - \alpha_i}{\alpha_n - \alpha_i} \right| \log \left| \frac{\alpha_n - \alpha_i}{y^n f'(\alpha_n)(t - \alpha_n)(t - \alpha_i)} \right| \\
&= (2n-2) \sum_{i=1}^{n-1} \log^2 \left| \frac{t - \alpha_i}{\alpha_n - \alpha_i} \right| - 2 \log \left| \frac{1}{y^n f'(\alpha_n)(t - \alpha_n)} \right| \left| \sum_{i=1}^{n-1} \log \left| \frac{t - \alpha_i}{\alpha_n - \alpha_i} \right| \right| \\
&= (2n-2) \sum_{i=1}^{n-1} \log^2 \left| \frac{t - \alpha_i}{\alpha_n - \alpha_i} \right| - 2 \log^2 \left| \frac{1}{y^n f'(\alpha_n)(t - \alpha_n)} \right|
\end{aligned}$$

On the other hand, it follows from the proof of Lemma 8.1 that the distance between $\phi(x, y)$ and the line

$$\mathbf{L}_n = \sum_{i=1}^{n-1} \log \frac{|\alpha_n - \alpha_i|}{|f'(\alpha_i)|^{\frac{1}{n-1}}} \mathbf{c}_i + z \mathbf{b}_n, \quad z \in \mathbb{R},$$

is equal to $\left\| \sum_{i=1}^{n-1} \log \frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \mathbf{c}_i \right\|$. Further, by the definition of \mathbf{c}_i in (24), we have

$$\begin{aligned}
& \left\| \sum_{i=1}^{n-1} \log \frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \mathbf{c}_i \right\|^2 \\
&= \left\| \sum_{i=1}^{n-1} \log \left(\frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \right) - \frac{1}{n-1} \left| \log \frac{1}{y^n f'(\alpha_n)(t - \alpha_n)} \right| \mathbf{e}_i \right\|^2 \\
&= \sum_{i=1}^{n-1} \log^2 \left(\frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \right) - \frac{1}{n-1} \left| \log \frac{1}{y^n f'(\alpha_n)(t - \alpha_n)} \right| \\
&= \sum_{i=1}^{n-1} \log^2 \left| \frac{t - \alpha_i}{\alpha_n - \alpha_i} \right| - \frac{1}{n-1} \log \left| \frac{1}{y^n f'(\alpha_n)(t - \alpha_n)} \right| \left| \sum_{i=1}^{n-1} \log \left| \frac{t - \alpha_i}{\alpha_n - \alpha_i} \right| \right|
\end{aligned}$$

where $\{\mathbf{e}_i\}$ is the standard basis for \mathbb{R}^{n-1} . So, there must be a pair (i, j) , for which the following holds:

$$\begin{aligned} & \log^2 \left| \frac{(t - \alpha_i)(\alpha_n - \alpha_j)}{(t - \alpha_j)(\alpha_n - \alpha_i)} \right| \\ & < \frac{1}{(n-1)(n-2)} \sum_{k=1}^{n-2} \sum_{i=1}^{n-1} \log^2 \left| \frac{(t - \beta_i)(\alpha_n - \beta_{i+k})}{(\alpha_n - \beta_i)(t - \beta_{i+k})} \right| = \\ & = \frac{2(n-1)}{(n-1)(n-2)} \left\| \sum_{i=1}^{n-1} \log \frac{|t - \alpha_i|}{|\alpha_n - \alpha_i|} \mathbf{e}_i \right\|^2. \end{aligned}$$

Therefore, by Lemma 8.1

$$|T_{i,j}(x, y)| = \left| \log \left| \frac{(t - \alpha_i)(\alpha_n - \alpha_j)}{(t - \alpha_j)(\alpha_n - \alpha_i)} \right| \right| < \frac{\sqrt{\frac{2}{n-2}}}{M(F)^{(n-2)(n-3)}} \exp \left(\frac{-4 \|\phi(x, y)\|}{(n+1)^2} \right).$$

□

9. EXPONENTIAL GAP PRINCIPLE

Here our goal is to prove

Theorem 9.1. *Suppose that (x_1, y_1) , (x_2, y_2) and (x_3, y_3) are three pairs of non-trivial solutions to (1) with*

$$|x_j - \alpha_n y_j| \leq 1,$$

for $j \in \{1, 2, 3\}$. If $r_1 \leq r_2 \leq r_3$ then

$$r_3 > M(F)^{n(n-1)} \exp \left(\frac{4r_1}{(n+1)^2} \right) \frac{\sqrt{3}}{256} \left(\frac{\log \log n}{\log n} \right)^6,$$

where $r_j = \|\phi(x_j, y_j)\|$.

Proof. Suppose that (x_1, y_1) , (x_2, y_2) and (x_3, y_3) are three pairs of non-trivial solutions to (1). We note that three point $\phi_1 = \phi(x_1, y_1)$, $\phi_2 = \phi(x_2, y_2)$ and $\phi_3 = \phi(x_3, y_3)$ form a triangle Δ . The length of each side of Δ is less than $2r_3$. Lemma 8.1 shows that the height of Δ is at most

$$\frac{2}{M(F)^{n(n-1)}} \exp \left(\frac{-4r_1}{(n+1)^2} \right).$$

Therefore, the area of Δ is less than

$$(29) \quad \frac{4}{M(F)^{n(n-1)}} r_3 \exp \left(\frac{-4r_1}{(n+1)^2} \right).$$

To estimate the area of Δ from below, we note that $x - \alpha_i y$ is a unit in $\mathbb{Q}(\alpha_i)$ when (x, y) is a pair of solution to (1). This is because

$$F(x, y) = (x - \alpha_1 y)(x - \alpha_2 y) \dots (x - \alpha_n y) = \pm 1.$$

Define the vector $\tilde{\mathbf{e}}$ as follows

$$\tilde{\mathbf{e}} = \phi(x_1, y_1) - \phi(x_2, y_2) = \left(\log \left| \frac{x_1 - \alpha_1 y_1}{x_2 - \alpha_1 y_2} \right|, \dots, \log \left| \frac{x_1 - \alpha_n y_1}{x_2 - \alpha_n y_2} \right| \right).$$

Since $x_1 - \alpha_i y_1$ and $x_2 - \alpha_i y_2$ are units in $\mathbb{Q}(\alpha_i)$, by Lemma 4.3 we have

$$\|\tilde{\mathbf{e}}\| \geq nh(\alpha_1) > \frac{1}{4} \left(\frac{\log \log n}{\log n} \right)^3.$$

Now we can estimate each side of Δ from below to conclude that the area of the triangle Δ is greater than

$$\frac{\sqrt{3}}{64} \left(\frac{\log \log n}{\log n} \right)^6.$$

Comparing this with (29) we conclude that

$$\frac{4}{M(F)^{n(n-1)}} r_3 \exp\left(\frac{-4r}{(n+1)^2}\right) > \frac{\sqrt{3}}{64} \left(\frac{\log \log n}{\log n} \right)^6.$$

The result is immediate from here. \square

Remark. If all the roots of polynomial $F(x, 1)$ are real then we can use the following lower bound for the size of vector $\tilde{\mathbf{e}}$:

$$\|\tilde{\mathbf{e}}\| \geq n \log^2 \frac{1 + \sqrt{5}}{2}$$

(see exercise 2 on page 367 of [21]). Now an argument similar to the proof of Theorem 9.1 shows that in this case,

$$r_3 > \frac{M(F)^{n(n-1)}}{2} \exp\left(\frac{4r_1}{(n+1)^2}\right) \frac{\sqrt{3}}{8} n^2 \log^4 \frac{1 + \sqrt{5}}{2}.$$

10. LINEAR FORMS IN LOGARITHMS

Let σ be the \mathbb{Q} -isomorphism from $\mathbb{Q}(\alpha_i)$ to $\mathbb{Q}(\alpha_j)$ such that $\sigma(\alpha_i) = \alpha_j$. Suppose that there are three solutions (x_1, y_1) , (x_2, y_2) , (x_3, y_3) to (1) satisfying the following conditions

$$\begin{aligned} (x_l, y_l) &\notin \mathfrak{A}, \\ y_l &> M(F)^{1+(n-1)^2} \end{aligned}$$

and

$$|x_l - \alpha_n y_l| = \min_{1 \leq i \leq n} |x_l - \alpha_i y_l| \quad l \in \{1, 2, 3\}.$$

Assume that $r_1 \leq r_2 \leq r_3$, where $r_j = \|\phi(x_j, y_j)\|$. We will apply Matveev's lower bound to

$$\begin{aligned} T_{i,j}(x_3, y_3) = T_{i,j}(t_3) &= \log \left| \frac{\alpha_n - \alpha_i}{\alpha_n - \alpha_j} \right| + \log \left| \frac{t_3 - \alpha_j}{t_3 - \alpha_i} \right| \\ &= \log |\lambda_{i,j}| + \sum_{k=2}^{r+s} n_k \log \left| \frac{\lambda_k}{\lambda'_k} \right|, \end{aligned}$$

where (i, j) is chosen according to Lemma 8.2, $t_3 = \frac{x_3}{y_3}$ and $n_k = n_k(x_3, y_3) \in \mathbb{Z}$. In order to apply Proposition 4.6, we shall find appropriate values for the quantities A_k and B in the Proposition. Since Proposition 4.6 gives a better lower bound for linear forms in fewer number of logarithms, we will assume that $\lambda_{i,j}$ and $\frac{\lambda_k}{\lambda'_k}$ are multiplicatively independent and $T_{i,j}(x_5, y_5)$ is a linear form in $r + s$ logarithms. Recall that $r + s \leq r + 2s = n$.

Let λ be a unit in the number field $\mathbb{Q}(\alpha_i)$ and λ' be its corresponding algebraic conjugate in $\mathbb{Q}(\alpha_j)$. Let d be the degree of $\mathbb{Q}(\alpha_i, \alpha_j)$ over \mathbb{Q} . Then λ/λ' is a unit in $\mathbb{Q}(\alpha_i, \alpha_j)$ and

$$\begin{aligned} dh\left(\frac{\lambda}{\lambda'}\right) &= \frac{1}{2} \left| \log\left(\tau\left(\frac{\lambda}{\lambda'}\right)\right) \right|_1 \\ &= \frac{1}{2} |\log(\tau(\lambda))|_1 + \frac{1}{2} |\log(\tau(\lambda'))|_1 \\ &= nh(\lambda) + nh(\lambda'). \end{aligned}$$

We also have

$$h(\lambda') = h(\lambda) = \frac{1}{2n} |\log(\tau(\lambda))|_1.$$

Here $|\cdot|_1$ is the L_1 norm on \mathbb{R}^{s+t-1} and mappings τ and \log are defined in (14) and (15). So we have

$$h(\lambda) = \frac{1}{2n} |\log(\tau(\lambda))|_1 \leq \frac{\sqrt{2}}{2n} \|\log(\tau(\lambda))\|,$$

where $\|\cdot\|$ is the L_2 norm on \mathbb{R}^{r+s-1} . So when λ is a unit

$$(30) \quad \max\left\{dh\left(\frac{\lambda}{\lambda'}\right), \left|\log\left(\frac{\lambda}{\lambda'}\right)\right|\right\} \leq \sqrt{2} \|\log(\tau(\lambda))\|.$$

Therefore, by Corollary 7.8 we may choose the values A_k so that

$$A_k \leq 2\sqrt{2}r_1, \quad \text{for } 2 \leq k \leq r+s.$$

Let d_1 be the degree of $\mathbb{Q}(\alpha_i, \alpha_j, \alpha_n)$ over \mathbb{Q} . Then $d_1 \leq n(n-1)(n-2)$. We shall find a value for A_1 that is at least $\max\{dh(\gamma_1), |\log \gamma_1|\}$ (see the statement of Proposition 4.6). The following Lemma allows us to take

$$\frac{A_1}{d_1} = 2 \log 2 + \frac{4}{\sqrt{n}} r_1.$$

Lemma 10.1. *Let F be a binary form of degree n at least 3 and with integral coefficients. Assume (x, y) is a solution to (1) with $y > M(F)^{1+(n-1)^2}$. Then we have*

$$h\left(\frac{\alpha_k - \alpha_i}{\alpha_k - \alpha_j}\right) \leq 2 \log 2 + \frac{4}{\sqrt{n}} \|\phi(x, y)\|.$$

Proof. Let, $\beta_i = x - y\alpha_i$. We have

$$\frac{\alpha_k - \alpha_i}{\alpha_k - \alpha_j} = \frac{\beta_k - \beta_i}{\beta_k - \beta_j}.$$

Thus, Lemma 4.2 implies that

$$(31) \quad h\left(\frac{\alpha_k - \alpha_i}{\alpha_k - \alpha_j}\right) \leq 2 \log 2 + 4h(\beta_k).$$

Set $v_i = \log |\beta_i| = \phi_i(x, y) - \phi_i(1, 0)$ for $i = 1, 2, \dots, n$ and $\vec{v} = (v_1, v_2, \dots, v_n)$. Since β_k is a unit, we have

$$h(\beta_k) = \frac{1}{2n} \sum_{i=1}^n |v_i| = \frac{1}{2n} (s_1, s_2, \dots, s_n) \cdot \vec{v}$$

for some $s_1, s_2, \dots, s_n \in \{+1, -1\}$. Noting that $\|(s_1, s_2, \dots, s_n)\| = \sqrt{n}$, we get

$$h(\beta_k) \leq \frac{1}{2\sqrt{n}} \|\vec{v}\|.$$

On the other hand, by Lemma 6.3 we have

$$\|\vec{v}\| \leq \|\phi(x, y)\| + \|\phi(1, 0)\| \leq 2\|\phi(x, y)\|.$$

This, together with (31), completes the proof. \square

Put

$$B = \max\{1, \max\{b_k A_k / A_1 : 1 \leq k \leq r + s\}\}.$$

To estimate B , we note that since we have chosen $\tau(\lambda_k)$ ($2 \leq k \leq r + s$) so that they are successive minima for the lattice Λ (see Section 6), we have

$$m_k \|\log \tau(\lambda_k)\| \leq \|\phi(x_3, y_3)\| + \|\phi(1, 0)\| < 2\|\phi(x_3, y_3)\|.$$

Hence, we may take $B \leq r_3$, since $A_1 > 2$. We estimate other values of the quantities in Proposition 4.6 as follows:

$$\begin{aligned} d &\leq n!, \\ C_n &\leq \frac{60 \exp(n)(n+1)^{n+1} 2^{2n+2} (n+2)(n+5/2)n^2}{n!}, \\ C_0 &\leq 4 \log n!, \\ W_0 &\leq 2 \log r_3. \end{aligned}$$

Proposition 4.6 implies that

$$\begin{aligned} \log T_{i,j}(x_3, y_3) &> -K \log r_3 r_1^{r+s} \\ &> -K \log r_3 r_1^n, \end{aligned}$$

where the constant K can be taken equal to

$$(32) \quad 480 \exp(n)(n+1)^{n+1} 2^{7n+3/2} (n+2)(n+5/2)n^{5/2} (n-1)(n-2)n! \log(n!).$$

Comparing this with Lemma 8.2, we have

$$-\log \left(M(F)^{n(n-1)} \right) + \log \left(\sqrt{\frac{2}{n-2}} \right) + \frac{-4r_3}{(n+1)^2} > -K \log r_3 r_1^{n-1},$$

By Lemma 7.9 and since $|D| > D_0(n)$, the value r_3 is large enough to satisfy

$$r_3^{\frac{e-1}{e}} < \frac{r_3}{\log r_3}.$$

So we may find a constant K_1 depending only on n (see the values of $C(n)$, C_0 and W_0 in Proposition 4.6) so that

$$r_3 < K_1 r_1^{\frac{e}{e-1}n}.$$

Notice that K_1 may be chosen equal to

$$\left(\frac{(n+1)^2}{4} K \right)^{\frac{e}{e-1}}$$

By Lemma 9.1, we have

$$\begin{aligned} &M(F)^{n(n-1)} \exp \left(\frac{4r_1}{(n+1)^2} \right) \frac{\sqrt{3}}{256} \left(\frac{\log \log n}{\log n} \right)^6 \\ &< K_1 r_1^{1.6n} \end{aligned}$$

This is a contradiction, as in the above inequality the left hand side is greater than the right hand side. Hence, related to a root of $F(x, 1) = 0$, there are at most 2 solutions $(x, y) \notin \mathfrak{A}$, with $y > M(F)^{1+(n-1)^2}$. To see the contradiction, one can consider two different cases. If $\frac{4r_1}{(n+1)^2} > \frac{e}{e-1}n^{\frac{e}{e-1}}$ then $\exp\left(\frac{4r_1}{(n+1)^2}\right) > \frac{4r_1}{(n+1)^2}$ and by (8) and since $|D| \geq 2^{22}(n+1)^{10}n^n$, the value $M(F)^{n(n-1)}\frac{\sqrt{3}}{256}\left(\frac{\log \log n}{\log n}\right)^6$ exceeds the rest of right hand side. If $\frac{4r_1}{(n+1)^2} \leq \frac{e}{e-1}n^{\frac{e}{e-1}}$ then the value $M(F)^{n(n-1)}\frac{\sqrt{3}}{256}\left(\frac{\log \log n}{\log n}\right)^6$ alone exceeds the right hand side.

Remark. To estimate the value of A_1 we proved Lemma 10.1. Having the inequality

$$h\left(\frac{\alpha_n - \alpha_i}{\alpha_n - \alpha_j}\right) \leq 2 \log 2 + 4h(\alpha_n)$$

in hand, one may attempt to bound the logarithmic height of α , a root of $F(x, 1) = 0$, in terms of the discriminant of F . To do so recall that we have assumed that the binary form F has the smallest Mahler measure among all equivalent forms that are monic. We need this assumption to obtain an upper bound for the number of small solutions (see (20)). We also have

$$h(\alpha) = \frac{1}{n} \log M(\alpha) \leq \frac{1}{n} \log \left((n+1)^{1/2} H(\alpha) \right).$$

Therefore, we can apply Proposition 4.1 to our selected form F and assume that for each root α of $F(x, 1) = 0$, we have

$$h(\alpha) \leq \frac{1}{n} \log \left((n+1)^{1/2} \exp\{n^{4n^{12}} |D|^{6n^8}\} \right).$$

This will provide an explicit value for A_1 . Should one wish to use this to establish a contradiction similar as above, one has to start with 5 solutions (instead of 3) and after the contradiction, concludes that there are at most 4 solutions (instead of 2) with large y related to each root.

11. ACKNOWLEDGMENTS

Part of this work has been done while I was supported by Hausdorff Institute for Mathematics in Bonn. I would like to thank Professor Yann Bugeaud, Professor Jan-Hendrik Evertse, Professor Andrew Granville, Professor Kálmán Győry and Professor Ryotaro Okazaki for their helpful suggestions and comments. The content of this manuscript is improved due to the referee's care over the details and presentation.

REFERENCES

- [1] S. Akhtari, The method of Thue-Siegel for binary quartic forms, *Acta. Arith.*(2010).
- [2] S. Akhtari, R. Okazaki, Quartic Thue equations, *J. Number Theory.*130 (2010) no. 1, 40-60.
- [3] A. Baker. Contributions to the theory of Diophantine equations. I. On the representation of integers by binary forms, *Philos. Trans. Roy. Soc. London Ser. A* 263 (1967/1968), 173-191.
- [4] M.A. Bennett. On the representation of unity by binary cubic forms. *Trans. Amer. Math. Soc.* 353 (2001), 1507-1534.
- [5] E. Bombieri, W. M. Schmidt, On Thue's equation, *Invent. Math.* 88 (1987), 69-81.
- [6] J.W.S. Cassels, *An Introduction to the Geometry of Numbers.* Springer-Verlag, (1959).
- [7] H. Cohen, *Number Theory Volume II: Analytic and Modern Tools.* Graduate Texts in Mathematics, 240. Springer, New York, (2007).
- [8] J.H. Evertse, Estimates for reduced binary forms. *J. Reine Angew. Math.* 434 (1993), 159-190.

- [9] J.H. Evertse, Upper Bounds for the Numbers of Solutions of Diophantine Equations. Mathematical Centre Tracts, 168, Amsterdam, 1983.
- [10] J. H. Evertse and K. Györy, Effective finiteness results for binary forms with given discriminant, *Compositio Math*, 79(1991), 169-204.
- [11] J. H. Evertse and K. Györy, Thue inequalities with a small number of solutions, in: *The mathematical heritage of C.F. Gauss*, World Scientific Publ. Co., Singapore, 1991, 204-224.
- [12] K. Györy, Polynomials and binary forms with given discriminant, *Pub. Math. Debrecen*. 69 (2006), no. 4, 473-499.
- [13] K. Györy, Sur les polynome a coefficients entiers et the discriminant donne II, *Pub. Math. Debrecen*. 21 (1974), 125-144.
- [14] K. Györy, Thue inequalities with a small number of primitive solutions, *Period. Math. Hungar.* 42 (2001), no. 1-2, 199-209.
- [15] K. Mahler, An application of Jensen's formula to polynomials. *Mathematika*, 7 (1960) , 98-100.
- [16] K. Mahler, An inequality for the discriminant of a polynomial, *Michigan Math. J.* 11 (1964), 257-262.
- [17] K. Mahler, Zur Approximation algebraischer Zahlen II. Über die Anzahl der Darstellungen ganzer Zahlen durch Binärformen, *Math. Ann.* 108 (1933), 37-55.
- [18] E. M. Matveev, An explicit lower bound for a homogeneous rational linear form in logarithms of algebraic numbers, *Izv. Ross. Akad. Nauk Ser. Mat.* 62 (1998), 81-136, translation in *Izv. Math.* 62 (1998), 723-772.
- [19] E. M. Matveev, An explicit lower bound for a homogeneous rational linear form in logarithms of algebraic numbers, *Izv. Ross. Akad. Nauk Ser. Mat.* 64 (2000), 125-180, translation in *Izv. Math.* 64 (2000), 1217-1269.
- [20] R. Okazaki. Geometry of a cubic Thue equation, *Publ. Math. Debrecen*. 61 (2002), 267-314.
- [21] M. Pohst and H. Zassenhaus. *Algorithmic Algebraic Number Theory*, Cambridge University Press (1989).
- [22] C. L. Stewart, On the number of solutions of polynomial congruences and Thue equations, *J. Amer. Math. Soc.* 4 (1991), 793-835.
- [23] A. Thue, Berechnung aller Lösungen gewisser Gleichungen von der form $ax^r - by^r = f$. *Vid. Skrifter I Mat.-Naturv. Klasse* (1918), 1-9.
- [24] P. Voutier, An effective lower bound for the height of algebraic numbers, *Acta Arith.* 74 (1996), 8195.

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111 BONN, GERMANY

E-mail address: akhtari@mpim-bonn.mpg.de